

DOCUMENT RESUME

ED 042 487

PS 003 315

AUTHOR Cross, Marie Z.
TITLE A Nutritional Survey of Children in Head Start Centers in Central United States.
INSTITUTION Kansas Univ., Lawrence. Head Start Evaluation and Research Center.
SPONS AGENCY Office of Economic Opportunity, Washington, D.C.
PUB DATE 30 Nov 67
NOTE 6p.
EDRS PRICE MF-\$0.25 HC-\$0.40
DESCRIPTORS Eating Habits, Lunch Programs, *Nutrition, Preschool Programs, *Questionnaires, *Validity
IDENTIFIERS Project Head Start

ABSTRACT

The purpose of this study was to investigate the effectiveness of a questionnaire in evaluating the nutritional status of Head Start children in rural, small city, and urban areas in central United States. The questionnaire, which was primarily concerned with the food preferences of the children, was filled out by the mother of each of the 154 children in the study. Nutritional data was also obtained from analysis of one week's menus from each of the Head Start centers involved in the study. The only quantitative data obtained from the questionnaires was the amount of milk consumed by each child. Answers were so nutritionally favorable that it is suspected that the mothers may have given answers they felt they should give, rather than actual milk amounts. The foods reported as being preferred by the children are valuable sources of nutrients. The Head Start food programs, as reported in the menus, were nutritionally adequate for the one meal served. There were no significant differences in food preferences of the children in the three different areas studied nor in the types of food served in the Centers in these areas. The questionnaires proved to be of questionable value because of the paucity of data they generated concerning quantities of food, but the data was helpful in determining food preferences. (MH)

A Nutritional Survey of Children in Head Start Centers in Central United States₁

Marie Z. Cross

Introduction

This study was undertaken to evaluate the effectiveness of using a questionnaire to evaluate the nutritional status of children in Head Start centers in rural, small city and metropolitan areas in central United States. Evaluation of nutritional status on the basis of data obtained from questionnaires is hazardous at best and doing this without having a nutritionist obtain the history made it even more difficult. Since parents usually speak more freely about food likes and dislikes of their children than about actual quantities of food consumed, we used this approach to obtain knowledge of foods which the children were familiar with and liked. The only quantitative data that we attempted to obtain was with respect to the quantities of milk consumed daily.

In addition to the questionnaire data, we obtained and analyzed one weeks menus from each of the Head Start centers involved in this study to determine their contribution to the nutritional status of these children.

Procedure

A nutritional questionnaire was answered by the mother of each of the 154 children that were included in this study. The research team members were instructed in the manner in which the questionnaire should be used and in ways to establish rapport with the parent during the interview. The questionnaire (see Figure 1) was designed to determine the childrens food likes, dislikes and preferences. It was our purpose to see if this approach might result in our obtaining an accurate and honest answer to questions involving quantities of food consumed by the child. In this particular study the only quantity asked for was the amount of milk consumed.

Insert Figure 1 here

The questionnaire data was analyzed according to the type of area the children lived in namely, rural, small city or metropolitan area.

The breakfast or lunch and snack menus were obtained from each center for the week of April 10th through April 14th. These menus were analyzed for nutrient content assuming average pre-school age servings. The Agricultural Handbook No. 8 entitled Composition of Foods, and published by the United States Department of Agriculture was used as the source for determining the nutrient content of the menus. The entire five day food intake was analyzed and the total values were divided by five to get a value for an average daily intake of each of the nutrients. Using the National Research Council's recommended daily allowances as a standard we calculated the percentage of the child's daily nutritional needs which were supplied by the Head Start center's lunch program.

ED042487

PS003315

Results and Discussion

Table 1. summarizes the data obtained from the questionnaires. The results are expressed as percentages of children who liked certain of the nutritionally important foods and in the case of milk, those who received adequate amounts each day. The high percentage of children who were reported to receive three or more glasses of milk per day led us to suspect that the mothers may have given the answer they thought they should give rather than an honest value for the amount of milk consumed by their child.

Insert Table 1 here

In most cases where a child was reported to like meat, fish or poultry he liked all three but if a preference was cited it was most frequently chicken. In the fruit group bananas, apples and oranges were most frequently cited as preferences with oranges occurring less frequently than the other two.

The questions relating to vegetables preferences indicated carrots as the most frequently liked raw vegetables. In the cooked vegetable group potatoes, green beans and corn were cited most often as favorites.

If the children received their preferred foods frequently it would indicate that their nutritional needs were being met quite well. The foods most frequently reported as favorites are foods which are valuable sources of nutrients needed by children of pre-school age.

There were no significant differences in preferences of foods between children in a small city, rural or metropolitan area. As Table 1 shows most mothers felt their children received enough food but some of them indicated it was not always the right food.

About one-half of the children's favorite snacks would be classified as nutritious including such foods as milk, fruit and cheese while the other one-half preferred sweet snacks such as cookies and candy. The reported consumption of candy by these children was quite high with the children from the rural area consuming slightly less candy than those in the small city or metropolitan area.

Table 2 shows the amounts of the different essential nutrients provided by the lunch programs in the Head Start centers in the three areas studied. In addition this table shows the percent of the total days nutritional needs for the 3-6 year old that was provided by the school lunch program. The three areas differed in the type of meal and or snacks offered which in turn effected the type and amount of food served and hence the amounts of certain nutrients supplied. The children in the rural area received only lunch while those in the small city area received lunch plus a snack. In the metropolitan area the children received breakfast and a snack. In spite of these differences, all of the children received at least one-third of their days requirements of all but one of the nutrients and in most cases over one-half of their nutritional needs. In some cases the daily requirements were exceeded due to certain foods being included in the diet which were exceptionally high in those nutrients.

Insert Table 2 here

The data in Table 2 can be used only as a general indication of what the children actually received since we do not know the actual amounts of foods which were served and we don't know how much each individual child ate. However, these data do indicate that the Head Start centers which were studied were providing meals which gave generous amounts of the nutrients needed by children in this age group. This should provide good nutritional training for these children and hopefully some of this experience and training would be carried to their homes.

Conclusions

The data obtained from the questionnaires are of questionable value as far as quantities of food consumed is concerned, but these data do give some indication of the foods which these children would prefer to eat.

The nutritional analysis of the Head Start center menus indicated that the centers which we studied were providing adequate amounts and varieties of food for children of this age.

There were no significant differences in the food preferences of the children in the three different areas studied nor in the types of food served in the centers in these areas.

FOOTNOTES

The research reported herein was performed pursuant to a contract with the office of Economic Opportunity, Executive Office of the President, Washington, D.C., 20506. The opinions expressed herein are those of the author and should not be construed as representing the opinions or policy of any agency of the United States Government.

Bibliography

Composition of Foods, Agricultural Handbook No. 8, USDA publication, 1963

Eppright, E. S., M. B. Patton, A. L. Marlatt, M. L. Hathaway: Dietary Study methods, 5. Some problems in collecting dietary information about groups of children. J. Am. Dietet. A. 28: 43, 1952

Wilson, G. S. : A review of methods used in nutrition surveys conducted by the Interdepartmental Committee on Nutrition for National Defense (ICNND). Am. J. Clin. Nutrition 15: 29. 1964

Young, C. M.: Methodology for dietary studies in epidemiological survey. 2. Strengths and weaknesses of existing methods. Am. J. Public Health 50: 803, 1960

Table 1.

Summary of questionnaire data from the three areas studied

	<u>Small City</u>	<u>Rural</u>	<u>Metropolitan</u>
Total No. of children studied	46	51	57
Percent of children receiving 3 or more glasses of milk daily.	80.4	82.3	89.5
Percent of children who liked meat, fish, or poultry.	84.8	90.0	87.7
Percent of children who liked fruit.	100.0	96.0	96.5
Percent of children who liked vegetables.	84.8	92.1	66.6
Percent of mothers who reported their children ate enough.	91.3	78.4	77.0
Percent of children who ate candy frequently.	75.0	78.5	64.0

Table 2.

Evaluation of the menus from the Head Start
Centers in the three areas studied.

Nutrient	Small City (lunch & snack)		Rural (lunch only)		Metropolitan (breakfast & snack)	
	Am't per day	% of day's requirement supplied	Am't per day	% of day's requirement supplied	Am't per day	% of day's requirement supplied
Calories	677	42.3	769.0	47.6	604.0	37.6
Protein(gms.)	31.3	78.2	33.0	82.5	17.3	43.2
Calcium(mgs.)	665.0	83.0	382.0	47.7	359.0	44.9
Iron(mgs.)	3.6	36.0	4.4	44.0	3.2	32.0
Vit. A(IU's)	2767.	110.0	3274.0	131.0	1322.0	52.8
Ascorbic Acid(mgs)	87.6	175.0	44.0	88.0	47.0	94.0
Thiamin(mgs)	.43	71.0	.38	63.3	.34	56.0
Riboflavin(mgs)	.80	80.0	.78	78.0	.58	58.0
Niacin(mgs)	7.46	67.8	5.86	53.2	3.1	28.0

THE UNIVERSITY OF KANSAS HEAD START EVALUATION & RESEARCH CENTER

QUESTIONNAIRE

R-50 Cross-Bass Research Project

(To be carried out following the OEO Parent Interview)

Dr. Marie Cross, Investigator, Dept. Human Development

Dr. William Bass, Department of Anthropology

Does he (she) like milk? _____

About how much milk does he drink each day? _____

Does he like meat, chicken, or fish? _____

Would he rather eat: Baked beans? _____

Ham and beans? _____

Peanut butter sandwich? _____

Maccaroni and cheese? _____

Does he like fruit? _____

What kind of fresh fruit does he eat most often? _____

If not mentioned above, ask:

If he likes and eats: Apples _____

Oranges _____

Bananas _____

How often does he eat canned or dried fruit? _____

Does he like vegetables? _____

What is his favorite raw vegetable? _____

What are his favorite cooked vegetables? _____

If not mentioned above, ask:

Does he like? Green peas? _____

Green beans? _____

Cabbage? _____

Potatoes? _____

Do you feel the child eats enough? _____

Too much? _____

Not enough? _____

If he snacks between meals, what are his favorite snacks? _____

Does he eat much candy? _____ Does he drink much pop? _____

CHILD'S NAME _____ CHILD'S NO. _____

CENTER NAME _____ CLASS NO. _____

CENTER NO. _____

Figure 1