

DOCUMENT RESUME

ED 117 284

UD 015 696

AUTHOR Malina, Robert M.; And Others
 TITLE Body Dimensions and Proportions, White and Negro Children, 6-11 Years; United States. National Health Survey Series 11, No. 143
 INSTITUTION Health Resources Administration (DHEW/PHS), Bethesda, Md.
 REPORT NO DHEW-HRA-75-1625
 PUB DATE Dec 74
 NOTE 72p.
 AVAILABLE FROM Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402 Order Publication No. DHEW (HRA) 75-1625; \$1.30

EDRS PRICE MF-\$0.76 HC-\$3.32 Plus Postage
 DESCRIPTORS Age Groups; Body Height; Body Weight; Caucasians; Child Development; Children; Comparative Analysis; *Growth Patterns; Maturation; Negro Youth; *Physical Characteristics; Physical Development; Racial Characteristics; *Racial Differences; *Sex Differences; Statistical Data; *Statistical Surveys

ABSTRACT

The fifth in a series of reports presenting analyses and discussions of data on height, weight, and 28 other body measurements taken from a probability sample of noninstitutionalized children in the United States aged 6-11 years, this document compares the growth patterns of white and Negro children for 20 body measurements. Emphasis is placed on comparisons of age and sex specific means and medians for Negro and white children. These national estimates are based on cross-sectional data, which are said to limit the analyses to attained size rather than velocities of growth. All dimensions are said to increase almost linearly with age from 6 through 11 years in Negro and white children, both males and females. Conclusions derived from these data are considered to agree generally with most other anthropometric comparisons of American white and Negro children with these statistics providing figures held to be current, comprehensive, and reliable national estimates. Twenty-six detailed tables of statistical material are provided. (Author/AM)

 * Documents acquired by ERIC include many informal unpublished *
 * materials not available from other sources. ERIC makes every effort *
 * to obtain the best copy available. Nevertheless, items of marginal *
 * reproducibility are often encountered and this affects the quality *
 * of the microfiche and hardcopy reproductions ERIC makes available *
 * via the ERIC Document Reproduction Service (EDRS). EDRS is not *
 * responsible for the quality of the original document. Reproductions *
 * supplied by EDRS are the best that can be made from the original. *

ED117284

UD015696

Data from the
NATIONAL HEALTH SURVEY

Series 11
Number 143

Body Dimensions and Proportions, White and Negro Children 6-11 Years United States

Presents and discusses data on 20 anthropometric dimensions of children 6-11 years of age in the United States, 1963-65. The data are analyzed by race as well as age and sex and compare the patterns of body growth in Negro and white children in the United States.

U.S. DEPARTMENT OF HEALTH,
EDUCATION & WELFARE
NATIONAL INSTITUTE OF
EDUCATION

THIS DOCUMENT HAS BEEN REPRODUCED EXACTLY AS RECEIVED FROM THE PERSON OR ORGANIZATION ORIGINATING IT. POINTS OF VIEW OR OPINIONS STATED DO NOT NECESSARILY REPRESENT OFFICIAL NATIONAL INSTITUTE OF EDUCATION POSITION OR POLICY.

DHEW Publication No. (HRA) 75-1625

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service

Health Resources Administration
National Center for Health Statistics
Rockville, Md. December 1974

UD 015 696

Series 11 reports present findings from the National Health Examination Survey, which obtains data through direct examination, tests, and measurements of samples of the U.S. population. Reports 1 through '38 relate to the adult program, Cycle I of the Health Examination Survey. The present report is one of a number of reports of findings from the children and youth programs, Cycles II and III of the Health Examination Survey. These latter reports from Cycles II and III are being published in Series 11 but are numbered consecutively beginning with 101. It is hoped this will guide users to the data in which they are interested.



Library of Congress Cataloging in Publication Data

Malina, Robert M.

Body dimensions and proportions, white and Negro children, 6-11 years, United States.

(National Center for Health Statistics. Vital and health statistics. Series 11: Data from the National Health Survey, no. 143) (DHEW publication no. (HRA) 75-1625)

Supt. of Docs. no.: HE 20.2210: 11/143.

Includes bibliographical references.

1. Children in the United States—Anthropometry. 2. Children—Growth. 3. Negro children—Anthropometry. I. Hamill, Peter V., joint author. II. Lemeshow, Stanley, joint author. III. Title. IV. Series: United States. National Center for Health Statistics. Vital and health statistics. Series 11: Data from the National Health Survey, Data from the health examination survey, no. 143. V. Series: United States. Dept. of Health, Education, and Welfare. DHEW publication no. (HRA) 75-1625. [DNLM: 1. Anthropometry—In infancy and childhood. 2. Body height—In infancy and childhood. 3. Body weight—In infancy and childhood. W2AN148vk no. 143 1974]

RA407.3.A347 no. 143 [GN63] 312'.0973s [572] 74-4239

ISBN 0-8406-0011-9

NATIONAL CENTER FOR HEALTH STATISTICS

EDWARD B. PERRIN, Ph.D., *Director*

PHILIP S. LAWRENCE, Sc.D., *Deputy Director*

JACOB J. FELDMAN, Ph.D., *Acting Associate Director for Analysis*

GAJL F. FISHER, *Associate Director for the Cooperative Health Statistics System*

ELIJAH L. WHITE, *Associate Director for Data Systems*

IWAO M. MORIYAMA, Ph.D., *Associate Director for International Statistics*

EDWARD E. MINTY, *Associate Director for Management*

ROBERT A. ISRAEL, *Associate Director for Operations*

QUENTIN R. REMEIN, *Associate Director for Program Development*

PAUL E. LEAVERTON, *Acting Associate Director for Research*

ALICE HAYWOOD, *Information Officer*

DIVISION OF HEALTH EXAMINATION STATISTICS

ARTHUR J. McDOWELL, *Director*

JEAN-PIERRE HABICHT, M.D., Ph.D., *Special Assistant to Director*

PETER V. V. HAMILL, M.D., *Medical Advisor*

JEAN ROBERTS, *Chief, Medical Statistics Branch*

SIDNEY ABRAHAM, *Chief, Nutrition Statistics Branch*

ROBERT S. MURPHY, *Chief, Survey Planning and Development Branch*

COOPERATION OF THE BUREAU OF THE CENSUS

In accordance with specifications established by the National Health Survey, the Bureau of the Census, under a contractual agreement, participated in the design and selection of the sample, and carried out the first stage of the field interviewing and certain parts of the statistical processing.

Vital and Health Statistics Series 11-No. 143

DHEW Publication No. (HRA) 75-1625

Library of Congress Catalog Card Number 74-4239

CONTENTS

	Page
Method-----	3
Results-----	4
Weight and Height-----	4
Ponderal Index-----	5
Components of Stature: Sitting Height and Subischial Length-----	5
Sitting Height/Stature Ratio-----	6
Partitioning the Lower Extremity-----	7
Partitioning the Upper Extremity-----	8
Biacromial Breadth-----	9
Bicristal Breadth-----	10
Biacromial Breadth/Bicristal Breadth Relationship-----	10
Chest Dimensions-----	11
Bicondylar Breadth of the Femur-----	11
Limb Girths-----	11
Estimated Mid-Arm Muscle Circumference-----	12
Torso Girths-----	13
Discussion-----	14
Summary-----	17
References-----	18
List of Detailed Tables-----	20
Appendix I. Statistical Notes-----	48
The Survey Design-----	48
Notes on Response Rates-----	49
Parameter and Variance Estimation-----	49
Standards of Reliability and Precision-----	50
Hypothesis Testing-----	50
Appendix II. Techniques of Measurement and Quality Control-----	51
Techniques of Measurement-----	51
Equipment-----	52
Measuring Procedures and Definitions-----	53
Derived Measurements-----	56
Indexes and Ratios-----	56
Quality Control and Estimation of Residual Measurement Error-----	57
Monitoring Systems-----	57
Biases and Controls in Replicate Measurements-----	57
Selection of Replicate Examinees-----	58
The Analysis of Replicate Data on Body Measurements-----	59
Cycle III Systematic Replicate Procedure-----	59
Results of the Replicate Analysis-----	60
Discussion and Summary of Replicate Analysis-----	65

BODY DIMENSIONS AND PROPORTIONS, WHITE AND NEGRO CHILDREN 6-11 YEARS

Robert M. Malina, Ph.D., Peter V. V. Hamill, M.D., M.P.H., and Stanley Lemeshow, M.S.P.H.^a

This report compares the growth patterns of white and Negro children for 20 body measurements selected from a survey of U.S. children 6-11 years of age, Cycle II of the Health Examination Survey. It is the fifth in a series of reports presenting analyses and discussions of data on height, weight, and 28 other body measurements taken in Cycle II. The first two reports^{1,2} analyzed height and weight by age, sex, race, geographic region, and various socioeconomic indicators. The third report³ presented data on skinfold thickness. The fourth⁴ considered data, by age and sex, on 21 body measurements performed in Cycle II which are specifically useful to those concerned with human factors in equipment and safety design and in the manufacture of furniture and clothing.

The Health Examination Survey (HES) is conducted by the National Center for Health Statistics to collect and analyze health-related data on the American people through direct examination of selected subjects. It is a succession of separate programs, each referred to as a "cycle," and each cycle lasts from 2 to 4 years.⁵

Cycle I of HES, conducted from 1959 to 1962, obtained information on the prevalence of certain chronic diseases and the distribution of a number of anthropometric and sensory characteristics in the civilian, noninstitutionalized population of the conterminous United States aged 18-79 years. The general plan and operation of the survey and Cycle I are described in two previous reports,^{5,6}

and most of the results are published in other Series 11 reports of *Vital and Health Statistics*.

Cycle II, conducted from July 1963 to December 1965, involved selection and examination of a probability sample of noninstitutionalized children in the United States aged 6-11 years. This program succeeded in examining 96 percent of the 7,417 children selected for the sample. The examination had two emphases. The first concerned factors related to healthy growth and development as determined by a physician, a nurse, a dentist, and a psychologist; the second concerned a variety of somatic and physiologic measurements performed by specially trained technicians. The detailed plan and operation of Cycle II and the response results are described in *Vital and Health Statistics*, Series 1, Number 5.⁷ A comparable examination of data collection for Cycle III, youths aged 12-17, was completed in 1970, and the plan and operation are described in Series 1, Number 8.⁸

The present report and Series 11, Number 123⁴, together consider 27 of the 30 body measurements taken in Cycle II, leaving out only the three skinfold measurements. Although companion pieces, these two reports are very dissimilar in purpose and method of analysis. As stated in the introduction to Series 11, Number 123:

The main purpose of the numerous body measurements collected in Cycle II was to define a normal pattern of growth and development in children in the United States in the middle 1960's (and to describe some of the modifying factors). However, the opportunity to obtain data on this uniquely representative sample of U.S. children for more utilitarian

^a Associate Professor of Anthropology, University of Texas, Austin, Texas; Medical Advisor, Children and Youth Program, Division of Health Examination Statistics; and formerly Analytical Statistician, DHES, respectively.

purposes as well was not disregarded. In Cycle I (adults aged 18-79), 18 body measurements were obtained not only as medical and anthropologic correlates to the rest of the examination, but also as data for use in the consideration of anthropometric factors in equipment and safety design and manufacture (furniture, clothing, etc.)... The 21 anthropometric dimensions in this report were included in the group of body measurements partly for their descriptive value in the growth and development battery and partly for their use in "human engineering" or "human factors" work. Some of the measures have limited value in describing growth and development because they comprise multiple layers of tissue, multiple organ systems, and/or multiple loci of growth (e.g., waist and chest size, thigh clearance, all girths, seat breadth). However, many of these dimensions were selected primarily to achieve continuity with those measurements taken on adults in Cycle I of the HES...⁴

That report was descriptive and utilitarian in its purposes. Percentile distributions of each measurement by age and sex were presented as found in the total population of U.S. children irrespective of race. In contrast, the present report is biologically oriented, and the data are analyzed separately for white and Negro children. In addition to the 20 separate body measurements taken in Cycle II, which are listed below, three indexes and two derived measurements (each based on two or more of the separate measurements) are used in the present comparative analysis.

The 20 separate dimensions presented in this report by age, sex, and race are weight, stature, sitting height, buttock-knee length, popliteal height, foot length, acromion-olecranon length, elbow-wrist length, hand length, biacromial breadth, bicristal breadth, bicondylar breadth of the femur, chest breadth, chest depth, upper arm girth, lower arm girth, calf girth, chest girth, waist girth, and hip girth.

In addition to gross body size, these dimensions provide measures of upper and lower extremity lengths, body breadths across bony landmarks, and various extremity and torso girths. Extremity length measurements make it

possible to assess the relative contribution of different segments (for instance, the upper arm and forearm to total arm length) and thus to better describe body proportions. Breadth measurements are indicators of skeletal breadth; when taken at several body sites they indicate the contribution of skeletal framework to body build. Limb circumferences provide an estimate of relative muscularity and thus an insight into the body's composition. The arm, for example, is comprised of successive layers of bone, muscle, and fat. When arm circumference is corrected for the thickness of the outer layer of subcutaneous fat at the triceps site, an estimate of the lean component of the arm's composition is obtained. (See below.) Trunk circumferences are of limited value in estimating body composition except perhaps in extreme cases of undernutrition or overnutrition. They may, however, contribute to a general appraisal of physique, e.g., the ratio of chest and waist circumference. Each of the dimensions is defined and the technique of measurements described in detail in appendix II.

Two dimensions were derived from the available measurements. First, sitting height was subtracted from stature to provide an estimate of subischial length. Subischial, or leg, length enables us to partition stature into two major components: the trunk, head, and neck, which comprise sitting height, and the lower extremity length. Second, an estimate of the mid-arm muscle circumference (lean component) was obtained by correcting the upper arm circumference for the thickness of the triceps skinfold as follows: LMC (estimated muscle circumference) = upper arm circumference - π triceps skinfold. In recent years the estimated upper arm muscle circumference is often used as an anthropometric index of nutritional status.⁹⁻¹¹

In addition, three ratios were derived from the available measurements: (1) the ponderal index, $height^3/weight$,^b to provide an approximation of physique on a linearity/laterality continuum; (2) the ratio of sitting height to stature, $sitting\ height/stature \times 100$, to indicate the relative contribution of sitting height (head, neck, and trunk) to

^bIn computing this ratio height is expressed in inches and weight in pounds, which produces a different result than would the use of metric measures.

total stature; and (3) the ratio of biacromial to bicristal breadths, biacromial breadth/bicristal breadth X 100, to indicate the relative proportions of shoulder and hip width. The shoulder, hip ratio is used frequently in studies of body proportions and physique.

Estimates of the usual parameters for distributions (i.e., mean and standard deviation) are used throughout the report except where distributions deviate from normal. The percentile values indicate the general nature of the distributions. In general, with the exception of body weight and several girths, the dimensions discussed in the present report are normally distributed, or show only slight and inconsistent deviations over the age span under study. Normally distributed dimensions are primarily length and breadth measurements, which are measured between well-defined bony landmarks—bone-to-bone measurements. The distributions of body weight and of limb and trunk circumference measurements deviate from the normal over the ages studied and are skewed to the right. The positive skewness is more apparent for body weight and trunk circumferences. Of the three limb circumferences, lower arm circumference is normally distributed, while both upper arm and calf circumferences are slightly skewed to the right. The skewing of weight and girth measurements reflects the contribution of subcutaneous fat to these dimensions. Subcutaneous fat measured via skinfold thickness is positively skewed in distribution.³ Hence in this report medians of weight and girth measurements are used for comparative purposes; however, means and their standard errors are also included in the detailed tables because these statistics are widely used by others.

The six new body measurements reported here complete the presentation of percentile distributions of all 30 body measurements taken in Cycle II. Five of the 30 measures (height, weight, and three skinfolds) were examined by socioeconomic and demographic variables, in addition to percentile distributions specific for age, sex, and race.¹³ Of the 25 remaining body dimensions, seven were considered useful only for Series 11, Number 123⁴ (foot, hand, seat, and elbow-elbow breadths; thigh clearance; knee height; and buttock-popliteal length). Six are presented for the first time in this report (the three

extremity girths: calf, upper arm, and lower arm; and three bone-to-bone breadths: biacromial, bicristal, and bicondylar). Twelve measurements are examined in both reports (sitting height, popliteal height, buttock-knee length, acromion-olecranon length, elbow-wrist length, foot length, hand length, chest breadth and depth, chest girth, waist girth, and hip girth). Height and weight, of course, are included in many of these reports.

METHOD

At each of 40 locations preselected randomly throughout the United States,^c the children were brought to the centrally located mobile examination center for an examination which lasted about 2 1/2 hours. Six children were examined in the morning and six in the afternoon. They were transported to and from school or home.

When the children entered the examination center, their oral temperature was taken and a cursory screening for acute illness was made; if illness was detected in a child, he was sent home and examined at a later date. The examinees changed, into shorts, cotton sweat socks, and a light, sleeveless top and proceeded to different stages of the examination, each one following a different route. There were six different stations where examinations were conducted simultaneously, and the stations were exchanged, somewhat like musical chairs, so that by the end of 2 1/2 hours each child had had essentially the same examinations by the same examiners but in a different sequence. At three of these stations a pediatrician, a dentist, and a psychologist made examinations, and at the other three stations highly trained technicians performed a number of other examinations—chest and hand-wrist X-rays, hearing and vision tests, respiratory function tests and electrocardiography, a bicycle exercise test, a battery of body measurements, and a grip strength test.

The recording form for the battery of 30 body measurements is reproduced in appendix II, which also gives details on equipment and measuring technique. All lateral measurements were performed on the subject's right side and recorded by a trained observer. Periodic quality control

^cSee appendix I for sample design.

observation and training sessions were conducted by the supervisory medical staff and outside consultants to insure continued proficiency and to obtain replicate data for the purpose of quantifying observer error. The results are presented in detail in appendix II.

As in all the HES reports, age is basically defined as age attained at last birthday (verified from a copy of the birth certificate in 95 percent of the Cycle II examinees). The mean age of each category therefore approximates the midpoint of the whole year; for instance, the 8-year-old male group consists of a 1-year cohort whose mean age is 8.51 years, while the corresponding female sample averages 8.49 years.

"Race" was recorded as "white," "Negro," and "other races." White children comprised 85.69 percent of the total, Negro children 13.87 percent, and children of other races only 0.45 percent.^c The differential response rate by age, sex, and race is discussed in appendix I.

RESULTS

Weight and Height

Weight and height in the Cycle II sample have been discussed at length in previous reports.^{1,2} They are included here to provide a more complete picture of the anthropometry of American

^aThe classification scheme used in the 1960 census was employed here. As described in the report on the operation of HES Cycle II,⁷ this information was obtained at the initial household interview by the U.S. Bureau of the Census field worker. Its accuracy was checked at the subsequent home visit by the HES representative and again at the examination in the trailer. A final record check by birth certificate turned up only seven inconsistencies, and these were mostly pertaining to the category "other races." Hence the possible extent of misclassification of the variable race is so minimal that it could have no effect on the data analyzed in this report. However, when comparing the HES findings to those on other variously defined racial groupings in the world, the degrees of genetic admixture, as discussed first by Hershkowitz in 1928¹² and later by Glass and Li,¹³ by Roberts,^{14,15} and by Reed¹⁶ should be taken into consideration.

^cChildren of other races were included in Series 11, Number 123, when all the data were analyzed independently of race but are omitted, because they are so few, when a white-Negro dichotomy is used, as in the present report.




Figure 1. Median weight of white and Negro children by sex and age.

Negro and white children during middle childhood. Both weight and height are measures of gross body size and, as expected, increase linearly with age between 6 and 12 years (figures 1 and 2, tables 1 and 2).

Median body weights for white and Negro boys differ only slightly from 6 through 8 years, but by 9, 10, and 11 years differences in median body weights have become greater and are consistently larger in white boys. White girls have slightly greater median body weights than Negro girls at 6, 7, 8, and 9 years of age, but at 10 and 11 years the median weights for Negro girls are slightly greater.

Racial differences in average stature are negligible for males, mean statures for Negro boys being slightly greater at 6, 7, and 8 years of age and those for white boys slightly greater at 9 and 10 years. On the other hand, Negro girls have consistently greater mean statures than white girls in all age groups except the 8-year-old group, so that at ages 10 and 11 they are both taller and heavier than white girls.




Figure 2. Mean height of white and Negro children by sex and age.

Ponderal Index

Viewing the relationship of height and weight (height/ $\sqrt[3]{\text{weight}}$) as expressed in the ponderal index (figure 3, table 3) indicates consistently higher indexes for Negro children of both sexes than for whites. High ponderal indexes suggest linearity of physique, while low indexes suggest laterality. Thus the foregoing ratio of height and weight indicates a linear physique in Negro children of both sexes. The difference between mean ponderal indexes of Negro and white boys at each age is rather consistent from 6 through 11 years. For girls this difference is similarly consistent from 6 through 9 years, but at 10 and 11 years the in-




Figure 3. Mean ponderal index of white and Negro children by sex and age.

dexes for Negro and white girls approach each other. Note, however, that it is at these two ages that Negro girls are heavier than white girls. Nevertheless, the mean ponderal indexes for Negro girls at every age except 10 are higher than those for white girls.

Components of Stature: Sitting Height and Subischial Length

Partitioning standing height into sitting height (figure 4, table 4) and subischial length, or stature minus sitting height (figure 5, table 5), illustrates the well-established racial difference in the components of stature: Negroes are longer legged and shorter trunked; conversely, whites are shorter legged and longer trunked. This is true for both sexes, the difference between racial groups being generally consistent over the age range studied.

Sitting Height/Stature Ratio

The ratio of sitting height to stature (sitting height/standing height X 100) is consistently higher in white children of both sexes (figure 6, table 6). This ratio indicates that a significantly greater percentage of standing height is contributed by the sitting height (head, neck, and trunk), and less by subischial or leg length in white than in Negro children, while of course the opposite is true for Negro children. The ratio of sitting height to stature decreases in a parallel manner with increasing age from 6 to 11 years, indicating a greater contribution of the lower extremities to stature with advancing age during middle childhood in children of both racial groups.




Figure 4. Mean sitting height of white and Negro children by sex and age.




Figure 5. Mean subischial length of white and Negro children by sex and age.




Figure 6. Mean sitting height/stature ratio of white and Negro children by sex and age.

Partitioning the Lower Extremity

Although traditional bone-to-bone landmark measurements of the lower extremity were not taken, two measurements used primarily for "human engineering" or "human factors" research purposes provide an approximation of racial differences in the components of lower extremity length. Buttock-knee length, though affected by fatty tissue deposits over the buttocks, provides a rough approximation of thigh or upper leg length; popliteal height provides an approximation of lower leg length. *Buttock-knee length* is consistently longer in Negro girls than in white girls, the mean differences being smaller at 6, 7, and 8 years than at 9, 10, and 11 (figure 7, table 7). The buttock-knee length of boys shows no consistent pattern of differences over the age range studied. Although Negro boys have longer buttock-knee length measurements at 6, 7, and 8

years, differences between Negro and white boys are negligible at 9, 10, and 11 years. *Popliteal height* is consistently greater in Negro children of both sexes over the age span, with the difference between means at each age group being very consistent (figure 8, table 8).

The data for buttock-knee length and popliteal height, though both are only approximate measurements, suggest that the greater length of the lower extremity in Negro children is due especially to a longer lower leg, the differences in upper leg length being minor and inconsistent. This generalization is in agreement with recent observations on Negro and white children utilizing traditional bony landmarks, which give more precise measurements.¹⁷

Foot length is likewise consistently greater in Negro children of both sexes over the age range studied (figure 9, table 9). Again, the difference between means at each age is very consistent.




Figure 7. Mean buttock knee length of white and Negro children by sex and age.




Figure 8. Mean popliteal height of white and Negro children by sex and age.

Partitioning the Upper Extremity.

A generally similar racial pattern to that observed for the lower extremity is evident for the upper extremity. *Acromion-olecranon length* (upper arm length) is consistently greater in Negro girls from 6 to 11 years (figure 10, table 10). In boys, however, the pattern of differences is not consistent. Negro boys have longer upper arms at 6, 7, 8, and 11 years of age, while white boys have longer upper arms at 10 years.




Figure 9. Mean foot length of white and Negro children by sex and age.




Figure 10. Mean acromion-olecranon length of white and Negro children by sex and age.




Figure 11. Mean elbow wrist length of white and Negro children by sex and age.

Mean acromion-olecranon lengths in Negro and white boys are identical at 9 years of age. It should be noted that a similar pattern of mean differences between Negro and white boys and girls is apparent for buttock-knee length, *Elbow-wrist length* (lower arm or forearm length) is consistently greater in Negro children of both sexes from 6 through 11 years (figure 11, table 11), and the difference between means is rather consistent at each age group.

Hand length is also consistently longer in Negro children of both sexes (figure 12, table 12). The difference between means at each age is very consistent.

Negro children of both sexes have longer upper extremities than do whites. This difference is largely due to the longer forearms and hands of Negro children, just as their longer lower extremities are especially due to longer lower legs. (This is considered in more detail in the discussion.)




Figure 12. Mean hand length of white and Negro children by sex and age.

Biacromial Breadth

Negro boys have slightly wider measurements of bony breadth of the shoulders at each age except 9 years (figure 13, table 13). Biacromial breadths of Negro girls are only slightly higher than those of white girls at 6 and 7 years (and are the same at age 8). At 9, 10, and 11 years, however, the difference between Negro and white girls is greater, perhaps only reflecting the larger body size of the Negro girls at these ages (see figures 1 and 2).




Figure 13. Mean biacromial breadth of white and Negro children by sex and age.

Bicristal Breadth

The bony breadth across the iliac crests is consistently larger in whites of both sexes over the age span studied (figure 14, table 14). Also, the difference between means at each age group is rather consistent from 6 through 11 years in males and 6 through 9 years in females. At 10 and 11 years the difference between bicristal breadth means in Negro and white females becomes slightly smaller than at the younger ages, although average bicristal breadth is still larger in white females. Thus, even though Negro girls are generally larger in overall body size than white girls, especially at 10 and 11 years of age, white girls have wider bicristal breadths over the entire age span studied.




Figure 14. Mean bicristal breadth of white and Negro children, by sex and age.

Biacromial Breadth/Bicristal Breadth Relationship

The ratio of shoulder to hip breadth (biacromial breadth/bicristal breadth X 100) is consistently higher in Negro children of both sexes over the age range studied (figure 15, table 15). The higher ratio indicates that Negro children have more slender pelvis relative to their shoulders than white children. Conversely, the lower ratio indicates that white children have broader hips relative to their shoulders than Negro children.

The magnitude of the biacromial/bicristal ratio decreases with age in an almost parallel manner in Negro and white girls, indicating a greater widening of the breadth across the iliac crests relative to the breadth of the shoulders. The ratio in white boys is almost constant between 6 and 11 years. In Negro boys the ratio at 6, 7, 8, and 11 years of age is almost constant, but at 9 and 10 years of age it shows a sharp increase. Whether this fluctuation in the ratio is only due to sampling variation in the smaller Negro sample is not clear.




Figure 15. Mean biacromial breadth/bicristal breadth ratio of white and Negro children by sex and age.

Chest Dimensions

Chest breadth is, on the average, slightly but consistently greater in white children of both sexes from 6 through 11 years (figure 16, table 16). *Chest depth*, on the other hand, shows no consistent pattern of racial differences over the age span studied (figure 17, table 17). Interestingly, Negro children of both sexes have slightly higher mean values at 6 years of age; thereafter mean values for the chest depth measurement are generally slightly larger in white children.




Figure 16. Mean chest breadth of white and Negro children by sex and age.




Figure 17. Mean chest depth of white and Negro children by sex and age.

Bicondylar Breadth of the Femur

This measurement, across the condyles of the femur shows only negligible differences between Negro and white children of both sexes from 6 through 11 years (table 18).

Limb Girths

Median values for *upper arm girth* are consistently larger in white boys and girls from 6 through 11 years (figure 18, table 19), and the difference between medians at each age is rather consistent. The significance of differences between Negro and white children for this girth measurement will be considered in the discussion. In *lower arm girth*, medians for white boys are slightly though consistently larger than those for Negro boys from 6 through 11 years of age. White girls have larger lower arm girth medians at 6, 7, and 8 years of age, but from 9 through 11 years




Figure 18. Median upper arm girth of white and Negro children by sex and age.




Figure 19. Median lower arm girth of white and Negro children by sex and age.

no differences between Negro and white girls are apparent in this measurement (figure 19, table

Calf girth shows a pattern similar to that for upper arm girth. At all ages white boys and girls have larger medians for calf girth than their Negro counterparts (figure 20, table 21). The differences between medians for Negro and white boys are relatively small at 6, 7, and 8 years but greater at 9, 10, and 11 years. Differences between the medians for Negro and white girls are generally consistent over the age span under study.

Estimated Mid-Arm Muscle Circumference

The upper arm girth corrected for the thickness of the triceps skinfold¹ provides an estimate of the mid-arm muscle or lean circumference of the arm. As noted earlier, this estimated measurement is widely used in public health surveys. Although the estimated circumference is generally indicated as being a muscle circumference, it should be noted that bone tissue, the humerus, is also included. The significance of the contribution of bone to this measurement, which is important in racial comparisons, is considered at length in the discussion.

¹Comparison of skinfold thicknesses in the Negro and white sample discussed here is treated in detail in another report.³




Figure 20. Median calf girth of white and Negro children by sex and age.

Differences in estimated muscle or lean circumference of the arm between Negro and white children of both sexes are small; nevertheless, they are consistent. Negro boys and girls have, on the average, consistently larger estimated muscle or lean circumference of the arm than whites from 6 through 11 years (figure 21, table 22). Thus the larger upper arm girth of white children noted earlier appears to be due to more subcutaneous fat at the triceps site in white children from 6 through 11 years.

Torso Girths

White children of both sexes have consistently larger median values for *chest girth* than Negro children from 6 through 11 years of age (figure 22; table 23). With the exception of the




Figure 21. Mean estimated mid-arm muscle circumference of white and Negro children by sex and age.




Figure 22. Median chest girth of white and Negro children by sex and age.




Figure 23. Median waist girth of white and Negro children by sex and age.




Figure 24. Median hip girth of white and Negro children by sex and age.

9-year-old sample, white boys also have larger *waist girth* measurements than Negro boys (figure 23, table 24). Differences in median values for waist girth between Negro and white girls, however, are not consistently apparent over the ages studied. *Hip girth* shows a pattern similar to that for chest girth: white boys and girls have consistently larger medians for this measurement from 6 through 11 years of age (figure 24, table 25). Median values for both chest and hip girths in girls show a pattern of decreasing racial differences with increasing age; so at 10 and 11 years of age differences between medians are small. These observations are probably related to the greater overall body size of Negro girls at these ages.

DISCUSSION

Patterns of growth as shown in a series of anthropometric dimensions and indexes were analyzed in a large, representative sample of U.S. Negro and white children 6 through 11 years of age. The data are cross-sectional (i.e., different children are represented at each age level) and are thus affected by the limitations of cross-sectional studies. In addition, emphasis is placed on comparisons of mean or median values for Negro and white children, despite considerable overlap between races in each age and sex group. Nevertheless, the present data provide updated anthropometric information which are reliable estimates of American Negro and white children of both sexes during middle childhood.

The Cycle II data for Negro and white children generally agree with existing data for contemporary as well as for earlier samples. It is, however, difficult to make precise comparisons with most other studies on account of sampling and other methodological differences. One difference is that the HES data are grouped by chronological age, and so the average age is the midpoint of the year. For example, 6-year-old children range from 6.00 to 6.99 years, with the mean at approximately 6.5 years. In two recent studies of Negro and white children which include the age period under study,^{17,18} children are grouped into yearly categories, with the whole year as the midpoint. In those studies 6-year-olds are classified as children aged 5.50 to 6.49 years, with the mean

at approximately 6.0 years. Each of the two procedures refers to the 6-year-old age group, but the definitions are different.

Other technical difficulties in making comparisons relate to definition of sites and techniques of measurement. For example, upper extremity measurements used in the present report were made between specific landmarks (e.g., acromion to olecranon). Krogman's measurements of upper extremity segments were derived by subtraction of measurements made at specific landmarks above floor level; for instance, acromial height minus radial height gives upper arm length.¹⁷

It should be further noted that buttock-knee length and popliteal height, two of the measurements used in this report to partition the segments of the lower extremity, are essentially human engineering measurements. Because these measurements are made from surface to surface with light contact, they are confounded to some extent by variation in soft tissues. Therefore buttock-knee length and popliteal height provide only an approximation of actual segmental lengths and are not as precise as measurements using traditional bony landmarks.

Although there are some overall body size differences between Negro and white children during middle childhood, especially in girls, the major anthropometric differences between American Negro and white children are essentially in the proportions of the trunk and limbs. In comparison with white children, Negro children have shorter trunks, more slender hips and chests, longer lower extremities (especially a longer lower leg), and longer upper extremities (especially a longer forearm and hand). Conversely, in comparison with Negro children, white children have longer, thicker trunks, wider hips, shorter lower extremities, and shorter upper extremities. These dimensional and proportional differences between American Negro and white children were noted as far back as 1929 by Todd,¹⁹ and are well documented in other studies of children and adults.^{17,20-24} Further, such proportional differences have been reported to occur prenatally during the first trimester.²⁵⁻²⁷ This implies that genetic factors affect skeletal dimensions: one's racial background predisposes the skeleton to certain proportions.

In an attempt to assess racial differences in the limb segments comprising the upper and lower extremities, the normal deviates for these segments were analyzed. For each age and sex group, the normal deviate

$$z = \frac{\bar{X}_W - \bar{X}_N}{\sqrt{S_{\bar{X}_W}^2 + S_{\bar{X}_N}^2}}$$

was computed. Instead of its usual use in statements of probability levels, this measure is used here as an indicator of relative magnitudes. The larger the deviate, the greater the difference between whites and Negroes, since by dividing by the standard error of the difference, the problem of difference in relative magnitude is eliminated. Results of this analysis are presented in table 26.

A distinct pattern suggesting a proximal-distal gradient emerges for the three segments of the upper extremity. Differences between Negro and white children are least marked for the upper arm segment (acromion-olecranon length) and most marked for hand length, with the values for the forearm segment (elbow-wrist length) being intermediate. This pattern of racial differences for the upper limb segments is apparent in all age groups except the 8-year-old boys and girls, in whom the deviation from a proximal-distal pattern was very slight.

In contrast to the proximal-distal gradient of racial differences in the upper limb segments, there is no consistently apparent pattern for the segments of the lower extremity in boys. The pattern for girls, however, is similar to that noted for the upper extremity (table 26). Differences between Negro and white children of both sexes are consistently least marked for thigh length (buttock-knee length). The pattern of differences for the lower leg and foot is not clear in boys. In all but one age group of girls (the 9-year-old group), however, racial differences are most marked for foot length, with the value for lower leg length (popliteal height) being intermediate.

The preceding observations on the lower extremity are in part a function of the measurements used and in part a function of the mechanics of the lower extremity. It should be carefully noted that two of the measurements of lower ex-

limb segments are only indirect measures of segment length. Buttock-knee length, for example, is affected to some extent by the deposition of subcutaneous fat on the buttocks and also includes the thickness of the patella or knee cap. (See Appendix II for discussion of measurement methods.) Hence it is not a true measure of thigh or femoral length. Similarly, popliteal height is an indirect measure of lower leg length. Because it is measured as the distance from the footrest to the popliteal fossa, it includes not only the tibia, the major bone of the lower leg, but also the bones of the ankle, especially the talus and calcaneus. Thus this measurement is not an accurate indicator of lower leg length per se.

In addition to measurement technique, the functional mechanics of the upper and lower extremities limit direct comparisons of their respective segment lengths. The foot, for example, is adapted as a weight-bearing unit in the upright posture, while the hand is adapted as primarily a prehensile organ. Thus the foot joins the remainder of the lower extremity at a 90-degree angle, while the hand is a continuous extension of the forearm.

Racial differences in girths are affected to a large extent by variation in amount of subcutaneous fat. This is especially apparent for upper arm girth. When corrected for the thickness of the triceps skinfold, the difference between Negro and white children in upper arm girth is seen to be due to an excess of triceps fat in white children. Comparing figures 18 (upper arm girth) and 21 (estimated muscle circumference of the arm), one can note a reversal in the positions of Negro and white children. White children have consistently larger arm girths, but Negro children have consistently larger estimated muscle or lean circumferences of the arm. Similar observations have been made by Malina,¹⁸ for a sample of Negro and white Philadelphia elementary school children studied longitudinally over a 1-year period.

As indicated earlier, correcting the upper arm girth for the thickness of the triceps skinfold results in what is generally termed the estimated mid-arm muscle circumference. The significance of the contribution of bone tissue to this estimated circumference is important in making racial comparisons. Data comparing the

breadth of the humerus (determined from a radiograph) in Negro and white children are lacking. Garn²⁸ and Smith and Risek,²⁹ however, reported little difference in periosteal diameters of the second metacarpal, but did note greater cortical bone thickness among adult Negroes of both sexes than among the corresponding white groups. This seems to suggest that adult Negroes have more cortical bone for the same periosteal diameter of bone. However, it is difficult to make inferences from measurements of second metacarpal width to the humerus; results of a study comparing radiographic bone breadth measurements of the second metacarpal, second metatarsal, humerus, and tibia indicated little predictive relationship from one bone to the other in a sample of white children aged 6 to 16.^{30,31}

It is difficult to make inferences from one skeletal area to another. The available data on bone width measurements of the second metacarpal indicate little consistent racial difference, if any, in total width at the midshaft level. If the same is true for the humerus, the corrected arm circumference data would then indicate slightly but consistently greater muscularity in the upper arms of Negro compared with white children. Data from samples of Olympic athletes indicate a generally similar trend. Using radiographic techniques, Tanner²¹ found that Negro athletes had slightly more muscle and bone tissue and less fat in the upper arm and thigh compared to white athletes; although in the calf they also had less fat and more bone tissue, they had less muscle than whites. Thus this sample of Negro athletes had substantially smaller calf muscles relative to muscular development in the arm and thigh than did white athletes.

Special consideration should be given to the comparison of Negro and white girls. On the average, Negro girls are taller than white girls from 6 through 11 years of age and are heavier at 10 and 11 years of age. Cycle II data also show a reduction in the difference between median girths of Negro and white girls at ages 10 and 11. It would be interesting to find out if there is a maturational difference between Negro and white girls at these ages. Malina,³² for example, noted Negro girls advanced over white girls in skeletal maturity from 9 through 12 years of age. Todd³³ in 1931 reported somewhat similar observations,

noting greater variability between Negro and white girls than between boys. Todd's small sample of Negro girls ($n=72$) were frequently skeletally advanced compared to his larger sample of white girls.

The report assessing skeletal maturity of the Cycle II sample of children 6-11 years of age, which is currently in preparation, will contribute substantially to this question.

SUMMARY

Patterns of growth in 20 measured anthropometric dimensions, two derived dimensions, and three ratios are reported and discussed for U.S. white and Negro children 6 through 11 years of age. The 20 dimensions reported by age, sex, and race are weight, stature, sitting height, buttock-knee length, popliteal height, foot length, acromion-olecranon length, elbow-wrist length, hand length, biacromial breadth, bicristal breadth, bicondylar breadth of the femur, chest breadth, chest depth, upper arm girth, lower arm girth, calf girth, chest girth, waist girth, and hip girth. The two dimensions derived from available measurements are estimated leg (subischial) length and estimated mid-arm muscle circumference. The three ratios are the ponderal index, the ratio of sitting height to stature, and the ratio of biacromial (shoulder) to bicristal (hip) width.

These national estimates are based on cross-sectional data, which limit the analyses to attained size rather than velocities of growth. Emphasis is placed upon comparisons of age- and sex-specific means and medians for Negro and white children.

All dimensions reported here increase almost linearly with age from 6 through 11 years in Negro and white children, both male and female. Although there are some overall body size differences (height and weight) between Negro and

white children during middle childhood, especially in girls, the major anthropometric differences between Negro and white children are differences in the proportions of the trunk and limbs. Negro children, on the average, have shorter trunks, more slender hips and chests, longer lower extremities (especially longer lower legs), and longer upper extremities (especially longer forearms and hands) than white children. White children, conversely, tend to have longer, thicker trunks, wider hips, and shorter lower and upper extremities than Negro children.

Limb (arm, forearm, and calf) and torso (chest, waist, and hip) girths are generally larger in white children over the age range studied. There are, nevertheless, exceptions, and the differences between Negro and white children are not apparent over all ages from 6 through 11 years. For instance, Negro girls are taller and heavier than white girls, especially at 10 and 11 years, and this is reflected in a reduction of median girth differences between Negro and white girls, at these ages. Much of the racial difference in girths is explained by differences in the amount of subcutaneous fat. This is especially apparent for arm girth; when the arm circumference is corrected for the thickness of the triceps skinfold, it is obvious that the girth differences between Negro and white children are due essentially to a larger amount of triceps fat in white children. In fact, when corrected for the thickness of the triceps skinfold, estimated mid-arm muscle (including bone) circumference is consistently larger in Negro children.

The conclusions derived from these HES data generally agree with most other anthropometric comparisons of American white and Negro children; the statistics provide current, comprehensive, and reliable national estimates of body dimensions for American Negro and white children of both sexes during middle childhood.

REFERENCES

- ¹National Center for Health Statistics: Height and weight of children, United States. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 11-No. 104. Public Health Service. Washington. U.S. Government Printing Office, Sept. 1970.
- ²National Center for Health Statistics. Height and weight of children: socioeconomic status, United States. *Vital and Health Statistics*, Series 11-No. 119. DHEW Pub. No. (HSM) 73-1601. Washington. U.S. Government Printing Office, Oct. 1972.
- ³National Center for Health Statistics: Skinfold thickness of children 6-11 years, United States. *Vital and Health Statistics*. Series 11-No. 120. DHEW Pub. No. (HSM) 73-1602. Washington. U.S. Government Printing Office, Oct. 1972.
- ⁴National Center for Health Statistics: Selected body measurements of children 6-11 years, United States. *Vital and Health Statistics*. Series 11-No. 123. DHEW Pub. No. (HSM) 73-1605. Washington. U.S. Government Printing Office, Jan. 1973.
- ⁵National Center for Health Statistics: Plan and initial program of the Health Examination Survey. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 4. Public Health Service. Washington. U.S. Government Printing Office, July 1965.
- ⁶National Center for Health Statistics: Cycle I of the Health Examination Survey, sample and response, United States, 1960-1962. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 11-No. 1. Public Health Service. Washington. U.S. Government Printing Office, Apr. 1964.
- ⁷National Center for Health Statistics: Plan, operation, and response results of a program of children's examinations. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 5. Public Health Service. Washington. U.S. Government Printing Office, Oct. 1967.
- ⁸National Center for Health Statistics: Plan and operation of a Health Examination Survey of U.S. youths 12-17 years of age. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 8. Public Health Service. Washington. U.S. Government Printing Office, Sept. 1969.
- ⁹Jelliffe, D. B.: The assessment of the nutritional status of the community. *WHO Monogr. Ser.* No. 53, 1966.
- ¹⁰Jelliffe, D. B.: The arm circumference as a public health index of protein-caloric malnutrition of early childhood, in E. F. Jelliffe and D. B. Jelliffe, eds., *J. Trop. Pediatr.* Monograph No. 8. 15:177-260, 1969.
- ¹¹Robinow, M.: Field measurement of growth and development, in N. S. Scrimshaw and J. E. Gordon, eds., *Malnutrition, Learning and Behavior*. Cambridge, Mass. The M.I.T. Press, 1968. pp. 409-424.
- ¹²Herskowitz, M. J.: *The American Negro, a Study in Racial Crossing*. New York. Alfred A. Knopf, 1928.
- ¹³Glass, B., and Li, C. C.: The dynamics of racial intermixture - an analysis based on the American Negro. *Am. J. Hum. Genet.* 5:1-20, 1953.
- ¹⁴Roberts, D. F.: The dynamics of racial intermixture in the American Negro - some anthropological considerations. *Am. J. Hum. Genet.* 7:361-367, 1955.
- ¹⁵Roberts, D. F., and Hiorns, R. W.: The dynamics of racial intermixture. *Am. J. Hum. Genet.* 14:261-277, 1962.
- ¹⁶Reed, T. E.: Caucasian genes in American Negroes. *Science* 165(3895):762-768, Aug. 22, 1969.
- ¹⁷Krogman, W. M.: Growth of head, face, trunk, and limbs in Philadelphia white and Negro children of elementary and high school age. *Monogr. Soc. Res. Child Dev.* 35(3), Serial No. 136, 1970.
- ¹⁸Malina, R. M.: Growth, Maturation, and Performance of Philadelphia Negro and White Elementary School Children. Unpublished doctoral dissertation, University of Pennsylvania, Philadelphia, 1968.
- ¹⁹Todd, T. W.: Entrenched Negro physical features. *Hum. Biol.* 1:57-69, 1929.
- ²⁰Metheny, E.: Some differences in bodily proportions between American Negro and white male college students as related to athletic performance. *Res. Q.* 10:41-53, 1939.
- ²¹Tanner, J. M.: *The Physique of the Olympic Athlete*. London. George Allen and Unwin, 1964.
- ²²Malina, R. M.: Growth and physical performance of American Negro and white children. *Clin. Pediatr.* 8(8):476-483, Aug. 1969.
- ²³Malina, R. M.: Biological substrata, in K. S. Miller and R. M. Dreger, eds., *Comparative Studies of Blacks and Whites in the United States*. New York. Seminar Press, 1973. pp. 53-123.
- ²⁴Long, L., and Churchill, E.: Anthropometry of USAF Male Basic Trainees - Contrasts of Several Subgroups. Paper prepared for a meeting of the American Association of Physical Anthropologists, 1968. Mimeographed.
- ²⁵Schultz, A. H.: Fetal growth in man. *Am. J. Phys. Anthropol.* 6:389-399, 1923.
- ²⁶Schultz, A. H.: Fetal growth of man and other primates. *Q. Rev. Biol.* 1:465-521, 1926.
- ²⁷Schultz, A. H.: Proportions, variability and asymmetries of the long bones of the limbs and the clavicles in man and apes. *Hum. Biol.* 9:281-328, 1937.
- ²⁸Garn, S. M.: The course of bone gain and the phases of bone loss. *Orthop. Clin. North Am.* 3:503-520, 1972.
- ²⁹Smith, R. W., Jr., and Risek, J.: Epidemiologic studies of osteoporosis in women of Puerto Rico and Southeastern Michigan with special reference to age, race, national origin and to other related or associated findings. *Clin. Orthop.* 45:31-48, 1966.

³⁰ Johnston, F. L., and Malina, R. M.: Correlations of mid-shaft breadths and compact bone thickness among bones of the upper and lower extremities of children aged 6 to 16 years. *Am. J. Phys. Anthropol.* n.s. 32:32-327, 1970.

³¹ Malina, R. M.: Quantification of fat, muscle and bone in man. *Clin. Orthop.* 65:9-38, 1969.

³² Malina, R. M.: Skeletal maturation studied longitudinally over one year in American whites and Negroes six through thirteen years of age. *Hum. Biol.* 42(3):377-390, Sept. 1970.

³³ Todd, T. W.: Differential skeletal maturation in relation to sex, race, variability and disease. *Child Dev.* 2:49-65, 1931.

³⁴ National Center for Health Statistics. Replication. An approach to the analysis of data from complex surveys. *Vital and Health Statistics*, PHS Pub. No. 1000-Series 2-No. 14. Public Health Service, Washington, U.S. Government Printing Office, Apr. 1966.

³⁵ Hertzberg, H. T. E.: The Conference on Standardization of Anthropometric Techniques and Terminology. *Am. J. Phys. Anthropol.* 28(1):1-16, Jan. 1968.

³⁶ National Center for Health Statistics: Quality control in a National Health Examination Survey. *Vital and Health Statistics*, Series 2-No. 44. DHEW Pub. No. (HSM) 72-1023. Washington, U.S. Government Printing Office, Feb. 1972.

LIST OF DETAILED TABLES

	Page
Table 1. Weight of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	22
2. Height of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	23
3. Ponderal index of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	24
4. Sitting height of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	25
5. Subischial length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	26
6. Sitting height/stature ratio (times 100) of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	27
7. Buttock-knee length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	28
8. Popliteal height of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	29
9. Foot length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	30
10. Acromion-olecranon length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	31
11. Elbow-wrist length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	32
12. Hand length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	33
13. Biacromial breadth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	34
14. Bicristal breadth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	35
15. Biacromial breadth/bicristal breadth ratio (times 100) of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	36
16. Chest breadth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	37

LIST OF DETAILED TABLES—Con.

	Page
Table 17. Chest depth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	38
18. Bicondylar breadth of the femur of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	39
19. Upper arm girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	40
20. Lower arm girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	41
21. Calf girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	42
22. Estimated mid-arm muscle circumference of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	43
23. Chest girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	44
24. Waist girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	45
25. Hip girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65-----	46
26. Normal deviates of differences between measurements of various components of the extremities of white and Negro children and ranking of normal deviates within sex and age groups-----	47

Table 1. Weight of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	s_x	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
<u>Boys</u>												
6 years-----	489	1,787	22.0	3.39	0.18	17.4	18.2	19.9	21.7	23.7	25.9	27.9
7 years-----	551	1,781	24.8	4.11	0.21	19.4	20.5	22.2	24.1	26.7	29.8	31.7
8 years-----	537	1,739	27.8	5.03	0.25	21.3	22.5	24.5	27.1	29.9	34.1	36.7
9 years-----	525	1,730	31.4	6.91	0.47	23.5	24.6	26.9	29.9	34.0	38.7	44.0
10 years-----	509	1,692	33.9	6.63	0.30	25.5	26.8	29.6	32.7	36.7	42.3	45.7
11 years-----	542	1,662	38.6	8.32	0.40	28.8	30.1	33.2	36.7	42.3	49.1	53.5
<u>Girls</u>												
6 years-----	461	1,722	21.6	3.79	0.25	16.4	17.5	19.3	21.1	23.3	25.9	28.9
7 years-----	512	1,716	24.3	4.17	0.20	18.7	19.5	21.4	23.5	26.5	29.9	31.6
8 years-----	498	1,674	27.6	5.21	0.26	20.8	21.9	23.9	26.8	30.1	34.4	37.9
9 years-----	494	1,663	31.4	6.84	0.42	22.8	24.3	26.6	29.9	34.6	41.8	45.7
10 years-----	505	1,632	35.0	7.97	0.44	24.8	26.2	29.2	34.1	39.4	45.4	49.3
11 years-----	477	1,605	39.8	8.75	0.36	28.6	30.0	33.5	38.2	44.9	51.8	57.5
NEGRO												
<u>Boys</u>												
6 years-----	84	289	21.8	3.21	0.37	17.5	18.1	19.5	21.3	23.4	26.4	28.2
7 years-----	79	286	24.0	2.89	0.32	19.4	20.1	21.9	24.0	26.2	28.1	28.8
8 years-----	79	279	27.5	3.67	0.42	22.4	22.9	24.9	27.2	29.4	31.9	34.4
9 years-----	74	269	29.4	5.57	0.77	21.8	22.9	25.7	28.5	32.2	37.3	39.7
10 years-----	65	264	32.4	5.36	0.72	25.4	26.4	28.7	31.6	35.4	40.1	42.2
11 years-----	83	255	36.8	6.29	0.50	27.6	29.5	32.4	36.1	39.6	46.2	48.7
<u>Girls</u>												
6 years-----	72	281	21.1	2.95	0.36	15.9	17.8	19.0	21.2	22.7	24.8	26.3
7 years-----	93	284	23.7	3.97	0.47	19.2	19.7	21.1	22.8	24.6	28.4	30.3
8 years-----	113	281	27.0	6.05	0.37	19.3	20.4	22.4	26.1	29.7	34.5	38.3
9 years-----	84	265	31.2	6.74	0.62	23.0	24.6	26.5	29.4	34.6	42.0	44.8
10 years-----	77	266	35.7	9.02	0.89	25.2	26.3	29.0	34.4	39.5	48.1	53.4
11 years-----	84	253	41.1	11.51	1.45	27.4	28.6	33.3	38.4	46.0	59.3	63.7

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; s_x = standard error of the mean.

Table 2. Height of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
In centimeters												
Boys												
6 years-----	489	1,787	118.5	5.15	0.30	110.4	111.7	115.0	118.4	121.8	125.7	127.8
7 years-----	551	1,781	124.5	5.52	0.38	115.5	117.7	120.8	124.3	127.8	131.8	134.4
8 years-----	537	1,739	129.8	5.70	0.29	120.0	123.1	126.2	129.7	133.6	137.1	138.8
9 years-----	525	1,730	135.5	6.77	0.50	124.5	126.9	131.6	135.7	140.2	143.4	145.4
10 years-----	509	1,692	140.3	6.62	0.37	129.4	131.6	136.4	140.6	144.6	148.5	151.3
11 years-----	542	1,662	145.7	6.69	0.30	134.6	137.3	141.1	145.8	150.5	154.2	158.9
Girls												
6 years-----	461	1,722	117.7	5.47	0.32	108.3	110.4	114.4	117.6	121.5	124.8	126.4
7 years-----	512	1,716	123.4	5.86	0.17	113.5	116.2	119.6	123.5	127.2	130.5	132.7
8 years-----	498	1,674	129.4	6.19	0.39	119.2	121.4	125.6	129.7	133.5	137.1	138.9
9 years-----	494	1,663	135.1	6.72	0.36	124.2	127.1	130.6	135.3	139.6	144.4	145.9
10 years-----	505	1,632	140.8	7.00	0.34	129.6	132.1	135.7	140.7	145.6	149.7	152.7
11 years-----	477	1,605	147.3	7.89	0.27	135.1	138.7	142.7	147.3	152.5	157.4	159.4
NEGRO												
Boys												
6 years-----	84	289	119.1	5.11	0.72	111.3	112.7	115.4	118.6	122.8	125.5	129.5
7 years-----	79	286	125.2	5.50	0.59	116.3	118.4	121.3	124.8	129.9	131.9	134.5
8 years-----	79	279	131.3	5.33	0.57	122.7	125.0	127.7	130.8	134.8	139.4	140.7
9 years-----	74	269	135.0	6.46	0.67	125.3	127.1	130.5	135.0	139.8	143.5	144.7
10 years-----	65	264	139.6	7.92	0.97	127.8	129.7	133.4	140.6	144.5	148.4	151.6
11 years-----	83	255	145.7	8.08	0.53	134.5	136.1	141.6	146.0	149.8	154.5	159.2
Girls												
6 years-----	72	281	118.5	5.75	0.86	106.5	111.7	114.6	118.5	122.5	126.5	127.5
7 years-----	93	284	124.6	5.55	0.59	115.7	117.6	120.3	125.1	128.2	131.6	132.7
8 years-----	113	281	129.4	6.69	0.52	118.7	120.8	124.5	129.3	132.8	138.8	142.4
9 years-----	84	265	137.5	7.80	0.90	125.8	127.4	132.8	136.4	142.6	149.8	150.8
10 years-----	77	266	141.8	9.25	0.65	129.2	131.6	138.3	142.1	146.4	155.0	155.9
11 years-----	84	253	149.2	7.42	0.69	136.2	140.3	144.5	148.4	154.8	160.4	161.7

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 3. Ponderal index of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
6 years-----	489	1,787	12.8	0.41	0.02	12.0	12.1	12.4	12.7	13.3	13.7	13.8
7 years-----	551	1,781	12.9	0.44	0.02	12.0	12.1	12.4	12.9	13.5	13.8	13.9
8 years-----	537	1,739	13.0	0.48	0.02	12.0	12.2	12.6	13.1	13.6	13.8	13.9
9 years-----	525	1,730	13.1	0.60	0.05	11.9	12.1	12.6	13.2	13.6	13.9	13.9
10 years-----	509	1,692	13.2	0.53	0.02	12.1	12.2	12.8	13.3	13.7	13.9	14.0
11 years-----	542	1,662	13.1	0.60	0.03	12.0	12.2	12.7	13.3	13.6	13.9	14.0
Girls												
6 years-----	461	1,722	12.8	0.45	0.02	12.0	12.1	12.4	12.8	13.3	13.7	13.9
7 years-----	512	1,716	12.9	0.50	0.03	12.0	12.1	12.4	12.9	13.5	13.8	13.9
8 years-----	498	1,674	13.0	0.55	0.03	12.0	12.1	12.5	13.1	13.6	13.8	13.9
9 years-----	494	1,663	13.0	0.58	0.03	11.9	12.1	12.5	13.2	13.6	13.9	14.0
10 years-----	505	1,632	13.1	0.63	0.03	11.9	12.2	12.6	13.2	13.6	13.9	14.0
11 years-----	477	1,605	13.1	0.68	0.03	11.9	12.1	12.6	13.2	13.7	14.0	14.4
NEGRO												
Boys												
6 years-----	84	289	12.9	0.36	0.03	12.1	12.2	12.5	13.0	13.4	13.6	13.7
7 years-----	79	286	13.1	0.37	0.05	12.5	12.6	12.8	13.2	13.6	13.9	14.0
8 years-----	79	279	13.2	0.36	0.03	12.5	12.6	12.9	13.3	13.7	13.9	14.0
9 years-----	74	269	13.3	0.50	0.07	12.2	12.4	13.0	13.4	13.7	13.9	14.1
10 years-----	65	264	13.3	0.64	0.06	12.0	12.3	12.9	13.4	13.8	14.0	14.3
11 years-----	83	255	13.3	0.65	0.06	12.1	12.4	13.0	13.4	13.8	14.0	14.2
Girls												
6 years-----	72	281	13.0	0.34	0.05	12.3	12.4	12.6	13.0	13.4	13.6	13.7
7 years-----	93	284	13.2	0.48	0.04	12.1	12.2	12.8	13.3	13.6	13.8	13.9
8 years-----	113	281	13.1	0.57	0.05	11.8	12.1	12.6	13.2	13.6	13.9	14.0
9 years-----	84	265	13.3	0.60	0.06	12.1	12.4	13.0	13.4	13.8	14.0	14.2
10 years-----	77	266	13.1	0.80	0.07	11.2	12.0	12.7	13.3	13.8	14.1	14.3
11 years-----	84	253	13.2	0.74	0.09	11.6	12.1	12.8	13.4	13.8	14.1	14.3

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

In computing this ratio height is expressed in inches and weight in pounds, which produces a different result than would the use of metric measures.

Table 4. Sitting height of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	s_x	Percentile						
						5th	10th	25th	50th	75th	90th	95th
<u>WHITE</u>												
Boys												
6 years-----	489	1,787	65.0	2.68	0.13	60.4	61.4	63.2	65.1	66.7	68.5	69.7
7 years-----	551	1,781	67.2	2.74	0.15	63.0	64.0	65.4	67.2	68.9	70.8	71.8
8 years-----	537	1,739	69.5	2.94	0.12	65.1	65.7	67.6	69.6	71.5	73.3	74.3
9 years-----	525	1,730	71.6	3.15	0.19	66.4	67.4	69.6	71.6	73.7	75.6	76.7
10 years-----	509	1,692	73.3	3.07	0.20	68.3	69.5	71.4	73.3	75.4	77.3	78.7
11 years-----	542	1,662	75.6	3.10	0.13	70.6	71.5	73.5	75.5	77.7	79.6	80.7
Girls												
6 years-----	461	1,722	64.2	3.00	0.18	59.2	60.4	62.3	64.3	66.1	68.2	69.1
7 years-----	512	1,716	66.4	2.99	0.12	61.5	62.6	64.3	66.5	68.4	70.4	71.4
8 years-----	498	1,674	68.8	2.89	0.13	63.7	64.8	67.1	68.9	70.8	72.4	73.3
9 years-----	494	1,663	71.1	3.19	0.17	65.8	67.1	68.9	71.1	73.4	75.3	76.3
10 years-----	505	1,632	73.5	3.39	0.14	68.2	69.2	71.1	73.5	75.7	77.6	79.1
11 years-----	477	1,605	76.5	3.96	0.15	70.4	72.0	74.2	76.2	78.8	81.6	83.7
<u>NEGRO</u>												
Boys												
6 years-----	84	289	63.2	2.48	0.35	59.2	59.7	61.4	63.2	64.8	66.5	68.1
7 years-----	79	286	65.6	2.58	0.29	61.6	62.5	63.5	65.6	67.5	69.1	70.1
8 years-----	79	279	67.8	2.81	0.29	63.5	64.3	66.2	67.5	69.4	71.8	72.7
9 years-----	74	269	69.3	3.69	0.40	63.7	64.5	66.7	68.6	71.7	74.5	75.8
10 years-----	65	264	70.9	3.54	0.33	64.8	66.1	68.3	70.6	73.2	75.4	77.5
11 years-----	83	255	73.4	3.36	0.37	67.8	68.7	71.4	73.6	74.8	77.5	79.3
Girls												
6 years-----	72	281	62.3	2.78	0.35	57.4	58.8	60.4	62.3	64.5	66.2	66.6
7 years-----	93	284	64.9	3.05	0.40	60.0	61.2	62.7	64.7	67.0	69.3	70.5
8 years-----	113	281	66.6	3.50	0.26	61.5	62.4	64.5	66.3	68.6	71.5	73.4
9 years-----	84	265	69.6	3.43	0.44	64.2	65.4	67.4	69.5	71.6	74.4	75.5
10 years-----	77	266	71.9	3.77	0.48	66.3	67.6	69.3	71.5	74.4	76.8	78.9
11 years-----	84	253	75.1	4.08	0.38	68.3	69.3	72.5	75.3	78.4	80.3	81.4

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; s_x = standard error of the mean.

Table 5. Subischial length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
In centimeters												
WHITE												
Boys												
6 years-----	489	1,787	53.6	3.24	0.20	48.5	49.6	51.5	53.3	55.7	57.8	59.4
7 years-----	551	1,781	57.2	3.53	0.25	51.6	53.1	55.0	57.1	59.2	61.9	63.7
8 years-----	537	1,739	60.3	3.44	0.18	54.5	55.9	58.2	60.3	62.5	64.7	65.7
9 years-----	525	1,730	63.9	4.52	0.33	57.3	59.1	61.3	64.1	66.6	69.0	70.3
10 years-----	509	1,692	66.9	4.33	0.21	60.2	61.6	64.3	67.0	69.7	72.2	73.7
11 years-----	542	1,662	70.1	4.43	0.21	63.0	64.8	67.5	70.1	73.1	75.4	77.0
Girls												
6 years-----	461	1,722	53.5	3.47	0.18	48.2	49.3	51.1	53.5	55.7	57.7	59.1
7 years-----	512	1,716	57.0	3.81	0.16	51.1	52.4	54.8	56.9	59.3	61.3	62.7
8 years-----	498	1,674	60.5	4.29	0.31	54.5	56.0	58.2	60.5	63.2	65.6	67.0
9 years-----	494	1,663	64.0	4.32	0.23	57.4	59.0	61.4	63.7	66.7	70.1	71.2
10 years-----	505	1,632	67.3	4.32	0.24	60.5	61.7	64.2	67.1	70.3	73.1	74.5
11 years-----	477	1,605	70.9	5.21	0.23	63.8	65.6	68.3	70.9	73.8	76.6	77.8
NEGRO												
Boys												
6 years-----	84	289	56.0	3.49	0.41	51.4	52.1	53.3	55.5	58.2	60.5	61.8
7 years-----	79	286	59.6	3.46	0.37	54.5	55.3	56.8	59.3	61.8	63.8	65.2
8 years-----	79	279	63.5	3.34	0.34	57.7	59.0	60.7	64.0	65.9	67.7	68.7
9 years-----	74	269	65.7	3.83	0.57	59.5	60.6	63.6	65.7	68.3	71.0	71.8
10 years-----	65	264	68.7	5.82	0.72	60.1	63.1	66.4	69.5	72.1	75.3	76.1
11 years-----	83	255	72.2	6.03	0.50	65.4	66.9	70.1	72.4	75.3	78.3	79.2
Girls												
6 years-----	72	281	56.2	3.59	0.57	50.3	51.4	53.5	56.5	59.0	60.5	62.1
7 years-----	93	284	59.7	3.22	0.25	54.9	55.5	57.2	59.5	62.1	63.7	65.5
8 years-----	113	281	62.7	4.14	0.32	56.1	58.0	60.0	62.8	65.3	68.2	71.1
9 years-----	84	265	67.9	5.16	0.56	60.1	62.0	64.5	67.5	71.1	75.1	77.4
10 years-----	77	266	69.9	7.06	0.54	62.4	65.2	67.2	70.8	73.6	77.2	79.1
11 years-----	84	253	74.0	4.18	0.41	67.3	68.5	71.1	73.6	77.3	79.8	81.3

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 6. Sitting height/stature ratio (times 100) of children by race, sex, and age at last birthday; sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	s_x	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
<u>Boys</u>												
6 years-----	489	1,787	54.8	1.33	0.06	53.1	53.6	54.4	55.3	56.2	56.7	57.2
7 years-----	351	1,781	54.0	1.34	0.07	52.2	52.8	53.7	54.5	55.4	56.0	56.5
8 years-----	537	1,739	53.6	1.20	0.05	52.1	52.4	53.2	54.1	54.7	55.5	55.9
9 years-----	525	1,730	52.9	1.66	0.10	51.2	51.6	52.4	53.3	54.2	54.8	55.4
10 years-----	509	1,692	52.3	1.41	0.06	50.8	51.2	51.9	52.7	53.6	54.3	54.7
11 years-----	542	1,662	51.9	1.36	0.06	50.4	51.0	51.5	52.3	53.2	53.8	54.5
<u>Girls</u>												
6 years-----	461	1,722	54.6	1.54	0.08	52.8	53.3	54.2	55.1	55.9	56.8	57.3
7 years-----	512	1,716	53.8	1.51	0.08	52.1	52.6	53.4	54.4	55.2	55.8	56.5
8 years-----	498	1,674	53.2	1.79	0.12	51.4	52.1	52.8	53.6	54.5	55.4	55.8
9 years-----	494	1,663	52.7	1.45	0.08	51.1	51.4	52.2	53.2	54.1	54.7	55.2
10 years-----	505	1,632	52.2	1.28	0.06	50.5	51.1	51.8	52.6	53.5	54.3	54.7
11 years-----	477	1,605	52.0	1.92	0.10	50.2	50.6	51.4	52.4	53.3	53.9	54.6
NEGRO												
<u>Boys</u>												
6 years-----	84	289	53.0	1.39	0.10	51.2	51.7	52.6	53.6	54.5	55.0	55.0
7 years-----	79	286	52.4	1.19	0.14	51.0	51.3	52.1	52.8	53.7	54.5	54.8
8 years-----	79	279	51.6	1.24	0.10	50.0	50.4	51.3	52.2	52.8	53.7	54.4
9 years-----	74	269	51.4	1.51	0.27	49.7	50.2	50.9	51.7	52.6	53.5	53.9
10 years-----	65	264	50.8	2.26	0.26	49.0	49.3	50.2	50.8	52.2	52.9	53.9
11 years-----	83	255	50.5	2.35	0.29	48.8	49.3	50.1	50.7	51.5	52.2	52.7
<u>Girls</u>												
6 years-----	72	281	52.6	1.23	0.17	51.2	51.6	52.2	52.8	53.8	54.8	55.5
7 years-----	93	284	52.1	1.18	0.10	50.7	51.2	51.7	52.5	53.4	54.1	54.6
8 years-----	113	281	51.6	1.52	0.10	49.6	50.2	51.0	52.2	53.2	53.7	53.9
9 years-----	84	265	50.7	1.49	0.19	48.8	49.3	50.2	51.2	52.3	52.9	53.6
10 years-----	77	266	50.8	2.77	0.35	48.8	49.2	49.9	51.1	51.8	52.6	53.4
11 years-----	84	253	50.4	1.23	0.14	49.0	49.3	50.1	50.7	51.6	52.6	53.3

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; s_x = standard error of the mean.

Table 7. Buttock-knee length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
In centimeters												
Boys												
6 years-----	489	1,787	37.1	2.79	0.20	31.7	33.7	35.6	37.3	38.9	40.7	41.5
7 years-----	551	1,781	39.6	3.00	0.26	33.7	36.2	38.0	39.8	41.5	43.2	44.4
8 years-----	537	1,739	41.6	3.21	0.26	35.7	37.7	40.2	41.7	43.6	45.2	46.3
9 years-----	525	1,730	44.0	3.46	0.27	37.8	40.0	41.9	44.1	46.1	47.8	49.5
10 years-----	509	1,692	45.9	3.48	0.25	40.0	41.4	44.2	46.2	48.2	49.9	50.9
11 years-----	542	1,662	48.2	3.45	0.24	42.2	44.1	46.2	48.3	50.5	52.5	53.7
Girls												
6 years-----	461	1,722	37.5	2.86	0.23	32.3	33.7	36.1	37.8	39.5	40.8	41.7
7 years-----	512	1,716	39.7	3.07	0.24	34.4	35.8	38.2	39.9	41.7	43.3	44.1
8 years-----	498	1,674	42.3	3.09	0.19	37.2	38.7	40.5	42.5	44.4	46.1	47.4
9 years-----	494	1,663	44.6	3.49	0.26	38.6	40.4	42.6	44.6	47.1	48.9	50.2
10 years-----	505	1,632	46.8	3.62	0.28	40.4	42.2	44.4	47.2	49.4	51.1	52.5
11 years-----	477	1,605	49.4	3.47	0.22	43.8	45.3	47.3	49.4	51.7	53.8	55.6
NEGRO												
Boys												
6 years-----	64	289	37.7	3.38	0.72	30.6	31.8	36.2	38.2	40.3	41.5	41.9
7 years-----	79	286	40.3	3.48	0.57	32.8	34.6	38.9	40.6	42.4	44.3	45.2
8 years-----	79	279	42.3	3.54	0.69	35.6	36.8	40.3	42.8	44.8	46.4	47.2
9 years-----	74	269	43.8	3.75	0.70	36.7	38.6	41.4	44.2	46.6	48.5	49.5
10 years-----	65	264	46.1	4.02	0.74	39.2	41.5	44.1	46.6	48.7	50.6	51.3
11 years-----	83	255	48.3	3.91	0.67	40.2	44.3	46.7	48.6	50.7	52.5	53.8
Girls												
6 years-----	72	281	37.8	3.76	0.81	30.4	32.5	35.2	38.6	41.0	42.2	42.7
7 years-----	93	284	40.4	3.44	0.56	33.1	35.3	39.2	40.7	42.9	44.1	45.1
8 years-----	113	281	42.8	3.78	0.50	35.6	37.7	40.6	43.1	45.3	47.5	48.9
9 years-----	84	265	45.9	4.10	0.74	38.6	40.5	43.2	46.1	48.7	51.6	53.1
10 years-----	77	266	48.4	3.96	0.82	41.7	44.5	46.3	48.4	50.5	52.8	56.2
11 years-----	84	253	51.0	4.67	0.78	41.8	44.6	48.3	51.1	54.4	57.4	57.9

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 8. Popliteal height of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
In centimeters.												
Boys												
6 years-----	489	1,787	29.1	1.77	0.08	26.3	26.8	27.8	29.2	30.3	31.4	32.1
7 years-----	551	1,781	30.9	1.93	0.13	27.9	28.5	29.6	30.9	32.2	33.5	34.3
8 years-----	537	1,739	32.4	1.94	0.13	29.1	29.9	31.2	32.5	33.7	34.8	35.6
9 years-----	525	1,730	34.1	2.16	0.13	30.7	31.4	32.7	34.2	35.5	36.8	37.7
10 years-----	509	1,692	35.7	2.20	0.12	32.1	32.8	34.3	35.7	37.3	38.8	39.6
11 years-----	542	1,662	37.2	2.30	0.12	33.5	34.3	35.6	37.2	38.8	40.2	41.0
Girls												
6 years-----	461	1,722	28.8	1.75	0.07	25.8	26.5	27.6	28.8	30.1	31.0	31.7
7 years-----	512	1,716	30.5	1.93	0.08	27.4	28.1	29.2	30.4	31.7	32.9	33.7
8 years-----	498	1,674	32.3	2.01	0.13	29.0	29.5	31.1	32.4	33.6	34.7	35.6
9 years-----	494	1,663	34.0	2.26	0.12	30.2	31.2	32.5	33.9	35.4	37.2	37.9
10 years-----	505	1,632	35.5	2.48	0.15	31.6	32.5	34.0	35.4	37.1	38.8	39.7
11 years-----	477	1,605	37.4	2.58	0.14	33.2	34.1	35.7	37.4	39.1	40.5	41.4
NEGRO												
Boys												
6 years-----	84	289	30.2	1.92	0.31	27.1	28.1	28.8	30.1	31.4	33.2	33.8
7 years-----	79	286	32.1	1.98	0.29	29.1	29.7	30.6	31.9	33.6	34.7	35.5
8 years-----	79	279	33.9	1.95	0.18	31.0	31.4	32.5	33.9	35.3	36.7	37.5
9 years-----	74	269	35.4	2.12	0.22	32.0	33.0	34.0	35.2	36.9	38.4	38.8
10 years-----	65	264	36.8	2.26	0.28	33.1	34.1	35.3	36.8	38.5	39.6	40.3
11 years-----	83	255	38.4	2.16	0.18	35.1	35.6	36.8	38.5	39.8	41.3	42.2
Girls												
6 years-----	72	281	29.9	2.03	0.30	26.4	27.2	28.7	29.7	31.5	32.7	33.4
7 years-----	93	284	31.8	1.96	0.19	29.2	29.9	30.4	31.7	33.2	34.4	35.2
8 years-----	113	281	33.2	2.35	0.25	29.7	30.5	31.7	33.1	34.5	36.3	37.4
9 years-----	84	265	35.7	2.39	0.24	31.9	32.5	34.0	35.7	37.6	38.8	39.6
10 years-----	77	266	36.9	2.37	0.35	33.1	33.7	35.3	36.9	38.4	39.6	40.3
11 years-----	84	253	38.5	2.66	0.38	34.2	35.1	36.4	38.4	40.4	41.7	43.3

NOTE: n = sample size, N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 9. Foot length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	s_x	Percentile						
						5th	10th	25th	50th	75th	90th	95th
<u>WHITE</u>												
In centimeters												
<u>Boys</u>												
6 years-----	489	1,787	17.9	1.01	0.06	16.2	16.5	17.2	17.8	18.6	19.4	19.8
7 years-----	551	1,781	18.8	1.05	0.06	17.1	17.4	18.1	18.8	19.6	20.4	20.8
8 years-----	537	1,739	19.6	1.20	0.07	17.7	18.1	18.7	19.6	20.5	21.3	21.7
9 years-----	525	1,730	20.7	1.23	0.07	18.5	19.1	19.9	20.6	21.5	22.4	22.8
10 years-----	509	1,692	21.4	1.30	0.06	19.2	20.0	20.4	21.4	22.4	23.2	23.7
11 years-----	542	1,662	22.3	1.34	0.07	20.1	20.4	21.3	22.3	23.2	24.1	24.7
<u>Girls</u>												
6 years-----	461	1,722	17.7	1.07	0.08	15.7	16.2	17.1	17.7	18.5	19.2	19.6
7 years-----	512	1,716	18.5	1.06	0.04	16.6	17.1	17.7	18.5	19.3	19.9	20.5
8 years-----	498	1,674	19.4	1.14	0.06	17.4	18.1	18.5	19.5	20.3	21.0	21.5
9 years-----	494	1,663	20.3	1.20	0.06	18.2	18.6	19.5	20.4	21.1	21.9	22.5
10 years-----	505	1,632	21.1	1.33	0.06	19.1	19.4	20.3	21.2	22.1	22.8	23.3
11 years-----	477	1,605	22.0	1.27	0.06	20.0	20.3	21.1	21.9	22.9	23.7	24.2
<u>NEGRO</u>												
<u>Boys</u>												
6 years-----	84	289	18.5	1.07	0.17	16.7	17.1	17.7	18.5	19.2	19.9	20.5
7 years-----	79	286	19.5	1.02	0.12	17.5	18.0	18.7	19.5	20.2	20.8	21.3
8 years-----	79	279	20.4	1.06	0.14	18.8	19.1	19.6	20.4	21.2	22.0	22.4
9 years-----	74	269	21.2	1.30	0.12	18.9	19.4	20.3	21.2	21.9	22.8	24.1
10 years-----	65	264	22.0	1.22	0.15	20.1	20.3	20.9	22.1	22.9	23.6	23.9
11 years-----	83	255	22.9	1.20	0.12	20.7	21.2	22.1	23.1	23.6	24.3	24.8
<u>Girls</u>												
6 years-----	72	281	18.4	1.10	0.14	16.4	17.1	17.6	18.4	19.3	19.8	20.2
7 years-----	93	284	19.3	0.97	0.10	17.6	17.8	18.6	19.4	20.1	20.7	21.0
8 years-----	113	281	20.0	1.26	0.10	18.2	18.4	19.0	20.1	21.0	21.8	22.4
9 years-----	84	265	21.2	1.40	0.18	18.8	19.2	20.2	21.0	22.4	23.2	23.7
10 years-----	77	266	22.2	1.20	0.18	20.1	20.5	21.3	22.2	23.1	23.8	24.4
11 years-----	84	253	22.7	1.52	0.20	20.1	20.6	21.5	22.7	23.8	24.9	25.3

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean, s = standard deviation; s_x = standard error of the mean.

Table 10. Acromion-olecranon length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
6 years-----	489	1,787	23.7	1.31	0.08	21.5	22.1	22.7	23.6	24.6	25.6	25.9
7 years-----	551	1,781	25.0	1.37	0.08	22.6	23.2	24.1	25.1	25.9	26.7	27.4
8 years-----	537	1,739	26.2	1.39	0.08	23.9	24.4	25.3	26.3	27.2	28.1	28.7
9 years-----	525	1,730	27.6	1.60	0.11	25.1	25.5	26.4	27.5	28.6	29.8	30.5
10 years-----	509	1,692	28.6	1.66	0.07	25.8	26.4	27.6	28.7	29.8	30.8	31.4
11 years-----	542	1,662	30.0	1.85	0.08	27.2	27.7	28.8	30.1	31.2	32.3	33.0
Girls												
6 years-----	461	1,722	23.4	1.41	0.09	21.2	21.5	22.5	23.5	24.5	25.3	25.8
7 years-----	512	1,716	24.6	1.41	0.04	22.3	22.8	23.6	24.6	25.7	26.5	26.8
8 years-----	498	1,674	26.1	1.57	0.08	23.5	24.2	25.2	26.2	27.2	28.3	29.0
9 years-----	494	1,663	27.5	1.61	0.09	25.1	25.4	26.4	27.5	28.6	29.7	30.4
10 years-----	505	1,632	28.8	1.88	0.09	25.9	26.5	27.5	28.8	30.1	31.3	31.8
11 years-----	477	1,605	30.4	1.88	0.08	27.4	28.2	29.2	30.4	31.8	32.9	33.6
NEGRO												
Boys												
6 years-----	84	289	24.1	1.32	0.16	22.2	22.4	23.2	24.1	25.1	25.9	26.6
7 years-----	79	286	25.2	1.58	0.15	22.7	23.1	24.0	25.3	26.3	27.6	28.2
8 years-----	79	279	26.9	1.16	0.15	24.9	25.3	26.2	27.0	27.6	28.4	28.8
9 years-----	74	269	27.6	1.52	0.24	24.7	25.4	26.6	27.6	28.6	29.6	29.9
10 years-----	65	264	28.2	1.88	0.22	25.2	26.2	27.3	28.3	29.7	30.6	30.8
11 years-----	83	255	30.2	1.64	0.15	27.8	28.2	29.0	30.1	31.3	32.7	33.5
Girls												
6 years-----	72	281	23.9	1.44	0.16	21.3	22.0	23.1	24.1	24.9	25.7	26.2
7 years-----	93	284	25.4	1.35	0.14	23.3	24.0	24.4	25.2	26.4	27.4	27.8
8 years-----	113	281	26.4	1.68	0.15	23.8	24.3	25.2	26.2	27.5	28.8	29.6
9 years-----	84	265	28.2	1.74	0.16	24.7	26.1	27.1	28.1	29.4	30.6	31.1
10 years-----	77	266	29.7	1.85	0.22	26.6	27.3	28.4	29.6	30.9	32.2	32.8
11 years-----	84	253	31.1	2.00	0.24	27.7	28.3	30.0	31.0	32.6	33.8	34.3

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 11. Elbow-wrist length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
In centimeters												
Boys												
6 years-----	489	1,787	18.2	0.97	0.05	16.5	17.0	17.4	18.2	18.8	19.6	19.9
7 years-----	551	1,781	19.2	1.11	0.07	17.4	17.8	18.4	19.2	19.8	20.7	21.1
8 years-----	537	1,739	20.0	1.15	0.08	18.1	18.5	19.3	20.1	20.8	21.6	21.9
9 years-----	525	1,730	21.1	1.22	0.08	19.1	19.4	20.2	21.1	21.8	22.7	23.3
10 years-----	509	1,692	21.8	1.24	0.07	19.7	20.2	21.1	21.8	22.8	23.6	23.9
11 years-----	542	1,662	22.8	1.40	0.05	20.5	21.1	22.0	22.8	23.7	24.7	25.3
Girls												
6 years-----	461	1,722	17.7	1.02	0.06	16.0	16.3	17.1	17.7	18.5	19.1	19.6
7 years-----	512	1,716	18.7	1.16	0.05	17.0	17.3	18.0	18.6	19.5	20.2	20.7
8 years-----	498	1,674	19.7	1.15	0.05	17.7	18.2	18.8	19.7	20.6	21.4	21.7
9 years-----	494	1,663	20.7	1.21	0.07	18.8	19.2	19.8	20.6	21.6	22.5	22.9
10 years-----	505	1,632	21.7	1.35	0.05	19.5	20.1	20.7	21.7	22.7	23.6	24.0
11 years-----	477	1,605	22.9	1.44	0.06	20.5	21.1	21.9	22.8	23.8	24.8	25.4
NEGRO												
Boys												
6 years-----	84	289	19.0	1.05	0.16	17.1	17.6	18.3	18.8	19.7	20.5	20.8
7 years-----	79	286	20.1	1.16	0.12	18.3	18.6	19.2	19.9	20.8	21.8	22.4
8 years-----	79	279	21.0	1.13	0.13	19.2	19.5	20.2	21.0	21.8	22.7	23.1
9 years-----	74	269	21.9	1.43	0.14	20.1	20.3	20.9	21.7	22.7	23.7	24.5
10 years-----	65	264	22.6	1.23	0.20	20.6	21.2	21.8	22.6	23.5	24.4	24.8
11 years-----	83	255	23.8	1.23	0.11	21.7	22.2	23.1	23.8	24.6	25.3	25.8
Girls												
6 years-----	72	281	18.6	1.11	0.17	16.4	17.1	18.1	18.7	19.5	20.0	20.5
7 years-----	93	284	19.6	1.10	0.10	18.0	18.2	18.8	19.6	20.5	21.0	21.6
8 years-----	113	281	20.6	1.34	0.11	18.6	19.1	19.6	20.5	21.4	22.4	23.1
9 years-----	84	265	22.0	1.42	0.13	19.6	20.1	21.1	21.9	23.1	23.9	24.5
10 years-----	77	266	23.1	1.42	0.18	21.1	21.3	22.1	22.9	24.1	25.2	25.9
11 years-----	84	253	24.3	1.59	0.22	21.7	22.2	22.9	24.3	25.4	26.5	26.9

NOTE: n = sample size, N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 12. Hand length of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	s_x	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	12.9	0.68	0.04	11.7	12.1	12.4	13.1	13.6	13.9	16.5
7 years-----	551	1,781	13.5	0.71	0.04	12.2	12.4	13.1	13.5	14.0	14.6	14.3
8 years-----	537	1,739	14.1	0.81	0.04	12.5	13.1	13.5	14.2	14.7	15.3	14.8
9 years-----	525	1,730	14.6	0.78	0.04	13.2	13.5	14.2	14.6	15.2	15.7	15.6
10 years-----	509	1,692	15.1	0.82	0.04	13.7	14.1	14.5	15.1	15.7	16.3	15.9
11 years-----	542	1,662	15.7	0.87	0.03	14.2	14.4	15.1	15.6	16.3	16.8	17.3
Girls												
6 years-----	461	1,722	12.8	0.70	0.04	11.4	11.8	12.2	12.7	13.4	13.8	14.2
7 years-----	512	1,716	13.3	0.76	0.03	12.1	12.3	12.8	13.4	13.8	14.4	14.7
8 years-----	498	1,674	13.9	0.76	0.05	12.4	13.0	13.3	13.8	14.5	14.9	15.4
9 years-----	494	1,663	14.6	0.77	0.04	13.2	13.4	14.1	14.5	15.1	15.7	16.0
10 years-----	505	1,632	15.1	0.86	0.04	13.5	14.0	14.4	15.2	15.7	16.4	16.7
11 years-----	477	1,605	15.9	0.91	0.04	14.3	14.6	15.2	15.8	16.6	17.2	17.6
NEGRO												
Boys												
6 years-----	84	289	13.6	0.66	0.09	12.3	12.6	13.1	13.6	14.1	14.6	14.8
7 years-----	79	286	14.2	0.71	0.09	13.1	13.2	13.6	14.2	14.7	15.3	15.7
8 years-----	79	279	14.7	0.73	0.10	13.5	13.7	14.2	14.7	15.4	15.7	15.8
9 years-----	74	269	15.2	0.83	0.07	13.8	14.1	14.5	15.2	15.7	16.5	16.9
10 years-----	65	264	15.7	0.84	0.11	14.2	14.4	15.1	15.7	16.4	16.8	17.0
11 years-----	83	255	16.3	0.76	0.07	14.9	15.2	15.8	16.4	16.8	17.4	17.7
Girls												
6 years-----	72	281	13.6	0.85	0.09	12.1	12.3	13.0	13.6	14.3	14.7	14.8
7 years-----	93	284	14.1	0.72	0.09	13.0	13.1	13.5	14.2	14.6	15.2	15.5
8 years-----	113	281	14.6	0.89	0.08	13.1	13.3	13.8	14.5	15.3	15.9	16.4
9 years-----	84	265	15.5	1.01	0.08	13.9	14.2	14.6	15.4	16.2	16.8	17.3
10 years-----	77	266	16.0	1.00	0.08	14.3	15.0	15.3	15.8	16.6	17.3	17.8
11 years-----	84	253	16.6	1.06	0.12	15.0	15.2	15.8	16.6	17.5	18.3	18.5

NOTE: n = sample size; N = estimated number of children in population in thousands, \bar{X} = mean; s = standard deviation; s_x = standard error of the mean.

Table 13. Biacromial breadth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
In centimeters												
Boys												
6 years-----	489	1,787	26.0	1.59	0.09	23.4	24.1	25.1	26.1	27.2	28.1	28.8
7 years-----	551	1,781	27.3	1.57	0.09	24.7	25.3	26.3	27.3	28.4	29.5	30.1
8 years-----	537	1,739	28.4	1.76	0.10	25.4	26.2	27.3	28.4	29.6	30.7	31.4
9 years-----	525	1,730	29.6	1.92	0.11	27.0	27.4	28.4	29.5	30.8	32.2	32.9
10 years-----	509	1,692	30.4	1.97	0.12	27.3	28.2	29.4	30.5	31.7	32.7	33.6
11 years-----	542	1,662	31.8	2.06	0.13	28.8	29.4	30.5	31.8	33.1	34.5	35.3
Girls												
6 years-----	461	1,722	25.8	1.53	0.09	23.3	24.0	24.7	25.7	26.7	27.7	28.5
7 years-----	512	1,716	27.0	1.54	0.06	24.5	25.1	26.0	27.0	27.8	28.8	29.6
8 years-----	498	1,674	28.1	1.62	0.09	25.4	26.2	27.1	28.1	29.4	30.5	30.8
9 years-----	494	1,663	29.2	1.87	0.09	26.3	27.1	28.1	29.3	30.5	31.6	32.3
10 years-----	505	1,632	30.4	1.93	0.09	27.4	28.1	29.1	30.4	31.7	33.1	33.7
11 years-----	477	1,605	31.8	2.23	0.10	28.6	29.3	30.5	31.7	33.3	34.6	35.5
NEGRO												
Boys												
6 years-----	84	289	26.3	1.95	0.27	23.7	24.2	25.0	26.2	27.4	29.0	29.5
7 years-----	79	286	27.4	1.58	0.17	24.5	25.3	26.4	27.3	28.4	29.5	30.0
8 years-----	79	279	28.9	1.61	0.22	25.7	26.9	27.7	29.1	30.1	30.9	31.6
9 years-----	74	269	29.6	2.00	0.25	26.3	27.0	28.3	29.5	31.1	31.9	33.0
10 years-----	65	264	30.8	1.57	0.27	28.4	29.1	29.6	30.8	32.1	33.1	33.6
11 years-----	83	255	32.0	1.90	0.16	28.9	30.0	30.8	32.1	33.0	34.3	35.6
Girls												
6 years-----	72	281	26.0	1.73	0.23	23.2	23.7	24.8	25.7	27.3	28.5	28.9
7 years-----	93	284	27.2	1.52	0.18	25.1	25.3	26.1	27.2	28.3	29.1	30.1
8 years-----	113	281	28.1	2.01	0.25	25.5	26.1	26.7	27.8	29.6	30.7	31.5
9 years-----	84	265	29.8	2.00	0.22	26.6	27.3	28.3	29.6	31.2	32.5	33.4
10 years-----	77	266	31.1	2.14	0.28	27.1	28.4	29.7	31.2	32.6	33.6	34.1
11 years-----	84	253	32.3	2.47	0.29	28.8	29.4	30.5	32.2	34.4	35.6	35.9

NOTE: n = sample size; N = estimated number of children in population in thousands, \bar{X} = mean, s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 14. Biorystal breadth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	s_x	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	18.2	1.28	0.08	16.1	16.5	17.3	18.2	19.2	19.8	20.4
7 years-----	551	1,781	19.0	1.36	0.07	16.8	17.3	18.1	19.1	19.8	20.7	21.2
8 years-----	537	1,739	19.8	1.61	0.07	17.3	17.9	18.7	19.7	20.6	21.6	22.4
9 years-----	525	1,730	20.7	1.90	0.11	18.2	18.7	19.6	20.6	21.6	22.7	23.9
10 years-----	509	1,692	21.3	2.02	0.09	18.4	19.2	20.2	21.2	22.3	23.4	24.4
11 years-----	542	1,662	22.3	2.16	0.08	19.5	20.2	21.1	22.2	23.4	24.8	25.9
Girls												
6 years-----	461	1,722	18.2	1.44	0.08	16.1	16.5	17.3	18.2	19.2	20.1	20.7
7 years-----	512	1,716	19.0	1.47	0.06	16.5	17.1	18.1	19.1	20.0	20.9	21.6
8 years-----	498	1,674	20.0	1.86	0.11	17.2	18.0	18.8	20.1	21.1	22.4	23.0
9 years-----	494	1,663	21.0	2.01	0.12	18.1	18.7	19.7	20.8	22.3	23.8	24.8
10 years-----	505	1,632	21.9	2.13	0.12	18.7	19.4	20.5	21.7	23.1	25.0	26.2
11 years-----	477	1,605	23.3	2.46	0.11	20.1	20.5	21.6	23.2	24.7	26.6	27.7
NEGRO												
Boys												
6 years-----	84	289	17.1	1.05	0.19	15.3	15.6	16.3	17.0	17.7	18.5	19.1
7 years-----	79	286	17.8	1.07	0.14	16.1	16.3	17.0	17.7	18.5	19.3	20.0
8 years-----	79	279	18.8	1.21	0.17	16.5	17.0	18.0	19.0	19.6	20.2	20.7
9 years-----	74	269	19.1	1.60	0.37	16.4	17.1	18.1	18.8	20.3	21.3	21.9
10 years-----	65	264	19.6	1.66	0.31	17.2	17.6	18.5	19.5	20.6	21.8	22.7
11 years-----	83	255	20.9	1.56	0.19	18.6	19.2	20.1	20.7	21.8	22.8	23.5
Girls												
6 years-----	72	281	17.0	1.27	0.14	14.9	15.4	16.2	17.0	17.8	18.7	19.1
7 years-----	93	284	17.7	1.43	0.22	15.7	16.2	17.0	17.7	18.6	19.5	20.3
8 years-----	113	281	18.6	1.81	0.18	16.2	16.6	17.3	18.3	19.7	21.3	22.6
9 years-----	84	265	19.7	1.78	0.27	17.1	17.5	18.4	19.5	20.8	21.9	22.8
10 years-----	77	266	21.3	2.39	0.25	17.8	18.6	19.6	21.1	22.8	24.5	25.1
11 years-----	84	253	22.4	2.65	0.33	19.0	19.4	20.5	21.7	23.9	26.3	27.6

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; s_x = standard error of the mean.

Table 15. Biacromial breadth/bicristal breadth ratio (times 100) of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
6 years	489	1,787	143.6	8.65	0.41	131.4	134.1	138.4	143.9	149.1	154.1	158.3
7 years	551	1,781	144.0	8.26	0.41	131.6	134.2	138.9	144.7	149.5	154.6	157.4
8 years	537	1,739	144.2	9.98	0.48	131.1	134.3	139.5	144.8	149.7	155.9	158.8
9 years	525	1,730	143.6	9.39	0.41	130.6	134.2	138.8	144.0	148.9	154.3	157.7
10 years	509	1,692	144.1	12.64	0.86	129.5	133.3	138.5	144.2	150.2	155.7	159.8
11 years	542	1,662	143.1	11.12	0.60	126.8	131.3	137.2	144.2	149.8	155.0	158.8
Girls												
6 years	461	1,722	141.9	9.02	0.53	128.2	131.7	136.7	141.8	147.7	153.2	157.2
7 years	512	1,716	142.1	8.64	0.41	128.6	131.9	137.1	142.1	147.8	152.9	156.6
8 years	498	1,674	141.4	10.19	0.57	126.7	130.3	135.7	141.3	147.2	152.2	155.9
9 years	494	1,663	139.5	10.72	0.47	123.4	128.1	133.3	140.2	146.1	151.8	156.6
10 years	505	1,632	139.5	9.69	0.68	123.4	128.2	133.7	140.1	146.5	152.4	155.1
11 years	477	1,605	137.7	12.62	0.52	120.5	125.6	131.7	137.8	143.8	150.6	155.3
NEGRO												
Boys												
6 years	84	289	154.7	11.27	1.79	140.3	144.2	149.3	155.3	159.4	164.4	166.0
7 years	79	286	154.5	8.94	0.74	140.0	144.3	148.9	154.3	160.7	167.2	170.4
8 years	79	279	153.9	7.91	0.88	142.0	146.0	148.2	153.3	160.4	164.8	167.5
9 years	74	269	155.7	9.19	1.98	142.1	144.5	150.8	156.3	162.1	166.5	174.0
10 years	65	264	157.9	11.85	1.49	143.8	145.6	150.8	156.4	162.4	182.1	199.0
11 years	83	255	153.2	9.23	1.29	136.8	140.4	147.2	153.8	161.2	165.2	166.4
Girls												
6 years	72	281	153.2	9.41	0.65	141.4	142.5	145.7	153.6	158.2	166.6	168.6
7 years	93	284	153.8	10.54	1.68	139.4	140.6	146.1	154.4	160.4	168.4	175.9
8 years	113	281	151.6	11.11	1.38	133.8	138.9	145.2	152.9	159.6	165.5	169.5
9 years	84	265	152.0	8.80	1.69	138.4	141.5	147.1	151.6	158.4	164.7	168.9
10 years	77	266	146.9	12.10	1.19	129.9	134.3	137.9	145.8	156.7	162.7	166.8
11 years	84	253	145.2	11.03	1.33	127.6	132.6	139.2	147.0	154.1	158.3	162.2

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 16. Chest breadth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	18.4	1.16	0.09	16.4	17.0	17.6	18.4	19.2	19.9	20.5
7 years-----	551	1,781	19.0	1.17	0.08	17.1	17.5	18.3	19.1	19.8	20.6	21.1
8 years-----	537	1,739	19.8	1.36	0.11	18.0	18.2	18.9	19.7	20.7	21.6	22.1
9 years-----	525	1,730	20.6	1.60	0.10	18.3	18.8	19.6	20.5	21.6	22.6	23.3
10 years-----	509	1,692	21.0	1.41	0.07	19.0	19.3	20.2	21.0	21.9	22.8	23.6
11 years-----	542	1,662	22.1	1.60	0.10	19.9	20.3	21.1	21.9	23.1	24.4	24.9
Girls												
6 years-----	461	1,722	17.9	1.23	0.08	16.1	16.3	17.1	17.8	18.6	19.5	19.9
7 years-----	512	1,716	18.6	1.29	0.09	16.6	17.1	17.8	18.5	19.4	20.0	20.7
8 years-----	498	1,674	19.2	1.22	0.07	17.3	17.7	18.4	19.3	20.1	20.9	21.6
9 years-----	494	1,663	20.0	1.59	0.08	18.0	18.2	18.9	19.8	20.8	21.9	22.9
10 years-----	505	1,632	20.8	1.82	0.11	18.3	18.7	19.6	20.7	21.8	23.2	24.3
11 years-----	477	1,605	21.7	1.77	0.07	19.1	19.6	20.5	21.6	22.8	24.1	25.2
NEGRO												
Boys												
6 years-----	84	289	18.3	1.11	0.18	16.3	16.8	17.4	18.3	19.0	19.9	20.5
7 years-----	79	286	18.7	1.31	0.17	17.2	17.4	18.0	18.6	19.4	20.1	21.2
8 years-----	79	279	19.6	1.01	0.13	18.1	18.4	19.1	19.6	20.3	20.8	21.2
9 years-----	74	269	20.2	1.30	0.13	18.1	18.4	19.2	20.3	21.3	22.1	22.6
10 years-----	65	264	20.7	1.24	0.19	19.1	19.2	19.7	20.6	21.6	22.8	23.3
11 years-----	83	255	21.7	1.42	0.18	19.4	20.0	20.6	21.6	22.6	23.5	24.0
Girls												
6 years-----	72	281	17.7	1.05	0.14	16.0	16.2	16.8	17.7	18.5	19.1	19.6
7 years-----	93	284	18.2	1.58	0.21	16.3	16.8	17.3	18.1	19.1	19.7	20.1
8 years-----	113	281	19.1	1.66	0.21	17.1	17.3	18.1	18.9	19.9	21.2	21.7
9 years-----	84	265	19.7	1.43	0.16	17.7	18.2	18.8	19.6	20.6	21.7	22.6
10 years-----	77	266	20.5	1.48	0.17	18.2	18.6	19.6	20.6	21.5	22.5	22.8
11 years-----	84	253	21.6	2.05	0.25	19.0	19.4	20.3	21.3	22.9	24.4	26.7

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 17. Chest depth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	13.3	1.19	0.11	11.5	12.0	12.4	13.2	14.1	14.8	15.3
7 years-----	551	1,781	13.7	1.29	0.12	11.8	12.2	13.0	13.6	14.5	15.4	15.9
8 years-----	537	1,739	14.2	1.43	0.10	12.2	12.5	13.3	14.2	14.8	15.8	16.7
9 years-----	525	1,730	14.7	1.52	0.08	12.6	13.1	13.7	14.6	15.6	16.6	17.6
10 years-----	509	1,692	15.0	1.43	0.12	13.0	13.3	14.1	14.9	15.8	16.8	17.6
11 years-----	542	1,662	15.7	1.58	0.10	13.4	14.0	14.7	15.6	16.7	17.8	18.7
Girls												
6 years-----	461	1,722	12.8	1.08	0.07	11.2	11.5	12.2	12.7	13.6	14.4	14.8
7 years-----	512	1,716	13.3	1.30	0.09	11.4	12.0	12.4	13.3	14.0	14.8	15.5
8 years-----	498	1,674	13.7	1.31	0.07	11.8	12.2	12.7	13.7	14.6	15.6	16.3
9 years-----	494	1,663	14.2	1.63	0.09	12.1	12.4	13.2	14.1	15.2	16.5	17.4
10 years-----	505	1,632	14.7	1.70	0.12	12.3	12.8	13.5	14.5	15.7	16.9	17.9
11 years-----	477	1,605	15.4	1.81	0.10	13.1	13.3	14.2	15.3	16.7	17.8	18.6
NEGRO												
Boys												
6 years-----	84	289	13.5	1.13	0.30	11.7	12.1	12.6	13.5	14.4	14.9	15.7
7 years-----	79	286	13.5	1.12	0.17	12.1	12.2	12.8	13.5	14.4	15.3	15.7
8 years-----	79	279	14.1	1.18	0.16	12.2	12.6	13.4	14.2	14.9	15.7	16.0
9 years-----	74	269	14.5	1.44	0.31	12.3	12.9	13.6	14.5	15.5	16.5	16.8
10 years-----	65	264	14.5	1.42	0.23	13.1	13.2	13.7	14.4	15.2	16.2	16.6
11 years-----	83	255	15.2	1.30	0.18	13.2	13.6	14.4	15.3	16.3	16.8	17.4
Girls												
6 years-----	72	281	13.1	1.43	0.21	11.4	11.8	12.3	12.8	13.6	14.4	14.8
7 years-----	93	284	13.1	1.14	0.20	11.1	11.5	12.3	13.1	13.8	14.7	15.2
8 years-----	113	281	13.5	1.26	0.18	11.6	12.1	12.5	13.4	14.4	14.9	16.3
9 years-----	84	265	13.9	1.38	0.14	12.1	12.3	13.0	13.8	14.8	15.7	16.3
10 years-----	77	266	14.7	1.64	0.16	12.5	12.8	13.6	14.5	15.6	16.8	18.3
11 years-----	84	253	15.1	1.96	0.26	12.3	12.7	14.1	14.8	16.3	17.7	20.0

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 18. Bicondylar breadth of the femur of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	7.6	1.14	0.06	6.60	6.90	7.20	7.54	7.88	8.56	8.76
7 years-----	551	1,781	7.9	0.58	0.05	7.04	7.12	7.37	7.78	8.41	8.76	8.87
8 years-----	537	1,739	8.2	0.64	0.05	7.09	7.23	7.62	8.24	8.65	9.00	9.51
9 years-----	525	1,730	8.5	0.73	0.04	7.20	7.44	8.09	8.48	8.87	9.63	9.88
10 years-----	509	1,692	8.7	0.78	0.05	7.36	7.77	8.22	8.62	9.25	9.75	10.09
11 years-----	542	1,662	9.1	0.80	0.06	7.84	8.10	8.43	9.07	9.64	10.31	10.71
Girls												
6 years-----	461	1,722	7.3	0.53	0.05	6.17	6.34	6.84	7.34	7.70	8.13	8.53
7 years-----	512	1,716	7.6	0.58	0.04	6.40	6.80	7.19	7.54	7.88	8.57	8.78
8 years-----	498	1,674	7.9	0.63	0.04	7.01	7.09	7.35	7.78	8.45	8.82	9.23
9 years-----	494	1,663	8.2	0.74	0.05	7.08	7.19	7.53	8.18	8.71	9.40	9.73
10 years-----	505	1,632	8.5	0.77	0.06	7.17	7.38	8.05	8.49	9.07	9.69	9.89
11 years-----	477	1,605	8.8	0.79	0.04	7.40	7.83	8.25	8.68	9.39	9.87	10.44
NEGRO												
Boys												
6 years-----	84	289	7.7	0.54	0.16	6.65	6.81	7.20	7.62	8.20	8.56	8.68
7 years-----	79	286	7.8	0.51	0.10	6.87	7.06	7.28	7.65	8.23	8.65	8.79
8 years-----	79	279	8.1	0.60	0.09	7.20	7.30	7.59	8.16	8.60	8.86	9.50
9 years-----	74	269	8.5	0.66	0.09	7.24	7.48	8.09	8.47	8.85	9.51	9.74
10 years-----	65	264	8.6	0.62	0.09	7.76	7.83	8.13	8.49	8.84	9.54	9.78
11 years-----	83	255	8.9	0.63	0.08	8.01	8.09	8.36	8.81	9.50	9.88	10.36
Girls												
6 years-----	72	281	7.2	0.43	0.07	6.41	6.52	6.85	7.30	7.64	7.84	8.01
7 years-----	93	284	7.5	0.60	0.10	6.41	6.62	7.12	7.47	7.82	8.49	8.75
8 years-----	113	281	7.9	0.78	0.10	6.82	7.07	7.34	7.79	8.52	9.10	9.57
9 years-----	84	265	8.2	0.78	0.10	6.86	7.12	7.54	8.23	8.69	9.32	9.71
10 years-----	77	266	8.6	0.89	0.09	7.20	7.40	8.04	8.49	9.09	9.82	10.59
11 years-----	84	253	8.9	0.95	0.13	7.65	8.00	8.25	8.65	9.48	10.51	11.02

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 19. Upper arm girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
6 years-----	489	1,787	17.6	1.61	0.10	15.3	15.8	16.5	17.5	18.5	19.7	20.6
7 years-----	551	1,781	18.3	1.90	0.12	15.7	16.2	17.1	18.2	19.3	20.7	21.7
8 years-----	537	1,739	19.0	2.24	0.12	16.2	16.7	17.6	18.7	20.3	21.8	23.5
9 years-----	525	1,730	19.9	2.66	0.16	16.5	17.2	18.3	19.5	21.1	23.4	25.8
10 years-----	509	1,692	20.4	2.46	0.13	17.3	18.0	18.8	20.1	21.7	23.9	25.5
11 years-----	542	1,662	21.6	2.85	0.14	18.2	18.8	19.7	21.2	23.3	25.6	27.3
Girls												
6 years-----	461	1,722	17.7	1.86	0.15	15.1	15.6	16.4	17.5	18.8	19.9	20.9
7 years-----	512	1,716	18.3	2.04	0.10	15.7	16.2	17.0	18.2	19.5	21.2	22.1
8 years-----	498	1,674	19.3	2.32	0.12	16.2	16.8	17.7	19.0	20.8	22.4	23.8
9 years-----	494	1,663	20.2	2.70	0.15	16.6	17.3	18.4	19.7	21.8	24.1	25.7
10 years-----	505	1,632	20.9	2.93	0.20	16.9	17.5	18.8	20.6	22.8	24.9	26.5
11 years-----	477	1,605	21.7	2.99	0.17	17.6	18.3	19.6	21.5	23.6	26.1	27.4
NEGRO												
Boys												
6 years-----	84	289	17.4	1.55	0.28	15.1	15.4	16.3	17.3	18.5	19.6	20.5
7 years-----	79	286	17.7	1.28	0.18	15.6	16.1	16.9	17.6	18.5	19.4	19.9
8 years-----	79	279	18.5	1.53	0.16	16.1	17.0	17.5	18.5	19.6	20.5	21.2
9 years-----	74	269	19.1	2.21	0.28	16.3	17.1	17.6	18.7	20.3	22.8	23.8
10 years-----	65	264	19.8	2.07	0.27	16.6	17.5	18.5	19.5	20.9	22.2	24.2
11 years-----	83	255	20.6	2.27	0.26	17.2	18.1	19.3	20.4	22.3	23.7	24.7
Girls												
6 years-----	72	281	17.1	1.41	0.20	14.7	15.2	16.0	17.2	17.9	18.7	20.1
7 years-----	93	284	17.9	2.26	0.25	15.3	15.6	16.5	17.5	18.8	20.7	21.8
8 years-----	113	281	18.8	2.70	0.17	15.3	15.7	16.8	18.3	20.7	22.3	23.7
9 years-----	84	265	19.5	2.65	0.24	15.9	17.0	17.7	19.3	20.8	22.8	24.8
10 years-----	77	266	20.6	3.22	0.42	16.6	17.2	18.4	20.1	22.1	24.8	27.5
11 years-----	84	253	21.5	3.54	0.51	17.2	17.7	19.3	20.8	23.2	27.0	27.9

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 20. Lower arm girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$S_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	17.4	1.32	0.08	15.4	16.0	16.6	17.4	18.2	18.9	19.8
7 years-----	551	1,781	18.0	1.40	0.08	16.0	16.3	17.2	18.0	18.8	19.8	20.6
8 years-----	537	1,739	18.7	1.59	0.09	16.3	16.9	17.6	18.6	19.6	20.8	21.6
9 years-----	525	1,730	19.4	1.74	0.11	16.9	17.3	18.3	19.3	20.4	21.7	22.8
10 years-----	509	1,692	19.9	1.61	0.08	17.4	18.0	18.7	19.7	20.9	22.1	22.9
11 years-----	542	1,662	20.8	1.83	0.09	18.2	18.7	19.6	20.6	21.8	23.5	24.3
Girls												
6 years-----	461	1,722	17.1	1.30	0.10	15.1	15.5	16.3	17.1	17.8	18.8	19.6
7 years-----	512	1,716	17.5	1.42	0.07	15.5	16.1	16.6	17.5	18.5	19.5	20.1
8 years-----	498	1,674	18.3	1.47	0.07	16.1	16.5	17.4	18.3	19.3	20.4	21.2
9 years-----	494	1,663	19.0	1.78	0.10	16.4	17.1	17.9	18.8	20.2	21.5	22.4
10 years-----	505	1,632	19.7	1.91	0.12	17.1	17.4	18.4	19.7	20.9	22.1	22.9
11 years-----	477	1,605	20.5	1.90	0.10	17.5	18.2	19.1	20.4	21.8	23.1	23.9
NEGRO												
Boys												
6 years-----	84	289	17.2	1.30	0.26	15.2	15.5	16.3	17.3	18.2	19.3	19.7
7 years-----	79	286	17.7	1.09	0.14	15.8	16.2	17.1	17.6	18.6	19.4	19.7
8 years-----	79	279	18.4	1.26	0.17	16.3	16.7	17.5	18.4	19.3	20.2	20.8
9 years-----	74	269	19.0	1.59	0.20	17.0	17.2	17.9	18.8	20.2	21.3	21.8
10 years-----	65	264	19.6	1.50	0.20	17.4	18.0	18.6	19.5	20.6	21.9	22.5
11 years-----	83	255	20.4	1.70	0.20	17.6	18.3	19.3	20.3	21.7	22.7	23.3
Girls												
6 years-----	72	281	16.6	1.11	0.14	14.9	15.2	15.9	16.6	17.4	18.0	18.6
7 years-----	93	284	17.4	1.78	0.18	15.2	15.6	16.4	17.4	18.3	19.0	20.2
8 years-----	113	281	18.0	1.70	0.14	15.3	15.7	16.8	17.8	19.2	20.4	21.3
9 years-----	84	265	19.0	1.82	0.20	15.9	17.1	17.7	18.8	20.1	21.5	22.7
10 years-----	77	266	19.7	2.02	0.24	16.6	17.3	18.4	19.7	20.8	23.0	23.8
11 years-----	84	253	20.7	2.47	0.37	17.4	17.9	18.8	20.4	22.0	24.3	25.6

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $S_{\bar{x}}$ = standard error of the mean.

Table 21. Calf girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
<u>Boys</u>												
6 years-----	489	1,787	23.6	1.87	0.12	20.8	21.3	22.3	23.5	24.7	26.1	27.1
7 years-----	551	1,781	24.7	2.14	0.12	21.6	22.3	23.3	24.6	25.8	27.3	28.4
8 years-----	537	1,739	25.7	2.35	0.14	22.3	23.1	24.3	25.6	27.0	28.7	29.9
9 years-----	525	1,730	26.8	2.65	0.16	23.4	24.1	25.1	26.5	28.2	30.4	31.8
10 years-----	509	1,692	27.6	2.49	0.14	24.1	24.7	26.0	27.5	29.3	30.7	32.1
11 years-----	542	1,662	29.0	2.82	0.16	25.2	26.0	27.2	28.6	30.6	32.8	34.6
<u>Girls</u>												
6 years-----	461	1,722	23.7	1.93	0.14	20.6	21.3	22.4	23.6	24.9	26.3	27.3
7 years-----	512	1,716	24.6	1.98	0.11	21.7	22.3	23.3	24.5	25.9	27.4	28.2
8 years-----	498	1,674	25.8	2.21	0.11	22.6	23.3	24.3	25.6	27.3	28.8	30.0
9 years-----	494	1,663	27.0	2.53	0.14	23.4	24.1	25.2	26.7	28.6	30.5	31.7
10 years-----	505	1,632	27.9	2.70	0.17	24.0	24.5	26.1	27.9	29.8	31.5	32.7
11 years-----	477	1,605	29.2	2.87	0.16	25.1	25.6	27.3	28.9	31.3	33.3	34.3
NEGRO												
<u>Boys</u>												
6 years-----	84	289	23.2	1.78	0.30	20.3	20.7	22.1	23.3	24.3	27.8	26.6
7 years-----	79	286	24.2	2.13	0.32	21.3	22.0	22.7	24.2	25.4	26.6	27.4
8 years-----	79	279	25.5	1.80	0.21	22.9	23.3	24.2	25.5	26.8	28.1	28.9
9 years-----	74	269	26.2	2.62	0.30	22.7	23.4	24.5	25.7	27.7	29.6	30.7
10 years-----	65	264	27.2	2.30	0.30	23.7	24.3	25.6	26.8	28.5	30.3	31.7
11 years-----	83	255	28.3	2.42	0.27	24.5	25.2	26.8	27.9	29.9	31.7	32.5
<u>Girls</u>												
6 years-----	72	281	23.2	1.49	0.17	21.0	21.3	22.2	23.1	24.3	25.3	26.1
7 years-----	93	284	24.2	1.86	0.24	21.5	22.1	22.8	24.3	25.2	26.3	26.7
8 years-----	113	281	25.4	2.71	0.21	21.5	22.2	23.5	24.9	27.1	29.3	30.4
9 years-----	84	265	26.4	2.54	0.25	22.5	23.2	24.8	26.2	28.1	30.4	31.2
10 years-----	77	266	27.8	3.24	0.28	23.6	24.2	25.4	27.6	30.1	31.8	33.9
11 years-----	84	253	29.1	3.55	0.38	24.3	24.9	26.5	28.6	31.3	34.8	35.9

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 22. Estimated mid-arm muscle circumference of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	15.0	1.21	0.06	13.1	13.3	14.0	14.8	15.8	16.7	17.2
7 years-----	551	1,781	15.6	1.30	0.09	13.4	13.9	14.6	15.5	16.4	17.3	17.8
8 years-----	537	1,739	16.1	1.49	0.07	14.0	14.3	15.1	16.0	16.9	17.9	18.7
9 years-----	525	1,730	16.6	1.57	0.08	14.1	14.7	15.6	16.6	17.6	18.6	19.3
10 years-----	509	1,692	17.1	1.54	0.08	15.0	15.3	16.1	17.0	18.0	18.9	19.8
11 years-----	542	1,662	18.0	1.73	0.09	15.5	16.1	16.0	17.9	19.0	20.3	21.1
Girls												
6 years-----	461	1,722	14.5	1.20	0.10	12.5	13.0	13.7	14.5	15.3	16.2	16.8
7 years-----	512	1,716	14.9	1.38	0.06	13.0	13.3	14.1	14.8	15.7	16.7	17.3
8 years-----	498	1,674	15.6	1.51	0.07	13.4	13.8	14.5	15.5	16.5	17.5	18.2
9 years-----	494	1,663	16.2	1.62	0.09	13.7	14.2	15.1	16.0	17.1	18.2	19.1
10 years-----	505	1,632	16.8	1.85	0.12	14.1	14.6	15.6	16.6	17.9	19.1	19.9
11 years-----	477	1,605	17.7	2.01	0.14	14.8	15.2	16.2	17.4	18.8	20.3	21.5
NEGRO												
Boys												
6 years-----	84	289	15.2	1.34	0.31	13.1	13.4	14.1	15.2	16.2	17.1	17.7
7 years-----	79	286	15.7	1.18	0.17	13.5	14.1	14.9	15.7	16.5	17.1	17.8
8 years-----	79	279	16.2	1.19	0.12	14.3	14.7	15.4	16.2	16.9	17.9	18.5
9 years-----	74	269	16.8	1.58	0.19	14.6	15.1	15.6	16.6	18.1	18.8	19.7
10 years-----	65	264	17.4	1.43	0.22	15.0	15.6	16.5	17.4	18.3	19.3	19.8
11 years-----	83	255	18.1	1.61	0.17	15.3	15.9	17.2	18.1	19.1	20.4	20.9
Girls												
6 years-----	72	281	14.6	1.25	0.17	12.5	12.8	13.6	14.7	15.6	16.3	16.7
7 years-----	93	284	15.3	1.73	0.24	13.0	13.3	14.2	15.2	16.2	17.2	17.7
8 years-----	113	281	15.8	1.74	0.16	13.2	13.5	14.4	15.6	16.8	18.2	18.8
9 years-----	84	265	16.3	1.61	0.14	14.0	14.3	15.2	16.3	17.2	18.6	19.3
10 years-----	77	266	17.4	2.16	0.29	14.2	15.0	15.8	17.0	18.7	20.3	21.5
11 years-----	84	253	18.1	2.34	0.35	15.1	15.6	16.6	17.7	19.2	20.8	23.1

NOTE: n = sample size; N = estimated number of children in population in thousands \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 23. Chest girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	58.7	3.50	0.17	54.2	54.9	56.4	58.4	60.6	63.1	64.4
7 years-----	551	1,781	60.9	4.08	0.23	55.5	56.5	58.3	60.5	62.9	65.8	68.2
8 years-----	537	1,739	63.3	4.90	0.29	57.3	58.4	60.1	62.5	65.7	68.8	71.8
9 years-----	525	1,730	66.3	5.92	0.41	59.3	60.4	62.5	65.4	68.6	72.8	78.1
10 years-----	509	1,692	67.6	5.33	0.29	60.4	61.8	64.3	67.2	70.1	73.9	76.8
11 years-----	542	1,662	71.3	6.23	0.26	63.8	65.2	67.3	70.1	74.2	80.3	83.3
Girls												
6 years-----	461	1,722	57.0	3.94	0.20	51.8	52.8	54.5	56.7	59.1	61.3	63.8
7 years-----	512	1,716	59.0	4.21	0.23	53.3	54.4	56.3	58.5	61.4	64.6	67.3
8 years-----	498	1,674	61.9	5.16	0.20	55.5	56.5	58.3	61.4	64.4	68.1	71.8
9 years-----	494	1,663	64.6	5.93	0.33	57.3	58.3	60.6	63.5	68.0	72.7	76.9
10 years-----	505	1,632	67.2	6.67	0.37	58.6	60.0	62.5	66.4	71.0	76.6	80.0
11 years-----	477	1,605	70.6	7.12	0.33	60.4	62.6	65.6	69.5	75.5	80.3	83.2
NEGRO												
Boys												
6 years-----	84	289	58.0	3.22	0.61	53.4	54.7	55.5	57.8	60.3	62.6	63.7
7 years-----	79	286	59.8	2.75	0.38	55.9	56.5	57.9	59.7	61.6	63.5	64.9
8 years-----	79	279	62.4	2.89	0.38	58.0	59.2	60.4	62.2	64.2	65.7	67.5
9 years-----	74	269	64.1	4.48	0.46	57.5	58.7	60.4	64.0	67.2	70.8	71.7
10 years-----	65	264	66.0	4.17	0.46	60.8	61.4	63.2	65.5	68.2	72.4	74.1
11 years-----	83	255	68.3	4.77	0.46	61.4	63.0	65.7	67.5	70.1	73.7	79.4
Girls												
6 years-----	72	281	55.9	2.91	0.38	51.2	52.2	54.1	55.8	58.3	59.7	61.3
7 years-----	93	284	57.8	3.88	0.53	53.3	53.8	55.0	57.1	60.3	62.3	63.7
8 years-----	113	281	60.3	5.01	0.42	53.8	54.9	56.6	59.2	63.4	66.9	68.6
9 years-----	84	265	63.2	5.61	0.53	56.5	57.3	58.9	63.1	64.9	70.9	75.6
10 years-----	77	266	66.2	6.64	0.73	56.6	60.2	61.8	65.4	69.0	73.2	81.0
11 years-----	84	253	69.7	7.09	0.97	60.5	61.6	64.3	69.1	74.2	80.7	83.8

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 24. Waist girth of children by race, sex, and age at last birthday; sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
<u>WHITE</u>												
<u>Boys</u>												
In centimeters												
6 years-----	489	1,787	53.0	4.20	0.20	47.4	48.4	50.3	52.5	54.9	58.3	60.2
7 years-----	551	1,781	54.5	4.64	0.26	47.9	49.6	51.7	54.1	56.7	59.7	61.7
8 years-----	537	1,739	56.4	5.64	0.30	49.5	50.7	53.0	55.8	58.7	62.5	66.1
9 years-----	525	1,730	58.3	6.33	0.43	51.3	52.3	54.4	57.3	60.5	66.3	71.5
10 years-----	509	1,692	59.5	6.08	0.29	52.3	53.4	55.3	58.4	62.3	67.2	72.3
11 years-----	542	1,662	62.4	7.27	0.30	54.2	55.5	57.7	60.6	65.5	71.6	77.5
<u>Girls</u>												
6 years-----	461	1,722	51.8	4.58	0.24	45.4	46.8	49.1	51.6	53.9	56.9	59.2
7 years-----	512	1,716	53.1	4.54	0.29	47.2	48.3	50.2	52.5	55.5	59.0	61.5
8 years-----	498	1,674	54.9	5.34	0.24	47.7	49.4	51.6	54.0	57.4	61.7	66.1
9 years-----	494	1,663	57.1	5.87	0.34	50.1	50.9	53.3	56.1	59.9	65.3	68.4
10 years-----	505	1,632	58.2	6.38	0.42	50.4	51.5	54.1	57.3	61.3	65.8	71.4
11 years-----	477	1,605	60.1	6.45	0.31	52.1	53.1	55.5	59.3	63.6	69.0	72.5
<u>NEGRO</u>												
<u>Boys</u>												
6 years-----	84	289	52.4	4.28	0.63	46.9	48.3	49.7	51.7	54.7	57.8	60.6
7 years-----	79	286	53.4	3.15	0.58	47.9	49.4	51.8	53.6	55.2	57.4	59.0
8 years-----	79	279	55.3	3.85	0.45	49.7	50.6	52.9	54.9	57.8	61.1	63.0
9 years-----	74	269	56.2	4.71	0.58	49.3	50.8	53.1	57.8	58.8	61.6	64.3
10 years-----	65	264	57.7	4.29	0.44	52.3	52.6	55.2	57.4	59.6	63.9	67.4
11 years-----	83	255	60.0	5.25	0.44	53.1	55.1	57.1	59.5	62.3	65.3	72.5
<u>Girls</u>												
6 years-----	72	281	51.5	3.28	0.38	45.5	46.7	49.6	51.8	53.9	55.2	56.4
7 years-----	93	284	52.6	4.11	0.55	47.5	48.4	49.6	52.2	54.6	57.6	61.7
8 years-----	113	281	54.8	5.47	0.36	47.8	49.1	50.8	53.7	58.3	61.7	63.9
9 years-----	84	265	56.8	5.27	0.52	51.0	51.6	53.4	56.7	58.5	64.2	67.8
10 years-----	77	266	58.3	5.57	0.59	51.1	52.7	54.7	56.8	60.9	64.7	71.5
11 years-----	84	253	60.9	8.80	1.22	53.0	53.5	56.2	58.8	63.1	68.6	78.4

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 25. Hip girth of children by race, sex, and age at last birthday: sample size, mean, standard deviation, standard error of the mean, and selected percentiles, United States, 1963-65

Race, sex, and age	n	N	\bar{X}	s	$s_{\bar{x}}$	Percentile						
						5th	10th	25th	50th	75th	90th	95th
WHITE												
Boys												
In centimeters												
6 years-----	489	1,787	58.3	4.82	0.37	51.4	52.5	55.1	58.0	61.1	64.5	66.6
7 years-----	551	1,781	61.1	5.24	0.35	53.7	55.4	57.8	60.5	63.8	67.7	69.8
8 years-----	537	1,739	64.2	5.93	0.42	56.5	57.7	60.4	63.7	67.2	71.6	75.1
9 years-----	525	1,730	67.3	7.23	0.53	58.3	59.6	62.6	66.1	70.6	76.2	81.7
10 years-----	509	1,692	69.4	6.54	0.47	60.6	62.4	65.0	68.5	73.4	78.4	80.9
11 years-----	542	1,662	73.5	7.49	0.43	63.0	65.6	68.3	72.4	77.5	83.8	88.1
Girls												
6 years-----	461	1,722	59.1	5.21	0.34	51.1	53.1	55.6	58.8	62.3	65.6	67.6
7 years-----	512	1,716	62.3	5.55	0.32	54.6	56.1	58.3	61.6	65.8	69.6	72.5
8 years-----	498	1,674	65.7	6.14	0.36	56.2	57.7	61.7	65.4	69.3	73.5	76.7
9 years-----	494	1,663	68.9	7.12	0.49	59.1	61.0	63.7	68.0	73.7	79.2	81.6
10 years-----	505	1,632	72.4	7.89	0.57	61.2	63.2	66.7	71.8	76.9	82.7	86.3
11 years-----	477	1,605	76.6	8.25	0.46	64.6	66.5	71.0	75.7	82.2	87.4	92.0
NEGRO												
Boys												
6 years-----	84	289	56.2	3.86	0.39	51.1	51.5	54.4	56.3	58.4	61.3	63.1
7 years-----	79	286	58.4	3.34	0.42	52.8	54.1	56.2	58.5	61.4	62.5	63.1
8 years-----	79	279	62.8	4.74	0.76	55.2	56.5	60.1	62.8	65.8	68.6	71.0
9 years-----	74	269	64.9	6.14	1.04	57.3	58.1	60.3	64.7	67.6	72.5	77.9
10 years-----	65	264	67.4	5.79	0.84	58.8	59.9	64.3	66.7	70.1	75.3	78.9
11 years-----	83	255	70.9	6.32	0.88	61.1	62.0	66.8	70.8	74.7	78.0	80.2
Girls												
6 years-----	72	281	57.2	4.44	0.64	49.7	51.1	54.4	57.3	60.3	63.5	64.5
7 years-----	93	284	60.0	5.67	0.71	53.3	54.2	56.4	59.4	62.4	66.6	70.5
8 years-----	113	281	64.3	6.90	0.53	54.5	55.9	59.3	63.6	69.0	73.4	76.9
9 years-----	84	265	67.8	7.53	0.85	58.1	59.6	62.6	66.7	71.4	77.6	79.8
10 years-----	77	266	72.4	8.55	0.88	60.1	61.9	66.9	71.5	76.8	83.8	86.7
11 years-----	84	253	77.1	10.56	1.64	63.4	64.9	70.4	74.9	81.8	91.5	100.7

NOTE: n = sample size; N = estimated number of children in population in thousands; \bar{X} = mean; s = standard deviation; $s_{\bar{x}}$ = standard error of the mean.

Table 26. Normal deviates of differences between measurements of various components of the extremities of white and Negro children and ranking of normal deviates within sex and age groups

Sex and age	Lower extremity			Upper extremity		
	Foot length	Popliteal height	Buttock-knee length	Hand length	Elbow-wrist length	Acromion-olecranon length
<u>Boys</u>						
Normal deviate						
6 years -----	-3.13	-3.52	-0.81	-6.14	-4.80	-2.61
7 years -----	-4.64	-3.65	-1.12	-7.45	-6.47	-1.42
8 years -----	-5.14	-6.44	-0.98	-6.24	-6.64	-4.08
9 years -----	-3.66	-4.87	+0.17	-7.61	-5.42	-
10 years -----	-3.57	-3.32	-0.22	-5.01	-3.61	+1.93
11 years -----	-4.50	-5.82	-0.20	-8.77	-7.96	-1.47
<u>Girls</u>						
6 years -----	-4.44	-3.66	-0.31	-8.23	-4.98	-2.50
7 years -----	-7.12	-6.18	-1.15	-8.41	-8.38	-5.37
8 years -----	-5.14	-3.35	-0.97	-6.96	-7.17	-1.49
9 years -----	-4.58	-6.46	-1.58	-10.57	-8.78	-3.75
10 years -----	-5.37	-3.54	-1.92	-9.79	-7.41	-3.86
11 years -----	-3.59	-2.74	-1.98	-6.48	-6.09	-2.50
<u>Boys</u>						
Rank of normal deviate						
6 years -----	2	3	1	3	2	1
7 years -----	3	2	1	3	2	1
8 years -----	2	3	1	2	3	1
9 years -----	2	3	1	3	2	1
10 years -----	3	2	1	3	2	1
11 years -----	2	3	1	3	2	1
<u>Girls</u>						
6 years -----	3	2	1	3	2	1
7 years -----	3	2	1	3	2	1
8 years -----	3	2	1	2	3	1
9 years -----	2	3	1	3	2	1
10 years -----	3	2	1	3	2	1
11 years -----	3	2	1	3	2	1

APPENDIX I STATISTICAL NOTES

The Survey Design

The sampling plan of the second cycle of the HES followed a highly stratified, multistage probability design in which a sample of the U.S. population (including Alaska and Hawaii) from the ages of 6 through 11 years inclusive was selected. Excluded were those children confined to an institution or residing upon any of the reservation lands set up for the American Indians.

In the first stage of this design, the nearly 2,000 primary sampling units (PSU's), geographic units into which the United States was divided, were grouped into 357 strata for the use of the Health Interview Survey and the Current Population Survey of the U.S. Bureau of the Census and were then further grouped into 40 superstrata for use in Cycle II of the HES.

The average size of each Cycle II stratum was 4.5 million persons, and all strata fell between the limits of 3.5 and 5.5 million. Grouping into 40 strata was done in a way that maximized homogeneity of the PSU's included in each stratum, particularly with regard to the degree of urbanization, geographic proximity, and degree of industrialization. The 40 strata were classified into four broad geographic regions (each with 10 strata) of approximately equal population and cross-classified into four broad population density groups (each having 10 strata). Each of the resultant 16 cells contained either two or three strata. A single stratum might include only one PSU (or only part of a PSU as, for example, New York City, which represented two strata) or several score PSU's.

To take account of the possible effect that the rate of population change between the 1950 and the 1960 Census might have had on health, the 10 strata within each region were further classified into four classes ranging from those with no increase to those with the greatest relative increase. Each such class contained two or three strata.

One PSU was then selected from each of the 40 strata. A controlled selection technique was used in which the probability of selection of a particular PSU was proportional to its 1960 population. In the controlled selection an attempt was also made to maximize the spread of the PSU's among the States. While not every one of the 64 cells in the 4x4x4 grid contributes

a PSU to the sample of 40 PSU's, the controlled selection technique ensured the sample's matching the marginal distributions in all three dimensions and being closely representative of all cross-classifications.

Generally, within a particular PSU, 20 ED's (census enumeration districts) were selected with the probability of selection of a particular ED proportional to its population in the age groups 5-9 years in the 1960 Census, which by 1963 roughly approximated the population in the target age group for Cycle II. A similar method was used for selecting one segment (cluster of households) in each ED. Each of the resultant 20 segments was either a bounded area or a cluster of households (or addresses). All of the children in the age range properly resident at the address visited were EC's (eligible children). Operational considerations made it necessary to reduce the number of prospective examinees at any one location to a maximum of 200. The EC's to be excluded for this reason from the SC (sample child) group were determined by systematic subsampling. If one of the sample children had a twin who was not a sample child, this other twin was brought in for examination, and while the results were recorded for use in a special substudy of twins, this twin was not included in the 7,119 children under the present analysis.

The total sample included 7,417 children 6-11 years of age of whom 96 percent were finally examined. These 7,119 examined children represented the roughly 24 million children in the United States who met the general criteria for inclusion in the sampling universe as of mid-1964.

All data presented in this publication are based on "weighted" observations. That is, data recorded for each sample child are inflated in the estimation process to characterize the larger universe of which the sample child is representative. The weights used in this inflation process are a product of the reciprocal of the probability of selecting the child, an adjustment for nonresponse cases, and a poststratified ratio adjustment which increases precision by bringing survey results into closer alignment with known U.S. population figures by color and sex for single years of age 6-11.

In the second cycle of the HES the sample was the result of three stages of selection—the single PSU

from each stratum, the 20 segments from each sample PSU, and the sample children from the eligible children. The probability of selecting an individual child is the product of the probabilities of selection at each stage.

Since the strata are roughly equal in population size and a nearly equal number of sample children were examined in each of the sample PSU's, the sample design is essentially self-weighting with respect to the target population; that is, each child 6-11 years old had about the same probability of being drawn into the sample.

The adjustment upward for nonresponse is intended to minimize the impact of nonresponse on final estimates by imputing to nonrespondents the characteristics of "similar" respondents. Here "similar" respondents were judged to be examined children in a sample PSU having the same age (in years) and sex as children not examined in that sample PSU.

The poststratified ratio adjustment used in the second cycle achieved most of the gains in precision which would have been attained if the sample had been drawn from a population stratified by age, color, and sex and made the final sample estimates of population agree exactly with independent controls prepared by the U.S. Bureau of the Census for the noninstitutional population of the United States as of August 1, 1964 (approximate midsurvey point) by color and sex for each single year of age 6-11. The weights of every responding sample child in each of the 24 age, color, and sex classes are adjusted upward or downward so that the weighted total within the class equals the independent population control.

A more detailed description of the sampling plan and estimation procedures is included in earlier reports of the *Vital and Health Statistics* series.^{6,7} Series 11, No. 1,⁸ describes the techniques used in Cycle I, which are similar to those of Cycle III.

Notes on Response Rates

There were 7,417 children aged 6-11 years selected for examination. Of these, 7,119 were actually examined, which made an overall response rate of just under 96 percent. The response rate by sex and 1-year age group is shown below.

Age	Boys	Girls
6-----	96.5	94.9
7-----	96.5	95.5
8-----	95.2	97.0
9-----	97.6	94.8
10-----	97.0	95.1
11-----	96.2	95.6
Total-----	96.5	95.5

NOTE: The list of references follows the text.

It can be seen that only at age 8 years was the response rate for girls better than that for boys.

A similar analysis can be done by age, race, and sex as shown below:

Age	Boys		Girls	
	Negro	White	Negro	White
6-----	97.7	96.3	97.3	94.7
7-----	97.5	96.3	98.9	94.8
8-----	97.5	95.0	99.1	96.5
9-----	98.7	97.4	100.0	93.9
10-----	98.5	96.8	97.5	94.7
11-----	98.8	95.8	98.8	95.0
Total-----	98.1	96.2	98.7	94.9

A striking difference in response is readily seen. Negro children responded better than their white counterparts at every age group, and 9-year-old Negro girls had an extraordinary 100-percent response rate.

Parameter and Variance Estimation

As each of the 7,119 sample children has an assigned statistical weight, all estimates of population parameters presented in HES publications are computed taking this weight into consideration. Thus, \bar{X} , the estimate of a population mean, " μ ," is computed as follows:

$$\bar{X} = \frac{\sum_{i=1}^n w_i X_i}{\sum w_i}$$
 where X_i is the observation or measurement taken on the i^{th} person and w_i is the statistical weight assigned to that person.

The HES has an extremely complex sampling plan, and obviously the estimation procedure is, by the very nature of the sample, complex as well. A method is required for estimating the reliability of findings which "reflects both the losses from clustering sample cases at two stages and the gains from stratification, ratio estimation, and poststratification."¹⁵

The method for estimating variances in the HES is the half-sample replication technique. The method was developed at the U.S. Bureau of the Census prior to 1957 and has at times been given limited use in the estimation of the reliability of results from the Current Population Survey. This half-sample replication technique is particularly well suited to the HES because the sample, although complex in design, is relatively small (7,119 cases) and is based on but 40 strata. This feature permitted the development of a variance estimation computer program which produces tables containing desired estimates of aggregates, means, or distributions, together with a table identical in format but with the estimated variances instead of the estimated statistics. The computations required by the method are simple, and the internal storage require-

ments are well within the limitation of the IBM 360-50 computer system utilized at the National Center for Health Statistics.

Variance estimates computed for this report were based on 20 balanced half-sample replications. A half sample was formed by choosing one sample PSU from each of 20 pairs of sample PSU's. The composition of the 20 half samples was determined by an orthogonal plan. To compute the variance of any statistic, this statistic is computed for each of the 20 half samples. Using the mean as an example, this is denoted \bar{X}_i . Then, the weighted mean of the entire, undivided sample (\bar{X}) is computed. The variance of the mean is the mean square deviation of each of the 20 half-sample means about the overall mean. Symbolically,

$$\text{Var.}(\bar{X}) = \frac{\sum_{i=1}^{20} (\bar{X}_i - \bar{X})^2}{20}$$

and the standard error of the mean is the square root of this. In a similar manner, the standard error of any statistic may be computed.

A detailed description of this replication process by Phillip J. McCarthy, Ph.D., has been published.³⁵

Standards of Reliability and Precision

All means, variances, and percentages appearing in this report met defined standards before they were considered acceptably precise and reliable.

The rule for reporting means and percentiles consisted of two basic criteria. The first criterion was that a sample size of at least five was required. If this first criterion was met, then the second criterion, that the coefficient of variation [i.e., the standard error of the mean divided by the mean (s_x/\bar{X})] was to be less than 25 percent, must have been demonstrated. Thus, if either the sample size was too small, or the variation with respect to the mean was too large, the estimate was considered neither precise nor reliable enough to meet the standards established for publication.

NOTE: The list of references follows the text.

Hypothesis Testing

Classically, to test the difference between two means (or, put differently, to test whether two samples could have been drawn from the same population), one could set up a test statistic which would utilize the means and standard errors of the means as computed from the samples. The statistic

$$z = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{s_{\bar{X}_1}^2 + s_{\bar{X}_2}^2}}$$

would then be compared to a table of normal deviates, to determine the probability of obtaining values of the test statistic as extreme or more extreme than that computed, if in fact the two population means were equal.

Because of the many breakdowns of the HES sample, innumerable tests of this nature could be performed. With each new test, the probability of rejecting a hypothesis incorrectly may be .05; but if 10 such tests are performed, the probability of making at least one mistake somewhere in those 10 tests is closer to 0.60. This last "overall error rate" will get increasingly large as the number of such tests increases. Therefore, while the data necessary to do z tests are provided in the tables of this report, no such tests were performed by the authors.

It was decided to place the greatest emphasis on a relationship remaining consistent over both sexes and all ages under consideration. In other words, to say, for instance, that "girls have buttock-knee lengths greater than boys for all ages between 6-11 years" has far more meaning and interpretability than to say "the mean buttock-knee length for 6-year-old girls is significantly greater than the corresponding mean for 6-year-old boys," as determined by a normal deviate. In these analyses *consistency*, rather than a statement about a succession of individual probability levels, is the factor considered most important in demonstrating relationship.

— 0 0 0 —

APPENDIX II

TECHNIQUES OF MEASUREMENT AND QUALITY CONTROL

Techniques of Measurement

Trained observers made all measurements, reading them to the nearest millimeter (tenth of a centimeter). All measurements were read aloud to a recorder, who repeated aloud each number back to the observer as it was recorded in the proper space on the record form.

This repetition served both as a doublecheck on the measurement and to reduce recording errors.

Measurements were performed in a regular sequence to minimize the number of position changes the child was required to make. The sequence is illustrated on the measurement recording form (figure 1). It should be noted that not all the measurements taken in

OBSERVER (6-7)		RECORDER	
CARD 06 COL. NO.	SITTING*	CARD 07 COL. NO.	STANDING (FLOOR)*
0-10	FOOT LENGTH	0-10	ACROMIAL DIAM.
11-13	FOOT BREADTH	11-13	ACROMION TO OLECRANON
14-17	KNEE HEIGHT	14-16	CHEST BREADTH 4TH ICS
18-21	POPLITEAL HEIGHT	17-19	CHEST DEPTH 4TH ICS
22-25	THIGH CLEARANCE	20-22	CRISTAL DIAM.
26-28	SEAT BREADTH	23-25	CHEST GIRTH
29-31	ELBOW-ELBOW BREADTH	26-28	WAIST GIRTH
32-35	SITTING HEIGHT-ERECT	29-31	HIP GIRTH
36-38	BUTTOCK-POPLIT LENGTH	32-34	R. UPPER ARM GIRTH
39-41	BUTTOCK-KNEE LENGTH	35-37	R. LOWER ARM GIRTH
42-44	ELBOW-WRIST LENGTH	SKIN FOLDS	
45-47	HAND LENGTH	38-40	R. UPPER ARM (MM)
48-50	HAND BREADTH	41-43	R. INFRASCAPULAR (MM)
STANDING (ON STEP)*		44-46	R. LAT. WRIST WALL (MM)
51-53	R. BICONSYLAR DIAM	WEIGHT (LBS)	
54-56	R. CALF GIRTH		
57-60	STANDING HEIGHT	END CARD 07	
ANTHRO. NO.			
61-62	COLS. 14-28	47-50	WEIGHT (LBS)
63-64	COLS. 29-31	79-80	END CARD 07
79-80	END CARD 06		

$\frac{1}{8}$ in cm

MEASUREMENTS NOT DONE OR SIDE VARIED—specify which and give reason

PHS-4811-3
REV. 7-64

SAMPLE NO (1-1)

Figure 1. Body measurement recording form.

the survey are reported in the present report. The three skinfolds, for example, have been analyzed and reported already, while other specific reports are still in preparation.

All of the technicians were experienced X-ray technicians who had been trained in anatomy and the identification of specific body landmarks. In addition, X-ray technicians, both by disposition and training, tend to work well with people and are skilled in giving the examinee verbal orders along with the necessary handling to achieve proper positioning.

Each technician received more than a month of intensive training before being considered minimally proficient in making body measurements. In this training, he became skilled with the equipment, the precise locations of the body at which the measurements were to be taken, and the technique of measurement itself. The major sources of measurement error are improper positioning of subject's body, improper selection of specific body landmarks, and improper application of instrument (for instance, not perpendicular when measuring a diameter or circumference, or improperly compressing the soft tissue over bony landmarks). Incorrect reading of the instrument (usually transposition of numbers) also occurs with discouraging frequency. When these errors were mostly overcome, the new technician's data were carefully compared with those of the other three technicians and the two supervisors before they were officially accepted as recorded data.

As was emphatically stated by Hertzberg when summarizing the Conference on Standardization of Anthropometric Techniques and Terminology in 1968,¹⁶ every effort must be made to insure accuracy of measurement and standardization of procedure if the data are to be useful. The preceding discussion sketches the chief procedures used to reduce both systematic and variable measurement error. As discussed in the lengthy subsequent section, "Quality Control and Estimation of Residual Measurement Error," the absolute amount of systematic error can never truly be known unless one agrees on the "perfect measurer with perfect equipment perfectly applied, etc." A good estimate of the residual variable measurement error can, however, be achieved by replicate examinations for both inter- and intra-observer variability.

In the subsequent pages, the equipment, measurements, and specific procedures used in the survey are described and illustrated. Next the quality control procedures which were used to monitor the body measurements are discussed extensively.

Equipment

The measuring equipment consisted of several anthropometers, small sliding calipers, steel tapes, and a measuring table with an adjustable footrest.

NOTE: The list of references follows the text.




Figure 11. Anthropometric instruments used in Health Examination Survey, United States, 1963-65; A: anthropometer; B: sliding caliper; C: steel tape.

The *anthropometer* (figure 11) was used to measure various body lengths, heights, and breadths. It is a rod consisting of four sections and two crossbars, or measuring arms. One of the crossbars is fixed, while the other is movable. The anthropometer is calibrated in centimeters and millimeters. It has two scales, one reading from the top down and the other from the bottom up. In this survey, a section of one anthropometer was fitted with a base for stabilizing purposes (to avoid tilting when making various height measurements), while another was fitted into the sliding backboard of the measuring table.

The *small sliding caliper* (figure 11) was used to measure hand length and breadth. It consists of a flat metal bar upon which a slide moves. One of the crossbars is fixed, while the other is movable. The small sliding calipers are calibrated in centimeters and millimeters.

The *steel tape* (figure 11) was used to measure various body circumferences. It is a flexible tape with a spring rewind and it is scaled in centimeters and millimeters on one side, in inches on the other.

The *measuring table* was such that it could accommodate children of varying sizes and proportions. It was equipped with an adjustable footrest in order to maintain a standardized position of the lower extrem-

ties during the measurement process. The surface of the table was also equipped with a measurement scale in centimeters and millimeters and with a sliding backboard at right angles to the scale.

Measuring Procedures and Definitions

Weight was measured on a Toledo self-balancing weight scale which mechanically printed the body weight directly onto a permanent record. It was recorded to the nearest 0.5 pounds.¹

Height was measured as the distance from the standing surface to the top of the head. The child was in stocking feet with feet together, back and heels against the upright bar of the height scale, head in the Frankfort plane (looking directly forward), and standing erectly ("standing up tall").¹

General position for sitting measurements. The child sat on the measuring table with the popliteal fossae at the front edge of the table. The footrest was adjusted so that the child sat with his knees and feet together, heels against the heel rests, feet at right angles to the lower legs, and lower legs at right angles to the thighs. Elbows were held at the sides with forearms at right angles, hands open, and palms facing each other, or with hands resting on knees. Arm positions were adjusted when necessary to meet the requirements of specific measurements.

General position for standing measurements. The child stood erectly with the head oriented in the Frankfort plane, i.e., looking directly ahead and feet together. Arms were held relaxed at the sides. Postural adjustments were made to meet the requirements of specific measurements.

Sitting height was measured as the vertical distance from the sitting surface to the top of the head. With the subject seated as described above, the backboard on the measuring table was brought up firmly against the buttocks. The movable arm of the anthropometer (which was inserted into the backboard) was brought down firmly to the midline of the top of the head.

Buttock-knee length was measured as the distance from the rearmost projection of the buttock to the front of the right kneecap. With the subject seated as previously described, the fixed crossbar of the anthropometer was placed in *light contact* with the rearmost projection of the buttock, and the movable crossbar was brought into *light contact* with the front surface of the right kneecap (patella).

Popliteal height was measured as the distance from the surface of the footrest to the underside of the right knee. With the subject seated as previously described, the anthropometer with its attached base was placed on the footrest adjacent to the right foot and the movable arm was brought to the level of the table surface on which the child was seated. This is the level at which

the under side of the right knee (tendon of the biceps femoris muscle) comes into contact with the table surface.

Foot length was measured as the distance from the back of the right heel to the tip of the longest toe. With the child seated as previously described, the fixed arm of the anthropometer was *lightly* applied behind the heel with the rod parallel to the long axis of the foot. The movable bar of the anthropometer was then brought into *light contact* with the tip of the longest toe.

Acromion-olecranon length was measured as the distance from the acromial process of the right scapula (outer point of the shoulder) to the olecranon process of the ulna (elbow). With the subject standing, right arm at his side and elbow bent at a 90-degree angle, the fixed crossbar of the anthropometer was placed *firmly* at the right acromial process and the movable crossbar was brought into *firm contact* with the olecranon process (tip of the elbow).

Elbow-wrist length was measured as the distance from the olecranon process (elbow) to the distal end of the styloid process of the ulna. With the subject seated as previously described but with palm facing downward, the fixed arm of the anthropometer was *firmly* placed at the olecranon process (tip of the elbow) and the movable arm was *firmly* placed at the distal end of the styloid process of the ulna.

Hand length was measured as the distance from the wrist (midpoint of most distal crease or groove) to the tip of the middle finger. With the right hand fully extended, palm up and thumb straight but relaxed, the fixed end of the sliding caliper was placed at the midpoint of the distal crease at the wrist (located by having the child flex the hand at the wrist), and the movable crossbar of the caliper was placed in *light contact* with the distal tip of the middle finger.

Biacromial breadth was measured as the maximum distance between the right and left acromial processes of the scapula. With the subject standing and the observer standing behind him, the fixed arm of the anthropometer was placed at the most lateral point of the left acromial process and the movable bar brought to the most lateral point of the right acromial process. The measurement was made with *firm contact*.

Bicristal breadth was measured as the distance between the most lateral points of the iliac crests. With the subject standing and the observer standing behind him, the crossbars of the upper segment of the anthropometer were brought into *firm contact* with the edges of the iliac crests on each side. This measurement was made with *firm pressure*.

Chest breadth was measured as the breadth of the rib cage under *firm pressure*. With the subject standing and breathing normally, the fixed crossbar of the anthropometer was applied firmly at one side of the rib cage and the movable crossbar was applied firmly to the other side at the level of the nipples. The crossbars were angled slightly downward to avoid slipping

NOTE. The list of references follows the text.












Figure III. Schematic illustration of anthropometric dimensions taken on children aged 6-11 years in the Health Examination Survey, United States, 1963-65.













Figure III. Schematic illustration of anthropometric dimensions taken on children aged 6-11 years in the Health Examination Survey, United States, 1963-65--Con.

into the spaces between the ribs. At all times the rod of the anthropometer was parallel to the floor. In older girls who had a noticeable degree of breast development, the level of the junction of the fourth rib with the sternum was used as the measurement landmark.

Chest depth was measured as the distance or depth from the front to the back of the rib cage under *firm pressure* during normal breathing. With the subject in the same position as for the chest breadth measurement and with the observer approaching the child from the right side, the fixed arm of the anthropometer was applied firmly to the back of the chest and the movable arm was applied firmly to the sternum at the level of the nipples. At all times the measuring instrument was parallel to the floor. In older girls who had a noticeable degree of breast development, the level of the junction of the fourth rib with the sternum was used as the measurement level.

Bicondylar breadth of the femur was measured as the maximum width between the condyles of the right femur. With the subject standing, the fixed bar of the anthropometer was placed firmly on the medial condyle and the movable bar brought to the lateral condyle with *firm pressure*.

Upper arm girth was measured at the level midway between the acromial and olecranon processes of the right arm. The midlevel was located while the subject held the forearm at a right angle to the upper arm. The measurement was made while the right arm hung loosely, with the tape horizontal and in contact with the skin without deforming the skin contours, i.e., without compressing the underlying tissues. Note that this measurement was made at the same level as the triceps skinfold measurement.

Lower arm girth was measured as the maximum circumference of the right forearm just below the elbow joint. The girth was measured just below the elbow at the widest part of the forearm while the arm hung loosely. The tape was applied horizontally in contact with the skin without deforming the skin contours.

Calf girth was measured as the maximum circumference of the right calf at right angles to the long axis of the leg. With the subject standing, legs several inches apart and weight equally distributed on both feet, the girth was measured at the level of maximum circumference. The tape was placed in contact with the skin without depressing the skin contours.

Chest girth was measured as the circumference of the chest during normal breathing at the level of the fourth intercostal space. With the subject standing as for the measurement of chest breadth and depth, the steel tape was applied *firmly but without depressing the skin*. Special care was taken to make certain that the tape was horizontal.

Waist girth was measured as the circumference of the waist, abdomen relaxed, at the level midway between the lower edge of the ribs and the iliac crests. With the subject standing and breathing normally, the

steel tape was applied *firmly but without depressing the skin*. Special care was taken to make certain the tape was horizontal.

Hip girth was measured as the circumference of the hips at the level of the greater trochanters (the widest bony part of the hips). With the child standing, feet together, the steel tape was applied *firmly* to compensate for clothing (this girth was measured over shorts). Special care was taken to make certain the tape was horizontal.

Each dimension measured is schematically illustrated in figure III.

Derived Measurements

Subischial length was obtained by subtracting sitting height from standing height. It provides an estimate of the length of the lower extremities.

Estimated mid-arm muscle circumference was derived from the upper arm circumference and the triceps skinfold, both of which were measured at the same level of the arm midway between the acromial and olecranon processes. The arm in cross section consists of skin, subcutaneous fat, muscle, and bone. The skinfold measurement is a double fold of skin and underlying subcutaneous fat. If it is assumed that the upper arm is a cylinder and the principles of circle geometry are applied, the arm circumference can be corrected for the thickness of the triceps skinfold, leaving an estimated mid-arm muscle-bone circumference or, for the sake of simplicity, estimated mid-arm muscle circumference. Thus,

$$EMC = AC - \pi S_t$$

where *EMC* is the estimated mid-arm muscle circumference in cm., *AC* is the upper arm circumference in cm., and *S_t* is the triceps skinfold in cm. Since the estimated mid-arm muscle circumference for large numbers of children is generally calculated via computers and skinfold thicknesses are usually measured in millimeters, the following formula was used:

$$EMC \text{ (cm)} = \pi \left[\frac{AC \text{ (cm)}}{\pi} - \frac{S_t \text{ (mm)}}{\sqrt{10}} \right]$$

Indexes and Ratios

Ponderal index provides an approximation of physique or body build on a linearity-laterality continuum. It was obtained as follows: standing height in inches divided by the cube root of body weight in pounds (height/ $\sqrt[3]{\text{weight}}$).

Sitting height/stature ratio indicates the relative contribution of sitting height (i.e., head, neck, and trunk) to total stature. It was obtained as follows: sitting height divided by standing height times 100. This ratio is expressed as a percentage.

Biacromial breadth, bicristal breadth ratio indicates the relative proportion of shoulder width to hip width. It was obtained as follows. biacromial breadth divided by bicristal breadth times 100. This ratio is expressed as a percentage.

Quality Control and Estimation of Residual Measurement Error⁹

Monitoring Systems

Despite efforts to reduce measurement errors, residual errors of a magnitude large enough to warrant concern occur with some regularity in any anthropometric survey. There is, therefore, a real and urgent need to have a system whereby these residual errors can be monitored. The concept of quality control is based on the desire to obtain end products of a specified quality. Thus, one of the main purposes of a monitoring system is to indicate whether the measurements produced by a certain measurement process have attained the desired quality. A second major purpose is to make possible quantitative summary descriptions of residual measurement errors to aid in the interpretation of survey data.

Perhaps the most direct monitoring system used in the Health Examination Survey was *observation of the measurement process* as it was being applied to an examinee. Medical, dental, and psychological advisors from HES and other advisors and consultants regularly visited the examination center to observe examination procedures and to retrain examiners if necessary. A good example of how routine observation was used as a monitoring system can be found in the taking of body measurements. One member of the examining team, a trained anthropometrist, acted as a recorder and aided in positioning of the examinees, while he was additionally responsible for observing and correcting any errors in measurement technique.

As a careful and thoughtful quality control program tends to be an evolving process, the most extensive systematic monitoring of body measurements performed in any of the cycles of the Health Examination Survey was achieved in Cycle III (youths 12-17 years, data collection 1966-70). The formal system of replicate examinations which was finally instituted in Cycle III is described later in this appendix along with a discussion of its applicability to Cycle II.

Replicate measurements are useful for a variety of reasons, e.g., as a means of increasing precision of individual measurement estimates, as a training technique, and as a monitoring system which includes the objective of final evaluation of measurement errors. These three objectives are compatible, and replicate

data collected primarily for one of them often indirectly, if not directly, accomplish one or both of the remaining two. For this reason replicate data are most often collected with a combination of these objectives in mind. The single most important source of replicate data in Cycle III was the replicate examination procedure, in which approximately 5 percent of the regular examinees were returned to the examination center for a second complete examination (except for drawing blood and taking X-rays).

Biases and Controls in Replicate Measurements

A major source of uncertainty in estimates derived from replicate measurements is inability to make the replicate measurement under precisely the same conditions and in the same manner as the original measurement. This uncertainty is difficult to evaluate, and most attempts to do so are restricted to subjective statements concerning the direction and/or size of the bias and the need for concern in the analysis of data.

Several policies regarding Cycle III replicate examinations were designed specifically to obtain measurements under the same conditions and in the same manner as the initial (original) exam. Replicate examinations were not conducted at a specific time. Whenever possible, they were interspersed among the regular examinations. An original examination was given priority over a replicate examination in that none would be scheduled if it occupied time needed for a regular examination. There was often space to interject replicate examinations in the schedule without interfering with regular examinations, but this priority, plus the fact that replicates were drawn from those previously examined, increased the likelihood that a replicate examination would be scheduled toward the end of the examination period. Nevertheless, the attempt to space replicate examinations throughout the regular schedule was a valuable policy in that the interspacing of replicate and original examinations created an atmosphere more conducive to both examinations being conducted in essentially the same manner.

The examiners were informed of the purpose and importance of the reexaminations. It was emphasized that they should not vary their procedures on a replicate examination or in any way try to collect "better" data than they would normally. Thereafter, instructions on the conduct of replicate examinations were not given greater emphasis than any other instruction because overemphasizing "sameness" might have created more bias than it would have eliminated.

At the time of the original examination neither the observer nor the examinee knew whether or not the examinee would be returned for a replicate examination. During the replicate examination, observers were not specifically informed that an examinee was a replicate, although no attempt was made to conceal this

⁹This section is in part based upon Schaible's lucid and systematic discussion of quality control and error estimation in the HES, Series 2, Number 44.³⁷

fact since in an examination as lengthy as that given in HES the examinee would undoubtedly be remembered by several, if not all, examiners. Even though an examinee might be remembered, it was extremely unlikely that an examiner would remember a specific measurement after a time lapse of 2 or 3 weeks. Some bias might be introduced by the examiner's knowledge of the replicate status of an examinee, but generally this bias would seem quite small when compared to the measurement error and in some cases to the biases associated with the knowledge and familiarity gained by the examinee during the original examination. Examinee bias can be important, especially when a response is elicited or when the true value of the measurement has changed because of a time lapse. Since the time lapse was usually 2 or 3 weeks, some appreciable changes might occur in certain measurements such as weight. However, for most of the data collected, the actual change over this short period of time can only be very small and this effect may usually be neglected. Previous experience is much more likely to affect the true replicability of psychological tests and those physiologic tests requiring high levels of subject participation (such as the treadmill and spirometry), with procedures in which the subject is passive and very little learning is involved, such as EKG and body measurements, the effect of previous experience is almost zero.

In Cycle III replicate data were obtained on approximately 70 percent of those selected for such examinations. One explanation for this low rate is that persuasion and followup efforts were not as intensive as for regular examinees. This is partially because regular examinees were given priority if interviewer or examination time was limited. There also appeared to be an increased frequency of objection to returning for a second examination, as demonstrated in the most frequent reasons for refusal. "One time is enough" and "I can't miss school again."

Selection of Replicate Examinees

The selection of Cycle III youths for replicate examinations was random within certain restrictions imposed by practical considerations. One restriction was that replicates were selected only from those examined during the first week and a half of the approximately 3½ weeks of examinations at any one location. This time period was chosen to facilitate the interspersing of replicate examinations with originals in the examining schedule without interfering with the time allotted for original examinations and without scheduling additional time to accommodate replicates. In a voluntary survey it is obviously impossible to follow a statistically random process in scheduling subjects, so those scheduled during the first week and a half are not, in the strict sense, a random sample of all those scheduled, though they may be randomly distributed for those features which are significant.

Evidence that replicates might be considered "representative" is found in the fact that youths of certain ages, locations, incomes, etc., were not routinely more likely to be scheduled during any particular segment of the examination schedule. However, the availability and desires of the subjects do influence the composition of the replicate sample. For instance, an examinee whose participation in an original examination was achieved only after repeated contacts by survey personnel was less likely to have been included in a replicate examination since it is unlikely that he would have received an original examination during the first week and a half. The schedule of locations, time of year, sequence of examinations, and other related factors which might make subjects more or less readily available show no obvious discriminatory effect in the selection of replicate examinees. After examining these and other relatively minor considerations, there appears to be no reason to believe that subjects scheduled and examined during the first part of a stand differed from those scheduled and examined during the latter portion with respect to the data gathered.

Another restriction on complete randomness in the selection of youths for replicate examinations was the exclusion of those examinees living somewhere "geographically inconvenient" to the examination center. "Geographically inconvenient" was arbitrarily defined as a distance of 30 miles or more although exceptions were sometimes allowed if conditions dictated. A primary consideration in choosing a site for the examination center was the centrality of the location in relation to the sample segments (a segment is a cluster of households). Since segments were drawn with probability proportional to population, most segments were in relatively populated areas, so the examination center was also in or adjacent to a relatively populated area. Therefore, the subjects deleted by this 30-mile restriction usually resided in relatively less populated areas. Thus this restriction may create a bias in replicate data if, in fact, characteristics differed with population density. Even if differences did exist, the total effect of this restraint would not be great since it excluded only approximately 10 percent of the eligible examinees. There were other minor restrictions of a medical and operational nature imposed on the complete randomness of the replicate sample. They were not, however, readily associated with large differences since at most only 1-2 percent of the eligible examinees were deleted for these reasons.

Since the purpose of replicate examinations is to give information about errors, the matter of concern between those excluded and those eligible for selection is not possible differences in measurement values but possible differences in the errors associated with measurements as shown by the discrepancy between two measurements on the same subject. For example, measurements may vary markedly by some demographic

classification, but this is not as relevant as the question of whether or not the measurement errors vary by this classification. A similar differential in the *active* and *passive* participation of subjects (e.g., spirometry versus body measurements) is assumed to operate here also, but in a different way. That is, it must be assumed that the most cooperative subjects, by and large, self-select themselves, and that their scores are true estimates of the variable being tested. It is thus likely that their test-retest difference would be *smaller*. On the other hand, although subjects did influence measurement errors, it should also be noted that the environment, procedures, and examiners were also highly influential in the final measurement. Consideration of these additional influences causes a completely random selection of subjects to be of somewhat less concern.

The Analysis of Replicate Data on Body Measurements

Although a variety of monitoring systems for body measurements were in effect in HES from the beginning of Cycle I, it was not until Cycle III that a formal system was instituted of recalling approximately 5 percent of the subjects already examined for a replicate examination. However, during Cycle II, which is the concern of the present report, several "in-field" attempts at assessing replicate body measurements were made. These included the following:

- 1) Several formal training sessions were held in which the examining technicians performed duplicate sets of measurements on a small group of subjects producing data for immediate examination of intra- and inter-examiner differences.
- 2) The two Cycle II examining caravans converged from the east and west for a measurement stand in the Greater Chicago area. After scheduled examinations were completed in the normal manner, one of the caravans (Caravan I) re-examined (for our purposes, remeasured) approximately 50 children who had been initially examined by the staff of the other caravan (Caravan II), and vice versa. This operation permitted the technicians an "in-field" examination and discussion of the replicate measurements.
- 3) Finally, a total of five intensive 2-day sessions were conducted by the supervisors in the field examination centers.

No formal, detailed analysis of the data in the statistical sense was carried out, primarily because the above attempts were more training than evaluation sessions.

In Cycle III, on the other hand, a systematic attempt at analysis of replicate body measurements was made. A total of 301 replicate examinations from Cycle III were collected and subjected to an extensive analysis of

intra- and inter-examiner variation in body measurements, i.e., variation within the same observer and variation between different observers. Since the *conditions* under which the body measurements were made were *essentially identical* in Cycles II and III, there is reason to believe that the results of the quantitative assessment of replicate measurements of data collected in Cycle III can be effectively applied to Cycle II. In other words, should the analysis indicate a reasonably good degree of accuracy within and between examiners in Cycle III, it can be safely assumed that a similar degree of measurement accuracy was apparent in Cycle II.

Although the anthropometry in Cycles II and III was very similar, there were four relatively minor differences. First, the children in Cycle II were younger and smaller in size. (There is, however, no reason to assume that the *relative measurement error* is different for younger and smaller individuals.) Second, four of the human engineering measurements taken in Cycle II were not measured in Cycle III; they were replaced by several segmental length measurements of greater biological significance and interest. Third, a total of 11 technicians made measurements during Cycle III, but in Cycle II, the same four technicians participated in equal degrees throughout the entire cycle. Fourth, a more elaborate, systematic collection of replicate data with greater numbers of subjects was utilized in Cycle III. Other factors—the instruments and their calibration, techniques of measurement, methods of training, selection of technicians, examination environment, and the chief medical examiner and the physical anthropologic consultant—were the same. It should be noted further that two of the four technicians who participated in Cycle II of the HES continued for several years into Cycle III. In summary, the only significant differences in quality control considerations for body measurements between Cycles II and III were the addition of the systematic collection of replicate data and the use of a greater number of technicians in Cycle III. The authors have concluded that these two differences approximately counterbalance one another, resulting in equivalent degrees of measurement variation.

Cycle III Systematic Replicate Procedure

Body measurements were taken on 6,768 youths and these data comprise the HES findings. Replicate body measurements were obtained on 301 youths at 30 of the 40 locations (or stands) visited throughout the United States. That is, an average of 10 youths were reexamined at each stand. Of the 301 youths, 224 were reexamined by a technician other than the one initially measuring the youth, while the remaining 77 were reexamined by the same technician. Altogether during the 4 years, 11 technicians participated in replicate measurements for this phase of the quality control program.

It is of interest to ascertain whether each of the examiners had a representative number of replicate measurement sessions with respect to the number of examina-

Table I. Percentage of regular and replicate examinations performed by each technician

Technician number	Percentage of regular Cycle III examinations	Replicate examinations	
		Percentage of intra-examinations	Percentage of inter-examinations
1	0.8	1.3	0.9
2	13.4	2.7	10.2
3	22.8	21.3	21.4
4	6.1	4.0	2.7
5	13.5	10.7	16.7
6	6.1	5.3	6.5
7	3.7	5.3	4.9
8	15.1	24.0	16.4
9	11.3	16.0	13.3
10	3.0	2.7	3.6
11	4.1	6.7	3.6

tions he performed during the survey. It should be carefully noted that it was not possible to insure that each technician had equal chances to measure replicate examinees since the length of time technicians were associated with the survey team varied. Table I presents the percentages of total examinations, intra-examiner replicates, and inter-examiner replicates participated in by each of the 11 technicians.

Table I clearly indicates some possible sources of bias which may affect the analysis of replicate data. For example, assume technician No. 9 was able to replicate his own measurements well but his readings were very different from the other examiners. Obviously, his results would be overrepresented in the replicate analysis since he examined only 11.3 percent of all youths in the actual survey but did 16 percent of the intra-examiner replicate examinations and 13.3 percent of the inter-examiner replicate examinations. Because of this technician's overrepresentation, the distribution of intra-examiner differences would cluster closer to zero than it really should have since this examiner self-replicated well. On the other hand, the inter-examiner distribution of differences would be considerably more skewed than it should have been since this technician did not agree well with the other technicians' measurements. Similar discrepancies are obvious for other technicians. An example of an opposite effect to that cited above is technician No. 2, who did only 2.7 percent of the intra-examiner replicate measurements and 10.2 percent of the inter-examiner replicate measurements, but did 13.4 percent of all examinations in Cycle III.

Thus, the various combinations of observers for the inter-examiner replicates and the proportions of intra-examiner replicates were not controlled so as to be balanced among the observers. In the survey proper the examinations were similarly not proportionately distributed among the observers, since the length of time

the various technicians were associated with the survey varied.

The foregoing indicates that the distribution of numbers of replicate examinations done by each technician is not the same as the distribution of the total number of survey examinations done by each in Cycle III. This represents one of the inherent problems of the present replicate data and limits to some extent implications to the survey as a whole. Nevertheless, the reader should be aware of the many problems confronting those who conduct large-scale health surveys, and in this context, the present systematic approach to the collection of replicate body measurement data is adequate.

Results of the Replicate Analysis

The absolute differences between the first and second measurements of the same child were computed for each dimension measured during Cycle III. The present analysis concerns itself with all body measurements except skinfold thicknesses, which have been reported separately with the results of the analysis of skinfold data.³

A distribution of absolute differences was compiled for each body measurement for the intra- and inter-examiner groups separately. The median and mode for each body measurement were extracted from the distribution of absolute differences. The mean absolute difference (\bar{X}_d) was computed by summing the differences and dividing by either 77 or 224, depending on which group (intra- and inter-examiner, respectively) was being considered.

A widely used measure of replicability is the statistic σ_e , the "technical error of measurement." It is defined as $\sigma_e = \sqrt{\sum d^2 / 2n}$, the square root of the sum of the squared differences of replicates divided by twice the number of pairs. This statistic assumes that the distribution of replicate differences is normal and that errors of all pairs can be pooled.

Since squaring a technical error of measurement

NOTE. The list of references follows the text

yields a variance, and since the ratio of two variances has the *F* distribution, a very simple test exists for comparing intra- and inter-examiner replicability. In table II the final column gives, for each variable, the *F* ratio (i.e., the ratio of the squares of the inter-examiner σ_e to the intra-examiner σ_i). As will be noted later, in three instances the variance for the intra-examiner group was larger and in these cases the ratios were reversed. A significant *F* statistic indicates the presence of a "technician-effect" or some characteristic which makes a particular measurement more easily replicated by the same technician than by another.

The coefficient of variation (CV), σ_e/\bar{X} , the technical error of measurement divided by the overall mean (the mean of all subjects) for the particular variable under study, was also calculated. The coefficient of variation is a measure of relative variability, i.e., variation in

replicability relative to the overall magnitude of the measure.

In the context of the present analysis, great care must be used in dealing with this statistic. It is not a coefficient of variation in the traditional sense since the numerator contains a measure of dispersion of differences (between replicates) whereas the denominator contains a mean—not a mean difference but a mean magnitude of the measurement taken.

The value of this statistic lies in its adjustment of the technical error by the magnitude of the original measurement. It attempts to answer the argument that replicability is likely to be much better for a variable of small magnitude than for one of great magnitude. As will be expanded later, dividing by the mean measurement may overadjust for such biases.

In the presentation of results of the replicate ob-

Table II. Results of intra-examiner and inter-examiner replicate analysis

Measuring device and dimension measured	Intra-examiner results					Inter-examiner results					<i>F</i> value
	\bar{X}_d	Median	Mode	σ_e	CV	\bar{X}_d	Median	Mode	σ_e	CV	
Automated recording											
Height-----	0.549	0.5	0.1	0.494	0.302	0.563	0.4	0.1	0.681	0.417	1.90
Weight-----	1.325	1.0	1.0	1.173	2.119	1.335	1.0	1.0	1.228	2.218	1.10
Anthropometer—height measurement											
Standing											
Cervicale height-----	0.714	0.6	0.1	0.692	0.500	1.054	0.9	0.5	0.953	0.689	1.90
Acromial height-----	0.752	0.6	0.2	0.795	0.601	0.875	0.75	0.1	0.891	0.673	1.26
Radial height-----	0.890	0.7	0.1	1.063	1.044	0.916	0.7	0.2	0.949	0.932	1.25
Styilion height-----	1.114	0.7	0.1	1.424	1.819	1.032	0.8	0.5 0.7	1.010	1.290	1.99
Iliac crest height-----	0.700	0.6	0.3	0.646	0.644	1.134	0.9	0.3	1.059	1.055	2.69
Trochanteric height-----	1.413	1.0	0.6	1.466	1.672	1.600	1.3	0.1	1.510	1.722	1.06
Tibial height-----	0.613	0.5	0.1	0.565	1.229	0.769	0.6	0.3	0.719	0.564	1.62
Sphyrion height-----	0.266	0.2	0.1	0.247	3.815	0.380	0.3	0.1	0.343	5.298	1.93
Sitting											
Sitting height-----	0.578	0.4	0.2	0.535	0.631	0.767	0.7	0.2	0.705	0.832	1.74
Thigh clearance-----	0.495	0.4	0.2	0.439	2.853	0.595	0.5	0.2	0.544	3.535	1.54
Anthropometer—length and breadth measurement											
Foot measurements											
Foot length-----	0.238	0.2	0.1	0.264	1.087	0.296	0.2	0.2	0.524	2.158	3.94
Foot breadth-----	0.138	0.1	0.1	0.122	1.329	0.226	0.2	0.1	0.202	2.200	2.74
Across bony landmarks on torso											
Biacromial breadth-----	0.553	0.4	0.1	0.544	1.529	0.807	0.5	0.1	0.915	2.571	2.54
Bicristal breadth-----	0.775	0.6	0.1	0.711	2.926	1.590	1.1	0.1	1.545	6.358	4.72
Bitrochanteric breadth-----	0.552	0.4	0.1	0.523	1.778	1.760	0.5	0.1	0.836	2.843	2.56
Across torso											
Seat breadth-----	0.610	0.4	0.3	0.921	2.835	0.909	0.7	0.8	0.993	3.057	1.16
Elbow-elbow breadth-----	1.104	0.8	0.8	1.131	3.415	1.725	1.2	0.3	1.346	4.065	1.42
Sliding caliper											
Knee breadth-----	0.112	0.1	0.1	0.106	1.165	0.183	0.1	0.1	0.244	2.683	5.30
Elbow breadth-----	0.105	0.1	0.1	0.117	1.799	0.152	0.1	0.1	0.154	2.368	1.73
Ankle breadth-----	0.097	0.1	0.1	0.092	1.367	0.186	0.2	0.1	0.171	2.540	3.45
Wrist breadth-----	0.108	0.1	0.1	0.115	2.208	0.150	0.1	0.1	0.139	2.669	1.46
Spreading caliper											
Bizygomatic breadth-----	0.075	0.1	0.0	0.076	0.589	0.158	0.1	0.1	0.162	1.255	4.54
Bigonial breadth-----	0.147	0.1	0.1	0.186	1.575	0.295	0.2	0.2	0.272	2.746	3.04
Steel tape											
Torso girths											
Chest girth-----	1.297	1.1	0.8	1.096	1.362	1.970	1.6	0.6	1.816	2.256	2.75
Waist girth-----	1.470	1.2	0.1	1.308	1.927	1.621	1.3	0.6	1.561	2.300	1.42
Hip girth-----	1.168	0.9	0.3	1.234	1.398	1.514	1.3	1.7	1.375	1.558	1.24
Extremity circumference											
Upper arm girth-----	0.339	0.3	0.1	0.347	1.358	0.458	0.4	0.3	0.425	1.664	1.50
Forearm girth-----	0.319	0.2	0.2	0.304	1.281	0.404	0.3	0.2	0.582	2.453	3.67
Calf girth-----	0.491	0.3	0.2	0.872	2.588	0.353	0.3	0.2	0.340	1.009	6.58

NOTE: For definition of symbols, see above.

ervation analysis, data were grouped according to the measuring instrument used in order to facilitate comparison since there is the possibility that differences between or within certain examiners might be peculiar to the particular measuring device used. First, height and weight were treated as a single group because they were machine-recorded. Note, however, that height measurement can be affected by variations in positioning. The second group was comprised of various height measurements which include the distance from the standing or sitting surface to the specific landmark. In most instances, the anthropometer was used to its full extent; nevertheless, as the landmarks approached the leg and ankle, the measuring distance was shorter. The third group of measurements included those made with the upper portion of the anthropometer. These measurements were made with the fixed arm of the anthropometer at one landmark while the free end was moved to the other landmark, which defined the measurement. This group included two foot measurements, three bony breadth measurements across the torso which required firm pressure, human engineering measurements which required light surface contact of the anthropometer. The fourth group included those made with a small sliding caliper. As a group these measurements represent the distance across a single bone or two bones at specific extremity joints. Compared to the height measurements mentioned above, the distance traversed by these measurements is rather small. The fifth group comprised only two facial breadths made with a spreading caliper. The sixth group consisted of measurements made with a steel tape and included six circumferences, three on the torso and three on the extremities.

Clearly body weight differs from all other values here since it was measured to the nearest half pound, while all others were measured to the nearest tenth of a centimeter, i.e., the nearest millimeter. Body weight is the only variable in which there is no chance of either intra- or inter-observer error. All weights were taken on a Toledo self-balancing scale which mechanically printed the child's weight directly onto the permanent record. It was not even important that the technician position the examinee rigidly, which was a significant factor in other measurements, for example, height. Any variability evident in replicate readings would thus be due to a gain or loss of body weight by the subject between examination sessions. Note that the F ratio for body weight was not significant, thus underlining the lack of technician effect in obtaining this measurement.

There were a total of 77 intra-examiner replications, i.e., the same technician re-examining the subject on two different occasions, and 224 inter-examiner replications, i.e., two different technicians doing the initial examination and replicate examination respectively, performed during Cycle III. Intra-examiner and inter-examiner results are presented separately in

table II, and all analyses were done within the group under consideration.

Taking the data in table II as a whole, the technical error of measurement was, with three exceptions, consistently less within examiners than between examiners. This was not entirely unexpected, for experience indicated greater intra-examiner consistency, i.e., there was greater consistency within the same technician than between different technicians. The three exceptions were radial height, stylium height, and calf circumference. Since each value was squared in calculating the technical error of measurement, this statistic can be greatly distorted by one or two highly divergent replicate values. That seems to be the case with these three divergent values.

Results of the variance analyses indicated that 25 of 31 F ratios were significant at the .05 level (or conversely, only 6 of 31 F ratios were not significant at the .05 level). Thus, in 25 measurements, intra-examiner differences were significantly smaller than inter-examiner differences. On the surface, such a tendency in the results might appear discouraging. However, such a tendency might function to eliminate or reduce systematic bias in large-scale surveys by eliminating or reducing the effects of individual idiosyncrasies (biases) of individual examiners.

Of 29 out of 31 measurements, the mean differences for intra-examiner observations were less than those for inter-examiner observations. These results were in the same general direction as those reported above for the technical error of measurement. The two measurements in which intra-examiner mean differences were the greater of the two were stylium height and calf circumference, both of which, as indicated above, had discrepant replicate readings which functioned to inflate the intra-examiner mean differences.

The median represents the midpoint of the distribution, i.e., 50 percent of the cases in the distribution are above and 50 percent are below this point. As such, it is not affected by the extremes of isolated discrepant values, as is the technical error of measurement. An examination of the median differences between replicate readings on an intra- and inter-examiner basis indicated eight instances in which the median differences between replicate measurements were identical within and between examiners. In 22 instances, median differences were less within examiners than between examiners, while in one instance the median difference was less between examiners than within examiners. In this last mentioned case, the difference between medians was only 0.1 cm. Thus, these observations are in general agreement with those indicated by comparison of σ and \bar{x} .

The magnitude of the differences between medians of replicate readings within and between examiners was only 0.1 cm. for 13 measurements, 0.15 cm. for one measurement, 0.3 cm. for five measurements, 0.4 cm.

for two measurements, and 0.5 cm. for two measurements. Incorporating the eight measurements in which median differences for replicate readings were identical within and between examiners with the above distribution indicated, that in 22 of the 31 measurements the difference in median differences of replicate readings within and between examiners was 0.15 cm. or less. This indicates a reasonable degree of consistency in the replicate measurements. It does not, however, consider the magnitude of the actual differences between replicate readings by the same observer, and by different observers.

Before going into a discussion of specific groups of measurements, the limitations of the technical error of measurement and the coefficient of variation should again be noted. As indicated earlier, the σ is generally an important and revealing statistic. By itself, however, it can be somewhat misleading at times. Consider, for example, the variables of standing height and knee breadth in table II for the intra-examiner group of data. Just considering σ would lead one to believe that knee breadth is a much better replicated measurement than is standing height since the variation for knee breadth is markedly smaller. It should be carefully noted, however, that the magnitude of standing height is far greater than that of knee breadth, and the margin of error is far greater for the greater measurement. To adjust for this factor, the coefficient of variation (σ/X) can be used. Examination of the coefficients of variation for these two variables indicates that standing height is more closely duplicated by the same examiner than is knee breadth.

On the other hand, coefficients of variation must be used with great caution. To divide σ for standing height by the entire mean for standing height is a bit drastic. For example, if an individual is 172 cm. tall, repeated measurements cannot vary by the whole 172 cm. Even if a technician makes a markedly discrepant replicate measurement of 10 cm., for example, this represents only 5.8 percent of the total height measurement. On the other hand, an error of 1.0 cm. for knee breadth, which for the sake of example is assumed to be 12.0 cm., represents 8.3 percent of the measurement. What is being suggested here is that there is noway errors of sufficiently large magnitude can be made for large measurements (of the order, say, of 170 cm. for height). Thus, to divide σ by the full mean for the particular measurement distorts the reality of the situation. This is why it is best to compare coefficients of variation within variables measured by the same instrument and within variables of about equal magnitudes.

Results of the replicate analysis for specific measurements and/or groups of measurements are now considered. As noted earlier, the data were grouped according to measurement instrument used.

Although body weight showed some variation within and between observers, the F ratio was not significant, indicating that all observers did comparable jobs in

measuring this variable. It should be noted, however, that there was no chance for individual idiosyncrasies of a given observer to affect the body weight measurement. All weights were taken on a Toledo self-balancing scale which mechanically printed the weight directly onto the child's permanent record. Hence the variation between observation sessions is due to the weight gain or loss occurring during the time lapse. Mean differences for body weight within and between examiners are well within the range of variation associated with diurnal changes in body weight.

As a group, measurements made with the sliding caliper had a high degree of replicability. This category included two measurements across single bones, i.e., knee breadth across the condyles of the femur and elbow breadth across the epicondyles of the humerus; and two measurements across two bones, i.e., ankle breadth across the distal aspects of the tibia and fibula and wrist breadth across the distal aspects of the radius and ulna. As a group the mean, median, and modal differences for the four extremity breadth measurements were the lowest relative to other variables measured during Cycle III. The technical errors of measurement were also lowest, indicating that these four measurements were quite accurately replicated. For example, these measurements averaged about 0.1 cm. difference for intra-examiner replications and about 0.10 cm. for inter-observer replications. Comparing the average differences for these four extremity breadth measurements to values for other body measurements in table II clearly indicates that precision was greater in these than in any other group of measurements considered in this report.

Attempting to compare coefficients of variation of these measurements with any others is misleading, as discussed earlier. Thus, the coefficient of variation statistics should be used only within the groups of measurements considered. For intra-examiner differences, knee breadth was best replicated, followed by ankle, elbow, and wrist breadths. For inter-examiner differences, elbow breadth had the smallest coefficient of variation, followed by ankle, wrist, and knee breadths. Testing at the .05 level, the F ratios indicated that in all instances intra-examiner differences were significantly smaller than inter-examiner differences.

The two measurements made with the spreading caliper, bizygomatic breadth and bigonial breadth, were likewise well replicated. The mean, median, and modal differences for these two facial breadth measurements were of approximately the same magnitude as those for the extremity breadth measurements. In fact, bizygomatic breadth had the smallest intra-examiner difference of all measurements considered, an average difference of 0.075 cm. and σ of 0.076. On an intra- and inter-examiner basis, bizygomatic breadth had a smaller coefficient of variation than bigonial breadth. The greater variability in replicating the latter might be related to variations in pressure in applying the

spreading caliper (slight variations producing an error of 0.1 cm.) and to variations in palpating the measuring landmark, the gonial angles of the mandible. Experience indicates that some observers allow the calipers to "slip" off the landmark. Similarly, if a child tenses his lower jaw, this also alters the measurement to some extent. In contrast to the measurement of bigonial breadth, bizygomatic breadth is a maximum measurement, in which the technician moves the spreading calipers until he notes the maximum reading. For both facial breadth measurements, the intra-examiner differences were significantly smaller than the inter-examiner differences at the .05 level.

The group of dimensions measured with the upper segment of the anthropometer included two foot measurements (length and breadth), three bony breadth measurements (biacromial, bicristal, and bitrochanteric breadths), and two human engineering breadth measurements (elbow-elbow and seat breadths). In making these measurements, the fixed arm of the anthropometer is set at one landmark, while the free arm is moved to the other landmark defining the particular measurement.

The two foot dimensions showed a high degree of replicability. Mean, median, and modal differences for foot breadth were less than or equal to those for foot length and were of the same magnitude as those for measurements made with the spreading and sliding calipers. This might be a function of the overall size of the dimensions being measured. The technical errors of measurement for both foot dimensions were smaller within than between examiners, and the intra-examiner differences were significantly smaller than the inter-examiner differences. The two foot measurements had consistently smaller technical errors of measurement and coefficients of variation than the other measurements made with the upper segment of the anthropometer.

The bony breadth measurements across the shoulders (biacromial breadth) and across the hips (bicristal and bitrochanteric breadths) also appeared to be reasonably well replicable measurements. Biacromial breadth and bitrochanteric breadth had essentially identical mean differences in the intra-examiner comparisons—0.553 and 0.552 cm., respectively. Bicristal breadth, on the other hand, had a larger average error in the intra-examiner comparisons, 0.775 cm. On an inter-examiner basis, biacromial breadth had the smallest average difference (0.807 cm.), while bitrochanteric breadth had the largest (1.760 cm.), with bicristal breadth very similar to it (1.590 cm.). These average differences are misleading and are perhaps influenced by extreme readings. Median differences in the inter-examiner comparisons are identical for both biacromial and bitrochanteric breadths (0.5 cm.), while that for bicristal breadth is much greater (1.1 cm.). All mean, median, and modal differences, as well as the technical errors of measurement for the three bony

breadth measurements, were smaller for intra-examiner comparisons than for inter-examiner comparisons. The intra-examiner differences were also significantly smaller than the inter-examiner differences. Within this group of three bony breadth measurements, biacromial breadth had the least relative variation, as indicated by the lower coefficients of variation on both an intra- and inter-examiner basis. Bitrochanteric breadth was close to biacromial breadth but larger in relative variation in both intra- and inter-examiner comparisons. Bicristal breadth had the largest coefficients of variation. The relative variability for the inter-examiner replicates was more than twice that noted for the intra-examiner replicates, indicating that different observers had difficulty in replicating this measurement with accuracy. These observations might be related to the nature and location of the bony landmarks involved in making these three measurements. The acromial processes are relatively close to the surface and easily located. The same applies in general to the greater trochanters of the femur. The iliac crests, though rather easily identified, are perhaps difficult to accurately replicate because of their irregular shape. Contributing to the overall variation in bony breadth measurements is the need for firm pressure in applying the arms of the anthropometer to the bony landmarks. Any inadvertent alteration of pressure applied can increase the error of measurement.

The two human engineering breadth measurements, elbow-elbow and seat breadths, appeared to be only moderately replicable when compared to other measurements made with the upper segment(s) of the anthropometer. Of the two measurements, elbow-elbow breadth had larger mean, median, and modal differences as well as larger technical errors of measurement in both the intra- and inter-examiner comparisons than did seat breadth. Elbow-elbow breadth also had a larger coefficient of variation than seat breadth. All statistics were smaller for the intra-examiner replications than for the inter-examiner replications. These two breadth measurements also had the lowest F ratios, the ratio for seat breadth being insignificant and that for elbow-elbow breadth barely significant at the .05 level, which would seem to suggest that in both measurements the individual idiosyncrasies of specific examiners had small effects. This interpretation is offset, however, by the fact that the magnitude of the differences between replicate readings in both the intra- and inter-examiner comparisons was rather large. This is perhaps a function of the specific measurements, since both require only light surface contact (the slightest pressure might distort replicate readings). Also, in measuring elbow-elbow breadth rather rigid positioning is required, and inadvertent alterations in positioning by the subject from one measurement session to the next might affect the replicate readings.

The six circumference measurements taken in Cycle III can be divided into those made on the torso

and those made on the extremities. The three torso girths—chest, waist, and hip girths—are essentially human engineering-type measurements, and the replicate analysis is similar to that noted for the two human engineering breadth measurements above. Chest, waist, and hip girths appeared only moderately replicable. Testing at the .05 level, the F ratios indicated no significant differences for hip girth, just barely significant differences for waist girth ($F = 1.42$), and significant differences for chest girth ($F = 2.75$) between intra- and inter-examiner replicates. These observations suggest that in such girth measurements individual idiosyncrasies of specific examiners had small effects. This interpretation is offset, however, by the magnitude of the differences between replicate examinations in both the intra- and inter-examiner comparisons, which were among the largest for the entire series of 31 measurements. Clearly, the same observer as well as different observers had difficulty replicating these three circumference measurements.

The three extremity circumferences had considerably smaller average differences between replicate readings, both within and between examiners, than did the three torso circumferences. This is perhaps a function of the magnitude of the circumferences measured. All but calf circumference appeared to be highly replicable measurements. Mean, median, and modal differences as well as the technical errors of measurement were slightly smaller for the intra-examiner than for the inter-examiner analysis. Observations for calf circumference were in the opposite direction, the average difference and the technical error of measurement were larger for the intra-examiner than for the inter-examiner analysis. However, the median and modal differences were identical on an intra- and inter-examiner basis. The effects of two or three discrepant replicate readings were responsible for inflating the intra-examiner mean difference value and the technical error of measurement. This is contrary to general measurement experience, for calf circumference is generally a highly replicable measurement. The present observations are probably a chance occurrence.

Although standing height was grouped with body weight on the basis of the automated measuring procedures used, the replicate observations for height will be considered here with other height measurements. Of all the height measurements, including standing height, sitting height, and segmental height measurements, it appeared that, both within and between examiners, standing height was best replicated. While sphygion height and thigh clearance (really height above the sitting surface) had smaller technical errors of measurement, this can be attributed to the smaller margin of error in taking the measurement. Problems encountered in radial and stylium heights have been discussed earlier. In these two measurements, the technical error of measurement was larger for the intra-examiner replicates than for the inter-examiner replicates. This was en-

tirely a function of one or two discrepant replicate readings, which distorted the technical error of measurement. Median differences between intra- and inter-examiner replicates were negligible for radial and stylium heights.

Examination of the F ratios for the various height measurements indicated that for all measurements except acromial height, radial height, and trochanteric height, there were significantly larger differences when two different observers made the measurements than when a single one did them. It should be noted in table II that the three height measurements for which the F ratio was not significant had among the largest mean differences both within and between examiners. For example, trochanteric height, which had the smallest F ratio ($F = 106$), had the largest mean differences on both intra-examiner replicates (1.413 cm.) and inter-examiner replicates (1.600 cm.). These observations perhaps depend on the measurements involved and factors affecting the taking of these measurements. In addition to the location of landmarks, acromial and radial height are greatly affected by slight changes in the posture and attitude of the subjects, while in the case of trochanteric height, location of the trochanteric landmark can be difficult in individuals with a lot of soft tissue over this area.

Discussion and Summary of Replicate Analysis

The preceding discussion of results of the replicate analysis of Cycle III body measurements was not aimed at determining which measurements were easiest or most difficult to perform but at evaluating the use of single and multiple examiners in a large-scale survey. Reports of large-scale surveys generally do not include discussions of replicate analyses of multiple examiner effects. One general impression derived from the analysis of the present data is that there is an obvious need to publish replicate studies in anthropometric surveys. This would insure better comparability of surveys and would aid in establishing tolerance limits for various body dimensions.

It should be emphasized that many of the measurements comprising the Cycle III (12-17 years) replicate analysis were taken in Cycle II (6-11 years). For example, six of the dimensions utilized in this report of Cycle II data and 11 of the 21 dimensions described in the previous report (Series II, No. 123) are included among the measurements discussed in the replicate analysis. Hence, of the 31 measurements used in the replicate analysis, 17 were also taken in Cycle II. The primary difference is in the replacement of traditional human engineering dimensions in Cycle II (buttock-knee length, buttock-popliteal length, popliteal height, knee height) and specific segmental lengths (acromion-olecranon length, elbow-wrist length, hand length) by eight segmental heights in Cycle III. Specific segmental lengths are estimated in the Cycle III data by subtrac-

tion. For example, acromial height minus olecranon height provides an estimate of upper arm length similar to that provided by direct measurement of acromion-olecranon length.

In addition, conditions under which the various anthropometric dimensions were measured were essentially identical in Cycles II and III, although several of the measurements were different. Instrumentation, instruction, and measurement technique were likewise basically the same in both cycles. Hence, *the observations derived from the Cycle III replicate analysis are generally applicable to the Cycle II data.*

Measurement of various body dimensions presents a unique situation. There are a large number of variables (sources of error) that must be controlled in the measurement environment in general and at the moment of measurement in particular. General sources of error can be grouped into three categories, the subject, the instrument, and the observer. Subject position, though carefully standardized, is difficult to control precisely. Postural attitude, phase of the breathing cycle, degree of tension and/or relaxation, and soon are factors which make it almost impossible to fully control the examinee so as to permit identical conditions during each of two measurement sessions that comprise replicate studies. In, for example, measurement of segmental heights, an inadvertent shifting of body weight from one leg to another can alter the height of a specific landmark from the standing surface, or tensing of the shoulders might make accurate location of acromiale difficult to replicate.

Instruments are carefully calibrated and checked out during the course of the survey. Hence, instrument variability is reasonably controlled. It is difficult, however, to control completely the observers' use and application of instruments to specific body landmarks in addition to the problem of consistently locating these landmarks. Differences between observers are inevitable, as the present replicate analysis indicates. Training, both prior to and in the field, helps reduce differences between observers, but it will not eliminate them completely. In light of this reality, there is an obvious need to establish tolerance limits within which two or more observers are permitted to vary in making a particular measurement. Similarly, the same observer varies to some extent within his own replicate measurements, although intra-observer variation, as expected, is consistently less than variation between observers. Perhaps the results of the Cycle III replicate analysis can be used to establish tolerance limits within which a single observer is permitted to vary in an intra-examiner replication and within which two or more observers are permitted to vary in an inter-examiner replication.

Since variation between observers is inevitable, what can be concluded from this analysis? In general, measurements made with the sliding and spreading callipers are highly replicable. These instruments are used in making bone-to-bone measurements requiring firm pressure and traversing relatively small distances. Further, the landmarks for these measurements are

rather easily located. Measurements made with the upper segment(s) of the anthropometer appear to vary with the specific measurement. The two foot measurements, breadth and length, are highly replicable. The three bony breadth measurements across the torso—biacromial, bicristal, and bitrochanteric breadths—are reasonably replicable. The apparent problem with these measurements relates to the consistent location of the landmarks, especially the iliac crests, and the application of firm pressure to compress underlying soft tissues, especially in the case of bitrochanteric breadth. It would be interesting to see a replicate analysis of the two hip breadth measurements by sex, since adolescent girls tend to accumulate adipose tissue over these sites. The two human engineering breadth measurements, elbow-elbow breadth and seat breadth, which are made with the upper segment of the anthropometer, are somewhat difficult to replicate, perhaps because light surface contact is required in making these measurements. Girth measurements on the torso are also difficult to replicate. Like the two human engineering breadth measurements, these dimensions require light surface contact with no soft tissue compression. Girth measurements on the extremities are, in general, well replicated. The discrepancy noted for calf circumference in the present analysis is somewhat of a surprise and is probably a chance occurrence. Calf circumference is generally a well-replicated girth measurement, and the result of the present analysis can be overlooked to some extent.

Height measurements, standing or sitting, are reasonably well replicated, there is, however, considerable variation in the replicability of the series of measurements evaluated. This variation is probably related to both subject and observer variation. Although the subject's position is standardized, inadvertent change in his postural attitude can alter the height of the segment landmark from the standing surface. It is almost impossible to control for this. Inter-observer variation is present for all measurements. Interestingly, it was least for standing height.

As indicated earlier, differences between examiners are inevitable in a large-scale anthropometric survey. This is true regardless of efforts at control and/or elimination. The extent of variation between observers should, however, be noted and reported? Error introduced by multiple observers, i.e., differences between examiners, have two apparent effects: first, they increase variable error, but second, they reduce the probability of a systematic error being introduced into the measuring process by idiosyncrasies of individual observers. An increase in the variable error must be tolerated to achieve a reduction of probable systematic error. Although variation is apparent in the present analysis of replicate measurements, the general impression is one of reasonable consistency in the measurement process utilizing multiple examiners. Comparative data from other large-scale anthropometric surveys of children are apparently not available.

VITAL AND HEALTH STATISTICS PUBLICATION SERIES

Originally Public Health Service Publication No. 1000

- Series 1. Programs and collection procedures.*—Reports which describe the general programs of the National Center for Health Statistics and its offices and divisions, data collection methods used, definitions, and other material necessary for understanding the data.
- Series 2. Data evaluation and methods research.*—Studies of new statistical methodology including experimental tests of new survey methods, studies of vital statistics collection methods, new analytical techniques, objective evaluations of reliability of collected data, contributions to statistical theory.
- Series 3. Analytical studies.*—Reports presenting analytical or interpretive studies based on vital and health statistics, carrying the analysis further than the expository types of reports in the other series.
- Series 4. Documents and committee reports.*—Final reports of major committees concerned with vital and health statistics, and documents such as recommended model vital registration laws and revised birth and death certificates.
- Series 10. Data from the Health Interview Survey.*—Statistics on illness, accidental injuries, disability, use of hospital, medical, dental, and other services, and other health-related topics, based on data collected in a continuing national household interview survey.
- Series 11. Data from the Health Examination Survey.*—Data from direct examination, testing, and measurement of national samples of the civilian, noninstitutional population provide the basis for two types of reports: (1) estimates of the medically defined prevalence of specific diseases in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics, and (2) analysis of relationships among the various measurements without reference to an explicit finite universe of persons.
- Series 12. Data from the Institutional Population Surveys.*—Statistics relating to the health characteristics of persons in institutions, and their medical, nursing, and personal care received, based on national samples of establishments providing these services and samples of the residents or patients.
- Series 13. Data from the Hospital Discharge Survey.*—Statistics relating to discharged patients in short-stay hospitals, based on a sample of patient records in a national sample of hospitals.
- Series 14. Data on health resources, manpower and facilities.*—Statistics on the numbers, geographic distribution, and characteristics of health resources including physicians, dentists, nurses, other health occupations, hospitals, nursing homes, and outpatient facilities.
- Series 20. Data on mortality.*—Various statistics on mortality other than as included in regular annual or monthly reports—special analyses by cause of death, age, and other demographic variables, also geographic and time series analyses.
- Series 21. Data on natality, marriage, and divorce.*—Various statistics on natality, marriage, and divorce other than as included in regular annual or monthly reports—special analyses by demographic variables, also geographic and time series analyses, studies of fertility.
- Series 22. Data from the National Natality and Mortality Surveys.*—Statistics on characteristics of births and deaths not available from the vital records, based on sample surveys stemming from these records, including such topics as mortality by socioeconomic class, hospital experience in the last year of life, medical care during pregnancy, health insurance coverage, etc.

For a list of titles of reports published in these series, write to:

Office of Information
National Center for Health Statistics
Public Health Service, HIRA
Rockville, Md. 20852