#### DOCUMENT RESUME

ED 225 032

CG 016 378

AUTHOR TITLE

Black, David R.; And Others

Developing a Couples Application Form to Predict

Weight Loss Success.

PUB DATE

NOTE

Aug 82 5p.; Paper presented, at the Annual Convention of the American Psychological Association (90th, Washington, DC, August 23-27, 1982). For related document, see CG.

016 379.

PUB TYPE

Reports - Research/Technical (143) -- Speeches/Conference Papers (150)

EDRS PRICE DESCRIPTORS

MF01/PC01 Plus Postage.

\*Behavior Change; \*Behavior Modification; Body Weight; Health Programs; Intervention; Obesity; \*Performance Factors; Predictor Variables; Spouses

**IDENTIFIERS** 

\*Marital Satisfaction; \*Weight Loss

#### **ABSTRACT**

Individual variability in response to behavioral weight loss programs remains an unresolved and perplexing issue for health psychologists. Determining in advance who will succeed or fail at losing weight would be useful in devising programs that produce a more homogeneous and dramatic response to treatment. In order to examine which responses to an application and a relationship quastionnaire relate to weight loss success, 23 couples participated in a l-year correspondence weight reduction program. One person in each couple enrolled to lose weight and the other agreed to act as helper. Reducers completed an application form recording demographic information, reactions to past attempts at weight loss, and number of previous methods used. Helpers rated their willingness to perform. weight-related tasks to help the reducer, Both reducers and helpers completed the Locke-Wallace Marital Adjustment Scale. Analyses of results indicated that better educated reducers and helpers with fewer negative attitudes about past weight loss attempts and fewer prior attempts at losing weight performed better, and that those with partners less willing to aid them or with unsatisfying marital relationships lost the most weight. The findings suggest that the application items and the measure of pelationship satisfaction may be useful predictors of weight loss. (PAS)

Developing A Couples Application Form to Predict

Weight Loss Success

David R. Black

University of Nebraska Medical Center

William E. Threlfall

Stanford University

Judy K. Black

Millard Public Schools, Omaha, Nebraska

Paper 'Presented at the Annual Meeting of the American Psychological Association, Washington, D.C., August, 1982

U.S. DEPARTMENT OF EDUCATION
NATIONAL INSTITUTE OF EDUCATION
EOUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

This document has been reproduced as received from the person or organization originating it

 Minor changes have been made to improve reproduction quality

 Points of view or opinions stated in this document do not necessarily represent official NIE position or policy, \*PERMISSION TO REPRODUCE THIS
MATERIAL HAS BEEN GRANTED BY

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)."

ERIC

- Developing A Couples Application Form to Predict Weight Loss Success

Individual variability in response to behavioral weight loss programs remains an unresolved and perplexing issue for health psychologists. While enrolled in behavioral weight loss programs, some individuals perform remarkably well and achieve their weight loss objectives and others perform quite poorly and may even gain weight. Therefore, determining in advance who would succeed or have difficulty losing weight would be useful in devising programs that would produce a more homogeneous and dramatic response to the treatment intervention. The present study was designed to meet this objective by examining which responses to an application and relationship questionnaire relate to weight loss success. A secondary purpose was to present questionnaire items that could be employed and further validated by other researchers working with couples who are interested in identifying prognostic indicators of weight loss success.

## Method

# Subjects

Twenty-three couples participated in a one year correspondence weight reduction program offered by the Stanford Heart Disease Prevention Program, Stanford School of Medicine. One person in each couple enrolled to lose weight and the other agreed to act as their "helper." The reducers included 6 men and 17 women between 28 and 62 years of age (M = 44.82 yrs.), who initially weighed 151 to 232 pounds (M = 186.14 lbs.) and were 37.65 to 84.25 percent overweight, (M = 56.67%).

# Procedure

As part of enrollment process, reducers were asked to complete an application form. They recorded demographic information, reactions to past

ERIC

weight loss attempts and the number of previous weight loss methods used. Helpers, on the other hand, rated their willingness to perform various weight-related tasks designed to aid the reducer in losing weight. In addition, both the helper and reducer completed the Locke-Wallace (1959) Marital Adjustment Scale (MAS).

# Results

Table 1 presents correlations between items on the application form and measures of body change at the end of the one year program. The majority of these correlations were statistically significant.

# Insert Table 1 about here

## Discassion

Because of the growing interest and relative success of utilizing spouses. in weight loss programs, an effort was made to isolate items on an initial application form and the MAS that would be helpful in predicting weight loss success. The findings of the present study show that better educated reducers (and helpers) with fewer negative attitudes about past weight loss attempts and fewer prior attempts at losing weight performed better. Also, those whose partners were less willing to aid the reducer in losing weight or whose marital relationship was rated less satisfying lost the most weight. One explanation for this counter-intuitive finding may be that the "uncooporative" attitude of the spouse and/or feelings of being moderately dissatisfied with the relationship serve as a motivator to reduce.

Although the findings are encouraging for identifying application items that may be useful predictors, sample size limitations suggest the need for further research. In particular, the application items as well as the measure of relationship satisfaction should be incorporated and further validated by other researchers in order to identify reliable and valid prognostic indicators of weight loss success for participants enrolled in couples programs.



Table 1
Correlations Between Application Items and Client Performance

•	•	Performance Measure			
	Application Item	Lbs. Lost	. RQ <sup>a</sup>	% Fat Lost	Fat .RQ <sup>b</sup>
1.	Educational level of reducer	.37*	.23	.25	.23
	Educational level of helper	.36*	.45*	.46*	.47*
2.	Reactions about past wt. loss efforts	· - 460*	÷.61**	59**	,55**
(3.	Number of previous wt. loss . attempts	55**	40*	40*	35*
4.	Helper's ratings of willingness to perform wt. loss tasks	42*	41*	56** ,	53**
5.	MAS scores for reducers	55**	56**	44*	43*
· ·	MAS <sup>C</sup> scores for helpers	34	20	51**	52**

 $<sup>^{</sup>a}$ RQ =  $\frac{Lbs. Lost}{Initial Lbs. Overwt.} \times 100.$ 

bFat RQ = Fat Lost Initial Amt. Overfat x 100.

CMAS = Locke-Wallace (1959) .
Marital Adjustment Scale.,

<sup>\*</sup>p < .05.

<sup>\*\*</sup> p < .01. ·