

DOCUMENT RESUME

ED 267 314

CG 018 894

AUTHOR Abidin, Richard R.; Wilfong, Ellen M.
TITLE Parenting Stress: Maternal & Child Health Correlates.
PUB DATE [Aug 85]
NOTE 16p.; Paper presented at the Annual Convention of the American Psychological Association (93rd, Los Angeles, CA, August 23-27, 1985).
PUB TYPE Reports - Research/Technical (143) -- Speeches/Conference Papers (150)
EDRS PRICE MF01/PC01 Plus Postage.
DESCRIPTORS *Child Rearing; *Diseases; *Health; Medical Services; *Mother Attitudes; Mothers; Parent Child Relationship; Parents; *Stress Variables; *Toddlers
IDENTIFIERS *Medical Utilization; Parenting Stress Index; Traumas

ABSTRACT

A study was undertaken to examine the relationship among parenting stress, maternal and child health, and medical utilization. Participants were patients from a large group pediatric practice. Mothers (N=56) of patients completed the Parenting Stress Index. Twelve months after completion of the Parenting Stress Index, subjects were assessed using the: (1) General Health Rating, which measured subjects' perception of their health status; (2) Medical Utilization for the Child, which gave frequency counts from the child's medical chart; (3) Mother's Health Care Seeking Behaviors, which measured the mother's report of her use of medical services; (4) Mother's Illness-Related Behaviors, which consisted of mother's monthly mean report of days in bed, non-prescription medicines used, etcetera; and (5) Traumatic Injuries, which gave frequency counts from child's medical charts. The results showed that maternal level of parenting stress appeared unrelated to obtaining medical services for young children. Inappropriate medical utilization rates based on maternal stress did not occur for very young children. Maternal level of parenting stress was related to high rates of maternal use of medical services, illness-related behaviors, and maternal perceptions of health status. Low and moderate stress levels were not related to illness behaviors. Parenting stress was not shown to be related to traumatic injuries of the children, possibly due to the young age of this group of children, who were 12-24 months old when the observations were made. The Parenting Stress Index may be used in identifying parents under excessive stress who may later exhibit illness behavior. (ABL)

 * Reproductions supplied by EDRS are the best that can be made *
 * from the original document. *

ED267314

PARENTING STRESS:
MATERNAL & CHILD HEALTH CORRELATES

ABIDIN, R. R. & WILFONG, E.
INSTITUTE OF CLINICAL PSYCHOLOGY
UNIVERSITY OF VIRGINIA

CG 018894

U.S. DEPARTMENT OF EDUCATION
NATIONAL INSTITUTE OF EDUCATION
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it
- Minor changes have been made to improve reproduction quality
-
- Points of view or opinions stated in this document do not necessarily represent official NIE position or policy

"PERMISSION TO REPRODUCE THIS MATERIAL HAS BEEN GRANTED BY

Richard R. Abidin
Ellen M. Wilfong

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC) "

ABSTRACT

THE PRESENT STUDY EXAMINED THE RELATIONSHIPS BETWEEN AND AMONG THE VARIABLES OF PARENTING STRESS, MATERNAL AND CHILD HEALTH, AND MEDICAL UTILIZATION.

THE DEMOGRAPHIC CHARACTERISTICS OF THE SAMPLE ARE DESCRIBED IN TABLE #1.

METHODS & RESULTS

I. DATA COLLECTION

ALL PATIENTS WERE DRAWN FROM A LARGE GROUP PEDIATRIC PRACTICE (PEDIATRIC ASSOCIATES OF CHARLOTTESVILLE, VIRGINIA). THE MOTHERS HAD ALL TAKEN THE PARENTING STRESS INDEX (PSI) "INITIALLY" AS A PART OF THE PRACTICE'S NORMAL COMPREHENSIVE PEDIATRIC HEALTH CARE. THE PHYSICIANS INVITED PATIENTS TO VOLUNTEER FOR THE PRESENT STUDY.

THE SUBJECTS COMPLETED THE "SUBSEQUENTLY" USED MEASURES APPROXIMATELY 12 MONTHS FOLLOWING THE INITIAL COMPLETION OF THE PSI AND GRANTED PERMISSION FOR THEIR MEDICAL RECORDS TO BE REVIEWED.

II. MEASURES USED

1. PARENTING STRESS INDEX (PSI)=101 ITEM MEASURE OF PARENTING STRESS DIVIDED INTO TWO DIVISIONS OF CHILD CHARACTERISTIC STRESSORS AND PARENTING CHARACTERISTICS--ABIDIN, R.R. (1983). PARENTING STRESS INDEX--MANUAL, PEDIATRIC PSYCHOLOGY PRESS, 320 TERRELL ROAD WEST, CHARLOTTESVILLE, VA 22901.
2. GENERAL HEALTH RATING (GHR)=35 ITEM MEASURE OF THE SUBJECTS' PERCEPTIONS OF THEIR HEALTH STATUS--DAVIES, A.R. AND WARE JR., J.E. (1981). MEASURING HEALTH PERCEPTIONS IN THE HEALTH INSURANCE EXPERIMENT. RAND CORPORATION, SANTA MONICA, CA 90406.
3. MEDICAL UTILIZATION FOR THE CHILD (MUC)=FREQUENCY COUNTS FROM THE CHILD'S MEDICAL CHART.
4. MOTHER'S HEALTH CARE SEEKING BEHAVIORS (MHCSB)=MOTHER'S REPORT OF HER USE OF MEDICAL SERVICES.
5. MOTHER'S ILLNESS-RELATED BEHAVIORS (MIRB)=MOTHER'S MONTHLY MEAN REPORT OF: DAYS IN BED, NON-PRESCRIPTION MEDICATIONS USED, NUMBER OF DIFFERENT SYMPTOMS, ETC.
6. TRAUMATIC INJURIES (Ti)=FREQUENCY COUNTS FROM CHILD'S MEDICAL CHARTS OF TRAUMATIC INJURIES--LACERATIONS, BURNS, ACCIDENTAL POISONING, ETC.

III. QUESTIONS ASKED

1. ARE DIFFERENT LEVELS OF MATERNAL STRESS RELATED TO DIFFERENT RATES OF MEDICAL SERVICES UTILIZATION FOR THE CHILD?

ANSWER: NO (SEE FIGURE #1).

2. ARE DIFFERENT LEVELS OF MATERNAL STRESS RELATED TO DIFFERENT RATES OF MEDICAL SERVICES UTILIZATION FOR THE MOTHER?

ANSWER: YES (SEE FIGURES #2 & 3).

3. ARE DIFFERENT LEVELS OF MATERNAL STRESS RELATED TO DIFFERENT RATES OF MATERNAL ILLNESS-RELATED BEHAVIORS (MIRB)?

ANSWER: YES (SEE FIGURE #4).

4. ARE DIFFERENT LEVELS OF MATERNAL STRESS RELATED TO DIFFERENT GENERAL HEALTH RATINGS (GHR)?

ANSWER: YES (SEE FIGURES #5 & 6).

5. BETWEEN THE VARIABLES OF MATERNAL STRESS (PSI) AND MATERNAL GENERAL HEALTH RATING (GHR), WHICH IS A BETTER PREDICTOR OF MOTHER'S HEALTH CARE SEEKING BEHAVIOR?

ANSWER: MATERNAL STRESS (SEE TABLE #2).

6. CAN WE REPLICATE ABIDIN'S 1982 FINDING OF DIFFERENT RATES OF TRAUMATIC INJURIES FOR A YOUNGER GROUP OF CHILDREN WITH MOTHERS AT DIFFERENT LEVELS OF STRESS?

ANSWER: NO (SEE TABLES #3 & 4).

ABIDIN, R.R., PARENTING STRESS AND THE UTILIZATION OF PEDIATRIC SERVICES. CHILDREN'S HEALTH CARE, 1982, 11.2, 70-73.

7. ARE THE PARENTING STRESS INDEX SCORES AND GENERAL HEALTH RATINGS SCORES OF THE PARENTS IN THIS STUDY DIFFERENT FROM THE NORMS OF BOTH INSTRUMENTS?

ANSWER: NO (SEE TABLE #5).

DISCUSSION


MATERNAL LEVEL OF PARENTING STRESS APPEARS UNRELATED TO OBTAINING MEDICAL SERVICES FOR YOUNG CHILDREN, AND THEREFORE INAPPROPRIATE MEDICAL UTILIZATION RATES BASED ON MATERNAL STRESS DO NOT APPEAR TO OCCUR FOR VERY YOUNG CHILDREN.

MATERNAL LEVEL OF PARENTING STRESS IS HIGHLY RELATED TO HIGH RATES OF MATERNAL USE OF MEDICAL SERVICES, TO ILLNESS-RELATED BEHAVIORS, AND TO MATERNAL PERCEPTIONS OF THEIR HEALTH STATUS. LOW TO MODERATE STRESS LEVELS DO NOT SEEM TO BE RELATED TO "ILLNESS BEHAVIORS," WHILE HIGH STRESS RESULTS IN A GREAT DEAL OF "ILLNESS BEHAVIOR" IN THE YEAR FOLLOWING THE STRESS MEASUREMENT. IT APPEARS THAT THE PARENTING STRESS INDEX MAY BE USEFUL IN IDENTIFYING PARENTS UNDER EXCESSIVE LEVELS OF STRESS WHICH MAY RESULT IN ILLNESS OR "ILLNESS BEHAVIOR."

THE RELATIONSHIP PREVIOUSLY REPORTED OF LEVELS OF PARENTING STRESS TO TRAUMATIC INJURIES IN CHILDREN WAS NOT REPLICATED IN THIS STUDY. THE YOUNGER AGE OF THIS SAMPLE (12-24 MONTHS AT THE TIME THE OBSERVATIONS WERE MADE) PROBABLY REFLECTS THE REDUCED OPPORTUNITY FOR TRAUMATIC INJURIES WHEN COMPARED TO THE PREVIOUS SAMPLE OF CHILDREN 24 TO 48 MONTHS OF AGE.

FIGURE #1

MEDICAL UTILIZATION* FOR CHILD
BY INITIAL AND SUBSEQUENT PARENTING STRESS LEVEL




	LOW	MODERATE	HIGH
INITIAL	N=10	N=31	N=15
SUBSEQUENT	N=16	N=34	N=6

*MEAN MONTHLY: SICK VISITS, EMERGENCY ROOM VISITS, VISITS TO OTHER PHYSICIANS, AND PHONE CALLS

FIGURE #2


MOTHERS' INITIAL STRESS LEVEL AND MOTHERS' SUBSEQUENT 12 MONTH,
MEAN¹ HEALTH CARE SEEKING BEHAVIORS



¹MEAN DIFFERENCES: LOW STRESS VS. MODERATE--N.S.
LOW STRESS OR MODERATE STRESS VS.
HIGH STRESS P=,00

FIGURE #3


MOTHERS' SUBSEQUENT STRESS CATEGORY AND MOTHERS' MEAN¹ HEALTH CARE SEEKING BEHAVIORS FOR 12 MONTHS PRIOR TO STRESS MEASUREMENT



¹MEAN DIFFERENCES: LOW STRESS VS. MODERATE--N.S.
LOW STRESS OR MODERATE STRESS VS.
HIGH STRESS P=.00

FIGURE #5

INITIAL MATERNAL STRESS AND
MEAN¹ GENERAL HEALTH RATINGS INDEX (RAND SCALE)²




¹MEAN DIFFERENCES AMONG STRESS LEVELS SIGNIFICANT AT .00

²PROBABILITY OF A PHYSICAL ILLNESS 98.0=10%; 76.8=22%
PROBABILITY OF AN EMOTIONAL IMPAIRMENT 98.0=7%; 76.8=20%

PROBABILITIES BASED ON THE RAND-HEALTH INSURANCE EXPERIMENT DATA--
RAND CORPORATION, SANTA MONICA, CALIFORNIA 90406

FIGURE #6

SUBSEQUENT MATERNAL STRESS CATEGORY
AND MEAN¹ GENERAL HEALTH RATINGS INDEX²



¹MEAN DIFFERENCE SIGNIFICANT AT .00

²PROBABILITY OF A PHYSICAL ILLNESS 92.4=15%; 64.5=50%
PROBABILITY OF AN EMOTIONAL IMPAIRMENT 92.4=10%; 64.5=50%

PROBABILITIES BASED ON THE RAND-HEALTH INSURANCE EXPERIMENT DATA--
RAND CORPORATION, SANTA MONICA, CALIFORNIA 90406

TABLE #1

PARENTING STRESS--MATERNAL AND CHILD HEALTH CARE CORRELATION

DEMOGRAPHIC CHARACTERISTICS OF SAMPLE N=56

CHILDREN

SEX

34 MALE
22 FEMALE

AGE IN YEARS AT TIME OF SECOND MEASURE

MEAN 2.35
STANDARD DEVIATION 1.036
MINIMUM 1.4
MAXIMUM 5.3

BIRTH ORDER

44 FIRSTBORN

RACE

55 WHITE
1 BLACK

MOTHERS

AGE AT TIME OF SECOND MEASURE

MEAN 31.4 YEARS
STANDARD DEVIATION 3.8
MINIMUM 23.4
MAXIMUM 39.6

MARITAL STATUS

52 MARRIED
4 DIVORCED

INCOME LEVEL

29% UNDER \$30,000
14% OVER \$50,000
MEAN INCOME CATEGORY: \$30 TO \$35,000

MOTHER'S EDUCATION

HIGH SCHOOL	8
SOME COLLEGE OR VOCATIONAL	11
COLLEGE GRADUATE	22
GRADUATE OR PROFESSIONAL SCHOOL	15

FATHER'S EDUCATION

HIGH SCHOOL	10
SOME COLLEGE OR VOCATIONAL	8
COLLEGE GRADUATE	16
GRADUATE OR PROFESSIONAL SCHOOL	22

TABLE #2

MULTIPLE REGRESSION ANALYSIS
OF MOTHERS' HEALTH CARE SEEKING BEHAVIORS

<u>VARIABLES IN THE EQUATION</u>	<u>DF</u>	<u>F LEVEL</u>	<u>R² CHANGE</u>
INITIAL STRESS LEVEL-- TAKEN PRIOR TO RECORDING PERIOD FOR MOTHERS' HEALTH CARE SEEKING BEHAVIORS		4.025*	.147
GENERAL HEALTH RATINGS INDEX-- OBTAINED AT THE END OF 12 MONTHS' RECORDING PERIOD		2.395	.037
REGRESSION	2	5.984**	.184 (TOTAL)
RESIDUAL	53		

*P < .05

** < .01

TABLE #3

ANALYSIS OF VARIANCE
TRAUMATIC INJURIES BY INITIAL STRESS SCORE & CHILD AGE

<u>SOURCE OF VARIANCE</u>	<u>MS</u>	<u>DF</u>	<u>F</u>	<u>P</u>
COVARIATES CHILD AGE	.002	1	.356	.55 n.s.
MAIN EFFECT INITIAL STRESS	.006	2	1.242	.30 n.s.
RESIDUAL (ERROR TERM)	.005	52		

TABLE #4

ANALYSIS OF VARIANCE
TRAUMATIC INJURIES BY SUBSEQUENT STRESS SCORE & CHILD AGE

<u>SOURCE OF VARIANCE</u>	<u>MS</u>	<u>DF</u>	<u>F</u>	<u>P</u>
COVARIATES CHILD AGE	.002	1	.351	.56 n.s.
MAIN EFFECT SUBSEQUENT STRESS	.004	2	.810	.45 n.s.
RESIDUAL (ERROR TERM)	.005	52		

TABLE #5

MEANS AND STANDARD DEVIATIONS FOR MOTHERS'
 PARENTING STRESS SCORES (PSI) AND GENERAL HEALTH RATINGS (GHR)

	<u>PSI</u>			<u>GHR</u>	
	INITIAL SCORES	SUBSEQUENT SCORES	NORMS	SUBSEQUENT	NORMS
MEAN	222.3	211.7	221.0	MEAN 84.5	83.4
S.D.	45.0	47.1	38.9	S.D. 14.5	13.5