

ED 352 568

CG 024 669

AUTHOR Winters, Ken C.  
 TITLE A Comparison of Various DSM Criteria for Alcohol Use Disorders in Adolescents.  
 INSTITUTION Minnesota Univ., Minneapolis. Center for Adolescent Substance Abuse.  
 PUB DATE [Jul 92]  
 NOTE 6p.; Paper presented at the Annual Convention of the American Psychological Association (100th, Washington, DC, August 14-18, 1992).  
 PUB TYPE Reports - Research/Technical (143) -- Speeches/Conference Papers (150)  
 EDRS PRICE MF01/PC01 Plus Postage.  
 DESCRIPTORS \*Adolescents; \*Alcohol Abuse; \*Clinical Diagnosis; Evaluation Criteria; \*Marijuana; \*Substance Abuse  
 IDENTIFIERS \*Diagnostic Statistical Manual of Mental Disorders

## ABSTRACT

In recent years the pace has been accelerating by which changes in the Diagnostic and Statistical Manual of Mental Disorders (DSM) systems are made. Revisions occur before extensive empirical knowledge can be drawn upon to guide these changes. For substance use disorders, the criteria have undergone changes from DSM-III to DSM-III-R, and from DSM-III-R to DSM IV. The extent and impact of these revisions have not been formally investigated in an adolescent population. This study compared the three DSM systems in an adolescent clinical sample with respect to how youth were sorted into alcohol diagnostic groups and in terms of their relative concurrent validity. Adolescents (N=130) receiving drug abuse assessments at an outpatient evaluation clinic participated in the study. Subjects received ratings on alcohol and cannabis diagnostic symptoms, recent (past 12 months) drug use frequency, depth of addiction, and recommendation for level of drug treatment. Analysis indicated: (1) the DSM-III-R and DSM-IV have a broader, more inclusive definition of alcohol use disorders than DSM-III; (2) DSM-IV has a broader definition of abuse and a narrower definition of dependence than DSM-III-R; (3) the validity data indicate that all three DSM systems are comparably related to alternate measures of problem severity; and (4) the results were replicated when subjects were analyzed according to their cannabis use diagnoses. (ABL)

\*\*\*\*\*  
 \* Reproductions supplied by EDRS are the best that can be made \*  
 \* from the original document. \*  
 \*\*\*\*\*

ED352568

A Comparison of Various DSM Criteria for  
Alcohol Use Disorders in Adolescents

Ken C. Winters

CG024669

**BEST COPY AVAILABLE**

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE THIS  
MATERIAL HAS BEEN GRANTED BY

Ken Winters

TO THE EDUCATIONAL RESOURCES  
INFORMATION CENTER (ERIC).

## **A Comparison of Various DSM Criteria for Alcohol Use Disorders in Adolescents**

**Ken C. Winters, Ph.D.**  
Center for Adolescent Substance Abuse  
University of Minnesota

### **Abstract**

Three criteria for alcohol use disorders (DSM-III, DSM-III-R, and DSM-IV, proposed) were compared in 130 adolescents receiving drug abuse assessments. An analysis of the distribution of subjects indicated that both the DSM-III-R and DSM-IV criteria are broader than the DSM-III, and that the DSM-IV produces a broader definition of abuse and a narrower definition of dependence than the DSM-III-R. The concurrent validity analysis indicated that all three systems yielded comparable validity coefficients.

### **Introduction**

In recent years, the pace has been accelerating by which changes in the DSM systems are made. Revisions occur before extensive empirical knowledge can be drawn upon to guide the changes. For substance use disorders, the criteria have undergone changes from DSM-III to DSM-III-R, and from DSM-III-R to DSM-IV. The extent and impact of these revisions have not been formally investigated in an adolescent population.

The study compares the three DSM systems in an adolescent clinical sample with respect to how youth were sorted into alcohol diagnostic groups and in terms of their relative concurrent validity.

### **Method**

One-hundred and thirty adolescents receiving drug abuse assessments at an outpatient evaluation clinic in St. Paul, Minnesota participated in the study. Subjects had been referred for assessments by local schools or the juvenile court. Subjects received ratings on alcohol and cannabis diagnostic symptoms, recent (past 12-months) drug use frequency, depth of addiction, and recommendation for level of drug treatment. Diagnostic symptom ratings were based on results from a structured diagnostic interview (ADI; Winters and Henly, 1992); drug use frequency and depth of addiction were obtained from a self-report questionnaire (PEI; Winters and Henly, 1989); and drug treatment recommendations were based on consensus clinical judgements. The diagnostic interviewers were blind to scores on the other three variables.

**TABLE 1**  
**SAMPLE CHARACTERISTICS (N=130)**

<u>Age</u>	<u>Percent</u>
12-15	42.3
16-18	57.7
Male	63.8
Female	36.2
African American	4.6
American Indian	5.4
European American	86.9
Mexicano/Latino/Chicano	3.1

**TABLE 2**  
**COMPARISON OF SAMPLE CHANGES**  
**BETWEEN THE ALCOHOL GROUPS**

		<u>DSM-III-R</u>			
		No Dx	Abuse	Dependence	
<u>DSM-III</u>	No Dx	81	11	5	97 (74.6%)
	Abuse	0	1	9	10 (7.7%)
	Dependence	0	3	20	23 (17.7%)
		81 (62.3%)	15 (11.5%)	34 (26.2%)	130

		<u>DSM-IV</u>			
		No Dx	Abuse	Dependence	
<u>DSM-III</u>	No Dx	81	16	0	97 (74.6%)
	Abuse	0	10	0	10 (7.7%)
	Dependence	1	0	22	23 (17.7%)
		82 (63.1%)	26 (20.0%)	22 (16.9%)	130

		<u>DSM-IV</u>			
		No Dx	Abuse	Dependence	
<u>DSM-III-R</u>	No Dx	76	5	0	81 (62.3%)
	Abuse	6	7	2	15 (11.5%)
	Dependence	0	14	20	34 (26.2%)
		82 (63.1%)	26 (20.0%)	22 (16.9%)	130

**TABLE 3**  
**VALIDITY COEFFICIENTS FOR THE**  
**DSM-III ALCOHOL GROUPS**

	<u>12-Month</u> <u>Alcohol Frequency</u>	<u>Problem</u> <u>Severity Scale</u>	<u>Treatment</u> <u>Referral</u>
DSM-III	.51	.55	.31
DSM-III-R	.50	.59	.34
DSM-IV	.49	.55	.31

## Results

- On the basis of comparing sample distributions between the DSM alcohol groups...
  - a) DSM-III-R and DSM-IV have a broader, more inclusive definition of alcohol use disorders than DSM-III. Averaging across DSM-III-R and DSM-IV, 37.3% of the subjects received a substance use disorder diagnosis, while 25.4% received a substance use disorder diagnosis with DSM-III.
  - b) DSM-IV has a broader definition of abuse and a narrower definition of dependence than DSM-III-R. Based on DSM-IV, 20.0% had alcohol abuse and 16.9% had alcohol dependence ratings; for DSM-III-R, the percentages were 11.5% (abuse) and 26.2% (dependence).
- The validity data indicate that all three DSM systems are comparably related to alternate measures of problem severity.
- The results were replicated when subjects were analyzed according to their cannabis use diagnoses (data not shown).

## **Conclusions**

- **DSM-IV may be more appropriate for adolescent populations because it has a broader definition of abuse and a narrower definition of dependence than its predecessor, DSM-III-R.**
- **However, because DSM-IV maintains a relatively broad definition of substance use disorders, the validity of the minimum criteria for substance use disorders is still a relevant question for adolescent researchers.**
- **It would be ideal to longitudinally test the predictive validity of the various DSM groupings.**
- **Study limitations include: some subgroups had small sample sizes; the client was the sole informant for the diagnostic ratings; and little is known how to best measure substance use disorders in adolescents.**

## **Contact...**

**Ken Winters, Ph.D., Director, Center for Adolescent Substance Abuse, Box 721 - UMHC, Harvard Street at East River Road, Minneapolis, MN 55455.**