

DOCUMENT RESUME

ED 395 228

CG 026 959

AUTHOR Munzer, Jane; And Others  
 TITLE Pathways of Adolescent Suicidal Behavior.  
 PUB DATE [95]  
 NOTE 10p.; Poster presented at the Annual Meeting of the American Psychological Association (103rd, New York, NY, August 11-15, 1995).  
 PUB TYPE Reports - Research/Technical (143)  
 EDRS PRICE MF01/PC01 Plus Postage.  
 DESCRIPTORS \*Adolescents; Behavior; Family Influence; \*Path Analysis; Predictor Variables; Psychopathology; Secondary Education; \*Self Injurious Behavior; \*Sex Differences; \*Suicide  
 IDENTIFIERS Internalization; \*Suicide Attempts; \*Suicide Ideation

ABSTRACT

Suicide attempts and suicidal ideation among adolescents have been increasing faster than those for adults. This study addresses three questions on adolescent suicidal behavior: (1) Why do some adolescents with psychiatric disorders have a history of suicidal behaviors and some do not?; (2) How do intrapsychic and interpersonal underpinnings of adolescent suicidal behavior differ from patterns of similar behavior among children and adults?; and (3) How do pathways of adolescent suicidal behavior differ for boys and for girls? It was hypothesized that differences in expression of pathology between males and females might underlie the relationship between gender, self-esteem, social support from family, and suicidal behavior. Psychopathology was classified according to whether a dysfunctional behavior originated from inner turmoil ("internalizing"), or from outer turmoil (focusing on others' role in creating pain--"externalizing"). After examining 171 boys' and girls' suicidal behaviors versus adolescents with no history of such behavior, statistical analysis revealed the influence of gender but no significant gender-by-group interaction. Findings indicate that a low level of perceived family support was the only significant predictor of history of suicide attempts for girls. Boys' suicide attempts were predicted by perceived family support and by higher levels of externalizing behavior and internalizing behavior, and by greater age, when compared to girls. Four tables present statistical analysis. Contains 12 references. (RJM)

\*\*\*\*\*  
 \* Reproductions supplied by EDRS are the best that can be made \*  
 \* from the original document. \*  
 \*\*\*\*\*

## Pathways of Adolescent Suicidal Behavior

Jane Munzer, M.Ed., Ph.D.<sup>1</sup>; Adrienne Shelton-Keller, Ph.D.<sup>2</sup>; Peter Sheras, Ph.D.<sup>1</sup>;

Malcolm West, Ph.D.<sup>3</sup>; Ken Adam, M.D.<sup>4</sup>

<sup>1</sup>Curry Program in Clinical and School Psychology, University of Virginia; <sup>2</sup>Department of Psychiatric Medicine, University of Virginia; <sup>3</sup>Department of Psychiatry, University of Calgary, Alberta, Canada; <sup>4</sup>Department of Psychiatry, University of Toronto, Ontario, Canada.

Poster presentation at the American Psychological Association Annual Convention, New York, 1995.

*Background.* The rate of suicide attempt and suicidal ideation among adolescents has been steadily increasing at a rate greater than that for adults (Centers for Disease Control, 1985; National Center for Health Statistics, U.S. Department of Health and Human Services, 1988; Garland & Zigler, 1993). Despite these findings, relatively little empirical attention has been given to better understanding the personal and interpersonal characteristics specifically associated with adolescent suicidal behavior. The purpose of the present study was threefold. First, it was designed to address the question "Why do some adolescents with psychiatric disorders have a history of suicidal behaviors and some do not?" Second, the study explored how the intrapsychic and interpersonal underpinnings of adolescent suicidal behavior might differ from patterns of suicidal behavior among children and adults. Third, the study considered how such pathways of adolescent suicidal behavior might be different for boys and for girls.

*Hypothesis.* We hypothesized that differences in expression of psychopathology between males and females might underlie the relationship we had found between gender, self-esteem, social support from family, and suicidal behavior (Munzer, 1993). Expression of psychopathology was classified according to a common theoretical model which holds that dysfunctional behavior is a manifestation of inner turmoil; the manner in which individuals express this turmoil determines their style of psychopathology. Individuals who turn their turmoil inward, focusing on themselves as the cause of their

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.

- Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE AND  
DISSEMINATE THIS MATERIAL  
HAS BEEN GRANTED BY

A. Sheldon-Keller

1

2

TO THE EDUCATIONAL RESOURCES  
INFORMATION CENTER (ERIC)

BEST COPY AVAILABLE

pain and minimizing its direct expression to those around them are "Internalizing." Individuals who turn their turmoil outward, focusing on the role of others in its cause and taking these feelings out on others are "Externalizing."

*Methods.* Participants were 171 adolescents (95 males and 76 females), recruited upon admission to psychiatric treatment. The mean age is 14.9 years (s.d. = 1.6), with no significant difference between males and females ( $t(169) = .65$ ,  $p = .513$ ). The sample is predominantly white (84.5%). The study was designed as a case/comparison study of adolescents with a history of suicidal behaviors versus adolescents with no history of suicidal behaviors. History of suicidal behaviors was determined by semi-structured interview (Linehan & Nielsen, 1981). Among the male participants, 39 (41%) had a history of suicidal behaviors; 45 (59%) of the female participants had a history of suicidal behaviors.

The protocol was a single point-in-time design, with all instruments administered as individual interviews. Levels of internalizing and externalizing behavior were measured using items from the Youth Self-Report (Achenbach & Edelbrock, 1987), the Rosenberg Self-Esteem Scale (Rosenberg, 1965), the Beck Hopelessness Scale (Beck, Weissman, Lester & Trexler, 1974.) and the Demographic and Descriptive Interview (Chiles, Miller & Cox, 1980.). These revised measures of internalizing and externalizing behavior were developed to reflect the specific theoretical model of psychopathology stated above (Table 1). A measure of perceived family support (Procidano, Heller & Swindle, 1980) was also included. Multivariate analysis of variance (MANOVA) with two factors (sex and suicidal behaviors) was used to determine multivariate significance and variables to include in the model. Best-fit models differentiating adolescents with a history of suicidal behaviors were determined using logistic regression analyses.

*Results.* MANOVA with perceived social support, internalizing behaviors scale, externalizing behaviors scale and age revealed a significant effect for gender [ $F(4,122) = 6.28$ ,  $p < .001$ ] and for group [ $F(4, 122) = 12.14$ ,  $p < .001$ ] but no significant gender-by-group interaction. Table 2 summarizes the post-hoc univariate analyses. Based on logistic regression analysis of various possible models, the study revealed two distinct pathways of adolescent suicidal behavior which seem to be a reflection of gender differences rather than differences in style of psychopathology, as initially hypothesized (Table 3). A low level of perceived family support was the only significant predictor of history of suicide attempt for girls. For boys, lower perceived family support joined higher levels of externalizing behavior, higher levels

of internalizing behavior, and greater age as predictors of history of suicide attempt.

*Conclusions.* The findings of this study offer only a glimpse into the complex dynamic of family relationships and personality among adolescents. Future exploration of the specific directional pathways of cause and effect among the variables of suicidal behavior, family support, and degree and style of psychopathology would increase the clinician's ability to design and implement more successful interventions with troubled youth. The findings of this study clearly indicate, however, that any such exploration must consider different pathways of suicidal behavior for adolescent girls and boys. The study suggests that girls at this developmental stage may think about, evaluate, and internalize family interactions more than their male counterparts. The study also has strong implications for suicide risk screening. While statistical findings can not lead us to direct predictions of risk in individuals, the assessment of perceived family support is strongly indicated in screening for suicide risk among all psychologically troubled adolescents, but is especially salient in screening for risk among girls. Researchers and clinicians must address the influential role that social interactions, and more importantly, family relationships, play in the psychological development of adolescents of both genders.

## References

- Achenbach, T. M., & Edelbrock, C. (1987). Manual for the Youth Self-Report and Profile. Burlington, VT: University of Vermont Department of Psychiatry.
- Beck, A.T., Weissman, A., Lester, D. & Trexler, L. (1974). The measurement of pessimism: The hopelessness scale. Jl. Cons. Clin. Psychol. 42: 861-865.
- Centers for Disease Control. (1985). Suicide Surveillance: Summary 1970-80. Atlanta, GA: U.S. Department of Health and Human Services, Center for Health Promotion and Education.
- Chiles, J., Miller, M. & Cox, G. (1980). Depression in an adolescent delinquent population. Arch. Gen. Psychia. 37: 1179-1184.
- Garland, A. F., & Zigler, E. (1993). Adolescent suicide prevention: Current research and social policy implementation. American Psychologist, 48, 169-182.
- Linehan, M.M. & Nielsen, S.L. (1981). Assessment of suicidal ideation and parasuicide: Hopelessness and social desirability. Jl. Cons. Clin. Psychol. 49: 773-775.
- Munzer, J. S. (1993). Patterns of suicidal behavior, perceived social support, and self-esteem among adolescents with psychiatric disorders. Unpublished manuscript.
- National Center for Health Statistics, U.S. Department of Health and Human Services. (1988). Vital Statistics of the United States. Hyattsville, MD: Public Health Service.
- Procidano, M.E., Heller, K. & Swindle, R.W. (1980). Experimental studies of the nature and effects of social support. In New Developments in Research on Life Stress and Social Support. symposium presented at the meeting of the American Psychological Association, Montreal.
- Rosenberg, M. (1965). Society and the Adolescent Self-image. Princeton, N.J.: Princeton University Press.

**Table 1:****Items in Scales Created to Measure Internalizing and Externalizing Behavior**

<b>SCALE</b>	<b>ITEM</b>	<b>SOURCE</b>
<b>Externalizing Behaviors (<math>\alpha = .76</math>)</b>		
	I threaten to hurt people.	YSR <sup>a</sup>
	I physically attack people.	YSR
	I show off or clown.	YSR
	I destroy things belonging to others.	YSR
	I get in many fights.	YSR
	I steal things from places other than home.	YSR
	I have been expelled or suspended from school.	DADI <sup>b</sup>
	I ease others a lot.	YSR
	I run away from home.	YSR
	I am mean to others.	YSR
	I swear or use dirty language.	YSR
	I steal things at home.	YSR
	I cut classes or skip school.	YSR

Table 1 (continued):

Items in Scales Created to Measure Internalizing and Externalizing Behavior

SCALE	ITEM	SOURCE
<b>Internalizing Behaviors (<math>\alpha = .74</math>)</b>		
	I feel that I have to be perfect.	YSR
	I feel too guilty.	YSR
	I am unhappy, sad, or depressed.	YSR
	I am nervous or tense.	YSR
	I feel lonely.	YSR
	I have trouble sleeping.	YSR
	I don't have much energy.	YSR
	I feel that no one loves me.	YSR
	I can't get my mind off certain thoughts.	YSR
	I take a positive attitude toward myself. (reverse scored)	SES <sup>c</sup>
	On the whole, I am satisfied with myself. (reverse scored)	SES
	My future seems dark to me.	HS <sup>d</sup>
	Things just won't work out the way I want them to.	HS

<sup>a</sup>Achenbach Youth Self Report; <sup>b</sup>Demographic & Descriptive Interview; <sup>c</sup>Rosenberg Self-Esteem Scale; <sup>d</sup>Beck Hopelessness Scale

**Table 2**  
**Means and Univariate Statistics of Model Variables**  
**for Study Groups, by Gender**

Variable	MALES		FEMALES		Univariate F for group effect (1,122)
	HSB <sup>1</sup> (n=39)	NoHSB <sup>2</sup> (n=56)	HSB <sup>1</sup> (n=45)	NoHSB <sup>2</sup> (n=31)	
	Mean	Mean	Mean	Mean	
Perceived family support	9.24	10.61	9.50	10.71	12.478*
Internalizing Behaviors Scale	11.66	9.02	14.60	11.00	17.84*
Externalizing Behaviors Scale	13.90	7.66	10.15	7.46	19.92*
Age	15.55	14.50	15.18	14.50	8.83

\*p < .01 (with Bonferroni's correction on post-hoc analyses)

**Table 3**  
**Best Fit Models, Based on Logistic Regression Analyses,**  
**for Predicting History of Suicidal Behaviors**  
**In a Sample of Adolescents in Psychiatric Treatment**

Sample	Terms in Model	Accuracy	Specificity	Sensitivity
<b>Males &amp; Females</b>	Gender; Age Externalizing Behaviors; Perceived Support from Family	76%	79%	73%
<b>Males Only</b>	Age; Externalizing Behaviors; Internalizing Behaviors; Perceived Support from Family	81%	83%	72%
<b>Females Only</b>	Perceived Support from Family	70%	60%	76%

**Table 4**

**Relative Risk Estimates For Significant Parameters of Logistic Model**

Sample	Parameter	Relative Risk	95% Confidence Interval Bounds for Relative Risk	
			Lower	Upper
<b>Males &amp; Females</b>				
	Female	1.62	1.01	2.59
	Older Age	1.46	1.11	3.03
	Higher Externalizing Behaviors	1.17	1.06	1.30
	Lower Perceived Support from Family	1.38	1.11	1.71
<b>Males Only</b>				
	Older Age	1.43	1.00	2.08
	Higher Externalizing Behaviors	1.20	1.06	1.35
	Higher Internalizing Behaviors	1.14	0.96	1.36
	Lower Perceived Support from Family	1.46	1.07	2.00
<b>Females Only</b>				
	Lower Perceived Support from Family	1.41	1.03	1.92