

DOCUMENT RESUME

ED 400 104

PS 024 650

TITLE Guidelines for School Health Programs To Promote Lifelong Healthy Eating. Morbidity and Mortality Weekly Report: Recommendations and Reports, Volume 45, June 14, 1996.

INSTITUTION Centers for Disease Control and Prevention (DHHS/PHS), Atlanta, GA.

PUB DATE 14 Jun 96

NOTE 48p.

AVAILABLE FROM Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402-9325.

PUB TYPE Guides - Non-Classroom Use (055)

JOURNAL CIT MMWR; v45 nRR-9, 14 Jun 1996

EDRS PRICE MF01/PC02 Plus Postage.

DESCRIPTORS Adolescents; *Child Health; Community Support; Diseases; *Eating Habits; Health Education; Learning Activities; *Nutrition Instruction; *Policy Formation; Prevention; *School Role; Teaching Methods; Young Children

ABSTRACT

School-based nutrition programs can play an important role in promoting lifelong healthy eating. Healthy eating patterns in childhood and adolescence improve childhood health and prevent health problems. School health programs can help children attain full educational potential and good health by providing them with the skills, social support, and environmental reinforcement they need to adopt long-term, healthy eating behaviors. This report summarizes strategies most likely to be effective in promoting healthy eating among school-age youths and provides nutrition education guidelines for a comprehensive school health program. These guidelines, based on a review of research, theory, and current practices, were developed by Centers for Disease Control (CDC) in collaboration with experts from universities and from national, federal, and voluntary agencies. The guidelines include recommendations on seven aspects of a school-based program to promote healthy eating: (1) school nutrition policy; (2) nutrition education; (3) appropriate instruction for students; (4) integration of school food service and nutrition education; (5) staff training; (6) family and community involvement; and (7) program evaluation. Contains 229 references. (MOK)

 * Reproductions supplied by EDRS are the best that can be made *
 * from the original document. *

MMWR™

Recommendations and Reports

MORBIDITY AND MORTALITY WEEKLY REPORT

ED 400 104

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

This document has been reproduced as received from the person or organization originating it.

Minor changes have been made to improve reproduction quality.

Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

Guidelines for School Health Programs to Promote Lifelong Healthy Eating

PS 024650

BEST COPY AVAILABLE

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
Centers for Disease Control
and Prevention (CDC)
Atlanta, Georgia 30333



The *MMWR* series of publications is published by the Epidemiology Program Office, Centers for Disease Control and Prevention (CDC), Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

Centers for Disease Control and Prevention. Guidelines for school health programs to promote lifelong healthy eating. *MMWR* 1996;45(No. RR-9): [inclusive page numbers].

Centers for Disease Control and Prevention David Satcher, M.D., Ph.D.
Director

The material in this report was prepared for publication by:

National Center for Chronic Disease Prevention
and Health Promotion James S. Marks, M.D.
Director

Division of Adolescent and School Health Lloyd J. Kolbe, Ph.D.
Director

Division of Nutrition and Physical Activity Frederick L. Trowbridge, M.D.
Director

The production of this report as an *MMWR* serial publication was coordinated in:

Epidemiology Program Office Stephen B. Thacker, M.D., M.Sc.
Director

Richard A. Goodman, M.D., M.P.H.
Editor, MMWR Series

Scientific Information and Communications Program

Recommendations and Reports Suzanne M. Hewitt, M.P.A.
Managing Editor

Elizabeth L. Hess
Project Editor

Peter M. Jenkins
Visual Information Specialist

Use of trade names and commercial sources is for identification only and does not imply endorsement by the Public Health Service or the U.S. Department of Health and Human Services.

Copies can be purchased from Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402-9325. Telephone: (202) 783-3238.

Contents

| | |
|---|----|
| Introduction..... | 1 |
| Effects of Diet on the Health, Growth, and Intellectual Development of Young Persons..... | 2 |
| Effects of Childhood Eating Patterns on Chronic Disease Risks of Adults..... | 5 |
| Guidelines for Healthy Eating..... | 7 |
| Eating Behaviors of Children and Adolescents in the United States..... | 8 |
| The Need for School-Based Nutrition Education..... | 9 |
| Promoting Healthy Eating Through a Comprehensive School Health Program..... | 10 |
| Recommendations for School Health Programs Promoting Healthy Eating..... | 11 |
| Conclusion..... | 23 |
| References..... | 24 |
| Appendix A: Nutrition Education Resource List..... | 34 |
| Appendix B: Youth Risk Behavior Surveillance System and School Health Policies and Programs Study..... | 36 |
| Appendix C: Selected School-based Strategies to Promote Healthy Eating..... | 37 |

**Technical Advisors for
Guidelines for School Health Programs
to Promote Lifelong Healthy Eating Patterns**

Tom Baranowski, Ph.D.
University of Texas M.D. Anderson
Cancer Center
Houston, TX

Isobel Contento, Ph.D.
Teachers College, Columbia University
New York, NY

Susan J. Crockett, Ph.D., R.D.
Syracuse University
Syracuse, NY

Shelley Evans, M.A., M.Ed., R.D.*
Pennsylvania State University
University Park, PA

Gail C. Frank, Dr.P.H., R.D.
California State University, Long Beach
Long Beach, CA

Leslie A. Lytle, Ph.D., R.D.
University of Minnesota
Minneapolis, MN

Amanda Dew Manning
U.S. Department of Agriculture
Alexandria, VA

Jeannie McKenzie, Dr.P.H., R.D.*
Pennsylvania State University
University Park, PA

Deanna H. Montgomery, Ph.D., R.D.
University of Texas-Houston
Houston, TX

Rebecca M. Mullis, Ph.D., R.D.
Georgia State University
Atlanta, GA

Christine M. Olson, Ph.D., R.D.
Cornell University
Ithaca, NY

Cheryl L. Perry, Ph.D.
University of Minnesota
Minneapolis, MN

Ken Resnicow, Ph.D.
Emory University
Atlanta, GA

Thomas N. Robinson, M.D., M.P.H.
Stanford University
Palo Alto, CA

Barbara Shannon, Ph.D., R.D.*
Pennsylvania State University
University Park, PA

Howell Wechsler, Ed.D., M.P.H.*
Centers for Disease Control and
Prevention
Atlanta, GA

Participating Agencies and Organizations

| | |
|--|--|
| American Academy of Pediatrics | National Association of Elementary School Principals |
| American Association of Family and Consumer Sciences | National Association of School Nurses |
| American Association of School Administrators | National Association of Secondary School Principals |
| American Cancer Society | National Association of State Boards of Education |
| American Dietetic Association | National Association of State NET Coordinators |
| American Heart Association | National Cancer Institute (USDHHS) |
| American Public Health Association | National Congress of Parents and Teachers |
| American School Food Service Association | National Education Association |
| American School Health Association | National Food Service Management Institute |
| Association for the Advancement of Health Education | National Heart, Lung, and Blood Institute (USDHHS) |
| Association of State and Territorial Directors of Health Promotion and Public Health Education | National School Boards Association |
| Association of State and Territorial Health Officials | National School Health Education Coalition |
| Association of State and Territorial Public Health Nutrition Directors | Office of Disease Prevention and Health Promotion (USDHHS) |
| Council of Chief State School Officers | Society for Nutrition Education |
| Health Resources and Services Administration (U.S. Department of Health and Human Services [USDHHS]) | Society of State Directors of Health, Physical Education, and Recreation |
| Indian Health Service (USDHHS) | U.S. Department of Agriculture |
| Maternal and Child Health Interorganizational Nutrition Group | U.S. Department of Education |

Guidelines for School Health Programs to Promote Lifelong Healthy Eating

Summary

Healthy eating patterns in childhood and adolescence promote optimal childhood health, growth, and intellectual development; prevent immediate health problems, such as iron deficiency anemia, obesity, eating disorders, and dental caries; and may prevent long-term health problems, such as coronary heart disease, cancer, and stroke. School health programs can help children and adolescents attain full educational potential and good health by providing them with the skills, social support, and environmental reinforcement they need to adopt long-term, healthy eating behaviors.

This report summarizes strategies most likely to be effective in promoting healthy eating among school-age youths and provides nutrition education guidelines for a comprehensive school health program. These guidelines are based on a review of research, theory, and current practice, and they were developed by CDC in collaboration with experts from universities and from national, federal, and voluntary agencies.

The guidelines include recommendations on seven aspects of a school-based program to promote healthy eating: school policy on nutrition, a sequential, coordinated curriculum, appropriate instruction for students, integration of school food service and nutrition education, staff training, family and community involvement, and program evaluation.

INTRODUCTION

School-based programs can play an important role in promoting lifelong healthy eating. Because dietary factors "contribute substantially to the burden of preventable illness and premature death in the United States," the national health promotion and disease prevention objectives encourage schools to provide nutrition education from preschool through 12th grade (1). The U.S. Department of Agriculture's (USDA) Nutrition Education and Training (NET) Program urges "nutrition education [to] be a major educational component of all child nutrition programs and offered in all schools, child care facilities, and summer sites" by the year 2000 (2). Because diet influences the potential for learning as well as health, an objective of the first national education goal is that children "receive the nutrition and health care needed to arrive at school with healthy minds and bodies" (3).

The recommendations in this report are intended to help personnel and policymakers at the school, district, state, and national levels meet the national health objectives and education goals by implementing school-based nutrition education policies and programs. This report may also be useful to students, to parents, and to personnel in local and state health departments, community-based health and nutrition programs, pediatric clinics, and training institutions for teachers and public health professionals. These recommendations complement CDC guidelines for school health programs to

prevent the spread of acquired immunodeficiency syndrome (AIDS) (4), to prevent tobacco use and addiction (5), and to promote physical activity (6).

In this report, *nutrition education* refers to a broad range of activities that promote healthy eating behaviors. The nutrition education guidelines focus largely on classroom instruction, but they are relevant to all components of a comprehensive school health program—health education; a healthy environment; health services; counseling, psychological, and social services; integrated school and community efforts; physical education; nutrition services; and school-based health promotion for faculty and staff (7). Although the meals served by school food service programs are an important part of a school health program, this report does not provide specific recommendations related to purchasing and preparing food for school meals. Detailed information on this topic is available from many other publications (8–19) and information sources (see Appendix A). These guidelines also do not address the specific nutrition education and counseling needs of pregnant adolescents (20,21) or young persons with special needs (22–28).

These guidelines are based on a synthesis of research, theory, and current practice and are consistent with the principles of the national health education standards (29), the opportunity-to-learn standards for health education (29), the position papers of leading voluntary organizations involved in child nutrition (30), and the national action plan to improve the American diet (31). To develop these guidelines, CDC convened meetings of experts in nutrition education, reviewed published research, considered the recommendations of national policy documents (1,32–35), and consulted with experts from national, federal, and voluntary organizations.

EFFECTS OF DIET ON THE HEALTH, GROWTH, AND INTELLECTUAL DEVELOPMENT OF YOUNG PERSONS

School-based nutrition education can improve dietary practices that affect young persons' health, growth, and intellectual development. Immediate effects of unhealthy eating patterns include undernutrition, iron deficiency anemia, and overweight and obesity.

Undernutrition

Even moderate undernutrition can have lasting effects on children's cognitive development and school performance (36). Chronically undernourished children attain lower scores on standardized achievement tests, especially tests of language ability (37). When children are hungry or undernourished, they have difficulty resisting infection and therefore are more likely than other children to become sick, to miss school, and to fall behind in class (36,37); they are irritable and have difficulty concentrating, which can interfere with learning (38); and they have low energy, which can limit their physical activity (38). Some reports have estimated that millions of children in the United States experience hunger over the course of a year (39), but no scientific consensus currently exists on how to define or measure hunger (1).

Skipping breakfast can adversely affect children's performance in problem-solving tasks (40–42). A study of low-income elementary school students indicated that those participated in the School Breakfast Program had greater improvements in stand-

ardized test scores and reduced rates of absence and tardiness than did children who qualified for the program but did not participate (43). Twelve percent of students reported skipping breakfast the day before one national survey was taken (44); 40% of 8th- and 10th-grade students in another study reported having eaten breakfast on ≤ 2 days the week before the survey (45). Strategies to encourage adequate nutrition among young persons include the following:

- Promote participation in USDA food assistance programs (e.g., the School Breakfast Program and School Lunch Program, the Summer Food Service Program, and the Child and Adult Care Food Program).
- Advise parents and guardians about community-based food supplementation programs (e.g., food stamps; local food pantries; and the Special Supplemental Nutrition Program for Women, Infants, and Children [WIC]).
- Educate students and their families about the importance of eating breakfast.

Iron Deficiency Anemia

Iron deficiency anemia is the most common cause of anemia in the United States (33). Iron deficiency hampers the body's ability to produce hemoglobin, which is needed to carry oxygen in the blood. This deficiency can increase fatigue, shorten attention span, decrease work capacity, reduce resistance to infection, and impair intellectual performance (33,46). Among school-age youths, female adolescents are at greatest risk for iron deficiency. Approximately 1% of elementary school-age children and 2%–4% of adolescent girls ages 12–19 years show evidence of iron deficiency anemia (47). To prevent iron deficiency, children and adolescents should eat adequate amounts of foods high in iron and in vitamin C, which helps the body absorb iron efficiently (33).

Overweight and Obesity*

Overweight and obesity are increasing among children and adolescents in the United States (48–52). The prevalence of overweight among youths ages 6–17 years in the United States has more than doubled in the past 30 years; most of the increase has occurred since the late 1970s (52). Approximately 4.7 million, or 11%, of youths ages 6–17 years are seriously overweight (52). Obesity in young persons is related to elevated blood cholesterol levels (53–56) and high blood pressure (57–59), and some very obese youths suffer from immediate health problems (e.g., respiratory disorders, orthopedic conditions, and hyperinsulinemia) (60). Being overweight during childhood and adolescence has been associated with increased adult mortality (61,62). Furthermore, obese children and adolescents are often excluded from peer groups and discriminated against by adults, experience psychological stress, and have a poor

*Obesity refers to an excess of total body fat. Body fat content is usually estimated by one of two techniques, measuring skinfold thickness or computing the ratio of body weight to height. Researchers who use weight-to-height ratios tend to use the term "overweight" instead of "obesity." Although weight-to-height ratios correlate highly with body fat, they do not distinguish between body fat and lean body tissue: excess fat tissue is generally assumed to account for the additional weight, but excess weight can also include lean body mass or a large body

body image and low self-esteem (63,64). Increased physical activity and appropriate caloric intake are recommended for preventing and reducing obesity (35). CDC's guidelines for school and community health programs to promote physical activity among youths address strategies for increasing physical activity among young persons (6).

Unsafe Weight-Loss Methods

Many young persons in the United States practice unsafe weight-loss methods. Deliberately restricting food intake over long periods can lead to poor growth and delayed sexual development (65). Data from one study indicated that the rate of smoking initiation is higher for adolescent girls who diet or who are concerned about their weight than for nondieters or girls having few weight concerns (66), and another study indicated that many white female high school students who smoke report using smoking to control their appetite and weight (67). Harmful weight loss practices have been reported among girls as young as 9 years old (68,69). Young persons involved in certain competitive sports and dancing are particularly at risk for unsafe weight control practices (70). A national survey of 8th- and 10th-grade students found that 32% skipped meals, 22% fasted, 7% used diet pills, 5% induced vomiting after meals, and 3% used laxatives to lose weight (45). Children and adolescents should learn about the dangers of unsafe weight-loss methods and about safe ways to maintain a healthy weight. The emphasis of society in the United States on thinness should be challenged, and young persons need to develop a healthy body image (71).

Eating Disorders

Eating disorders (e.g., anorexia nervosa and bulimia nervosa) are psychological disorders characterized by severe disturbances in eating behavior. Anorexia nervosa is characterized by a refusal to maintain a minimally normal body weight, and bulimia nervosa is characterized by repeated episodes of binge eating followed by compensatory behaviors such as self-induced vomiting (72). Eating disorders often start in adolescence, and >90% of cases occur among females (72). Anorexia nervosa and bulimia nervosa affect as many as 3% of adolescent and young adult females, and the incidence of anorexia nervosa appears to have increased in recent decades (72). Compared with adolescents who have normal eating patterns, adolescents who have eating disorders tend to have lower self-esteem; a negative body image; and feelings of inadequacy, anxiety, social dysfunction, depression, and moodiness (73). Eating disorders can cause many severe complications, and mortality rates for these disorders are among the highest for any psychiatric disorder (74). Persons who have eating disorders should receive immediate medical and psychological treatment.

Dental Caries

Dental caries is perhaps the most prevalent of all diseases (1). It affects 50.1% of youths ages 5–17 years and 84.4% of youths age 17 years (75). More than 50 million hours of school time are lost annually because of dental problems or dental visits (76). Dental caries is a progressive disease, which, if left untreated, can result in acute infections, pain, costly treatment, and tooth loss. A strong link exists between sugar

consumption and dental caries (33). To prevent dental caries, children and adolescents should drink fluoridated water, use fluoridated toothpaste, brush and floss their teeth regularly, have dental sealants applied to the pits and fissures of their teeth, and consume sugars in moderation (7).

EFFECTS OF CHILDHOOD EATING PATTERNS ON CHRONIC DISEASE RISKS OF ADULTS

Nutrition education also should focus on preventing children and adolescents from developing chronic diseases during adulthood. Some of the physiological processes that lead to diet-related chronic disease begin in childhood. For example, autopsy studies have demonstrated that early indicators of atherosclerosis (the hardening of the arteries that is the most common cause of coronary heart disease [CHD]) begin in youth (77–83) and are related to blood cholesterol levels in young persons (79,81–83). Unhealthy eating practices that contribute to chronic disease are established early in life; young persons having unhealthy eating habits tend to maintain these habits as they age (84). Thus, it is efficacious to teach persons healthy eating patterns when they are young; high-risk eating behaviors and physiological risk factors are difficult to change once they are established during youth.

Diet-related risk factors for cardiovascular disease (e.g., high blood cholesterol level, high blood pressure, and overweight) are common in youths in the United States (34,52,85–90). Compared with their peers, children and adolescents who have high blood cholesterol (34,91–96), have high blood pressure (97,98), or are obese (91,99–103) are more likely to have these risk factors during adulthood. Poor diet and inadequate physical activity together account for at least 300,000 deaths in the United States annually and are second only to tobacco use as the most prominent identifiable contributor to premature death (104). Interventions that promote healthy eating and physical activity behaviors during childhood and adolescence may not only prevent some of the leading causes of illness and death but also decrease direct health-care costs and improve quality of life.

Diet is a known risk factor for the development of the nation's three leading causes of death: CHD, cancer, and stroke (33). Other health problems of adulthood associated with diet are diabetes, high blood pressure, overweight, and osteoporosis.

Coronary Heart Disease

CHD kills more persons in the United States than any other disease does (7). Diet-related risk factors for CHD include high blood cholesterol, high blood pressure, and obesity. These risk factors can be reduced by consuming less fat (particularly saturated fat) and cholesterol and by increasing physical activity (105).

Cancer

One out of every five deaths in the United States is attributable to cancer (106). Dietary factors have been associated with several types of cancer, including colon, breast, and prostate (33). All cancer deaths in the United States might be reduced as much as 35% through dietary changes (107,108). The risk for some types of cancer reduced by maintaining a healthy weight; limiting consumption of fat, alcohol,



and salt-cured, salt-pickled, or smoked foods; and eating more foods that protect the body against cancer (fruits, vegetables, whole grain cereals, and other high-fiber foods) (109). The National Cancer Institute advises eating at least five servings of fruits and vegetables each day (110).

Stroke

Cerebrovascular disease, or stroke, is the third leading cause of death in the United States and a major cause of illness and disability (111). The most important risk factor for stroke is high blood pressure, which often can be controlled or prevented by adopting a healthy diet and maintaining a healthy weight (112). The risk for stroke can be reduced by consuming less sodium, increasing physical activity, and maintaining a healthy body weight.

Diabetes

Diabetes is the seventh leading cause of death in the United States (104). CHD is two to four times more common and stroke is two to six times more common in persons who have diabetes than in persons who do not have diabetes (113). Diabetes can lead to blindness, kidney disease, and nerve damage (113). Non-insulin-dependent diabetes mellitus, which affects approximately 90% of persons who have diabetes, is often associated with obesity (114). Maintaining a desirable body weight through physical activity and modest caloric restriction is important in preventing diabetes and controlling its complications (114).

High Blood Pressure

High blood pressure is a major cause of CHD, stroke, and kidney failure. About one in four adults in the United States has high blood pressure (115). Persons who have high blood pressure have three to four times the risk of developing CHD and as much as seven times the risk of stroke as do those who have normal blood pressure (116). Persons can reduce their risk for high blood pressure by consuming less sodium, increasing physical activity, and maintaining a healthy body weight. A diet high in potassium may help reduce the risk of high blood pressure (117).

Overweight

In the United States, about one in three adults is overweight (118), and these persons are at increased risk for CHD, some types of cancer, stroke, diabetes mellitus, high blood pressure, and gallbladder disease (33). Overall risk for premature death is increased by excess weight; the risk increases as severity of overweight increases (33). The best way to lose weight is to increase physical activity and control caloric intake, preferably by adopting a diet that is low in fat and high in vegetables, fruits, and grains (35).

Osteoporosis

Osteoporosis is a decrease in the amount of bone so severe that the bone fractures easily. About 1.3 million bone fractures, including many fatal hip fractures, occur per

year in persons ≥ 45 years of age (33). Low dietary calcium, a mineral essential for bone growth, may be associated with an increased risk for osteoporosis (33). For females especially, eating enough calcium is particularly important during childhood, adolescence, and young adulthood—when bones approach their maximum density—to reduce the risk for osteoporosis later in life (1,119–122). Regular weight-bearing exercises also can help prevent osteoporosis (33).

GUIDELINES FOR HEALTHY EATING

To prevent certain diseases and to promote good health, persons >2 years of age should follow the seven recommendations that constitute the Dietary Guidelines for Americans (35). These guidelines are developed by the USDA and USDHHS and are published every 5 years. They are based on extensive reviews of hundreds of studies conducted over many years and represent the best current advice that nutrition scientists can give. The guidelines are consistent with dietary recommendations made by major health promotion organizations, including the National Research Council (32), the National Cholesterol Education Program of the National Institutes of Health (34,105), the National Cancer Institute (109), the American Cancer Society (123), and the American Heart Association (124).

The principles contained in the Dietary Guidelines for Americans should be the primary focus of school-based nutrition education. By enabling young persons to adopt practices consistent with the guidelines, schools can help the nation meet its health objectives (1), which were designed to guide health promotion and disease prevention policy and programs at the federal, state, and local level throughout the 1990s. Objective 2.19 is to “increase to at least 75 percent the proportion of the Nation’s schools that provide nutrition education from preschool through 12th grade, preferably as part of quality school health education” (1). The six relevant dietary guidelines are (a) eat a variety of foods; (b) balance the food you eat with physical activity—maintain or improve your weight; (c) choose a diet with plenty of grain products, vegetables, and fruits; (d) choose a diet low in fat, saturated fat, and cholesterol; (e) choose a diet moderate in sugars; and (f) choose a diet moderate in salt and sodium. (The seventh recommendation concerns adults and alcoholic beverages.) Enabling children and adolescents to follow these guidelines can help the nation achieve these national health objectives for the year 2000 (1):

- 2.3** Reduce overweight to a prevalence of $\leq 20\%$ among people aged 20 and older and $\leq 15\%$ among adolescents aged 12 through 19.
- 2.5** Reduce average dietary fat intake to $\leq 30\%$ of calories and average saturated fat intake to $\leq 10\%$ of calories among people aged two and older.
- 2.6** Increase complex carbohydrates and fiber-containing foods in the diets of adults to five or more daily servings for vegetables (including legumes) and fruits and to six or more daily servings for grain products.
- 2.7** Increase to $\geq 50\%$ the proportion of overweight people aged 12 and older who have adopted sound dietary practices combined with regular physical activity to attain an appropriate body weight.
- 2.8** Increase calcium intake so $\geq 50\%$ of youth aged 12 through 24 and $\geq 50\%$ of pregnant and lactating women consume three or more servings daily of foods rich in calcium, and $\geq 50\%$ of people aged ≥ 25 consume two or more servings daily.


2.9 Decrease salt and sodium intake so that $\geq 65\%$ of home meal preparers prepare foods without adding salt, $\geq 80\%$ of people avoid using salt at the table, and $\geq 40\%$ of adults regularly purchase foods modified or lower in sodium.

The Food Guide Pyramid (Figure 1) was designed by the USDA and USDHHS to help persons follow the Dietary Guidelines for Americans. Schools can use the pyramid to illustrate the concepts of variety (eat different foods from among and within the food groups), moderation (limit the consumption of foods high in fat and added sugars), and proportionality (eat relatively greater amounts of foods from the groups that are lower in the pyramid: grains, vegetables, and fruits) (125). Other educational materials supplement the pyramid by listing low-fat choices within each food group (35).

EATING BEHAVIORS OF CHILDREN AND ADOLESCENTS IN THE UNITED STATES

Many young persons in the United States do not follow the recommendations of the Dietary Guidelines for Americans or the Food Guide Pyramid. On average, children and adolescents consume too much fat, saturated fat, and sodium and not enough fruits, vegetables, or calcium (44,126-129 ; CDC, unpublished data). Children and adolescents obtain 33%-35% of their calories from fat and 12%-13% from saturated fat

FIGURE 1. The Food Guide Pyramid — a guide to daily choices



Source: U.S. Department of Agriculture/U.S. Department of Health and Human Services

(above the recommended levels of 30% and 10%, respectively) (44,128,129). Only 16% of children ages 6–11 years and 15% of adolescents ages 12–19 years meet the recommendation for total fat intake; only 9% of children and 7% of adolescents meet the recommendation for saturated fat intake (130). Almost one-half of 8th- and 10th-grade students eat three or more snacks a day, and most of these snacks are high in fat, sugar, or sodium (45).

Unpublished data from CDC's 1993 Youth Risk Behavior Survey indicated that, on the day before the survey, 41% of high school students in the United States ate no vegetables and 42% ate no fruits (127) (Appendix B). An analysis of a nationally representative sample of youths ages 2–18 years indicated that, over a 3-day period, the youths ate only 3.6 servings of fruits and vegetables daily and that fried potatoes accounted for a large proportion of the vegetables consumed, 20.4% of the youths ate the recommended five or more servings of fruits and vegetables daily, 50.8% ate fewer than one serving of fruit per day, and 29.3% ate fewer than one serving per day of vegetables that were not fried (131). Adolescent females eat considerably less calcium and iron than recommended by the Food and Nutrition Board of the National Research Council (126,129).

Children and adolescents appear to be familiar with the general relationship between nutrition and health but are less aware of the relationship between specific foods and health. For example, young persons understand the importance of limiting fat, cholesterol, and sodium in one's diet, but they do not know which foods are high in fat, cholesterol, sodium, or fiber (45,132,133). One study indicated that adolescents were well-informed about good nutrition and health but did not use their knowledge to make healthy food choices (134).

THE NEED FOR SCHOOL-BASED NUTRITION EDUCATION

Young persons need nutrition education to help them develop lifelong eating patterns consistent with the Dietary Guidelines for Americans and the Food Guide Pyramid. Schools are ideal settings for nutrition education for several reasons:

- Schools can reach almost all children and adolescents.
- Schools provide opportunities to practice healthy eating. More than one-half of youths in the United States eat one of their three major meals in school, and 1 in 10 children and adolescents eats two of three main meals in school (135).
- Schools can teach students how to resist social pressures. Eating is a socially learned behavior that is influenced by social pressures. School-based programs can directly address peer pressure that discourages healthy eating and harness the power of peer pressure to reinforce healthy eating habits.
- Skilled personnel are available. After appropriate training, teachers can use their instructional skills and food service personnel can contribute their expertise to nutrition education programs.
- Evaluations suggest that school-based nutrition education can improve the eating behaviors of young persons (136–138).

School-based nutrition education is particularly important because today's children and adolescents frequently decide what to eat with little adult supervision (139). The increase in one-parent families or families having two working parents and the availability of convenience foods and fast-food restaurants inhibit parents' monitoring of their children's eating habits.

Young persons' food choices are influenced by television advertisements for low-nutritive foods. Young persons see about one food advertisement for every 5 minutes of Saturday morning children's shows (140). Most of the foods advertised during children's programming are high in fat, sugar, or sodium; practically no advertisements are for healthy foods such as fruits and vegetables (140-142). Studies have indicated that, compared with those who watch little television, children and adolescents who watch more television are more likely to have unhealthy eating habits and unhealthy conceptions about food (143), ask their parents to buy foods advertised on television (144), and eat more fat (145). Some studies of young persons have found that television watching is directly associated with obesity (146-149). Because youths in the United States spend, on average, more than 20 hours a week watching television (150)—more time over the course of the year than they are in school (141)—school-based programs should help counter the effect of television on young persons' eating habits.

Schools are a critical part of the social environment that shapes young persons' eating behaviors and can therefore play a large role in helping improve their diet. However, schools cannot achieve this goal on their own when the cultural milieu has a large influence on food-related beliefs, values, and practices (30,138). Families, food stores, restaurants, the food industry, religious institutions, community centers, government programs, and the mass media must also support the principles of the Dietary Guidelines for Americans. The USDA's Team Nutrition (see Appendix A) seeks to gain the support of many sectors of society for improving the diet of young persons by creating innovative public and private partnerships that promote healthy food choices through the media, schools, families, and community (151).

PROMOTING HEALTHY EATING THROUGH A COMPREHENSIVE SCHOOL HEALTH PROGRAM

In the school environment, classroom lessons alone might not be enough to effect lasting changes in students' eating behaviors (30); students also need access to healthy food and the support of persons around them (137). The influence of school goes beyond the classroom and includes normative messages from peers and adults regarding foods and eating patterns. Students are more likely to receive a strong, consistent message when healthy eating is promoted through a comprehensive school health program.

A comprehensive school health program empowers students with not only the knowledge, attitudes, and skills required to make positive health decisions but also the environment, motivation, services, and support necessary to develop and maintain healthy behaviors (152). A comprehensive school health program includes health education; a healthy environment; health services; counseling, psychological, and social services; integrated school and community efforts; physical education; nutrition services; and a school-based health program for faculty and staff (7). Each compo-

ment can contribute to integrated efforts that promote healthy eating. For example, classroom lessons on nutrition can be supported by

- schools providing appealing, low-fat, low-sodium foods in vending machines and at school meetings and events;
- school counselors and nurses providing guidance on health and, if necessary, referrals for nutritional problems;
- community organizations providing counseling or nutrition education campaigns;
- physical education instructors helping students understand the relationship between nutrition and physical activity;
- school food service personnel serving healthy, well-balanced meals in the cafeteria; and
- school personnel acting as role models for healthy eating (153).

The USDA is promoting health-enhancing changes in the food service component of the school health program by requiring schools to serve meals that comply with the Dietary Guidelines for Americans (154) and by providing technical support to schools through Team Nutrition (151).

RECOMMENDATIONS FOR SCHOOL HEALTH PROGRAMS PROMOTING HEALTHY EATING

Based on the available scientific literature, national nutrition policy documents, and current practice, these guidelines provide seven recommendations for ensuring a quality nutrition program within a comprehensive school health program. These recommendations address school policy on nutrition, a sequential, coordinated curriculum, appropriate and fun instruction for students, integration of school food service and nutrition education, staff training, family and community involvement, and program evaluation. Strategies that schools can use to achieve these recommendations are available (Appendix C). However, local school systems need to assess the nutrition needs and issues particular to their communities, and they need to work with key school- and community-based constituents, including students, to develop the most effective and relevant nutrition education plans for their communities. Vigorous, coordinated, and sustained support from communities, local and state education and health agencies, institutions of higher education, and national organizations also is necessary to ensure success (29).

1. **Policy:** Adopt a coordinated school nutrition policy that promotes healthy eating through classroom lessons and a supportive school environment.
2. **Curriculum for nutrition education:** Implement nutrition education from pre-school through secondary school as part of a sequential, comprehensive school health education curriculum designed to help students adopt healthy eating behaviors.

3. **Instruction for students:** Provide nutrition education through developmentally appropriate, culturally relevant, fun, participatory activities that involve social learning strategies.
4. **Integration of school food service and nutrition education:** Coordinate school food service with nutrition education and with other components of the comprehensive school health program to reinforce messages on healthy eating.
5. **Training for school staff:** Provide staff involved in nutrition education with adequate preservice and ongoing in-service training that focuses on teaching strategies for behavioral change.
6. **Family and community involvement:** Involve family members and the community in supporting and reinforcing nutrition education.
7. **Program evaluation:** Regularly evaluate the effectiveness of the school health program in promoting healthy eating, and change the program as appropriate to increase its effectiveness.

Recommendation 1. Policy: Adopt a coordinated school nutrition policy that promotes healthy eating through classroom lessons and a supportive school environment.

Rationale for the Policy

A coordinated school nutrition policy, particularly as part of an overall school health policy, provides the framework for implementing the other six recommendations and ensures that students receive nutrition education messages that are reinforced throughout the school environment. For example, such a policy would address nutrition education classes; school lunch and breakfast; classroom snacks and parties; use of food to reward or discipline; and food sold in vending machines, at school stores, snack bars, sporting events, and special activities, and as part of fundraising activities. The school environment can powerfully influence students' attitudes, preferences, and behaviors related to food (137). Without a coordinated nutrition policy, schools risk negating the health lessons delivered in the classroom and cafeteria by allowing actions that discourage healthy eating behaviors.

Developing the Policy

A school nutrition policy should be a brief document that incorporates input from all relevant constituents of the school community: students, teachers, coaches, staff, administrators, food service personnel, nurses, counselors, public health professionals, and parents. The policy should meet local needs and be adapted to the health concerns, food preferences, and dietary practices of different ethnic and socioeconomic groups. Technical assistance for assessing nutrition education needs is available through the state NET Program (155). Schools might consider using one or more of the following techniques to assess their particular needs:

- Interview nutrition professionals to learn more about local eating habits and to identify materials and services available for youths and adults. Schools might interview representatives from the school food service program; the state NET Program; the nutrition unit within the State Department of Health; the district or

state school health coordinator; the local WIC program and Cooperative Extension nutrition education program; the state or local chapters of the American Cancer Society, American Dietetic Association, and the American Heart Association; nutrition councils or coalitions; university research programs; organizations with special insights into the particular nutrition education needs of cultural and ethnic minorities; or businesses that offer nutrition-related services or food products.

- Interview food service staff about students' eating practices in the school cafeteria.
- Observe the school cafeteria, the teachers' lunchroom, and other areas in the school where food is available.
- Review nutrition curricula used by teachers.
- Survey teachers to determine how nutrition is taught, whether teachers use food for reward or punishment, and the level of interest of teachers in nutrition or wellness programs for themselves.
- Survey students to determine their dietary preferences and what types of healthy changes in school food they most want.

The policy plan should include means of obtaining follow-up input from all parties and means of revising the plan as needed. Student involvement is critical to the success of a nutrition policy. A nutrition advisory committee or a nutrition subcommittee of the school health advisory council having student members can develop and promulgate a coordinated school nutrition policy. Technical assistance in forming a school nutrition advisory committee is available from the American School Food Service Association (Appendix A). Successful implementation of a nutrition policy also requires the active support of school and district educational leadership.

Content of the Policy

The written policy should describe the importance of the nutrition component within the comprehensive school health program. This section can briefly describe the role of good nutrition in promoting childhood growth, health, and learning; discuss the role of child and adolescent nutrition in reducing the risk for chronic diseases of adulthood; identify the importance of establishing a school environment that supports healthy eating choices by young persons; and generate support for the policy by identifying how improvements in student nutrition can satisfy the needs of different constituents of the school community (e.g., students, teachers, and food service personnel). An optimal policy on nutrition should publicly commit the school to providing adequate time for a curriculum on nutrition, serving healthy and appealing foods at school, developing food-use guidelines for teachers, supporting healthy school meals, and establishing links with nutrition service providers.

Curriculum. Adequate time should be allocated for nutrition education throughout the preschool, primary, and secondary school years as part of a sequential, comprehensive school health education program. In addition, teachers should be adequately trained to teach nutrition and be provided with ongoing in-service training.

Healthy[†] and appealing foods. Healthy and appealing foods should be available in meals, a la carte items in the cafeteria, snack bars, and vending machines (Exhibit 1); as classroom snacks; and at special events, athletic competitions, staff meetings, and parents' association meetings. In addition, schools should discourage the sale of foods high in fat, sodium, and added sugars (e.g., candy, fried chips, and soda) on school grounds and as part of fundraising activities. Although selling low-nutritive foods may provide revenue for school programs, such sales tell students that it is acceptable to compromise health for financial reasons (158). The school thereby risks contradicting the messages on healthy eating given in class. If schools contract with food service management companies to supply meals, the contractors should be required to serve appealing, low-fat, low-sodium meals that comply with the standards of the Dietary Guidelines for Americans.

EXHIBIT 1: Sample List of Vending Machine Foods Low in Saturated Fat (34)

Canned fruit
 Fresh fruit (e.g., apples and oranges)*
 Fresh vegetables (e.g., carrots)
 Fruit juice and vegetable juice
 Low-fat crackers and cookies, such as fig bars and gingersnaps
 Pretzels
 Bread products (e.g., bread sticks, rolls, bagels, and pita bread)
 Ready-to-eat, low-sugar cereals
 Granola bars made with unsaturated fat
 Low-fat (1%) or skim milk*
 Low-fat or nonfat yogurt*
 Snack mixes of cereal and dried fruit with a small amount of nuts and seeds[†]
 Raisins and other dried fruit[†]
 Peanut butter and low-fat crackers[§]

*These foods are appropriate if the vending machine is refrigerated.

[†]Some schools might not want to offer these items because these foods can contribute to dental caries.

[§]Some schools might not want to offer peanut butter; although it is low in saturated fatty acids, peanut butter is high in total fat.

[†]As defined by the U.S. Food and Drug Administration (156,157) in its food label regulations, a "healthy" food must be low in fat (≤ 3 g per serving), be low in saturated fat (≤ 1 g per serving), contain limited amounts of cholesterol (≤ 60 mg per serving for a single-item food), and contain limited amounts of salt (≤ 480 mg per serving until 1998, when the criterion for a single-item food will decrease to ≤ 360 mg per serving). In addition, single-item foods that are not raw fruits or vegetables must provide $\geq 10\%$ of the daily value of one or more of the following nutrients: vitamin A, vitamin C, iron, calcium, protein, and fiber. Criteria for products that include more than one type of food (e.g., macaroni and cheese) vary depending on the food.

Food use guidelines for teachers. Schools should discourage teachers from using food for disciplining or rewarding students. Some teachers give students low-nutritive foods, such as candy, as a reward for good behavior, and punish misbehaving students by denying a low-nutritive treat (159). These practices reinforce students' preferences for low-nutritive foods and contradict what is taught during nutrition education. Schools should recommend that both teachers and parents serve healthy party snacks and treats (160).

Support for healthy school meals. Starting with the 1996–1997 school year, schools will be required to serve meals that comply with the standards of the Dietary Guidelines for Americans (154). To encourage students to participate in school meal programs and to make healthy choices in cafeterias, schools can use marketing-style incentives and promotions (13, 14, 135, 161); use healthy school meals as examples in class; educate parents about the value of healthy school meals; involve students and parents in planning meals; and have teachers, administrators, and parents eat in the cafeteria and speak favorably about the healthy meals available there. Students should also be given adequate time and space to eat meals in a pleasant and safe environment (162).

Links with nutrition service providers. Schools should establish links with qualified public health and nutrition professionals who can provide screening, referral, and counseling for nutritional problems (30, 163); inform families about supplemental nutrition services available in the community, such as WIC (164), food stamps, local food pantries, the Summer Food Service Program, and the Child and Adult Care Food Program; and implement nutrition education and health promotion activities for school faculty, other staff, school board members, and parents. These links can help prevent and resolve nutritional problems that can impair a student's capacity to learn, demonstrate the value placed on good nutrition for the entire school community, and help adults serve as role models for school-age youths.

Recommendation 2. Curriculum for nutrition education: Implement nutrition education from preschool through secondary school as part of a sequential, comprehensive school health education curriculum designed to help students adopt healthy eating behaviors.

Nutrition Education as Part of a Comprehensive School Health Education Program

Nutrition education should be part of a comprehensive health education curriculum that focuses on understanding the relationship between personal behavior and health. This curriculum should give students the knowledge and skills they need to be "health literate," as delineated by the national health education standards (29) (Exhibit 2). The comprehensive health education approach is important to nutrition education because

EXHIBIT 2: National Health Education Standards (29)

1. Students will comprehend concepts related to health promotion and disease prevention.
2. Students will be able to access valid health information and health-promoting products and services.
3. Students will be able to practice health-enhancing behaviors and reduce health risks.
4. Students will analyze the influence of culture, media, technology, and other factors on health.
5. Students will be able to use interpersonal communication skills to enhance health.
6. Students will be able to use goal-setting and decision-making skills to enhance health.
7. Students will be able to advocate for personal, family, and community health.

- unhealthy eating behaviors may be interrelated with other health risk factors (e.g., cigarette smoking and sedentary lifestyle) (165),
- nutrition education shares many of the key goals of other health education content areas (e.g., raising the value placed on health, taking responsibility for one's health, and increasing confidence in one's ability to make health-enhancing behavioral changes), and
- state-of-the-art nutrition education uses many of the social learning behavioral change techniques used in other health education domains.

Therefore, nutrition education activities can reinforce, and be reinforced by, activities that address other health education topics as well as health in general.

Linking nutrition and physical activity is particularly important because of the rising proportion of overweight youths in the United States. Nutrition education lessons should stress the importance of combining regular physical activity with sound nutrition as part of an overall healthy lifestyle. Physical education classes, in turn, should include guidance in food selection (6).

Sequential Lessons and Adequate Time

Students who receive more lessons on nutrition have more positive behavioral changes than students who have fewer lessons (166,167). To achieve stable, positive changes in students' eating behaviors, adequate time should be allocated for nutrition education lessons. The curriculum should be sequential from preschool through secondary school; attention should be paid to scope and sequence. When designing the curriculum, schools should assess and address their students' needs and concerns. A curriculum targeted to a limited number of behaviors might make the most effective use of a scarce instructional time available for nutrition education (136).

To maximize classroom time, nutrition education can be integrated into the lesson plans of other school subjects; for example, math lessons could analyze nutrient intake or reading lessons could feature texts on nutrition (168). Little research on the integrated approach has been conducted (137), but embedding information on nutrition in other courses probably reinforces the goals of nutrition education. However, the exclusive use of an integrative approach might sacrifice key elements of an effective nutrition education program (e.g., adequate time, focusing on behaviors and skill-building, attention to scope and sequence, and adequate teacher preparation) (137). Therefore, integration into other courses can complement but should not replace sequential nutrition education lessons within a comprehensive school health education curriculum. Classroom time can be maximized also by having nutrition education lessons use skills learned in other classes (e.g., math or language arts) (169–171).

Organizations and agencies can supply information on specific nutrition education curricula and materials (Appendix A). The USDA's NET Program provides technical assistance in school-based nutrition education (2,172). The Food and Nutrition Information Center of USDA's National Agricultural Library provides information on nutrition education evaluation and resources and serves as a national depository and lending library for NET materials. Nutritionists at some organizations can also answer specific nutrition content questions (Appendix A).

Focusing on Promoting Healthy Eating Behaviors

The primary goal of nutrition education should be to help young persons adopt eating behaviors that will promote health and reduce risk for disease. Knowing how and why to eat healthily is important, but knowledge alone does not enable young persons to adopt healthy eating behaviors (137). Cognitive-focused curricula on nutrition education typically result in gains in knowledge but usually have little effect on behavior (173–178).

Behaviorally based education encourages specific healthy eating behaviors (e.g., eating less fat and sodium and eating more fruits and vegetables) (136,179); however, it does not detail the technical and scientific knowledge on which dietary recommendations are based and, therefore, might not fulfill science education requirements (180). The strategies listed in Appendix C can be used as central concepts in a behaviorally based nutrition education program.

Several programs using a behavioral approach have achieved significant ($p < 0.05$), positive changes in students' eating behaviors (167,181–190). Compared with students in control schools, students in some behaviorally based health and nutrition education programs had significant ($p < 0.05$), favorable changes in serum cholesterol levels (167,188,191), blood pressure level (167,191), and body mass index (184). Although most of the behaviorally oriented programs did not achieve all their behavioral aims—perhaps because of the limited amount of curriculum time (136)—current scientific knowledge indicates that a focus on behavior is a key determinant in the success of nutrition education programs (136–138).

Recommendation 3. Instruction for students: Provide nutrition education through developmentally appropriate, culturally relevant, fun, participatory activities that involve social learning strategies.

Developmentally Appropriate and Culturally Relevant Activities

Different educational strategies should be used for young persons at different stages of cognitive development. Regardless of the amount and quality of teaching they receive, young elementary schoolchildren might not fully understand abstract concepts (e.g., the nutrient content of foods or the classification of foods into groups) (192–194). Nutrition education for young children should focus on concrete experiences (e.g., increasing exposure to many healthy foods and building skills in choosing healthy foods) (169).

More abstract associations between nutrition and health become appropriate as children approach middle school. By this age, children can understand and act on the connection between eating behaviors and health (137,194). Nutrition education for middle and high school students should focus on helping students assess their own eating behaviors and set goals for improving their food selection (138,195). Lessons for older children should emphasize personal responsibility, decision-making skills, and resisting negative social pressures (183,185,187,189).

Nutrition education presents opportunities for young persons to learn about and experience cultural diversity related to food and eating. Students from different cultural groups have different health concerns, eating patterns, food preferences, and food-related habits and attitudes. These differences need to be considered when designing lesson plans or discussing food choices. Nutrition education can succeed only when students believe it is relevant to their lives.

Active Learning and an Emphasis on Fun

The context in which students learn about healthy eating behaviors and the feelings students associate with healthy foods are key factors in determining their receptivity to nutrition education. Students are more likely to adopt healthy eating behaviors when

- they learn about these behaviors through fun, participatory activities rather than through lectures (138,196,197);
- lessons emphasize the positive, appealing aspects of healthy eating patterns rather than the negative consequences of unhealthy eating patterns;
- the benefits of healthy eating behaviors are presented in the context of what is already important to the students; and
- the students have repeated opportunities to taste foods that are low in fat, sodium, and added sugars and high in vitamins, minerals, and fiber during their lessons.⁵

⁵When serving food, teachers must use hygienic food handling practices and consider possible allergies and religious prohibitions; the food service director can help in this area.

Computer-based lessons on nutrition can also be effective (198), especially when teacher time is limited or when student self-assessment is appropriate. Interactive, highly entertaining, and well-designed computer programs are now available to help young persons learn healthy food selection skills and assess their own diets (199,200). Computer-based lessons allow students to move at their own pace and can capture their attention.

Social Learning Techniques

Most of the nutrition education programs that have resulted in behavioral change have used teaching strategies based on social learning theory (195,201–205). In such lessons, increasing student knowledge is only one of many objectives. Social learning instruction also emphasizes

- raising the value students place on good health and nutrition and identifying the benefits of adopting healthy eating patterns, including short-term benefits that are important to young persons (e.g., physical appearance, sense of personal control and independence, and capacity for physical activities);
- giving students repeated opportunities to taste healthy foods, including foods they have not yet tasted;
- working with parents, school personnel, public health professionals, and others to overcome barriers to healthy eating;
- using influential role models, including peers, to demonstrate healthy eating practices;
- providing incentives (e.g., verbal praise and small prizes) to reinforce messages;
- helping students develop practical skills for and self-confidence in planning meals, preparing foods, reading food labels, and making healthy food choices through observation and hands-on practice;
- enabling students to critically analyze sociocultural influences, including advertising, on food selection, to resist negative social pressures, and to develop social support for healthy eating; and
- helping students analyze their own eating patterns, set realistic goals for changes in their eating behaviors, monitor their progress in reaching those goals, and reward themselves for achieving their goals.

Nutrition education strategies include social learning techniques (Appendix C).

Recommendation 4. Integration of school food service and nutrition education: Coordinate school food service with nutrition education and with other components of the comprehensive school health program to reinforce messages on healthy eating.

The school cafeteria provides a place for students to practice healthy eating. This experience should be coordinated with classroom lessons to allow students to apply critical thinking skills taught in the classroom (2,8,9,11–15,18,169,178,206). School food service personnel can

- visit classrooms and explain how they make sure meals meet the standards of the Dietary Guidelines for Americans,
- invite classes to visit the cafeteria kitchen and learn how to prepare healthy foods,
- involve students in planning the school menu and preparing recipes,
- offer foods that reinforce classroom lessons (e.g., whole wheat rolls to reinforce a lesson on dietary fiber),
- post in the cafeteria posters and fliers on nutrition, and
- display nutrition information about available foods and give students opportunities to practice food analysis and selection skills learned in the classroom.

In addition, classroom teaching can complement the goals of the school food service. For example, teachers can help food service managers by teaching students about the importance of nutritious school meals and getting feedback from students on new menu items developed to meet the goals set by USDA's School Meals Initiatives for Healthy Children (154).

To ensure consistent nutrition messages from the school, food service personnel should work closely with those responsible for other components of the school health program. For example, the personnel can

- help develop and implement school policies that make healthful foods available;
- educate parents about the value of school meals (e.g., put health messages in monthly menus sent home to parents or make periodic presentations at parents' association meetings) (11,13);
- help schools access and assess community public health and nutrition services; and
- keep classroom teachers, physical education teachers, coaches, counselors, health-service providers, and other staff informed about the importance of healthy school meals.

Recommendation 5. Training for school staff: Provide staff involved in nutrition education with adequate preservice and ongoing in-service training that focuses on teaching strategies for behavioral change.

Training in nutrition and health education can increase the extent to which teachers implement a curriculum (207–209), which in turn affects the likelihood that students' eating behaviors will change (167,207). All elementary school teachers as well as secondary school teachers in disciplines such as home economics, family and consumer sciences, language arts, physical education, and science should receive nutrition education training. State NET Programs can provide technical assistance for training teachers in nutrition education (Appendix A).

Training should address content and teaching strategies. Because classroom teachers often need more help with innovative nutrition teaching techniques than with content (195,210), training should focus on giving teachers the skills they need to use the nonlecture, active learning methods discussed previously (195). Training programs are most effective if they

- are designed to meet the specific needs of the teachers and are based on the teachers' level of nutrition knowledge and experience with the suggested teaching strategies,
- model behavioral change techniques and give teachers practice in using them,
- involve multiple sessions spaced across time so that teachers can try out the newly learned techniques in their classrooms and report on their experiences to the training group, and
- provide posttraining sessions so that teachers can share experiences with their peers (211,212).

Teachers should understand the importance of fully implementing the selected curriculum and become familiar with its underlying theory and concepts. Training should also help teachers assess and improve their own eating practices and make them aware of the behavioral messages they give as role models (213).

Continuing education activities in nutrition education should be offered to food service personnel so this staff can reinforce classroom instruction through the school meal program and help shape the school's nutrition policy. State NET Programs and the National Food Service Management Institute provide technical assistance and training seminars for school nutrition professionals (see Appendix A). Administrative support is also critical to implementing a new program (214). Training for school administrators can help gain their support for nutrition education. Health promotion services for all school staff can positively affect their eating behaviors and their effectiveness in teaching healthy eating behaviors (180,215,216).

Recommendation 6. Family and community involvement: Involve family members and the community in supporting and reinforcing nutrition education.

The attitudes and behaviors of parents and caretakers directly influence children's and adolescents' choice of foods (217,218). Parents control most of the food choices available at home, so changing parents' eating behaviors may be one of the most effective ways to change their children's eating behaviors. Involving parents in a nutrition education curriculum at the elementary school level can enhance the eating behaviors of both the students (181,219–221) and the parents (181,219,222). Although parental involvement can enhance the effects of nutrition education programs at the elementary school level, it is not known whether involving parents at the secondary school level helps improve the students' eating behaviors. For older youths, self-assessment (185,189,198) and peer educators (187) might be more influential than parental involvement (137).

Parents are usually more receptive to activities that can be done at home than to those that require their attendance at the school (223,224). To involve parents and other family members in nutrition education, schools can

- send nutrition education materials and cafeteria menus home with students,
- ask parents to send healthy snacks to school,
- invite parents and other family members to periodically eat with their children in the cafeteria,
- invite families to attend exhibitions of student nutrition projects or health fairs (217),
- offer nutrition education workshops and screening services, and
- assign nutrition education homework that students can do with their families (e.g., reading and interpreting food labels, reading nutrition-related newsletters, and preparing healthy recipes).

Through school health advisory councils or through direct contact with community organizations, schools can engage community resources and services to respond to the nutritional needs of students (225,226). Schools can also participate in community-based nutrition education campaigns sponsored by public health agencies or voluntary organizations. Students are most likely to adopt healthy eating behaviors if they receive consistent messages through multiple channels (e.g., home, school, community, and the media) and from multiple sources (e.g., parents, peers, teachers, health professionals, and the media) (225).

Recommendation 7. Program evaluation: Regularly evaluate the effectiveness of the school health program in promoting healthy eating, and change the program as appropriate to increase its effectiveness.

Policymakers should regularly review the effectiveness of the school nutrition program. All groups affected by the program should have the opportunity to provide input. Assessment of nutrition programs and policies should include whether

- a comprehensive school nutrition policy exists and is implemented as written;
- nutrition education is provided throughout the preschool, primary, and secondary school years as part of comprehensive school health education;
- teachers deliver nutrition education through developmentally appropriate, culturally relevant, fun, participatory activities that involve social learning strategies;
- teachers and school food service personnel have undertaken joint project planning and action;
- teachers have received curriculum-specific training; and
- families and community organizations are involved in nutrition education.

Schools might also consider measuring the effects of their programs and policies on self-reported eating behaviors; key variables that influence behavior, such as knowledge, attitudes, self-confidence, and behavioral intentions; and in-school eating behaviors that are easy to assess, such as participation in school food service programs and the number of students choosing healthy alternatives in the cafeteria (e.g., salad bars or low-fat milk).

Schools can consult with the state NET Program or with evaluation specialists at universities, school districts, or the state departments of education or health to identify methods and materials for evaluating the effectiveness of their program (227,228). Valid evaluations can increase parent and community support for school programs, help schools reward teachers for exceptional work, and support grant applications for enhancing school health programs.

CONCLUSION

To ensure a healthy future for our children, school-based nutrition education programs must become a national priority. These programs should be part of comprehensive school health programs and reach students from preschool through secondary school. School leaders, community leaders, and parents must commit to implementing and sustaining nutrition education programs within the schools. Such support is crucial to promoting healthy eating behaviors.

The seven recommendations for school-based nutrition education presented in this report provide the framework for establishing such programs. By adopting these recommendations, schools can help ensure that all school-age youths attain their full educational potential and good health.

References

1. Public Health Service. Healthy people 2000: national health promotion and disease prevention objectives. Full report, with commentary. Washington, DC: US Department of Health and Human Services, Public Health Service, 1991. DHHS publication no. (PHS) 91-50212.
2. Mandell RJ, ed. The strategic plan for nutrition education: promoting healthy eating habits for our children. Washington, DC: US Department of Agriculture, Food and Nutrition Service, Nutrition and Technical Services Division, 1993.
3. National Education Goals Panel. The national educational goals report: building a nation of learners. Washington, DC: US Department of Education, National Education Goals Panel, 1992.
4. CDC. Guidelines for effective school health education to prevent the spread of AIDS. *MMWR* 1988;37(S-2):1-14.
5. CDC. Guidelines for school health programs to prevent tobacco use and addiction. *MMWR* 1994;43(RR-2):1-18.
6. CDC. Guidelines for school and community health programs to promote physical activity among youth. *MMWR* (in press).
7. Allensworth DD, Kolbe LJ. The comprehensive school health program: exploring an expanded concept. *J Sch Health* 1987;57(10):409-12.
8. American Cancer Society. Changing the course: a manual for school foodservice providers. Atlanta: American Cancer Society, 1990.
9. American Heart Association. Lower fat and cholesterol easily and economically. . .with Hearty School Lunch. Dallas, TX: American Heart Association, 1992.
10. Ellison RC, Capper AL, Goldberg RJ, Witschi JC, Stare FJ. The environmental component: changing school food service to promote cardiovascular health. *Health Educ Q* 1989;16(2):285-97.
11. Gregoire MB, Sneed J. Standards for nutrition integrity. *Sch Food Serv Res Rev* 1994;18(2):106-11.
12. Nicklas TA, Forcier JE, Farris RP, Hunter SM, Webber LS, Berenson GS. Heart Smart School Lunch Program: a vehicle for cardiovascular health promotion. *Am J Health Promotion* 1989;4(2):91-100.
13. Nicklas TA, Stone E, Montgomery D, et al. Meeting the dietary goals for school meals by the year 2000: The CATCH Eat Smart school nutrition program. *J Health Educ* 1994;25(5):299-307.
14. Raizman DJ, Montgomery DH, Osganian SK, et al. CATCH: food service program process evaluation in a multicenter trial. *Health Educ Q* 1994;(suppl 2):S51-S71.
15. School Food Service Foundation. The healthy E.D.G.E. in schools: eating, dietary guidelines and education. *Sch Food Serv J* 1991;45(3 suppl):1-21.
16. Simons-Morton BG, Parcel GS, Baranowski T, Forthofer R, O'Hara NM. Promoting physical activity and a healthful diet among children: results of a school-based intervention study. *Am J Public Health* 1991;81(8):986-91.
17. Snyder MP, Story M, Lytle Trenkner L. Reducing fat and sodium in school lunch programs: the LUNCHPOWER! Intervention Study. *J Am Diet Assoc* 1992;92:1087-91.
18. US Department of Agriculture. Building for the future: nutrition guidance for the child nutrition programs. Washington, DC: US Department of Agriculture, Food and Nutrition Service, 1992.
19. Whitaker RC, Wright JA, Finch AJ, Psaty BM. An environmental intervention to reduce dietary fat in school lunches. *Pediatrics* 1993;91(6):1107-11.
20. Worthington-Roberts B, Endres J. Position of the American Dietetic Association: nutrition management of adolescent pregnancy. *J Am Diet Assoc* 1989;89(1):104-9.
21. National Agricultural Library. Nutri-Topics: nutrition during pregnancy [resource list]. Beltsville, MD: US Department of Agriculture, National Agricultural Library, Food and Nutrition Information Center, 1994.
22. Alabama State Department of Education. CARE: special nutrition for kids workbook. Montgomery, AL: Alabama State Department of Education, 1995.
23. Blyler EM, Lucas BL. Position of the American Dietetic Association: nutrition in comprehensive program planning for persons with developmental disabilities. *J Am Diet Assoc* 1992;92(5):613-5.
24. Cross B. Nutrition management for children with special needs in child nutrition programs: workshop proceedings. University, MS: National Food Service Management Institute, 1993.

25. Cross B. Annotated bibliography: nutrition management for children with special needs. 2nd ed. University, MS: National Food Service Management Institute, 1993.
26. US Department of Agriculture, the University of Alabama at Birmingham, and the Sparks Clinics. Meeting their needs: training manual for food service personnel caring for children with special needs. Birmingham, AL: University of Alabama at Birmingham, Department of Nutrition Sciences, and the Sparks Clinics, 1994.
27. Rokusek C, Heinrichs E. Nutrition and feeding for persons with special needs: a practical guide and resource manual. 2nd ed. Pierre, SD: South Dakota Department of Education and Cultural Affairs, Office of Child and Adult Nutrition, 1992.
28. Wellman N, Sinofsky J, Crawford L, et al. Feeding for the future: exceptional nutrition in the IEP. Tallahassee, FL: Florida Nutrition Education and Training Program, State of Florida, 1995.
29. Joint Committee on National Health Education Standards. National health education standards: achieving health literacy. An investment in the future. Atlanta: American Cancer Society, 1995.
30. Society for Nutrition Education. Joint position of Society for Nutrition Education (SNE), The American Dietetic Association (ADA), and American School Food Service Association (ASFS): school-based nutrition programs and services. *J Nutr Educ* 1995;27(2):58-61.
31. Trumpfheller W, Foerster SB, Palombo R, eds. The national action plan to improve the American diet: a public/private partnership. Washington, DC: Association of State and Territorial Health Officials, 1993.
32. National Research Council. Diet and health: implications for reducing chronic disease risk. Washington, DC: National Academy Press, 1989.
33. Public Health Service. The Surgeon General's report on nutrition and health. Washington, DC: US Department of Health and Human Services, Public Health Service, 1988. DHHS publication no. (PHS) 88-50210.
34. National Heart, Lung, and Blood Institute. Report of the expert panel on blood cholesterol levels in children and adolescents. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, 1991. NIH publication no. 91-2732.
35. US Department of Agriculture and US Department of Health and Human Services. Nutrition and your health: dietary guidelines for Americans. 4th ed. Washington, DC: US Department of Agriculture and US Department of Health and Human Services, 1995.
36. Center on Hunger, Poverty and Nutrition Policy. Statement on the link between nutrition and cognitive development in children. Medford, MA: Tufts University School of Nutrition, 1995.
37. Hinton AW, Heimindinger J, Foerster SB. Position of the American Dietetic Association: domestic hunger and inadequate access to food. *J Am Diet Assoc* 1990;90(10):1437-41.
38. Troccoli KB. Eat to learn, learn to eat: the link between nutrition and learning in children. Washington, DC: National Health/Education Consortium, 1993. (National Health/Education Consortium occasional paper no. 7.)
39. Food Research and Action Center. Community Childhood Hunger Identification Project: a survey of childhood hunger in the United States. Washington, DC: Food Research and Action Center, 1991.
40. Pollitt E, Leibel RL, Greenfield D. Brief fasting, stress, and cognition in children. *Am J Clin Nutr* 1981;34:1526-33.
41. Pollitt E, Lewis NL, Garza C, Shulman RJ. Fasting and cognitive function. *J Psychiatr Res* 1982-83;17(2):169-74.
42. Pollitt E. Does breakfast make a difference in school? *J Am Diet Assoc* 1995;95(10):1134-9.
43. Meyers AF, Sampson AE, Weitzman M, Rogers BL, Kayne H. School Breakfast Program and school performance. *Am J Dis Child* 1989;143:1234-9.
44. Devaney BL, Gordon AR, Burghardt JA. Dietary intakes of students. *Am J Clin Nutr* 1995;61(1 suppl):205S-212S.
45. American School Health Association, Association for the Advancement of Health Education, and the Society for Public Health Education. The National Adolescent Student Health Survey: a report on the health of America's youth. Oakland, CA: Third Party Publishing, 1989.
46. Pollitt E. Iron deficiency and cognitive function. *Annu Rev Nutr* 1993;13:521-37.
47. Dallman PR, Looker AC, Carroll M, Johnson CL. Influence of age on laboratory criteria for the diagnosis of iron deficiency and iron deficiency anemia in infants and children. In: Hallburg

- L, ed. Proceedings of the symposium on iron nutrition in health and disease. London: John Libbey & Co. (in press).
48. CDC. Prevalence of overweight among adolescents—United States, 1988–91. *MMWR* 1994;43(44):818–21.
 49. Nicklas TA, Webber LS, Srinivasan SR, Berenson GS. Secular trends in dietary intakes and cardiovascular risk factors of 10-y-old children: the Bogalusa Heart Study (1973–1988). *Am J Clin Nutr* 1993;57:930–7.
 50. Ross JG, Pate RR, Lohman TG, Christenson GM. Changes in the body composition of children. *J Physical Educ Recreation Dance* 1987;58(9):74–7.
 51. Shear CL, Freedman DS, Burke GL, Harsha DW, Webber LS, Berenson GS. Secular trends of obesity in early life: the Bogalusa Heart Study. *Am J Public Health* 1988;78(1):75–7.
 52. Troiano RP, Flegal KM, Kuczumski RJ, Campbell SM, Johnson CL. Overweight prevalence and trends for children and adolescents: the National Health Examination Surveys, 1963–1991. *Arch Pediatr Adolesc Med* 1995;149:1085–91.
 53. Freedman DS, Burke GL, Harsha DW, et al. Relationship of changes in obesity to serum lipid and lipoprotein changes in childhood and adolescence. *JAMA* 1985;254(4):515–20.
 54. Kikuchi DA, Srinivasan SR, Harsha DW, Webber LS, Sellers TA, Berenson GS. Relation of serum lipoprotein lipids and apolipoproteins to obesity in children: the Bogalusa Heart Study. *Prev Med* 1992;21:177–90.
 55. Laskarzewski P, Morrison JA, Mellies MJ, et al. Relationships of measurements of body mass to plasma lipoproteins in schoolchildren and adults. *Am J Epidemiol* 1980;111(4):395–406.
 56. Resnicow K, Morabia A. The relation between body mass index and plasma total cholesterol in a multiracial sample of US schoolchildren. *Am J Epidemiol* 1990;132(6):1083–90.
 57. Aristimuño GG, Foster TA, Voors AW, Srinivasan SR, Berenson GS. Influence of persistent obesity in children on cardiovascular risk factors: the Bogalusa Heart Study. *Circulation* 1984;69(5):895–904.
 58. Clarke WR, Woolson RF, Lauer RM. Changes in ponderosity and blood pressure in childhood: the Muscatine Study. *Am J Epidemiol* 1986;124(2):195–206.
 59. Shear CL, Freedman DS, Burke GL, Harsha DW, Berenson GS. Body fat patterning and blood pressure in children and young adults: the Bogalusa Heart Study. *Hypertension* 1987;9:236–44.
 60. Dietz WH Jr. Obesity in infants, children, and adolescents in the United States. I. Identification, natural history, and aftereffects. *Nutr Res* 1981;1:117–37.
 61. Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH. Long-term morbidity and mortality of overweight adolescents. *N Engl J Med* 1992;327:1350–5.
 62. Nieto FJ, Szklo M, Comstock GW. Childhood weight and growth rate as predictors of adult mortality. *Am J Epidemiol* 1992;136(2):201–13.
 63. Brownell KD. The psychology and physiology of obesity: implications for screening and treatment. *J Am Diet Assoc* 1984;84(4):406–14.
 64. Wadden TA, Stunkard AJ. Social and psychological consequences of obesity. *Ann Intern Med* 1985;103(6 pt 2):1062–7.
 65. Lifshitz F, Moses N. Nutritional dwarfing: growth, dieting, and fear of obesity. *J Am Coll Nutr* 1988;7(5):367–76.
 66. French SA, Perry CL, Leon GR, Fulkerson JA. Weight concerns, dieting behavior, and smoking initiation among adolescents: a prospective study. *Am J Public Health* 1994;84(11):1818–20.
 67. Camp DE, Klesges RC, Relyea G. The relationship between body weight concerns and adolescent smoking. *Health Psychol* 1993;12(1):24–32.
 68. Maloney MJ, McGuire J, Daniels SR, Specker B. Dieting behavior and eating attitudes in children. *Pediatrics* 1989;84(3):482–9.
 69. Mellin LM. Responding to disordered eating in children and adolescents. *Nutr News* 1988;51(2):5–7.
 70. Beals KA, Manore MM. The prevalence and consequences of subclinical eating disorders in female athletes. *Int J Sport Nutr* 1994;4:175–95.
 71. Collins ME. Promoting healthy body image through the comprehensive school health program. *J Health Educ* 1991;22(5):297–302.
 72. American Psychiatric Association Task Force on DSM-IV. Diagnostic and statistical manual of mental disorders: DSM-IV. Washington, DC: American Psychiatric Association, 1994.

73. Trowbridge F, Collins B. Measuring dietary behaviors among adolescents. *Public Health Rep* 1993;108(suppl 1):37-41.
74. Herzog DB, Copeland PM. Eating disorders. *N Engl J Med* 1985;313(5):295-303.
75. National Institute of Dental Research. Oral health of United States children: the national survey of dental caries in U.S. school children: 1986-87. National and regional findings. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, 1989. NIH publication no. 89-2247.
76. Gift HC, Reisine ST, Larach DC. The social impact of dental problems and visits. *Am J Public Health* 1992;82(12):1663-8.
77. Enos WF Jr, Beyer JC, Holmes RH. Pathogenesis of coronary disease in American soldiers killed in Korea. *JAMA* 1955;158(11):912-4.
78. McNamara JJ, Molot MA, Stremple JF, Cutting RT. Coronary artery disease in combat casualties in Vietnam. *JAMA* 1971;216(7):1185-7.
79. Newman WP III, Freedman DS, Voors AW, et al. Relation of serum lipoprotein levels and systolic blood pressure to early atherosclerosis: the Bogalusa Heart Study. *N Engl J Med* 1986;314:138-44.
80. Strong JP. Coronary atherosclerosis in soldiers: a clue to the natural history of atherosclerosis in the young. *JAMA* 1986;256(20):2863-6.
81. Freedman DS, Newman WP III, Tracy RE, et al. Black-white differences in aortic fatty streaks in adolescence and early adulthood: the Bogalusa Heart Study. *Circulation* 1988;77(4):856-64.
82. Newman WP III, Wattigney W, Berenson GS. Autopsy studies in United States children and adolescents: relationship of risk factors to atherosclerotic lesions. *Ann NY Acad Sci* 1991;623:16-25.
83. Pathobiological Determinants of Atherosclerosis in Youth Research Group. Relationship of atherosclerosis in young men to serum lipoprotein cholesterol concentrations and smoking: a preliminary report from the Pathobiological Determinants of Atherosclerosis in Youth (PDAY) Research Group. *JAMA* 1990;264(23):3018-24.
84. Kelder SH, Perry CL, Klepp K-I, Lytle LL. Longitudinal tracking of adolescent smoking, physical activity, and food choice behaviors. *Am J Public Health* 1994;84(7):1121-6.
85. Berenson GS, Epstein FA. Conference on blood lipids in children: optimal levels for early prevention of coronary artery disease. Workshop report: Epidemiological Section. *Prev Med* 1983;12:741-97.
86. Frerichs RR, Srinivasan SR, Webber LS, Berenson GS. Serum cholesterol and triglyceride levels in 3,446 children from a biracial community: the Bogalusa Heart Study. *Circulation* 1976;54(2):302-9.
87. Kwiterovich PO Jr. Plasma lipid and lipoprotein levels in childhood. *Ann NY Acad Sci* 1991;623:90-107.
88. Lauer RM, Connor WE, Leaverton PE, Reiter MA, Clarke WR. Coronary heart disease risk factors in school children: the Muscatine Study. *J Pediatr* 1975;86(5):697-706.
89. Webber LS, Osganian V, Luepker RV, et al. Cardiovascular risk factors among third grade children in four regions of the United States: the CATCH study. *Am J Epidemiol* 1995;141(5):428-39.
90. Wynder EL. An American Health Foundation monograph: coronary artery disease prevention: cholesterol, a pediatric perspective. *Prev Med* 1989;18:323-409.
91. Ernst ND, Obarzanek E. Child health and nutrition: obesity and high blood cholesterol. *Prev Med* 1994;23:427-36.
92. Lauer RM, Lee J, Clarke WR. Factors affecting the relationship between childhood and adult cholesterol levels: the Muscatine Study. *Pediatrics* 1988;82(3):309-18.
93. Orchard TJ, Donahue RP, Kuller LH, Hodge PN, Drash AL. Cholesterol screening in childhood: does it predict adult hypercholesterolemia? The Beaver County experience. *J Pediatr* 1983;103(5):687-91.
94. Porkka KVK, Viikari JSA, Akerblom HK. Tracking of serum HDL-cholesterol and other lipids in children and adolescents: the Cardiovascular Risk in Young Finns Study. *Prev Med* 1991;20:713-24.
95. Webber LS, Srinivasan SR, Wattigney WA, Berenson GS. Tracking of serum lipids and lipoproteins from childhood to adulthood: the Bogalusa Heart Study. *Am J Epidemiol* 1991;133(9):884-99.

96. Wynder EL. Summary and recommendations of the conference on blood lipids in children: optimal levels for early prevention of coronary artery disease. *Prev Med* 1983;12:728-40.
97. Lauer RM, Clarke WR. Childhood risk factors for high adult blood pressure: the Muscatine Study. *Pediatrics* 1989;84(4):633-41.
98. Shear CL, Burke GL, Freedman DS, Berenson GS. Value of childhood blood pressure measurements and family history in predicting future blood pressure status: results from 8 years of follow-up in the Bogalusa Heart Study. *Pediatrics* 1986;77(6):862-9.
99. Casey VA, Dwyer JT, Coleman KA, Valadian I. Body mass index from childhood to middle age: a 50-y follow-up. *Am J Clin Nutr* 1992;56:14-8.
100. Coates TJ, Thoresen CE. Treating obesity in children and adolescents: a review. *Am J Public Health* 1978;68(2):143-51.
101. Freedman DS, Shear CL, Burke GL, et al. Persistence of juvenile-onset obesity over eight years: the Bogalusa Heart Study. *Am J Public Health* 1987;77(5):588-92.
102. Garn SM, LaVelle M. Two-decade follow-up of fatness in early childhood. *Am J Dis Child* 1985;139:181-5.
103. Guo SS, Roche AF, Chumlea WC, Gardner JD, Siervogel RM. The predictive value of childhood body mass index values for overweight at age 35 y. *Am J Clin Nutr* 1994;59:810-9.
104. McGinnis JM, Foege WH. Actual causes of death in the United States. *JAMA* 1993;270(18):2207-12.
105. National Heart, Lung, and Blood Institute. Report of the expert panel on population strategies for blood cholesterol reduction. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, 1990. NIH publication no. 90-3046.
106. American Cancer Society. Cancer facts & figures—1995. Atlanta: American Cancer Society, 1995.
107. Doll R. The lessons of life: keynote address to the Nutrition and Cancer Conference. *Cancer Res* 1992;52(suppl):2024S-2029S.
108. Doll R, Peto R. The causes of cancer: quantitative estimates of avoidable risks of cancer in the United States today. New York: Oxford University Press, 1981.
109. Butrum RR, Clifford CK, Lanza E. NCI dietary guidelines: rationale. *Am J Clin Nutr* 1988;48:888-95.
110. National Cancer Institute. Eat more fruits & vegetables—5 a day for better health. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, 1991. NIH publication no. 92-3248.
111. American Heart Association. Heart and stroke facts: 1995 statistical supplement. Dallas, TX: American Heart Association, 1994.
112. American Heart Association. Heart and stroke facts. Dallas, TX: American Heart Association, 1994.
113. National Institute of Diabetes and Digestive and Kidney Diseases. Diabetes statistics. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, 1994. Publication no. 94-3822.
114. American Diabetes Association. Medical management of non-insulin-dependent (Type II) diabetes. 3rd ed. Alexandria, VA: American Diabetes Association, 1994.
115. National Heart, Lung, and Blood Institute. Working group report on primary prevention of hypertension. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, 1993. NIH publication no. 93-2669.
116. Dawber TR. The Framingham Study: the epidemiology of atherosclerotic disease. Cambridge, MA: Harvard University Press, 1980.
117. Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure. The fifth report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure (JNC V). *Arch Intern Med* 1993;153:154-83.
118. Kuczmarski RJ, Flegal KM, Campbell SM, Johnson CL. Increasing prevalence of overweight among US adults. *JAMA* 1994;272(3):205-11.
119. NIH Consensus Development Panel on Optimal Calcium Intake. Optimal calcium intake. *JAMA* 1994;272(24):1942-8.
120. Matkovic V. Calcium and peak bone mass. *J Intern Med* 1992;231:151-60.
121. Sandler RB, Slemenda CW, LaPorte RE, et al. Postmenopausal bone density and milk consumption in childhood and adolescence. *Am J Clin Nutr* 1985;42:270-4.

122. Sentipal JM, Wardlaw GM, Mahan J, Matkovic V. Influence of calcium intake and growth indexes on vertebral bone mineral density in young females. *Am J Clin Nutr* 1991;54:425-8.
123. Weinhouse S, Bal DG, Adamson R, et al. American Cancer Society guidelines on diet, nutrition, and cancer. *CA Cancer J Clin* 1991;41(6):334-8.
124. Chait A, Brunzell JD, Denke MA, et al. Rationale of the diet-heart statement of the American Heart Association: report of the Nutrition Committee. *Circulation* 1993;88(6):3008-29.
125. Human Nutrition Information Service. The food guide pyramid. Hyattsville, MD: US Department of Agriculture, Human Nutrition Information Service, 1992. (Home and Garden bulletin no. 252.)
126. Alaimo K, McDowell MA, Briefel RR, et al. Dietary intake of vitamins, minerals, and fiber of persons ages 2 months and over in the United States: third National Health and Nutrition Examination Survey, Phase 1, 1988-91. Hyattsville, MD: US Department of Health and Human Services, Public Health Service, CDC, 1994. (Advance data from vital and health statistics; no. 258.) DHHS publication no. (PHS) 95-1250.
127. Kann L, Warren CW, Harris WA, et al. Youth risk behavior surveillance—United States, 1993. *MMWR* 1995;44(SS-1):1-56.
128. McDowell MA, Briefel RR, Alaimo K, et al. Energy and macronutrient intakes of persons ages 2 months and over in the United States: third National Health and Nutrition Examination Survey, Phase 1, 1988-91. Hyattsville, MD: US Department of Health and Human Services, Public Health Service, CDC, 1994. (Advance data from vital and health statistics; no. 255.) DHHS publication no. (PHS) 95-1250.
129. Tippet KS, Mickle SJ, Goldman JD, et al. Food and nutrient intakes by individuals in the United States, 1 day, 1989-91. Beltsville, MD: US Department of Agriculture, Agricultural Research Service, Beltsville Human Nutrition Research Center, 1995. Nationwide Food Surveys report no. 91-2.
130. Lewis CJ, Crane NT, Moore BJ, Hubbard VS. Healthy People 2000: report on the 1994 nutrition progress review. *Nutr Today* 1994;29(6):6-14.
131. Krebs-Smith SM, Cook DA, Subar AF, Cleveland L, Friday J, Kahle LL. Fruit and vegetable intakes of children and adolescents in the United States. *Arch Pediatr Adolesc Med* 1996;150:81-6.
132. Murphy AS, Youatt JP, Hoerr SL, Sawyer CA, Andrews SL. Nutrition education needs and learning preferences of Michigan students in grades 5, 8, and 11. *J Sch Health* 1994;64(7):273-8.
133. Resnicow K, Reinhardt J. What do children know about fat, fiber, and cholesterol? A survey of 5,116 primary and secondary school students. *J Nutr Educ* 1991;23(2):65-71.
134. Story M, Resnick MD. Adolescents' views on food and nutrition. *J Nutr Educ* 1986;18(4):188-92.
135. Dwyer J. The School Nutrition Dietary Assessment Study. *Am J Clin Nutr* 1995;61(1 suppl):173S-177S.
136. Contento IR, Manning AD, Shannon B. Research perspective on school-based nutrition education. *J Nutr Educ* 1992;24:247-60.
137. Contento I, Balch GI, Bronner YL, et al. Nutrition education for school-aged children. *J Nutr Educ* 1995;27(6):298-311.
138. Lytle L, Achterberg C. Changing the diet of America's children: what works and why? *J Nutr Educ* 1995;27(5):250-60.
139. Crockett SJ, Sims LS. Environmental influences on children's eating. *J Nutr Educ* 1995;27(5):235-49.
140. Kotz K, Story M. Food advertisements during children's Saturday morning television programming: are they consistent with dietary recommendations? *J Am Diet Assoc* 1994;94:1296-1300.
141. Cotugna N. TV ads on Saturday morning children's programming—what's new? *J Nutr Educ* 1988;20(3):125-7.
142. Taras HL, Gage M. Advertised foods on children's television. *Arch Pediatr Adolesc Med* 1995;149:649-52.
143. Signorielli N, Lears M. Television and children's conceptions of nutrition: unhealthy messages. *Health Commun* 1992;4(4):245-57.
144. Taras HL, Sallis JF, Patterson TL, Nader PR, Nelson JA. Television's influence on children's diet and physical activity. *J Dev Behav Pediatr* 1989;10(4):176-80.

145. Robinson TN, Killen JD. Ethnic and gender differences in the relationships between television viewing and obesity, physical activity, and dietary fat intake. *J Health Educ* 1995;26(2 suppl):S91-S98.
146. Pate RR, Ross JG. Factors associated with health-related fitness. *J Physical Educ Recreation Dance* 1987;58(9):93-5.
147. Dietz WH Jr, Gortmaker SL. Do we fatten our children at the television set? Obesity and television viewing in children and adolescents. *Pediatrics* 1985;75(5):807-12.
148. Obarzanek E, Schreiber GB, Crawford PB, et al. Energy intake and physical activity in relation to indexes of body fat: the National Heart, Lung, and Blood Institute Growth and Health Study. *Am J Clin Nutr* 1994;60:15-22.
149. Shannon B, Peacock J, Brown MJ. Body fatness, television viewing and calorie-intake of a sample of Pennsylvania sixth grade children. *J Nutr Educ* 1991;23(6):262-8.
150. Nielsen Media Research. Nielsen report on television. New York: Nielsen Media Research, 1990.
151. Team Nutrition. A healthier tomorrow. Alexandria, VA: Team Nutrition, 1995.
152. Resnicow K. Cancer prevention and comprehensive school health education: the role of the American Cancer Society. Paper presented at the American Cancer Society National Conference, Atlanta, GA, May 28-30, 1991.
153. Allensworth D, Wyche J, Lawson E, Nicholson L, eds. Defining a comprehensive school health program: an interim statement. Washington, DC: National Academy Press, 1995.
154. US Department of Agriculture, Food and Consumer Service. 7 CFR Parts 210 and 220. National School Lunch Program and School Breakfast Program: school meal initiatives for healthy children. *Federal Register* 1995;60(113):31188-222.
155. US Department of Agriculture. Needs assessment guide for the Nutrition Education and Training Program. Washington, DC: US Department of Agriculture, Food and Nutrition Service, 1994.
156. US Department of Health and Human Services, Food and Drug Administration. 21 CFR Part 101. Food labeling: nutrient content claims, definition of term: healthy. *Federal Register* 1994;59(89):24232-50.
157. US Department of Health and Human Services. The new food label. Washington, DC: US Department of Health and Human Services, Food and Drug Administration, 1995. Publication no. BG 95-14.
158. Caldwell DR, Pilant VB. Position of the American Dietetic Association: competitive foods in schools. *J Am Diet Assoc* 1991;91(9):1123-5.
159. Wolfe WS, Campbell CC. Nutritional health of school-aged children in upstate New York: what are the problems and what can schools do? Ithaca, NY: Cornell University, Division of Nutritional Sciences, 1991.
160. West Virginia Department of Education. Let's party: party ideas for school and home. 2nd ed. Charleston, WV: West Virginia Department of Education, 1993.
161. Snyder P, Lytle L, Pellegrino T, Anderson M, Selk J. Commentary on school meals from school food service personnel and researchers. *Am J Clin Nutr* 1995;61(suppl):247S-249S.
162. American School Food Service Association. Creating policy for nutrition integrity in schools. Alexandria, VA: American School Food Service Association, 1994.
163. McConnell PE, Shaw JB. Position of the American Dietetic Association: child nutrition services. *J Am Diet Assoc* 1993;93(3):334-6.
164. Steinschneider J, Coyne AH. Bringing WIC to school. Washington, DC: Center on Budget and Policy Priorities, 1995.
165. Lytle LA, Kelder SH, Perry CL, Klepp K-I. Covariance of adolescent health behaviors: the Class of 1989 Study. *Health Educ Res* 1995;10(2):133-46.
166. Devine CM, Olson CM, Frongillo EA Jr. Impact of the Nutrition for Life program on junior high students in New York State. *J Sch Health* 1992;62(8):381-5.
167. Resnicow K, Cohn L, Reinhardt J, et al. A three-year evaluation of the Know Your Body Program in inner-city schoolchildren. *Health Educ Q* 1992;19(4):463-80.
168. Weiss EH, Kien CL. A synthesis of research on nutrition education at the elementary school level. *J Sch Health* 1987;57(1):8-13.
169. Contento IR, Kell DG, Keiley MK, Corcoran RD. A formative evaluation of the American Cancer Society Changing the Course nutrition education curriculum. *J Sch Health* 1992;62(9):411-6.

170. Bagby R, Campbell VS, Achterberg C. Every day, lots of ways: an interdisciplinary nutrition curriculum for kindergarten–sixth grade. Harrisburg, PA: Pennsylvania State Department of Education, 1993.
171. Zeller PK, Jacobson MF. Eat, think, and be healthy! Creative nutrition activities for children. Washington, DC: Center for Science in the Public Interest, 1987.
172. Kalina BB, Philipps CA, Minns HV. The NET Program: a ten-year perspective. *J Nutr Educ* 1989;21(1):38–42.
173. Byrd-Bredbenner C, O'Connell LH, Shannon B. Junior high home economics curriculum: its effect on students' knowledge, attitude, and behavior. *Home Econ Res J* 1982;11(2):123–33.
174. Byrd-Bredbenner C, O'Connell LH, Shannon B, Eddy JM. A nutrition curriculum for health education: its effect on students' knowledge, attitude, and behavior. *J Sch Health* 1984;54(10):385–8.
175. Byrd-Bredbenner C, Shannon B, Hsu L, Smith DH. A nutrition education curriculum for senior high home economics students: its effect on students' knowledge, attitudes, and behaviors. *J Nutr Educ* 1988;20(6):341–6.
176. German MJ, Pearce J, Wyse BW, Hansen RG. A nutrition component for high school health education curriculums. *J Sch Health* 1981;51(3):149–53.
177. Lewis M, Brun J, Talmage H, Rasher S. Teenagers and food choices: the impact of nutrition education. *J Nutr Educ* 1988;20(6):336–40.
178. Shannon B, Chen AN. A three-year school-based nutrition education study. *J Nutr Educ* 1988;20(3):114–24.
179. Havas S, Heimendinger J, Damron D, et al. 5 a day for better health—nine community research projects to increase fruit and vegetable consumption. *Public Health Rep* 1995;110(1):68–79.
180. DeFriese GH, Crossland CL, MacPhail-Wilcox B, Sowers JG. Implementing comprehensive school health programs: prospects for change in American schools. *J Sch Health* 1990;60(4):182–7.
181. Coates TJ, Jeffery RW, Slinkard LA. Heart healthy eating and exercise: introducing and maintaining changes in health behaviors. *Am J Public Health* 1981;71(1):15–23.
182. Coates TJ, Barofsky I, Saylor KE, et al. Modifying the snack food consumption patterns of inner-city high school students: the Great Sensations Study. *Prev Med* 1985;14:234–47.
183. Howison D, Niedermeyer F, Shortridge R. Field testing a fifth-grade nutrition education program designed to change food-selection behavior. *J Nutr Educ* 1988;20(2):82–6.
184. Killen JD, Telch MJ, Robinson TN, Maccoby N, Taylor CB, Farquhar JW. Cardiovascular disease risk reduction for tenth graders: a multiple-factor school-based approach. *JAMA* 1988;260(12):1728–33.
185. King AC, Saylor KE, Foster S, et al. Promoting dietary change in adolescents: a school-based approach for modifying and maintaining healthful behavior. *Am J Prev Med* 1988;4(2):68–74.
186. Perry CL, Mullis RM, Maile MC. Modifying the eating behavior of young children. *J Sch Health* 1985;55(10):399–402.
187. Perry CL, Klepp K-I, Halper A, et al. Promoting healthy eating and physical activity patterns among adolescents: a pilot study of "Slice of Life." *Health Educ Res* 1987;2(2):93–103.
188. Walter HJ. Primary prevention of chronic disease among children: the school-based "Know Your Body" intervention trials. *Health Educ Q* 1989;16(2):201–14.
189. White AA, Skinner JD. Can goal setting as a component of nutrition education effect behavior change among adolescents? *J Nutr Educ* 1988;20(6):327–35.
190. Luepker RV, Perry CL, McKinlay SM, et al. Outcomes of a field trial to improve children's dietary patterns and physical activity: the Child and Adolescent Trial for Cardiovascular Health (CATCH). *JAMA* 1996;275:768–76.
191. Bush PJ, Zuckerman AE, Taggart VS, Theiss PK, Peleg EO, Smith SA. Cardiovascular risk factor prevention in black school children: the "Know Your Body" Evaluation Project. *Health Educ Q* 1989;16(2):215–27.
192. Contento I. Children's thinking about food and eating—a Piagetian-based study. *J Nutr Educ* 1981;13(1 suppl):S86–S90.
193. Lieberman LD, Clark NM, Krone KV, Orlandi MA, Wynder EL. The relationship between cognitive maturity and information about health problems among school age children. *Health Educ Res* 1992;7(3):391–401.

194. Lytle Trenkner L, Kelder SH. Nutrition education and school food service intervention as components of comprehensive school health education. Report to the American Cancer Society's Advisory Committee on Technology Transfer of Behavioral Research, November 11, 1991.
195. Killen JD, Robinson TN. School-based research on health behavior change: the Stanford Adolescent Heart Health Program as a model for cardiovascular disease risk reduction. In: Rothkopf EZ, ed. Review of research in education. Vol. 15. Washington, DC: American Educational Research Association, 1988:171-200.
196. Jaycox S, Baranowski T, Nader PR, Dworkin R, Vanderpool NA. Theory-based health education activities for third to sixth grade children. *J Sch Health* 1983;53(10):584-8.
197. Rickard KA, Gallahue DL, Gruen GE, Tridle M, Bewley N, Steele K. The play approach to learning in the context of family and schools: an alternative paradigm for nutrition and fitness education in the 21st century. *J Am Diet Assoc* 1995;95:1121-6.
198. Burnett KF, Magel PE, Harrington S, Taylor CB. Computer-assisted behavioral health counseling for high school students. *J Counseling Psychol* 1989;36(1):63-7.
199. National Agricultural Library. Database of food and nutrition software and multimedia programs. Beltsville, MD: US Department of Agriculture, National Agricultural Library, Food and Nutrition Information Center, 1995.
200. Kolasa KM, Miller MG. New developments in nutrition education using computer technology. *J Nutr Educ* 1996;28:7-14.
201. Bandura A. Social learning theory. Englewood Cliffs, NJ: Prentice-Hall, 1977.
202. Bandura A. Social foundations of thought and action: a social cognitive theory. Englewood Cliffs, NJ: Prentice-Hall, 1986.
203. Baranowski T. Reciprocal determinism at the stages of behavior change: an integration of community, personal and behavioral perspectives. *Int Q Community Health Educ* 1989-90;10(4):297-327.
204. Perry CL, Baranowski T, Parcel GS. How individuals, environments, and health behavior interact: social learning theory. In: Glanz K, Lewis FM, Rimer BK, eds. Health behavior and health education: theory, research, and practice. San Francisco: Jossey-Bass, 1990:161-86.
205. Perry CL, Jessor R. The concept of health promotion and the prevention of adolescent drug abuse. *Health Educ Q* 1985;12(2):169-84.
206. Smith SF, James MA. School lunch as a nutrition education resource for fourth graders. *J Nutr Educ* 1980;12(2):46-9.
207. Connell DB, Turner RR, Mason EF. Summary of findings of the School Health Education Evaluation: health promotion effectiveness, implementation, and costs. *J Sch Health* 1985;55(8):316-21.
208. Olson CM, Devine CM, Frongillo EA Jr. Dissemination and use of a school-based nutrition education program for secondary school students. *J Sch Health* 1993;63(8):343-8.
209. Ross JG, Luepker RV, Nelson GD, Saavedra P, Hubbard BM. Teenage Health Teaching Modules: impact of teacher training on implementation and student outcomes. *J Sch Health* 1991;61(1):31-4.
210. Rye JA, Hunt BN, Nicely R Jr, Shannon B. The development of a nutrition inservice course for teachers of young children. *J Nutr Educ* 1982;14(3):93-6.
211. Gingiss PL. Enhancing program implementation and maintenance through a multiphase approach to peer-based staff development. *J Sch Health* 1992;62(5):161-6.
212. Contento I, Balch GI, Bronner YL, et al. Inservice preparation in nutrition education for professionals and paraprofessionals. *J Nutr Educ* 1995;27(6):347-54.
213. Downey AM, Virgilio SJ, Serpas DC, Nicklas TA, Arbeit ML, Berenson GS. "Heart Smart"—a staff development model for a school-based cardiovascular health intervention. *Health Educ* 1988;19(5):64-71.
214. Smith DW, McCormick LK, Steckler AB, McLeroy KR. Teachers' use of health curricula: implementation of Growing Healthy, Project SMART, and the Teenage Health Teaching Modules. *J Sch Health* 1993;63(8):349-54.
215. Baranowski T, Hearn MD, Baranowski JC, et al. Teach Well: the relation of teacher wellness to elementary student health and behavior outcomes: baseline subgroup comparisons. *J Health Educ* 1995;26(suppl 2):S61-S71.

216. Johnson CC, Powers CR, Bao W, Harsha DW, Berenson GS. Cardiovascular risk factors of elementary school teachers in a low socio-economic area of a metropolitan city: the Heart Smart Program. *Health Educ Res* 1994;9(2):183-91.
217. Johnson CC, Osganian SK, Budman SB, et al. CATCH: family process evaluation in a multicenter trial. *Health Educ Q* 1994;(suppl 2):S91-S106.
218. Crockett SJ, Mullis RM, Perry CL. Parent nutrition education: a conceptual model. *J Sch Health* 1988;58(2):53-7.
219. Hearn MD, Bigelow C, Nader PR, et al. Involving families in cardiovascular health promotion: the CATCH Feasibility Study. *J Health Educ* 1992;23(1):22-31.
220. Kirks BA, Hendricks DG, Wyse BW. Parent involvement in nutrition education for primary grade students. *J Nutr Educ* 1982;14(4):137-40.
221. Perry CL, Luepker RV, Murray DM, et al. Parent involvement with children's health promotion: the Minnesota Home Team. *Am J Public Health* 1988;78(9):1156-60.
222. Crockett SJ, Mullis R, Perry CL, Luepker RV. Parent education in youth-directed nutrition interventions. *Prev Med* 1989;18:475-91.
223. Crockett SJ, Perry CL, Pirie P. Nutrition intervention strategies preferred by parents: results of a marketing survey. *J Nutr Educ* 1989;21(2):90-4.
224. Perry CL, Crockett SJ, Pirie P. Influencing parental health behavior: implications of community assessments. *Health Educ* 1987;18(5):68-77.
225. Killip DC, Lovick SR, Goldman L, Allensworth DD. Integrated school and community programs. *J Sch Health* 1987;57(10):437-44.
226. Kirby D. Comprehensive school health and the larger community: issues and a possible scenario. *J Sch Health* 1990;60(4):170-7.
227. US Department of Agriculture. Evaluation guide for the Nutrition Education and Training Program. Washington, DC: US Department of Agriculture, Food and Nutrition Service, 1995.
228. Stone EJ, McGraw SA, Osganian SK, Elder JP, eds. Process evaluation in the multicenter Child and Adolescent Trial for Cardiovascular Health (CATCH). *Health Educ Q* 1994;(suppl 2):S1-S142.
229. CDC. School Health Policies and Programs Study (SHPPS): a summary report. *J Sch Health* 1995;65(8):281-353.

APPENDIX A: NUTRITION EDUCATION RESOURCE LIST

Nutrition education curricula and print, audiovisual, and computer-based materials are available from government agencies, voluntary organizations, corporations, and commodity organizations. State Nutrition Education and Training Program coordinators can help schools identify the most appropriate nutrition education curricula and materials. National clearinghouses that can help schools identify a wide range of nutrition education and school food service resources are sponsored by the U.S. Department of Agriculture's Food and Nutrition Information Center and the National Food Service Management Institute; the former also serves as a lending library.

Food and Nutrition Information Center
National Agricultural Library
U.S. Department of Agriculture
10301 Baltimore Blvd., Room 304
Beltsville, MD 20705
301-504-5719

National Food Service Management
Institute
P.O. Box 188
University of Mississippi
University, MS 38677
800-321-3054

At the local and state levels, educational materials or curricula may be available from affiliates of voluntary health promotion organizations (e.g., the American Cancer Society or the American Heart Association), commodity organizations or national boards for specific food industries, county cooperative extension services, local and state health departments, school districts, state education agencies, and universities. At the national level, nutrition education materials can also be obtained from the following voluntary organizations and federal government agencies:

American Cancer Society
1599 Clifton Road, NE
Atlanta, GA 30328
800-ACS-2345 (800-227-2345)

American Dietetic Association
National Center for Nutrition
and Dietetics
216 W. Jackson Blvd., Suite 800
Chicago, IL 60606-6995
800-745-0775 ext. 5000

American Heart Association
7272 Greenville Ave.
Dallas, TX 75231-4596
800-AHA-USA1 (800-242-8721)

American School Food Service
Association
1600 Duke St., 7th Floor
Alexandria, VA 22314
800-877-8822 ext. 116

Consumer Information Center
Pueblo, CO 81009
719-948-4000 (call for catalog)

International Food Information Council
1100 Connecticut Ave., NW, Suite 430
Washington, DC 20036
202-296-6540

National Cancer Institute
Office of Cancer Communications
Building 31, Room 10A16
31 Center Drive MSC-2580
Bethesda, MD 20892-2580
800-4-CANCER (800-422-6237)

National Heart, Lung, and Blood
Institute Information Center
P.O. Box 30105
Bethesda, MD 20824-0105
301-251-1222

Team Nutrition
U.S. Department of Agriculture
3101 Park Center Drive, Room 802
Alexandria, VA 22302
703-305-1624

APPENDIX B: YOUTH RISK BEHAVIOR SURVEILLANCE SYSTEM AND SCHOOL HEALTH POLICIES AND PROGRAMS STUDY

In 1990, CDC established the Youth Risk Behavior Surveillance System to help monitor progress in attaining national health and education objectives by periodically measuring the prevalence of behaviors in six health risk categories. These behaviors, which are usually established during youth, contribute to the leading causes of death and disease in the United States. Dietary behaviors are one of the six health risk categories. CDC conducts the Youth Risk Behavior Survey (YRBS) biennially in a national probability sample of high school students and enables interested state and local education agencies to conduct the survey in comparable probability samples in states and cities (127). The specific dietary behaviors and attitudes monitored by the YRBS include consumption of fruits and vegetables, consumption of foods high in fat, perceptions of body weight, and attempted weight loss and weight-loss techniques used. The YRBS also obtains information about specific physical activity behaviors.

In 1994, CDC conducted the School Health Policies and Programs Study (SHPPS), which is a national study of school policies and programs at the school, district, and state levels that support comprehensive school health programs. The study also provides baseline data on national health and education objectives that can be attained through school health and physical education, school food service, and school health services and policies (229).

SHPPS included a mail survey of local and state education agencies' policies related to school health in grades kindergarten through 12. The survey was conducted in all states and in a nationally representative sample of districts. The study also included on-site, structured interviews with school principals, health education teachers, physical education teachers, school food service directors, school nurses, counselors, and other personnel in a nationally representative sample of middle schools and high schools. The questionnaire included the following: school nutrition education requirements for students; the content of nutrition education curricula; training and joint activities of food service staff and teachers responsible for nutrition education; school policies related to foods sold in vending machines and for fundraising; food service practices related to purchasing and preparing food; involvement of parents, staff, and students in planning food service meals; and involvement of fast-food or food service management companies in school meals.

Single copies of YRBS and SHPPS reports are available from CDC's Division of Adolescent and School Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Mailstop K-33, 4770 Buford Highway, NE, Atlanta, GA 30341-3724; telephone: (770) 488-5330.

APPENDIX C: SELECTED SCHOOL-BASED STRATEGIES TO PROMOTE HEALTHY EATING

Different, developmentally appropriate activities are listed for lower elementary school, upper elementary school, and middle and high school students (194). This list is not intended to be comprehensive. However, it does include many of the concepts critical to improving the diet and health of young persons in this country. Schools should review these educational activities in relation to their students' needs and abilities to determine which activities are appropriate at each grade level.

Interventions that promote healthy changes in eating behaviors need to target three interacting spheres of influence: (a) the environment, which influences the likelihood that healthy eating behaviors will be adopted through social norms, influential role models, cues to action, reinforcements, and opportunities for action; (b) personal characteristics (e.g., knowledge, attitudes, beliefs, values, confidence in one's ability to change eating behaviors, and expectations about the consequences of making those changes); and (c) behavioral skills and experience, which are related to selecting or preparing specific foods, dietary self-assessment, and decision-making (186, 194, 203, 204).

The strategies listed here require the involvement of teachers, administrators, food service personnel, other school staff, and parents (194). Classroom teachers play the lead role in most of these activities, but many activities would be most effective if they were reinforced by other persons; all adults in the school community can help by serving as role models. Each school or district should determine the policies it needs to guide its nutrition-related activities and who is responsible for the tasks.

For lower elementary students

Strategies to make the food environment more health-enhancing

- Make healthy foods (e.g., fruits, vegetables, and whole grains) widely available at school, and discourage the availability of foods high in fat, sodium, and added sugars.
- Involve parents in nutrition education through homework.
- Provide role models (e.g., teachers, parents, other adults, older children, and celebrities or fictional characters) for healthy eating.
- Provide cues, through posters and marketing-style incentives, that encourage students to make healthy choices about eating and physical activity.
- Use incentives, such as verbal praise or token gifts, to reinforce healthy eating and physical activity. Do not use food for reward or punishment of any behavior.

Strategies to enhance personal characteristics that will support healthy eating

- Make basic connections between food and health (e.g., "You need food to feel good and to grow").

- Teach the importance of balancing food intake and physical activity.
- Identify healthy snacks (e.g., fruits, vegetables, and low-fat milk).
- Increase students' confidence in their ability to make healthy eating choices by gradually building up their food selection and preparation skills and giving them practice.

Strategies to enhance behavioral capabilities that will support healthy eating

- Provide many healthy foods for students to taste in an enjoyable social context.
- Let students prepare simple snacks.
- Have students try unfamiliar and culturally diverse foods that are low in fat, sodium, and added sugars.

For upper elementary students

Strategies to make the food environment more health-enhancing

- Make healthy foods (e.g., fruits, vegetables, and whole grains) widely available at school, and discourage the availability of foods high in fat, sodium, and added sugars.
- Involve parents in nutrition education through homework.
- Provide role models (e.g., teachers, parents, other adults, adolescents, and celebrities or fictional characters) for healthy eating.
- Through class discussions and small-group exercises, provide social support for making healthy changes in eating and physical activity.
- Provide cues, through posters and marketing-style incentives that students design, that encourage students to make healthy choices about eating and physical activity.
- Use incentives, such as verbal praise or token gifts, to reinforce healthy eating and physical activity. Do not use food as a reward or punishment of any behavior.

Strategies to enhance personal characteristics that will support healthy eating

- Explain the effects that diet and physical activity have on future health as well as on immediate concerns (e.g., current health, physical appearance, obesity, sense of well-being, and capacity for physical activity).
- Teach the principles of the Dietary Guidelines for Americans and the Food Guide Pyramid. Instill pride in choosing to eat meals and snacks that comply with these principles.

- Help students identify foods high and low in fat, saturated fat, cholesterol, sodium, added sugars, and fiber.
- Teach the importance of balancing food intake and physical activity.
- Teach the importance of eating adequate amounts of fruits, vegetables, and whole grains.
- Help students increase the value they place on health and their sense of control over food selection and preparation.
- Increase students' confidence in their ability to make healthy eating choices by gradually building up their food selection and preparation skills and giving them practice.
- Have students analyze food preferences and factors that trigger eating behaviors.

Strategies to enhance behavioral capabilities that will support healthy eating

- Provide opportunities for students to taste many healthy foods in an enjoyable social context.
- Let students prepare healthy snacks or simple meals.
- Encourage students to try unfamiliar and culturally diverse foods that are low in fat, sodium, and added sugars and that are high in fiber.
- Have students select healthy foods from a fast-food restaurant menu.
- Teach students how to recognize the fat, sodium, and fiber contents of foods by reading nutrition labels.
- Help students record and assess their food intake.
- Teach students how to use the Food Guide Pyramid to assess their diet for variety, moderation, and proportionality.
- Have students set simple goals for changes in eating and physical activity, and devise strategies for implementing these changes and monitoring progress in reaching their goals.
- When appropriate, let students practice (through role plays) encouraging parents to make healthy choices about eating and physical activity at home.
- Have students examine media and social influences on eating and physical activity; teach students how to respond to these pressures.

For middle and high school students

Strategies to make the food environment more health-enhancing

- Make healthy foods (e.g., fruits, vegetables, and whole grains) widely available at school, and discourage the availability of foods high in fat, sodium, and added sugars.
- Provide role models (e.g., teachers, parents, other adults, and celebrities) for healthy eating.
- Use peers as role models, and use peer-led nutrition education activities.
- Through class discussions and small-group exercises, provide social support for making healthy changes in eating and physical activity.
- Provide cues, through posters and marketing-style incentives that students design, that encourage students to make healthy choices about eating and physical activity.

Strategies to enhance personal characteristics that will support healthy eating

- Explain the effects that diet and physical activity have on future health as well as on immediate concerns (e.g., current health, physical appearance, obesity, eating disorders, sense of well-being, and capacity for physical activity).
- Have students identify reasons to adopt healthy eating and physical activity patterns.
- Teach the principles of the Dietary Guidelines for Americans. Instill in the students pride in choosing to eat meals and snacks that comply with these principles.
- Teach students how to identify foods high and low in fat, saturated fat, cholesterol, sodium, and added sugars.
- Teach students how to identify foods that are excellent sources of fiber, complex carbohydrates, calcium, iron, vitamin A, vitamin C, and folate.
- Teach the importance of balancing food intake and physical activity.
- Teach the effects of unsafe weight-loss methods and the characteristics of a safe weight-loss program.
- Help students increase the value they place on health and their sense of control over food selection and preparation.
- Increase students' confidence in their ability to eat healthily by gradually building up their skills and giving them practice.
- Help students examine what motivates persons to adopt particular eating habits. Have students keep a food diary noting what cues their own eating behavior (e.g., mood, hunger, stress, or other persons).

Strategies to enhance behavioral capabilities that will support healthy eating

- Let students plan and prepare healthy meals.
- Have students select healthy foods from restaurant and cafeteria menus.
- Teach students how to use nutrition labels to make healthy food choices.
- Teach students ways to modify recipes and prepare foods to reduce fat and sodium content and to increase fiber content.
- Help students identify incentives and reinforcements for their current eating and physical activity behaviors.
- Have students examine media and social inducements to adopt unhealthy eating and physical activity patterns, teach them how to respond to these pressures, and let them use their new knowledge to identify their own resistance strategies.
- Have students analyze environmental barriers to healthy eating and physical activity; explore strategies for overcoming these barriers.
- When appropriate, give students practice in encouraging parents to make healthy choices about eating and physical activity at home.
- Teach students to record their food intake, then have them assess and compare their diets with the standards set forth in the Dietary Guidelines for Americans and the Food Guide Pyramid. Have them assess and compare their intake of key nutrients (e.g., calcium and iron) with the intake recommended by the Public Health Service.
- Have students set goals for healthy changes in eating and physical activity, identify barriers and incentives, and assess alternative strategies for reaching their goals and decide which to follow. Show students how to monitor their progress, revise their goals if necessary, and reward themselves for successfully attaining their goals.
- Teach students how to evaluate nutrition claims from advertisements and nutrition-related news stories.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to lists@list.cdc.gov. The body content should read *subscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/> or from CDC's file transfer protocol server at <ftp.cdc.gov>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (404) 332-4555.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

☆U.S. Government Printing Office: 1996-733-175/47007 Region IV

UNITED STATES GOVERNMENT PRINTING OFFICE
SUPERINTENDENT OF DOCUMENTS
Washington, D.C. 20402

OFFICIAL BUSINESS
Penalty for Private Use, \$300

BULK RATE
POSTAGE & FEES PAID
GPO
Permit No. G-26



U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement (OERI)
Educational Resources Information Center (ERIC)



NOTICE

REPRODUCTION BASIS

This document is covered by a signed "Reproduction Release (Blanket)" form (on file within the ERIC system), encompassing all or classes of documents from its source organization and, therefore, does not require a "Specific Document" Release form.

This document is Federally-funded, or carries its own permission to reproduce, or is otherwise in the public domain and, therefore, may be reproduced by ERIC without a signed Reproduction Release form (either "Specific Document" or "Blanket").