

DOCUMENT RESUME

ED 400 494

CG 027 381

TITLE Substance Abuse and the American Woman.
 INSTITUTION Columbia Univ., New York, NY. National Center on
 Addiction and Substance Abuse.
 SPONS AGENCY Bristol-Myers Squibb Foundation, Inc., New York, NY.;
 Pew Charitable Trusts, Philadelphia, PA.
 PUB DATE Jun 96
 NOTE 308p.
 PUB TYPE Reports - Research/Technical (143)

EDRS PRICE MF01/PC13 Plus Postage.
 DESCRIPTORS Alcoholism; At Risk Persons; *Behavior; Drug Abuse;
 Drug Addiction; *Drug Use; Family Environment;
 *Females; Literature Reviews; Sex Differences; *State
 of the Art Reviews; *Substance Abuse

ABSTRACT

The first comprehensive assessment of substance abuse and women, this report arose from an analysis of more than 1,700 scientific and technical articles, surveys, government reports and books. Results show that American women are closing the gap with men in that they are increasingly likely to abuse substances at the same rate as men. Findings show that women are starting to smoke, drink, and use drugs at earlier ages than ever before. Unlike men, though, women get drunk faster, become addicted quicker, and develop substance abuse-related diseases sooner. Furthermore, at least one of every five pregnant women uses drugs, drinks, or smokes. An enormous gap exists between what experts know about women's substance abuse and what is known and acted on by women and those who care for them. It is recommended that women be made aware of the dangers of substance abuse. Prevention programs must address the reasons why women abuse substances and such programs must identify girls at highest risk. Health professionals, too, must recognize that women will manifest symptoms of substance abuse that are different from those of men, which may include a woman trying harder to hide her substance abuse due to intense shame. Contains approximately 750 references. (RJM)

 * Reproductions supplied by EDRS are the best that can be made *
 * from the original document. *



The National Center on
Addiction and Substance Abuse
at Columbia University

ED 400 494

152 West 57th Street
New York, NY 10019-3310

phone 212 841 5200
fax 212 956 8020

Board of Directors

Joseph A. Califano, Jr.
Chairman and President

James E. Burke

James Dimon

Mary Fisher

Betty Ford

Douglas A. Fraser

Donald R. Keough

LaSalle D. Leffall, Jr., M.D.

Manuel T. Pacheco, Ph.D.

Nancy Reagan

E. John Rosenwald, Jr.

George Rupp, Ph.D.

Michael P. Schulhof

Barbara C. Jordan (1936-1996)

Frank G. Wells (1932-1994)

Substance Abuse and The American Woman

June 1996

"PERMISSION TO REPRODUCE THIS
MATERIAL HAS BEEN GRANTED BY

D. MAN

TO THE EDUCATIONAL RESOURCES
INFORMATION CENTER (ERIC)."

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.

° Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

**Funded by: Bristol-Myers Squibb Foundation, Inc.
The Pew Charitable Trusts**

027381



Board of Directors

Joseph A. Califano, Jr.

Chairman and President of CASA

James E. Burke

Chairman of the Partnership for a Drug-Free America

James Dimon

President, Chief Operating Officer and Director of Travelers Group

Mary Fisher

Founder of Family AIDS Network

Betty Ford

Chief Executive Officer of the Betty Ford Center in Rancho Mirage, California

Douglas A. Fraser

Professor of Labor Studies at Wayne State University (former President of United Auto Workers)

Donald R. Keough

Chairman of the Board of Allen and Company Incorporated
(former President of The Coca-Cola Company)

LaSalle D. Leffall, Jr., M.D., F.A.C.S.

Charles R. Drew Professor of Surgery, Howard University Hospital

Manuel T. Pacheco, Ph.D.

President of The University of Arizona

Nancy Reagan

Former First Lady

E. John Rosenwald, Jr.

Vice Chairman of The Bear Stearns Companies Inc.

George Rupp, Ph.D.

President of Columbia University

Michael P. Schulhof

Barbara C. Jordan (1936-1996)

Frank G. Wells (1932-1994)

Advisory Board

SHEILA B. BLUME, M.D.

Medical Director
Alcoholism, Chemical Dependency and
Compulsive Gambling Programs
South Oaks Hospital

WILLIAM A. VEGA, PH.D.

Professor
University of California
at Berkeley
School of Public Health

SHIRLEY D. COLETTI

President
Operation Parental Awareness and Responsibility

EVELYN DAVIS, M.D.

Assistant Clinical Professor of Pediatrics
Department of Child Psychiatry
Harlem Hospital Center

LORETTA P. FINNEGAN, M.D.

Director
Women's Health Initiative
National Institutes of Health

MINDY T. FULLILOVE, M.D.

Associate Professor of Clinical Psychiatry
and Public Health
Columbia University

THEODORE JOYCE, PH.D.

Professor of Economics
Baruch College, City University of New York
and
Research Associate
National Bureau of Economic Research

SISTER DOROTHY ANN KELLY, O.S.U., PH.D.

President
College of New Rochelle

DOROTHY P. RICE, SC.D.

Professor Emeritus
University of California
at San Francisco
Institute of Health and Aging

ANNE M. SEIDEN, M.D.

Chairman
Department of Psychiatry
Cook County Hospital

ZILI SLOBODA, SC.D.

Director
Division of Epidemiology and
Prevention Research
National Institute of Drug Abuse

TABLE of CONTENTS

Executive Summary and Foreword 1

Chapter I -- The Wrong Way 17

Chapter II -- The Difference 27

Chapter III -- Damage and Death: The Truth and Consequences for Women 45

Chapter IV -- Drugs, Alcohol and Cigarettes During Pregnancy:
A Lethal Combination for Mother and Child 69

Chapter V -- Combatting the Problem 95

Chapter VI -- Next Steps: Acting on What We Know and Building Knowledge 131

References 143

Executive Summary

and

Foreword

Substance Abuse and The American Woman is the first comprehensive assessment of the impact on women of all substance abuse--illegal drugs, alcohol, tobacco and prescription drugs. The National Center on Addiction and Substance Abuse at Columbia University (CASA) has conducted an analysis of the available data and more than 1,700 scientific and technical articles, surveys, government reports and books over the past two years in an effort to lift the curtains of denial that have hidden the major public health problem of women's substance abuse for so long.* As a result of this study, CASA will assemble an issue of the *American Journal of Public Health* dedicated to women and substance abuse to be published in the spring of 1997.

The results are disturbing. In the worst way, American women are closing the gap with men: Women are increasingly likely to abuse substances at the same rate as men and women are starting to smoke, drink and use drugs at earlier ages than ever before. Women get drunk faster than men, become addicted quicker and develop substance abuse-related diseases sooner. At least one of every five pregnant women uses drugs, drinks or smokes, putting herself and her newborn in great and avoidable danger.

There is an enormous chasm between what the medical and scientific experts know about women's substance abuse and what is known and acted on by women, their husbands

* A copy of the complete bibliography is available from CASA. To order call (212) 841-5227 or fax (212) 956-8020.

and friends, their doctors and other health professionals. Too many of us do not understand the different ways in which women exhibit symptoms of abuse, the distinct risk factors and reasons why women begin to abuse substances, what makes it difficult for them to stop, and that women need treatment tailored to their specific needs. Until we recognize that what's good for the gander may not help the goose, we will not have effective prevention and treatment programs for women.

First, every woman should be aware of the danger in which substance abuse places her. Women are more vulnerable than men to the immediate and long-term consequences of alcohol and drug abuse. Drinking like men will cause women to develop diseases faster than men. Drug and alcohol abuse make women more likely to be victims of rape and other violent crimes and abuse. Abusing alcohol while taking prescription drugs places women in the fast lane to dependence. One of every two female smokers will die of a tobacco-related disease.

Second, prevention programs for girls and women must be designed around the reasons why they abuse substances. Preventive programs must identify girls at highest risk, such as those who have been physically or sexually abused. These programs must recognize the link women see between being thin and smoking or using drugs like cocaine and heroin and the overriding importance women attach to being thin. (On any given day, 40 percent of American women are on diets.) They must consider that women may smoke or abuse alcohol and drugs to cope with stress or sexual dysfunction and the likelihood that girls are more responsive to peer pressure than boys. Only after accounting for these and other

differences in the reasons why women and men use tobacco, alcohol and drugs will this nation be able to mount an effective substance abuse prevention campaign.

Third, health professionals (female and male), as well as husbands, friends, teachers--each of us--must become better able to identify women who have drug and alcohol abuse problems. Family doctors accustomed to diagnosing drug and alcohol problems in men by external manifestations, such as drunk-driving and frequent fighting, should become better aware that a woman's symptoms are usually inner-directed--depression, anxiety, low self-esteem--and that a woman's often intense shame can lead her to try harder than a man to hide her substance abuse from family, friends and physicians.

Fourth, these two years of work have uncovered a tremendous void in research on women and substance abuse. Much past research was conducted on men only; the results were assumed to apply to men and women alike. The more recent recognition that women's problems require separate attention has served to underscore how much we do not know. We need better understanding of how to motivate women to avoid substance abuse, why women abuse substances and what treatments work, for which women, under what circumstances. This is especially true with respect to elderly women. We need a women's agenda for substance abuse research.

Closing the Gap

In the best of ways, American women are closing the gap with men--filling posts as corporate officers, law firm partners, doctors, academics and in other professions once not open to them. At the same time, in the worst of ways, women are becoming like men--in

the extent to which they abuse alcohol, tobacco, illegal drugs and prescription medication, and in the high price they pay for it. Today, 21.5 million women smoke, 4.5 million are alcoholics or alcohol abusers, 3.5 million misuse prescription drugs and 3.1 million regularly use illicit drugs.** While these numbers are lower than the corresponding figures for men, the gap between the sexes has been closing.

From 1965 to 1994, the percentage of men who smoke dropped 46 percent (from 52 percent to 28 percent), but among women it dropped only 35 percent (from 34 percent to 22 percent). If this trend continues, the United States will be the first nation in the world where the number of women who smoke equals the number of men who smoke.

The percent of drug addicts who are women doubled between 1960 and the late 1970s. Today, some 40 percent of crack addicts are women. The percentage of women (3.7 percent) and men (3.9 percent) who abuse prescription drugs is already equal. Women receive two-thirds of prescriptions for tranquilizers and anti-depressants. Women taking these medications often do not understand or accept the great danger in using them while drinking alcohol. Washing down a few tranquilizers or anti-depressants with wine, beer or cocktails invites dependence and danger.

Among adolescents, the gender gap is gone. Since 1975, girls have been equally or more likely than boys to smoke. Adolescent boys and girls are now equally likely to drink. Among adults men are more likely to have used illegal drugs, but among teens girls and boys are equally likely to have used illicit drugs. For adults over age 30, the ratio of men to women who ever used illicit drugs other than marijuana is 1.5 to 1; for 12- to 18-year-olds,

** These numbers cannot be added because many women abuse more than one substance.

the ratio is 1 to 1. Age of first drug use is also about the same, having dropped for both boys and girls. For adults age 19 to 44, the average age of first marijuana use is 17.2 for women and 16.7 for men; among adults age 45 and older it is 29.5 for women and 24.9 for men. Today's daughters are 15 times likelier than their mothers to have begun using illegal drugs by age 15. In substance abuse and addiction, women have come the wrong way.

What's Use for Men Can Be Abuse for Women

As serious as it is, the shrinking gender gap is only half the problem.

What's moderate drinking for a man spells big trouble for a woman. Women become intoxicated after drinking roughly half as much as men and suffer more quickly the long-term effects of alcohol. They metabolize alcohol differently, get drunk faster, become addicted more easily and develop cirrhosis of the liver more readily. Female alcoholics are up to twice as likely to die as male alcoholics in the same age group. A greater percentage of alcoholic women than men die from alcohol-related accidents, violence and suicide. Among teens in alcohol treatment, most girls had considered suicide, while most boys had not. The risk of liver cirrhosis becomes significant for women at less than two drinks a day; for men this risk becomes significant after six drinks per day.

The consequences of drug abuse strike women with a special vengeance. While fewer than half of men with AIDS acquired HIV from injection drug use, 70 percent of women did so--either by sharing dirty needles or having sex with an injection drug user (among crack addicts, often in exchange for drugs). The added feminine twist is that a woman is more likely to get AIDS from an infected man than a man is to get AIDS from an infected woman.

Women are paying an increasingly frightful toll for smoking. Since 1986, more women have died from lung cancer than from breast cancer. Every year, at least 140,000 women die from illnesses attributable to cigarettes: 62,000 from heart disease, 36,000 from lung cancer, 25,000 from chronic lung disease, 10,000 from other cancers including those of the esophagus, larynx and cervix, and 8,000 from lower respiratory infections.

These and other illnesses and accidents attributable to female substance abuse accounted for \$68 billion in health care costs in 1995, or 12.3 percent of all health care spending for women.

Diseases are just one part of the onslaught which daily threatens substance abusing-women. The link between substance abuse, sex and violence is particularly sinister for women, starting at early ages. Teenage girls who drink more than five times a month are five times more likely to have sex and a third less likely to have it with a condom than girls who do not drink. Women who have been drinking are likelier to become the objects of unwanted sexual advances on the part of men; 60 percent of women who had been drinking reported that another drinker had become more sexually aggressive to them. The link between assaults, including rape, and alcohol abuse is tight: in 75 percent of rapes and 70 percent of domestic violence, either the victim or the assailant has been drinking.

The changing role for women in today's society has given them unprecedented opportunities, but it has also planted land mines on their road to success. Women who work in male-dominated environments are more likely to abuse alcohol than those who don't. Working women are 89 percent likelier to abuse alcohol than homemakers (1.7 percent

versus 0.9 percent). The most likely to be substance abusers are working women who are without work (7.2 percent).

Risk factors for substance abuse among women are quite different from those among men. Female alcoholics are more likely than male alcoholics to have a mental health disorder. Women who drink or smoke are more likely to be depressed than men who do so. Nearly 70 percent of female substance abusers in treatment were sexually abused as children; only 12 percent of men were. Eating disorders like bulimia and anorexia occur more frequently among alcoholic women than other women. Alcoholic women are almost five times likelier to attempt suicide than non-alcoholic women of similar age and income.

Illicit drug use and sales account for much of the sharp rise in female prison populations since 1980. Although there are far more men than women in federal and state prisons, from 1980 to 1994 the number of women prisoners quintupled, while the number of men only tripled. By 1991, a third of female state inmates had violated drug laws, compared to a fifth of male inmates, and this does not include all women who committed their crimes while under the influence of drugs or alcohol, or who did so in order to buy drugs.

Threatening the Health of Babies

More than one in every five pregnant women--at least 800,000 women a year--endangers her own and her newborn's health by smoking, drinking and/or using drugs during pregnancy. Drug, alcohol and tobacco use poses one of the greatest threats to the health of newborn babies in this country.

The term "crack baby" makes for great television soundbites and tabloid headlines, but it masks a far more complex and common reality. Virtually every child exposed to crack or cocaine in the womb has also been exposed to other illegal drugs, tobacco and/or alcohol. Far more newborns are Virginia Slims babies, Newport babies or other cigarette babies (820,000 or 20.4 percent) and beer babies, wine babies or other alcohol babies (757,000 or 18.8 percent) than crack babies or other drug babies (500,000 or 13 percent).

Prenatal exposure to tobacco, alcohol and drugs can be tragic for the fetus. Cigarettes account for 20 percent (61,000) of all low birth weight babies, making tobacco the number one preventable cause of this frequent precursor to infant death. Smoking causes up to 141,000 miscarriages, 4,800 newborn deaths and 2,200 deaths from Sudden Infant Death Syndrome. Even after adjusting for factors such as education and income, smoking during pregnancy has been found to lead to lower scores on intelligence, hearing and language tests.

Like tobacco, alcohol poses a great threat to the health of newborns, accounting for 10 percent of all mental retardation. Prenatal alcohol exposure is the single greatest preventable cause of mental retardation. Estimates of Fetal Alcohol Syndrome, which is characterized by behavioral abnormalities, physical deformities and mental retardation, run as high as 12,000 babies a year. Thousands more newborns suffer the less severe birth abnormalities termed Fetal Alcohol Effects.

Much less is known about the specific effects of particular illegal drugs on newborns, but the 1,370 babies born each day exposed to these drugs make clear the pressing need to close this research gap. Of all pregnant drug users, 53 percent smoke marijuana, 20 percent

use cocaine, and 14 percent use heroin, methamphetamines or other illegal drugs. Some 27 percent misuse prescription drugs.

Injection drug use, usually heroin, signs death warrants for thousands of babies before they've received birth certificates. From 1985 to 1995, 16,000 babies were born HIV-positive, infected by their mothers, most of whom had acquired the virus by sharing needles, trading sex for drugs or having sex with an injection drug user. For these babies, the issue is whether they will die before they become orphans.

Care for alcohol-, tobacco- and drug-exposed babies does not come cheaply. The health care costs for babies exposed to alcohol, tobacco and illegal drugs in the womb came to \$6 billion in 1995. After release from the hospital costs continue to mount. It can cost \$1 million in health care, social services and education to raise a drug-exposed child to age 18.

Illegal drug use not only puts the health of the fetus in danger, it signals more trouble to come. As the number of children in foster care jumped from 280,000 in 1986 to 429,000 in 1991, the percentage of those children under age four exposed prenatally to drugs doubled from 29 percent to 62 percent. From 1985 to 1994, reports of child abuse and neglect rose 64 percent from just under 2 million to more than 3 million; in some cities, more than 75 percent of such cases are linked to drugs and alcohol. In 1994, in New York City, caring for 36,000 children whose parents abuse drugs or alcohol triggered 77 percent--\$595 million of the \$775 million--of the total foster care costs.

Women, Alcohol and Sexual Dysfunction

As far back as ancient Greece and Rome, women and men have believed that alcohol facilitates sexual enjoyment. However, though alcohol can be disinhibiting for men and women, it depresses physical sexual response. The contrast between expectations and physical reality can be devastating for women. Alcoholic women who report problems of sexual dysfunction (which usually preceded their heavy drinking) often use alcohol to "treat" their problem. Since they end up aggravating it, they find themselves in a vicious circle, medicating their sexual dysfunction with more alcohol, only to make both their addiction and dysfunction worse. It is important for women to understand this link between alcohol and sexual dysfunction.

Tailoring Prevention and Treatment for Women

The closing gender gap, particularly among teenagers and young adults, and increasing health care, social service and criminal justice consequences of substance abuse as the gap closes, cry out for increased prevention and treatment. But to be effective these prevention and treatment programs must take into account the ways in which a woman is not like a man.

Prevention. We should launch a major campaign targeted at pre-teen and teenage girls, tailored to their needs and attacking all substance abuse. Based on everything we know, a young woman or man who gets to age 21 without smoking, abusing alcohol or using illicit drugs is virtually certain never to do so.

Any attempt to fight the deadly consequences of tobacco use by women must address the link between smoking and thinness, a relationship the nicotine pushers cynically exploit. Many women hesitate to quit, or quit only to relapse, because they fear the 11 pound weight gain that often accompanies cessation.

Women's magazines, as well as television and the movies, should take more responsibility here. Fulfilling this responsibility will require many of these magazines to break their own addiction to tobacco advertising revenue. Magazines violate their relationship of trust with their readers when they fail to report fully the health consequences of smoking. Entertainment executives should take care that they do not encourage girls to begin abusing substances by creating television shows and movies that glamorize women who abuse alcohol and smoke without ever showing the diseases, violence and other disastrous consequences of this abuse. The fashion industry should ease off its adulation of social x-ray thinness and display their wares on healthier women.

Treatment. The first step is diagnosis--and the earlier the better. Unfortunately, however, doctors are more likely to miss a substance abuse diagnosis in women than in men, despite the fact that women see physicians more often. Better physician training is essential here, in medical school and continuing medical education programs. Working with Columbia University's College of Physicians and Surgeons, CASA has begun testing the use of recovering female substance abusers in a continuing education program for family practitioners, pediatricians and psychiatrists who specialize in treating adolescents.

We need more treatment programs and we need to learn more about how to encourage girls and women to enter them. In 1989, the year of the most recent

comprehensive survey, less than 14 percent of all women and 12 percent of pregnant women who needed treatment received it. It is a commentary on the tendency to ignore the problem of women's substance abuse that a more recent survey has not been taken, but there is no indication that the situation has improved since then.

Even when slots are available, getting women to treatment is especially difficult. Doctors and nurses are slow to recognize the markers of female substance abuse and often mistake these signs--depression, low self-esteem--for emotional disturbances. Friends, accustomed to seeing the public displays of male abuse, such as crime and violence, often overlook developing inner-directed symptoms among women. Families often discourage women from leaving the home to get help because they feel they need mother to run the household and keep the family together. The stigma surrounding female substance abuse, which for cultural and historical reasons is greater than that for men, has discouraged many women from seeking help.

Treatment programs should provide for the child care needs of women abusers. Those that have a child care component or allow the child to stay with the mother tend to have higher retention rates, a factor associated with greater success. Because women's substance abuse is often tied to physical and sexual abuse during childhood and depression, treatment programs which do not address these underlying issues court failure. More than for men, for women substance abuse is intertwined with the drug and alcohol abuse of their partners: thus, dealing with a husband's abuse is part and parcel of addressing a wife's substance abuse.

Because pregnancy can be a window of opportunity for getting a woman into treatment and because substance abuse during this time harms both the woman and the fetus, effectively treating expecting addicts is a matter of special importance.

Rethinking Conventional Wisdom

Drug and alcohol abuse, particularly the explosion of crack cocaine, forces all of us to rethink our assumptions about families, our foster care system and how to insure a child's well-being. Foster care is no longer about Little Orphan Annie and Daddy Warbucks; it has become a system to salvage the wreckage left in the wake of parental drug and alcohol abuse.

As a society, we must begin to ask, and answer, some difficult questions: Does the assumption that a child is best left with the family hold true when drugs and alcohol have been related to so many cases of neglect, physical beating, sexual abuse and even murder? Do our family court judges have adequate time and expertise to make judgements on what's best for the child? What kind of foster care system do we need today? Are adoptive homes or orphanages better for children than parents with drug or alcohol problems? Can temporary separation from the child motivate substance abusing-parents to seek treatment and stay off drugs and alcohol? How can we identify parents who will be so motivated? How can we identify parents who are more likely to shake their addiction if their children remain with them?

Conclusion

The origins, patterns and consequences of substance abuse are different--and often far more devastating--for women than for men. What motivates a woman to seek--or not to seek--treatment is likely to be different from what motivates a man. Treatment programs must take into account the distinct needs of women.

Those are the differences. But there is a grim consequence common to women and men: women who smoke, abuse alcohol and use illegal drugs like men, will die like men who smoke, abuse alcohol and use illegal drugs--from cancers and heart disease, from violence and AIDS.

This study was made possible by the support of the Bristol-Myers Squibb Foundation, Inc., an extraordinary example of corporate responsibility especially sensitive to women's issues, and The Pew Charitable Trusts, one of CASA's initial supporters. The law firm of Cadwalader, Wickersham and Taft generously conducted, pro bono, a state-by-state survey of laws and court decisions regarding pregnant women and substance abuse.

Jeanne L. Reid was the principal investigator with the guidance of Jeffrey Merrill, Vice President and Director of Policy Research and Analysis. Sharon Gray, CASA's Librarian, Susan Lewis and Russell Jones helped to research the report, which I reviewed and edited. Dr. Herbert Kleber, Executive Vice President and Medical Director, Hila Richardson, Deputy Director of Medical Research and Practice Policy, and John Demers, my Research Assistant, reviewed drafts of the report.

We greatly appreciate the help of outside professionals who made up our advisory board. These talented and dedicated individuals, with extensive experience with the problem of substance abuse among women, provided valuable guidance in setting the course of study and reviewing the report.

We are grateful for the support of our funders and guidance of our advisors, but CASA alone is responsible for the report's contents and conclusions.

Joseph A. Califano, Jr.

I.

The Wrong Way

It's time to face the facts about substance abuse and the American woman.

While more men than women abuse alcohol, tobacco, and licit and illicit drugs, the gender gap has been closing since World War II.¹ Today, one in five American women (compared to one in three men) abuses or becomes dependent on alcohol and/or drugs at some point in her life.²

At least 4.5 million women are alcohol abusers or alcoholics;³ 3.1 million regularly use illicit drugs; 3.5 million misuse prescription drugs; and 21.5 million smoke cigarettes.^{**} From lung cancer, heart disease, liver cirrhosis, osteoporosis and AIDS, to domestic violence, lost jobs and shattered families, to the damage done to children when mothers use drugs, alcohol or tobacco during pregnancy, the consequences of substance abuse hit women with tragic and deadly force.

Each year, women give birth to some 500,000 babies who have been exposed prenatally to illicit drugs.^{***} More than five of 10 pregnant drug users smoke marijuana;

* In this study, women and men are ages 15 to 54. "Abuse" and "dependence" are defined according to the Diagnostic and Statistical Manual of Mental Disorders, edition III.

** Unless otherwise noted, all data in this report are from the U.S. Department of Health and Human Services' 1993 National Household Survey on Drug Abuse (NHSDA); women and men are adults over 18 years old. "Abuse" means problem use or dependence. "Alcoholic" is used to connote dependence. Regular illicit drug users took drugs at least monthly during the past year. Prescription drug misusers took stimulants, sedatives, tranquilizers or analgesics non-medically during the past year. Current smokers had at least one cigarette during the past month. Smoking data only are from the 1994 NHSDA, which measured adults ages 18 and over. The numbers in the text above cannot be added because many women use, abuse and misuse more than one substance.

*** National estimates range from 212,000 to 736,000 prenatally exposed newborns each year.

six of 10 use an illicit drug other than marijuana; one of five snorts or smokes cocaine; and one of four misuses prescription drugs.⁵ Prenatal use of tobacco and alcohol is much more common; each year, roughly 820,000 women smoke and 760,000 women drink alcohol during pregnancy. These numbers cannot be added because most women use more than one substance.

Smoking

Because the long-term decline in smoking that began in 1965 has been less dramatic among women than men, the gender gap in smoking rates has narrowed. From 1965 to 1994, the percentage of women who smoke dropped 35 percent (from 33.9 percent to 22.1 percent); among men it fell 46 percent (from 51.9 percent to 28.0 percent).^{*} Adolescent girls and boys are now equally likely to smoke. If this gender equity continues, shortly after the turn of the century the U.S. could become the first nation in the world where equal numbers of women and men smoke.⁶

Tobacco companies began targeting women in the 1920s, pitching cigarettes as a route to thinness.⁷ "Reach for a Lucky instead of a sweet," said the Lucky Strike ad.⁸ The American Tobacco Company described the female market as "a new gold mine right in our front yard." Indeed, Lucky Strike sales almost tripled from 13.7 billion cigarettes in 1925 to more than 40 billion in 1930, making it the leading brand nationwide.⁹

^{*} 1965 National Health Interview Survey and 1994 NHSDA; data from both surveys measure adults age 18 and over.

After World War II, tobacco companies heralded the modern woman who--still skinny--smoked cigarettes as a sign of her spunk. From Virginia Slims' rallying cry, "You've come a long way baby!" to the updated, "A woman's place is any place her feet will take her," or Misty cigarettes' "Slim 'n Sassy" slogan, cigarette makers continued to trumpet the link between thinness and smoking, merely adding the rebellious spirit of the independent woman who might flout the evidence that nicotine is addictive and deadly.

From the early 1960s to the mid-1980s, as the first large group of women who smoke heavily reached their 50s and 60s, the death rate from lung cancer among female smokers soared 496 percent (from 26 to 155 per 100,000), six times the rate of increase among male smokers (from 187 to 341 per 100,000).¹⁰ In 1986, for the first time, more women died of lung cancer than of breast cancer.¹¹ A woman who smokes is up to four times likelier to get heart disease, the leading killer of women.¹² Smoking also increases a woman's risk of osteoporosis, producing brittle bones that can break in old age, robbing her of her independence.¹³ Such facts prompted Ms. magazine to ask, "If we've come such a long way, why are we still smoking?"¹⁴

Alcohol Abuse

Women are especially susceptible to the ill-effects of alcohol. They develop alcohol-related illnesses such as liver cirrhosis, hypertension, anemia and malnutrition more rapidly than men.¹⁵ Women who drink heavily are more likely than those who don't to get breast cancer, to be infertile and to suffer violent abuse at the hands of their partners.¹⁶ Teenage girls who drink are more likely than boys who drink--and girls who don't drink--to attempt

suicide, to have sex and have it without a condom, which can lead to unplanned pregnancies and sexually transmitted diseases such as AIDS and gonorrhea.¹⁷

Illegal Drugs

Women move more quickly than men from trying a drug to getting hooked, and female addicts face more barriers to treatment than male addicts.

Since the advent of crack cocaine in the mid-1980s, the number of crack-exposed newborns and "boarder" babies left in hospitals by cocaine-addicted parents or held there by child welfare investigators has jumped sharply.¹⁸ Reports of child abuse and neglect, most related to drug and alcohol abuse, rose 64 percent, from 1.92 million in 1985 to 3.14 million in 1994.¹⁹ The percentage of children in foster care under age four who had been exposed prenatally to drugs more than doubled, from 29 percent in 1986 to 62 percent in 1991.²⁰ From 1985 to 1995, some 16,000 children contracted the AIDS virus in the wombs of their mothers, most of whom were infected through their or their sexual partner's injection drug use.²¹ Seventy percent of AIDS cases among women stem from such illicit drug use.²²

The criminal justice consequences of women's involvement with illegal drugs are also increasing at alarming rates. Illicit drug use and sales have sparked the 386 percent rise in the federal and state female prison population, from 12,331 in 1980 to 59,878 in 1994, while the number of men rose 214 percent from 303,643 to 952,585.²³ Among state inmates, a third of women had violated drug laws in 1991, compared to a fifth of men.²⁴ Many more had committed their crimes under the influence of drugs or alcohol, or to get money to buy drugs.

Stigma

Despite the scope of the problem, a woman's substance abuse has traditionally been tucked behind the curtains of private homes, while a man's is often a public event at bars, athletic events and fraternity parties.²⁵ Given the different standards to which women traditionally have been held, a female who slips into addiction invites more scorn than her male counterpart.²⁶ This stigma has discouraged many women from seeking help for their substance abuse, deterred them from entering treatment and contributed to a scarcity of research, prevention and treatment efforts that address the unique needs of women.²⁷

How Far Have We Come?

American women have a long history of substance abuse and addiction. In the 19th century, most of the nation's 250,000 opiate addicts were women.²⁸ Surveys in Michigan in 1878, Chicago in 1880 and Iowa in 1885 found that 61 percent to 72 percent of opiate users were women, typically upper- or middle-class white housewives or socialites.²⁹

Women had easy access to drugs from stores that sold over-the-counter patent medicines containing opiates. Doctors commonly recommended them to women for "nervousness," "melancholy" and "female troubles," such as menstrual pain.³⁰ A 1914 Tennessee study found that most female opiate users were ages 25 to 45, "when the stresses of life begin to make themselves felt with women, and [when menopause begins]. . . . It appears reasonable, therefore, to ascribe to this part of female life, no small portion of the addiction among women."³¹ Manufacturers gave opiate "medicines" comforting names such

as "Mrs. Winslow's Soothing Syrup," and advertised them as "women's friends." By the early 1900s, Americans were spending \$100 million a year for such potions.³²

The popularity of opiates stemmed in part from the cultural disdain for women who drank alcohol in public.³³ Women could take opiate "medicines" in the privacy of their own home. Opiate use became so fashionable that Macy's department store sold "exquisitely jewelled gold morphine sets," which contained miniature morphine and cocaine syringes.³⁴ In contrast, drinking was a social event for men in bars and saloons, where women's presence was considered inappropriate. As one commentator put it in 1894, "Twenty years ago I rarely ever saw a female drinking at the bar of a public house or beerhouse. Now I see numbers so engaged from an early hour in the morning, not a few of these early risers and early drinkers having had an infant at the breast, and giving the child a share of the morning dram."³⁵ Women leading the Temperance Movement reinforced the belief that those who drank to the point of intoxication were "fallen women," often thought to be promiscuous.³⁶

Concerned about the spread of opiate and cocaine addiction, Congress passed the Harrison Act in 1914 to restrict narcotic prescriptions.³⁷ While many women shied away from illicit sales, some continued to get opiates fraudulently from their doctors; others sought prescriptions for hypnotic and sedative drugs such as barbiturates.³⁸ The Public Health Service Hospital in Lexington, Kentucky--one of two Federal facilities to treat addicts and the only one to accept women--reported that from 1941 (when it began accepting female opiate addicts) to 1965, 16.5 percent of its admissions--14,866 clients--were women.³⁹

With the passage of Prohibition in 1920, speakeasies welcomed both male and female customers. Women's drinking appears to have flourished in the devil-may-care era of the flapper.⁴⁰ Yet the stigma endured, particularly for women who got drunk.⁴¹ When women advocated repeal of Prohibition in the 1930s, the American Independent, a Kentucky newspaper, declared that a woman who opposed Prohibition was "either a drunkard," or had an immoral "home life. . . . Most of them are no more than the scum of the earth, parading around in skirts, and possibly late at night flirting with other women's husbands."⁴²

To tap the women's market, alcohol companies countered this stigma by depicting the modern women's drinking as a sign of sexy assertiveness. In 1975, the Distilled Spirits Council allowed its members to use women in ads that were "dignified, modest and in good taste," which opened the door to attractive women draped over drinking men's shoulders.⁴³ Dewar's Scotch updated this image in the 1990s with ads aimed at young female professionals in which a woman puts on her work clothes while a bare-chested man sleeps in the bed beside her.⁴⁴ The slogan: "You finally have a real job, a real place and a real boyfriend. How about a real drink?"

New Light On An Old Problem

Over the last 30 years, women's growing presence in factories, offices and legislatures has given visibility to their battles with drugs and alcohol. The entertainment industry dramatized the stereotype of the unhappy housewife who pops pills and empties the liquor closet in the privacy of her home.⁴⁵ And the courageous leadership of Betty Ford,

who publicly acknowledged her own alcohol and pill problem, challenged the belief that only "a certain kind" of woman could be an addict.⁴⁶

In 1976 the National Council on Alcoholism created a special office on women and held its first national conference on women and alcoholism.⁴⁷ Led by Senator William Hathaway, a recovering alcoholic, the U.S. Senate held hearings on the issue of female drinking, setting the stage for more research on women and alcohol, tobacco and illicit drugs. In 1980, the Department of Health, Education and Welfare issued a Surgeon General's report on women and smoking.

We now know that women are subject to the deadly effects of smoking and are more vulnerable than men to liver cirrhosis and other alcohol-related illnesses. Studies of domestic violence have exposed the extent of spousal abuse when alcohol is used by either victim or perpetrator. Growing evidence of the long-term effects of Fetal Alcohol Syndrome--the mental retardation and birth defects caused by drinking during pregnancy--has inspired a cultural sea-change against the use of alcohol by pregnant woman. Recent studies reveal that most cocaine users also use alcohol and/or tobacco, and suggest that cocaine, tobacco and alcohol can each be severely damaging to the fetus.⁴⁸

The American woman is many people. She is a lawyer, doctor, nurse, engineer, secretary, teacher, housewife, student, mother, daughter and grandmother. She is unemployed, married, single, rich, poor, black, white, Asian, Hispanic and Native American. She is a teenager in high school and a senior citizen in a nursing home. She is a cop on the beat and an inmate on Rikers Island. She is straight and gay, thin and obese,

living in a city and a rural hamlet. She is Catholic, Jewish, Protestant and Muslim. The typical female substance abuser or addict can be any or all of these women.

This report seeks to document what we know about substance abuse and addiction among all women: how their patterns of use and reasons for abuse differ from men and among themselves; the particularly severe health and social consequences they can suffer; their unique responsibilities during pregnancy; their vulnerabilities when drunk; the policy implications that result from these differences; and the importance of recognizing the special needs of women in crafting efforts to prevent and treat substance abuse and addiction.

II.

The Difference

Most substance abuse research has focused on men.¹ Until recently, researchers generally assumed that the nature of substance abuse by women and men is identical or that the size of the problem among women is small, eliminating the need to recruit female subjects. In fact, the problem is widespread, and though women and men often experience similar patterns and consequences of substance abuse, they differ in critical ways.

Alcohol

Although women usually drink less frequently and heavily than men, they may get drunk just as often. On average, women who drink only half of what men drink reach the same level of intoxication.² Studies that compare women's and men's average daily alcohol intake have found a ratio of one to two in the amount they drink, but women metabolize alcohol differently than men.³ As a result, less alcohol can induce a state of intoxication among women. The Department of Health and Human Services and the Department of Agriculture recommend that a woman have no more than one drink a day, while a man should have no more than two.⁴

¹ Differences in body weight, body water and the stomach enzymes that metabolize alcohol give alcohol a more powerful effect on women. Oral contraceptives further reduce a woman's ability to metabolize alcohol.

² A standard drink is 12 oz. of regular beer, 5 oz. of wine or 1.5 oz. of 80-proof distilled spirits; each contains 0.5 oz. of pure alcohol.

Almost half of all women (40.5 million) and more than half of all men (51.2 million) drink.* The 1993 National Household Survey on Drug Abuse (NHSDA), based on self-reported alcohol use, found that 2.5 million women (2.6 percent of all women) drank at least 60 drinks per month--the measure of heavy drinking established by the National Institute of Alcohol Abuse and Alcoholism (CHART 1). But that measure is based on the male standard for heavy drinking: two drinks a day. Since the recommended maximum for women is half that--one drink a day--alcohol abuse and heavy drinking among women is far more widespread than these reports of alcohol intake suggest. The NHSDA found the percentage of women who drank at least 30 drinks per month to be 6.6 percent (about 6.3 million). (The percentage of men who have 60 or more drinks a month is 13.1 (about 11.4 million).) A study that applied mental health diagnostic criteria to drinking-related behavior among adults concluded that 4.5 million women (4.6 percent) abuse or depend on alcohol, and about 10.6 million men (12 percent) do so.⁵

Age

While women, like men, are less likely to drink as they get older, the proportion who drink heavily changes little.⁶ Since tolerance for alcohol decreases with age, this may signal a serious problem among elderly women.⁷

* Current drinkers had at least one drink in the last month.

Chart 1

Adult Women and Men Who Drink Heavily by Age*

Age	Heavy Drinkers**	
	Women	Men
19-30	2.4%	13.0%
31-40	2.4	11.7
41-59	1.8	11.9
60+	4.1	16.5
All	2.6%	13.1%

* Adults are age 19 and older.

** Drinking heavily is having at least 60 drinks in the past month.

Source: 1993 National Household Survey on Drug Abuse

Economic Status

Women and men with lower income are likelier to drink heavily: women in households with income below \$15,000 are about twice as likely to be heavy drinkers as women in households with incomes of \$15,000 or more (S)* (CHART 2). Among homemakers, those with household incomes below \$30,000 are more than four times as likely to drink heavily as those with higher incomes (S).

Education

Heavy drinking rates decline as individuals attain higher levels of education. The exception is binge drinking, which peaks among adults with some college education but without a degree (S) (CHART 3).** Women in college are more likely to binge drink than those of the same age who are not in college (33 percent vs. 25 percent), an indication that binge drinking by women on college campuses is a particularly important target for prevention efforts.⁸

Race and Ethnicity

White women are more likely to drink than African American women (S), but their rates of heavy drinking differ only slightly (NS) (CHART 4). Black women tend to be

* Comparisons using 1993 National Household Survey Data have been tested for statistical significance. Those that reach a level of significance ($p < 0.5$) are noted (S). Those that do not are noted (NS). Data that are not statistically significant may still reflect meaningful comparisons, but CASA could not rule out the possibility that they stem from random chance. This report only includes non-statistically-significant data when it is consistent with other research.

** Binge drinkers have had five or more drinks in a row at least twice in the past month.

concentrated at the extremes of either abstinence or heavy drinking; those who participate actively in religion are more likely to abstain.⁹ Unlike white women, heavy drinking by black women does not drop as their income rises.¹⁰ Native American women have high rates of heavy drinking, but they vary widely by tribal origin.¹¹


Relatively few Asian American or Hispanic women report any heavy drinking, although drinking rates rise as they adopt the attitudes, behaviors and other markers of American culture.¹² In some Hispanic and Asian communities, cultural disapproval of public drunkenness discourages women from drinking, but it may also encourage them to keep any drinking secret.¹³

Employment

It is not clear whether women's migration into the paid labor force over the past 30 years led to an increase in their drinking and alcohol-related problems.¹⁴ New roles in the labor force could nurture a woman's self-confidence and personal fulfillment, diminishing her tendency to abuse alcohol.¹⁵ But her responsibilities in factories and offices could create pressures that led to heavy drinking, or simply provide more opportunities to drink.¹⁶

Surveys have not found a dramatic rise in drinking among women during the last 30 years. In fact, the percentage of women and men who drink decreased in the 1980s.¹⁷ But there is a sharp difference between the older and younger generation of women. Heavy drinking by women ages 18 to 29 did not drop during the 1980s, and their reports of alcohol-

Chart 2
Adult Women Who Drink Heavily*
by Household Income




*Drinking heavily is having at least 60 drinks in the past month.

Source: 1993 National Household Survey on Drug Abuse

BEST COPY AVAILABLE


Chart 3

Adult Women Who Binge Drink* by Education



* Binge drinkers have had 5 or more drinks in one sitting at least twice in the past month.
Source: 1993 National Household Survey on Drug Abuse

Chart 4 Adult Women Who Drink Heavily* by Race and Ethnicity



* Drinking heavily is having at least 60 drinks in the past month. **BEST COPY AVAILABLE**
Source: 1993 National Household Survey on Drug Abuse

related problems rose.^{*18} Today, the ratio of alcohol abuse or dependence among men and women ages 65 and older is 4.4 to 1, but the ratio is only 2.2 to 1 among men and women ages 18 to 29.¹⁹

Women who work outside the home are 67 percent more likely to drink heavily than homemakers (S).^{**} However, the most important work-related risk factor for heavy drinking among adult women is unemployment (S) (CHART 5). A working woman who is unemployed is more than 400 percent likelier to drink heavily than a woman who is working full- or part-time (S). Drinking problems may lead to unemployment for these women and, in turn, unemployment may cause or aggravate drinking problems.

Marital Status

Marital status is another factor in drinking patterns. Women who have never been married (particularly those living with a partner) are 50 percent likelier to drink heavily than married women (NS) (CHART 6).²⁰ Widows are three times more likely than married women to drink heavily (6.2 percent vs. 1.9 percent) (S), which suggests that they are an important target for prevention.²¹ The impact of divorce on women's drinking is complex. When women with alcohol problems divorce, their drinking often subsides, especially if their partner is also a problem drinker.²² But when women without alcohol problems divorce, their drinking often increases.²³

* This study (Wilsnack, et al, 1986) defines "heavy drinking" as having five or more drinks in a row at least weekly during the past year.

** Homemakers are assumed to be employed full-time at home.

Genetics, Family and Personal Experience

Genetics play a strong role in alcoholism among men, but the research among women is sparse and inconsistent.²⁴

So far, the influence of genes appears equally or less powerful among women. Nevertheless, female alcoholics are *more* likely than male alcoholics to have a family history of alcoholism.²⁵ The extent to which family, other environmental factors and genetics influence women demands intensive research since assessing the significance of each factor is important in crafting prevention and treatment strategies.²⁶

A history of childhood abuse--and especially sexual abuse--may play as powerful a role in women's alcoholism as genetics.²⁷ Alcoholic women are twice as likely as non-alcoholic women to have been beaten or sexually assaulted as a child.²⁸ In one study, 69 percent of women in treatment reported being sexually abused as children, compared to 11.6 percent of men.²⁹ Roughly half of these women are victims of incest.³⁰

Adding complexity to the diagnosis and treatment of their substance abuse problems, female alcoholics are much likelier than male alcoholics to have a mental health disorder in


◆ *Alcoholic women are more likely to have been beaten recently by a partner than non-alcoholic women. Women in treatment for alcoholism are four times more likely to have been abused by a partner.*¹

◆ *Women who drink heavily are likely to have a partner who drinks heavily as well. Alcoholic women are five times more likely to have a spouse with alcohol problems than non-alcoholic women.*² *Unless her partner also seeks treatment, separation or divorce may be critical to a woman's ability to stay sober.*³

¹ Miller, B. A. and Downs, W. R. (1993). The impact of family violence on the use of alcohol by women. Alcohol, Health and Research World, 17(2), 137-143; ² Miller, B. A., Downs, W. R. and Gondoli, D. M. (1989). Spousal violence among alcoholic women as compared to a random household sample of women. Journal of Studies on Alcohol, 50(6), 533-540.; ³ Blume, S. B. (1994). Gender differences in alcohol-related disorders. Harvard Review of Psychiatry, 2(1), 7-14.


Chart 5

Adult Women Who Drink Heavily* by Employment Status



* Drinking heavily is having at least 60 drinks in the past month.
Source: 1993 National Household Survey on Drug Abuse

Adult Women Who Drink Heavily* by Marital Status



* Drinking heavily is having at least 60 drinks in the past month.
Source: 1993 National Household Survey on Drug Abuse

BEST COPY AVAILABLE 47

addition to their alcoholism (65 percent vs. 44 percent).³¹ For women, the most frequent problem is depression, while for men, anti-social personality disorder (linked to aggression and criminality) is more common. Women's depression usually precedes the onset of their alcoholism; men's depression tends to follow it. Eating disorders such as bulimia or anorexia are also found more frequently in female alcoholics than in women without alcohol problems.³²

Both female and male alcoholics report drinking to escape life pressures, but women more often report feeling powerless and inadequate before problem drinking.³³ Alcoholic women are also more likely to say their heavy drinking followed a crisis, such as a miscarriage, divorce, unemployment or the recent departure of a child from the home (an "empty nest").³⁴ It is unclear how often these events precede women's drinking and how often drinking causes or contributes to them. But women who lack a personally fulfilling and socially accepted role in life are at much higher risk for alcoholism than women who are happily employed, whether inside or outside the home, or both.³⁵

Tobacco

In 1994, 21.5 million women (22.1 percent) and 24.6 million men (28.0 percent) age 18 and older smoked cigarettes (CHART 7). Most want to quit; in a 1993 survey, almost three-fourths of female smokers (72.7 percent) and two-thirds of male smokers (67.1 percent)

said they would like to stop.³⁶ They are about equally likely to succeed; 69 percent of women and 65 percent of men who have ever smoked managed to quit.*

Why Women Smoke

Though smoking became popular among men early in the 1900s, women did not take up the habit in large numbers until World War II.³⁷ The rate of smoking among men peaked at more than 60 percent in the 1950s, when evidence of smoking's harmful effects began to emerge. But the rate among women did not peak until 1965--at 33.9 percent--when the Surgeon General issued his report on smoking, initiating a long-term decline in smoking rates that has been less dramatic among women than men.³⁸

One reason for the slower decline among women is that soon after the Surgeon General's report, tobacco companies introduced new advertising campaigns to recruit female smokers.³⁹ They tailored ads to the modern woman, most saliently in the Virginia Slims campaign, and placed them in women's magazines, which then rarely reported new evidence that smoking could be deadly for women.⁴⁰ These ads have been linked to sharp increases--110 percent among 12-year-olds and 35 percent among 17-year-olds--in the rate at which girls started smoking from 1967 to 1973; in contrast, the initiation rate among boys did not change.⁴¹ By 1975, more girls than boys were smokers.⁴² Because of a surge in boy's smoking during the late 1980s, girls and boys are now equally likely to smoke (12.0 percent vs. 11.9 percent).⁴³

* The percentage of smokers who have quit is the number of former smokers divided by the number of current and former smokers. Former smokers have not had a cigarette during the past month.

Chart 7

Adult Women and Men Who Smoke*

	<u>Current**</u>	<u>Former**</u>	<u>Never</u>
Women	22.1%	48.9%	29.0%
Men	28.0	52.6	19.4

* The 1994 NHSDA data classifies adults as 18 and older.

** Current smokers have smoked in the past month; former smokers smoked more than a month ago.
Source: 1994 National Household Survey on Drug Abuse

Both girls and boys cite their peers' smoking as a reason they start to smoke.⁴⁴ Receptivity to tobacco advertising may also affect their decision.⁴⁵ But girls in particular are likely to smoke to control their weight.⁴⁶ Among white teens who smoke, girls are three times likelier than boys to smoke to reduce their appetite.⁴⁷

Such worries about weight gain persist in

adulthood; women who smoke are more than twice as likely as men to cite weight concerns as a reason not to quit.⁴⁸ The belief that quitting smoking will lead to weight gain is accurate.⁴⁹ One study found that over 10 years, women who quit gained 11 pounds more on average than those who continued to smoke.⁵⁰

Economic Status and Education

For women and men, smoking rates decrease as income rises. A woman whose income is below \$15,000 is almost twice as likely to smoke as a woman with income of \$75,000 or more (S) (CHART 8). Smoking is also less common among more educated individuals.⁵¹ Women who drop out of high school are three times likelier to smoke than those with graduate degrees (S) (29.7 percent vs. 9.3 percent).⁵² Adults of any income and education are equally likely to start smoking--but as they gain income or education, they are

Andrea Lampson, age 13, believes effective antismoking ads must feature real-life stories. If she could design one herself, it would show children talking about a parent who died from smoking-- "not actors, but real people who know how much it hurts." The quiet eighth-grader expresses concern about her mother's smoking. Yet Andrea confesses she recently considered taking up smoking herself to lose weight. "It speeds up your metabolism," she says.¹

¹ Ono, Yumiko. Teenagers tell which antismoking ads work. *Wall Street Journal*, 8/30/95, p. B1.

more likely to quit. As a result, primary prevention efforts should reach all income and education levels, while quitting campaigns can be more focused.

Race and Ethnicity


Native American women are most likely of all women to smoke (S) (CHART 9); seven out of eight have smoked, and less than half (43 percent) quit. Asian American women are least likely to smoke (S); three-quarters of them never even start, and among those who do, more than half (54 percent) quit. Smoking is slightly more common among African American than white women (27.0 percent vs. 23.8 percent) (NS).⁵³ Though black women are less likely than white women to start, fewer black women quit than white women (57 percent vs. 68 percent) (S). Black women are also less likely to quit than black men (63 percent), which suggests a need for cessation programs that focus on black women.⁵⁴

While smoking declined moderately among white high school seniors during the 1980s,^{*} it appears to have dropped dramatically from 22.3 percent to 7.1 percent among black females and from 23.6 percent to 8.6 percent among black males.⁵⁵ In 1993, the National Household Survey on Drug Abuse reported that only 5.0 percent of black girls and 4.8 percent of black boys smoked; rates among white teenagers were almost three times higher (S) (CHART 10). Because this survey is self-reported and does not include those not in the home, and since it may not accurately reflect rates among the most disadvantaged, these numbers for smoking among blacks may be too low. Nevertheless, religious participation, perception of smoking as a "white" social norm, reaction against the blatant

^{*} Smoking rates in this study (Bachman, Wallace, et al, 1991) are averages for 1976-1979 and 1985-1989.

Chart 8

Adult Women Who Smoke by Household Income



BEST COPY AVAILABLE

* Current smokers have smoked in the past month.
Source: 1993 National Household Survey on Drug Abuse


Chart 9

Adult Women Who Smoke by Race and Ethnicity

	Current*	Former*	Never
White	23.8%	50.9%	25.3%
African American	27.0	35.3	37.6
Hispanic	19.2	30.8	50.1
Asian American	12.6	14.7	72.8
Native American	49.6	37.5	12.9

* Current smokers have smoked in the past month; former smokers smoked more than a month ago.
Source: 1993 National Household Survey on Drug Abuse

Chart 10
Adolescents Who Smoke
by Race and Ethnicity



* Current smokers have smoked in the past month.
Source: 1993 National Household Survey on Drug Abuse

targeting of black youth by tobacco companies and the lower priority black girls give to thinness may discourage many black teens from smoking.⁵⁶ The decline among black teenagers now may be enduring among black women ages 18 to 24. One study reported a decline in smoking rates in this group from 27.1 percent in 1978 to 8.3 percent in 1993.⁵⁷

But the jury is still out. Black women appear to be more likely than white women to take up the habit as adults. More than one in six (17.8 percent) black female smokers say they began after age 20, while fewer than one out of ten (9.8 percent) white female smokers say they began after that age (S). Almost one out of ten (9.1 percent) black female smokers say they started after age 25, but few (2.5 percent) white female smokers say they started after that age (S).

Illicit Drugs

Fewer women than men have used illicit drugs in their lifetime (S) (CHART 11). Women and men ages 19 to 40 are equally likely to have used marijuana (and no other illicit drug) (26.3 percent vs. 25.5 percent). But women in this age group are still less likely than men to have used other illicit drugs such as cocaine and heroin (27.6 percent vs. 38.5 percent) (S), and to be regular users of any illicit drug (5.3 percent vs. 12.6 percent) (S).

However, women are not immune to the lure of "hard drugs." The surge of cocaine use among middle and upper class women during the 1970s and the crack cocaine epidemic among poor women in the 1980s narrowed the gender gap in illicit drug use.⁵⁸ From 1960 to the late-1970s, the percent of drug addicts who were women doubled (from 14 to 30).⁵⁹ During the 1980s, their numbers rose even more rapidly. From 1985 to 1989, the number of

women seeking treatment for crack cocaine at Phoenix House in New York City almost tripled.⁶⁰

Some 40 percent of all crack addicts are women.⁶¹ Experts attribute its popularity among women to its cheap price, their preference for smoking a drug rather than injecting one, and the sense of confidence that crack temporarily delivers.⁶² Since the beginning of the crack epidemic, women's involvement in drug trafficking also has risen.⁶³

Women with higher incomes are most likely to experiment with illicit drugs, but regular drug use is most common among those in poverty (S) (CHART 12). Almost half (45.6 percent) of women with incomes of at least \$75,000 have ever used illicit drugs, but relatively few (1.9 percent) report regular use. White women are more likely than black or Hispanic women to have ever used illicit drugs (S), but the

◆ *Female drug addicts are more likely than male addicts to have a partner who uses illicit drugs.*¹

◆ *Many women say a man introduced them to drugs, while men more often began with male peers. In one study, 33 percent of female heroin addicts said a male friend, spouse or partner influenced their decision to use narcotics. Only 2 percent of male addicts said a woman influenced their decision.*²

◆ *Female addicts are more likely to be depressed.*³ *A survey of cocaine users found women twice as likely (37 percent vs. 14 percent) to say cocaine made them feel confident and sociable. Men were more likely (42 percent vs. 26 percent) to cite physical energy and the sense of a controlled high as cocaine's appeal.*⁴

◆ *Female drug abusers and addicts are four times more likely to have suffered sexual assaults than women without drug problems.*⁵

¹ Lex, B. W. (1995). Alcohol and other psychoactive substance dependence in women and men. In M. V. Seeman (Ed.), Gender and psychopathology (pp. 311-358). Washington, DC: American Psychiatric Press. ² Hser, Y. et al. (1987). Sex differences in addict careers: Initiation of use. American Journal of Drug and Alcohol Abuse, 13(1), 33-57. ³ Lex, B. W. (1995). ⁴ Erickson, P. G. et al. (1989). Sex differences in cocaine use and experiences. American Journal of Drug and Alcohol Abuse, 15(2), 135-152. ⁵ Winfield, I. et al. (1990). Sexual assault and psychiatric disorders among a community sample of women. American Journal of Psychiatry, 147(3), 335-341.

Chart 11


Adult Women and Men Who Have Ever Used Illicit Drugs* by Age

Age	Women	Men
19-30	53.5%	59.2%
31-40	54.3	69.2
41-59	26.9	39.9
60+	4.1	10.9
All	33.9%	45.6%

* Cocaine (including crack), heroin, hallucinogens (including LSD), marijuana and hashish, and the non-medical use of sedatives, tranquilizers, stimulants, analgesics and inhalants.
Source: 1993 National Household Survey on Drug Abuse

Chart 12

Adult Women Who Use Illicit Drugs by Household Income



BEST COPY AVAILABLE

* Regular users have used an illicit drug at least monthly in the past year.
Source: 1993 National Household Survey on Drug Abuse

rates of regular drug use are similar among the three groups (NS) (CHART 13). Again, black women are more likely to be clustered at the extremes of abstaining or becoming regular users.⁶⁴

The crack epidemic has been particularly devastating to some African American communities, snaring women and men alike and destroying families.⁶⁵ Black women are more than twice as likely as white women to be regular cocaine users (S) (CHART 14) and are 14 times likelier to use crack cocaine during pregnancy.⁶⁶ In terms of total numbers, white women who use drugs regularly (2.2 million) outnumber all those who are black, Hispanic or from other minority groups (846,000).

Women progress more quickly than men from trying a drug to getting hooked.⁶⁷ Female cocaine addicts are also twice as likely as male addicts to be unemployed, diminishing their access to treatment.⁶⁸ Thus, the female addict is more likely than her male counterpart to be sicker, and to lack the resources to get treatment and to support herself once she is drug-free.

Prescription Drugs

Despite the conventional view that women misuse prescription drugs to relieve anxiety, concrete data on the extent of the problem are hard to find.⁶⁹

What evidence does exist indicates that women and men are equally likely to misuse four types of prescription drugs: stimulants, tranquilizers, sedatives and analgesics--

* Misuse of prescription drugs, also called "non-medical use," is use beyond or without a doctor's recommendation.

psychoactive drugs commonly prescribed for depression, anxiety, weight loss, insomnia and pain (CHART 15). Misuse is most common among young, white, low-income adults who are unemployed or working part-time. Contrary to the stereotype, among women with household incomes of \$30,000 or more, homemakers are less likely to misuse prescription drugs than women who work full-time outside the home (NS).⁷⁰


Non-medical use is only a small slice of the prescription drug problem. Women may suffer unintended or harmful consequences of chronic or dependent use of prescription drugs (even legally obtained), such as fainting, stupor or confusion, but not consider it "misuse" or "abuse." Women may also use drugs according to a doctor's advice, but combine them with alcohol, which may accelerate the onset of dependence and amplify the effects of the drug, leading to impaired motor skills and accidents.⁷¹ Female alcoholics often combine their alcohol with prescription drugs like tranquilizers and sedatives.⁷²

Women receive two-thirds of all prescriptions for tranquilizers and anti-depressants.⁷³ This is largely because women account for almost two-thirds of all visits to doctors, and because they are almost twice as likely as men to suffer depression.⁷⁴ One of eight women suffers depression each year, compared to one of 13 men.⁷⁵ Elderly women are especially vulnerable; 4 million of those over age 65 are depressed, compared to 2 million men.⁷⁶

Because female alcoholics are more likely than male alcoholics to suffer depression in addition to their addiction, doctors may medicate a woman's depression but miss the signs of alcohol abuse, and they may contribute to such abuse by adding to the array of drugs at her


Chart 13

Adult Women Who Use Illicit Drugs by Race and Ethnicity



* Regular users have used an illicit drug at least monthly in the past year. **BEST COPY AVAILABLE**
Source: 1993 National Household Survey on Drug Abuse

Chart 14 Drug-Using Adult Women Who Regularly Use Drugs



70

71

* Regular users have used the drug at least monthly in the past year.
Source: 1993 National Household Survey on Drug Abuse

BEST COPY AVAILABLE

Chart 15
Adult Women and Men Who
Use Prescription Drugs Non-Medically*
by Age

Age	Women	Men
19-30	8.1%	9.9%
31-40	4.6	5.6
41-59	3.2	1.7
60+	1.7	1.2
All	3.7%	3.9%

72

73

* Use of stimulants, tranquilizers, sedatives and analgesics at least once in the past year.
Source: 1993 National Household Survey on Drug Abuse

disposal.⁷⁷ Education about this danger and training to detect the signs of substance abuse in women would help doctors avoid this costly mistake.

Doctors can overprescribe drugs to women if they too casually attribute women's medical complaints to emotional causes.⁷⁸ Doctors have been found to attribute headaches to mental and nervous conditions in women more than in men, and to decide that women who complain of digestive, reproductive, nervous and mental symptoms are not really sick, compared with men who make similar complaints.⁷⁹ Even after controlling for patient symptoms and physician diagnosis, women have been found 37 percent more likely to get a prescription for an anti-anxiety drug and 82 percent more likely to get an anti-depressant.⁸⁰

The risk of inappropriate prescriptions or uses of prescription drugs is especially high among elderly women.⁸¹ Two-thirds of all long-term users of tranquilizers are women and a third of them are over age 64.⁸² One-fifth of nursing home residents--75 percent of whom are women--receive anti-psychotic drugs, usually to ease behavioral problems related to dementia, often when non-pharmaceutical actions would suffice.⁸³

Alcohol abuse compounds the danger of prescription drug use. Many experts believe that alcohol abuse by the elderly is widely under-reported.⁸⁴ At Johns Hopkins Hospital in Baltimore, 21 percent of hospital inpatients age 60 and older met criteria for alcohol dependence. Yet doctors had diagnosed only 37 percent of these patients--and they failed to diagnose a single case of alcoholism in an elderly woman.⁸⁵

* Long-term users have taken the drug for 12 months or more.

Adolescence: Where It All Begins

Unlike adult women and men, girls and boys are often indistinguishable in their rates of alcohol and drug use. Since 1975, adolescent girls have been equally or more likely than boys to smoke.⁸⁶ Today girls and boys are just as likely to smoke (12.0 percent vs. 11.9 percent), to quit (26.2 percent and 26.0 percent), or to avoid smoking (61.8 percent vs. 62.1 percent).⁸⁷

Adolescent girls and boys are also equally likely to drink (18.5 percent vs. 17.8 percent).⁸⁸ Though girls are still less likely than boys to binge drink, the gap has narrowed since 1975.⁸⁹ Most of the remaining difference stems from less beer drinking by females. Girls are less likely than boys to binge on beer (16 percent vs. 36 percent) and hard liquor (13 percent vs. 22 percent), but equally likely to binge on wine (5 percent). Girls are more likely to binge on wine coolers (10 percent vs. 7

Patterns of drug and alcohol abuse differ among younger and older women in ways that may signal important cultural changes. A study of 572 women in treatment for alcohol and/or drug addiction found:¹

◆ *Women under age 35 are more likely to use an array of drugs, while those age 35 and over commonly use only one or two, such as alcohol and painkillers. On a weekly basis, younger women are nine times likelier to use marijuana, six times likelier to use stimulants and five times likelier to use cocaine.*

◆ *Younger women are four times likelier to have begun drinking before age 16, and 15 times likelier to have tried illicit drugs by age 15.*

◆ *79 percent of older women usually or always use alcohol or drugs by themselves, compared to less than half (47 percent) of younger women.*

¹ Harrison, P. A. (1989). Women in treatment: Changing over time. *International Journal of the Addictions*, 24(7), 655-673.

* NHSDA data on adolescents, collected in homes with a parent present, understate the overall problem. For example, the 1990 Youth Risk Behavior Survey, conducted in schools, found that among students in grades 9 through 12, 61 percent of boys and 55 percent of girls drink.

percent). The degree of similarity varies by race and ethnicity.⁹⁰ Hispanic girls are less likely than Hispanic boys to drink at all (14.7 percent vs. 22.2 percent) (S), though Mexican and Cuban American girls are catching up.⁹¹

The gap between girl's and boy's illicit drug use has closed in the last 30 years, but boys are still more likely to become regular users.⁹² While men ages 31 to 59 are 1.5 times more likely than women in that age group to have used illicit drugs other than marijuana (S), in the 19 to 30 age group, the rate is only 1.3 times higher for men, and the rates are equal (12.7 percent vs. 12.6 percent) for adolescent girls and boys (S) (CHART 16).

The age at which tobacco, alcohol and illicit drug use begins has declined significantly in the last 30 years.⁹³ For example, among adult women and men who have used marijuana, those who are 45 years old or above started smoking marijuana on average during their middle or late 20s, while those 19 to 44 years old began at about age 17 (CHART 17). On average, adolescents now start to smoke at age 13 and to drink at age 15. By age 12, 15 percent of all girls and boys have tried cigarettes; by age 14, the percentage has doubled to 30 percent, and by age 18 it has doubled again to almost 60 percent.


Experimentation at such young ages is a concern for all adolescents, but it is most ominous for girls.⁹⁴ In older generations, women were more likely to try their first cigarette, drink or illicit drug at a later age than men. This is one reason women were less likely to become regular users;⁹⁵ they started when they were more mature and less vulnerable to the temptation and pressure to smoke and get high or drunk. But over the last

30 years, this protective factor has vanished. Girls and boys are now wading into drugs, alcohol and tobacco at the same early ages.

Chart 16

The Declining Gender Gap in Rates of Illicit Drug Use

Ratio of Male to Female Rates of Use of Illicit Drugs Other Than Marijuana*




* Those who have ever used illicit drugs other than marijuana.
Source: 1993 National Household Survey on Drug Abuse

BEST COPY AVAILABLE

Age at Which Women and Men First Used Marijuana

Chart 17



80

Current Age of Adults 81

BEST COPY AVAILABLE

Source: 1993 National Household Survey on Drug Abuse

III.

Damage and Death: The Truth and Consequences for Women

Women who smoke, abuse alcohol or use illegal drugs suffer the same consequences as men, but for women the harsh truths of addiction, illness, violence and death come more quickly, and the social stigma is more severe.

Alcohol

Illness and Death

Women feel the physical effects of alcohol, become addicted and develop alcohol-related illnesses, such as liver damage, hypertension, anemia, malnutrition, peptic ulcer, brain and heart damage, more rapidly than men.¹ Women who drink are more likely than men to develop liver cirrhosis, to develop it at an earlier stage of problem drinking and at lower levels of alcohol consumption.² Female alcoholics are up to twice as likely to die as male alcoholics in the same age group (who in turn die at rates three times above the general population), and a greater percentage of alcoholic women than men die from alcohol-related accidents, violence and suicide.³

A follow-up of 100 women 11 years after being hospitalized for alcoholism found that a third of them had died, primarily of alcohol-related causes such as pancreatitis, hepatic

cirrhosis and other liver disorders, violence and accidents.⁴ On average, their lives were 15 years shorter than women in the general population.

More than 9,000 women die each year from liver cirrhosis, accounting for one third of such deaths; alcohol is the cause or a major contributor in virtually all cases.⁵ According to the NIAAA, the risk of liver cirrhosis becomes significant for men who drink more than six drinks per day (80 grams of pure alcohol), but that risk becomes significant for women at less than two drinks per day (20 grams of alcohol).⁶ This phenomenon, known as "telescoping," may be due in part to the body weight and metabolic differences that, on average, cause women who drink roughly half of what men drink to have the same amount of alcohol in their blood.⁷ That's why one expert concluded, "It is now clear that an alcohol intake that may be considered moderate and innocuous in men is not necessarily so in women."⁸

Although the rate of heavy drinking is about the same for black and white women, cirrhosis mortality is 73 percent higher among black females than white females (8.3 vs. 4.8 per 100,000 individuals), suggesting that black women are more susceptible to alcohol's deadly consequences.⁹ Black alcoholic women suffer death rates that are nearly double those of white women.¹⁰ In the general population, black women's mortality from alcohol-induced causes in 1990 was 7.7 per 100,000, compared to 2.8 for white women.¹¹

A possible reason for the higher alcohol-related death rate among black women is that among women who frequently drink heavily, black women may have more drinks per month than do white women.¹² The proportion of heavy drinkers among black women peaks at a

* In this study, frequent heavy drinkers have five or more drinks at a sitting at least once a week.

later age (45-59) than among white women (25-44), which could indicate that black women are sustaining their heavy drinking for more years of their life.¹³ The higher concentration of poverty among African Americans is also a contributing factor.

For all alcoholics, drinking affects the brain in ways that impair learning, memory, abstract thinking, problem-solving, perceptual-motor skills (such as eye-hand coordination) and ability to analyze spatial relationships.¹⁴ But women suffer these impairments after fewer years of drinking.¹⁵

Some studies have found that women who have two or more drinks per day are 1.3 to 2.0 times more likely to get breast cancer than abstainers.¹⁶ But these studies rely on statistical correlations; research has not yet established a causal link.¹⁷

Heavy drinking can also impair a woman's endocrine system and lead to menstrual irregularities, amenorrhea (the absence of menstrual cycles), infertility and early menopause.¹⁸ This underlines the importance of preventing alcohol abuse among teenage girls, whose bodies are just developing.¹⁹

Moderate drinking appears to protect some women against heart disease, possibly stemming from an increase in estrogen that such drinking may cause.²⁰ Caution is important, however, in drawing conclusions about the benefits of drinking by women. While light to moderate drinking by men reduces mortality due to heart disease, the increased risk of liver cirrhosis (and possibly breast cancer) among women complicates the equation for them.²¹ In a study of 85,709 women, alcohol consumption appeared to lower mortality among women who drank one to 16 drinks per week.²² Above that, liver cirrhosis and breast cancer increased mortality rates. Moreover, any protective effect was limited to

women over age 50 and those at high risk of heart disease due to hypertension, diabetes, smoking, high cholesterol or a family history of heart attacks.

Women who drink less than men and do not consider themselves heavy drinkers may nevertheless experience a variety of health and social consequences.²³ In a study of binge drinking among college students, women were 30 percent more likely than men to experience a hangover, but they were less likely to describe themselves as heavy drinkers (8 percent vs. 22 percent).²⁴ And even though some women drank less than men, they were just as likely as men to experience alcohol-related problems such as missing classes, falling behind in their studies and feeling regret about drinking.²⁵ They were almost as likely to get injured while drinking, to have unplanned sex and to have it without a condom.²⁶

Sex

Among teenage girls, alcohol use is strongly associated with unprotected sex.²⁷ Adolescent girls who drink more than five times a month are five times likelier to be sexually active and a third less likely to have sex with a condom than girls who do not drink (S) (CHARTS 18 and 19). In contrast, adolescent boys' drinking appears to have little relationship to whether they use condoms (S). The link between female drinking and unprotected sex is more pronounced with first-time encounters.²⁸ Two-thirds of women who drank at the time of their first sexual intercourse said they did not use contraception, while nearly half of those who did not drink used contraception.²⁹

* Female binge drinkers consumed four or more drinks in a row at least once during the past two weeks. Male binge drinkers had five or more drinks in a row at least once during the past two weeks.


Chart 18

Sexual Activity Among Adolescents Ages 12-19

Used Alcohol:

Never
 Ever
 >5 days past month

Percentage Who Reported
Having Sexual
Intercourse



Female

Male


87

Source: 1993 Youth Risk Behavior Survey

BEST COPY AVAILABLE

Condom Use Among Adolescents Ages 12-19

Chart 19



Source: 1993 Youth Risk Behavior Survey

BEST COPY AVAILABLE

This connection is perilous for females, given that they may get pregnant and have a greater chance of contracting the AIDS virus, which appears to be more easily transmitted from a man to a woman during heterosexual contact than from a woman to a man.³⁰ They may also catch other sexually transmitted diseases (STDs) such as syphilis, genital herpes, chlamydia, gonorrhea, human papillomavirus and chancroid.³¹ Females 15 to 19 years old have the highest rates of gonorrhea among women.³² The potential consequences of STDs include pelvic infections, tubal infertility, and genital and cervical cancer--and if pregnancy occurs, ectopic pregnancy or neonatal transmission of STDs to the fetus, which can cause perinatal illness and even death.³³

Whatever the combination of factors--the disinhibiting effect of alcohol, a romantic vision, the excitement of drinking and risky sex--the connection between alcohol and unsafe sex is so compelling that a successful effort to prevent alcohol abuse seems sure to reduce high-risk sexual behavior as well.³⁴

The belief that alcohol is a disinhibitor or aphrodisiac has roots as far back as ancient Greece and Rome.³⁵ Yet while both genders tend to believe that alcohol facilitates sexual enjoyment, studies find that alcohol depresses physical sexual response.³⁶ Although women and men alike say that alcohol reduces tension in social situations, women often experience the opposite effect: their anxiety may increase when they drink and interact with men, possibly because they worry about the stigma attached to a woman who drinks.³⁷

This gap between women's expectations regarding alcohol and sex and the physical reality can have negative consequences. Many alcoholic women in treatment report problems of sexual dysfunction, which usually preceded the onset of their heavy drinking.³⁸ They

may have used alcohol to "treat" the sexual problem--only to find out that it failed and probably aggravated the problem. Thus, women who expect alcohol to ease problems of sexual dysfunction may get into a self-reinforcing cycle in which they medicate the problem with alcohol and, finding that the problem has only grown worse, drink more.

Even when alcohol does disinhibit a woman, it is unclear how often this translates into promiscuity.³⁹ In one survey, 60 percent of women said that drinking made them feel less inhibited, but only 22 percent said that drinking made them become "sexually forward," and even fewer (8 percent) said they became less particular in choosing a partner.⁴⁰ Yet 60 percent said that when they had been

A young woman told of how she agreed to go back to her date's house after a party. "We played quarter bounce [a drinking game]. I got sick drunk; I was slumped over a toilet vomiting. He grabbed me and dragged me into his room and raped me. I had been a virgin and felt it was all my fault for going back to his house when no one else was at home." Describing a separate incident, a young man said, "Alcohol loosened us up and the situation occurred by accident. If no alcohol was consumed, I would never have crossed that line."⁴¹

⁴¹ Watson, R. R. (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press.

drinking, someone *else* who had also been drinking had been sexually aggressive towards them.⁴¹ These and other studies suggest that more than creating a sexually aggressive woman, a woman's drinking may encourage men's sexual assertiveness towards her.⁴² Men tend to believe that a woman who is drinking is more sexually available than a woman who is not, and to view alcohol as a surefire social and sexual facilitator--that "candy is dandy, but liquor is quicker."⁴³

Violence

The consequences become brutal when assertiveness turns into sexual assault or other violence. The link between sexual assault, rape, marital abuse and other violence towards women when either the woman or man drinks is well-documented.⁴⁴ In up to three-quarters of all rapes and up to 70 percent of all incidents of domestic violence, either the victim, the perpetrator or both have been drinking.⁴⁵ An extensive national survey found that 74 percent of college students who had raped a woman and 55 percent of rape victims had been drinking prior to the incident.⁴⁶

Alcohol may make violence more likely because a woman who drinks heavily may seem sexually available and more vulnerable to attack, and/or the man may think intoxication will provide an excuse for his aggression.⁴⁷ Both college women and

men were found to consider a man who beats his wife less culpable if he'd been drinking-- but they held a woman who had been beaten more responsible if she had been drinking.⁴⁸ Women and men in college also attribute less blame to a man who raped if he'd been drinking and more responsibility to a women for being raped if she'd been drinking.⁴⁹

The link between women's drinking and spousal abuse may also stem from the fact that women who abuse alcohol often have male partners who drink heavily. Male drinking is a powerful predictor of wife abuse.⁵⁰ In 44 percent of marital assaults involving alcohol,

A study of spousal abuse among alcoholic and non-alcoholic women found alcoholic women nine times more likely to be slapped by their husband, five times more likely to be kicked or hit, five times more likely to be beaten and four times more likely to have their lives threatened.¹

¹ Miller, B. A. et al. (1989). Spousal violence among alcoholic women as compared to a random household sample of women. Journal of Studies on Alcohol, 50(6), 533-540.

both spouses had been found to be drinking; in another 44 percent, only the violent spouse had been drinking; and in the remaining 13 percent, only the victim had been drinking.⁵¹ Women often drink to cope with the trauma of being beaten, setting up a spiral of assault and alcohol abuse.⁵²

Personal and Social Problems

Females are much more likely than males to report internal and personal correlates of alcohol abuse, such as depression, feeling isolated, anxious or irritable, and conflicts in marital or family relationships.⁵³ Males are likelier to cite outer-directed problems, such as trouble at work, financial difficulties and arrests for drunk driving.⁵⁴ Among adolescents in treatment for alcoholism, girls most often report histories of depression and suicide attempts, while boys commonly report being arrested or suspended from school.⁵⁵

These differences in the way men and women see the consequences of their alcohol abuse have significant implications for prevention and treatment efforts. Not only do women have different issues to address in treatment, but their abuse or addiction may be harder to nip in the bud or recognize, given the lower visibility of its symptoms. Moreover,

"I married an alcoholic. I married him because I thought he was the only guy who would ever show me any attention. In those seven years, I put up with mental abuse, physical abuse and somewhat sexual abuse. The way that alcohol and marijuana played a role in my marriage—that was the only way of getting along with him. He was mean as a snake. He run me down onto the ground and start beating me. To relax, I drank, I smoked pot and go on into my own dream world. That's what the marijuana did, was put you in another dream state, where you would forget about all the pain and all the anger and all the abuse and get high. Until the next morning, and then it started all over again."¹

¹ Smith, P. H. et al. (1995). Battered substance-abusing women. Raleigh, NC: North Carolina Governor's Institute for Alcohol and Substance Abuse.

researchers who attempt to gauge the extent of alcohol problems among women and men--and doctors who seek to detect them--may miss a substantial number of female cases if they focus solely on external manifestations such as getting arrested or job difficulties.⁵⁶

Suicide. Suicide attempts are another deadly problem common among women who abuse alcohol.⁵⁷ Alcoholic women are almost five times more likely to attempt suicide than women of similar age and income who are not alcoholics (40 percent vs. 8.8 percent).⁵⁸ The dance of death between alcohol and suicide begins in adolescence, often before parents or other adults sense any danger.⁵⁹ Teenage girls who drink more than five times a month are almost six times likelier to attempt suicide than those who never drink (25.9 percent vs. 4.5 percent) (S) (CHART 20). Among adolescents in treatment for alcoholism and/or drug addiction, 30 percent of the girls report attempting suicide, compared to 15 percent of the boys; most girls (57 percent) had at least thought about suicide, while most boys (66 percent) had not.⁶⁰

The typical teenage suicide attempt is by a girl who tries to overdose on drugs.⁶¹ Boys are more likely to commit suicide, but the gap has been closing since 1980 as girls have used more lethal methods such as guns.⁶² Among adult alcoholics, the suicide rate for women already equals that for men.⁶³

An important factor in the link between suicide attempts and alcohol use among adolescent girls may be a history of sexual abuse. Some 35 percent of teenage girls in treatment for alcohol and drug abuse have been found to be sexually abused. Those who had been abused by a family member were 1.5 times more likely than those who had not to feel ashamed of themselves, more than twice as likely to express self-hate and almost three times as likely to attempt suicide.¹

¹ Edwall, G. E. et al. (1989). Psychological correlates of sexual abuse in adolescent girls in chemical dependency treatment. *Adolescence*, 24(94), 279-288.

The Wrong Way

Gender differences in substance abuse-related problems are diminishing.⁶⁴ As female drinking becomes a more public event among younger women, drunk driving, problems with legal authorities and vulnerability to assault have risen as well.⁶⁵ In a comparison of female alcoholics ages 20 to 29 and those ages 40 to 49, the younger women are three times likelier to report getting drunk in public more than once (72 percent vs. 23 percent) and driving while drunk (68 percent vs. 22 percent), and five times likelier to experience blackouts (61 percent vs. 12 percent).⁶⁶ Among women in treatment for alcohol or drug abuse, those under age 35 are more likely than those age 35 and older to report problems at work, such as troubles with co-workers or supervisors, mistakes, absenteeism, tardiness or injury.⁶⁷

While deaths among drunk drivers declined from 1982 to 1990, they dropped much less among women (4 percent) than men (15 percent).⁶⁸ Worse, drunk driving appears to be increasing among young women.⁶⁹ In North Carolina, the rate of alcohol-related crashes soared 74 percent between 1976 and 1985 among female drivers ages 18 to 20 (compared to a 27 percent drop for males), and almost doubled (up 93 percent) among those ages 21 to 24 (compared to a 7 percent drop for males).⁷⁰ In New York state, from 1980 to 1988, the proportion of adults convicted for drunk driving who were female rose from 7.2 percent to 11.6 percent; the number more than doubled from 3,124 to 7,050.⁷¹ The drunk driving recidivism rate of women also rose, almost equaling that of men.


Chart 20

Suicide Attempts by Adolescents Ages 12-19

Used Alcohol:

Percentage Who Reported
Having Attempted
Suicide

□ Never □ Ever >5 days past month



Tobacco

Because smoking by women was not widespread until World War II, only recently has the full spectrum of illness and death wrought by tobacco emerged. In 1980, the U.S. Surgeon General pointed out, "The first signs of an epidemic of smoking-related disease among women are now appearing," and an "epidemic of chronic obstructive lung disease among women has also begun."⁷² The leap of lung cancer beyond breast cancer as the leading cause of cancer death among women in 1986 was a grim milestone in this morbid trend.⁷³

Today, half the women who smoke (47.4 percent) will die from tobacco-related causes; in the early 1960s, only one out of six (16.7 percent) were expected to die from such causes.⁷⁴ Now at least 140,000 women die each year from illnesses

Women who live with a smoking spouse have a 30 percent higher risk of lung cancer than women who are not chronically exposed to passive smoke.¹

¹ Fontham, E. T. H. et al. (1994). Environmental tobacco smoke and lung cancer in nonsmoking women: A multicenter study. *JAMA*, 271(22), 1752-1759.

attributable to smoking cigarettes: 62,000 from heart disease, 36,000 from lung cancer, 25,000 from chronic lung disease, 10,000 from pancreatic, oral, esophageal, laryngeal, urinary and cervical cancers, and 8,000 from lower respiratory infections.⁷⁵ Since 1983, heart disease has killed more women than men each year (478,179 vs. 447,900 in 1991); smoking accounts for 13 percent of these deaths due to heart disease, the leading killer of women.⁷⁶

Mortality from tobacco-related illnesses is higher among black women than white women.⁷⁷ No one is sure why, but explanations include: indications that black women

retain nicotine in their bodies longer than do white women;⁷⁸ preference among many black women for menthol cigarettes, which have a higher tar and nicotine content than other cigarettes and are easier to inhale deeply;⁷⁹ and higher rates of poverty.⁸⁰

Women who smoke 20 or more cigarettes a day are up to four times likelier to get heart disease than non-smoking women.⁸¹ Women who smoke 25 or more cigarettes a day are 5.4 times likelier to get fatal heart disease.⁸² Women who smoke and use oral contraceptives increase their risk of a heart attack by 1,000 percent.⁸³

Smoking can sabotage an elderly woman's ability to live independently. It raises a women's risk of osteoporosis, apparently by reducing her estrogen levels.⁸⁴ Women who smoke a pack a day throughout adulthood cut their bone density up to 10 percent by the time they reach menopause.⁸⁵ Because of osteoporosis, even minor falls can result in hip, forearm and vertebrae fractures.⁸⁶ Hip fractures, which strike some 200,000 women a year, push up a women's mortality by 20 percent during the first year after the accident; most of those

At age 16, Trudy Grover moved out of her parents house. "I loved the feeling of taking care of myself. I had my own apartment, worked as a waitress and always had a pack of Marlboros in my pocket. One day when I was 35, I was running back and forth between tables and suddenly couldn't catch my breath. It was like somebody was holding me underwater. I told myself I was just getting older. At age 42, I was diagnosed with emphysema. The doctor put me on an oxygen tank, an ugly thing with green rubber tubes running into my nose. In the struggle to breathe, you can lose control of your bladder. I cried in shame. In 1993, I had a lung transplant—my only option, said the doctor. Now, a year and a half later, I take 23 pills a day. To pay bills, I was forced to sell my home and possessions. And I made the saddest journey of my life: back to my mother's house. 'Mom,' I said, averting my eyes, 'I'm broke. Will you take me in?' I was begging my 73-year-old mother to take care of me again. I couldn't help crying."

¹ Ecenbarger, W. So you think you want to smoke: The stories of four who did. *Reader's Digest*, 9/94, pp. 61-7.

who survive are permanently disabled.⁸⁷ Smoking has also been linked to weak muscles, poor balance and impaired neuromuscular function among women over age 65; one study found the decrease in muscle function comparable to five years of aging.⁸⁸

Smoking may also cause infertility and early menopause.⁸⁹ Women who smoke are 1.3 to 3.4 times likelier to suffer infertility, and women ages 44 to 54 who smoke have early menopause at about twice the rate of those who never smoked.⁹⁰ Early menopause, in turn, has been linked to a higher risk of osteoporosis-related fractures, heart disease and cancers of the reproductive system.⁹¹ A woman who smokes raises her risk of cervical cancer at least 50 percent and can double her risk of having ovarian cysts.⁹² Smoking may also increase the severity and effects of hypothyroidism, a hormonal disorder.⁹³

Although most smokers want to quit, only 2.5 percent succeed each year.⁹⁴ Of women who smoke daily, 42 percent tried to quit at least once in the previous year--and failed.⁹⁵ It is little surprise that fear of weight gain is a significant reason for failure since so many American women chronically worry about their weight. On any given day, four out of 10 women in the U.S. are trying to lose weight.⁹⁶ A third of female smokers are overweight.⁹⁷

*Susan started smoking at age 22. It seemed the thing to do. Now 42 and a college journalism professor, she can't kick the habit, despite a chronic asthma condition that requires daily medication--and once landed her in the hospital. "I maintain this fantasy," she said in the gravelly voice of a 20-year smoker, "that if I really wanted to quit, I could. I actually have a conversation with myself in which I say, 'Okay, I have to stop smoking and lose weight, so which should I do first? Quit eating or quit smoking? Clearly, I'd be better off if I didn't smoke, but I'm afraid I won't be able to lose weight then.'"*⁹¹

⁹¹ Morain, C. Still a long way to go, baby...
American Medical News, 7/4/94.

Female smokers who think the health risks of gaining weight cancel out the benefits of quitting smoking are misguided.⁹⁸ Over a 10-year period, women who quit smoking gain an average of 11 pounds more than women who continue to smoke.⁹⁹ But, as one expert insists, "You would have to gain more than 100 pounds to equal the health risks of smoking two packs a day."¹⁰⁰

Many female smokers clearly care more about appearance than health.¹⁰¹ Yet, by reducing blood flow to the skin, smoking also increases facial wrinkling, particularly crow's feet around the eyes.¹⁰² In time, "smoker's face"--lines or wrinkles, gaunt facial features, grayish skin and a plethoric complexion--can be found among 46 percent of smokers.¹⁰³

Quitting produces results. Women who stop smoking cut their risk of stroke in half.¹⁰⁴ Within a year, their smoking-related risk of heart disease drops 50 percent.¹⁰⁵ After three years, the risk of a heart attack is no greater than for women who never smoked.¹⁰⁶ Within five years, their smoking-related risk of heart disease can disappear altogether.¹⁰⁷ The higher risk of cervical cancer also declines in former smokers.¹⁰⁸

Illicit Drugs

In both women and men, cocaine use can cause heart attacks, hypertension, strokes, pulmonary disorders, seizures, tremors, motor and visual problems, malnutrition and infections from intravenous injection sites.¹⁰⁹ AIDS, syphilis, tuberculosis, hepatitis B, pneumonia and suppressed immune function are also common ills among cocaine and other illicit drug users.¹¹⁰ Some consequences of illicit drug use, however, strike women with particular force.

Because women may develop dependence on drugs more quickly than men, they may experience the ill-health effects of drug use faster.¹¹¹ Female drug addicts are also at higher risk of contracting the AIDS virus than men, not only because of the easier transmission of the virus from a man to a woman, but because they are more likely to have a drug-using partner with whom they have unprotected sex or share dirty needles and to have unprotected sex with multiple partners in order to finance their addiction.¹¹²

Seventy percent of all AIDS cases among women stem from illicit drug use.¹¹³ Almost half are the result of injection-drug use; another quarter come from having sex with an injection-drug user.¹¹⁴ The 80,000 women with HIV in 1992 will leave behind some 150,000 children when they die.¹¹⁵ Of these children, some 38,000 will themselves have AIDS.¹¹⁶

Initially a disease that raged primarily among homosexual men, AIDS is now spreading most rapidly among drug users and their partners, and particularly women of color.¹¹⁷ Almost one out of five new AIDS cases (18 percent) in 1995 were women, compared to one out of 14 (7 percent) in 1985.¹¹⁸ In 1994, more than three-fourths (77 percent) of new AIDS cases among women were blacks (8,000) and Hispanics (2,800).¹¹⁹ AIDS is now the leading cause of death for black women ages 25 to 44 (22 percent of deaths) and the fifth leading cause of death for white women in this age group (6 percent of deaths).¹²⁰

Even if the female drug user escapes HIV infection, she is at high risk of catching other sexually transmitted diseases such as syphilis, genital herpes, gonorrhea, human papillomavirus and chancroid.¹²¹ The incidence of these diseases among women has risen

in tandem with the crack epidemic.¹²² In Connecticut, a 422 percent jump in syphilis cases among women from 1986 to 1988 was linked to cocaine use and prostitution.¹²³ Tuberculosis (TB) has also surged among drug users. From 1985 to 1992, the number of TB cases among women ages 15 to 44 rose 41 percent in the U.S.¹²⁴

Adolescents who use cocaine are 31 times more likely to be sexually active, 27 times more likely to have four or more partners and twice as likely to fail to use a condom than teens who do not use alcohol, tobacco or illicit drugs.¹


¹ Lowry, R. et al. (1994). Substance use and HIV-related sexual behaviors among U.S. high school students: Are they related? *American Journal of Public Health*, 84(7), 1116-1120.

Many non-drug using women put themselves at risk by having unprotected sex with men who may be drug-users. Among sexually active unmarried women, less than one in six always has her partner use a condom to avoid sexually transmitted diseases.¹²⁵ Women may be reluctant to suggest a man wear a condom for fear of signaling distrust or "ruining the romance."¹²⁶ Hispanic women may be particularly unwilling to ask for a condom for fear of suggesting they are savvy about the risks of sex, which a man might consider inappropriate.¹²⁷ African American women may not want to challenge men who might feel powerless in every domain of their life except their sexuality.¹²⁸ Some women fear violent reprisals.¹²⁹

Like alcohol, marijuana heightens the likelihood of unprotected sex.¹³⁰ Teenage girls who used marijuana at least three times in the past month are more than twice as likely to be sexually active and 25 percent less likely to use condoms than those who never use marijuana (S) (CHARTS 21 and 22).

Chart 21

Sexual Activity Among Adolescents Ages 12-19




Source: 1993 Youth Risk Behavior Survey

BEST COPY AVAILABLE

Condom Use Among Adolescents Ages 12-19

Chart 22



Illicit drugs may interfere with a woman's reproductive system, though most studies are limited by a failure to account for concurrent alcohol use. Heroin, marijuana and cocaine have each been linked to amenorrhea, anovulation and spontaneous abortion.¹³¹ Cocaine and heroin can depress sexual functioning as well: women are 66 percent more likely than men to say that cocaine negatively affected sexual experience (48 percent vs. 29 percent).¹³²

Weight

Because cocaine, heroin and methamphetamines can suppress appetite, some women may use them (or continue to use them) to lose weight.¹³³ One study found that in contrast to male cocaine-users, female users often reported skipping four or more regular meals in a row.¹³⁴ Malnourishment is a particular concern during adolescence, an important period of development, and pregnancy, when a woman's nutritional needs rise dramatically to assure healthy fetal growth.¹³⁵

According to staff at the Farm Place rehabilitation center in Surrey, England, many fashion models are addicted to drugs. While caffeine, nicotine and laxatives are among the substances frequently abused, they are not the most dangerous. To keep their weight down, models often take amphetamines or other stimulants—such as cocaine—that increase their heart rate and decrease their appetite. Heroin also has become more popular as a means of weight control, because it can be snorted rather than injected.¹

¹ Hardy, R. Drugs, dieting and my fear of falling in love. Daily Mail, 4/18/95, p. 18; Gandel, D. Fashion inside out: It's an ad of a different color, perspective. Los Angeles Times, 7/7/94, p. E4; Blundy, A. Real lives: When looks can kill. The Independent, 8/20/95, p. 3.

Prostitution and Crime

While the link between drugs and prostitution is hardly new, trading sex for money to buy drugs became a common and deadly practice among crack-addicted women.¹³⁶ Among 84 crack-using women who were not in treatment, three out of four (73 percent) had been arrested for prostitution.¹³⁷ In New York City, as the price of cocaine dropped with the arrival of crack "rocks" at \$2 each, in some areas the charge for sex acts also fell from \$25 to \$2.¹³⁸

Crack houses have become not only the hovels of addicts, but centers of the sex trade, where women desperate for drugs sell their bodies to finance their addiction. Despite the physical abuse, violence, unintended pregnancies, AIDS and other sexually transmitted diseases that come with it, female addicts become dependent not only on the drug, but also on the crack house for what, in the web of addiction, appears to them to be their only way to survive.¹³⁹ One researcher concluded,

"The sex-for-crack phenomenon and the incredible degradation of women surrounding much of its routine enactment is like nothing ever seen in the annals of drug use, street life, prostitution or domestic wife-battering."¹⁴⁰

The declining price for sex acts and the violent environment of crack addiction has pushed some women into robbery,

For the Chambers brothers, millionaire entrepreneurs of crack, "unparalleled customer service" was their watchword. At one of their first all-night crack palaces, the smallest rocks were sold on the first floor, slightly larger rocks on the second, escalating to "\$20 boulders" on the fourth. Private rooms were available for smoking and drinking, and one apartment was furnished with a bed frame, box springs and mattress for "female customers who craved crack but had no money."¹

¹ Olsen, J. Brains and Industry. New York Times Book Review, 4/23/95, p. 6.

burglary and drug dealing.¹⁴¹ Although women are still less likely than men to commit violent crimes, female addicts commonly shoplift, pickpocket, sell stolen goods and forge checks.¹⁴²

The crack epidemic has fueled an explosion in the female prison population since 1980.¹⁴³ From 1984 to 1993, the number of women arrested for drug offenses jumped 82 percent, compared to 52 percent for men.¹⁴⁴ By 1991, a third of female state inmates (32.8 percent) were serving time for a drug offense, compared to one in eight (12 percent) just five years earlier.¹⁴⁵ While men outnumber women in state and federal prisons by a ratio of 16 to 1 (952,585 men vs. 59,878 women in 1994), the number of female inmates is growing faster than the number of male inmates.¹⁴⁶

Women who commit crimes are more likely than men to have used drugs regularly, to have used them daily in the month before the crime, to have committed the offense to raise money to buy drugs and to have been under the influence of drugs at the time of the crime.¹⁴⁷ Of those women under the influence, 10.1 percent were using crack; 10.8 percent, heroin.*

In 1988, of 2,280 inmates that the New York City Department of Corrections identified as addicted to crack, 56 percent were women.¹⁴⁸ In 1989, among women imprisoned for drug offenses in New York State, 78.6 percent cited crack (44.7 percent) or cocaine (33.9 percent) as the drug leading to their crime and conviction.¹⁴⁹ Nationwide, more than half the women arrested in 1994 tested positive for drugs; in some cities, including Cleveland, New York and St. Louis, more than three-quarters tested positive.¹⁵⁰

* These numbers cannot be added because some women were under the influence of more than one drug.

Two-thirds of female inmates are mothers of at least one child; between 6 and 15 percent are pregnant when they enter prison.¹⁵¹ A 1989 survey of jail inmates found that 64 percent of those who were mothers had used cocaine in the month before arrest; 45 percent of those who were fathers had done so.¹⁵² Fifty-eight percent of the mothers and 39 percent of the fathers reported recent heroin use.¹⁵³

Violence

While most research has focused on the link between violence and alcohol use, violence is also the deadly companion of illicit drug use--and it strikes women with magnum force. Most female drug addicts have been violently assaulted.¹⁵⁴ Nine out of 10 women (87 percent) in a New York City drug treatment program were found to be victims of violence.¹⁵⁵ Seventy percent of drug-addicted, low-income pregnant women in methadone maintenance have been beaten--86 percent of them by their husbands or partners.¹⁵⁶ Many partners are also drug-users, which raises the woman's risk of getting beaten.¹⁵⁷

Women who use cocaine are at high risk of being murdered.¹⁵⁸ In New York City from 1990 to 1991, black female homicide victims, ages 15 to 24, were twice as likely as black male victims of the same age to test positive for cocaine (35.6 percent vs. 18.9 percent).¹⁵⁹ In the 25 to 34 age group, 71.8 percent of black women (vs. 44.4 percent of black men) and 59.1 percent of white women (vs. 37.5 percent of white men) tested positive for cocaine.¹⁶⁰

Homicide, suicide, overdoses or other injuries that prove fatal often follow cocaine use for both women and men.¹⁶¹ In New York City from 1990 to 1992, fatal injuries after

cocaine use exceeded the number of deaths from AIDS, cancer or heart disease among black and Hispanic women ages 15 to 24.

Prescription Drugs

Alcohol and most prescription drugs don't mix. Any prescribed drug carries the risk of an adverse reaction if the woman is using other drugs or alcohol at the same time.¹⁶² Stupor or confusion from sedatives, lowered blood pressure from anti-psychotics, fainting following the use of antidepressants, sedatives or tranquilizers--all are the unintended and perilous results of a prescription drug used in combination with other drugs and/or alcohol.¹⁶³ Tranquilizers, such as Valium or Xanax, are particularly dangerous in combination with alcohol because both depress the central nervous system, leading to falls, hip fractures and car accidents that are potentially fatal.¹⁶⁴

In the 1970s, the replacement of barbiturates with the sedative drugs known as benzodiazepines (Valium, Librium, Xanax, Tranxene, Ativan, Halcion and others) greatly reduced the risk of accidental overdoses and mental impairment that can lead to debilitating and even deadly falls.¹⁶⁵ But benzodiazepines still carry a risk of unwanted drowsiness, confusion and ataxia, which can incapacitate the elderly and lead to falls and broken bones.¹⁶⁶

The consequences of prescription drug misuse affect elderly women more severely than men because these women use more psychoactive prescription drugs and doctors are less likely to recognize an alcohol problem in an elderly woman than in a man.¹⁶⁷ Use of sedating and antidepressant prescription drugs almost doubles the risk of falls and fractures

among the elderly.¹⁶⁸ In a prospective study of 336 people age 75 or older, 32 percent fell within a year, 24 percent with serious injuries and 6 percent with fractures.¹⁶⁹ Sedative use--more so even than dementia--was the most important factor in predicting who would fall.¹⁷⁰ Use of antidepressants increases the risk of hip fracture by 1.6 in adults ages 65 and older, after accounting for factors such as dementia and general functional status.¹⁷¹

Health and Welfare Costs

Women's substance abuse triggers substantial health care and welfare costs. In 1995, women's use of alcohol, tobacco and illicit drugs accounted for \$68 billion in health care costs, 12.3 percent of all health care spending for women.¹⁷² Men's substance use and abuse was responsible for \$72 billion in costs, 19.2 percent of all spending for men. Among women, 52 percent of the costs were due to cigarette smoking, 30 percent to alcohol and 18 percent to illicit drugs. For men, 58 percent of the costs were for cigarettes, 26 percent for alcohol and 16 percent for illicit drugs.

In addition, one million women--at least 20 percent of those receiving welfare benefits from the Aid to Families with Dependent Children program--are frequent binge drinkers or regular drug users.¹⁷³ These numbers may be low. A Maryland study found substance abuse a serious problem for 37 to 52 percent of the women in a welfare-to-work program; the prevalence of alcoholism and drug abuse ranged from 16 to 21 percent, while an

* In this analysis, frequent binge drinkers have 5 or more drinks at a sitting at least weekly; frequent drug users use hallucinogens at least monthly, cocaine or heroin at least weekly, or marijuana more than 2 days a week.

additional 21 to 31 percent of these women experienced problems related to substance use.¹⁷⁴ Without treatment and other help, these women are likely to be trapped on welfare.

IV.

Drugs, Alcohol and Cigarettes During Pregnancy:

A Lethal Combination for Mother and Child

Smoking, drinking alcohol and using drugs during pregnancy can have devastating consequences for the fetus and newborn child. Cigarette smoking accounts for up to 20 percent of all low birth weight in America, making tobacco the leading preventable cause of this frequent precursor to infant mortality.^{*1} Alcohol use is responsible for 10 percent of all mental retardation, making it the leading preventable cause of this birth defect.² Cocaine and heroin use during pregnancy may severely impair the newborn.

Nicotine and alcohol are the drugs most widely used during pregnancy. Of the 4 million women who get pregnant each year, some 820,000 (20.4 percent) smoke cigarettes and 757,000 (18.8 percent) drink alcohol during pregnancy.³ Roughly 500,000 (13 percent) use illicit drugs.^{**4} Of them, more than five of 10 smoke marijuana; six of 10 use an illicit drug other than marijuana; one-fifth snort or smoke cocaine; and one of four misuse prescription drugs, such as Librium and Valium (CHARTS 23 and 24).^{***5}

* Low birth weight babies weigh 5.5 pounds (2,500 grams) or less at birth. Such small babies are 40 times more likely to die in the first month of life and have a higher risk of retardation, blindness and learning problems. Disorders relating to prematurity and low birth weight were the third leading cause of infant death in 1991.

** National estimates range from 212,000 to 736,000 prenatally exposed newborns each year. See pg. 68.

*** These numbers cannot be added because most women use more than one substance.

Drug use during pregnancy prompted public concern when the crack cocaine epidemic swept cities nationwide in the 1980s.⁶ From 1979 to 1990, the percentage of pregnant women who used drugs, as reported on hospital discharge records, soared 576 percent.⁷ Thousands of newborns arrived in hospitals affected by crack cocaine and other drugs. Hundreds became "boarder" babies who stayed in the hospital for months, often for no medical reason, because their drug-using mothers and fathers were unable or unwilling to care for them or because child welfare investigations delayed discharge of the child.

Cities and states across the country are struggling to respond to this influx of drug-exposed babies. In New York City from 1980 to 1988, the recorded number of newborns whose mothers had used illicit drugs while pregnant jumped from one in 137 to one in 33, which in 1996 puts about one drug-exposed baby in every average lower-school classroom of 30 students.⁸ In California, drug-exposed births at three hospitals doubled from 1986 to 1989.⁹ Both the Los Angeles and Chicago school districts have developed a special preschool curriculum for babies exposed to drugs in the womb.¹⁰


Foster Care

The crack epidemic, as well as alcohol and other drug use, has sparked a surge in child abuse and neglect that has overwhelmed child welfare agencies.¹¹ From 1985 to 1994, reports of abuse and neglect nationwide rose 64 percent from 1.92 million to 3.14 million.¹² Drug and alcohol abuse is a significant factor in more than 75 percent of child

* Based on the National Hospital Discharge Survey, the rate of drug use among pregnant women rose from 10.1 per 10,000 (0.1%) in 1979 to 68.3 per 10,000 (0.7%) in 1990. While the magnitude of the rise provides evidence of an epidemic, the rates of drug use appear low. Hospital discharge records are widely believed to under-report drug use substantially.

Drug Use During Pregnancy

Of the 4 million women who give birth each year:



117

NOTE: These numbers cannot be added because most women use more than one substance.

Source: NIDA, 1994; Vega, 1993; Center for the Future of Children, 1991;


South Carolina State Council on Maternal, Infant & Child Health, 1991

118

BEST COPY AVAILABLE

Illicit Drug Use During Pregnancy

Of the 4 million women who give birth each year, roughly 500,000 use illicit drugs:



NOTE: These numbers cannot be added because most women use more than one substance.

* Non-medical use of psychotherapeutics (amphetamines, sedatives, tranquilizers or analgesics).

Source: NIDA, 1994; Vega, 1993; Center for the Future of Children, 1991;

South Carolina State Council on Maternal, Infant & Child Health, 1991

welfare cases in some cities, such as New York.¹³ With so many of these parents unable to raise their children, the number of children in foster care nationally jumped 53 percent from 280,000 in 1986 to 429,000 in 1991.¹⁴ Over the same period, the percentage of children in foster care age three or younger, who have been exposed prenatally to drugs, doubled from 29 to 62; the percentage of those with documented prenatal cocaine exposure more than tripled from 17 to 55.¹⁵ Critical shortages of foster care families have forced child welfare authorities to shuffle children among homes, separate siblings and place thousands of children in group homes--with little or no hope of being adopted.¹⁶

Juggling individual caseloads that have ballooned up to 160, case workers who investigate reports of abuse and neglect must make delicate judgment calls, often with little or no training in substance abuse and scarce resources or treatment slots at their disposal.¹⁷ Yet without swift action, reports of abuse can become certificates of death. In New York City, the deaths of 74 children in 1994 had been reported for suspected abuse or neglect. In 17 of the 25 cases (68 percent) investigated by the Child Fatality Review Panel, the mother had a history of substance abuse, and in 15 of these she was using drugs at the time the child died.¹⁸ Seven of the children who died had tested positive for drugs at birth.¹⁹

In a 1989 survey of 12 metropolitan areas nationwide, child welfare officials said that standards for foster parents have declined to meet demand created by the crack epidemic. A caseworker on the West Coast said that applicants with "marginal" qualifications who would have been turned away a few years ago are now accepted. Nevertheless, another caseworker said that there are not enough people willing to adopt "these kind of children."

¹ Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.

AIDS

AIDS is another deadly consequence of substance abuse for both mother and child. A quarter of infants born to women with HIV are themselves infected with the virus.²⁰ From 1985 to 1995, some 16,000 babies were born infected with HIV.²¹ These children contracted the virus from their mothers, most of whom were infected by sharing dirty needles to inject drugs or having sex with an HIV-positive injection drug user.²² Some mothers contracted the virus by trading sex for drugs--usually crack cocaine.²³

Though AIDS is the most serious threat to newborns, syphilis is another common scourge passed from pregnant drug users to their children.²⁴ As crack cocaine use surged during the 1980s, so did the number of newborns with syphilis. In 1991, 4,410 babies were born with syphilis, compared to 107 in 1980.²⁵ Women who have syphilis can be tested during pregnancy and treated to prevent transmission to the newborn, but many drug-addicted women get no prenatal care or get it too late.²⁶

Illicit Drug Use

No single survey has definitively determined how many women use illicit drugs during pregnancy. Studies vary in

On November 11, 1992, a baby girl was born at Kings County Hospital in Brooklyn with the matchstick arms and legs and underdeveloped lungs typical of babies born 11 weeks early. But this newborn also had an enlarged liver and spleen and peeling skin on the palms of her hands and soles of her feet, signs that she had contracted syphilis in her mother's womb. Most pregnant women who have syphilis are screened at their first prenatal visit, get early treatment and have no problem. But most of the babies with syphilis are born to mothers who get no prenatal care. Many of them are addicted to crack and contracted syphilis trading sex for drugs.¹

¹ Lewin, Tamar. Syphilis cases among newborns climb with the rise in crack use. New York Times, 11/23/92, p. A13.

their method for determining use (self-report, urine tests, tests of meconium (baby's first stool)), the demographics of the sample (race, income, residence (urban, suburban, rural)) and the hospital protocol for deciding whom to test. Some hospitals screen for substance use only when a woman who is giving birth appears to be high on drugs or if her newborn has problems that often stem from prenatal drug exposure. If hospital staff decide to ask about drug use, a woman can provide information on the quantity, frequency and timing of use throughout her pregnancy, but she may forget, withhold information or lie.²⁷ Urine tests are more reliable, but can detect cocaine for only up to 48 hours after use. Less commonly employed, meconium screening can detect cocaine and other drug use from a few weeks to several months after use.²⁸ Rarely employed, new technology to analyze hair can determine quantity and timing of some drug use throughout pregnancy.²⁹

This menu of methods produces varying estimates of the number of pregnancies that involve illicit drug use--from 5.3 percent to 23.9 percent (CHART 25). Vega *et al*'s study used urine tests from 29,494 women in California; applying the results nationally indicates a rate of illicit drug use during pregnancy of 5.3 percent (212,000 infants in 1995).³⁰ The National Health and Pregnancy Survey of the National Institute on Drug Abuse (NIDA), which asked women in face-to-face interviews about their drug use, found a rate of 5.5 percent (220,000 babies).³¹ But NIDA's 1994 National Household Survey on Drug Abuse, which also used face-to-face interviews, found that 9.4 percent of pregnant women had used an illicit drug during the past year (376,000).³² Chasnoff's 1989 survey of 36 urban

* CASA estimated the number of illicit drug-exposed infants by applying the percentages to the total number of live births in the U.S. in 1995 (approximately 4 million).

hospitals found a rate of 11.0 percent (440,000).³³ In those hospitals that tested for substance use only when they saw severe birth complications, average prevalence was 3.0 percent; but where they tested *every* woman, average prevalence was 15.7 percent.

Assuming that pregnant women use illicit drugs at the same rate as other women, Gomby and Shiono concluded that roughly 18.4 percent of pregnant women use illicit drugs (736,000 babies).³⁴ While this may overestimate use because some women abstain while pregnant, estimates that rely on more conservative measures, such as urine tests and self-reports, may be just the tip of the iceberg.³⁵ A state-wide study in South Carolina found that meconium and urine tests picked up twice as many drug users as urine tests alone (23.9 percent vs. 10.2 percent). Until meconium and other more reliable tests are widely employed, substance use by pregnant women could be the most missed diagnosis in obstetric and pediatric medicine.³⁶

Who Uses During Pregnancy?

Illicit drug, alcohol and tobacco use during pregnancy crosses racial, economic and educational lines, but the drug of choice varies depending on the woman's situation.

African American women are more than twice as likely as white women to use illicit drugs while pregnant (11.3 percent vs. 4.5 percent).³⁷ Prenatal cocaine use is far more common among black women than white women (4.5 percent vs. 0.4 percent). White women are more likely to drink alcohol (22.7 percent vs. 15.8 percent) and smoke cigarettes

* Gomby and Shiono estimate the rate of illegal drug use among pregnant women to fall between 15.8 and 21 percent; 18.4 percent is the mid-point.

Chart 25

Illicit Drug Use During Pregnancy

%	(Source)	1995*
5.3	(Vega, <i>et al</i> , 1992)	212,000
5.5	(NIDA, 1992-93)	220,000
9.4	(NHSDA, 1994)	376,000
11.0	(Chasnoff, 1989)	440,000
18.4	(Gomby & Shiono, 1990)	736,000
23.9	(South Carolina, 1990-91)	956,000

* CASA estimated the number of illicit drug-exposed infants by applying the percentages to the total number of live births in the U.S. in 1995 (approximately 4 million).

(24.4 percent vs. 19.8 percent) than black women during pregnancy. Some surveys have found white women more likely to use marijuana prenatally.³⁸

Unmarried pregnant women are nearly four times likelier to use illicit drugs than married pregnant women.³⁹ Cigarette and illicit drug use is more common among women with low levels of education.⁴⁰ Women with less than a high school degree are more than twice as likely to smoke cigarettes and more than three times as likely to use marijuana while pregnant as women who have been to college.⁴¹

Drinking rates during pregnancy tend to rise with age, education and income. Women age 20 and over are twice as likely to drink during pregnancy as those under age 20.⁴² Despite the Surgeon General's urging in 1981 that women abstain during pregnancy, women with 16 or more years of education and those with income of \$40,000 or more are likelier to drink during pregnancy.⁴³ However, women with incomes of \$10,000 or less are more likely to drink *frequently* while pregnant.⁴⁴

Survey results that have compared illicit drug use among publicly and privately insured women are inconsistent. Two statewide studies, as well as a random sample in Hillsborough County, Florida, found higher rates of illicit drug use among pregnant women who rely on public insurance or are uninsured.⁴⁵ But a study of all pregnant women receiving prenatal care in public health clinics or in selected private clinics in Florida's Pinellas County found illicit drug use among 16.3 percent of patients at public clinics and 13.1 percent of those at private doctors' offices.⁴⁶

* In this 1995 Centers for Disease Control and Prevention study, frequent drinkers had six or more drinks per week during pregnancy.

Damage Done to Children

Ideally, research on birth outcomes (such as rates of low birth weight, prematurity and birth defects) and child development would compare women who use a drug with non-drug using women who are otherwise similar in age, race, income, education and use of prenatal care. In practice, multiple drug use and factors such as poverty and inadequate prenatal care confound most assessments of the impact of using a particular drug.⁴⁷ Many, if not most, pregnant women who take drugs during pregnancy use more than one kind of drug, usually including alcohol or tobacco.⁴⁸ Tabloid and sound-bite journalism that spotlights "crack babies" is misleading because most such newborns have been exposed to at least one other illicit drug, as well as nicotine and alcohol.

Untangling the relative impact of each drug on a baby born to a woman who smoked cigarettes, drank alcohol and snorted cocaine while pregnant is extremely difficult. If she also lacked prenatal care and a nutritious diet, or was abused by her partner, the task of teasing out the impact of each of these factors is even more complex. If possible, research samples need to be large enough to separate and compare the effects of individual drugs, as well as drugs used in combination, and to control for demographic factors such as income, education, access to prenatal care and nutrition.

Due to formidable cost and practical obstacles, no study has met all these standards. Yet in recent years, researchers have begun to address urgent questions--although they sometimes come up with conflicting answers--about the damage to the fetus and babies exposed to tobacco, alcohol, and illicit and prescription drugs in the womb. The complexity of this issue requires much more research, but here is what we know.

Fetal Tobacco Syndrome

Each year, smoking during pregnancy causes up to 141,000 miscarriages, 61,000 low birth weight babies, 4,800 perinatal deaths (including stillborn infants and infants who die shortly after birth) and 2,200 infant deaths from Sudden Infant Death Syndrome (SIDS),* and may cause respiratory illness and delay a child's cognitive development.⁴⁹

An analysis of the 1991 National Center for Health Statistics Linked Birth/Infant Death Data Set reveals that smoking during pregnancy increases the risk that a child will die within the first year of life (CHART 26).^{**50} While the infant mortality rate among women who do not smoke during pregnancy is 8.0 per 1,000, it jumps to 12.2 per 1,000 among those who

A group of pregnant teenagers said controlling their weight gain was a chief reason they smoke.¹

◆ *"I smoked one pack a day before I got pregnant. I smoke more now since I'm pregnant. If I put on too much weight now, it would be hard to lose after delivery. I smoke now 'cause I have trouble with diets."*

◆ *"I know with smoking I will not have to diet postpartum. I smoke two packs a day now. Before I got pregnant I needed to lose 100 pounds. I can't think of having to lose more weight by dieting."*

◆ *"My friends poke fun at me and call me names 'cause I'm too fat with this pregnancy. I smoke to keep my weight down."*

◆ *"I won't get another boyfriend if I gain too much weight. I smoke so I'll be slim and boys will ask me out."*

¹ Lawson, E. J. (1994). The role of smoking in the lives of low-income pregnant adolescents: A field study. *Adolescence*, 29(113), 61-79.

* Lowinson, et al, (1992) define SIDS as the sudden and expected death of an infant between 1 week and 1 year of age, whose death remains unexplained after a complete autopsy examination, full history and death site investigation.

** Data do not account for use of alcohol and illicit drugs.

smoke (S). The risk of infant death is almost as high among lighter smokers (less than a pack a day) as among heavier smokers.

The damage tobacco inflicts on the fetus and newborn is so severe and widespread that experts talk of a "fetal tobacco syndrome" (FTS), and conclude, "The time has come to recognize cigarette smoking as the single most powerful determinant of poor fetal growth in the developed world."⁵¹

Smoking during pregnancy doubles the likelihood that a baby will be born underweight, after controlling for maternal alcohol and drug use, education and employment, and prenatal care.⁵² Even passive exposure of pregnant women to cigarette smoke can double the risk.⁵³ Blood levels of cotinine (a metabolite of nicotine) are higher in black pregnant women who smoke than in white women who do so--even after accounting for the number of cigarettes smoked, years of smoking and other factors. This suggests that African-American women may metabolize the chemicals in cigarettes less effectively, leading to more severe health consequences for them and their children.⁵⁴ Coupled with higher rates of poverty and lower rates of prenatal care, such biological factors help explain why African American smokers have rates of infant mortality and low birth weight twice those of whites.⁵⁵

The risk of SIDS is up to five times greater for infants born to women who smoke during the second trimester of pregnancy compared to those who don't smoke at all.⁵⁶ For children of parents who begin or resume smoking after birth, the risk of SIDS more than doubles compared to families without smokers. If the mother or father smokes in the same room as the infant, the risk of SIDS has been found to jump 4.6 to 8.5 times.⁵⁷ Passive

Chart 26

The Link Between Smoking During Pregnancy and Infant Death

	<u>Infant Mortality Rate (per 1,000)</u>
Non-Smoking Pregnant Women	8.0
All Smoking Pregnant Women	12.2
< 1 pack per day	12.0
≥ 1 pack per day	12.4

Source: Centers for Disease Control & Prevention, National Center for Health Statistics,
Linked Birth/Infant Death Data Set, 1991

exposure to tobacco smoke during pregnancy raises the risk of low birth weight; exposure after birth increases the baby's chances of suffering respiratory illness, middle-ear infections and asthma.⁵⁸

Smoking during pregnancy has also been linked to lower scores on intelligence tests.⁵⁹ Children prenatally exposed to half a pack or more of cigarettes per day have been found to have intelligence scores significantly lower at ages three and four than those with non-smoking mothers, even after adjusting for the mothers' education level.⁶⁰ But the cognitive consequences of tobacco exposure can dissipate over time. One study found a link between smoking during pregnancy and lower intelligence scores at one year, but not two, possibly due to intervening environmental factors, such as positive parent-child relationships.⁶¹ Another study did not find lower intelligence scores at age five for children prenatally exposed to smoke.⁶² (However, after adjusting for factors such as education, income, and prenatal alcohol use, children in this study whose parents smoked when their kids were in early childhood did have lower scores.)⁶³

Children of mothers who smoked while pregnant and/or after giving birth appear to have more behavior problems, such as anxiety, trouble relating to other children, disobedience and difficulty concentrating.⁶⁴ Daughters of mothers who smoked during pregnancy, regardless of whether their mother smoked after giving birth, have been found four times likelier to smoke by age 13 than daughters of mothers who did not smoke during pregnancy.⁶⁵ One possible explanation is that smoking during pregnancy can affect the brain of the female fetus at a critical point in development and predispose her to nicotine addiction as a teen.⁶⁶

Quitting or reducing smoking during pregnancy can improve birth weight and reduce the chances of premature birth.⁶⁷ A study of maternity clinics comparing pregnant smokers and women who either quit or cut back during pregnancy found that, after controlling for the mother's race, age and height and the child's gestational age at delivery, infants born to quitters were, on average, more than half a pound (241 g) heavier than infants born to smokers, and infants born to women who reduced their level of smoking averaged a fifth of a pound (92 g) more. Quitting altogether increases the length of pregnancy by one week, reducing the chance of complications and birth defects due to premature birth.⁶⁸ Even quitting during the third trimester can significantly improve a baby's birth weight because so much of fetal growth occurs at this time.⁶⁹

Alcohol

Drinking during pregnancy produces a range of consequences, from fetal death and miscarriage to no apparent problems for some children.⁷⁰ Compared to non-drinkers, the risk of spontaneous abortion during the second trimester has been found to be twice as high for those who had one to two drinks a day and 3.5 times as high for those who had more than three drinks daily.⁷¹ Drinking while pregnant increases the risk of death during the infant's first year by more than 50 percent. While the infant mortality rate among women who don't drink during

Drinking during pregnancy can more than double a child's risk of acute myeloid leukemia, a potentially fatal cancer.¹

¹ Shu, X. et al. (1996). Parental alcohol consumption, cigarette smoking and risk of infant leukemia: A children's cancer group study. Journal of the National Cancer Institute, 88(1), 24-31.

pregnancy is 8.6 per 1,000, it rises to 13.3 per 1,000 among those who do (S) (CHART 27).⁷² The infant death rate is much higher--23.5 per 1,000--among pregnant women who drink an average of 2 or more per day (S).

If the child survives, the most severe outcome is Fetal Alcohol Syndrome (FAS), characterized by central nervous system dysfunction, including abnormally small head size and mental retardation, small length and weight, and facial malformations.⁷³

The reported rate of FAS increased almost seven-fold from 1.0 per 10,000 live births in 1979 to 6.7 in 1993.⁷⁴ Hospitals recorded about 2,700 cases of newborns with FAS that year. Since overall drinking rates did not rise from 1979 to 1993, the increase in FAS is probably due to greater awareness of the syndrome and better diagnosis of it. Many experts believe FAS is still under-reported because so many of its symptoms are not obvious at birth.⁷⁵ The latest estimate for the U.S. is a rate of 19.5 per 10,000 births, although estimates run as high as 30 per 10,000--about 12,000 babies a year.⁷⁶ Thousands more suffer from central nervous system abnormalities that are called Fetal Alcohol Effects.⁷⁷ The number of children with Fetal Alcohol Effects may be triple the number with FAS.⁷⁸

Since American researchers first identified FAS in 1973, evidence has grown that its consequences endure for many years.⁷⁹ The Seattle Longitudinal Prospective Study on Alcohol and Pregnancy, which has followed children of women who drank while pregnant in 1974 and 1975, found that physical signs of FAS may become less distinctive with age, but developmental problems with intelligence, memory and attention persist at least to age 14.⁸⁰

* Data do not account for use of tobacco and illicit drugs.

Binge drinking and the number of drinks per occasion are strong predictors of problems, indicating that a high peak blood alcohol level is especially damaging.⁸¹

Some experts caution that consuming an average of two drinks a day may lead to spontaneous abortion or neurobehavioral problems after birth.⁸² Even lower levels of drinking have been linked to subtle learning and behavior impairment.⁸³ But

some experts say research has not

established that consuming two or fewer drinks per day poses any risk to the fetus of a well nourished pregnant woman who does not smoke and gets prenatal care.⁸⁴ Even so, because a safe level of drinking during pregnancy has not been established, the Surgeon General, the U.S. Departments of Agriculture and Health and Human Services, and many obstetricians recommend that women abstain from alcohol during pregnancy or if they are trying to get pregnant.⁸⁵

The timing of alcohol use during pregnancy appears significant.⁸⁶ Drinking early in pregnancy can lead to physical malformations that characterize FAS.⁸⁷ Drinking later in pregnancy may retard the baby's weight gain and overall growth (length and head circumference). As a result, women who stop drinking during pregnancy are less likely to have babies who are born underweight or with small heads.⁸⁸

As children with FAS become adults, they can achieve a wide range of self-sufficiency and intelligence. For example, some are illiterate, while others read at a high school level.¹ If more physicians learn how to detect the signs of FAS in infants, early identification will make it possible to provide the support they need—from parents, health care providers and educators—to help them reach their full potential.

¹ Streissguth, A. P. (1992). Fetal alcohol syndrome and fetal alcohol effects: A clinical perspective of later developmental consequences. In I. S. Zagon, T. A. Slotkin (Eds.), Maternal substance abuse and the developing nervous system (pp. 5-25). San Diego, CA: Academic Press.

Chart 27

The Link Between Drinking During Pregnancy and Infant Death

	<u>Infant Mortality Rate (per 1,000)</u>
Non-Drinking Pregnant Women	8.6
All Drinking Pregnant Women	13.3
< 2 drinks per day	12.7
≥ 2 drinks per day	23.5

Source: Centers for Disease Control & Prevention, National Center for Health Statistics,
Linked Birth/Infant Death Data Set, 1991

Many newborns who were prenatally exposed to alcohol do well, but we don't know enough about what factors protect some or put others at risk. About six percent of alcoholic women have a child with FAS.⁸⁹ Rates of FAS are much higher among African American and Hispanic alcoholics who live in poverty, have low levels of education and get late or no prenatal care than among white alcoholics who have higher incomes and levels of education, and who get timely prenatal care--even though their rates of heavy drinking are similar.⁹⁰

While the infant death rate among women who drink during pregnancy is 8.5 per 1,000 for whites, it jumps to 13.1 per 1,000 for Hispanics and 28.1 for African Americans (S) (CHART 28).⁹¹ The combination of heavy drinking and any smoking is particularly deadly. The infant death rate is 10.9 per 1,000 among white women who drink and smoke heavily during pregnancy; for black women who are both heavy smokers and drinkers, the death rate soars to 40.7 (S) (CHART 29).

One study of 84 alcoholic women found FAS at a rate of 70.9 percent among black and Hispanic women who drank an average of 12 drinks a day and 4.5 percent among white women of higher income and education who drank at similar levels.⁹² A quarter of black and Hispanic women, compared to 61 percent of the white women, were married; 23 percent of black and Hispanic women, compared to 97 percent of the white women, were high school graduates. Such wide disparities suggest the importance of environmental factors in FAS.

Genetic factors also appear to play a role. Reports of alcohol-exposed fraternal twins, one with FAS and one with less severe or no apparent effects, indicate that fetuses have

* Data do not account for use of tobacco and illicit drugs.

different levels of susceptibility to alcohol exposure, which are apparently genetic. This variance in susceptibility was not found in identical twins--who have identical genes.⁹³

Cocaine

In the mid 1980s, reports of birth complications and developmental problems associated with cocaine flooded the popular press and sparked concern over the fate of "crack babies."⁹⁴ Yet few early studies on cocaine considered that "crack babies" are often also alcohol, tobacco, Valium, marijuana and/or other drug-exposed babies born to malnourished women with little or no prenatal care. Subsequent research, which avoided many shortcomings of early work, has confirmed some fears of the serious damage cocaine can do to the developing fetus. But many cocaine-exposed infants are surprisingly resilient, overcoming initial signs of physical and cognitive impairment; others show no apparent effects from their mother's prenatal cocaine use.⁹⁵

But not all are so lucky. Cocaine use during pregnancy is linked to microcephaly (abnormally small head size) and prematurity.⁹⁶ Multiple drug use aggravates these

Marijuana, the illicit drug most commonly used by pregnant women, may cause intrauterine growth retardation, low birth weight, cognitive deficits and jitteriness in children.¹ Much more research is needed to assess the impact of marijuana, the illicit drug most commonly used during pregnancy.

¹ Fried, P. A. (1993). Prenatal exposure to tobacco and marijuana: Effects during pregnancy, infancy, and early childhood. Clinical Obstetrics and Gynecology, 36(2), 319-337; Khalsa, J. H. et al (1991). Epidemiology and health consequences of drug abuse among pregnant women. Seminars in Perinatology, 15(4), 265-270; Zuckerman, B. et al. (1992). Prenatal cocaine and marijuana exposure: Research and clinical implications. In Zagon and Slotkin (Eds.), Maternal substance abuse and the developing nervous system (pp. 125-153). San Diego, CA: Academic Press; Day, N. L. et al (1991). Prenatal marijuana use: Epidemiology, methodological issues and infant outcome; Clinics in Perinatology, 18(1), 77-91; Parker, S. et al. (1990). Jitteriness in full-term neonates: Prevalence and correlates. Pediatrics, 85(1), 17-23.

The Link Between Drinking During Pregnancy and Infant Death by Race and Ethnicity

	Infant Mortality Rate (per 1,000)		
	White	Black	Hispanic
Non-Drinking Pregnant Women	7.0	16.0	6.9
All Drinking Pregnant Women	8.5	28.1	13.1
< 2 drinks per day	8.5	27.0	13.7
≥ 2 drinks per day	9.5	35.1	*

* The sample was too small to provide a death rate.
 Source: Centers for Disease Control & Prevention, National Center for Health Statistics,
 Linked Birth/Infant Death Data Set, 1991

Chart 29

The Link Between Smoking and Drinking During Pregnancy and Infant Death

by Race

	Infant Mortality Rate (per 1,000)	
	White	Black
Non-Smoking, Non-Drinking	6.2	15.3
Smoking and Drinking:		
< 1 pack & < 2 drinks per day	8.8	30.5
≥ 1 pack & ≥ 2 drinks per day	10.9	40.7

Source: Centers for Disease Control & Prevention, National Center for Health Statistics, Linked Birth/Infant Death Data Set, 1991

consequences. Infants exposed to cocaine and other drugs have greater reductions in gestational age, head circumference and length than those exposed only to cocaine.⁹⁷

Because some women believe that cocaine initiates labor and eases the pain, they may use it intentionally to relieve the discomfort of pregnancy and induce labor, exacerbating the risk of prematurity.⁹⁸ Cocaine use has also been associated with abruption placentae (premature detachment of the placenta), which can cause miscarriage, premature birth, stillbirth and hemorrhaging that endangers the woman's life.⁹⁹

By triggering premature delivery, cocaine causes low birth weight.¹⁰⁰ Cocaine may also reduce a baby's weight by suppressing a woman's appetite or reducing blood flow to the fetus.¹⁰¹ A study at Harlem Hospital found that, controlling for demographic and lifestyle factors (including tobacco and alcohol use) and gestational age, cocaine use was associated with an average weight reduction of a quarter of a pound (125 g); when women used cocaine as well as other illicit drugs, the average decrease was a third of a pound (195 g).¹⁰²

Infants exposed prenatally to cocaine may be tremulous, irritable, erratic sleepers and unable to suck properly.¹⁰³ While these symptoms mimic signs of withdrawal from the drug, they more likely represent the effects of the drug itself on the infant. Although they do not require medication, such infants should be cared for in ways that are calming, such as being swaddled in a blanket, held firmly and rocked gently, in areas with dim lighting, low noise levels and soft music.¹⁰⁴

Cocaine has also been linked to cognitive and neurobehavioral problems, although study results have been mixed, which makes it difficult to draw definitive conclusions. Some infants exposed to cocaine: have difficulty coordinating movement, which can make it hard

for them to roll over, lift their heads up, put thumbs into their mouths and walk; are easily overstimulated (shaking when startled, or breathing quickly and crying easily when aroused); and have trouble tracking visual stimuli.¹⁰⁵

As cocaine-exposed infants grow up, they may be easily frustrated and act out because of their low tolerance for stimulation. They require stable home and school environments with predictable routines, rules, discipline and nurturing.¹⁰⁶ Behavior problems, such as antisocial and violent behavior, have been observed--but not scientifically researched--in children exposed to cocaine in the womb.¹⁰⁷

Follow-up studies at ages two and three have compared children prenatally exposed to cocaine and drugs other than opiates (such as marijuana, alcohol and/or tobacco), those exposed to drugs other than cocaine (marijuana, opiates, alcohol and/or tobacco), and those with no evidence of prenatal exposure to alcohol or illegal drugs.¹⁰⁸ Few significant

Because babies can receive any drug used by a woman through her breast milk, mothers who do not stop using drugs, alcohol or tobacco may do harm to their children.¹

◆ *Infants who are breast-fed by cocaine-using mothers may show signs of cocaine intoxication (such as hypertension, irritability and tremulousness).*

◆ *Cigarette smoking by nursing mothers has been associated with infant vomiting, diarrhea, restlessness and colic.*

◆ *The danger of drinking by breastfeeding women is unclear, but children of mothers who had at least one drink a day while breast-feeding have been found to have slightly poorer motor development at age one than women who had less than a drink a day.*

¹ Kharasch, S. et al. (1990). Unsuspected cocaine exposure in young children. American Journal of Diseases of Children, 144, 441; Chasnoff, I. J. et al (1987). Cocaine intoxication in a breast-fed infant. Pediatrics, 80(6), 836-838; Wilton, J. M. (1992). Breastfeeding and the chemically dependent woman. NAACOG's Clinical Issues in Perinatal and Women's Health Nursing, 3(4), 667-672; Little, R. E. et al. (1989). Maternal alcohol use during breast-feeding and infant mental and motor development at one year. New England Journal of Medicine, 321(7), 425-430.

differences in overall developmental scores (measuring mental and psychomotor skills) were found, and differences in weight and height that were apparent at birth had disappeared by age one. But children exposed to cocaine and other drugs continued to get overstimulated easily, and both drug-exposed groups had smaller head circumferences than unexposed children at age two. Cognitive functioning, measured by intelligence tests, was lower in the two drug-exposed groups, although three factors appeared to moderate this impact: quality of the home environment, child's head circumference and child's perseverance at tasks.¹⁰⁹

Because families in these studies received exceptional services, including prenatal care, psychotherapeutic interventions for the mothers and early physical and speech therapy for the children, they represent a "best case" scenario. This and other research suggests that many children exposed prenatally to cocaine are remarkably resilient, given a positive home environment and early interventions.¹¹⁰

Other consequences of prenatal cocaine use remain uncertain. Research conducted in the early years of the crack epidemic found a high risk of SIDS in a small sample of cocaine users.¹¹¹ However, subsequent research has not replicated this finding. As a result, it is not clear how much of the risk may stem from cocaine use and how much may be a consequence of other factors such as smoking.¹¹²

Research has also suggested that cocaine may cause birth defects.¹¹³ Studies have found an increased risk for congenital anomalies such as genitourinary tract malformations, but other research has found no significant impact of cocaine on birth defects after accounting for demographics such as income, education, race and the use of other drugs.¹¹⁴

Heroin

Heroin use during pregnancy has been linked to low birth weight, miscarriage, prematurity, small head size and intrauterine growth retardation.¹¹⁵

Here again, however, many studies on such use do not account for other factors such as tobacco, alcohol and other drug use, lack of prenatal care, and poor nutrition.

Newborns exposed to heroin in the womb often experience withdrawal from the drug after birth. They shake with tremors; cry in high-pitched tones; have tight or tense muscles and clenched fists; are irritable and inconsolable; sleep, feed and breathe irregularly; vomit; and may even have seizures.¹¹⁶ Using proper treatment and medication, doctors can stabilize these children.¹¹⁷

Some treatment providers maintain pregnant heroin addicts on methadone. They argue that methadone treatment can produce stable drug levels (preventing the peaks and valleys of heroin use), remove the woman from the dangerous environment of illegal drug sales, and lead to longer pregnancies than heroin use and normal developmental scores up to

Her baby cried and seemed out of sorts. Nothing would settle her down. So, Jacqueline Rosetta Edwards did the only thing she knew to do. She fed heroin to her own 4-month-old daughter. Edwards told the police that she had been hooked on heroin and cocaine for a dozen years and assumed her daughter was born addicted as well. Instead of seeking medical help, she decided that small amounts of heroin would help the baby through withdrawal. "She said she gave it to the child to try to ease the withdrawal and calm it down because it would cry a lot and carry on," said the police department's Lt. Albert Scott. Age 28, she appeared in court this morning to answer charges of drug distribution and felony neglect and abuse.¹

¹ Baker, Peter. Virginia mother allegedly gave heroin to her baby. Washington Post, 5/10/95, p. D3.

Edwards received a 10-year prison sentence and three years' probation. She will be required to serve at least 85 percent of the term.²

² Heroin fed to baby. Washington Post, 8/29/95, p. C4.

age four.¹¹⁸ But it is recognized as no panacea, and many pregnant women taking methadone continue to use other drugs.¹¹⁹ At best, methadone use here is the lesser of two evils.

Prescription Drugs

Benzodiazepines--the tranquilizers commonly used to treat anxiety, epilepsy and insomnia--have been linked to birth defects, but not in reliable studies that take the use of other drugs into account.¹²⁰ Before the advent of benzodiazepines in the 1960s, doctors often prescribed barbiturates for these symptoms, and many women used or misused them during pregnancy.¹²¹

An estimated 22 million children in the U.S. were born exposed to phenobarbital, a barbiturate, from 1950 to the late 1970s.¹²² Barbiturates double the risk of birth defects and can produce withdrawal in infants similar to that caused by heroin.¹²³ Some adult men prenatally exposed to phenobarbital had lower scores on intelligence tests than non-exposed men.¹²⁴

Today barbiturates, such as phenobarbital, are still used medically during pregnancy to control epilepsy because physicians consider the risks of discontinuing the medication more serious than the increased chance of birth defects.¹²⁵

The Next Drug

Like heroin in the 1970s and cocaine in the 1980s, another drug will come into fashion in the years ahead, hooking its users and spreading its damage to their children and

families. Amid indications that the crack cocaine epidemic may be subsiding, the next drug of choice may be methamphetamine, also known as "speed" or "ice."¹²⁶ Reports that amphetamine use is spreading rapidly have prompted concern over a new epidemic.¹²⁷

While little research has isolated the impact of amphetamine use during

pregnancy, amphetamines have been linked to intrauterine growth retardation and prematurity, and newborns may experience withdrawal from the drug.¹²⁸ If an amphetamine epidemic emerges on the scale of the cocaine epidemic of the 1980s, then research to assess its impact on pregnancies, as well as on the patterns and effects of use to inform prevention and treatment efforts, will be desperately needed. Even more urgent is a major education campaign to inform women of the hazards of amphetamines and all other substances, including tobacco and alcohol.

Child Abuse and Neglect

Drug and alcohol abuse by either or both parents is a top cause and exacerbator of child abuse and neglect.¹²⁹ Alcohol and drug disorders increase the risk of child abuse five times and neglect nine times, after controlling for factors such as depression and social support.¹³⁰ Alcohol abuse is also a significant factor in sexual abuse of children.¹³¹

A startling rise in the use of methamphetamine in California and the Southwest has raised fears among law enforcement officials that the trend toward dangerous stimulants could spread across the country much as crack did in the 1980s.... Young men use methamphetamine for sexual stimulation. Because the drug suppresses appetite, it has also been tried by teenage girls trying to lose weight.¹

¹ Wren, Christopher. Sharp rise in use of methamphetamines generates concern. New York Times, 2/14/96, p. A16.

In a study of 200 alcoholic or opiate addicted parents, all children were abused or neglected to some degree; nearly a third were seriously neglected (30.5 percent); nearly a quarter were physically or sexually abused (22.5 percent).¹³² Almost half (42 percent) of these parents had been abused by their own parents during childhood.

Perhaps more than any other drug, including heroin, crack cocaine becomes the primary "relationship" in the addict's life at the expense of any instincts about her welfare and that of her children.¹³³ Addicts experiencing intense cravings for crack may put fulfilling their desire for the drug above virtually everything else. Moreover, drug and alcohol abusing women are likely to have been victims of childhood abuse or neglect; caught in an intergenerational cycle, these women may be at high risk of becoming abusive or neglectful parents, easily frustrated by the irritability of children exposed to alcohol, tobacco and illicit drugs in the womb.¹³⁴

While many long-term developmental consequences of drug use during pregnancy are uncertain, the high risk of abuse and neglect after birth is not. Wherever you find prenatal drug and alcohol abuse, you will likely find child abuse or neglect.

Charlene, a 33-year-old recovering crack addict, prostituted herself for crack and coaxed her children to steal after she sold her food stamps to buy drugs. She would disappear for days, even weeks, at a time, forcing her older children to play hooky to care for the baby. "When you get high, you know your children are there, you just don't care," she said. "All you care about is that next hit."

¹ The crack in the system. Daily News, 12/4/95, p. 22; One sure way to save kids. Daily News, 12/5/95, p. 26.

It Takes Two: The Role of Dads

An important question is the role of a father's drug, alcohol or tobacco use on his newborn children. Only a handful of studies have sought the answer, but what research exists on animals and humans indicates that this area deserves further exploration.¹³⁵

Independent of maternal drinking, a third of a pound (137 g) reduction in birth weight has been found for infants born to fathers who, in the month before conception, drank two or more drinks per day or had at least five drinks on one occasion.¹³⁶

Smoking fathers have an indirect impact on fetal development. The risk of SIDS rises in infants born to women passively exposed to smoke by fathers during pregnancy, as well as to infants passively exposed to their fathers' smoke after birth.¹³⁷ Children of nonsmoking mothers and fathers who smoked more than a pack a day during the mother's pregnancy can weigh a fifth of a pound (88 g) less than children whose parents did not smoke.¹³⁸

Violence against pregnant women is often related to their and their partner's substance use. Male drug and alcohol use is linked to spousal violence, and pregnant women who are battered are more likely to use alcohol and drugs than other pregnant women.¹³⁹

Health and Social Welfare Costs

It is difficult to estimate the total health and social welfare costs of substance abuse to the U.S. because no such comprehensive study has been undertaken, and because smaller cost studies either overlap or produce inconsistent results. However, based on previous CASA analyses, we think the health and social welfare costs approach \$10 billion.

The health care costs of caring for babies prenatally exposed to alcohol, tobacco and/or illicit drugs came to some \$6 billion in 1995.¹⁴⁰ Most of the bill--\$5.8 billion--stems from hospital care at birth. After discharge, costs for treatment of congenital anomalies and low birth weight add another \$132 million; these are undoubtedly a low estimate of post-discharge costs because providers often fail to note that many other medical problems during infancy may be the result of prenatal alcohol, tobacco and drug use.

In 1990, the U. S. General Accounting Office estimated that hospital charges in 1989 for infants exposed to illicit drugs were up to four times greater than those for drug-free infants. At one hospital, the median charge for drug-exposed infants was \$5,500, compared to \$1,400 for non-exposed infants--an added cost of \$4,100.¹⁴¹ A study in New York City put the average cost of additional medical care in the hospital for newborns exposed to cocaine and possibly other drugs at nearly \$5,200 in 1990.¹⁴² In another study in New York City, infants exposed to cocaine and other drugs, including nicotine, stayed in the hospital about seven days longer than drug-free newborns at an additional cost of \$7,731.¹⁴³

Boarder babies are a tragedy that adds to health care costs. The cost of caring for these babies, even if they have no medical complications, ranges from \$200 to \$500 a day--or up to \$15,000 a month.¹⁴⁴ With an average stay of four months for these children, the bill can reach \$60,000.¹⁴⁵

After release from the hospital, costs continue to mount. In 1990, the Florida Department of Health and Rehabilitative Services estimated that by age 18, the cost of

services such as hospital costs, foster care and special education, for a child with physiologic or neurologic impairment would reach at least \$750,000.¹⁴⁶

Take foster care: A survey of 10 hospitals in 1989 found the cost of one year in foster care for 1,200 drug-exposed infants to be \$7.2 million.¹⁴⁷ Given that in 1991, some 49,000 children under age three, prenatally exposed to drugs, were in foster care, the annual cost of foster care for the youngest drug-exposed children was some \$325 million.¹⁴⁸ In New York City alone about 43,000 children are in foster care, and 77 percent of the foster care budget in 1994 (\$595 million of the \$775 million) was attributable to parental drug and alcohol abuse.¹⁴⁹ If parental substance abuse is a factor in three-quarters of all foster care placements across the nation, the cost would be about \$3 billion.¹⁵⁰

Licit drugs also take their toll. The total cost to the economy of FAS (both for treatment and from lost productivity) was about \$2.7 billion in 1995.^{**151} The cost of low birth-weight infants in neonatal intensive care because of their mother's cigarette smoking is estimated to be as high as \$792 million per year.¹⁵² The cost of caring for individuals with FAS, including medical care expenses and institutional stays for those who are mentally retarded, is at least \$90 million.^{**153}

* This cost was adjusted to 1991 dollars using the Consumer Price Index.

** These costs were adjusted to 1995 dollars using the Medical Consumer Price Index.

V.

Combatting the Problem

To be effective for women, any concerted effort of prevention, treatment, law enforcement and research must take into account how markedly women and men differ in their risk factors, physiology, psychology and patterns of abuse and addiction.¹ Each stage of a woman's life is also unique, and the impact of tobacco, alcohol or drug use varies during adolescence, when their development into a healthy, productive adult is at stake; during pregnancy, when their actions affect not only themselves but the health of their children; or later in life, when their health determines how much independence and opportunity they'll enjoy in their golden years. Each stage requires a tailored response.

Prevention

Public education campaigns should explain how substance abuse may lead to severe health and social consequences for women and the unique perils of tobacco, alcohol and drug use and abuse during pregnancy. Each woman should know why she should not drink as much as men, how differently from men she metabolizes alcohol, how much more rapidly she will suffer alcohol-related illnesses, how alcohol suppresses physical sexual response and causes or aggravates sexual dysfunction, how poorly heavy drinking serves to ease tension or stress, and how her alcohol or drug use can make her more vulnerable to unwanted sexual advances, rape and violence.²

The different risk factors that mark women's substance abuse and addiction create opportunities for targeted prevention. Antecedents such as a family history of alcohol or drug abuse; abuse and violence during childhood, adolescence or adulthood; depression; and substance abuse by a partner serve as warnings for women. These women may be found in physicians' and psychologists' offices, mental health clinics, support groups for children of alcoholics, domestic violence shelters, family planning clinics, jails, child care centers, supermarkets, aerobics classes and churches.³ All these sites and situations offer opportunities for prevention.

Children and Teenagers

Adolescence is a time of high susceptibility to substance abuse and a critical window for prevention. To be effective, prevention efforts here must recognize the different factors that motivate girls and boys.⁴ Girls who have been sexually abused should be taught the dangers of self-medicating with alcohol or drugs and offered alternative ways to cope with the trauma of such abuse.⁵ Women are more likely than men to say they began binge drinking and smoking marijuana as teenagers to ease stress or get away from their troubles.⁶ Strong family bonds and peer disapproval may discourage girls more than boys from alcohol or drug abuse.⁷ Girls whose peers disapprove of drinking are 80 percent less likely to be problem drinkers than boys with such peers.⁸

Smoking by girls deserves special attention. In addition to the deadly consequences of smoking, tobacco is more strongly linked to illicit drug use for girls than for boys.⁹ While male drug users often use alcohol as a gateway substance before moving onto illicit

drugs, female drug users are more likely to start smoking cigarettes (sometimes in addition to drinking) before such use.

Prevention efforts must start early because nicotine addiction takes hold quickly.¹⁰

Among females ages 12 to 24 who smoke five or fewer cigarettes a day, 52 percent already feel dependent; of those who smoke six to 15 cigarettes daily, 81 percent consider themselves hooked.¹¹ Women in their 20s who began smoking at or before age 13 have a harder time quitting smoking than those who started later--another indication that prevention campaigns focused at the preteen and early teen years are critical.¹²

An important factor in the relative ineffectiveness of public health campaigns against smoking during the 1960s and 1970s with respect to teenage girls was their failure to take into account the different reasons that girls and boys smoke--reasons that tobacco companies have exploited.¹³ Teenage girls are often obsessed with weight. The desire to be thin motivates girls far more than boys.¹⁴ Girls may also use smoking as a way to deal with depression or anxiety.¹⁵ For girls, teaching healthy methods of weight control, boosting

Warning signs of problem drinking by adolescent girls:¹

- ◆ *Peer use of alcohol*
- ◆ *Expectation that alcohol will "provide a high" or be disinhibiting*
- ◆ *Antisocial or aggressive behavior, temper tantrums, shoplifting or vandalism*
- ◆ *Frequent absences from school and low educational aspirations*
- ◆ *Alienation*
- ◆ *Early experience with alcohol intoxication*
- ◆ *Early use of marijuana*
- ◆ *Family history of alcohol abuse and dependence, marital conflict or inadequate parenting*

¹ Gomberg, E. S. L. (1994). Risk factors for drinking over a woman's life span. Alcohol Health and Research World, 18(3), 220-227.

self-esteem and finding alternative ways to cope with problems are integral to smoking prevention.¹⁶

Prevention campaigns targeting teenage girls should deglamorize smoking and show its immediate and distasteful effects to counter tobacco advertising that emphasizes thinness and sexy independence.¹⁷ For example, the bumper sticker that says, "Kissing a Smoker is Like Licking a Dirty Ashtray" emphasizes the short-term consequences of smoking, such as unpleasant smells, yellow teeth and less attractiveness to the opposite sex.¹⁸ Challenging the belief that smoking is socially acceptable and common among their peers (the idea that "everybody's doing it") may also be effective. It is imperative to address teenage girls' concerns about weight and body image and give them a better understanding of how to stay healthy as part of any anti-smoking effort directed at them.

Women's magazines bear a special responsibility here.¹⁹ Critics charge that magazines such as McCall's and Woman's Day have courted tobacco advertising by providing scant coverage of the dangers smoking poses to women.²⁰ Teenage girls may be particularly reliant on what they

Many leading women's magazines have a substantial readership among teenage girls. Glamour and Vogue are both read by more than 2.6 million girls ages 12 to 19. Other magazines, such as Family Circle and McCall's, reach the mothers of more than a million teenage girls.¹

¹ Kaiser Family Foundation. (1996). "Leading Women's, men's, teens' and minority magazines by circulation and readership."

learn from such media. High school girls name the mass media as the best source of information on alcohol and drugs (25.2 percent), more important than friends (17.2 percent), personal experience (16.6 percent), family (14.2 percent), their doctors (7.8 percent) and their teachers (6.4 percent).²¹ While alcohol and drug education by parents, teachers,

clergy and other adults is important, it must be reinforced by communication of the dangers of substance abuse in magazines, television shows and movies that girls read and look to for role models.

Girls appear to be more responsive than boys to warnings about the hazards of drug, alcohol or tobacco use; if they think a substance is dangerous to their health, they are less likely to use it.²² In a survey of 5,891 eleventh-graders in Ohio during 1977, 1980 and 1983, girls perceived most substances--but not cigarettes--as more harmful than boys did, and this perception translated into lower levels of use among girls.

Another element of prevention for children and adolescents is reducing the supply of the substance. In a 1994 study in Massachusetts, 12 youths (ages 12 to 17) made 480 attempts to buy tobacco products from vending machines and over-the-counter retailers.²³ Only 28 percent of vendors consistently did not sell to minors. Girls were one and a half times likelier to be successful at purchasing cigarettes, even when taking into account how old they looked.* "It's the Law" programs sponsored by the tobacco industry--where stickers with the legal age to buy tobacco products are displayed on tobacco vending machines and in store windows--had no impact; vendors in these programs were just as likely as other vendors to sell to children and teenagers.

Young Adult Women: At College and in the Workforce

Alcohol advertisers are targeting young women as a "growth market" by trying to make their drinking more socially acceptable.²⁴ Prevention of heavy drinking is particularly

* "Apparent age" was measured by asking individuals who did not know the youths to guess their ages.

important for women on college campuses. Nationally, women are still less likely than men to binge drink at college (39 percent vs. 50 percent); however, here again the gap appears to be closing. In Massachusetts the percentage of women in their first year of school who drink "to get drunk" tripled from 10 percent in 1977 to 34 percent in 1989.²⁵ At the University of Wisconsin in Milwaukee, beer consumption among women rose 34 percent over roughly the same period.²⁶ Eighty percent of sorority members and 50 percent of women involved in college athletics nationwide binge drink; both are important targets for prevention campaigns.²⁷ At the University of Alabama, women and men were found to be equally likely to get drunk whenever they drink (45.1 percent vs. 46.1 percent).²⁸ Some women feel pressured to keep up with men when drinking. "If you can't drink with the guys, people don't respect you as much," said a female student at William and Mary College.²⁹

Young employed women are another high risk group.³⁰ Relative to women who are homemakers, employment outside the home appears to increase the likelihood that a woman will drink heavily.³¹ There is some evidence that women who work in male-dominated occupations tend to drink more heavily than those in female-dominated occupations.³² Possible reasons

Karen Valenstein...the highest-ranking woman in the public-finance department of E.F. Hutton & Company and one of the pre-eminent women in investment banking...has an intuitive understanding of masculine culture, with all its aggressiveness, competition and politics; she also has a genuine enthusiasm for living by its fierce rules. By her own admission, she can trade locker-room vulgarities [and] belt back stingers until dawn....¹

¹ Gross, Jane. Having, getting, earning, eating. *New York Times Magazine*, 4/14/96, p. 136.

* Wechsler, et al, (1994) define binge drinking as four or more drinks in a row for women and five or more drinks in a row for men.

range from the stress of being a "token" female to the possibility that these women want to be--or feel pressured to be--"one of the boys."³³

Drinking, Smoking and Drug Use During Pregnancy

The best time to prevent smoking, drinking and drug use during pregnancy is before conception. During the first trimester, when many women do not even realize they are pregnant, the fetus is particularly susceptible to the effects of tobacco, alcohol and drugs.³⁴ According to the 1988 National Maternal and Infant Health Survey, 45.4 percent of recent mothers reported drinking in the three months before they learned they were pregnant.³⁵ The association of alcohol and illicit drug use with amenorrhea (ceasing menstruation) exacerbates this problem, because a woman using alcohol or drugs cannot rely on a missed period as a sign that she is pregnant.³⁶

Prevention of Fetal Alcohol Syndrome (FAS) requires educating all women of child-bearing age about the dangers of alcohol use during pregnancy, as well as identifying and treating alcoholic women before they become pregnant.³⁷ One way doctors and other health professionals can identify a woman at risk of having a child with FAS is to evaluate her older children.³⁸ The chance that an alcoholic woman will have a child with FAS is about 6 percent. But if her first child was born with FAS, the odds jump to 70 percent for the next one.³⁹

For the child, the first step is diagnosing FAS. Unfortunately, the disorder appears to be widely under-reported.⁴⁰ Even when physicians identify FAS, they are often reluctant to note the diagnosis on the child's records, thinking it's not necessary for appropriate

treatment.⁴¹ This attitude not only denies these children services they may need, but also hinders prevention efforts for future babies.

Warning Labels. Alcohol warning labels can help educate some women about the risks of drinking while pregnant. In the late 1970s and early 1980s, some state and local governments required the display of warnings about alcohol-related birth defects in bars and other locations where alcohol is sold.⁴² In 1988, the federal government mandated that alcohol bottles carry similar warning labels.⁴³ While these efforts may contribute to a drop in light or moderate drinking during pregnancy, their

efficacy in preventing the most serious alcohol-related birth defects, such as FAS, is dubious.

A study of 12,000 black women found that alcohol warning labels appeared to lead to a slight decrease in drinking among pregnant women who were light drinkers, but to have no significant impact on heavier drinkers.⁴⁴ Another study found that nearly three-quarters of

An obstacle to prevention is the fact that many women know someone who smoked or drank during pregnancy and delivered an apparently unharmed child.

◆ *Alcohol warning labels appear to inspire a small but significant decline in drinking among women pregnant with their first child, but to have no impact on pregnant women with at least one child, suggesting that they discounted the warning because of their own experience.¹*

◆ *In a study of pregnant teenagers who smoke, family members or friends who smoked during pregnancy and delivered "healthy" babies were an important model. A typical comment was, "Five of my friends smoked a pack a day. Their babies are healthy."²*

¹ Hankin, J. R. et al. (1996). Heeding the alcoholic beverage warning label during pregnancy: Multiparae versus nulliparae. Journal of Studies on Alcohol, 57(2), 171-177. ² Lawson, E. J. (1994). The role of smoking in the lives of low-income pregnant adolescents: A field study. Adolescence, 29(113), 61-79.

* The label says: "Government warning: According to the Surgeon General, women should not drink alcoholic beverages during pregnancy because of the risk of birth defects."

women of child-bearing age did not recall the information on labels; African American women were least likely to do so.⁴⁵ Warning labels do not deter heavy drinkers, who are at greatest risk of having a child with FAS; helping such women to stop drinking requires much more.⁴⁶

Prenatal Care. When efforts to deter alcohol, tobacco or drug use before conception fail, prevention efforts can reduce the risk of further damage to the child by discouraging continued substance use or abuse among women who are already pregnant and, if necessary, referring them to treatment.⁴⁷ Women who stop drinking, smoking or using drugs during their pregnancy can improve the overall health of their child and increase the birth weight.⁴⁸

Illicit drug users who receive prenatal care have healthier babies than those who do not in terms of growth and prematurity.⁴⁹ During prenatal visits, doctors and nurses can identify addicts, refer them to treatment and provide ongoing support and encouragement.⁵⁰

Because even minimal efforts by health care providers can help women quit smoking, routinely incorporating advice on smoking cessation into prenatal visits can have a substantial impact.⁵¹ Brief physician advice, counseling and encouragement to stop smoking increases the likelihood of quitting.⁵² At each visit, doctors should urge a smoker to quit and personalize the message by relating it to her individual health.⁵³ Cessation programs that reinforce physician's advice with printed materials, home visits and/or phone calls can increase success.⁵⁴

Yet doctors and nurses often miss these opportunities. In a sample of 56 pregnant drug users, while most got prenatal care, their physicians did not refer them for treatment.⁵⁵ Pregnant African American women are 20 percent less likely to be advised to quit smoking

and 29 percent less likely to be advised to quit drinking during a prenatal care appointment than are white pregnant women, after taking into account age, marital status, income, payment type and pregnancy history.⁵⁶ This is particularly disturbing given higher rates of FAS and infant mortality among black women.⁵⁷

Most addicted pregnant women receive no prenatal care at all.⁵⁸ Lack of access to prenatal care, long waits for appointments and the need to travel long distances to clinics deter pregnant women from seeking help--particularly those with jobs that pay by the hour.⁵⁹ For pregnant addicts, assuring access to prenatal care is not enough; she may require aggressive outreach, transportation and follow-up.⁶⁰

After Menopause

Alcoholism tends to take hold later in women's lives than in men's.⁶¹ Women at high risk because of "role deprivation" from recent divorce or separation have been the target of a prevention program in California that provides workshops in stress management and life and career planning, as well as alcohol education.⁶² Women who completed the program had higher self-esteem and used alcohol and drugs less than a similar group who didn't participate in it.

*Doctors often miss or ignore signs of drug use by pregnant women. One woman told of her experience with doctors when she was pregnant. "The first one--I got up my nerve and told him right out I was an addict. It didn't seem to affect his plans for me or my baby one bit. It was like he just did not want to hear me--he ignored what I said. So I went to another doctor. He never even asked [about my drug use] and I didn't tell him."*⁶¹

¹ U.S. General Accounting Office. (1991). ADMS Block Grant: Women's set-aside does not assure drug treatment for pregnant women. Washington, DC: GAO.

Among elderly women, the physician's role in preventive education and counseling is especially important because these women may be unaware of their increasing vulnerability to alcohol's effects and dangerous interactions that it can trigger with prescription drugs.⁶³ Elderly women may also be lonely and isolated from family and friends who otherwise might detect early signals of problem drinking and intervene.⁶⁴

To avoid dangerous interactions with other drugs or alcohol, doctors who prescribe psychoactive drugs to elderly women should:

- ◆ *Screen carefully for drug and alcohol abuse before prescribing any drug for depression.*
- ◆ *Discontinue as many drugs as possible before starting the psychoactive drug.*
- ◆ *Prescribe small doses and increase doses cautiously.*
- ◆ *Warn patients of the dangers of mixing sedating drugs with alcohol.¹*

¹ Richelson, E. (1984). Psychotropics and the elderly: Interactions to watch for. Geriatrics, 39(12), 30-42.

Treatment

Despite claims that addicted women are harder to treat than men, women do not fare worse in treatment.⁶⁵ Though most studies on treatment outcomes do not examine gender variations, those that do find little difference between the sexes for the success of alcohol, tobacco or drug treatment.⁶⁶ An analysis of 20 studies on alcohol treatment found only small differences: women did slightly better in the first 12 months after treatment, and men did slightly better after the year.⁶⁷

However, women who need treatment for alcohol problems are less likely than men to get it. The 1992 National Drug and Alcohol Treatment Utilization Survey (NDATUS) found the ratio of men to women in alcoholism treatment was 3.5:1, compared to a ratio of alcohol

abuse or dependence of 2.4:1.⁶⁸ On the other hand, the 1992 NDATAUS found a ratio of men to women in drug treatment of 1.8:1, compared to a drug abuse or dependence prevalence ratio of 2.3:1.⁶⁹ But these ratios miss a central point: the National Association of State Alcohol and Drug Abuse Directors estimated that in 1989 only 13.7 percent of all women and 11.9 percent of pregnant women who need treatment receive it.⁷⁰

In 1989, the U.S. House of Representatives Select Committee on Children, Youth and Families surveyed 18 hospitals nationwide and found that in Los Angeles, pregnant women had to wait 10 to 16 weeks for drug treatment, and in Boston, the 30 residential treatment slots for pregnant women addicted to cocaine throughout the city fell short of the 300 cocaine users who delivered at just one hospital.⁷¹ Two-thirds of the hospitals reported having no place to refer pregnant substance abusers for treatment.⁷² In California, despite the addition of nearly 200 programs to serve pregnant and parenting women, 272 pregnant women were on waiting lists for drug treatment in 1993.⁷³

One reason for the lack of treatment for pregnant women is fear of legal liability if treatment fails to protect the child, and/or because of the potential impact of some medications during pregnancy.⁷⁴ In New York State, the Office of Alcoholism and Substance Abuse Services argues that pregnancy "in and of itself" is not a justifiable basis to deny treatment.⁷⁵

The Ways and Odds of Referral

Even where slots are available, the ways in which women tend to be identified as candidates for treatment may hinder their entry to treatment.⁷⁶ Physicians, the legal system

and employers are less likely to refer women to treatment.⁷⁷ Three effective early intervention methods--drunk-driver rehabilitation, public-intoxicant intervention and employee assistance programs--reach higher proportions of male problem drinkers than female.⁷⁸ This may be because men tend to exhibit alcohol abuse publicly while women tend to be more inner-directed. Police appear less likely to arrest women than men who are caught driving drunk.⁷⁹

In a study of 3,234 public and private employees in work settings, these factors combined to result in fewer referrals to alcoholism programs for women: male supervisors were less likely to identify women who were alcoholics than female supervisors; men were uncomfortable discussing drinking with female subordinates; and female employees seemed to make greater efforts to conceal drinking than male employees.⁸⁰ Among those who do enter treatment, women and men are just as likely to return to work, indicating that effective employee assistance programs work just as well for both genders.⁸¹

Among adolescents, fewer girls than boys with alcohol or drug problems get involved with the courts, where referrals to treatment often occur; but more girls are referred by parents.⁸² Girls ages 12 to 19 in treatment for alcohol or drug problems also are less likely than boys to get into trouble at school.⁸³

School nurses are in a unique position to identify and offer support to teenage girls who are substance abusers. They can ask about risk factors, such as physical or sexual abuse, and provide crisis intervention and referrals. They also can help prevent relapses as teens who have been in treatment return to school and need assistance finding drug-free peers.¹

¹ Tuttle, J. (1993). Adolescent substance abuse: Psychosocial factors. *Journal of School Nursing*, 9(3), 18-25.

A study of 305,000 alcoholic patients in treatment found analogous referral patterns

among the elderly.⁸⁴ Elderly men are more likely to have a legal referral (29 percent vs. 13 percent) than women, while elderly women are more likely to have a medical (12 percent vs. 9 percent) or a personal referral (self, employer, church, school, spouse, family or friend; 51 percent vs. 42 percent).⁸⁵

Among female alcoholics, the top four reasons for entering treatment are psychological consequences such as deepening depression, the physical and medical effects of alcohol, troubles with husband or children and trouble with the police.⁸⁶ In contrast, alcoholic men identify trouble on the job and with the law as the top reasons they entered treatment.⁸⁷

Such differences point to important opportunities--too often missed--for identifying substance abusing women and helping them, as one expert put it, "in doctor's offices, in hospitals and health clinics, in obstetric practices and gynecological clinics, in family and social service agencies, and in the offices of divorce lawyers."⁸⁸

Physicians: See, Speak and Hear No Substance Abuse in Female Patients?

Women see physicians more often than men, and the prevalence of alcohol disorders among women who seek medical attention is at least double that in the general population.⁸⁹ This imposes a responsibility on family practice doctors, gynecologists, obstetricians, pediatricians and other health care professionals to spot signs of substance abuse in their female patients and provide referrals and ongoing support to get them into treatment.⁹⁰ Gynecologists are often a woman's only doctor; because sexual dysfunction is associated with

alcoholism, routine screening by gynecologists can be an effective way to identify women for treatment.⁹¹

Yet physicians are less likely to diagnose abuse and addiction in women than in men.⁹² In one study, hospital doctors were three times more likely to fail to diagnose a female alcoholic than a male alcoholic.⁹³ At Johns Hopkins Hospital,

physicians were found less likely to identify alcoholics who were female, especially those with private insurance or higher incomes and education.⁹⁴

The reasons for this are many. If women have a substance abuse problem, they are likely to see a doctor for the medical or psychiatric symptoms associated with their abuse, such as digestive difficulties, insomnia or depression, rather than for the problem itself; indeed, the stigma of substance abuse may cause them to hide their alcohol or drug abuse.⁹⁵ By emphasizing a woman's psychological symptoms over her physical symptoms, doctors may attribute her alcohol-related problems to depression.⁹⁶ These physicians often exacerbate a drug or alcohol problem by inappropriately prescribing psychoactive drugs and failing to refer women to treatment.⁹⁷

Screening instruments that measure the quantity and frequency of drinking often do not take into account women's different metabolism of alcohol and the fact that many female alcoholics drink less than male alcoholics.⁹⁸ Assessment tools tend to focus on alcohol-

A national panel of doctors, insurance company representatives, public officials, business leaders and leaders in medical education concluded, "The primary-care specialties have done an inadequate job of training and preparing physicians in the treatment of substance-abusing patients." Such training, the panel said, should be required for all residents in family practice, internal medicine, obstetrics and gynecology and pediatrics.¹

¹ Gilbert, Susan. Doctors found to fail in diagnosing addictions. *New York Times*, 2/14/96, p. C8.

related issues that are more common among men, such as legal and job troubles, rather than those more relevant to many women, such as family conflict, violence, sexual abuse and reproductive dysfunction.⁹⁹

Among the elderly, medical conditions such as depression, anxiety and insomnia can mask the signs of alcohol abuse, while dementia can substitute as a diagnosis for alcohol-related behavior.¹⁰⁰ Physicians may also fail to adjust for a woman's declining tolerance for alcohol.¹⁰¹ What might be considered light or moderate drinking in her 30s can have serious health effects in her 60s or 70s. Adding to these concerns is evidence that elderly women who abuse alcohol are more likely than men to use prescribed psychoactive drugs and less likely to seek treatment.¹⁰²

With these differences in mind, routinely taking a thorough history of use of alcohol, tobacco, and licit and illicit drugs is an important step toward identifying female substance abusers and getting them help.¹⁰³ Characteristics associated with women's alcohol abuse, such as depression, family problems, an alcoholic partner or a family history of alcoholism, should serve as red flags to physicians.¹⁰⁴ Questions to elicit more information include:

Some recovering female alcoholics report that doctors were unhelpful or even discouraged them from seeking help.¹

◆ *"The priest sent me to a doctor, and he referred me to a psychiatrist."*

◆ *"I wrote to the analyst and he wrote back, 'You're too smart to be an alcoholic. Just don't drink!'"*

◆ *"I was having all these medical problems. The doctor asked if I was a heavy or a light drinker. Well, I didn't want to lie. So I would mock him. 'What's heavy and what's light?' I asked. When he said a heavy drinker drank every day, I had it beat. I didn't drink every day. But of course, when I did drink, I just kept on until I was through!"*

¹ Smith, A. R. (1986). Alcoholism and gender: Patterns of diagnosis and response. *Journal of Drug Issues*, 16(3), 407-420.

"Do you ever carry an alcoholic beverage in your purse? How has your drinking changed during pregnancies? What effect do you feel your drinking has had on your children?"¹⁰⁵

Doctors can ask elderly women, "Did you find your drinking increased after someone close to you died? Does alcohol make you sleepy so that you often fall asleep in your chair?"¹⁰⁶

If a physician determines that the woman has alcohol-related problems, he should advise her to cut down or abstain.¹⁰⁷ For patients who are not dependent, strategies to help quit include setting a specific drinking goal and helping patients recognize and cope with triggers for drinking. For alcohol-dependent patients, physicians can outline the available treatment options and schedule an appointment for the woman while she is in the office. Physicians (and nurses) can continue to monitor their patients' drinking behavior at subsequent appointments or by keeping in contact with their treatment program.

A screening tool known as the "CAGE" questionnaire identifies up to 90 percent of alcoholics with four simple questions:

- ◆ *Have you ever felt you ought to Cut down on your drinking?*
- ◆ *Have you ever been Annoyed by criticism about your drinking?*
- ◆ *Have you ever felt Guilty about your drinking?*
- ◆ *Have you ever had a drink first thing in the morning to steady your nerves or to get rid of a hangover (an Eye opener)?*

If a woman answers yes once or more, or drinks either at least eight drinks per week or at least four drinks per occasion, her physician should learn more about her patterns of use, reasons for drinking, any alcohol-related medical problems and her family history of alcoholism.¹

¹ National Institute on Alcohol Abuse and Alcoholism. (1995). The physicians' guide to helping patients with alcohol problems. Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; Cyr, M. G. and Moulton, A. W. (1993). The physician's role in prevention, detection, and treatment of alcohol abuse in women. Psychiatric Annals, 23(8), 454-462.

Barriers to Treatment

The shortcomings of referral routes are only one of the barriers to treatment women face. They also face a paralyzing lack of child care options.¹⁰⁸ Only 33 percent of women-only treatment sites and 6 percent of other treatment units provide child care for clients.¹⁰⁹ A 1993 survey of drug treatment programs in five cities found that most accept pregnant women (83 percent of outpatient and 70 percent of residential programs), but only about 20 percent both accept pregnant women and provide child care. In three of the five cities surveyed, no residential treatment programs accepted pregnant women and provided child care.¹¹⁰

As a result, female addicts who seek treatment may be forced to choose between leaving their children alone or with inadequate caretakers, placing their children in foster care or losing custody of their children altogether.¹¹¹ With these choices, many decide not to enter treatment. Offering child care to these women eliminates the need to make such a

Barriers to treatment for women¹

External barriers:

- ◆ *Different referral patterns*
- ◆ *Inadequate training of health professionals*
- ◆ *Opposition by family or friends*
- ◆ *Inadequate financial resources and insurance*
- ◆ *Comprehensive services not in a single location*
- ◆ *Lack of child care facilities*
- ◆ *Male-oriented treatment services*

Internal barriers:

- ◆ *Denial of the problem*
- ◆ *Guilt and shame*
- ◆ *Fear of stigmatization*
- ◆ *Fear of leaving or losing children*

¹ Beckman, L. J. (1994). Treatment needs of women with alcohol problems. Alcohol Health and Research World, 18(3), 206-211.

choice. PAR Village, a Florida residential treatment program, found that women allowed to bring their children to the program were five times more likely to stay for a year and, overall, stayed about three times longer than women not offered on-site child care.¹¹²

Women who seek treatment are more likely than men to lack support or face opposition from their spouses and family.¹¹³ Alcoholic women are more often divorced in the course of their alcoholism than are alcoholic men.¹¹⁴ A study of 6,400 employees at 84 worksites found that women with alcohol problems who entered treatment through an employee assistance program were less likely than men to have family support in seeking help for their problem.¹¹⁵ Most women were unmarried (68 percent had never married or were divorced or separated).¹¹⁶ Even when the women were married, their spouses were less likely to play a role in their getting help.¹¹⁷ At the Betty Ford Clinic in California, female patients often report that they delay seeking treatment because their husbands and children pressed them to stay home and care for the family.¹¹⁸

The stigma of drinking as unfeminine and inappropriate for women may lead husbands to deny their wife's alcoholism.¹¹⁹ A study of 10 male and 10 female alcoholics found women were more likely to be the first to identify their alcoholism, while men were more often identified by wives or families.¹²⁰

A recovering alcoholic woman described her husband's denial of her situation:¹
"Even after I'd been through treatment, I'd have a hard time at home. My husband kept telling me, 'You're not an alcoholic. You can have a drink if you want to.' What he meant was if he wanted me to! Anyway, eventually, it was a choice I had to make: leave and live, or drink again."

¹ Smith, A. R. (1986). Alcoholism and gender: Patterns of diagnosis and response. *Journal of Drug Issues*, 16(3), 407-420.

Financial concerns and transportation problems are also barriers that affect women more often than men, if only because more of them are poor.¹²¹ Unemployment rates among drug-addicted women run as high as 80 percent;¹²² many rely on Medicaid, which treatment providers often do not accept.¹²³

Special Treatment Needs

Beyond removing barriers to access, successful treatment programs for women must address their psychological and social needs, such as childhood sexual abuse, eating disorders, domestic violence, depression, post-traumatic stress disorder, personal shame and lack of parenting skills.¹²⁴ Childhood sexual abuse can lead to distrust, depression, anxiety, shame and poor self-image, which can hinder recovery from addiction.¹²⁵ Female survivors of sexual abuse start using alcohol and drugs at younger ages and use them more to self-medicate and escape family problems than other girls.¹²⁶ Because they may continue to do so if they return to the same environment, addressing family problems in treatment may be essential to prevent relapse.¹²⁷ Similarly, screening for spousal violence is necessary not only because it affects how women respond to treatment, but because those who return to violent relationships tend to relapse.¹²⁸

Pregnant and parenting drug users, whose development may be arrested at age levels when their drug use began, often need parent education courses to teach them how to care for their children and to bolster their own recovery.¹²⁹ Many of these women lack adequate role models in their own parents and suffer low self-esteem and guilt over their substance use, which undermines relationships with their children. Often they have

unrealistic expectations for their children's cognitive and emotional development and for their ability to control their children's behavior.¹³⁰

Many female addicts also need basic medical care, which facilities for women frequently do not offer.¹³¹ Many lack basic skills needed to get jobs; pregnant drug abusers have been found to have an average of sixth grade math skills and seventh grade reading skills.¹³² In addition to treatment, these women need drug-free housing, education and job training.¹³³

Some researchers and treatment providers believe women do better in all-female settings. The environment may be more nurturing and supportive, and women may feel more comfortable speaking about issues such as domestic violence, sexual abuse and incest, shame and self-esteem in groups without men.¹³⁴ Female treatment providers can also serve as role models.¹³⁵ In co-ed settings, women are less likely to attend group therapy than men, and when they do, they are less likely to talk.¹³⁶

Founded in 1935, Alcoholics Anonymous (AA) initially was designed for men. As late as 1968, only 22 percent of its members were female.¹³⁷ Women for Sobriety was founded in 1976 as a program designed to address why females drink, with the belief that

Providing comprehensive services in one place may be necessary to retain women in treatment. A program at King/Drew Medical Center in Los Angeles found it nearly impossible to keep a substance abuser in both prenatal care and drug treatment if both services were not offered at the same location. When the Center incorporated prenatal care, primary care, parent-child programs and substance abuse treatment at one site, the savings from reduced prematurity among just 40 women were \$1 million.¹

¹ Haughton, J. G. (1992). The linkage: Substance abuse treatment and primary care. Paper presented at Strengthening the Linkages, a Secretarial Conference on Primary Care and Substance Abuse Linkage, Washington, DC; Haughton, J. G. (1993). Substance abuse prevention: A perinatal perspective. Paper presented at The Cost-benefit of Substance Abuse Prevention, Washington, DC.

traditional mutual-help groups such as AA inadequately served women.¹³⁸ During the 1970s and '80s, representation of women in AA grew and women-only AA groups spread across the country.¹³⁹ By 1992, at least 35 percent of all AA members were women.¹⁴⁰

Researchers have recently begun to investigate what treatment best suits women.¹⁴¹ One study, which compared women in a female-only program to women in a coed program, found those in the all-women group did better after 2 years on several measures: fewer deaths, less alcohol consumption, less need for inpatient care due to relapse, higher job stability, better relationships with children and maintenance of child custody.¹⁴² At the Betty Ford Center, women in their female-only program were found more likely to remain sober for 12 months than women in coed treatment programs.¹⁴³ After this evaluation, the Center discontinued coed treatment and moved all women into the female-only program.¹⁴⁴

Confrontational therapy programs, which aim to push addicts to shed their denial and assume responsibility for their behavior, may backfire on women by reinforcing feelings of shame, low self-esteem and depression.¹

¹ O'Connor, L. E., Berry, J. W., Inaba, D., Weiss, J. and Morrison, A. (1994). Shame, guilt, and depression in men and women in recovery from addiction. *Journal of Substance Abuse Treatment*, 11(6), 503-510; Burman, S. (1994). The disease concept of alcoholism: Its impact on women's treatment. *Journal of Substance Abuse Treatment*, 11(2), 121-126; Quinby, P. M. and Graham, A. V. (1993). Substance abuse among women. *Primary Care*, 20(1), 131-140; Tracy, C. E. and Williams, H. C. (1991). Social consequences of substance abuse among pregnant and parenting women. *Pediatric Annals*, 20(10), 548-552; Watson, R. R. (Ed.). (1994). *Addictive behaviors in women*. Totowa, NJ: Humana Press; Wells, D. V. B. and Jackson, J. F. (1992). HIV and chemically dependent women: Recommendations for appropriate health care and treatment services. *International Journal of the Addictions*, 27(5), 571-585.

Quitting Smoking

More than 80 percent of adults who try to quit fail on their first attempt, and more than 50 percent fail on their second.¹⁴⁵ While overall quit rates are nearly equal for women and men, women appear less likely to keep trying or stay with it.¹⁴⁶ Among those who had previously quit for at least a week on three or more occasions, only 24 percent of female smokers planned to try again within a year, compared to 61 percent of male smokers.¹⁴⁷ Up to 90 percent of women who quit smoking while pregnant relapse within a year of giving birth.¹⁴⁸

Women's fear of weight gain is a major reason for failing to quit.¹⁴⁹ In a survey of young adults (mean age 19.2), female smokers were twice as likely as male smokers to be concerned about gaining weight if they quit (57.9 percent vs. 26.3 percent).¹⁵⁰ Among quitters, more women than men reported wanting to eat more than usual (47.8 percent vs. 26.2 percent), and more women reported gaining weight (29.5 percent vs. 19.2 percent).

Weight concerns may be particularly strong

after a woman gives birth.¹⁵¹ One study found that postpartum women who resumed

One woman on her struggle to quit smoking: "I have quit smoking about five times, only to gain weight. As a result, I have returned to smoking cigarettes. On November 1, 1990, deciding once again to quit, I joined a weight-loss clinic. I was determined to diet vigorously. By January 1, 1991, I had gained 12 pounds. I am faithful to my diet and to my exercises, and I am most unhappy to say that I have not lost one pound. I am on the verge of resuming smoking even though it is truly not what I want to do. I am vain, my clothes don't fit, my self-image is poor and I think that no matter what I do I will not lose weight until I begin to smoke again."¹

¹ Shapiro, F. (1991). Correspondence: Smoking cessation and severity of weight gain. New England Journal of Medicine, 325(7), 517.

smoking were more likely to believe they would return to their former weight within six months of delivery than those who did not start smoking again.¹⁵²

Accepting the near inevitability of some weight gain may be an important step towards quitting.¹⁵³ Women who quit tend to gain more weight than women who relapse, probably because their tolerance of some extra weight allows them to persist in cessation.¹⁵⁴

High smoking rates and low quitting rates are also associated with depression, which is more common among women.¹⁵⁵ Individuals who are depressed are almost three times more likely to smoke, while depressed smokers are half as likely to quit as other smokers.¹⁵⁶ Stress is also cited as a factor in whether individuals start smoking, how much they smoke and whether they quit--despite the lack of evidence that smoking reduces stress.¹⁵⁷ This has led to the belief that smokers--particularly women--see cigarettes as a way to reduce stress and cope with nervousness, anxiety and emotions.¹⁵⁸

While doctors and other advisers to female smokers can suggest ways to minimize weight gain and ease stress, they can also underline that some weight gain is probably inevitable--and decidedly less important than the enormous health benefits of quitting.¹⁵⁹ Women assigned to a cessation program, plus three sessions of exercise per week, were found five times more likely to be smoke-free a year later than women smokers who only used the cessation program.¹⁶⁰ In another study, female smokers who quit exercised more than those who relapsed--even though the exercise did not translate into less weight gain.¹⁶¹ Nicotine gum may also help women trying to minimize weight gain.¹⁶² Even when the

weight control component has no impact on weight gain or cessation, it may make the challenge of quitting more palatable to women.¹⁶³

Using the Law

No aspect of women's substance use or abuse has generated more attention, concern, anger, outrage and controversy than prenatal drug use. Some public prosecutors have prosecuted women who test positive for drugs when they enter a hospital to deliver a baby. Some use the threat of jail or the loss of child custody to try to push women into treatment.¹⁶⁴

Prosecutors have used drug test results as evidence to charge women with child abuse, delivering drugs to a minor through the umbilical cord and drug possession.¹⁶⁵ Since 1985, at least 200 pregnant and postpartum women in 30 states have been criminally charged under these laws.¹⁶⁶

In 1995, as part of this CASA study, the New York City-based law firm Cadwalader, Wickersham and Taft completed a comprehensive, nationwide review of state statutes, case law and pending legislation pertaining to the legal issues surrounding substance-using pregnant or postpartum women. CASA updated this review in February 1996. Together this work found 35 reported decisions in 20 states stemming from the charges brought against such women since 1985.^{*167}

* States have prosecuted many more women; these were the only reported cases found. Most cases are not reported. Others may have been dismissed, may be in the process of adjudication or may involve indictments for an unrelated crime.

Among the 35 decisions, 20 women were tried for child abuse and neglect (including criminal neglect and endangerment), eight for delivery of drugs to a minor, six for possession and one for both delivery and possession.¹⁶⁸

Of the 20 child abuse cases, 12 were dismissed. Six defendants were found guilty, but these convictions were overturned. One woman pled guilty to child abuse and was sentenced to six months in jail. One was allowed to enter a treatment program in lieu of incarceration.¹⁶⁹

Of the eight cases of women charged with delivery of drugs to a minor, five were dismissed. One conviction was overturned. Two women entered no contest pleas and were sentenced (one to 18 months probation; the other to 18 months in jail and three years probation).¹⁷⁰

Of the six cases of women charged with possession, four were dismissed. One conviction was overturned. One woman pled guilty and was sentenced to 150 days in jail, five years probation and a \$225 fine.¹⁷¹

In the one case of the woman charged with delivery and possession, both charges were dismissed.¹⁷²

Teresa Reinesto was charged with child abuse after giving birth to a baby who tested positive for heroin and experienced withdrawal. The court ruled that her prenatal conduct did not fall within the "plain language of the [child abuse] statute.... Allowing the state to define the crime of child abuse according to the health or condition of the newborn child would subject many mothers to criminal liability for engaging in all sorts of legal or illegal activities during pregnancy." The court concluded, "We know of no other jurisdiction that has upheld a charge of child abuse against a woman for conduct that harmed her fetus, even if the resulting condition affected the child after birth."¹

¹ Teresa Lopez Reinesto v. Superior Court; 182 Ariz. 190; 894 P.2d 733, 1995.

In addition to prosecuting women criminally, child abuse has been alleged in civil proceedings. In these settings, the issue of whether prenatal drug use--as evidenced by a positive urine test at birth--is sufficient to prove child abuse and neglect in the absence of other evidence is hotly contested. While no state statute defines prenatal drug use as criminal child abuse or neglect, in eight states evidence of drug exposure in a civil case classifies a child as neglected or at risk of being neglected.¹⁷³ In at least 22 states and Washington D.C., prenatal drug use is considered relevant in civil child abuse or neglect cases.¹⁷⁴

The review of these cases reflects the difficulty of resorting to existing drug laws to deal with the frightful problem of illegal drug use during pregnancy and the controversy over the appropriateness of criminally prosecuting women who use such drugs while pregnant.¹⁷⁵ There's complete agreement on the objective: get the pregnant woman off drugs. Society struggles to "do something" to stop women from using illegal drugs during pregnancy. There is a desire to punish those who use drugs for reckless failure to fulfill their responsibility as childbearers; yet the difficulty of shaking addiction to drugs like crack cocaine and heroin is enormous. Abortion rights activists want to avoid any actions or

In 1991, Dianne Pfannestiel, who was pregnant, went to an emergency room out of concern for her fetus after her partner beat her severely. She was arrested and charged with child abuse for drinking while pregnant. No charges of child abuse were brought against her partner.¹

¹ Chavkin, W. et al. (1994). Finding common ground: The necessity of an integrated agenda for women's and children's health. Journal of Law, Medicine and Ethics, 22(3), 262-269.

* Drug exposed children are defined by states as follows: neglected or abused (Illinois and Kentucky); disabled (Alaska); harmed (Florida); at biological or environmental risk (Hawaii); in need of services (Indiana); in need of assistance (Iowa); deprived (Oklahoma).

policies that might legally recognize rights of the fetus, because that might inhibit a woman's freedom to have an abortion.

Right to life groups want the mother held responsible for any damage to the fetus, which they regard as the most vulnerable and innocent expression of human life.

Because so many of these women are black, many African-Americans charge that the exercise of prosecutorial discretion in these cases is tinged with racism. The issue is further complicated by the lack of readily available treatment for pregnant mothers and decent foster care systems and orphanages for placing children of drug abusing mothers in drug-free environments.

In 1988 Brenda Vaughan, who was pregnant, spent nearly four months in jail for forging \$700 worth of checks.

Though the prosecutor had recommended probation, the judge imposed a sentence to last the duration of her pregnancy because Vaughan had been using cocaine and he wanted to protect the fetus by making sure she gave birth in prison. While in prison, Vaughan received no prenatal care or drug treatment.¹

¹ Inciardi, J. A. et al. (1993). Women and crack-cocaine. New York: Macmillan Publishing Company; Garcia, S. A. (1993). Maternal drug abuse: Laws and ethics as agents of just balances and therapeutic interventions. International Journal of the Addictions, 28(13), 1311-1339; Krauss, D. (1991). Regulating women's bodies: The adverse effect of fetal rights theory on childbirth decisions and women of color. Harvard Civil Rights-Civil Liberties Law Review, 26(2), 523-548; Weber, E. M. (1992). Alcohol- and drug-dependent pregnant women: Laws and public policies that promote and inhibit research and the delivery of services. In M. M. Kilbey and K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children, NIDA research monograph 117 (pp. 349-365). Rockville, MD: NIDA.

Foster Care: The Era of Orphan Annie is Over

Americans may think of Orphan Annie and Daddy Warbucks when they think of foster care. But in the 1990s, foster care is a system to salvage children of drug and alcohol-abusing parents, which raises profound moral, ethical and legal questions about what is best for these children.

The current caseload of drug-exposed children has pushed foster care systems in cities and counties across the country to their limits and beyond, leading in some cases to children's deaths.¹⁷⁶ During the late-1980s, the number of children in foster care skyrocketed, largely as a result of drug and alcohol abuse.¹⁷⁷ The average age of children in foster care has declined, with the mother's use of crack cocaine before and after birth a primary cause.¹⁷⁸

Child welfare administrators say that scarce resources and insufficiently trained staff are preventing adequate investigation of many cases.¹⁷⁹ The need for training and support in making judgment calls related to substance abuse is acute.¹⁸⁰ In New York City, a family court judge often decides 40 to 50 cases each day.¹⁸¹

In some states, social service agencies are required by law to make every effort to preserve families.¹⁸² Only if parents abandon a child or demonstrate a chronic inability to care for them will a family court sever parental rights, making the child eligible for adoption.¹⁸³ But such proceedings often take at least three years to complete.¹⁸⁴

Children in foster care may be caught in limbo if their parents are not capable of caring for them, but have not done anything so severe as to warrant termination of parental rights.¹⁸⁵ In these cases, particularly if the woman refuses to enter treatment, some argue for easier and earlier termination of parental rights to get children into adoptive homes that may be able to compensate for any prenatal damage due to drug exposure.¹⁸⁶

Some public officials and prosecutors are trying to make a positive drug test at birth sufficient evidence to justify removing children from their families and placing them in foster care.¹⁸⁷ To date, judges have based custody decisions solely on prenatal drug use in at

least 22 reported cases; in 16 of them, evidence of prenatal drug use was adequate to terminate parental rights and in 6 cases it was not.¹⁸⁸

This shift away from keeping the child with the parents and avoiding foster care has been sharpened by the advent of crack cocaine. While the goal of family preservation has obvious appeal, the consuming power and destructive consequences of crack addiction have created concern that efforts to keep the family together can consign the child to a home of neglect, chaos, danger, violence and even death.

On the other hand, the chance to keep their children appears to be a strong motivator for many women to seek treatment and pull their lives together.¹⁸⁹ And the well-being of their children and their desire to be with them are often key reasons why addicted women seek treatment.¹⁹⁰

The question of what would best serve the child raises profound questions about who should care for them: Should a family court judge ever decide that an alcoholic or drug-addicted mother can care for her children, especially given the high rate of relapse that is characteristic of addiction? Is foster care or an orphanage a better place for the children? Are adoptive homes better for children than parents with drug or alcohol problems? How do you weigh the emotional costs to the child of separating him or her from parents and family? What kinds of investigations by case workers are required to make such delicate judgments?

Often, politicians, policy-makers and prosecutors must answer these questions under the harsh spotlight of intense media coverage and public demands for quick solutions that follow the death of an innocent child at the hands of a drug-crazed or drunk parent. Public costs also mount. Family preservation services cost less than half the bill for foster care

placement, with estimates ranging from \$3,000 to \$7,000 per family per year compared to \$10,000 to \$17,500 per child in foster care per year. But support for family preservation attempts has waned as doubts persist about its appropriateness in situations of parental drug and alcohol abuse.¹⁹¹ Crack cocaine, alcohol and other drugs have shattered old assumptions about parental rights and the sanctity of the family as society struggles to protect the children.

Mandatory or Coerced Treatment

Whether as an alternative to criminal prosecution or to the loss of child custody, mandatory treatment is one course of action increasingly in use. When faced with the prospect of incarceration or losing her children, the hope is that the drug-addicted mother will instead choose treatment. Because a woman could opt for jail, this policy is often called "coerced treatment."

Coerced treatment works as well as voluntary treatment in retaining men and women in treatment programs, an important predictor of success.¹⁹² But reports vary on the effectiveness of coerced treatment for pregnant or parenting women.¹⁹³ The lack of research on the efficacy of mandatory or coerced treatment makes it impossible to draw any conclusions.¹⁹⁴

The shortage of treatment renders much of this policy debate moot. The use of civil commitment for female addicts, in which judges assign individuals to treatment because they are considered a threat to themselves or someone else's safety, floundered in Massachusetts during the 1980s because of lack of available treatment.¹⁹⁵ Civilly-committed female

addicts were sent to prison until public outcry prompted the creation of more treatment slots for them; however, demand for treatment still exceeds supply.¹⁹⁶ New Mexico and Oklahoma make civil commitment contingent on the availability of appropriate treatment. Georgia, Hawaii, Minnesota and West Virginia require confinement in the "least restrictive alternative" treatment environment.¹⁹⁷

The shortage of treatment in prison is a particular concern in light of the rise in incarceration of pregnant addicts. Pregnant woman in prison often get no drug or alcohol treatment and little prenatal care. Not surprisingly, their rates of miscarriage are high.¹⁹⁸ In one California county, only 20 percent of pregnant inmates delivered live babies. The only drug treatment program for female inmates in the California state prison system serves 120 women and has a waiting list of 70 to 90.

Driving the lack of support for treatment are doubts about its effectiveness.¹⁹⁹ While research on treatment outcomes is desperately needed, there is evidence that treatment of pregnant drug users can improve the short-term health of the woman and child and reduce health care costs. Pregnant, drug-using women in treatment have been found to have more

In 1991, Jennifer Johnson was convicted of delivering drugs to a minor through the umbilical cord in the seconds after giving birth. She had tried to get treatment, but could not find a program that would admit her. Her conviction was unanimously overturned by the Supreme Court of Florida, which ruled that the drug trafficking law was not intended for use against a woman for giving birth to a drug-exposed baby.¹

¹ Chavkin, W. et al. (1990). Drug-using families and child protection: Results of a study and implications for change. *University of Pittsburgh Law Review*, 54(1), 295-324; Moss, K. L. (1991). Forced drug or alcohol treatment for pregnant and postpartum women: Part of the solution or part of the problem? *New England Journal on Criminal and Civil Confinement*, 17(1), 1-16; Hansen, M. (1992). Courts side with moms in drug cases. *ABA Journal*, 78, 18.

negative urine tests, infants with higher birth weights and larger heads, and more prenatal care than women who did not receive treatment.²⁰⁰

Two small studies of pregnant women in methadone maintenance for opiate addiction found that enhancing methadone programs with other services may improve the newborn's health.²⁰¹ Women on methadone who received weekly medical check-ups, group therapy to prevent relapse, rewards for clean urine screens and child care during treatment, in addition to standard services, had heavier infants and fewer positive urine screens than women in standard methadone treatment involving only daily methadone, group counseling and random urine screens.²⁰² In the other study, women with enhanced treatment had longer gestations and heavier babies, but not significantly fewer positive urine screens.²⁰³

Pregnancy: A Window of Opportunity

Pregnancy opens a window of opportunity when many female addicts can be motivated to enter treatment and stop using cigarettes, alcohol and illicit drugs.²⁰⁴ Indeed, most pregnant women and teenagers at least reduce their licit and

Dee Cantie gave birth to a baby boy in 1994 with cocaine in his blood. Four months later, she was pregnant again. She dismissed the fathers as flings; she was married to the pipe. Her second boy was born premature with narcotics in his blood. She cried in the delivery room, realizing the addiction was no longer just her problem. The devil was on her back. "It was like a snowball," she said. "I had guilt, and I would use more. The more I used, the more I had to have." Officials threatened to take her children, so she checked into detox and, later, a day treatment center designed to help women with felony arrests get clean and put their lives in order. The sober life, a "normal" life, is a strange existence for her. "It's boring sometimes, but I can live with it. It's better than dying." And she adds, "It's about the kids, man. Without their mother, what are they going to do?"¹

¹ LeDuff, Charlie. Cocaine daze. New York Times, 3/10/96, Sect. 13, p. 1.

illicit drug use while pregnant.²⁰⁵ The benefits of prenatal care are even greater for HIV-infected women because medical treatment can cut the chances that they will transmit the virus to the fetus by 68 percent.²⁰⁶

Yet some experts caution that while pregnancy appears to motivate many women to enter treatment, it also aggravates feelings of shame and guilt, which may discourage women from seeking help.²⁰⁷ Encouraging women to quit for both themselves and their future children may be more effective.²⁰⁸

The threat of prosecution or swift loss of child custody if a woman fails a drug test complicates this message. Some experts warn that women who fear having to submit to a drug test will stay away from prenatal care altogether.²⁰⁹ But sufficient research has not been done to resolve the issue.²¹⁰

Health care providers complain that having to report the results of drug tests, particularly those done prenatally, compromises their vow of confidentiality to the patient, hinders the candor and trust they require from patients, and places them in a role of law enforcement rather than care giver.²¹¹ However, the dangers of leaving a child with a drug-addicted or alcoholic parent has punctured old assumptions about the inviolability of the doctor-patient relationship in this situation.

One woman, charged with delivering drugs to a minor when she was pregnant, explains how getting help for a drug problem backfired: "I'm a perfect example of someone who tried to reach out, and it's all coming back in my face.... Everyone I talked to about my drug problem has been subpoenaed."¹

¹ Krauss, D. (1991). Regulating women's bodies: The adverse effect of fetal rights theory on childbirth decisions and women of color. Harvard Civil Rights-Civil Liberties Law Review, 26(2), 523-548.

Before the crack epidemic of the 1980s, hospitals did not systematically screen women or babies for drugs, either prenatally or at the time of delivery.²¹² If health care providers discovered a pregnant woman had used drugs, the response could range from ignoring it to referring her to treatment.²¹³ Since the advent of crack, hospitals have tested more frequently, most often at birth.²¹⁴ If a woman or newborn screens positive for illegal drugs at delivery, hospital clinicians generally report the mother to child protective services as a matter of hospital policy, even when not required by law.²¹⁵

In a survey of 49 hospitals in the Chicago metropolitan area, 94 percent always report a positive drug test to the child protection agency.²¹⁶ A Minnesota law requires physicians to test for drugs if they suspect substance abuse and report positive results to a child protective service agency.²¹⁷ In Florida, after a baby is born, any individual who learns that a mother has used drugs while pregnant must report her to the Florida Protective Service Systems' statewide child abuse registry, triggering an investigation of the infant's home.²¹⁸ If the home is deemed safe, case managers periodically visit to teach parents the skills to care for their child; if not, the agency begins legal proceedings to remove the child from the home.²¹⁹ Few women reported to the Florida register are offered treatment.²²⁰

Drug testing is far from universal. It is more common at public rather than private hospitals. Most look for certain indicators that trigger a drug test, which limits the number of women tested but sometimes sparks charges of selection bias and discrimination.²²¹ Adding to the controversy is the fact that some hospitals test without the mother's knowledge or consent.²²²

In the Chicago hospital survey, hospital personnel cited low income, nationality or ethnicity, and place of residence as cues for testing.²²³ In another study in a midwestern suburb, nurses were more likely to suspect that women who were patients of hospital physicians--a larger proportion of whom were black and publicly insured--would have a positive drug test than the mostly white, privately insured women who were patients of private physicians, even though the difference in positive drug tests between the two groups was not statistically significant.^{*224}

The thorny issues that surround testing and reporting of drug use during pregnancy add to the complexity of the moral, ethical and legal questions raised by women's substance abuse and addiction--questions our nation has not considered sufficiently.

* A study in Pinellas County, Florida found that, despite similar rates of illicit drug use, African American women were 10 times likelier to be reported to the Florida Department of Health and Rehabilitative Services than white women.

VI.

Next Steps:

Acting on What We Know and Building Knowledge

The first step is to act on what we know. The second is to learn a lot more. The foundation of solid prevention and treatment efforts is research.

Since the 1960s, when only a handful of researchers was devoted to the problem, our understanding of how women's patterns of use differ from men's, the severe health and social consequences women suffer and the unique dangers that arise during pregnancy has come a long way. The current base of knowledge, if put to use, can make a big difference.¹

Research and experience have taught us three critical lessons:

- Substance abuse and addiction among women is a major public health problem.
- If prevention and treatment efforts for women are to be effective, they must take into account their special needs.
- Most women with substance problems use a combination of alcohol, tobacco, and licit and illicit drugs.

With these lessons in mind, politicians and policy-makers, doctors and nurses, educators and parents, clergy and teen counselors--can mount an attack on this problem before another generation of women and their children are struck by its deadly consequences. For the knowledge gaps that remain, both basic and applied research is necessary. In particular, research, prevention and treatment efforts should focus on:

◆ **Smoking and weight loss.** As the number of women who die from smoking rises dramatically, the need for effective prevention and cessation strategies for female smokers becomes more urgent. Medical visits provide an ideal setting for primary prevention efforts. Doctors who routinely incorporate advice on smoking into prenatal and other medical visits increase the odds that a patient will quit.²

Yet we still know too little about how to break the strong link between weight concerns and smoking among women during adolescence, young adulthood and pregnancy.³ Preventive efforts must not only attempt to deglamorize smoking, but also include a strategy to put in perspective the cultural bias toward extreme thinness for women. More research is needed to examine the relationship between weight concerns and illicit substances, such as cocaine and methamphetamine, that suppress appetite.

◆ **The health consequences of substance use and abuse.** Failing to include women in most research on substance use and abuse has left health care and public health professionals, policy-makers and women themselves without the knowledge they need to prevent or treat the problem.⁴ Prevention efforts in doctor's offices, schools, businesses and community centers need to educate women about their greater susceptibility to the effects and health consequences of alcohol and illicit drugs.

Researchers need to investigate remaining questions about the consequences of alcohol, tobacco and drugs on women. The effect of alcohol on a woman's endocrine function, for example, is an important area for research, given evidence that consumption at levels many women consider "social drinking" might create serious health problems for women.⁵

Understanding the extent and nature of women's greater physiological vulnerability to the physical damage of alcohol, drugs and tobacco is also key.⁶ Preliminary research has suggested that women who smoke may be more susceptible to lung cancer than men who smoke.⁷ Investigating why women may become addicted to illicit drugs more quickly than men could also yield important results.

◆ **Substance abuse among young adult women.** The lack of a decline in heavy drinking among younger women during the 1980s, the popularity of binge drinking by college women and evidence suggesting that women's drinking patterns often solidify from ages 21 to 34 indicate that researchers should give greater attention to this time in a woman's life.⁸

The stresses and benefits of employment in young women's lives should be examined. To frame prevention and treatment efforts, researchers need to look at the relationship between alcohol and drug use and a woman's entry into the job market, how much she wants to work, how much support she receives from her spouse and how she perceives the occupational culture with regard to the sexes and drinking norms.⁹

One specific target for prevention efforts is the college campus, where many women drink heavily.¹⁰ Since the link between high risk sex and alcohol is so strong, information about the health and other consequences of drinking for women might effectively be presented along with information about safe sex practices.¹¹ Women also need to understand the relationship between alcohol use and violence against women on campus.

◆ **Older women.** Even though women today live a third of their lives after menopause, little research on female substance abuse focuses on this phase.¹² The

frequency of late onset of alcohol problems, often in combination with prescription drug misuse, among women after menopause and in retirement makes studying the prevalence, risk factors and consequences of substance abuse among them necessary to developing effective prevention and treatment strategies.¹³ Widowed women are a particularly important target for prevention, since they are three times more likely than married women to drink heavily.¹⁴

◆ **Prenatal exposure.** Prenatal visits provide physicians and nurses with an opportunity to identify substance abusers, encourage them to quit or reduce the frequency of their substance use, discuss "triggers" for substance use and refer dependent patients to treatment. Quitting or cutting down during pregnancy can reduce the adverse effects of prenatal substance abuse.¹⁵ However, the majority of addicted pregnant women do not receive prenatal care; they may require a concerted outreach effort, as well as transportation and follow-up services.

Research to assess the long-term effects of prenatal exposure to alcohol, tobacco and illicit drugs is critical in order to bolster prevention efforts and design interventions to help children overcome damage done during pregnancy.¹⁶ The Administration for Children, Youth and Families, the National Institutes of Health (National Institute of Child Health and Human Development, and National Institute on Drug Abuse) and the Substance Abuse and Mental Health Services Administration (Center for Substance Abuse Treatment) have funded a study to compare the birth outcomes of some 3,000 drug-exposed infants and 13,000 newborns whose mother did not use drugs, alcohol or tobacco during pregnancy.¹⁷

The 5-year study, begun in 1993, will also track the neurobehavioral development of 4,000 infants (half exposed, half non-exposed) for three years, taking into account the role of parenting and environmental factors.¹⁸ Other research is also needed, based on current knowledge, to design and test effective interventions for children with fetal alcohol syndrome and other disabilities caused by substance use during pregnancy.¹⁹

◆ **Child abuse and neglect.** Substance abuse is a pervasive factor in cases of child abuse and neglect. Protecting a woman's right to privacy and the confidentiality of the doctor-patient relationship is important. But the fact that newborns of drug- and alcohol-abusing parents are at high risk of child abuse and neglect makes it critical to identify these children and get their parents into treatment for the safety and welfare of the children. Indeed, pregnancy is a window of opportunity when a women may be especially likely to seek treatment for her own sake and that of her child.

States need to look at how they are handling children of drug-abusing mothers, including a review of the systems of treatment, adoption, foster care and orphanages. Rampant drug and alcohol abuse raises questions about assumptions taken for granted years ago.

◆ **Multiple drug use.** Combined abuse of tobacco, alcohol and other drugs is the norm among alcoholic and drug-addicted women.²⁰ In one study, 82 percent of alcoholic women vs. 34 percent of non-alcoholic women were current smokers.²¹ Separate funding agencies for alcohol research and drug research at the federal level, as well as distinct traditions and funding streams in the treatment community, have perpetuated a single-drug approach to research and clinical practice and thwarted coordinated responses to drug

epidemics. This approach ignores the pervasiveness of multiple substance abuse by women and renders single-drug strategies obsolete when new drugs come into fashion.²²

Different drug and alcohol combinations, such as alcohol and prescription drugs, or alcohol and cocaine, appear to characterize different populations of women. Research can help decipher the patterns and effects of multiple drug use, with important implications for targeted prevention and treatment.²³

Use of a combination of alcohol, illicit drugs and/or tobacco is also the norm among substance-using pregnant women. While some progress has been made to disentangle the relative impact of these drugs, more research is needed to understand the mix, dose and timing of the tobacco, alcohol and drugs that most damage the fetus. The combination of alcohol and cocaine appears to be more severe than either on its own, though little research has focused on this phenomenon and its implications for prevention and treatment.²⁴

Doctors may unwittingly encourage multiple drug use among women. Since female alcoholics are more likely than males to suffer depression, doctors may prescribe drugs for a woman's depression but fail to recognize her substance abuse.²⁵ Education about this potential danger and training in substance abuse recognition would help doctors avoid this mistake.

◆ **Racial and ethnic differences.** Studies should pay more attention to differences among women from racial and ethnic minorities and various income and educational backgrounds.²⁶ For example, white women are more likely to use drugs and alcohol, but black women are more likely to be regular or heavy users. Studies have found that while the rate of alcohol problems usually peaks for white women in their 20s, it continues to rise for

black women into their 40s and 50s, which suggests that the origin and course of these problems may be quite different.²⁷ Black women are also less likely than either white women or black men to quit smoking, revealing a need for cessation efforts that focus on black women.²⁸ Prevention programs targeting this group of smokers should challenge misconceptions about the relative safety of menthol cigarettes.²⁹

Racial and ethnic differences also surface in attitudes toward treatment. Physicians who detect the signs of a drinking problem in an Hispanic or Asian American woman should be aware that a family's denial or desire to "protect" the woman by keeping her problem a secret may thwart her motivation to seek treatment.³⁰

Most studies of African American women have concentrated on low-income females in urban America.³¹ This focus has highlighted barriers to treatment for low-income women: many can afford neither transportation nor child care, and they need job training to help them support themselves after they are substance-free. However, as a result of this focus, we know little about drinking patterns, alcohol abuse and addiction among women in the black middle class.³²

◆ **Treatment: What works?** Not enough research has been done on treatment outcomes for women.³³ To begin closing this gap, the National Center on Addiction and Substance Abuse at Columbia University is conducting an extensive study of 2,000 individuals at 200 treatment sites nationwide that will provide urgently needed data on what works for whom and why, including important insights by gender. CASA is also assessing the effectiveness of acupuncture as a treatment for cocaine addiction, and plans to include a sample of pregnant women. Furthermore, the National Institute on Drug Abuse funded 20

demonstration projects that treat pregnant and parenting women, called the Perinatal-20 projects, which have provided important data on what works for pregnant and postpartum women.³⁴

More work is needed to explore the influence of genetic and environmental factors on substance abuse by women. Although research on alcoholism to date is inconclusive, family and other external factors seem to have a more significant impact than genetics. Developing a clearer understanding of the role these factors play is an important step in crafting prevention and treatment strategies for women.³⁵

Additional investigations should focus on the role in treatment of addressing particular issues for women, such as the high rates of domestic violence, sexual abuse, eating disorders, primary depression, suicide attempts and family histories of substance abuse among female alcoholic and addicts. Women may be more comfortable discussing such personal issues in groups without men. Those who self-medicate to cope with these problems, past and present, should be taught alternative coping strategies and methods for building self-esteem.³⁶

◆ **Prevention strategies and the special role of doctors.** Though the focus on drug and alcohol use during pregnancy is understandable and important, primary prevention of damage to both mother and fetus requires avoiding substance use before conception, which means focusing on women of child-bearing age and younger. Doctors, nurses and other health professionals have an important role to play here, and research can help them by testing the effectiveness of various intervention strategies in the setting of a hospital, clinic or doctor's office.³⁷ The fact that women are more likely than men to visit a doctor, but less

likely to have a doctor diagnose their substance abuse disorders, suggests that physicians are missing many opportunities to refer women to treatment.

Effective prevention strategies require an understanding of the origins of the problem. More research is needed to assess the importance of the risk factors that characterize women and differ so significantly from those of men. A priority should be determining which risk factors are causal. Evidence that women and men respond differently to prevention messages underlines the importance of targeted interventions.³⁸

Doctors and school nurses also have an opportunity to detect the risk factors and early signs of substance abuse among teenage girls. Preventing substance abuse among adolescent girls is a challenge our nation cannot afford to neglect. Based on everything we know, if we can get our children to age 21 without smoking cigarettes, abusing alcohol or using illicit drugs, they are virtually certain never to do so. Research is needed to inform efforts that aim to seize this opportunity.³⁹

◆ **Measuring the problem.** Tools to detect substance abuse tend to rely on quantity consumed by men. These instruments should adjust for the lower quantities of alcohol and drug use that characterize women with alcohol- and drug-related problems and include more women-specific concerns, such as how their drinking is affecting their health and close relationships.⁴⁰

Traditional measures of alcohol-related problems focus on those that are typically male: drinking-related fights, drunk driving, alcohol-related arrests, financial problems and job impairment.⁴¹ While women appear to be experiencing these problems at an increasing rate, researchers and doctors trying to detect substance abuse problems among their female

patients solely with these measures will miss many cases and underestimate the scope of the problem.

Characteristics frequently associated with substance abuse among women, such as depression, anxiety and family violence, should alert physicians to the possibility of substance abuse. Doctors can use a screening tool known as the CAGE questionnaire, as well as the following questions, to elicit additional information: "Do you ever carry an alcoholic beverage in your purse? How has your drinking changed during pregnancies?"⁴² For elderly women, such questions might include: "Did you find your drinking increased after someone close to you died?"⁴³

The Double-edged Sword of Stigma

Too often the public response to women's substance abuse and addiction has vacillated between denial or benign neglect on one hand and disdain or scorn on the other. Often the stigma of women's substance abuse and addiction drives our thinking and hinders our ability to grasp and fight the problem effectively. The shame and secrecy that surrounds this issue pose a special challenge for treatment providers, who require an open and honest dialogue with their clients.⁴⁴

Yet stigma is a double-edged sword.⁴⁵ The cultural disapproval attached to female substance abuse may have a protective effect for women by discouraging them from taking the risk of using drugs and abusing alcohol or continuing to do so once they have experimented. This deterrent could account for a significant portion of the remaining gap between women and men's drinking and drug use.⁴⁶ Even as rates of drinking and drug use

among teenage girls and boys converge, girls are more likely than boys to "age out" of problem drinking or frequent drug use as they reach young adulthood, perhaps because they experience pressure to conform to gender expectations.⁴⁷ In a study of 31 women married to heavy drinkers, a woman's belief that such drinking among females is acceptable only in the teen and young adult years and at special occasions significantly reduced the likelihood that she would drink heavily.⁴⁸ Other research has found that women who support the goal of gender equality drink more, suggesting that freedom from gender-related expectations may prompt heavier use.⁴⁹

At the same time, the negative consequences of stigma are formidable.⁵⁰ The sharp social disdain that female addicts face can dampen their will to confront their problem in a public light that is very harsh.⁵¹ Often this disdain comes from within. One study found that 51 percent of alcoholic women and 36 percent of non-alcoholic women agreed that an intoxicated woman is "more obnoxious and disgusting" than an intoxicated man.⁵² More than half the women in the study believed that a mother's alcoholism is more damaging to children than a father's alcoholism.⁵³

Reducing the paralyzing consequences of stigma could reap substantial benefits. Women and their families would more readily recognize and acknowledge a substance abuse problem in its early stages and seek help if necessary.⁵⁴ Social stigma would no longer aggravate the debilitating lack of self-esteem that women must surmount to overcome their addiction.

Effective prevention and treatment of substance abuse and addiction among women requires eroding the special stigma of the female alcoholic and addict without suggesting that

the risks of smoking, alcohol abuse and illicit drug use among women and men are alike. As one expert points out, "The challenge is to preserve the protection afforded women by cultural expectations that they will drink less than men, while changing inaccurate perceptions about alcohol effects on sexual responsiveness," such as women's supposed promiscuity and consequent moral plunge.⁵⁵

With preventive education about women's greater susceptibility to the consequences of alcohol and drugs and the potential harm to the fetus of such use during pregnancy, we can preserve cultural attitudes that maintain differences in female and male drinking norms and fully inform women about the risks they take when they smoke cigarettes, abuse alcohol and use drugs.⁵⁶ For women, the path to equality may offer equal access to the tragic consequences of substance abuse and addiction. But they do not have to go as willing victims. With effective prevention strategies, they do not have to go at all.

CHAPTER I.

REFERENCES

1. Wilsnack, S. C., Wilsnack, R. W. and Klassen, A. D. (1984/1985). Drinking and drinking problems among women in a U.S. national survey. Alcohol Health and Research World, 9(2), 3-13; Midanik, L. T. and Clark, W. B. (1995). Drinking-related problems in the United States: Description and trends, 1984-1990. Journal of Studies on Alcohol, 56, 395-402; Midanik, L. T. and Clark, W. B. (1994). The demographic distribution of U.S. drinking patterns in 1990: Description and trends from 1984. American Journal of Public Health, 84(8), 1218-1222; Penniman, L. J. and Agnew, J. (1989). Women, work and alcohol. Occupational medicine, 4(2), 263-273; Reich, T., Cloninger, C. R., Van Eerdewegh, P., Rice, J. P. and Mullaney, J. (1988). Secular trends in the familial transmission of alcoholism. Alcoholism: Clinical and Experimental Research, 12, 458-464; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press; Taylor, M. E. and St. Pierre, S. (1986). Women and alcohol research: A review of current literature. Journal of Drug Issues, 16(4), 621-636.
2. Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C. B., Hughes, M., Eshleman, S., Wittchen, H. and Kendler, K. S. (1994). Lifetime and 12-month Prevalence of DSM-III-R Psychiatric Disorders in the United States: Results from the National Comorbidity Survey. Archives of General Psychiatry, 51(5), 8-19.
3. Williams, G. D., Grant, B. F., Harford, T. C. and Noble, J. (1989). Population projections using DSM-III criteria: Alcohol abuse and dependence, 1990-2000. Alcohol, health and research world, 13(4), 366-370; Grant, B. F., Harford, T. C., Dawson, D. A., Chou, P., Dufour, M. and Pickering, R. (1994). Prevalence of DSM-IV alcohol abuse and dependence, United States, 1992. Alcohol Health and Research World, 18(3), 243-248.
4. Vega, W. A., Kolody, B., Hwang, J., Porter, P., Bole, A. and Dimas, J. (1993). Profile of alcohol and drug use during pregnancy in California, 1992: Perinatal substance exposure study. Scientific Report to the State of California Department of Alcohol and Drug Programs, Sacramento, CA: Office of Perinatal Substance Abuse; National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute on Drug Abuse; National Household Survey on Drug Abuse, 1994; Center for the Future of Children (1991). The future of children: Drug exposed infants. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children.
5. National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute of Drug Abuse.

6. Amos, A. (1996). Women and smoking. British Medical Bulletin, 52(1), 74-89.
7. Pierce, J. P. and Gilpin, E. A. (1995). A historical analysis of tobacco marketing and the uptake of smoking by youth in the United States: 1890-1977. Health Psychology, 14(6), 500-508.
8. Surgeon General. (1994). Preventing tobacco use among young people. Washington, DC: U.S. Department of Health and Human Services, Office on Smoking and Health.
9. Pierce, J. P. and Gilpin, E. A. (1995). A historical analysis of tobacco marketing and the uptake of smoking by youth in the United States: 1890-1977. Health Psychology, 14(6), 500-508.
10. Thun, M. J., Day-Lilly, C. A., Calle, E. E., Flanders, W. D. and Heath, C. W. (1995). Excess mortality among cigarette smokers: Changes in a 20-year interval. American Journal of Public Health, 85(9), 1223-1230.
11. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. (1987). Vital Statistics of the United States, 1985, volume II: Mortality. Hyattsville, MD: National Center for Health Statistics; U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. (1988). Vital Statistics of the United States, 1986, volume II: Mortality. Hyattsville, MD: National Center for Health Statistics; U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. (1989). Vital Statistics of the United States, 1987, volume II: Mortality. Hyattsville, MD: National Center for Health Statistics.
12. Rich-Edwards, J. W., Manson, J. E., Hennekens, C. H. and Buring, J. E. (1995). The primary prevention of coronary heart disease in women. New England Journal of Medicine, 332(26), 1758-1766; Rosenberg, L., Palmer, J. R. and Shapiro, S. (1990). Decline in the risk of myocardial infarction among women who stop smoking. New England Journal of Medicine, 322(4), 213-217.
13. Hopper, J. L. and Seeman, E. (1994). The bone density of female twins discordant for tobacco use. New England Journal of Medicine, 330(6), 387-392; Baron, J. A., La Vecchia, C. and Levi, F. (1990). The antiestrogenic effects of cigarette smoking in women. American Journal of Obstetrics, 162(2), 502-514; Slemenda, C. W. (1994). Cigarettes and the skeleton. New England Journal of Medicine, 330(6), 430.
14. Surgeon General. (1980). The health consequences of smoking for women. Washington, DC: U. S. Department of Health and Human Services, Office on Smoking and Health; Lerner, S. (1995). If We've Come Such a Long Way, Why Are We Still Smoking? Ms., 5(6), 22-27.

15. Lex, B. W. (1995). Alcohol and other psychoactive substance dependence in women and men. In M. V. Seeman (Ed.), Gender and psychopathology, (pp. 311-357). Washington, DC: American Psychiatric Press; Blume, S. B. (1988). Alcohol/drug dependent women: New insights into their special problems, treatment, recovery. Minneapolis, MN: Johnson Institute.
16. Reichman, M. E. (1994). Alcohol and breast cancer. Alcohol Health and & Research World, 18(3), 182-184; National Institute on Alcohol Abuse and Alcoholism. (1993). Eighth special report to the U.S. Congress on alcohol and health. Alexandria, VA: National Institute on Alcohol Abuse and Alcoholism; Miller, B. A. (1990). The interrelationships between alcohol and drugs and family violence. In M. De La Rosa, E. Y. Lambert and B. Gropper (Eds.), Drugs and violence: Causes, correlates and consequences. NIDA research monograph 103 (pp. 177-207). Washington, DC: National Institute on Drug Abuse.
17. Bergmann, P. E., Smith, M. B. and Hoffman, N. G. (1995). Adolescent treatment: Implications for assessment, practice guidelines and outcome management. Pediatric Clinics of North America, 42(2), 453-472; Garrison, C. Z., McKeown, R. E., Valois, R. F. and Vincent, M. L. (1993). Aggression, substance use and suicidal behaviors in high school students. American Journal of Public Health, 83(2), 179-184; Silverman, W. H. (1991). Prevalence of substance use in a rural teenage population. Journal of Adolescent Chemical Dependency, 2(1), 107-117; Lowry, R., Holtzmann, D., Truman, B. I., Kann, L., Collins, J. L. and Kolbe, L. J. (1994). Substance use and HIV-related sexual behaviors among U.S. high school students: Are they related? American Journal of Public Health, 84(7), 1116-1120; Mott, F. L. and Haurin, R. J. (1988). Linkages between sexual activity and alcohol and drug use among American adolescents. Family Planning Perspectives, 20(3), 128-136; Ford, K. and Norris, A. (1994). Urban minority youth: Alcohol and marijuana use and exposure to unprotected intercourse. Journal of Acquired Immune Deficiency Syndromes, 7, 389-396.
18. U. S. General Accounting Office. (1994). Foster care: Parental drug abuse has alarming impact on young children. Washington, DC: U.S. General Accounting Office; U. S. House of Representatives, Committee on Ways and Means, Subcommittee on Human Resources. (1990). The enemy within: Crack-cocaine and America's families. Washington, DC: U.S. Government Printing Office.
19. Wiese, D., Daro, D. and National Center on Child Abuse Prevention Research. (1995). Current trends in child abuse reporting and fatalities: The results of the 1994 annual fifty state survey. Chicago, IL: National Committee to Prevent Child Abuse.
20. U.S. General Accounting Office. (1994). Foster care: Parental drug abuse has alarming impact on young children. Washington, DC: U.S. General Accounting Office.
21. Davis, S. F., Byers, R. H., Lindegren, M. L., Caldwell, M. B., Karon, J. M. and Gwinn, M. (1995). Prevalence and incidence of vertically-acquired HIV infection in the United States. Journal of the American Medical Association, 274(12), 952-955.

22. Centers for Disease Control. (1995). Update: AIDS among women--United States, 1994. Morbidity and Mortality Weekly Report, 44(5), 81-84; Melnick, S. L., Sherer, R., Louis, T. A., Hillman, D., Rodriguez, E. M., Lackman, C., Capps, L., Brown, L. S., Carlyn, M., Korvick, J. A. and Deyton, L. (1994). Survival and disease progression according to gender of patients with HIV infection. JAMA, 272(24), 1915-1922; U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Treatment. (1994). Practical approaches in the treatment of women who abuse alcohol and other drugs. Rockville, MD: U.S. Department of Health and Human Services; Cohen, J. B., Hauer, L. B. and Wofsy, C. B. (1989). Women and IV drugs: Parenteral and heterosexual transmission of human immunodeficiency virus. Journal of Drug Issues, 19(1), 39-56; Finnegan, L. P., Davenny, K. and Hartel, D. (1993). Drug use in HIV-infected women. In F. D. Johnstone and M. Johnson (Eds.), HIV in infected women (pp. 133-155). Edinburgh, Great Britain: Churchill Livingstone.
23. Inciardi, J. A., Lockwood, D., Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; Beck, A. J and Gilliard, D. K. (1995, August). Prisoners in 1994. Bureau of Justice Statistics Bulletin, 1-16.
24. U. S. Department of Justice. (1991). Survey of state prison inmates, 1991: Women in prison. Washington, DC: U. S. Department of Justice.
25. Lemle, R. and Mishkind, M. E. (1989). Alcohol and masculinity. Journal of Substance Abuse Treatment, 6(4), 213-222; Blume, S. B. (1992). Alcohol and other drug problems in women. In J. H. Lowinson, P. Ruiz and R. B. Millman (Eds.), Substance abuse: A comprehensive textbook, second edition (pp. 795-807). Baltimore, MD: Williams and Wilkins.
26. Leigh, B. C. (1995). A thing so fallen, and so vile: Images of drinking and sexuality in women. Contemporary Drug Problems, 22(3), 415-434; Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.
27. Lex, B.W. (1991). Some gender differences in alcohol and polysubstance users. Health Psychology, 10(2), 121-132.
28. Falco, M. (1991). Drug abuse: A national policy perspective. Bulletin of the New York Academy of Medicine, 67(3), 196-206.
29. Aldrich, M. R. (1994). Historical notes on women addicts. Journal of Psychoactive Drugs, 26(1), 61-64; Ball, J.C. and Chambers, C. D. (1970). The epidemiology of opiate addiction in the United States. Springfield, IL: Charles C. Thomas; Brecher, E. M. and the Editors of Consumer Reports. (1972). Licit and illicit drugs, Boston: Little, Brown and Company.

30. Aldrich, M. R. (1994). Historical notes on women addicts. Journal of Psychoactive Drugs, 26(1), 61-64; Ball, J.C. and Chambers, C. D. (1970). The epidemiology of opiate addiction in the United States Springfield, IL: Charles C. Thomas; Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: Tab Books.
31. Brecher, E. M. and the Editors of Consumer Reports. (1972). Licit and illicit drugs. Boston: Little, Brown and Company.
32. Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: Tab Books.
33. Aldrich, M. R. (1994). Historical notes on women addicts. Journal of Psychoactive Drugs, 26(1), 61-64; Leigh, B. C. (1995). A thing so fallen, and so vile: Images of drinking and sexuality in women. Contemporary Drug Problems, 22(3), 415-434; Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: Tab Books.
34. Aldrich, M. R. (1994). Historical notes on women addicts. Journal of Psychoactive Drugs, 26(1), 61-64.
35. Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.
36. Leigh, B. C. (1995). A thing so fallen, and so vile: Images of drinking and sexuality in women. Contemporary Drug Problems, 22(3), 415-434; Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: Tab Books; Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146; Gomberg, E. S. (1986). Women: Alcohol and other drugs. In B. Segal (Ed.), Perspectives on drug use in the United States (pp. 76-78). New York: Haworth Press.
37. Musto, D. F. (1987). The American disease: Origins of narcotic control. New York: Oxford University Press.
38. Aldrich, M. R. (1994). Historical notes on women addicts. Journal of Psychoactive Drugs, 26(1), 61-64; Falco, M. (1991). Drug abuse: A national policy perspective. Bulletin of the New York Academy of Medicine, 67(3), 196-206; Blume, S. B. (1988). Alcohol/drug dependent women: New insights into their special problems, treatment, recovery. Minneapolis, MN: Johnson Institute.
39. Ball, J.C. and Chambers, C. D. (1970). The epidemiology of opiate addiction in the United States. Springfield, IL: Charles C. Thomas.
40. Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems

in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.

41. Leigh, B. C. (1995). A thing so fallen, and so vile: Images of drinking and sexuality in women. Contemporary Drug Problems, 22(3), 415-434.

42. Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: Tab Books.

43. Jacobson, M., Atkins, R. and Hacker, G. (1983). The Booze Merchants: The Inebriating of America. Washington, DC: Center for Science in the Public Interest.

44. Elliott, S., "The media business: Advertising; In a whisky pitch to Generation X members, Leo Burnett tries to prove it is not their fathers' agency," The New York Times, 9/13/93, D7.

45. Wilsnack, R. W. and Wilsnack, S. C. (1978). Sex roles and drinking among adolescent girls. Journal of Studies on Alcohol, 39(11), 1855-1856.

46. Ford, B. and Chase, C. (1987). Betty: A glad awakening. Garden City, NY: Doubleday.

47. Gomberg, E. S. (1986). Women: Alcohol and other drugs. In B. Segal (Ed.), Perspectives on drug use in the United States (pp. 75-109). New York: Haworth Press.

48. Robins, L. N., Mills, J. L., Krulewitch, C. and Herman, A. A. (1993). Effects of in utero exposure to street drugs. American Journal of Public Health, 83 (Suppl.), 9-32; Mayes, L. C., Granger, R. H., Bornstein, M. H. and Zuckerman, B. (1992). The problem of prenatal cocaine exposure: A rush to judgment. Journal of the American Medical Association, 267(3), 406-408; Chasnoff, I. J. (1991). Cocaine and pregnancy: Clinical and methodological issues. Clinics in Perinatology, 18(1), 113-123; Cotton, P. (1994). Smoking cigarettes may do developing fetus more harm than ingesting cocaine, some experts say. Journal of the American Medical Association, 271(8), 576-577; DiFranza, J. R. and Lew, R. A. (1995). Effect of maternal cigarette smoking on pregnancy complications and sudden infant death syndrome. Journal of Family Practice, 40(4), 385-394; Streissguth, A. P., Barr, H. M., Olson, H. C., Sampson, P. D., Bookstein, F. L. and Burgess, D. M. (1994). Drinking during pregnancy decreases word attack and arithmetic scores on standardized tests: Adolescent data from a population-based prospective study. Alcoholism: Clinical and Experimental Research, 18(2), 248-254; National Institute on Alcohol Abuse and Alcoholism. (1993). Eighth special report to the U.S. Congress on alcohol and health. Alexandria, VA: National Institute on Alcohol Abuse and Alcoholism.

CHAPTER II.

REFERENCES

1. Brett, P. J., Graham, K. and Smythe, C. (1995). An analysis of specialty journals on alcohol, drugs and addictive behaviors for sex bias in research methods and reporting. Journal of Studies on Alcohol, 56(1), 24-34; Chavkin, W. (1994). Women and clinical research. Journal of the American Medical Women's Association, 49(4), 99-100; Wilke, D. (1994). Women and alcoholism: How a male-as-norm bias affects research assessment and treatment. Health and Social Work, 19(1), 29-35; Pittman, D. J. and White, H. R. (Eds.). (1991). Society, culture and drinking patterns reexamined. New Brunswick, NJ: Alcohol Research Documentation, Inc., Rutgers Center of Alcohol Studies.
2. Lieber, C. S., Professor of Medicine and Pathology, Mount Sinai School of Medicine of CUNY and Director of Alcohol Research and Treatment Center at Bronx VA Medical Center. (1995). Personal communication.
3. Lieber, C. S. (1994). Alcohol and the liver: 1994 update. Gastroenterology, 106(4), 1085-1105; Frezza, M., Padova, C., Pozzato, G., Terpin, M., Baraona, E. and Lieber, C. S. (1990). High Blood Alcohol Levels in Women. New England Journal of Medicine, 322(2), 95-99; Lieber, C. S. (1993). Women and alcohol: Gender differences in metabolism and susceptibility. In E. S. L. Gomberg, T. D. Nirenberg (Eds.), Women and substance abuse (pp. 1-17). Norwood, NJ: Ablex Publishing; Center for Substance Abuse Treatment. Practical approaches in the treatment of women who abuse alcohol and other drugs. Rockville, MD: U.S. Dept. of Health and Human Services, Center for Substance Abuse Treatment.
4. U.S. Department of Agriculture, Agricultural Research Service, Advisory Guidelines Committee. (1995). Report of the Dietary Guidelines Advisory Committee on the dietary guidelines for Americans, 1995, to the Secretary of Health and Human Services and the Secretary of Agriculture. [Washington, DC]: U.S. Department of Agriculture, Agricultural Research Service, Dietary Guidelines Advisory Committee.
5. Williams, G. D., Grant, B. F., Harford, T. C. and Noble, J. (1989). Population projections using DSM-III criteria: Alcohol abuse and dependence, 1990-2000. Alcohol, Health and Research World, 13(4), 366-370. See also Grant, B. F., Harford, T. C., Dawson, D. A., Chou, P., Dufour, M. and Pickering, R. (1994). Prevalence of DSM-IV alcohol abuse and dependence, United States, 1992. Alcohol and Research World, 18(3), 243-248; Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C. B., Hughes, M., Eshleman, S., Wittchen, H. and Kendler, K. S. (1994). Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: Results from the National Comorbidity Survey. Archives of General Psychiatry, 51(5), 8-19; Grant, Bridget F. (1993). IDC-10

- harmful use of alcohol and the alcohol dependence syndrome: prevalence and implications. Addiction 88, 413-20.
6. Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.
 7. Curtis, J. R., Geller, G., Stokes, E. J., Levine, D. M. and Moore, R. D. (1989). Characteristics, diagnosis and treatment of alcoholism in elderly patients. Journal of the American Geriatrics Society, 37(4), 310-316.
 8. Johnston, L. D., O'Malley, P. M. and Bachman, J. G. (1993). National survey results on drug use from the Monitoring the Future study, 1975-1992, volume II. Rockville, MD: National Institute on Drug Abuse.
 9. Caetano, R. C. (1994). Drinking and alcohol-related problems among minority women. In Welsh, D. M. (Ed.), Alcohol Health and Research World: Women and alcohol, 18(3), (pp. 233-241). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Weibel-Orlando, J. (1984). Women and alcohol: Special populations and cross-cultural variations. In Women and alcohol: Health-related issues: Proceedings of a conference, May 23-25, 1984: NIAAA research monograph 16 (pp. 161-187). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; Leland, J. (1984). Alcohol use and abuse in ethnic minority women. In S. C. Wilsnack and L. J. Beckman (eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 66-96). New York: Guilford Press.
 10. 1993 National Household Survey on Drug Abuse.
 11. Weibel-Orlando, J. (1984). Women and alcohol: Special populations and cross-cultural variations. In Women and alcohol: Health-related issues: Proceedings of a conference, May 23-25, 1984: NIAAA research monograph 16 (pp. 161-187). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism.
 12. Vega, W. A., Kolody, B., Hwang, J., Noble, A. and Porter, P. (1996). Perinatal drug abuse among immigrant and native born Latinas. International Journal of the Addictions, 31(in press); Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application (pp. 381-443). Washington, DC: American Psychological Association; Caetano, R. C. (1994). Drinking and alcohol-related problems among minority women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 233-241). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Austin, G. A. and Gilbert, M. J. (1989). Substance abuse among Latino youth. Portland, OR: Northwest Regional Educational Laboratory; Caetano, R. and Mora, M. E. M. (1988). Acculturation and drinking among people of Mexican descent in Mexico and the United States. Journal of Studies on Alcohol, 49(5), 462-471.

13. Weibel-Orlando, J. (1984). Women and alcohol: Special populations and cross-cultural variations. In Women and alcohol: Health-related issues: Proceedings of a conference, May 23-25, 1984: NIAAA research monograph 16 (pp. 161-187). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; Leland, J. (1984). Alcohol use and abuse in ethnic minority women. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 66-96). New York: Guilford Press; Ja, D. Y. and Aoki, B. (1993). Substance abuse treatment: Cultural barriers in the Asian-American community. Journal of Psychoactive Drugs, 25(1), 61-79.
14. Wilsnack, R. W. and Wilsnack, S. C. (1978). Sex roles and drinking among adolescent girls. Journal of Studies on Alcohol, 39(11), 1855-1874; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.
15. Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press; LaRosa, J. H. (1990). Executive women and health: Perceptions and practices. American Journal of Public Health, 80(12), 1450-1454.
16. LaRosa, J. H. (1990). Executive women and health: Perceptions and practices. American Journal of Public Health, 80(12), 1450-1454.
17. Gomberg, E. S. L. (1993). Gender issues. In M. Galanter (Ed.), Recent developments in alcoholism, volume 11: Ten years of progress, (pp. 95-107). New York: Plenum Press; Wilsnack, S. C. and Wilsnack, R. W. (1991). Epidemiology of women's drinking. Journal of Substance Abuse, 3, 133-157; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press; Wilsnack, S. C., Wilsnack, R. W. and Klassen, A. D. (1984/1985). Drinking and drinking problems among women in a U.S. national survey. Alcohol Health and Research World, 9(2), 3-13; Midanik, L. T. and Clark, W. B. (1995). Drinking-related problems in the United States: Description and trends, 1984-1990. Journal of Studies on Alcohol, 56, 395-402; Midanik, L. T. and Clark, W. B. (1994). The demographic distribution of U.S. drinking patterns in 1990: Description and trends from 1984. American Journal of Public Health, 84(8), 1218-1222.
18. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application (pp. 381-443). Washington, DC: American Psychological Association; Wilsnack, R. W. and Klassen, A. D. and Wilsnack, S. C. (1986). Retrospective analysis of lifetime changes in women's drinking behavior. Advances in Alcohol and Substance Abuse, 5(3), 9-28.

19. Grant, B. F., Harford, T. C., Dawson, D. A., Chou, P., Dufour, M. and Pickering, R. (1994). Prevalence of DSM-IV alcohol abuse and dependence, United States, 1992. Alcohol and Research World, 18(3), 243-248; Robins, L. N., Helzer, J. E., Pryzbeck, T. R. and Regier, D. A. (1988). Alcohol disorders in the community: A report from the epidemiological catchment area. In R. M. Rose and J. Barret (Eds.), Alcoholism: Origins and outcome (pp. 15-29). New York: Raven Press.
20. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application (pp. 381-443). Washington, DC; American Psychological Association; Wilsnack, S. C., Klassen, A. D., Schur, B. E. and Wilsnack, R. W. (1991). Predicting onset and chronicity of women's problem drinking: A five-year longitudinal analysis. American Journal of Public Health, 81(3), 305-318; Wilsnack, R. W., Wilsnack, S. C. and Klassen, A. D. (1984). Women's drinking and drinking problems: Patterns from a 1981 national survey. American Journal of Public Health, 74(11), 1231-1238; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.
21. Gomberg, E. S. L. (1995). Older women and alcohol use and abuse. In M. Galanter (Ed.), Alcoholism and Women, (pp. 61-79). New York: Plenum Press.
22. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association.
23. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association.
24. Hill, S. Y (1995). Vulnerability to alcoholism in women: Genetic and cultural factors. In M. Galanter (Ed.), Alcoholism and women (pp. 9-28). New York: Plenum Press; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application (pp. 381-443). Washington, DC: American Psychological Association; Svikis, D. S., Velez, M. L. and Pickens, R. W. (1994). Genetic aspects of alcohol use and alcoholism in women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 192-196). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Hill, S. (1993). Genetic vulnerability to alcoholism in women. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 42-61). Norwood, NJ: Ablex Publishing; Kendler, K. S., Heath, A. C., Neale, M. C., Kessler, R. C. and Eaves, L. J. (1992). A population-based twin study of alcoholism in women. Journal of the American Medical Association, 268(14), 1877-1882.

25. Gomberg, E. S. L. and Lisansky, J. M. (1984). Antecedents of alcohol problems in women. In S. C. Wilsnack and L. J. Beckman (eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 233-259). New York: Guilford Press; Cotton, N. S. (1979). The familial incidence of alcoholism: A review. Journal of Studies on Alcohol, 40(1), 89-116; Leckman, A. L., Umland, B. E. and Blay, M. (1984). Alcoholism in the families of family practice outpatients. Journal of Family Practice, 19(2), 205-207.
26. Friedman, A. S., Granick, S., Bransfield, S. and Kreishner, C. (1995). Gender differences in early life risk factors for substance use/abuse: A study of an African-American sample. American Journal of Drug and Alcohol Abuse, 21(4), 511-531; Kendler, K. S., Neale, M. C., Kessler, R. C., Heath, A. C., Corey, L. A. and Eaves, L. J. (1996). Childhood parental loss and alcoholism in women: A causal analysis using a twin-family design. Psychological Medicine, 26(1), 79-95; Hill, S. Y. and Smith, T. R. (1991). Evidence for genetic mediation of alcoholism in women. Journal of Substance Abuse, 3(2), 159-74; Lex, B. W. (1995). Alcohol and other psychoactive substance dependence on women and men. In M. V. Seeman (Ed.), Gender and psychopathology (pp. 311-358). Washington, DC: American Psychiatric Press; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application (pp. 381-443). Washington, DC.
27. Miller, B. A., Downs, W. R. and Testa, M. (1993). Interrelationships between victimization experiences and women's alcohol use. Journal of Studies on Alcohol, 11(Suppl.), 109-117; Miller, B. A., Downs, W. R., Gondoli, D. M. and Keil, A. (1987). The role of childhood sexual abuse in the development of alcoholism in women. Violence and Victims, 2(3), 157-172; Miller, B. A. and Downs, W. R. (1993). The impact of family violence on the use of alcohol by women. Alcohol, Health and Research World, 17(2), 137-143.
28. Hill, S. Y (1995). Vulnerability to alcoholism in women: Genetic and cultural factors. In M. Galanter (Ed.), Alcoholism and women (pp. 9-28). New York: Plenum Press; Miller, B. A., Downs, W. R. and Testa, M. (1993). Interrelationships between victimization experiences and women's alcohol use. Journal of Studies on Alcohol, 11(Suppl.), 109-117; Miller, B. A. and Downs, W. R. (1993). The impact of family violence on the use of alcohol by women. Alcohol, Health and Research World, 17(2), 137-143; Miller, B. A., Downs, W. R., Gondoli, D. M. and Keil, A. (1987). The role of childhood sexual abuse in the development of alcoholism in women. Violence and Victims, 2(3), 157-172; Winfield, I., George, L. K., Swartz, M. and Blazer, D. G. (1990). Sexual assault and psychiatric disorders among a community sample of women. American Journal of Psychiatry, 147(3), 335-341; Browne, A. and Finkelhor, D. (1986). Impact of child sexual abuse: A review of the research. Psychological Bulletin, 99(1), 66-77.
29. Windle, M., Windle, R. C., Scheidt, D. M. and Miller, G. B. (1995). Physical and sexual abuse and associated mental disorders among alcoholic inpatients. American Journal of Psychiatry, 152(9), 1322-1328.

30. Janikowski, T. P. and Glover, N. M. (1994). Incest and substance abuse: Implications for treatment professionals. Journal of Substance Abuse Treatment, 11(3), 117-183; Hurley, D. L. (1991). Women, alcohol and incest: An analytical review. Journal of Studies on Alcohol, 52(3), 253-268.
31. Helzer, J. E. and Pryzbeck, T. R. (1994). Psychiatric disorders in the addicted patient: The co-occurrence of alcoholism with other psychiatric disorders. In N. S. Miller (Ed.), Principles of addiction medicine (pp. 1-5). Chevy Chase, MD: American Society of Addiction Medicine.
32. Schuckit, M. A., Tipp, J. E., Anthenelli, R. M., Bucholz, K. K., Hesselbrock, V. M. and Nurnberger, J. I. (1996). Anorexia nervosa and bulimia nervosa in alcohol-dependent men and women and their relatives. American Journal of Psychiatry, 153(1), 74-82; Holderness, C. C., Brooks-Gunn, J. and Warren, M. P. (1994). Co-morbidity of eating disorders and substance abuse: Review of the literature. International Journal of Eating Disorders, 16(1), 1-34; Goldbloom, D. S. (1993). Alcohol misuse and eating disorders: Aspects of an association. Alcohol and Alcoholism, 28(4), 375-381; Krahn, D. D. (1993). The relationship of eating disorders and substance abuse. In E. S. L. Gomberg, T. D. Nirenberg (Eds.), Women and substance abuse (pp. 286-313). Norwood, NJ: Ablex Publishing.
33. Gomberg, E. S. L. and Lisansky, J. M. (1984). Antecedents of alcohol problems in women. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 233-259). New York: Guilford Press.
34. Wilsnack, R. W., Wilsnack, S. C. and Klassen, A. D. (1987). Antecedents and consequences of drinking and drinking problems in women: Patterns from a U.S. national survey. In R. A. Dienstbier (Ed.), Nebraska symposium on motivation: Alcohol and addictive behavior (pp. 85-158). Lincoln, NB: University of Nebraska Press; Wilsnack, S.C., Klassen, A. D., Schur, B. E. and Wilsnack, R. W. (1991). Predicting the onset and chronicity of women's problem drinking: A five-year longitudinal analysis. American Journal of Public Health, 81(3), 305-318; Johnson, V. and White, H. R. (1995). The relationship between work-specific and generalized stress and alcohol and marijuana use among recent entrants to the labor force. Journal of Drug Issues, 25(2), 237-251.
35. Allan, C. A. and Cooke, D. J. (1985). Stressful life events and alcohol misuse in women: A critical review. Journal of Studies on Alcohol, 46(2), 147-152; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application (pp. 381-443). Washington, DC: American Psychological Association; Blume, S. B. (1994). Gender differences in alcohol-related disorders. Harvard Review of Psychiatry, 2(1), 7-14.

36. Centers for Disease Control. (1994). Cigarette smoking among adults: United States, 1993. Morbidity and Mortality Weekly Report, 43(50), 925-930; Centers for Disease Control and Prevention. (1995). Indicators of nicotine addiction among women--United States, 1991-1992. Morbidity and Mortality Weekly Report, 44(6), 102-105.
37. Pierce, J. P. and Gilpin, E. A. (1995). A historical analysis of tobacco marketing and the uptake of smoking by youth in the United States: 1890-1977. Health Psychology, 14(6), 500-508.
38. Surgeon General's Advisory Committee on Smoking and Health. (1964). Smoking and health. Washington, DC: Public Health Service; Escobedo, L. G. and Peddocrod, J. P. (1996). Smoking prevalence in U.S. birth cohorts: The influence of gender and education. American Journal of Public Health, 86(2), 231-236.
39. Pierce, J. P., Lee, L and Gilpin, E. A. (1994). Smoking initiation by adolescent girls, 1944 through 1988: An association with targeted advertising. Journal of the American Medical Association, 271(8), 608-611.
40. Kessler, L. (1989). Women's magazines' coverage of smoking related health hazards. Journalism Quarterly, 66(2), 316-323; Warner, K. E. and Goldenhar, L. M. (1992). Targeting of cigarette advertising in U.S. magazines, 1959-1986. Tobacco Control, 1, 25-30.
41. Pierce, J. P., Lee, L and Gilpin, E. A. (1994). Smoking initiation by adolescent girls, 1944 through 1988: An association with targeted advertising. Journal of the American Medical Association, 271(8), 608-611; Pierce, J. P. and Gilpin, E. A. (1995). A historical analysis of tobacco marketing and the uptake of smoking by youth in the United States: 1890-1977. Health Psychology, 14(6), 500-508.
42. Gilpin, E. A., Lee, L. Evans, N. and Pierce, J. P. (1994). Smoking initiation rates in adults and minors: United States, 1944-1988. American Journal of Epidemiology, 140(6), 535-543.
43. Johnston, L. D., O'Malley, P. M. and Bachman, J. G. (1995). National survey results on drug use from the Monitoring the Future study, 1975-1994, volume I. Rockville, MD: National Institute on Drug Abuse; 1993 NHSDA.
44. Mermelstein, R. J. and Borrelli, B. (1995). Women and smoking. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application (pp. 309-348). Washington, DC; American Psychological Association; Feigelman, W. and Lee, J. A. (1995). Patterns of cigarette use among black and white adolescents. American Journal on Addictions, 4(3), 215-225.
45. Evans, N., Farkas, A., Gilpin, E., Berry C. and Pierce, J. P. (1995). Influence of tobacco marketing and exposure to smokers on adolescent susceptibility to smoking. Journal of National Cancer Institute, 87(20), 1538-1545.

46. Clayton, S. (1991). Gender differences in psychosocial determinants of adolescent smoking. Journal of School Health, 61(3), 115-120; Evans, N., Gilpin, E., Farkas, A. J., Shenassa, E. and Pierce, J. P. (1995). Adolescents' perceptions of their peers' health norms. American Journal of Public Health, 85(8), 1064-1069.
47. Camp, D. E., Klesges, R. C. and Relyea, G. (1993). The relationship between body weight concerns and adolescent smoking. Health Psychology, 12(1), 24-32.
48. Pirie, P. L., Murray, D. M. and Luepker, R. V. (1991). Gender differences in cigarette smoking and quitting in a cohort of young adults. American Journal of Public Health, 81(3), 324-327; Klesges, L. M. and Meyers, A. W. (1989). Smoking, body weight and their effects on smoking behavior: A comprehensive review of the literature. Psychological Bulletin, 106(2), 204-230.
49. Klesges, L. M. and Meyers, A. W. (1989). Smoking, body weight and their effects on smoking behavior: A comprehensive review of the literature. Psychological Bulletin, 106(2), 204-230.
50. Flegal, K. M., Troiano, R. P., Pamuk, E. R., Kuczmarska, R. J. and Campbell, S. M. (1995). The influence of smoking cessation on the prevalence of overweight in the United States. New England Journal of Medicine, 333(18), 1165-1170.
51. 1993 National Household Survey on Drug Abuse.
52. Escobedo, L. G. and Peddocro, J. P. (1996). Smoking prevalence in U.S. birth cohorts: The influence of gender and education. American Journal of Public Health, 86(2), 231-236.
53. Solomon, L. J. and Flynn, B. S. (1993). Women who smoke. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 339-349). New York: Oxford University Press.
54. Royce, J. M., Hymowitz, N., Corbett, K., Hartwell, T. D. and Orlandi, M. A. (1993). Smoking cessation factors among African Americans and whites. American Journal of Public Health, 83(2), 220-226; Solomon, L. J., and Flynn, B. S. (1993). Women who smoke. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 339-349). New York: Oxford University Press.
55. Bachman, J. G., Wallace, J. M., O'Malley, P. M., Johnston, L. D., Kurth, C. L. and Neighbors, H. W. (1991). Racial-ethnic differences in smoking, drinking and illicit drug use among American high school seniors, 1976-1989. American Journal of Public Health, 81(3), 372-377.
56. Nelson, D. E., Giovino, G. A., Shopland, D. R., Mowery, P. D., Mills, S. L. and Eriksen, M. P. (1995). Trends in cigarette smoking among U.S. adolescents, 1974 through 1991. American Journal of Public Health, 85(1), 34-40; McIntosh, H. (1995). Black teens not smoking in great numbers. Journal of the National Cancer Institute, 87(8), 564; Evans,

N., Gilpin, E., Farkas, A. J., Shenassa, E. and Pierce, J. P. (1995). Adolescents' perceptions of their peers' health norms. American Journal of Public Health, 85(8), 1064-1069; Royce, J. M., Hymowitz, N., Corbett, K., Hartwell, T. D. and Orlandi, M. A. (1993). Smoking cessation factors among African Americans and whites. American Journal of Public Health, 83(2), 220-226; Camp, D. E., Klesges, R. C. and Relyea, G. (1993), The relationship between body weight concerns and adolescent smoking. Health Psychology, 12(1), 24-32; Headen, S. W., Bauman, K. E., Deane, G. D. and Koch, G. G. (1991). Are the correlates of cigarette smoking initiation different for black and white adolescents? American Journal of Public Health, 81(7), 854-858; Center on Addiction and Substance Abuse at Columbia University. (1995). National survey of American attitudes on substance abuse. New York: CASA.

57. Husten, C. G., Chrismon, J. H. and Reddy, M. N. (1996). Trends and effects of cigarette smoking among girls and women in the United States, 1965-1993. Journal of the Medical American Women's Association, 51 (1), 1-8.

58. Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; Kandall, S. R. and Chavkin, W. (1992). Illicit drugs in America: History, impact on women and infants, and treatment strategies for women. Hastings Law Journal, 43(3), 615-643; U. S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: U. S. General Accounting Office.

59. Kandall, S. R. and Chavkin, W. (1992). Illicit drugs in America: History, impact on women and infants, and treatment strategies for women. Hastings Law Journal, 43(3), 615-643; Kandall, S. R. (1991). Perinatal effects of cocaine and amphetamine use during pregnancy. Bulletin of the New York Academy of Medicine, 67(3), 240-255.

60. Falco, M. (1991). Drug abuse: A national policy perspective. Bulletin of the New York Academy of Medicine, 67(3), 196-206.

61. Lillie-Blanton, M., Anthony, J. C. and Schuster, C. R. (1993). Probing the meaning of racial/ethnic group comparisons in crack cocaine smoking. Journal of the American Medical Association, 269(8), 993-997; Kandall, S. R. and Chavkin, W. (1992). Illicit drugs in America: History, impact on women and infants, and treatment strategies for women. Hastings Law Journal, 43(3), 615-643; Falco, M. (1991). Drug abuse: A national policy perspective. Bulletin of the New York Academy of Medicine, 67(3), 196-206; Wagner-Echeagaray, F. A., Schutz, C. G., Chilcoat, H. D. and Anthony, J. C. (1994). Degree of acculturation and the risk of crack cocaine smoking among Hispanic Americans. American Journal of Public Health, 84(11), 1825-1827; Maher, L. and Curtis, R. (1992). Women on the edge of crime: Crack cocaine and the changing contexts of street-level sex work in New York City. Crime, Law and Social Change, 18(3), 221-258.

62. Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO; Evelyn Davis, MD, Harlem Hospital, New York, NY. (1995). Personal Communication.
63. Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; McCoy, H. V., Inciardi, J. A., Letsch, L. R., Pottieger, A. E. and Saum, C. A. (1995). Women, crack and crime: Gender comparisons of criminal activity among crack cocaine users. Contemporary Drug Problems, 22(3), 435-451; Finnegan, L. P., Sloboda, Z., Haverkos, H. W., Mello, N. K., Kreek, M. J., Cottler, L. B. and Frank, D. A. (1995). Drug abuse and the health of women. In L. S. Harris (Ed.), Problems of drug dependence, 1994: Volume I, NIDA research monograph 152 (pp. 45-52). Rockville, MD: NIDA; Fagan, J. (1994). Women and drugs revisited: Female participation in the cocaine economy. Journal of Drug Issues, 24(2), 179-25; Mieczkowski, T. (1994). The experiences of women who sell crack: Some descriptive data from the Detroit crack ethnography project. The Journal of Drug Issues, 24(2), 227-248.
64. Liepman, M. R., Goldman, R. E., Monroe, A. D., Green, K. W., Sattler, A. L., Broadhurst, J. B. and Gomberg, E. S. L. (1993). Substance abuse by special populations of women. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 214-257). Norwood, NJ: Ablex Publishing.
65. Lillie-Blanton, M., Anthony, J. C. and Schuster, C. R. (1993). Probing the meaning of racial/ethnic group comparisons in crack cocaine smoking. Journal of the American Medical Association, 269(8), 993-997; Inciardi, J. A., Lockwood, D., Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; Booth, R. E. (1995). Gender differences in high-risk sex behaviors among heterosexual drug injectors and crack smokers. American Journal of Drug and Alcohol Abuse, 21(4), 419-432; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO; Maher, L. and Curtis, R. (1992). Women on the edge of crime: Crack cocaine and the changing contexts of street-level sex work in New York City. Crime, Law and Social Change, 18(3), 221-258.
66. National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute on Drug Abuse.
67. Lex, B. W. (1995). Alcohol and other psychoactive substance dependence in women and men. In M. V. Seeman (Ed.), Gender and psychopathology (pp.311-358). Washington, DC: American Psychiatric Press; Anglin, M. C. and Fisher, D. G. (1993). Alcohol and heroin use patterns of narcotics addicts: Gender and ethnic differences. American Journal of Drug and Alcohol Abuse, 19(2), 219-238; Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners, NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse; Griffin, M. L., Weiss, R. D., Mirin, S. M. and Lange, U. (1989). A comparison of male and female cocaine abusers. Archives of General Psychiatry, 46, 122-126.

68. Denier, C. A., Thevos, A. K., Latham, P. K. and Randall, C. L. (1991). Psychological and psychopathology differences in hospitalized male and female cocaine abusers: A retrospective chart review. Addictive Behaviors, 16(6), 489-496; Almog, Y. J., Anglin, M. C. and Fisher, D. G. (1993). Alcohol and heroin use patterns of narcotics addicts: Gender and ethnic differences. American Journal of Drug and Alcohol Abuse, 19(2), 219-238; Griffin, M. L., Weiss, R. D., Mirin, S. M. and Lange, U. (1989). A comparison of male and female cocaine abusers. Archives of General Psychiatry, 46, 122-126.
69. Robbins, C. and Clayton, R. R. (1989). Gender-related differences in psychoactive drug use among older adults. Journal of Drug Issues, 19(2), 207-219; The Rolling Stones. (1966). Mother's little helper. A. L. Oldham (Prod.).
70. Kaestner, E., Frank, B., Marel, R., and Schmediler, J. (1986). Substance use among females in New York State: Catching up with the males. Advances in Alcohol and Substance Abuse, 5(3), 29-49.
71. Senay, E. C. (1989). Addictive behaviors and benzodiazepines: Abuse liability and physical dependence. Advances in Alcohol and Substance Abuse, 8(1), 107-124; Richelson, E. (1984). Psychotropics and the elderly: Interactions to watch for. Geriatrics, 39(12), 30-42.
72. Gomberg, E. S. L. (1993). Women and alcohol: Use and abuse. Journal of Nervous and Mental Disease, 181(4), 211-219; Gomberg, E. S. L. (1989). Alcoholism in women: use of other drugs. Alcoholism: Clinical and Experimental Research, 13, 338.
73. Baum, C., Kennedy, D. L., Knapp, D. E., Juergens, J. P. and Faich, G. A. (1988) Prescription drug use in 1984 and changes over time. Medical Care, 26(2), 105-114; Clayton, R. R., Voss, H. L., Robbins, C. and Skinner, W. (1986). Gender differences in drug use: An epidemiological perspective. In Women and drugs: A new era for research: NIDA research monograph 65. Washington, DC: NIDA.
74. Williams, L. A., Burke, L. B. and Kennedy, D. L. (1991). Drug utilization in the United States, 1990: Poster presentation to the American Society of Hospital Pharmacists: Mid-year clinical meeting, December 1991. Rockville, MD: Food and Drug Administration, Division of Epidemiology and Surveillance; Lipkind, K. L. (1995). Advance data: National hospital ambulatory medical survey: 1993: Outpatient department summary. Hyattsville, MD: National Center for Health Statistics; Schappert, S. M. (1993). Advance data: Office visits to psychiatrists: United States, 1989-1990. National Center for Health Statistics; National Center for Health Statistics. (1995). Health, United States, 1994. Hyattsville, MD: Public Health Service; Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C. B., Hughes, M., Eshleman, S., Witcher, H. and Kendler, K. S. (1994). Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: Results from the National Comorbidity Survey. Archives of General Psychiatry, 51(5), 8-19; Verbrugge, L. M. (1982). Sex differences in legal drug use. Journal of Social Issues, 38(2), 59-76.

75. Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C. B., Hughes, M., Eshleman, S., Wittchen, H. and Kendler, K. S. (1994). Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: Results from the National Comorbidity Survey. Archives of General Psychiatry, 51(5), 8-19.
76. Liebowitz, B. (1995). Personal communication, November 8, Washington, DC: National Institute of Mental Health; Anthony, J. C. and Aboraya, A. (1992). Epidemiology of selected mental disorders in later life. In J. E. Birren, R.B. Sloane and G. D. Cohen (Eds.), Handbook of mental health and aging (pp. 27-73). San Diego, CA: Academic Press.
77. Helzer, J. E. and Pryzbeck, T. R. (1994). Psychiatric disorders in the addicted patient: The co-occurrence of alcoholism with other psychiatric disorders. In N. S. Miller (Ed.), Principles of addiction medicine (pp. 1-5). Chevy Chase, MD: American Society of Addiction Medicine; Anthenelli, R. M. and Schuckit, M. A. (1994). Psychiatric disorders in the addicted patient: Affective and anxiety disorders. In N. S. Miller (Ed.), Principles of addiction medicine (pp. 1-7). Chevy Chase, MD: American Society of Addiction Medicine; Drake, R. E., Alterman, A. I. and Rosenberg, S. R. (1993). Detection of substance use disorders in severely mentally ill patients. Community Mental Health Journal, 29(2), 175-192.
78. N. Van Den Bergh (Ed.), (1991). Feminist perspectives on addictions. New York: Springer Publishing Company; Travis, C. B. (1988). Women and health psychology. Hillsdale, NJ: Lawrence Erlbaum Associates; Morabia, A., Farbre, J. and Dunand, J. P. (1992). The influence of patient and physician gender on prescription of psychotropic drugs. Journal of Clinical Epidemiology, 45(2), 111-116.
79. Verbrugge, L. M. (1980). Sex differences in complaints and diagnoses. Journal of Behavioral Medicine, 3(4), 327-354.
80. Hohmann, A. A. (1989). Gender bias in psychotropic drug prescribing in primary care. Medical Care, 27(5), 478-490.
81. Gomberg, E. S. L. (1995). Older women and alcohol use and abuse. In M. Galanter (Ed.), Alcoholism and Women, (pp. 61-79). New York: Plenum Press.
82. Mellinger, G. D., Balter, M. B. and Uhlenhuth, E. H. (1984). Prevalence and correlates of the long-term regular use of anxiolytics. Journal of the American Medical Association, 251(3), 375-379.
83. Ray, W., Taylor, J. A., Meador, K. G., Lichtenstein, M. J., Griffin, M. R., Fought, R., Adams, M. L. and Blazer, D. G. (1993). Reducing antipsychotic drug use in nursing homes. Archives of Internal Medicine, 153, 713-721; U.S. Senate Special Committee on Aging, the American Association of Retired Persons, the Federal Council on the Aging and the U.S. Administration on Aging. (1991). Aging America: Trends and projections. Washington, DC: U.S. Department of Health and Human Services.

84. Morse, R. M. (1994). Elderly patients. In N. S. Miller (Ed.), Principles of addiction medicine (pp. 1-6). Chevy Chase, MD: American Society of Addiction Medicine; Szwabo, P. A. (1993). Substance abuse in older women. Clinics in Geriatric Medicine, 9(1), 197-208; Miller, N. S., Belkin, B. M. and Gold, M. S. (1991). Alcohol and drug dependence among the elderly: Epidemiology, diagnosis and treatment. Comprehensive Psychiatry, 32(2), 153-165; Curtis, J. R., Geller, G., Stokes, E. J., Levine, D. M. and Moore, R. D. (1989). Characteristics, diagnosis and treatment of alcoholism in elderly patients. Journal of the American Geriatrics Society, 37(4), 310-316; Glantz, M. D. and Backenheimer, M. S. (1988). Substance abuse among elderly women. Clinical Gerontologist, 8(1), 3-26.
85. Curtis, J. R., Geller, G., Stokes, E. J., Levine, D. M. and Moore, R. D. (1989). Characteristics, diagnosis and treatment of alcoholism in elderly patients. Journal of the American Geriatrics Society, 37(4), 310-316.
86. Gilpin, E. A., Lee, L., Evans, N. and Pierce, J. P. (1994). Smoking initiation rates in adults and minors: United States, 1944-1988. American Journal of Epidemiology, 140(6), 535-543.
87. 1993 National Household Survey on Drug Abuse; Breslau, N. and Peterson, E. L. (1996). Smoking cessation in young adults: Age at initiation of cigarette smoking and other suspected influences. American Journal of Public Health, 86(2), 214-220.
88. Windle, M. (1991). Alcohol use and abuse: Some findings from the National Adolescent Student Health Survey. Alcohol, Health and Research World, 15(1), 5-10; Wechsler, H. and McFadden, M. (1976). Sex differences in adolescent alcohol and drug use: A disappearing phenomenon. Journal of Studies on Alcohol, 37(9), 1291-1301.
89. Johnston, L. D., O'Malley, P. M. and Bachman, J. G. (1995). National survey results on drug use from the Monitoring the Future Study, 1975-1994, volume I. Rockville, MD: National Institute on Drug Abuse; Thompson, K. M. and Wilsnack, R. W. (1984). Drinking and drinking related problems among female adolescents: Patterns and influences. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 37-65). New York: Guilford Press.
90. Allen, O. and Page, R. M. (1994). Variance in substance use between rural black and white Mississippi high school students. Adolescence, 29(114), 401-404.
91. Khoury, E. L., Warheit, G. J., Zimmerman, R. S., Vega, W. A. and Gil, A. G. (1996). Gender and ethnic differences in the prevalence of alcohol, cigarette and illicit drug use over time in a cohort of young Hispanic adolescents in south Florida. Women and Health, 24(1), in press; Austin, G. A. and Gilbert, M. J. (1989). Substance abuse among Latino youth. Portland, OR: Northwest Regional Educational Laboratory; Gilbert, M. J. and Alcocer, A. M. (1988). Alcohol use and Hispanic youth: An overview. Journal of Drug Issues, 18(1), 33-48.

92. Johnston, L. D., O'Malley, P. M. and Bachman, J. G. (1995). National survey results on drug use from the Monitoring the Future study, 1975-1994, volume I. Rockville, MD: National Institute on Drug Abuse; Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners, NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse; Robins, L. N. and Pryzbeck, T. R. (1985). Age of onset of drug use as a factor in drug use and other disorders. In C. L. Jones and R. J. Battjes (Eds.), Etiology of drug abuse: Implications for prevention, NIDA monograph 56 (pp. 178-192). Rockville, MD: National Institute on Drug Abuse; Kaestner, E., Frank, B., Marel, R. and Schmediler, J. (1986). Substance use among females in New York State: Catching up with the males. Advances in Alcohol and Substance Abuse, 5(3), 29-49; Wechsler, H. and McFadden, M. (1976). Sex differences in adolescent alcohol and drug use: A disappearing phenomenon. Journal of Studies on Alcohol, 37(9), 1291-1301.
93. Clayton, R. R., Voss, H. L., Robbins, C. and Skinner, W. (1986). Gender differences in drug use: An epidemiological perspective. In Women and drugs: A new era for research: NIDA research monograph 65 (pp. 80-99). Rockville, MD: NIDA.
94. Windle, M. (1991). Alcohol use and abuse: Some findings from the National Adolescent Student Health Survey. Alcohol, Health and Research World, 15(1), 5-10; Robins, L. N. and Pryzbeck, T. R. (1985). Age of onset of drug use as a factor in drug use and other disorders. In C. L. Jones and R. J. Battjes (Eds.), Etiology of drug abuse: Implications for prevention, NIDA monograph 56 (pp. 178-192). Rockville, MD: National Institute on Drug Abuse.
95. Bodinger-de Uriarte, C. and Austin G. (1991). Substance abuse among adolescent females. Portland, OR: Northwest Regional Educational Laboratory.

CHAPTER III.

REFERENCES

1. Schuckit, M. A., Anthenelli, R. M., Bucholz, K. K. and Hesselbrock, V. M. (1995). The time course of development of alcohol-related problems in men and women. Journal of Studies on Alcohol, 56(2), 218-225; Urbano-Marquez, A., Estruch, R., Fernandez-Sola, J., Nicolas, J. M., Pare, J. C. and Rubin, E. (1995). The greater risk of alcoholic cardiomyopathy in women compared to men. Journal of the American Medical Association, 274(2), 149-154; Blume, S. B. (1994). Gender differences in alcohol-related disorders. Harvard Review of Psychiatry, 2(1), 7-14; National Institute on Alcohol Abuse and Alcoholism. (1993). Eighth special report to the U.S. Congress on alcohol and health. Alexandria, VA: National Institute on Alcohol Abuse and Alcoholism; Deal, S. R. and Gavaler, J. S. (1994). Are women more susceptible than men to alcohol-induced cirrhosis? In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 189-191). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Nixon, S. J. (1994). Cognitive deficits in alcoholic women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 228-232). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Mann, K., Batra, A., Gunther, A. and Schroth, G. (1992). Do women develop alcoholic brain damage more readily than men? Alcoholism: Clinical and Experimental Research, 16(6), 1052-1056; Parrish, K. M., Higuchi, S. and Dufour, M. C. (1991). Alcohol consumption and the risk of developing liver cirrhosis: Implications for future research. Journal of Substance Abuse, 3(3), 325-335; Hill, S. Y. (1984). Physiological effects of alcohol on women. In Women and alcohol: Health-related issues. Proceedings of a conference, May 23-25, 1984. NIDA research monograph 16 (pp. 199-214). Rockville, MD: National Institute on Drug Abuse.
2. Deal, S. R. and Gavaler, J. S. (1994). Are women more susceptible than men to alcohol-induced cirrhosis? In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 189-191). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association.
3. Smith, E. M., Cloninger, C. R. and Bradford, S. (1983). Predictors of mortality in alcoholic women: Prospective follow-up study. Alcohol and Clinical Experimental Research, 7, 237-243; Blume, S. B. (1994). Gender differences in alcohol-related disorders. Harvard Review of Psychiatry, 2(1), 7-14; Gomberg, E. S. and Nirenberg, T. D. (1993). Antecedents and consequences. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 118-141). Norwood, NJ: Ablex Publishing; Hill, S. Y. (1984). Physiological effects of alcohol on women. In Women and alcohol: Health-related issues. Proceedings of a

conference, May 23-25, 1984. NIDA research monograph 16 (pp. 199-214). Rockville, MD: National Institute on Drug Abuse.

4. Smith, E. M., Cloninger, C. R. and Bradford, S. (1983). Predictors of mortality in alcoholic women: A prospective follow-up study. Alcoholism: Clinical and Experimental Research, 7(2), 237-243.

5. Savage, G. E., Calvert, W. S., Rohde, F. C. and Grant, B. F. (1994). Liver cirrhosis mortality in the United States, 1970-91. Rockville, MD: National Institute of Alcohol Abuse and Alcoholism.

6. National Institute on Alcohol Abuse and Alcoholism. (1993). Eighth special report to the U.S. Congress on alcohol and health. Alexandria, VA: National Institute on Alcohol Abuse and Alcoholism; Grant, B. F., DuFour, M. C. and Harford, T. C. (1988). Epidemiology of alcoholic liver disease. Seminars in Liver Disease, 8(1), 12-25. Others have reported findings that range from 40 to 80 grams for men and 20 to 40 grams for women. See Norton, R., Batey, R., Dwyer, T. and MacMahon, S. (1987). Alcohol consumption and the risk of alcohol related cirrhosis in women. British Medical Journal, 295(6590), 80-82; Hill, S. Y. (1984). Physiological effects of alcohol on women. In Women and alcohol: Health-related issues, Proceedings of a conference, May 23-25, 1984, NIDA research monograph 16 (pp. 199-214). Rockville, MD: National Institute on Drug Abuse; Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Health Maintenance Strategies, 17(4), 905-925.

7. Deal, S. R. and Gavaler, J. S. (1994). Are women more susceptible than men to alcohol-induced cirrhosis? In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 189-191). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; Lieber, C. S. (1994). Alcohol and the liver: 1994 update. Gastroenterology, 106(4), 1085-1105; Lieber, C. S. (1993). Women and alcohol: Gender differences in metabolism and susceptibility. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 1-17). Norwood, NJ: Ablex Publishing; Frezza, M., Padova, C., Pozzato, G., Terpin, M., Baraona, E. and Lieber, C. S. (1990). High blood alcohol levels in women. New England Journal of Medicine, 322(2), 95-99.

8. Lieber, C. S. (1993). Women and alcohol: Gender differences in metabolism and susceptibility. In E. S. L. Gomberg, T. D. Nirenberg (Eds.), Women and substance abuse (pp. 1-17). Norwood, NJ: Ablex Publishing.

9. Savage, G. E., Calvert, W. S., Rohde, F. C. and Grant, B. F. (1994). Liver cirrhosis mortality in the United States, 1970-91. Rockville, MD: National Institute of Alcohol Abuse and Alcoholism; Goddard, L. L. (1993). Background and scope of the alcohol and other drug problem. In L. L. Goddard (Ed.), An African-centered model of prevention for African American youth at risk (pp. 11-18). Rockville, MD: U.S. Dept. of Health and Human

Services, Center for Substance Abuse Prevention; Leland, J. (1984). Alcohol use and abuse in ethnic minority women. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 66-96). New York, Guilford Press; Rebach, H. (1992). Alcohol and drug use among American minorities. In J. E. Tremble, C. S. Balek and S. J. Niemcryn (Eds.). Ethnic and multicultural drug abuse (pp. 23-57). Binghamton, NY: Haworth Press.

10. Smith, E. M., Cloninger, C. R. and Bradford, S. (1983). Predictors of mortality in alcoholic women: A prospective follow-up study. Alcoholism: Clinical and Experimental Research, 7(2), 237-243; Center for Substance Abuse Treatment. Practical approaches in the treatment of women who abuse alcohol and other drugs. Rockville, MD: U.S. Department of Health and Human Services, Center for Substance Abuse Treatment.

11. Center for Substance Abuse Treatment. Practical approaches in the treatment of women who abuse alcohol and other drugs. Rockville, MD: U.S. Department of Health and Human Services, Center for Substance Abuse Treatment.

12. Caetano, R. C. (1994). Drinking and alcohol-related problems among minority women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 233-241). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism.

13. Liepman, M. R., Goldman, R. E., Monroe, A. D., Green, K. W., Sattler, A. L., Broadhurst, J. B. and Gomberg, E. S. L. (1993). Substance abuse by special populations of women. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 214-257). Norwood, NJ: Ablex Publishing.

14. Nixon, S. J. (1994). Cognitive deficits in alcoholic women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 228-232). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism.

15. Nixon, S. J. (1994). Cognitive deficits in alcoholic women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 228-232). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism.

16. Hill, S. Y. (1995). Mental and physical health consequences of alcohol use in women. In M. Galanter (Ed.), Alcoholism and women (pp. 181-197). New York: Plenum Press; van den Brandt, P. A., Goldbohm, R. A. and van't Veer, P. (1995). Alcohol and breast cancer: Results from the Netherlands cohort study. American Journal of Epidemiology, 141(10), 907-915; Gavaler, J. S. (1993). Alcoholic beverage consumption and estrogenization in normal postmenopausal women. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 18-41). Norwood, NJ: Ablex Publishing; Reichman, M. E. (1994). Alcohol and breast cancer. Alcohol Health and Research World, 18(3), 182-184; Longnecker, M. P. (1994). Alcoholic beverage consumption in relation to risk of breast cancer: Meta-analysis and review. Cancer Causes and Control, 5(1), 73-82; Plant, M. L. (1992). Alcohol and breast cancer: A review. International Journal of Addictions, 27(2), 107-

128; Howe, J., Rohan, T., DeCarli, A., Iscovich, J., Kaldor, J., Katsouyanni, K., Marubini, E., Miller, A., Riboli, E., Toniolo, P. and Trichopoulos, D. (1991). The association between alcohol and breast cancer risk: Evidence from the combined analysis of six dietary case-control studies. International Journal of Cancer, 47(5), 707-710.

17. Longnecker, M. P. (1994). Alcoholic beverage consumption in relation to risk of breast cancer: Meta-analysis and review. Cancer Causes and Control, 5(1), 73-82; Gavaler, J. S. (1993). Alcoholic beverage consumption and estrogenization in normal postmenopausal women. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 18-41). Norwood, NJ: Ablex Publishing; Rosenberg, L., Metzger, L. S. and Palmer, J. R. (1993). Alcohol consumption and risk of breast cancer: A review of the epidemiological evidence. Epidemiological Reviews, 15(1), 133-144.

18. Blume, S. B. (1994). Women and addictive disorders. In N. S. Miller (Ed.), Principles of addiction medicine (pp.1-16). Chevy Chase, MD: American Society of Addiction Medicine; Grodstein, F., Goldman, M. B. and Cramer, D. W. (1994). Infertility in women and moderate alcohol use. American Journal of Public Health, 84(9), 1429-1432; Gavaler, J. S. (1993). Alcoholic beverage consumption and estrogenization in normal postmenopausal women. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 18-41). Norwood, NJ: Ablex Publishing; Mello, N. K., Mendelson, J. H. and Teoh, S. K. (1989). Neuroendocrine consequences of alcohol abuse in women. Annals of the New York Academy of Sciences, 562, 211-240; Wilsnack, S. C., Klassen, A. D. and Wilsnack, R. W. (1984). Drinking and reproductive dysfunction among women in a 1981 national survey. Alcoholism: Clinical and Experimental Research, 8(5), 451-458.

19. National Institute on Alcohol Abuse and Alcoholism. (1993). Eighth special report to the U.S. Congress on alcohol and health. Alexandria, VA: National Institute on Alcohol Abuse and Alcoholism.

20. National Institute on Alcohol Abuse and Alcoholism. (1993). Eighth special report to the U.S. Congress on alcohol and health. Alexandria, VA: National Institute on Alcohol Abuse and Alcoholism; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; Tivis, L. J. and Galaver, J. S. (1994). Alcohol, hormones and health in postmenopausal women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 185-188). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Gavaler, J. S. (1993). Alcoholic beverage consumption and estrogenization in normal postmenopausal women. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 18-41). Norwood, NJ: Ablex Publishing.

21. Fuchs, C. S., Stampfer, M. J., Colditz, G. A., Giovannucci, E. L., Manson, J. E., Kawachi, I., Hunter, D. J., Hankinson, S. E., Hennekens, C. H., Rosner, B., Speizer, F. E. and Willett, W. C. (1995). Alcohol consumption and mortality among women. New England Journal of Medicine, 332(19), 1245-1250; Gordis, E., Dufour, M. C., Warren, K.

- R., Jackson, R. J., Floyd, R. L. and Hungerford, D. W. (1995). Should physicians counsel patients to drink alcohol? JAMA, 273(18), 1415; Colsher, P. L. and Wallace, R. B. (1989). Is modest alcohol consumption better than none at all? An epidemiological assessment. Annual Review of Public Health, 10, 203-219.
22. Fuchs, C. S., Stampfer, M. J., Colditz, G. A., Giovannucci, E. L., Manson, J. E., Kawachi, I., Hunter, D. J., Hankinson, S. E., Hennekens, C. H., Rosner, B., Speizer, F. E. and Willett, W. C. (1995). Alcohol consumption and mortality among women. New England Journal of Medicine, 332(19), 1245-1250.
23. Sanchez-Craig, M., Wilkinson, D. A. and Davila, R. (1995). Empirically based guidelines for moderate drinking: 1-year results from three studies with problem drinkers. American Journal of Public Health, 85(6), 823-828.
24. Wechsler, H., Dowdall, G. W., Davenport, A. and Rimm, E. B. (1995). A gender-specific measure of binge drinking among college students. American Journal of Public Health, 85(7), 982-985.
25. Wechsler, H., Dowdall, G. W., Davenport, A. and Rimm, E. B. (1995). A gender-specific measure of binge drinking among college students. American Journal of Public Health, 85(7), 982-985.
26. Wechsler, H., Dowdall, G. W., Davenport, A. and Rimm, E. B. (1995). A gender-specific measure of binge drinking among college students. American Journal of Public Health, 85(7), 982-985.
27. Lowry, R., Holtzmann, D., Truman, B. I., Kann, L., Collins, J. L. and Kolbe, L. J. (1994). Substance use and HIV-related sexual behaviors among U.S. high school students: Are they related? American Journal of Public Health, 84(7), 1116-1120; Mott, F. L. and Haurin, R. J. (1988). Linkages between sexual activity and alcohol and drug use among American adolescents. Family Planning Perspectives, 20(3), 128-136; Ford, K. and Norris, A. (1994). Urban minority youth: Alcohol and marijuana use and exposure to unprotected intercourse. Journal of Acquired Immune Deficiency Syndromes, 7, 389-396; Robertson, J. A. and Plant, M. A. (1988). Alcohol, sex and risks of HIV infection. Drug and Alcohol Dependence, 22(1-2), 75-78.
28. Leigh, B. C. (1990). Alcohol and unsafe sex: An overview of research and theory. Progress in Clinical and Biological Research, 325, 35-46; Senf, J. H. and Price, C. Q. (1994). Young adults, alcohol and condom use: What is the connection? Journal of Adolescent Health, 15, 238-244; Flanigan, B. J. and Hitch, M. A. (1986). Alcohol use, sexual intercourse and contraception: An exploratory study. Journal of Alcohol and Drug Education, 31(3), 6-40.
29. Flanigan, B. J. and Hitch, M. A. (1986). Alcohol use, sexual intercourse and contraception: An exploratory study. Journal of Alcohol and Drug Education, 31(3), 6-40.

30. Norris, J. (1994). Alcohol and female sexuality: A look at expectancies and risks. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 197-201). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Finnegan, L. P., Davenny, K. and Hartel, D. (1993). Drug use in HIV-infected women. In F. D. Johnstone and M. Johnson (Eds.), HIV in infected women (pp. 133-155). Edinburgh, Great Britain: Churchill Livingstone; Boscarino, J. A., Avins, A. L., Woods, W. J., Lindan, C. P., Hudes, E. S. and Clark, W. (1995). Alcohol-related risk factors associated with HIV infection among patients entering alcoholism treatment: Implications for prevention. Journal of Studies on Alcohol, 56(6), 642-653; Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse; Lowry, R., Holtzmann, D., Truman, B. I., Kann, L., Collins, J. L. and Kolbe, L. J. (1994). Substance use and HIV-related sexual behaviors among U.S. high school students: Are they related? American Journal of Public Health, 84(7), 1116-1120; Leigh, B. C. and Stall, R. (1993). Substance use and risky sexual behavior for exposure to HIV: Issues in methodology, interpretation and prevention. American Psychologist, 48(10), 1035-1045; Leigh, B. C. (1991). Review of research on adolescents. Alcohol Health and Research World, 15(1), 59-63; Schilling, R. F., Schinke, S. P., Nichols, S. E., Zayas, L. H., Miller, S. O., Orlandi, M. A. and Botvin, G. J. (1989). Developing strategies for AIDS prevention research with black and Hispanic drug users. Public Health Reports, 104(1), 2-11.
31. Berenson, A. B., Wilkinson, G. S. and Lopez, L. A. (1995). Substance use during pregnancy and peripartum complications in a triethnic population. International Journal of Addictions, 30(2), 135-145; Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse.
32. U.S. Department of Health and Human Services, Public Health Service. (1995, September). Sexually transmitted disease surveillance, 1994. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.
33. Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse.
34. Leigh, B. C. and Stall, R. (1993). Substance use and risky sexual behavior for exposure to HIV: Issues in methodology, interpretation and prevention. American Psychologist, 48(10), 1035-1045.
35. Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146; Wilsnack, S. C. (1984). Drinking, sexuality and sexual dysfunction in women. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 189-227). New York: Guilford Press.

36. Norris, J. (1994). Alcohol and female sexuality: A look at expectancies and risks. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 197-201). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; George, W. J. and Norris, J. (1991). Alcohol, disinhibition, sexual arousal and deviant sexual behavior. Alcohol Health and Research World, 15(2), 133-138; Malatesta, V. J., Pollack, R. H., Crotty, T. D. and Peacock, L. J. (1982). Acute alcohol intoxication and female orgasmic response. Journal of Sex Research, 18(1), 1-17; Marlatt, G. A. and Rohsenow, D. J. (1980). Cognitive processes in alcohol use: Expectancy and the balanced placebo design. Advances in Substance Abuse, 1 (3-4), 159-199; Wilson, G. T. and Lawson, D. M. (1976). Effects of alcohol on sexual arousal in women. Journal of Abnormal Psychology, 85(5), 489-497; Marlatt, G. A. (1984). Sex differences in psychosocial alcohol research: Implications for prevention. In Women and alcohol: Health related issues: Proceedings of a conference May 23-25, 1984. NIDA research monograph 16 (pp. 260-271.). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism.
37. Gomberg, E. S. L. (1991). Women and alcohol: Psychosocial aspects. In D. J. Pittman and H. R. White (Eds.), Society, culture and drinking patterns reexamined (pp. 263-84). New Brunswick, NJ: Alcohol Research Documentation, Inc., Rutgers Center of Alcohol Studies; George, W. J. and Norris, J. (1991). Alcohol, disinhibition, sexual arousal and deviant sexual behavior. Alcohol Health and Research World, 15(2), 133-138; Kushner, M. G., Sher, K. J. and Beitman, B. D. (1990). The relation between alcohol problems and the anxiety disorders. American Journal of Psychiatry, 147(6), 685-695; Abrams, D. B. and Wilson, G. T. (1979). Effects of alcohol and social anxiety in women: Cognitive versus physiological processes. Journal of Abnormal Psychology, 88(2), 161-173.
38. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; Norris, J. (1994). Alcohol and female sexuality: A look at expectancies and risks. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 197-201). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Wilsnack, S. C. (1984). Drinking, sexuality and sexual dysfunction in women. In S.C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp.189-227). New York: Guilford Press.
39. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146; Norris, J. (1994). Alcohol and female sexuality: A look at expectancies and risks. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 197-201). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism.
40. Klassen, A. D. and Wilsnack, S. C. (1986). Sexual experience and drinking among women in a U. S. national survey. Archives of Sexual Behavior, 15(5), 363-392.

41. Klassen, A. D. and Wilsnack, S. C. (1986). Sexual experience and drinking among women in a U. S. national survey. Archives of Sexual Behavior, 15(5), 363-392.
42. Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146.
43. Leigh, B. C. (1995). A thing so fallen, and so vile: Images of drinking and sexuality in women. Contemporary Drug Problems, 22(3), 415-434; Norris, J. (1994). Alcohol and female sexuality: A look at expectancies and risks. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 197-201). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146; George, W. H., Gournic, S. J. and McAfee, M. (1988). Perceptions of postdrinking female sexuality: Effects of gender, beverage choice and drink payment. Journal of Applied Social Psychology, 18(5), 1295-1317.
44. Miller, B. A. and Downs, W. R. (1993). The impact of family violence on the use of alcohol by women. Alcohol, Health and Research World, 17(2), 137-143; Miller, B. A. (1990). The interrelationships between alcohol and drugs and family violence. In M. De La Rosa, E. Y. Lambert and B. Gropper (Eds.), Drugs and violence: Causes, correlates and consequences. NIDA research monograph 103 (pp. 177-207). Washington, DC: National Institute of Drug Abuse; Miller, B. A., Downs, W. R. and Gondoli, D. M. (1989). Spousal violence among alcoholic women as compared to a random household sample of women. Journal of Studies on Alcohol, 50(6), 533-540; Leonard, K. and Jacob, T. (1988). Alcohol, alcoholism and family violence. In V. B. Van Hasselt, R. K. Morrison, Bellack, A. S. and M. Hersen (Eds.), Handbook of family violence (pp. 383-406). New York: Plenum Press; Koss, M. P. (1988). Hidden rape: Sexual aggression and victimization in a national sample of students in higher education. In A. W. Burgess (Ed.), Rape and sexual assault, volume 2 (pp. 3-25). New York: Garland Publishing; Kantor, G. K. and Straus, M. A. (1989). Substance abuse as a precipitant of wife abuse victimizations. American Journal of Drug Abuse, 15(2), 173-189; Muehlenhard, C. L. and Linton, M. A. (1987). Date rape and sexual aggression in dating situations: Incidence and risk factors. Journal of Counseling Psychology, 34(2), 186-196; Frieze, I. H. and Schafer, P. C. (1984). Alcohol use and marital violence: Female and male differences in reactions to alcohol. In S.C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp.260-279). New York: Guilford Press.
45. Collins, J. J. and Messerschmidt, P. M. (1993). Epidemiology of alcohol-related violence. Alcohol Health and Research World, 17(2), 93-100; Koss, M. P. (1988). Hidden rape: Sexual aggression and victimization in a national sample of students in higher education. In A. W. Burgess (Ed.), Rape and sexual assault, volume 2 (pp. 3-25). New York: Garland Publishing; Rada, R. T. (1975). Alcoholism and forcible rape. American Journal of Psychiatry, 132(4), 444-446; Frieze, I. H. and Schafer, P. C. (1984). Alcohol use and marital violence: Female and male differences in reactions to alcohol. In S.C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and

- intervention (pp.260-279). New York: Guilford Press; Groth, A. N. (1978). The older rape victim and her assailant. Journal of Geriatric Psychiatry, 11(2), 203-215.
46. Koss, M. P. (1988). Hidden rape: Sexual aggression and victimization in a national sample of students in higher education. In A. W. Burgess (Ed.), Rape and sexual assault, volume 2 (pp. 3-25). New York: Garland Publishing.
47. Abbey, A., Ross, L. T. and McDuffie, D. (1994). Alcohol's role in sexual assault. In R. R. Watson (Ed.), Addictive behaviors in women (pp.97-124). Totowa, NJ: Humana Press; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; Norris, J. (1994). Alcohol and female sexuality: A look at expectancies and risks. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 197-201). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146; George, W. H., Gournic, S. J. and McAfee, M. P. (1988). Perceptions of postdrinking female sexuality: Effects of gender, beverage choice and drink payment. Journal of Applied Psychology, 18(5), 1295-1317; Frieze, I. H. and Schafer, P. C. (1984). Alcohol use and marital violence: Female and male differences in reactions to alcohol. In S.C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp.260-279). New York: Guilford Press.
48. Richardson, D. C. and Campbell, J. L. (1980). Alcohol and wife abuse: The effect of alcohol on attributions of blame for wife abuse. Personality and Social Psychology Bulletin, 6(1), 51-56.
49. Richardson, D., Campbell and J. L. (1982). Alcohol and rape: The effect of alcohol on attributions of blame for rape. Personality and Social Psychology Bulletin, 8(3), 468-477.
50. National Institute of Health and Human Services. (1990). Seventh special report to the U.S. Congress on alcohol and health. Rockville, MD: National Institute of Health and Human Services; Kantor, G. K. and Straus, M. A. (1989). Substance abuse as a precipitant of wife abuse victimizations. American Journal of Drug Abuse, 15(2), 173-189; Frieze, I. H. and Schafer, P. C. (1984). Alcohol use and marital violence: Female and male differences in reactions to alcohol. In S.C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp.260-279). New York: Guilford Press.
51. Frieze, I. H. and Schafer, P. C. (1984). Alcohol use and marital violence: Female and male differences in reactions to alcohol. In S.C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp.260-279). New York: Guilford Press.

52. Miller, B. A. (1990). The interrelationships between alcohol and drugs and family violence. In M. De La Rosa, E. Y. Lambert and B. Gropper (Eds.), Drugs and violence: Causes, correlates and consequences. NIDA research monograph 103 (pp. 177-207). Washington, DC: National Institute of Drug Abuse; Kilpatrick, D. G., Resnick, H. S., Saunders, B. E., Best, C. L. and Epstein, J. (1994, June). Violent assault and alcohol dependence among women: Results of a longitudinal study. Paper presented at the Annual Meeting of the Research Society on Alcoholism, Maui, HI; Haver, B. (1987). Female alcoholics: IV. The relationship between family violence and outcome 3-10 years after treatment. Acta Psychiatrica Scandinavia, 75(5), 449-455; Smith, P. H. and Chescheir, N. (1995). Battered substance-abusing women: A vulnerable and underserved population in North Carolina: Final report submitted to North Carolina Governor's Institute for Alcohol and Substance Abuse. Raleigh, NC: North Carolina Governor's Institute for Alcohol and Substance Abuse.
53. Gomberg, E. S. and Nirenberg, T. D. (1993). Antecedents and consequences. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 118-141). Norwood, NJ: Ablex Publishing; Robbins, C. (1989). Sex differences in psychological consequences of alcohol and drug abuse. Journal of Health and Social Behavior, 30(1), 117-130; Malin, H., Wilson, R., Williams, G. and Aitken, S. (1985). 1983 alcohol/health practices supplement. Alcohol Health and Research World, 10(1), 48-53; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press; Dunham, R. G. (1986). Noticing alcoholism in the elderly and women: A nationwide examination of referral behavior. Journal of Drug Issues, 16(3), 397-406.
54. Robbins, C. (1989). Sex differences in psychological consequences of alcohol and drug abuse. Journal of Health and Social Behavior, 30(1), 117-130; Gomberg, E. S. L. and Nirenberg, T. D. (1993). Antecedents and consequences. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 118-141). Norwood, NJ: Ablex Publishing; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press; Dunham, R. G. (1986). Noticing alcoholism in the elderly and women: A nationwide examination of referral behavior. Journal of Drug Issues, 16(3), 397-406.
55. Bergmann, P. E., Smith, M. B. and Hoffman, N. G. (1995). Adolescent treatment: Implications for assessment, practice guidelines and outcome management. Pediatric Clinics of North America, 42(2), 453-472.
56. Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.

57. Gomberg, E. S. L. (1989). Suicide risk among women with alcohol problems. American Journal of Public Health, 79(10), 1363-1365; Murphy, G. E. and Wtzel, R. D. (1990). The lifetime risk of suicide in alcoholism. Archives of General Psychiatry, 47(4), 383-392; Wilsnack, R. W., Klassen, A. D. and Wilsnack, S. C. (1986). Retrospective analysis of lifetime changes in women's drinking behavior. Advances in Alcohol and Substance Abuse, 5(3), 9-28.
58. Gomberg, E. S. L. (1989). Suicide risk among women with alcohol problems. American Journal of Public Health, 79(10), 1363-1365.
59. Bergmann, P. E., Smith, M. B. and Hoffman, N. G. (1995). Adolescent treatment: Implications for assessment, practice guidelines and outcome management. Pediatric Clinics of North America, 42(2), 453-472; Garrison, C. Z., McKeown, R. E., Valois, R. F. and Vincent, M. L. (1993). Aggression, substance use and suicidal behaviors in high school students. American Journal of Public Health, 83(2), 179-184; Silverman, W. H. (1991). Prevalence of substance use in a rural teenage population. Journal of Adolescent Chemical Dependency, 2(1), 107-117.
60. Bergmann, P. E., Smith, M. B. and Hoffman, N. G. (1995). Adolescent treatment: Implications for assessment, practice guidelines and outcome management. Pediatric Clinics of North America, 42(2), 453-472.
61. Centers for Disease Control and Prevention. (1995). Fatal and nonfatal suicide attempts among adolescents--Oregon, 1988-1993. Morbidity and Mortality Weekly Report, 44(16), 289-291; Downey, A. M. (1990-1991). The impact of drug abuse upon adolescent suicide. Omega, 22(4), 261-275.
62. Centers for Disease Control and Prevention. (1995). Suicide among children, adolescents and young adults--United States, 1980-1992. Morbidity and Mortality Weekly Report, 44(15), 289-291.
63. Hill, S. Y. (1984). Physiological effects of alcohol on women. In Women and alcohol: Health-related issues. Proceedings of a conference, May 23-25, 1984, NIDA research monograph 16 (pp. 199-214). Rockville, MD: National Institute on Drug Abuse.
64. Gomberg, E. S. L. (1994). Risk factors for drinking over a woman's life span. Alcohol Health and Research World, 18(3), 220-227; Windle, M. and Barnes, G. M. (1988). Similarities and differences in correlates of alcohol consumption and problem behaviors among male and female adolescents. International Journal of Addictions, 23(7), 707-728; Harrison, P. A. (1989). Women in treatment: Changing over time. International Journal of the Addictions, 24(7), 655-673.
65. Gomberg, E. S. L. (1994). Risk factors for drinking over a woman's life span. Alcohol Health and Research World, 18(3), 220-227.

66. Gomberg, E. S. and Nirenberg, T. D. (1993). Antecedents and consequences. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 118-141). Norwood, NJ: Ablex Publishing.
67. Harrison, P. A. (1989). Women in treatment: Changing over time. International Journal of the Addictions, 24(7), 655-673.
68. Centers for Disease Control. (1992). Trends in alcohol-related traffic fatalities, by sex--United States. Morbidity and Mortality Weekly Report, 41(11), 189, 195-197; Popkin, C. L. (1991). Drinking and driving by young females. Accident Analysis and Prevention, 23(1), 37-44.
69. Popkin, C. L. (1991). Drinking and driving by young females. Accident Analysis and Prevention, 23(1), 37-44; Gomberg S. L. and Nirenberg, T. D. (1993). Antecedents and consequences. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 118-141). Norwood, NJ: Ablex Publishing; Waller, P. F. and Blow, F. C. (1995). Women, alcohol and driving. In M. Galanter (Ed.), Alcoholism and women (pp. 103-123). New York: Plenum Press.
70. Popkin, C. L. (1991). Drinking and driving by young females. Accident Analysis and Prevention, 23(1), 37-44.
71. Yu, J., Essex, D. T. and Williford, W. R. (1992). DWI/DWAI offenders and recidivism by gender in the eighties: A changing trend? International Journal of Addictions, 27(6), 637-647.
72. Surgeon General. (1980). The health consequences of smoking for women. Washington, DC: U. S. Department of Health and Human Services, Office on Smoking and Health.
73. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. (1987). Vital Statistics of the United States, 1985, volume II: Mortality. Hyattsville, MD: National Center for Health Statistics; U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. (1988). Vital Statistics of the United States, 1986, volume II: Mortality. Hyattsville, MD: National Center for Health Statistics; U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. (1989). Vital Statistics of the United States, 1987, volume II: Mortality. Hyattsville, MD: National Center for Health Statistics.
74. Thun, M. J., Day-Lilly, C. A., Calle, E. E., Flanders, W. D. and Heath, C. W. (1995). Excess mortality among cigarette smokers: Changes in a 20-year interval. American Journal of Public Health, 85(9), 1223-1230.
75. Husten, C. G., Chrismon, J. H. and Reddy, M. N. (1996). Trends and effects of cigarette smoking among girls and women in the United States, 1965-1993. Journal of the

- Medical American Women's Association, 51 (1), 1-8; Centers for Disease Control and Prevention. (1995). Indicators of nicotine addiction among women: United States, 1991-1992. Morbidity and Mortality Weekly Report, 44(6), 103-105.
76. American Heart Association (1993). Heart and stroke facts: 1994 statistical supplement. Dallas, TX: American Heart Association; Gardner, P. and Hudson, B. L. (1996). Advance report of final mortality statistics, 1993. Monthly Vital Statistics Report, 44(7), 1-84.
77. Solomon, L. J. and Flynn, B. S. (1993). Women who smoke. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 339-349). New York: Oxford University Press; Ramirez, A. G. and Gallion, K. J. (1993). Nicotine dependence among blacks and Hispanics. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 350-364). New York: Oxford University Press.
78. English, P. B., Eskenazi, B. and Christianson, R. E. (1994). Black-white differences in serum cotinine levels among pregnant women and subsequent effects on infant birthweight. American Journal of Public Health, 84(9), 1439-1443.
79. Royce, J. M., Hymowitz, N., Corbett, K., Hartwell, T. D. and Orlandi, M. A. (1993). Smoking cessation factors among African Americans and whites. American Journal of Public Health, 83(2), 220-226; Solomon, L. J. and Flynn, B. S. (1993). Women who smoke. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 339-349). New York: Oxford University Press. But a recent study failed to link this preference to higher nicotine levels in black women. See Ahijevych, K., Gillespie, J., Demirchi, M. and Jagadeesh, J. (1996). Menthol and nonmenthol cigarettes and smoke exposure in black and white women. Pharmacology, Biochemistry and Behavior, 53(2), 355-360.
80. DeNavas, C. and U.S. Bureau of the Census. (1995). Income, poverty, and valuation of noncash benefits: 1993. Washington, DC: U.S. Department of Commerce, Economics and Statistics Administration, Bureau of the Census [Supt. of Docs., U.S. G.P.O., distributor].
81. Rich-Edwards, J. W., Manson, J. E., Hennekens, C. H. and Buring, J. E. (1995). The primary prevention of coronary heart disease in women. New England Journal of Medicine, 332(26), 1758-1766; Rosenberg, L., Palmer, J. R. and Shapiro, S. (1990). Decline in the risk of myocardial infarction among women who stop smoking. New England Journal of Medicine, 322(4), 213-217.
82. Willett, W. C., Green A., Stampfer, M. J., Speizer, F., et al. (1987). Relative and absolute excess risks of coronary heart disease among women who smoke cigarettes. New England Journal of Medicine, 317(21), 1303-1309.
83. Shopland, D. R. and U.S. Department of Health and Human Services. (1983). The health consequences of smoking: Cardiovascular disease: A report of the Surgeon General. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Office on Smoking and Health; Layde, P. M. and Beral, V. (1981). Further analysis of

mortality in oral contraceptive users: Royal College of General Practitioners' Oral Contraception Study. Lancet, 1(8219), 541-546.

84. Hopper, J. L. and Seeman, E. (1994). The bone density of female twins discordant for tobacco use. New England Journal of Medicine, 330(6), 387-392; Baron, J. A., La Vecchia, C. and Levi, F. (1990). The antiestrogenic effects of cigarette smoking in women. American Journal of Obstetrics, 162(2), 502-514; Slemenda, C. W. (1994). Cigarettes and the skeleton. New England Journal of Medicine, 330(6), 430.

85. Hopper, J. L. and Seeman, E. (1994). The bone density of female twins discordant for tobacco use. New England Journal of Medicine, 330(6), 387-392.

86. Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: U.S. Department of Health and Human Services, Office on Smoking and Health; Hopper, J. L. and Seeman, E. (1994). The bone density of female twins discordant for tobacco use. New England Journal of Medicine, 330(6), 387-392.

87. Katelaris, A. G. and Cumming, R. G. (1996). Health status before and mortality after hip fracture. American Journal of Public Health 86(4), 557-559.

88. Nelson, H. D., Nevitt, M. C., Scott, J. C., Stone, K. L. and Cummings, S. R. (1994). Smoking, alcohol and neuromuscular function of older women. JAMA, 272(23), 1825-1832.

89. Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: U.S. Department of Health and Human Services, Office on Smoking and Health; Midgette, A. S. and Baron, J. A. (1990). Cigarette smoking and the risk of natural menopause. Epidemiology, 1(6), 474-480; Willett, W., Stampfer, M. J., Bain, C., Lipnick, R., Speizer, F. E., Rosner, B., Cramer, D. and Hennekens, C. H. (1983). Cigarette smoking, relative weight and menopause. American Journal of Epidemiology, 117(6), 651-658; Slemenda, C. W. (1994). Cigarettes and the skeleton. New England Journal of Medicine, 330(6), 430.

90. Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: U.S. Department of Health and Human Services, Office on Smoking and Health.

91. Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: U.S. Department of Health and Human Services, Office on Smoking and Health.

92. Slattery, M. L., Robison, L. M., Schuman, K. L., French, T. K., Abbott, T. M., Overall, J. C. and Gardner, J. W. (1989). Cigarette smoking and exposure to passive smoke are risk factors for cervical cancer. JAMA, 261(11), 1593-1598; Sood, A. K. (1991). Cigarette smoking and cervical cancer: Meta-analysis and critical review of recent studies. American Journal of Preventive Medicine, 7(4), 208-213; Licciardone, J. C., Brownson, R. C., Chang, J. C. and Wilkens, J. R. (1990). Uterine cervical cancer risk in cigarette smokers: A meta-analytic study. American Journal of Preventive Medicine, 6(5), 274-281; Layde, P. M. and Broste, S. K. (1989). Carcinoma of the cervix and smoking. Biomedicine

and Pharmacotherapy, 43(3), 161-165; Layde, P. M. (1989). Smoking and cervical cancer: Cause or coincidence? JAMA, 261(11), 1631-1632; Brinton, L. A., Schairer, C., Haenszel, W., Stolley, P., Lehman, H. F., Levine, R. and Savitz, D. A. (1986). Cigarette smoking and invasive cervical cancer. JAMA, 255(23), 3265-3269; Winkelstein, W. (1986). Cigarette smoking and cancer of the uterine cervix. In D. Hoffman and C. C. Harris (Eds.), Mechanisms in tobacco carcinogenesis (pp. 329-341). Banbury Report, 23. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory; Winkelstein, W. (1990). Smoking and cervical cancer--current status: A review. American Journal of Epidemiology, 131(6), 945-957; Holt, V. L., Daling, J. R., McKnight, B., Moore, D. E., Stergachis, A. and Weise, N. S. (1994). Cigarette smoking and functional ovarian cysts. American Journal of Epidemiology, 139(8), 781-786.

93. Muller, B., Zulewski, H., Huber, P., Ratcliffe, J. G. and Staub, J. J. (1995). Impaired action of thyroid hormone associated with smoking in women with hypothyroidism. New England Journal of Medicine, 333(15), 964-969.

94. Centers for Disease Control and Prevention. (1995). Indicators of nicotine addiction among women--United States, 1991-1992. Morbidity and Mortality Weekly Report, 44(6), 102-105.

95. Brown, E. R., Wyn, R., Cumberland, W. G., Yu, H., Abel, E., Gelberg, L., and Ng, L. (1995). Women's health-related behaviors and use of clinical preventive services: A report to the Commonwealth Fund. Los Angeles: UCLA Center for Health Policy Research.

96. Brown, E. R., Wyn, R., Cumberland, W. G., Yu, H., Abel, E., Gelberg, L., and Ng, L. (1995). Women's health-related behaviors and use of clinical preventive services: A report to the Commonwealth Fund. Los Angeles: UCLA Center for Health Policy Research.

97. Flegal, K. M., Troiano, R. P., Pamuk, E. R., Kuczmarskia, R. J. and Campbell, S. M. (1995). The influence of smoking cessation on the prevalence of overweight in the United States. New England Journal of Medicine, 333(18), 1165-1170.

98. Flegal, K. M., Troiano, R. P., Pamuk, E. R., Kuczmarskia, R. J. and Campbell, S. M. (1995). The influence of smoking cessation on the prevalence of overweight in the United States. New England Journal of Medicine, 333(18), 1165-1170; Fisher, E. B., Lichtenstein, E. and Haire-Joshu, D. (1993). Multiple determinants of tobacco use and cessation. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 59-88). New York: Oxford University Press; Ricotti, N. A. (1989). Cigarette smoking and body weight. New England Journal of Medicine, 320(14), 931-933; Gritz, E. R. and St. Jeor, S. T. (1992). Task force 3: Implications with respect to intervention and prevention. Health Psychology, 11(Suppl.), 17-25.

99. Flegal, K. M., Troiano, R. P., Pamuk, E. R., Kuczmarskia, R. J. and Campbell, S. M. (1995). The influence of smoking cessation on the prevalence of overweight in the United States. New England Journal of Medicine, 333(18), 1165-1170.

100. Manson, J. E., Willett, W. C., Stampfer, M. J., Colditz, G. A., Hunter, D. J., Hankinson, S. E., Hennekens, C. H. and Speizer, F. E. (1995). Body weight and mortality among women. New England Journal of Medicine, 333(11), 677-685.
101. Fisher, E. B., Lichtenstein, E. and Haire-Joshu, D. (1993). Multiple determinants of tobacco use and cessation. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 59-88). New York: Oxford University Press.
102. Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: U.S. Department of Health and Human Services, Office on Smoking and Health.
103. Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: U.S. Department of Health and Human Services, Office on Smoking and Health.
104. Kawachi, I., Colditz, G. A., Stampfer, M. J., Willett, W. C., Manson, J. E., Rosner, B., Speizer, F. E. and Hennekens, C. H. (1993). Smoking cessation and decreased risk of stroke in women. JAMA, 269(2), 232-236.
105. Centers for Disease Control and Prevention. (1995). Indicators of nicotine addiction among women--United States, 1991-1992. Morbidity and Mortality Weekly Report, 44(6), 102-105.
106. Rosenberg, L., Palmer, J. R. and Shapiro, S. (1990). Decline in the risk of myocardial infarction among women who stop smoking. New England Journal of Medicine, 322(4), 213-217.
107. Willett, W. C., Green A., Stampfer, M. J., Speizer, F., et al. (1987). Relative and absolute excess risks of coronary heart disease among women who smoke cigarettes. New England Journal of Medicine, 317(21), 1303-1309; Rich-Edwards, J. W., Manson, J. E., Hennekens, C. H. and Buring, J. E. (1995). The primary prevention of coronary heart disease in women. New England Journal of Medicine, 332(26), 1758-1766.
108. Licciardone, J. C., Brownson, R. C., Chang, J. C. and Wilkens, J. R. (1990). Uterine cervical cancer risk in cigarette smokers: A meta-analytic study. American Journal of Preventive Medicine, 6(5), 274-281.
109. Trachtenberg, A. I. and Fleming, M. F. (1994). Diagnosis and treatment of drug abuse in family practice. Kansas City, Missouri: American Academy of Family Physicians; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO; Smart, R. G. (1991). Crack cocaine use: A review of prevalence and adverse affects. American Journal of Drug and Alcohol Abuse, 17(1), pp. 13-26; Finnegan, L. P., Davenny, K. and Hartel, D. (1993). Drug use in HIV-infected women. In F. D. Johnstone and M. Johnson (Eds.), HIV in infected women (pp. 133-155). Edinburgh, Great Britain: Churchill Livingstone; Lynn, L. A. and Mackler, S. A. (1992). Cocaine and the spread of AIDS. Hospital Practice, 105-113.

110. Trachtenberg, A. I. and Fleming, M. F. (1994). Diagnosis and treatment of drug abuse in family practice. Kansas City, Missouri: American Academy of Family Physicians; Finnegan, L. P., Davenny, K. and Hartel, D. (1993). Drug use in HIV-infected women. In F. D. Johnstone and M. Johnson (Eds.), HIV in infected women (pp. 133-155). Edinburgh, Great Britain: Churchill Livingstone; Glantz, J. C. and Woods, J. R. (1993). Cocaine, heroin and phencyclidine: Obstetric perspectives. Clinical Obstetrics and Gynecology, 36(2), 279-301; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO; Smart, R. G. (1991). Crack cocaine use: A review of prevalence and adverse affects. American Journal of Drug and Alcohol Abuse, 17(1), 13-26; Des Jarlais, D. C. (1991). Potential cofactors in the outcomes of HIV infection in intravenous drug users. In P. Hartsock and S. G. Genser (Eds.), Longitudinal studies of HIV infection in intravenous drug users: Methodological issues in natural history research. NIDA research monograph 109 (pp. 155-123). Rockville, MD: National Institute on Drug Abuse.
111. Lex, B. W. (1995). Alcohol and other psychoactive substance dependence in women and men. In M. V. Seeman (Ed.), Gender and psychopathology (pp.311-358). Washington, DC: American Psychiatric Press; Anglin, M. C. and Fisher, D. G. (1993). Alcohol and heroin use patterns of narcotics addicts: Gender and ethnic differences. American Journal of Drug and Alcohol Abuse, 19(2), 219-238; Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse; Griffin, M. L., Weiss, R. D., Mirin, S. M. and Lange, U. (1989). A comparison of male and female cocaine abusers. Archives of General Psychiatry, 46, 122-126.
112. Morokoff, P. J., Harlow, L. L. and Quina, K. (1995). Women and AIDS. In A. L. Stanton and S. J. Gallant (Eds.), Psychology of women's health: Progress and challenges in research and application (pp. 117-169). Washington, DC: American Psychological Association; Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse; Booth, R. E., Koester, S. K. and Pinto, F. (1995). Gender differences in sex-risk behaviors, economic livelihood and self-concept among drug injectors and crack smokers. American Journal of Addictions, 4(4), 313-322; Freeman, R. C., Rodriguez, G. M. and French, J. F. (1994). A comparison of male and female intravenous drug users' risk behaviors for HIV infection. American Journal of Drug and Alcohol Abuse, 20(2), 129-157; Booth, R. E. (1995). Gender differences in high-risk sex behaviors among heterosexual drug injectors and crack smokers. American Journal of Drug and Alcohol Abuse, 21(4), 419-432; Kail, B. L., Watson, D. D. and Ray, S. R. (1995). Needle-using practices within the sex industry. American Journal of Alcohol Abuse, 21(2), 241-255; Joe, G. W. and Simpson, D. D. (1995). HIV risks, gender and cocaine use among opiate users. Drug and Alcohol Dependence, 37, 23-28; Astemborski, J., Vlahov, D., Warren, D., Solomon, L. and Nelson, K. E. (1994). The trading of sex for drugs or money and HIV seropositivity among female intravenous drug users. American Journal of Public Health, 84(3), 382-387.

113. Centers for Disease Control. (1995). Update: AIDS among women--United States, 1994. Morbidity and Mortality Weekly Report, 44(5), 81-84; Melnick, S. L., Sherer, R., Louis, T. A., Hillman, D., Rodriguez, E. M., Lackman, C., Capps, L., Brown, L. S., Carlyn, M., Korvick, J. A. and Deyton, L. (1994). Survival and disease progression according to gender of patients with HIV infection. JAMA, 272(24), 1915-1922; U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Treatment. (1994). Practical approaches in the treatment of women who abuse alcohol and other drugs. Rockville, MD: U.S. Department of Health and Human Services; Cohen, J. B., Hauer, L. B. and Wofsy, C. B. (1989). Women and IV drugs: Parenteral and heterosexual transmission of human immunodeficiency virus. Journal of Drug Issues, 19(1), 39-56; Finnegan, L. P., Davenny, K. and Hartel, D. (1993). Drug use in HIV-infected women. In F. D. Johnstone and M. Johnson (Eds.), HIV in infected women (pp. 133-155). Edinburgh, Great Britain: Churchill Livingstone.
114. Centers for Disease Control. (1995). Update: AIDS among women--United States, 1994. Morbidity and Mortality Weekly Report, 44(5), 81-84.
115. Centers for Disease Control and Prevention. (1996). Update: Mortality attributable to HIV infection among persons aged 25-44--United States, 1994. Morbidity and Mortality Weekly Report, 45(6), 122-125.
116. Davis, S.F., Byers, R.H., Lindegren, M.L., Caldwell, M.B., Karon, J.M., and Gwinn, M. (1995). Prevalence and incidence of vertically acquired HIV infection in the United States. JAMA 274(12), 952-955.
117. Centers for Disease Control and Prevention. (1995). First 500,000 AIDS cases--United States, 1995. Morbidity and Mortality Weekly Report, 44(8), 849-853; Finnegan, L. P., Davenny, K. and Hartel, D. (1993). Drug use in HIV-infected women. In F. D. Johnstone and M. Johnson (Eds.), HIV in infected women (pp. 133-155). Edinburgh, Great Britain: Churchill Livingstone.
118. Centers for Disease Control and Prevention. (1995). Update: AIDS among women--United States, 1994. Morbidity and Mortality Weekly Report, 44(5), 81-84; Centers for Disease Control and Prevention. (1995). First 500,000 AIDS cases--United States, 1995. Morbidity and Mortality Weekly Report, 44(8), 849-853.
119. Centers for Disease Control. (1995). Update: AIDS among women--United States, 1994. Morbidity and Mortality Weekly Report, 44(5), 81-84.
120. Centers for Disease Control and Prevention. (1996). Update: Mortality attributable to HIV infection among persons aged 25-44--United States, 1994. Morbidity and Mortality Weekly Report, 45(6), 122-125.
121. Berenson, A. B., Wilkinson, G. S. and Lopez, L. A. (1995). Substance use during pregnancy and peripartum complications in a triethnic population. International Journal of

- Addictions, 30(2), 135-145; Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse.
122. Marx, R., Aral, S. O., Rolfs, R. T., Sterk, C. E. and Kahn, J. G. (1991). Crack, sex and STD. Sexually Transmitted Diseases, 18(2), 92-101; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO; Centers for Disease Control and Prevention. (1994). Sexually transmitted disease surveillance 1994. Atlanta, GA: Centers for Disease Control and Prevention; Lewin, T. (1992, January 11). Syphilis cases among newborns climb with the rise in crack use. New York Times, p. A13.
123. Farley, T. A., Hadler, J. L. and Gunn, R. A. (1990). The syphilis epidemic in Connecticut: Relationship to drug use and prostitution. Sexually Transmitted Diseases, 17(4), 163-168.
124. Cantwell, M. F., Shehab, Z. M., Costello, A. M., Sands, L., Green, W. F., Ewing, E. P., Valway, S. E. and Onorato, I. M. (1994). Brief report: Congenital tuberculosis. New England Journal of Medicine, 330(15), 1051-1054.
125. Mosher, W. D. and Pratt, W. F. (1993). AIDS-related behavior among women 15-44 years of age: United States, 1988 and 1990. In Advance Data No. 239 (pp. 1-16). Hyattsville, MD: National Center for Health Statistics.
126. Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse; Zweben, J. E. (1991). Counseling issues in methadone maintenance treatment. Journal of Psychoactive Drugs, 23(2), 177-190.
127. Wells, D. V. B. and Jackson, J. F. (1992). HIV and chemically dependent women: Recommendations for appropriate health care and treatment services. International Journal of the Addictions, 27(5), 571-585; Schilling, R. F., Schinke, S. P., Nichols, S. E., Zayas, L. H., Miller, S. O., Orlandi, M. A. and Botvin, G. J. (1989). Developing strategies for AIDS prevention research with black and Hispanic drug users. Public Health Reports, 104(1), 2-11; Hardesty, L. and Greif, G. L. (1994). Common themes in a group for female IV drug users who are HIV positive. Journal of Psychoactive Drugs, 26(3), 289-293.
128. Schilling, R. F., Schinke, S. P., Nichols, S. E., Zayas, L. H., Miller, S. O., Orlandi, M. A. and Botvin, G. J. (1989). Developing strategies for AIDS prevention research with black and Hispanic drug users. Public Health Reports, 104(1), 2-11.
129. Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse.

130. Diem, E. C., McKay, L. C. and Jamieson, J. L. (1994). Female adolescent alcohol, cigarette and marijuana use: Similarities and differences in patterns of use. International Journal of the Addictions, 29(8), 987-997; Mott, F. L. and Haurin, J.R. (1988). Linkages between sexual activity and alcohol and drug use among American adolescents. Family Planning Perspectives, 20(3), 128-136.
131. Lex, B. W. (1995). Alcohol and other psychoactive substance dependence in women and men. In M. V. Seeman (Ed.), Gender and psychopathology (pp. 311-358). Washington, DC: American Psychiatric Press; Blume, S. B. (1994). Women and addictive disorders. In N. S. Miller (Ed.), Principles of addiction medicine (pp.1-16). Chevy Chase, MD: American Society of Addiction Medicine; Mello, N. K., Mendelson, J. H. and Teoh, S. K. (1989). Neuroendocrine consequences of alcohol abuse in women. Annals of the New York Academy of Sciences, 562, 211-240.
132. Blume, S. B. (1994). Women and addictive disorders. In N. S. Miller (Ed.), Principles of addiction medicine (pp. 1-16), Chevy Chase, MS: American Society of Addiction Medicine; Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing.
133. Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Obstetrics and Gynecology Clinics of North America, 17(4), 905-25; Erickson, P. G. and Murray, G. F. (1989). Sex differences in cocaine use and experiences: A double standard revived? American Journal of Drug and Alcohol Abuse, 15(2), 135-152; Robbins, C. (1989). Sex differences in psychological consequences of alcohol and drug abuse. Journal of Health and Social Behavior, 30(1), 117-130; Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Health Maintenance Strategies, 17(4), 905-925; New York Times, 2/14/96, p. A16; Daily Mail, 4/18/95, p. 18; Los Angeles Times, 7/7/94, p. E4; The Independent, 8/20/95, p. 3.
134. Robbins, C. (1989). Sex differences in psychological consequences of alcohol and drug abuse. Journal of Health and Social Behavior, 30(1), 117-130.
135. Kandall, S. R. and Chavkin, W. (1992). Illicit drugs in America: History, impact on women and infants, and treatment strategies for women. Hastings Law Journal, 43(3), 615-643.
136. Fullilove, M. T., Weinstein, M., Fullilove, R. E., Crayton, E. J., Goodjoin, R. B., Bowser, B. and Gross, S. (1990). Race/gender issues in the sexual transmission of AIDS. AIDS Clinical Review, 25-62; Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; Booth, R. E. (1995). Gender differences in high-risk sex behaviors among heterosexual drug injectors and crack smokers. American Journal of Drug and Alcohol Abuse, 21(4), 419-432; Cohen, E., Navaline, H. and Metzger, D. (1994). High-risk behaviors for HIV: A comparison between crack-abusing and opioid-abusing African-American women. Journal of Psychoactive Drugs, 26(3), 233-241; Maher, L. and Curtis, R. (1992). Women on the edge of crime: Crack cocaine and the

- changing contexts of street-level sex work in New York City. Crime, Law and Social Change, 18(3), 221-258; Anglin, M. D. and Hser, Y. (1987). Addicted women and crime. Criminology, 25(2), 359-397.
137. McCoy, H. V., Inciardi, J. A., Metsch, L. R., Pottieger, A. E. and Saum, C. A. (1995). Women, crack and crime: Gender comparisons of criminal activity among crack cocaine users. Contemporary Drug Problems, 22(3), 435-451.
138. Maher, L. and Curtis, R. (1992). Women on the edge of crime: Crack cocaine and the changing contexts of street-level sex work in New York City. Crime, Law and Social Change, 18(3), 221-258.
139. Fullilove, M. T., Lown, E. A. and Fullilove, R. E. (1992). Crack 'hos and skeezers: Traumatic experiences of women crack users. Journal of Sex Research, 29(2), 275-287; Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; McCoy, H. V., Inciardi, J. A., Metsch, L. R., Pottieger, A. E. and Saum, C. A. (1995). Women, crack and crime: Gender comparisons of criminal activity among crack cocaine users. Contemporary Drug Problems, 22(3), 435-451; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO; Booth, R. E. (1995). Gender differences in high-risk sex behaviors among heterosexual drug injectors and crack smokers. American Journal of Drug and Alcohol Abuse, 21(4), 419-432.
140. Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing.
141. Lex, B. W. (1995). Alcohol and other psychoactive substance dependence on women and men. In M. V. Seeman (Ed.), Gender and psychopathology (pp. 311-358). Washington, DC: American Psychiatric Press; Maher, L. and Curtis, R. (1992). Women on the edge of crime: Crack cocaine and the changing contexts of street-level sex work in New York City. Crime, Law and Social Change, 18(3), 221-258; Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; McCoy, H. V., Inciardi, J. A., Metsch, L. R., Pottieger, A. E. and Saum, C. A. (1995). Women, crack and crime: Gender comparisons of criminal activity among crack cocaine users. Contemporary Drug Problems, 22(3), 435-451.
142. Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; McCoy, H. V., Inciardi, J. A., Letsch, L. R., Pottieger, A. E. and Saum, C. A. (1995). Women, crack and crime: Gender comparisons of criminal activity among crack cocaine users. Contemporary Drug Problems, 22(3), 435-451; Finnegan, L. P., Sloboda, Z., Haverkos, H. W., Mello, N. K., Kreek, M. J., Cottler, L. B. and Frank, D. A. (1995). Drug abuse and the health of women. In L. S. Harris (Ed.), Problems of drug dependence, 1994: Volume I, NIDA research monograph 152 (pp. 45-52). Rockville, MD: NIDA; Fagan, J. (1994). Women and drugs revisited: Female participation in the cocaine economy. Journal of Drug Issues, 24(2), 179-25; Mieczkowski, T. (1994).

The experiences of women who sell crack: Some descriptive data from the Detroit crack ethnography project. The Journal of Drug Issues, 24(2), 227-24; Erickson, P. G. and Watson, V. A. (1990). Women, illicit drugs and crime. In L. T. Kozlowski, H. M. Annis, et al (Eds.), Research advances in alcohol and drug problems, volume 10 (pp. 251-72). New York: Plenum Press; Anglin, M. D. and Hser, Y. (1987). Addicted women and crime. Criminology, 25(2), 359-397.

143. Prendergast, M. L., Wellisch, J. and Falkin, G. P. (1995). Assessment of and services for substance-abusing women offenders in community settings. Prison Journal, 75(2), 240-256; Kassebaum, G. and Chandler, S. M. (1994). Polydrug use and self control among men and women in prisons. Journal of Drug Education, 24(4), 333-350; Kemper, K. J. and Rivara, F. P. (1993). Parents in jail. Pediatrics, 92(2), 261-264.

144. U.S. Federal Bureau of Investigation. (1994). Uniform crime reports for the United States, 1993. Washington, DC: GPO.

145. U. S. Department of Justice. (1991). Survey of state prison inmates, 1991: Women in prison. Washington, DC: U. S. Department of Justice.

146. Beck, A. J and Gilliard, D. K. (1995, August). Prisoners in 1994. Bureau of Justice Statistics Bulletin, 1-16.

147. U. S. Department of Justice. (1991). Survey of state prison inmates, 1991: Women in prison. Washington, DC: U. S. Department of Justice.

148. Maher, L. and Curtis, R. (1992). Women on the edge of crime: Crack cocaine and the changing contexts of street-level sex work in New York City. Crime, Law and Social Change, 18(3), 221-258.

149. State of New York Department of Correctional Services. (1990, March). Female Drug Commitment Population, 1987-1989. Albany: State of New York Department of Correctional Services, Division of Program Planning, Research and Evaluation.

150. National Institute of Justice. (1995). Drug use forecasting: 1994 annual report on adult and juvenile arrestees. Washington, DC: U.S. Department of Justice, Office of Justice Programs, National Institute of Justice.

151. U. S. Department of Justice. (1991). Survey of state prison inmates, 1991: Women in prison. Washington, DC: U. S. Department of Justice; Prendergast, M. L., Wellisch, J. and Falkin, G. P. (1995). Assessment of and services for substance-abusing women offenders in community settings. Prison Journal, 75(2), 240-256; Wellisch, J., Prendergast, M. L. and Anglin, M. D. (1994). Jail: A found opportunity for intervening with drug-abusing pregnant women. Journal of Correctional Health Care, 1(1), 17-38; Farabee, D. (1995). Substance abuse among female inmates entering the Texas department of criminal justice--Institutional division: 1994. Austin, TX: Texas Commission on Alcohol and Drug Abuse; Maher, L. and

- Curtis, R. (1992). Women on the edge of crime: Crack cocaine and the changing contexts of street-level sex work in New York City. Crime, Law and Social Change, 18(3), 221-258.
152. Kemper, K. J. and Rivara, F. P. (1993). Parents in jail. Pediatrics, 92(2), 261-264.
153. Kemper, K. J. and Rivara, F. P. (1993). Parents in jail. Pediatrics, 92(2), 261-264.
154. Kantor, G. K. and Straus, M. A. (1989). Substance abuse as a precipitant of wife abuse victimizations. American Journal of Drug Abuse, 15(2), 173-189; Regan, D. O., Erlich, S. M. and Finnegan, L. P. (1987). Infants of drug addicts: At risk of child abuse, neglect and placement in foster care. Neurotoxicology and Teratology, 9(4), 315-319; Zweben, J. E. (1991). Counseling issues in methadone maintenance treatment. Journal of Psychoactive Drugs, 23(2), 177-190; Fullilove, M. T., Fullilove, R. E., Smith, M., Winkler, K., Michael, C., Panzer, P. G. and Wallace, R. (1993). Violence, trauma and post-traumatic stress disorder among women drug users. Journal of Traumatic Stress, 6(4), 533-543.
155. Fullilove, M. T., Fullilove, R. E., Smith, M., Winkler, K., Michael, C., Panzer, P. G. and Wallace, R. (1993). Violence, trauma and post-traumatic stress disorder among women drug users. Journal of Traumatic Stress, 6(4), 533-543.
156. Regan, D. O., Erlich, S. M. and Finnegan, L. P. (1987). Infants of drug addicts: At risk of child abuse, neglect and placement in foster care. Neurotoxicology and Teratology, 9(4), 315-319.
157. Kantor, G. K. and Straus, M. A. (1989). Substance abuse as a precipitant of wife abuse victimizations. American Journal of Drug Abuse, 15(2), 173-189; Regan, D. O., Erlich, S. M. and Finnegan, L. P. (1987). Infants of drug addicts: At risk of child abuse, neglect and placement in foster care. Neurotoxicology and Teratology, 9(4), 315-319; Zweben, J. E. (1991). Counseling issues in methadone maintenance treatment. Journal of Psychoactive Drugs, 23(2), 177-190.
158. Tardiff, K., Marzuk, P. M., Leon, A. C., Hirsch, C. S., Stajic, M., Portera, L. and Hartwell, N. (1994). Homicide in New York City: Cocaine use and firearms. JAMA, 272(1), 43-46.
159. Tardiff, K., Marzuk, P. M., Leon, A. C., Hirsch, C. S., Stajic, M., Portera, L. and Hartwell, N. (1994). Homicide in New York City: Cocaine use and firearms. JAMA, 272(1), 43-46.
160. Tardiff, K., Marzuk, P. M., Leon, A. C., Hirsch, C. S., Stajic, M., Portera, L. and Hartwell, N. (1994). Homicide in New York City: Cocaine use and firearms. JAMA, 272(1), 43-46.
161. Marzuk, P. M., Tardiff, K., Leon, A. C., Hirsch, C. S., Stajic, M., Portera, L., Hartwell, N. and Iqbal, M. I. (1995). Fatal injuries after cocaine use as a leading cause of

death among young adults in New York City. New England Journal of Medicine, 332(26), 1753-1757.

162. Braude, M. C. (1986). Drugs and drug interactions in the elderly woman. In B. A. Ray and M. C. Braude (Eds.). Women and drugs: A new era for research. NIDA research monograph 65 (pp. 58-65). Rockville, MD: National Institute on Drug Abuse.

163. Braude, M. C. (1986). Drugs and drug interactions in the elderly woman. In B. A. Ray and M. C. Braude (Eds.). Women and drugs: A new era for research. NIDA research monograph 65 (pp. 58-65). Rockville, MD: National Institute on Drug Abuse.

164. Galbraith, S. (1991). Women and legal drugs. In P. Roth (Ed.), Alcohol and drugs are women's issues (pp. 15-154). Metuchen, NJ and London: Women's Action Alliance and Scarecrow Press; Braude, M. C. (1986). Drugs and drug interactions in the elderly woman. In B. A. Ray and M. C. Braude (Eds.). Women and drugs: A new era for research. NIDA research monograph 65 (pp. 58-65). Rockville, MD: National Institute on Drug Abuse; Trachtenberg, A. I. and Fleming, M. F. (1994). Diagnosis and treatment of drug abuse in family practice. Kansas City, Missouri: American Academy of Family Physicians.

165. Galbraith, S. (1991). Women and legal drugs. In P. Roth (Ed.), Alcohol and drugs are women's issues (pp. 15-154). Metuchen, NJ and London: Women's Action Alliance and Scarecrow Press; Dupont, R. L. and Saylor, K. E. (1992). Depressant substances in adolescent medicine. Pediatrics in Review, 13(10), 381-386; Swift, C. G. (1981). Psychotropic drugs and the elderly. In G. Tognoni, C. Bellantuono and M. Lader (Eds.), Epidemiological impact of psychotropic drugs: Proceedings of the International Seminar on the Impact of Psychotropic Drugs Held in Milan, Italy, 24-26 June, 1981 (pp. 325-338). New York: Elsevier/North-Holland Biomedical Press.

166. Swift, C. G. (1981). Psychotropic drugs and the elderly. In G. Tognoni, C. Bellantuono and M. Lader (Eds.), Epidemiological impact of psychotropic drugs: Proceedings of the International Seminar on the Impact of Psychotropic Drugs Held in Milan, Italy, 24-26 June, 1981 (pp. 325-338). New York: Elsevier/North-Holland Biomedical Press; Ray, W. A., Griffin, M. R. and Downey, W. (1989). Benzodiazepines of long and short elimination half-life and the risk of hip fracture. JAMA, 262 (23), 3303-3307.

167. Mellinger, G. D., Balter, M. B. and Uhlenhuth, E. H. (1984). Prevalence and correlates of the long-term regular use of anxiolytics. JAMA, 251(3), 375-379; Ray, W., Taylor, J. A., Meador, K. G., Lichtenstein, M. J., Griffin, M. R., Fought, R., Adams, M. L. and Blazer, D. G. (1993). Reducing antipsychotic drug use in nursing homes. Archives of Internal Medicine, 153, 713-721; U.S. Senate Special Committee on Aging, the American Association of Retired Persons, the Federal Council on the Aging and the U.S. Administration on Aging. (1991). Aging America: Trends and projections. Washington, DC: U.S. Department of Health and Human Services; Curtis, J. R., Geller, G., Stokes, E. J., Levine, D. M. and Moore, R. D. (1989). Characteristics, diagnosis and treatment of alcoholism in elderly patients. Journal of the American Geriatrics Society, 37(4), 310-316.

168. Tinetti, M. E., Speechley, M. and Ginter, S. F. (1988). Risk factors for falls among elderly persons living in the community. New England Journal of Medicine, 319(26), 1701-1707; Ray, W. A., Griffin, M. R. and Malcom, E. (1991). Cyclic antidepressants and the risk of hip fracture. Archives of Internal Medicine, 151, 754-756; Lipsitz, L. A., Hirayama, T., Nakajima, I., Kelley, M., Ruthazer, R., Hirayama, T., Levine, D. and Izumo, H. (1991). Muscle strength, medications and falls in elderly Japanese and American nursing home residents: A cross-cultural study. Journal of the American Geriatric Society, 39(8), A10; Ray, W. A., Griffin, M. R. and Downey, W. (1989). Benzodiazepines of long and short elimination half-life and the risk of hip fracture. JAMA, 262 (23), 3303-3307; Ray, W. A., Griffin, M. R., Schaffner, W., Baugh, D. K. and Melton, L. J. (1987). Psychotropic drug use and the risk of hip fractures. New England Journal of Medicine, 316(7), 363-369.
169. Tinetti, M. E., Speechley, M. and Ginter, S. F. (1988). Risk factors for falls among elderly persons living in the community. New England Journal of Medicine, 319(26), 1701-1707.
170. Tinetti, M. E., Speechley, M. and Ginter, S. F. (1988). Risk factors for falls among elderly persons living in the community. New England Journal of Medicine, 319(26), 1701-1707.
171. Ray, W. A., Griffin, M. R. and Malcom, E. (1991). Cyclic antidepressants and the risk of hip fracture. Archives of Internal Medicine, 151, 754-756.
172. Center on Addiction and Substance Abuse at Columbia University. (Unpublished monograph). Substance abuse and health care costs: Its pervasive impact on morbidity and expenditures. New York: CASA.
173. Center on Addiction and Substance Abuse at Columbia University. (1994). Substance abuse and women on welfare. New York: CASA.
174. Sisco, C. B. and Pearson, C.L. (1994). Prevalence of alcoholism and drug abuse among female AFDC recipients. Health & Social Work, 19(1), 75-6.

CHAPTER IV.

REFERENCES

1. Center for the Future of Children. (1995). The future of children: Low birth weight. Los Altos, CA: Center for the Future of Children; DiFranza, J. R. and Lew, R. A. (1995). Effect of maternal cigarette smoking on pregnancy complications and sudden infant death syndrome. Journal of Family Practice, 40(4), 385-394; Nieburg, P. (1985). The fetal tobacco syndrome. Journal of the American Medical Association, 253(20), 2998-2999.
2. Clarren, S. K. (1986). Prepared statement in placing infants at risk: Parental addiction and disease. Hearing before the Select Committee on Children, Youth, and Families, George Miller, Chairman, U.S. House of Representatives, Ninety-Ninth Congress, Second Session, held in Washington, DC on May 21, 1986. Washington, DC: U.S. House of Representatives; Clarren, S. K. (1988). Fetal alcohol syndrome: Diagnosis, treatment and mechanisms of teratogenesis. In R. Bellisario, G. J. Mizejewski (eds.), Transplacental disorders: Perinatal detection, treatment and management (including pediatric AIDS) (pp. 37-55). Albany, NY: Alan R. Liss;; Streissguth, A. P., Aase, J. M., Clarren, S. K., Randels, S. P., LaDue, R. A. and Smith, D. F. (1991). Fetal alcohol syndrome in adolescents and adults. Journal of the American Medical Association, 265(15), 1961-1967; Abel, E. L. and Sokol, R. J. (1987). Incidence of fetal alcohol syndrome and economic impact of FAS-related anomalies. Drug and Alcohol Dependence, 19(1), 5-70.
3. National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute on Drug Abuse.
4. National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute on Drug Abuse; Vega, W. A., Kolody, B., Hwang, J., and Noble, A. (1993). Prevalence and magnitude of perinatal substance exposure in California. New England Journal of Medicine, 329(12), 850-854; Center for the Future of Children (1991). The future of children: Drug exposed infants. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children; South Carolina State Council on Maternal, Infant and Child Health, Office of the Governor of South Carolina. (1991). 1991 South Carolina prevalence study of drug use among women giving birth. South Carolina: State Council on Maternal, Infant and Child Health.
5. National Household Survey on Drug Abuse, 1994; National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute of Drug Abuse.

6. U. S. House of Representatives, Committee on Ways and Means, Subcommittee on Human Resources. (1990). The enemy within: Crack-cocaine and America's families. Washington, DC: U.S. Government Printing Office.
7. Dicker, M. and Leighton, E. A. (1994). Trends in the US prevalence of drug-using parturient women and drug-affected newborns, 1979 through 1990. American Journal of Public Health, 84(9), 1433-1438.
8. Kandall, S. R. and Chavkin, W. (1992). Illicit drugs in America: History, impact on women and infants, and treatment strategies for women. Hastings Law Journal, 43(3), 615-643; Moroney, J. T. and Allen, M. H. (1994). Cocaine and alcohol use in pregnancy. In O. Devinsky, E. Feldmann, B. Hainline (Eds.), Neurological complications of pregnancy (pp. 231-242). New York: Raven Press.
9. Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.
10. Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.
11. Wiese, D., Daro, D. and National Center on Child Abuse Prevention Research. (1995). Current trends in child abuse reporting and fatalities: The results of the 1994 annual fifty state survey. Chicago, IL: National Committee to Prevent Child Abuse; Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General; Miller, J. (1989). Drug initiatives. W-Memo, 1(10), 2-9.
12. Wiese, D., Daro, D. and National Center on Child Abuse Prevention Research. (1995). Current trends in child abuse reporting and fatalities: The results of the 1994 annual fifty state survey. Chicago, IL: National Committee to Prevent Child Abuse.
13. Miller, J. (1989). Drug initiatives. W-Memo, 1(10), 2-9; New York City Human Resources Administration. (1992). Preliminary report on kinship foster care family profile. Memorandum from George Gabel, Director, Office of Management Analysis to Robert L. Little, Executive Deputy Commissioner.
14. U.S. General Accounting Office. (1994). Foster care: Parental drug abuse has alarming impact on young children. Washington, DC: U.S. General Accounting Office.
15. U.S. General Accounting Office. (1994). Foster care: Parental drug abuse has alarming impact on young children. Washington, DC: U.S. General Accounting Office.
16. Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.

17. Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.
18. New York City Child Fatality Review Panel. (1995). Annual report for 1994. [New York]: New York City Child Fatality Review Panel.
19. New York City Child Fatality Review Panel. (1995). Annual report for 1994. [New York]: New York City Child Fatality Review Panel.
20. Davis, S. F., Byers, R. H., Lindegren, M. L., Caldwell, M. B., Karon, J. M. and Gwinn, M. (1995). Prevalence and incidence of vertically-acquired HIV infection in the United States. Journal of the American Medical Association, 274(12), 952-955.
21. Davis, S. F., Byers, R. H., Lindegren, M. L., Caldwell, M. B., Karon, J. M. and Gwinn, M. (1995). Prevalence and incidence of vertically-acquired HIV infection in the United States. Journal of the American Medical Association, 274(12), 952-955.
22. Centers for Disease Control. (1995). Update: AIDS among women--United States, 1994. Morbidity and Mortality Weekly Report, 44(5), 81-84; Brown, L. S., Mitchell, J. L. and DeVore, S. L. (1989). Female intravenous drug users and perinatal HIV transmission. New England Journal of Medicine, 320(22), 1493-1494.
23. Marx, R., Aral, S. O., Rolfs, R. T., Sterk, C. E., Kahn, J. G. (1991). Crack, sex, and STD. Sexually Transmitted Diseases, 18(2), 92-101; Booth, R. E., Koester, S. K. and Pinto, F. (1995). Gender differences in sex-risk behaviors, economic livelihood and self-concept among drug injectors and crack smokers. American Journal of Addictions, 4(4), 313-322; Battjes, R. J., Sloboda, Z. and Grace, W. C. (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA research monograph 143. Rockville, MD: National Institute on Drug Abuse; McCoy, H. V., Inciardi, J. A., Metsch, L. R., Pottieger, A. E. and Saum, C. A. (1995). Women, crack and crime: Gender comparisons of criminal activity among crack cocaine users. Contemporary Drug Problems, 22(3), 435-451; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO.
24. Marx, R., Aral, S. O., Rolfs, R. T., Sterk, C. E., Kahn, J. G. (1991). Crack, sex, and STD. Sexually Transmitted Diseases, 18(2), 92-101; Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO; Centers for Disease Control and Prevention. (1994). Sexually transmitted disease surveillance 1994. Atlanta, GA: Centers for Disease Control and Prevention.
25. Centers for Disease Control and Prevention, National Center for HIV, STD and TB Prevention, Division of STD Prevention. (1995). Sexually transmitted disease surveillance 1994. Atlanta, GA: Centers for Disease Control and Prevention.

26. Farley, T. A., Hadler, J. L. and Gunn, R. A. (1990). The syphilis epidemic in Connecticut: Relationship to drug use and prostitution. Sexually Transmitted Diseases, 17(4), 163-168; Marcenko, M. O., Spence, M. and Rohweder, C. (1994). Psychosocial characteristics of pregnant women with and without a history of substance abuse. Health and Social Work, 19(1), 17-22; Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing; U.S. General Accounting Office. (1991). Drug abuse: The crack cocaine epidemic: Health consequences and treatment. Washington, DC: GAO; Smart, R. G. (1991). Crack cocaine use: A review of prevalence and adverse affects. American Journal of Drug and Alcohol Abuse, 17(1), pp. 13-26; Lewin, Tamar. Syphilis cases among newborns climb with the rise in crack use. New York Times, 11/23/92, p. A13.
27. Ostrea, E. M. and Welch, R. A. (1991). Detection of prenatal drug exposure in the pregnant woman and her newborn infant. Clinics in Perinatology, 18(3), 629-645.
28. Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321; South Carolina State Council on Maternal, Infant and Child Health, Office of the Governor of South Carolina. (1991). 1991 South Carolina prevalence study of drug use among women giving birth. South Carolina: State Council on Maternal, Infant and Child Health; Current meconium tests often fail to detect cocaine exposure. (1995). Forensic Drug Abuse Advisor, 7(1), 4.
29. Forman, R., Schneiderman, J., Klein, J., Graham, K., Greenwald, M. and Koren, G. (1992). Accumulation of cocaine in maternal and fetal hair: The dose response curve. Life Sciences, 50(18), 1333-1341.
30. Vega, W. A., Kolody, B., Hwang, J., Porter, P., Bole, A. and Dimas, J. (1993). Profile of alcohol and drug use during pregnancy in California, 1992: Perinatal substance exposure study. Scientific Report to the State Department of Alcohol and Drug Programs, Sacramento, CA: Office of Perinatal Substance Abuse.
31. National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute on Drug Abuse.
32. Gfroerer, J. C. and Office of Applied Studies. (1995). Preliminary estimates from the 1994 National Household Survey on Drug Abuse. [Rockville, MD]: Office of Applied Studies.
33. Chasnoff, I. J. (1989). Drug use and women: Establishing a standard of care. Annals of the New York Academy of Sciences, 562, 208-210.
34. Center for the Future of Children (1991). The future of children: Drug exposed infants. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children.

35. Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321.
36. Chasnoff, I. J. (1992). Drug use and women: Establishing a standard of care. Annals of the New York Academy of Sciences, 562, 208-210; Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.
37. National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute on Drug Abuse.
38. South Carolina State Council on Maternal, Infant and Child Health, Office of the Governor of South Carolina. (1991). 1991 South Carolina prevalence study of drug use among women giving birth. South Carolina: State Council on Maternal, Infant and Child Health; Cartwright, P. S., Schorge, J. O. and McLaughlin, F. J. (1991). Epidemiologic characteristics of drug use during pregnancy: Experience in a Nashville hospital. Southern Medical Journal, 84(7), 867-870; Chasnoff, I. J., Landress, H. J. and Barrett, M. E. (1990). The prevalence of illicit-drug or alcohol use during pregnancy and discrepancies in mandatory reporting in Pinellas County, Florida. New England Journal of Medicine, 322(17), 1202-1206; Lake, M. F., Angel, J. L., Murphy, J. M. and Poekert, G. (1992). Patterns of illicit drug use at the time of labor in a private and public hospital. Journal of Perinatology, 12(2), 134-136; Robins, L. N., Mills, J. L., Krulewitch, C. and Herman, A. A. (1993). Effects of in utero exposure to street drugs. American Journal of Public Health, 83(Suppl.), 9-32.
39. Vega, W. A., Kolody, B., Hwang, J., Porter, P., Bole, A. and Dimas, J. (1993). Profile of alcohol and drug use during pregnancy in California, 1992: Perinatal substance exposure study. Scientific Report to the State of California Department of Alcohol and Drug Programs, Sacramento, CA: Office of Perinatal Substance Abuse; National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute on Drug Abuse.
40. Abma, J. C. and Mott, F. L. (1991). Substance use and prenatal care during pregnancy among young women. Family Planning Perspectives, 23(3), 117-122; National Household Survey on Drug Abuse, 1994; National Institute on Drug Abuse. (1994). National health and pregnancy survey. Rockville, MD: National Institute of Drug Abuse.
41. Abma, J. C. and Mott, F. L. (1991). Substance use and prenatal care during pregnancy among young women. Family Planning Perspectives, 23(3), 117-122.
42. Abma, J. C. and Mott, F. L. (1991). Substance use and prenatal care during pregnancy among young women. Family Planning Perspectives, 23(3), 117-122; South Carolina State Council on Maternal, Infant and Child Health, Office of the Governor of South Carolina. (1991). 1991 South Carolina prevalence study of drug use among women giving birth. South Carolina: State Council on Maternal, Infant and Child Health; Centers for Disease Control

and Prevention. (1995). Sociodemographic and behavioral characteristics associated with alcohol consumption during pregnancy: United States, 1988. Morbidity and Mortality Weekly Report, 44(13), 261-264; But a study in California found pregnant women age 24 and under more likely to use alcohol. See Vega, W. A., Kolody, B., Hwang, J., Porter, P., Bole, A. and Dimas, J. (1993). Profile of alcohol and drug use during pregnancy in California, 1992: Perinatal substance exposure study. Scientific Report to the State of California Department of Alcohol and Drug Programs, Sacramento, CA: Office of Perinatal Substance Abuse.

43. U.S. Department of Health and Human Services, Public Health Service (1981). Surgeon General's advisory on alcohol and pregnancy. FDA Bulletin, 11 (2), 9-10; Centers for Disease Control and Prevention. (1995). Sociodemographic and behavioral characteristics associated with alcohol consumption during pregnancy: United States, 1988. Morbidity and Mortality Weekly Report, 44(13), 261-264.

44. Centers for Disease Control and Prevention. (1995). Sociodemographic and behavioral characteristics associated with alcohol consumption during pregnancy: United States, 1988. Morbidity and Mortality Weekly Report, 44(13), 261-264; Streissguth, A. P., Bookstein, F. L., Sampson, P. D. and Barr, H. M. (1993). The enduring effects of prenatal alcohol exposure on child development: Birth outcomes through seven years, a partial least squares solution. Ann Arbor, MI: University of Michigan Press; Streissguth, A. P., Sampson, P. D., Olson, H. C., Bookstein, F. L., Barr, H. M., Scott, M., Feldman, J. and Mirsky, A. F. (1994). Maternal drinking during pregnancy: Attention and short-term memory in 14-year-old offspring: A longitudinal study. Alcoholism: Clinical and Experimental Research, 18(1), 202-218; Stratton, K., Howe, C., Battaglia, F. and Institute of Medicine, Division of Biobehavioral Sciences and Mental Disorders, Committee to Study Fetal Alcohol Syndrome. (1995). Fetal Alcohol Syndrome: Diagnosis, epidemiology, prevention and treatment. Washington, DC: National Academy Press.

45. South Carolina State Council on Maternal, Infant and Child Health, Office of the Governor of South Carolina. (1991). 1991 South Carolina prevalence study of drug use among women giving birth. South Carolina: State Council on Maternal, Infant and Child Health; Vega, W. A., Kolody, B., Hwang, J., Porter, P., Bole, A. and Dimas, J. (1993). Profile of alcohol and drug use during pregnancy in California, 1992: Perinatal substance exposure study. Scientific Report to the State of California Department of Alcohol and Drug Programs, Sacramento, CA: Office of Perinatal Substance Abuse; Lake, M. F., Angel, J. L., Murphy, J. M. and Poekert, G. (1992). Patterns of illicit drug use at the time of labor in a private and public hospital. Journal of Perinatology, 12(2), 134-136.

46. Chasnoff, I. J., Landress, H. J. and Barrett, M. E. (1990). Prevalence of illicit drug or alcohol use during pregnancy and discrepancies in mandatory reporting in Pinellas County, Florida. New England Journal of Medicine, 322(17), 1202-1206.

47. Lutiger, B., Graham, K., Einarson, T. R. and Koren, G. (1991). Relationship between gestational cocaine use and pregnancy outcome: A meta-analysis. Teratology, 44(4), 405-414; Mayes, L. C., Granger, R. H., Bornstein, M. H. and Zuckerman, B. (1992). The

problem of prenatal cocaine exposure: A rush to judgment. Journal of the American Medical Association, 267(3), 406-408.

48. Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321; Streissguth, A. P., Grant, T. M., Barr, H. M., Brown, Z. A., Martin, J. C., Mayock, D. E., Ramey, S. L. and Moore, L. (1991). Cocaine and the use of alcohol and other drugs during pregnancy. American Journal of Obstetrics and Gynecology, 164(5, Pt. 1), 1239-1243; Feig, L. (1990). Drug exposed infants and children: Service needs and policy questions. Washington, DC: U.S. Department of Health and Human Services; Robins, L. N., Mills, J. L., Krulewitch, C. and Herman, A. A. (1993). Effects of in utero exposure to street drugs. American Journal of Public Health, 83(Suppl.), 9-32; Coles, C. D. (1992). Discussion: Measurement issues in the study of effects of substance abuse in pregnancy. In M. M. Kilbey, K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 248-258). Rockville, MD: National Institute on Drug Abuse; Marcenko, M. O., Spence, M. and Rohweder, C. (1994). Psychosocial characteristics of pregnant women with and without a history of substance abuse. Health and Social Work, 19(1), 17-22.

49. Center for the Future of Children (1995). The future of children: Low birth weight. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children; DiFranza, J. R. and Lew, R. A. (1995). Effect of maternal cigarette smoking on pregnancy complications and sudden infant death syndrome. Journal of Family Practice, 40(4), 385-394; Drews, C. D., Murphy, C. C., Yeargin-Allsopp, M. and Decoufle, P. (1996). The relationship between idiopathic mental retardation and maternal smoking during pregnancy. Pediatrics 97(4), 547-53; Husten, C. G., Chrismon, J. H. and Reddy, M. N. (1996). Trends and effects of cigarette smoking among girls and women in the United States, 1965-1993. Journal of the American Medical Women's Association, 51(1), 1-8; Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: Office on Smoking and Health; Wang, X., Wypi, D., Gold, D. R., Speizer, F. E., Ware, J. H., Ferris, B. G. and Dockery, D. W. (1994). A longitudinal study of the effects of parental smoking on pulmonary function in children 6-18 years. American Journal of Respiratory and Critical Care Medicine, 149(6), 1420-1425; Gold, D. R., Rotnitzky, A., Damokosh, A. I., Ware, J. H., Speizer, F. E., Ferris, B. G. and Dockery, D. W. (1993). Race and gender differences in respiratory illness prevalence and their relationship to environmental exposures in children 7 to 14 years of age. American Review of Respiratory Diseases, 148(1), 10-18; Floyd, R. L., Rimer, B. K., Giovino, G. A., Mullen, P. D. and Sullivan, S. E. (1993). A review of smoking in pregnancy: Effects on pregnancy outcomes and cessation efforts. Annual Review of Public Health, 14, 379-411; Taylor, B. and Wadsworth, J. (1987). Maternal smoking during pregnancy and lower respiratory tract illness early in life. Archives of Disease in Childhood, 62(8), 786-791; Olds, D. L., Hendersen, C. R. and Tatelbaum, R. (1994). Intellectual impairment in children of women who smoke cigarettes during pregnancy. Pediatrics, 93(2), 221-227.

50. Centers for Disease Control and Prevention, National Center for Health Statistics, Linked Birth/Infant Death Data Set.
51. Nieburg, P., Marks, J. S., McLaren, N. M. and Remington, P. L. (1985). The fetal tobacco syndrome. JAMA, 253(20), 2998-2999.
52. Center for the Future of Children (1995). The future of children: Low birth weight. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children; DiFranza, J. R. and Lew, R. A. (1995). Effect of maternal cigarette smoking on pregnancy complications and sudden infant death syndrome. Journal of Family Practice, 40(4), 385-394.
53. Martin, T. R. and Bracken, M. B. (1986). Association of low birth weight with passive smoke exposure in pregnancy. American Journal of Epidemiology, 124(4), 633-641; Eskenazi, B., Prehn, A. W. and Christianson, R. E. (1995). Passive and active maternal smoking as measured by serum cotinine: The effect on birthweight. American Journal of Public Health, 85(3), 395-398; Haddow, J. E., Knight, G. J., Palomaki, G. E. and McCarthy, J. E. (1988). Second-trimester serum cotinine levels in nonsmokers in relation to birth weight. American Journal of Obstetrics and Gynecology, 159(2), 481-484.
54. English, P. B., Eskenazi, B. and Christianson, R. E. (1994). Black-white differences in serum cotinine levels among pregnant women and subsequent effects on infant birthweight. American Journal of Public Health, 84(9), 1439-1443; Knight, J. M., Eliopoulos, C., Klein, J., Greenwald, M. and Koren, G. (1996). Passive smoking in children: Racial differences in systematic exposure to cotinine by hair and urine analysis. Chest, 109(2), 446-450.
55. Ramirez, A. G. and Gallion, K. J. (1993). Nicotine dependence among blacks and Hispanics. In C. T. Orleans, J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 350-364). New York: Oxford University Press; Centers for Disease Control and Prevention. (1987). Cigarette smoking among blacks and other minority populations. Morbidity and Mortality Weekly Report, 36(25), 404-407; Barnett, E. (1995). Race differences in the proportion of low birth weight attributable to maternal cigarette smoking in a low-income population. American Journal of Health Promotion, 10(2), 105-110; Gardner, P. and Hudson, B. L. (1996). Advance report of final mortality statistics, 1993. Monthly Vital Statistics Report, 44(7), 1-84.
56. Klonoff-Cohen, H. S., Edelstein, S. L., Lefkowitz, E. S., Srinivasen, I. P., Kaegi, D., Chang, J. C. and Wiley, K. J. (1995). Effect of passive smoking and tobacco exposure through breast milk on sudden infant death syndrome. JAMA, 273(10), 795-798.
57. Klonoff-Cohen, H. S., Edelstein, S. L., Lefkowitz, E. S., Srinivasen, I. P., Kaegi, D., Chang, J. C. and Wiley, K. J. (1995). Effect of passive smoking and tobacco exposure through breast milk on sudden infant death syndrome. JAMA, 273(10), 795-798.
58. Mainous, A. G. and Hueston, W. J. (1994). Passive smoke and low birth weight. Archives of Family Medicine, 3(10), 875-878; Reed, B. D., and Lutz, L. J. (1988).

- Household smoking exposure: Association with middle ear effusions. Family Medicine, 20 (6), 426-430; Counsel of Scientific Affairs, American Medical Association. (1994).
- Environmental tobacco smoke: Health effects and prevention policies. Archives of Family Practice, 3(10), 865-871; Environmental Protection Agency. (1993). Respiratory health effects of passive smoking: Lung cancer and other disorders. Bethesda, MD: National Institutes of Health; Strachan, D. P., Jarvis, M. J. and Feyerabend, C. (1989). Passive smoking, salivary cotinine concentrations and middle ear effusion in 7 year old children. British Medical Journal, 298(6687), 1549-1552; Gold, D. R., Rotnitzky, A., Damokosh, A. I., Ware, J. H., Speizer, F. E., Ferris, B. G. and Dockery, D. W. (1993). Race and gender differences in respiratory illness prevalence and their relationship to environmental exposures in children 7 to 14 years of age. American Review of Respiratory Diseases, 148(1), 10-18.
59. Fried, P. A. (1993). Prenatal exposure to tobacco and marijuana: Effects during pregnancy, infancy, and early childhood. Clinical Obstetrics and Gynecology, 36(2), 319-337.
60. Olds, D. L., Hendersen, C. R. and Tatelbaum, R. (1994). Intellectual impairment in children of women who smoke cigarettes during pregnancy. Pediatrics, 93(2), 221-227.
61. Fried, P. A. and Watkinson, B. (1988). 12- and 24-month neurobehavioral follow-up of children prenatally exposed to marihuana, cigarettes and alcohol. Neurotoxicology and Teratology, 10(4), 305-313.
62. Eskenazi, B. and Trupin, L. S. (1995). Passive and active maternal smoking during pregnancy, as measured by serum cotinine, and postnatal smoke exposure. II. Effects on neurodevelopment at age 5 years. American Journal of Epidemiology, 142(9, Suppl.), S19-S29.
63. Eskenazi, B. and Trupin, L. S. (1995). Passive and active maternal smoking during pregnancy, as measured by serum cotinine, and postnatal smoke exposure. II. Effects on neurodevelopment at age 5 years. American Journal of Epidemiology, 142(9, Suppl.), S19-S29.
64. Weitzman, M., Gortmaker, S. and Sobol, A. (1992). Maternal smoking and behavior problems of children. Pediatrics, 90(3), 342-349.
65. Kandel, D. B., Wu, P. and Davies, M. (1994). Maternal smoking during pregnancy and smoking by adolescent daughters. American Journal of Public Health, 84(9), 1407-1413.
66. Kandel, D. B., Wu, P. and Davies, M. (1994). Maternal smoking during pregnancy and smoking by adolescent daughters. American Journal of Public Health, 84(9), 1407-1413.
67. Center for the Future of Children. (1995). The future of children: Low birth weight. Los Altos, CA: Center for the Future of Children.

68. Li, C. Q., Windsor, R. A., Perkins, L., Goldenberg, R. L. and Lowe, J. B. (1993). The impact on infant birth weight and gestational age of cotinine-validated smoking reduction during pregnancy. JAMA, 269(12), 1519-1524.
69. Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: Office on Smoking and Health.
70. National Institute on Alcohol Abuse and Alcoholism. (1993). Eighth special report to the U.S. Congress on alcohol and health. Alexandria, VA: National Institute on Alcohol Abuse and Alcoholism; Mello, N. K. (1988). Effects of alcohol abuse on reproductive function in women. Recent Developments in Alcoholism, 6, 253-276.
71. Harlap, S. and Shiono, P. H. (1980). Alcohol, smoking and incidence of spontaneous abortions in the first and second trimesters. Lancet, 2(8187), 173-6.
72. Centers for Disease Control and Prevention, National Center for Health Statistics, Linked Birth/Infant Death Data Set.
73. Clarren, S. K. (1988). Fetal alcohol syndrome: Diagnosis, treatment, and mechanisms of teratogenesis. In R. Bellisario, G. J. Mizejewski (Eds.), Transplacental disorders: Perinatal detection, treatment and management (including pediatric AIDS) (pp. 37-55). Albany, NY: Alan R. Liss; Aase, J. M. (1994). Clinical recognition of FAS: Difficulties of detection and diagnosis. Alcohol Health and Research World. Special Focus: Alcohol-Related Birth Defects, 18(1), 5-9; Streissguth, A. P. and LaDue, R. A. (1985). Psychological and behavioral effects in children prenatally exposed to alcohol. Alcohol Health and Research World, 10(1), 6-12; Stratton, K., Howe, C., Battaglia, F. and Institute of Medicine. Fetal alcohol syndrome: Diagnosis, epidemiology, prevention, and treatment. Washington, DC: National Academy Press.
74. Centers for Disease Control and Prevention. (1995). Update: Trends in fetal alcohol syndrome--United States, 1979-1993. Morbidity and Mortality Weekly Report, 44(13), 249-251.
75. Little, B. B., Snell, L. M., Rosenfeld, C. R., Gilstrap, L. C. and Gant, N. F. (1990). Failure to recognize fetal alcohol syndrome in newborn infants. American Journal of Diseases of Children, 144(10), 1142-1146; Clarren, S. K. (1988). Fetal alcohol syndrome: Diagnosis, treatment, and mechanisms of teratogenesis. In R. Bellisario, G. J. Mizejewski (Eds.), Transplacental disorders: Perinatal detection, treatment, and management (including pediatric AIDS) (pp. 37-55). Albany, NY: Alan R. Liss; Streissguth, A. P. (1994). Fetal alcohol syndrome: Understanding the problem: Understanding the solution: What Indian communities can do. American Indian Culture and Research Journal, 18(3), 45-83; Anderson, B., Novick, E. and U.S. Department of Health and Human Services, Division of Children and Youth Policy. (1992). Fetal alcohol syndrome and pregnant women who abuse alcohol: An overview of the issue and the federal response. [Washington, DC]: U.S. Department of Health and Human Services, Office of the Assistant Secretary for Planning

and Evaluation, Office of Human Services Policy, Division of Children and Youth Policy; Stratton, K., Howe, C., Battaglia, F. and Institute of Medicine. (1995) Fetal alcohol syndrome: Diagnosis, epidemiology, prevention, and treatment. Washington, DC: National Academy Press; Clarren, S. K. (1990). Maternal alcohol ingestion: Effects on the developing fetus. Current Pediatric Therapy, 13, 706-707.

76. Stratton, K., Howe, C., Battaglia, F. and Institute of Medicine. (1995). Fetal alcohol syndrome: Diagnosis, epidemiology, prevention, and treatment. Washington, DC: National Academy Press.

77. Caruso, K. and ten Benschel, R. (1993). Fetal alcohol syndrome and fetal alcohol effects. Minnesota Medicine, 76(4), 25-29; Stratton, K., Howe, C., Battaglia, F. and Institute of Medicine. (1995). Fetal alcohol syndrome: Diagnosis, epidemiology, prevention, and treatment. Washington, DC: National Academy Press.

78. Caruso, K. and ten Benschel, R. (1993). Fetal alcohol syndrome and fetal alcohol effects. Minnesota Medicine, 76(4), 25-29; Streissguth, A. P. and LaDue, R. A. (1985). Psychological and behavioral effects in children prenatally exposed to alcohol. Alcohol Health and Research World, 10(1), 6-12; Little, R. E. and Wendt, J. K. (1993). The effects of maternal drinking during the reproductive period. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 191-213). Norwood, NJ: Ablex Publishing.

79. Streissguth, A. P. (1994). A long-term perspective of FAS. Alcohol Health and Research World, 18(1), 74-81; Streissguth, A. P. (1992). Fetal alcohol syndrome and fetal alcohol effects: A clinical perspective of later developmental consequences. In I. S. Zagon, T. A. Slotkin (Eds.), Maternal substance abuse and the developing nervous system (pp. 5-25). San Diego: Academic Press; Streissguth, A. P., Aase, J. M., Clarren, S. K., Randels, S. P., LaDue, R. A. and Smith, D. F. (1991). Fetal alcohol syndrome in adolescents and adults. Journal of the American Medical Association, 265(15), 1961-1967; Stratton, K., Howe, C., Battaglia, F. and Institute of Medicine. Fetal alcohol syndrome: Diagnosis, epidemiology, prevention, and treatment. Washington, DC: National Academy Press.

80. Streissguth, A. P. (1992). Fetal alcohol syndrome and fetal alcohol effects: A clinical perspective of later developmental consequences. In I. S. Zagon, T. A. Slotkin (Eds.), Maternal substance abuse and the developing nervous system (pp. 5-25). San Diego, CA: Academic Press; Sampson, P. D., Bookstein, F. L., Barr, H. M. and Streissguth, A. P. (1994). Prenatal alcohol exposure, birthweight, and measures of child size from birth to age 14 years. American Journal of Public Health, 84(9), 1421-1428.

81. Streissguth, A. P., Sampson, P. D., Olson, H. C., Bookstein, F. L., Barr, H. M., Scott, M., Feldman, J. and Mirsky, A. F. (1994). Maternal drinking during pregnancy: Attention and short-term memory in 14-year-old offspring: A longitudinal study. Alcoholism: Clinical and Experimental Research, 18(1), 202-218; Streissguth, A. P., Bookstein, F. L., Sampson, P. D. and Barr, H. M. (1993). The enduring effects of prenatal alcohol exposure

on child development: Birth outcomes through seven years. a partial least squares solution. Ann Arbor, MI: University of Michigan Press.

82. Jacobson, J. L. and Jacobson, S. W. (1994). Prenatal alcohol exposure and neurobehavioral development: Where is the threshold? Alcohol Health and Research World 18(1), 30-6; Harlap, S. and Shiono, P. H. (1980). Alcohol, smoking and incidence of spontaneous abortions in the first and second trimesters. Lancet, 2(8187), 173-6; Stratton, K., Howe, C., Battaglia, F. and Institute of Medicine, Division of Biobehavioral Sciences and Mental Disorders, Committee to Study Fetal Alcohol Syndrome. (1995). Fetal Alcohol Syndrome: Diagnosis, epidemiology, prevention and treatment. Washington, DC: National Academy Press.
83. Centers for Disease Control (1995). Sociodemographic and behavioral characteristics associated with alcohol consumption during pregnancy: United States, 1988. Morbidity and Mortality Weekly Report, 44(13), 261-265; Jacobson, J. L. and Jacobson, S. W. (1994). Prenatal alcohol exposure and neurobehavioral development: Where is the threshold? Alcohol Health and Research World 18(1), 30-6; Little, R. E. and Wendt, J. K. (1993). The effects of maternal drinking in the reproductive period: An epidemiological review. In E. S. L. Gomberg, T. D. Nirenberg (Eds.), Women and substance abuse (pp. 191-213). Norwood, NJ: Ablex Publishing; Blume, S. B. (1986). Is social drinking during pregnancy harmless? There is reason not to think so. In B. Stimmel (Ed.), Controversies in alcoholism and substance abuse (pp. 209-219). New York: Hayworth Press; Smith, K. J. and Eckardt, M. J. (1991). The effects of prenatal alcohol on the central nervous system. Recent Developments in Alcoholism, 9, 151-164; Friedman, S. L. and Haywood, H. C. (Eds.). (1994). Developmental follow-up: Concepts, domains, and methods. San Diego: Academic Press; Day, N. L., Richardson, G. A., Geva, D. and Robles, N. (1994). Alcohol, marijuana and tobacco: Effects of prenatal exposure on offspring growth and morphology at age six. Alcoholism: Clinical and Experimental Research, 18(4), 786-794.
84. Alpert, J. J. and Zuckerman, B. (1991). Alcohol use during pregnancy: What is the risk? Pediatrics in Review, 12(12), 375-379; Hatfield, D. (1986). Is social drinking during pregnancy harmless? In B. Stimmel (Ed.), Controversies in alcoholism and substance abuse (pp. 221-226). New York: Haworth Press.
85. U.S. Department of Health and Human Services, Public Health Service (1981). Surgeon General's advisory on alcohol and pregnancy. FDA Bulletin, 11 (2), 9-10; The American College of Obstetricians and Gynecologists. (1994). ACOG patient education pamphlet: Drugs and pregnancy: Alcohol, tobacco, and other drugs; U.S. Department of Agriculture, Agricultural Research Service, Advisory Guidelines Committee. (1995). Report of the Dietary Guidelines Advisory Committee on the dietary guidelines for Americans, 1995, to the Secretary of Health and Human Services and the Secretary of Agriculture. [Washington, DC]: U.S. Department of Agriculture, Agricultural Research Service, Dietary Guidelines Advisory Committee; Grinspoon, L. and Bakalar, J. B. (1990). Harvard Medical School Mental Health Review: Alcohol abuse and dependence. Cambridge, MA: President and Fellows of Harvard College; Clarren, S. K. (1990). Maternal alcohol ingestion: Effects on

the developing fetus. Current Pediatric Therapy, 13, 706-707; Centers for Disease Control (1995). Sociodemographic and behavioral characteristics associated with alcohol consumption during pregnancy: United States, 1988. Morbidity and Mortality Weekly Report, 44(13), 261-265; Jacobson, J. L. and Jacobson, S. W. (1994). Prenatal alcohol exposure and neurobehavioral development: Where is the threshold? Alcohol Health and Research World 18(1), 30-6.

86. Little, R. E. and Wendt, J. K. (1993). The effects of maternal drinking in the reproductive period: An epidemiological review. In E. S. L. Gomberg, T. D. Nirenberg (Eds.), Women and substance abuse (pp. 191-213). Norwood, NJ: Ablex Publishing; Streissguth, A. P., Sampson, P. D., Olson, H. C., Bookstein, F. L., Barr, H. M., Scott, M., Feldman, J. and Mirsky, A. F. (1994). Maternal drinking during pregnancy: Attention and short-term memory in 14-year-old offspring: A longitudinal study. Alcoholism: Clinical and Experimental Research, 18(1), 202-218; Weiner, L. and Larsson, G. (1987). Clinical prevention of fetal alcohol effects--a reality: Evidence for the effectiveness of intervention. Alcohol Health and Research World, 11(4), 60-63, 92-94.

87. Springhouse Corp. (1994). Illustrated Manual of Nursing Practice. (Second ed.). Springhouse, PA: Springhouse Corp; Streissguth, A. P., Bookstein, F. L., Sampson, P. D. and Barr, H. M. (1993). The enduring effects of prenatal alcohol exposure on child development: Birth outcomes through seven years, a partial least squares solution. Ann Arbor, MI: University of Michigan Press; Larroque, B., Kaminski, M., Dehaene, P., Subtil, D., Delfosse, M. and Querleu, D. (1995). Moderate prenatal alcohol exposure and psychomotor development at preschool age. American Journal of Public Health, 85(12), 1654-1661; Jacobson, S. W. and Sokol, R. J. (1994). Effects of prenatal exposure to alcohol, smoking and illicit drugs on postpartum somatic growth. Alcoholism: Clinical and Experimental Research, 18(2), 317-323; Streissguth, A. P. (1994). Fetal alcohol syndrome: Understanding the problem: Understanding the solution: What Indian communities can do. American Indian Culture and Research Journal, 18(3), 45-83.

88. Weiner, L. and Larsson, G. (1987). Clinical prevention of fetal alcohol effects--a reality: Evidence for the effectiveness of intervention. Alcohol Health and Research World, 11(4), 60-63, 92-94; Day, N. L., Richardson, G. A., Geva, D. and Robles, N. (1994). Alcohol, marijuana and tobacco: Effects of prenatal exposure on offspring growth and morphology at age six. Alcoholism: Clinical and Experimental Research, 18(4), 786-794; Jacobson, J. L., Jacobson, S. W. and Sokol, R. J. (1994). Effects of prenatal exposure to alcohol, smoking and illicit drugs on postpartum somatic growth. Alcoholism: Clinical and Experimental Research, 18(2), 317-323; Streissguth, A. P. (1994). Fetal alcohol syndrome: Understanding the problem: Understanding the solution: What Indian communities can do. American Indian Culture and Research Journal, 18(3), 45-83.

89. Day, N. L. and Richardson, G. A. (1991). Prenatal alcohol exposure: A continuum of effects. Seminars in Perinatology, 15(4), 271-279; Abel, E.L. and Sokol, R.J. (1987). Incidence of fetal alcohol syndrome and economic impact of FAS-related anomalies. Alcohol and Drug Dependence 19, 51-70.

90. Bingol, N., Schuster, C., Fuchs, M., Iosub, S., Turner, G., Stone, R. K. and Gromisch, D. S. (1987). The influence of socioeconomic factors on the occurrence of fetal alcohol syndrome. Advances in Alcoholism and Substance Abuse, 6(4), 105-118.
91. Centers for Disease Control and Prevention, National Center for Health Statistics, Linked Birth/Infant Death Data Set.
92. Bingol, N., Schuster, C., Fuchs, M., Iosub, S., Turner, G., Stone, R. K. and Gromisch, D. S. (1987). The influence of socioeconomic factors on the occurrence of fetal alcohol syndrome. Advances in Alcoholism and Substance Abuse, 6(4), 105-118.
93. Clarren, S. K. (1986). Prepared statement in placing infants at risk: Parental addiction and disease. Hearing before the Select Committee on Children, Youth, and Families, George Miller, Chairman, U.S. House of Representatives, Ninety-Ninth Congress, Second Session, held in Washington, DC on May 21, 1986. Washington, DC: U.S. House of Representatives; Smith, K. J. and Eckardt, M. J. (1991). The effects of prenatal alcohol on the central nervous system. Recent Developments in Alcoholism, 9, 151-164; Streissguth, A. P. (1994). Fetal alcohol syndrome: Understanding the problem: Understanding the solution: What Indian communities can do. American Indian Culture and Research Journal, 18(3), 45-83.; Streissguth, A. P., and Dehaene, P. (1993). Fetal alcohol syndrome in twins of alcoholic mothers: Concordance of diagnosis and IQ. American Journal of Medical Genetics, 47(6), 857-861.
94. Hutchings, D. E. (1993). The puzzle of cocaine's effects following maternal use during pregnancy: Are there reconcilable differences? Neurotoxicology and Teratology, 15(5), 281-286.
95. Griffith, D. R. (1992). Prenatal exposure to cocaine and other drugs: Developmental and educational prognoses. Phi Delta Kappan, 74, 30-34; Griffith, D. R., Azuma, S. D. and Chasnoff, I. J. (1994). Three-year outcome of children exposed prenatally to drugs. Journal of the American Academy of Child Adolescent Psychiatry, 33(1), 20-27; Mayes, L. C., Granger, R. H., Bornstein, M. H. and Zuckerman, B. (1992). The problem of prenatal cocaine exposure: A rush to judgment. Journal of the American Medical Association, 267(3), 406-408; Robins, L. N., Mills, J. L., Krulewitch, C. and Herman, A. A. (1993). Effects of in utero exposure to street drugs. American Journal of Public Health, 83 (Suppl.), 9-32; Lutiger, B., Graham, K., Einarson, T. R. and Koren, G. (1991). Relationship between gestational cocaine use and pregnancy outcome: A meta-analysis. Teratology, 44(4), 405-414; Neuspil, D. R. (1993). Cocaine and the fetus: Mythology of severe risk. Neurotoxicology and Teratology, 15(5), 305-306; Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321.
96. Bateman, D. A., Ng, S. K. C., Hansen, C. A. and Heagarty, M. C. (1993). The effects of intrauterine cocaine exposure in newborns. American Journal of Public Health, 83(2), 190-193; Glantz, J. C. and Woods, J. R. (1993). Cocaine, heroin, and phencyclidine: Obstetric perspectives. Clinical Obstetrics and Gynecology, 36(2), 279-301; Chouteau, M.,

Namerow, P. B. and Leppert, P. (1988). The effect of cocaine abuse on birth weight and gestational age. Obstetrics and Gynecology, 72(3), 351-354; MacGregor, S. N., Keith, L. G., Chasnoff, I. J., Rosener, M. A., Chisum, G. M., Shaw, P. and Minogue, J. P. (1987). Cocaine use during pregnancy: Adverse perinatal outcome. American Journal of Obstetrics and Gynecology, 157(3), 686-690; Cohen, H. R., Green, J. R. and Crombleholme, W. R. (1991). Peripartum cocaine use: Estimating risk of adverse pregnancy outcome. International Journal of Gynecology and Obstetrics, 35(1), 51-54; Zuckerman, B., Frank, D. A., Hingson, R., Amaro, H., Levenson, S. M., Kayne, H., Parker, S., Vinci, R., Aboagye, K., Fried, L. E., Cabral, H., Timperi, R. and Bauchner, H. (1989). Effects of maternal marijuana and cocaine use on fetal growth. New England Journal of Medicine, 320(12), 762-768; Coles, C. D., Platzman, I. S., James, M. E. and Falek, A. (1992). Effects of cocaine and alcohol use in pregnancy on neonatal growth and neurobehavioral status. Neurotoxicology and Teratology, 14, 23-33; Cherukuri, R., Minkoff, H., Feldman, J., Parekh, A. and Glass, L. (1988). A cohort study of alkaloidal cocaine ("crack") in pregnancy. Obstetrics and Gynecology, 72(2), 147-151; Chasnoff, I. J. (1991). Cocaine and pregnancy: Clinical and methodologic issues. Clinics in Perinatology, 18(1), 113-123; Frank, D. A., Bauchner, H., Parker, S., Huber, A. M., Kyei-Aboagye, K., Cabral, H. and Zuckerman, B. (1990). Neonatal body proportionality and body composition after in utero exposure to cocaine and marijuana. Journal of Pediatrics, 117(4), 622-626.

97. Bateman, D. A., Ng, S. K. C., Hansen, C. A. and Heagarty, M. C. (1993). The effects of intrauterine cocaine exposure in newborns. American Journal of Public Health, 83(2), 190-193; Lutiger, B., Graham, K., Einarson, T. R. and Koren, G. (1991). Relationship between gestational cocaine use and pregnancy outcome: A meta-analysis. Teratology, 44(4), 405-414.

98. Besharov, D. J. (1989). The children of crack: Will we protect them? Public Welfare, 47(4), 6-11; Dixon, S. D. (1989). Effects of transplacental exposure to cocaine and methamphetamine on the neonate. Western Journal of Medicine, 150(4), 436-442.

99. Springhouse Corp. (1994). Illustrated Manual of Nursing Practice. (Second ed.). Springhouse, PA: Springhouse Corp; Dombrowski, M. P., Wolfe, H. M., Welch, R. A. and Evans, M. I. (1991). Cocaine abuse is associated with abruptio placentae and decreased birth weight, but not shorter labor. Obstetrics and Gynecology, 77(1), 139-141; Bingol, N., Fuchs, M., Diaz, V., Stone, R. K. and Gromisch, D. S. (1987). Teratogenicity of cocaine in humans. Journal of Pediatrics, 110(1), 93-96; Frank, D. A., Bresnahan, K. and Zuckerman, B. S. (1993). Maternal cocaine use: Impact on child health and development. Advances in Pediatrics, 40, 65-99; Glantz, J. C. and Woods, J. R. (1993). Cocaine, heroin, and phencyclidine: Obstetric perspectives. Clinical Obstetrics and Gynecology, 36(2), 279-301.

100. Frank, D. A. and Zuckerman, B. S. (1993). Children exposed to cocaine prenatally: Pieces of the puzzle. Neurotoxicology and Teratology, 15(5), 298-300; Jacobson, J. L., Jacobson, S. W., Sokol, R. J., Martier, S. S., Ager, J. W. and Shankaran, S. (1994). Effects of alcohol use, smoking, and illicit drug use on fetal growth in black infants. Journal of Pediatrics, 124(5, Pt. 1), 757-764.

101. Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Obstetrics and Gynecology Clinics of North America, 17(4), 905-25; Chasnoff, I. J. (1991). Cocaine and pregnancy: Clinical and methodological issues. Clinics in Perinatology, 18(1), 113-123; Frank, D. A. and Zuckerman, B. S. (1993). Children exposed to cocaine prenatally: Pieces of the puzzle. Neurotoxicology and Teratology, 15(5), 298-300; Robbins, C. (1989). Sex differences in psychological consequences of alcohol and drug abuse. Journal of Health and Social Behavior, 30(1), 117-130; Finnegan, L. P., Sloboda, Z., Haverkos, H. W., Mello, N. K. Kreek, M. J., Cottler, L. B. and Frank, D. A. (1995). Drug abuse and the health of women. In L. S. Harris (Ed.), Problems of drug dependence, 1994: Proceedings of the 56th Annual Scientific Meeting. The College on Problems of Drug Dependence, Inc., volume I, NIDA research monograph 152 (pp. 45-48); Robins, L. N., Mills, J. L., Krulewitch, C. and Herman, A. A. (1993). Effects of in utero exposure to street drugs. American Journal of Public Health, 83(Suppl.), 9-32.

102. Bateman, D. A., Ng, S. K., Hansen, C. A. and Heagarty, M. C. (1993). The effects of intrauterine cocaine exposure in newborns. American Journal of Public Health, 83(2) 190-193. But another study, which controlled for age, race, gestational age, marital status, tobacco, alcohol and illicit drug use, found no direct role for cocaine. See Richardson, G. A. and Day, N. L. (1994). Detrimental effects of prenatal cocaine exposure: Illusion or reality? Journal of the American Academy of Child and Adolescent Psychiatry, 33(1), 28-34.

103. Chasnoff, I. J. (1988). Newborn infants with drug withdrawal symptoms. Pediatrics in Review, 9(9), 273-277; Finnegan, L. P. and Weiner, S. M. (1993). Drug withdrawal in the neonate. In M. Gardner, G. B. Merenstein (Eds.), Neonatal intensive care (3rd ed.) (pp. 40-54). St. Louis, MO: Mosby-Year Book; Doberczak, T. M., Shanzer, S., Senie, R. T. and Kandall, S. R. (1988). Neonatal neurologic and electroencephalographic effects of intrauterine cocaine exposure. Journal of Pediatrics, 113(2), 354-8. But another study did not find jitteriness to be common among infants exposed to cocaine prenatally. See Parker, S., Zuckerman, B., Bauchner, H., Frank, D., Vinci, R and Cabral, H. (1990). Jitteriness in full-term neonates: prevalence and correlates. Pediatrics 85(1), 17-23.

104. White, E. (1992). Foster parenting the drug-affected baby. Zero to Three, 13(1), 13-17; Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321

105. Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321; Schneider, J. W. and Chasnoff, I. J. (1992). Motor assessment of cocaine/polydrug exposed infants at age 4 months. Neurotoxicology and Teratology, 14(2), 97-101; Griffith, D. R. (1992). Prenatal exposure to cocaine and other drugs: Developmental and educational prognoses. Phi Delta Kappan, 74, 30-34; Chasnoff, I. J., Griffith, D. R., MacGregor, S., Dirkes, K. and Burns, K. A. (1989). Temporal patterns of cocaine use in pregnancy: Perinatal outcome. JAMA, 261(12), 1741-1744; Mirochnick, M., Meyer, J., Cole, J., Herren, T. and Zuckerman, B. (1991). Circulating catecholamine concentrations in cocaine-exposed neonates: A pilot study. Pediatrics 88(3), 481-5. But see Hutchings, D. E. (1993). The puzzle of cocaine's effects following maternal use during pregnancy: Are there

- reconcilable differences? Neurotoxicology and Teratology, 15(5), 281-286; and Frank, D. A. and Zuckerman, B. S. (1993). Children exposed to cocaine prenatally: Pieces of the puzzle. Neurotoxicology and Teratology, 15(5), 298-300.
106. Griffith, D. R. (1992). Prenatal exposure to cocaine and other drugs: Developmental and educational prognoses. Phi Delta Kappan, 74, 30-34.
107. Griffith, D. R., Azuma, S. D. and Chasnoff, I. J. (1994). Three-year outcome of children exposed prenatally to drugs. Journal of the American Academy of Child Adolescent Psychiatry, 33(1), 20-27; Waller, M. B. (1994). Crack babies grow up. American School Board Journal, 181(6), 30-31.
108. Griffith, D. R., Azuma, S. D. and Chasnoff, I. J. (1994). Three-year outcome of children exposed prenatally to drugs. Journal of the American Academy of Child Adolescent Psychiatry, 33(1), 20-27; Azuma, S. D. and Chasnoff, I. J. (1993). Outcome of children prenatally exposed to cocaine and other drugs: A path analysis of three-year data. Pediatrics, 92(3), 396-402; Chasnoff, I. J., Griffith, D. R., Freier, C. and Murray, J. (1992). Cocaine/polydrug use in pregnancy: Two-year follow-up. Pediatrics, 89(2), 284-289.
109. Azuma, S. D. and Chasnoff, I. J. (1993). Outcome of children prenatally exposed to cocaine and other drugs: A path analysis of three-year data. Pediatrics, 92(3), 396-402.
110. Azuma, S. D. and Chasnoff, I. J. (1993). Outcome of children prenatally exposed to cocaine and other drugs: A path analysis of three-year data. Pediatrics, 92(3), 396-402; Hill, J. (Ed.). (1994). Vulnerable women and visionary programs: Safety net programs for drug-involved women and their children. Washington, DC: National Public Health and Hospital Institute; Malakoff, M. E., Mayes, L. C., and Schottenfeld, R. S. (1994). Language abilities of preschool-age children living with cocaine-using mothers. American Journal on Addictions, 3(4), 346-54; Ellis, J. E., Byrd, L. D., Sexson, W. R., and Patterson-Barnett, C. A. (1993). In utero exposure to cocaine: A review. Southern Medical Journal, 86(7), 725-31.
111. Chasnoff, I. J., Burns, K. A. and Burns, W. J. (1987). Cocaine use in pregnancy: Perinatal morbidity and mortality. Neurotoxicology and Teratology, 9, 291-293.
112. Lounsbury, B., Lifshitz, M. and Wilson, G. S. (1990). In utero exposure to cocaine and the risk of SIDS. In L. S. Harris (Ed.), Problems of drug dependence 1989: Proceeding of the 51st Annual Scientific Meeting of the Committee on Problems of Drug Dependence, Inc. NIDA Research Monograph (pp. 352). Rockville, MD: National Institute on Drug Abuse; Kandall, S. R. and Gaines, J. (1991). Maternal substance use and subsequent sudden infant death syndrome (SIDS) in offspring. Neurotoxicology and Teratology, 13(2), 235-240; Bauchner, H., Zuckerman, B., McClain, M., Frank, D., Fried, L. and Kayne, H. (1988). Risk of sudden infant death syndrome among infants with in utero exposure to cocaine. Journal of Pediatrics, 113(5), 831-834; Bauchner, H. and Zuckerman, B. (1990). Cocaine, sudden infant death syndrome, and home monitoring. Journal of Pediatrics, 117(6), 904-906.

113. Robins, L. N., Mills, J. L., Krulewitch, C. and Herman, A. A. (1993). Effects of in utero exposure to street drugs. American Journal of Public Health, 83(Suppl.), 9-32; Zuckerman, B. and Bresnahan, K. (1991). Developmental and behavioral consequences of prenatal drug and alcohol exposure. Pediatric Clinics of North America, 38(6), 1387-1406; Jones, T. B., Smith, R. S., Martier, S., Dombrowski, M. P. and Sokol, R. J. (1994). Women, children and addiction: Alcohol and drug use in pregnancy: Effects on the offspring. In N. S. Miller (Ed.), Principles of addiction medicine, (pp. 1-8). Chevy Chase, MD: American Society of Addiction Medicine; Zuckerman, B. and Frank, D. A. (1992). Prenatal cocaine and marijuana exposure: Research and clinical implications. In I. S. Zagon, T. A. Slotkin (Eds.), Maternal substance abuse and the developing nervous system (pp. 125-153). San Diego, CA: Academic Press.
114. Robins, L. N., Mills, J. L., Krulewitch, C. and Herman, A. A. (1993). Effects of in utero exposure to street drugs. American Journal of Public Health, 83(Suppl.), 9-32; Richardson, G. A. and Day, N. L. (1994). Detrimental effects of cocaine exposure: Illusion or reality? Journal of the American Academy of Child and Adolescent Psychiatry, 33(1), 28-34; Lutiger, B., Graham, K., Einarson, T. R. and Koren, G. (1991). Relationship between gestational cocaine use and pregnancy outcome: A meta-analysis. Teratology, 44(4), 405-414; Chavez, G. F., Mulinare, J. and Cordero, J. F. (1989). Maternal cocaine use during early pregnancy as a risk factor for congenital urogenital anomalies. JAMA, 262(6), 795-798.
115. Doberczak, T. M., Thornton, J. C., Bernstein, J. and Kandall, S. R. (1987). Impact of maternal drug dependency on birth weight and head circumference of offspring. American Journal of Diseases, 141(11), 1161-1167; Glantz, J. C. and Woods, J. R. (1993). Cocaine, heroin, and phencyclidine: Obstetric perspectives. Clinical Obstetrics and Gynecology, 36(2), 279-301; Chasnoff, I. J., Burns, K. A., Burns, W. J. and Schnoll, S. H. (1986). Prenatal drug exposure: Effects on neonatal and infant growth and development. Neurobehavioral Toxicology and Teratology, 8(4), 357-362; Chasnoff, I. J. (1989). Substance abuse: Pregnancy and the neonate. In M. Rathi (Ed.), Current perinatology (pp. 55-65). New York: Springer-Verlag.
116. Chasnoff, I. J. (1989). Substance abuse: Pregnancy and the neonate. In M. Rathi (Ed.), Current perinatology (pp. 55-65). New York: Springer-Verlag.; Finnegan, L. P., Hagan, T., and Kaltenbach, K. A. (1991). Scientific foundation of clinical practice: Opiate use in pregnant women. Bulletin of the New York Academy of Medicine, 67(3), 223-239; Finnegan, L. P. and Weiner, S. M. (1993). Drug withdrawal in the neonate. In M. Gardner, G. B. Merenstein (Eds.), Neonatal intensive care (3rd ed.) (pp. 40-54). St. Louis, MO: Mosby-Year Book; Chasnoff, I. J. (1988). Newborn infants with drug withdrawal symptoms. Pediatrics in Review, 9(9), 273-277; Finnegan, L. P. and Kaltenbach, K. (1992). Neonatal abstinence syndrome. In Hoekelman, R. A. and Nelson, N. (Eds.), Primary Pediatric Care, 2nd Ed. St. Louis: Mosby Year Book.

117. Chasnoff, I. J. (1988). Newborn infants with drug withdrawal symptoms. Pediatrics in Review, 9(9), 273-277; Finnegan, L. P. and Weiner, S. M. (1993). Drug withdrawal in the neonate. In M. Gardner, G. B. Merenstein (Eds.), Neonatal intensive care (3rd ed.) (pp. 40-54). St. Louis, MO: Mosby-Year Book.
118. Finnegan, L. P. (1991). Perinatal substance abuse: Comments and perspectives. Seminars in Perinatology, 15(4), 331-9; Zweben, J. E. and Payte, J. T. (1990). Methadone maintenance in the treatment of opioid dependence: A current perspective. Western Journal of Medicine, 152(5), 588-599; Kaltenbach, K. and Finnegan, L. P. (1987). Perinatal and developmental outcome of infants exposed to methadone in-utero. Neurotoxicology and Teratology, 9(4), 311-313; Kaltenbach, K. and Finnegan, L. P. (1986). Neonatal abstinence syndrome, pharmacotherapy and developmental outcome. Neurobehavioral Toxicology and Teratology, 8(4), 353-355; Kaltenbach, K. and Finnegan, L. P. (1984). Developmental outcome of children born to methadone maintained women: A review of longitudinal studies. Neurobehavioral Toxicology and Teratology, 6(4), 271-275; Doberczak, T. M., Thornton, J. C., Bernstein, J. and Kandall, S. R. (1987). Impact of maternal drug dependency on birth weight and head circumference of offspring. American Journal of Diseases, 141(11), 1161-1167; Finnegan, L. P. (1991). Treatment issues for opioid-dependent women during the perinatal period. Journal of Psychoactive Drugs, 23(1), 191-201.
119. Edelin, K. C., Gurganious, L., Golar, K., Oellerich, D., Kyei-Aboagye, K. and Hamid, M. A. (1988). Methadone maintenance in pregnancy: Consequences to care and outcome. Obstetrics and Gynecology, 71(3, Pt. 1), 399-403; Kaltenbach, K. and Finnegan, L. P. (1989). Children exposed to methadone in utero: Assessment of the developmental and cognitive ability. Annals of the New York Academy of Sciences, 562, 360-362; Finnegan, L. P. (1991). Treatment issues for opioid-dependent women during the perinatal period. Journal of Psychoactive Drugs, 23(1), 191-201.
120. Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Health Maintenance Strategies, 17(4), 905-925; DuPont, R. L. and Saylor, K. E. (1992). Depressant substances in adolescent medicine. Pediatrics in Review, 13(10), 381-386.
121. Reinisch, J. M., Sanders, S. A., Mortensen, E. L. and Rubin, D. B. (1995). In utero exposure to phenobarbital and intelligence deficits in adult men. JAMA, 274(19), 1518-1525; Senay, E. C. (1989). Addictive behaviors and benzodiazepines: 1. Abuse liability and physical dependence. Advances in Alcohol and Substance Abuse, 8(1), 107-124.
122. Reinisch, J. M., Sanders, S. A., Mortensen, E. L. and Rubin, D. B. (1995). In utero exposure to phenobarbital and intelligence deficits in adult men. JAMA, 274(19), 1518-1525.
123. Finnegan, L. P. (1986). Neonatal abstinence syndrome: Assessment and pharmacotherapy. In F. F. Rubaltelli, B. Granati (Eds.), Neonatal therapy: An update (pp. 122-146). New York: Elsevier.

124. Reinisch, J. M., Sanders, S. A., Mortensen, E. L. and Rubin, D. B. (1995). In utero exposure to phenobarbital and intelligence deficits in adult men. JAMA, 274(19), 1518-1525.
125. Reinisch, J. M., Sanders, S. A., Mortensen, E. L. and Rubin, D. B. (1995). In utero exposure to phenobarbital and intelligence deficits in adult men. JAMA, 274(19), 1518-1525; Wesson, D. R. and Smith, D. E. (1977). Barbiturates: Their use, misuse and abuse. New York: Human Sciences Press.
126. Greater New York March of Dimes Birth Defects Foundation, New York Community Trust, United Hospital Fund. (1993). Infants at risk: 1990-1991 statistics. New York: Greater New York March of Dimes Birth Defects Foundation, New York Community Trust, United Hospital Fund; Dillon, S. (1995, December 27). Mexican drug dealer pushes speed, helping set off an epidemic in U.S. New York Times, A7; Wren, C. S. (1996, February 14). Sharp rise in use of methamphetamines generates concern. New York Times, A16; U. S. Department of Health and Human Services, Public Health Service. (1994, December). Preliminary estimates from the drug abuse warning network. Rockville, MD: Substance Abuse and Mental Health Services Administration, Office of Applied Studies.
127. Dillon, S. (1995, December 27). Mexican drug dealer pushes speed, helping set off an epidemic in U.S. New York Times, A7; Wren, C. S. (1996, February 14). Sharp rise in use of methamphetamines generates concern. New York Times, A16; Trad, P. V. (1993). Substance abuse in adolescent mothers: Strategies for diagnosis, treatment and prevention. Journal of Substance Abuse Treatment, 10(5), 421-431.
128. Dixon, S. D. (1989). Effects of transplacental exposure to cocaine and methamphetamine on the neonate. Western Journal of Medicine, 150(4), 436-442; Oro, A. S. and Dixon, S. D. (1987). Perinatal cocaine and methamphetamine exposure: Maternal and neonatal correlates. Journal of Pediatrics, 11(4), 571-578.
129. Regan, D. O., Erlich, S. M. and Finnegan, L. P. (1987). Infants of drug addicts: AT risk of child abuse, neglect and placement in foster care. Neurotoxicology and Teratology, 9(4), 315-319; Davis, S. K. (1990). Chemical dependency in women: A description of its effects and outcome on adequate parenting. Journal of Substance Abuse Treatment, 7(4), 225-232; Kelleher, K., Chaffin, M., Hollenberg, J. and Fischer, E. (1994). Alcohol and drug disorders among physically abusive and neglectful parents in a community-based sample. American Journal of Public Health, 84(10), 1586-1590; Mitchel, L. and Savage, C. (1991). The relationship between substance abuse and child abuse. Chicago, IL: National Committee for Prevention of Child Abuse; Wiese, D., Daro, D. and National Center on Child Abuse Prevention Research. (1995). Current trends in child abuse reporting and fatalities: The results of the 1994 annual fifty state survey. Chicago, IL: National Committee to Prevent Child Abuse.
130. Kelleher, K., Chaffin, M., Hollenberg, J. and Fischer, E. (1994). Alcohol and drug disorders among physically abusive and neglectful parents in a community-based sample. American Journal of Public Health, 84(10), 1586-1590.

131. Frieze, I. H. and Schafer, P. C. (1984). Alcohol use and marital violence: Female and male differences in reactions to alcohol. In S.C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp.260-279). New York: Guilford Press; Leonard, K. E. and Jacob, T. (1988). Alcohol, alcoholism and family violence. In V. B. Van Hasselt, R. L. Morrison, A. S. Bellack and M. Hersen (Eds.), Handbook of family violence (pp. 383-406), New York: Plenum Press; Mitchel, L. and Savage, C. (1991). The relationship between substance abuse and child abuse. Chicago, IL: National Committee for Prevention of Child Abuse.
132. Black, R. and Mayer, J. (1980). Parents with special problems: Alcoholism and opiate addiction. Child Abuse and Neglect, 4(1), 45-54.
133. Brooks, C. S., Zuckerman, B., Bamforth, A., Cole, J. and Kaplan-Sanoff, M. (1994). Clinical issues related to substance-involved mothers and their infants. Infant Mental Health Journal, 15(2), 202-217; Fullilove, M. T., Lown, E. A. and Fullilove, R. E. (1992). Crack 'hos and skeezers: Traumatic experiences of women crack users. Journal of Sex Research, 29(2), 275-287; Mitchel, L. and Savage, C. (1991). The relationship between substance abuse and child abuse. Chicago, IL: National Committee for Prevention of Child Abuse.
134. Davis, S. K. (1990). Chemical dependency in women: A description of its effects and outcome on adequate parenting. Journal of Substance Abuse Treatment, 7(4), 225-232; Chasnoff, I. J. (Ed.). (1986). Drug use in pregnancy: Mother and child. Lancaster, England: MTP Press; Kaltenbach, K. A. and Finnegan, L. P. (1989). Prenatal narcotic exposure: Perinatal and developmental effects. Neurotoxicology, 10(3), 597-604; Freier, M. C., Griffith, D. R. and Chasnoff, I. J. (1991). In utero drug exposure: Developmental follow-up and maternal-infant interaction. Seminars in Perinatology, 15(4), 310-316; Miller, B. A. (1990). The interrelationships between alcohol and drugs and family violence. In M. De La Rosa, E. Y. Lambert, B. Gropper (Eds.), Drugs and violence: Causes, correlates, and consequences. NIDA Research Monograph 103 (pp. 177-207). Washington, DC: National Institute on Drug Abuse; Regan, D. O., Erlich, S. M. and Finnegan, L. P. (1987). Infants of drug addicts: At risk for child abuse, neglect, and placement in foster care. Neurotoxicology and Teratology, 9(4), 315-319.
135. Cicero, T. J., Nock, B., O'Connor, L., Adams, M. L., Sewing, B. N. and Meyer, E. R. (1994). Acute alcohol exposure markedly influences male fertility and fetal outcome in the male rat. Life Sciences, 55(12), 901-910; Coombs, R. H. and Ziedonis, D. (Eds.). (1995). Handbook on drug abuse prevention: A comprehensive strategy to prevent the abuse of alcohol and other drugs. Boston: Allyn and Bacon.
136. Little, R. E. and Sing, C. F. (1987). Father's drinking and infant birth weight: Report of an association. Teratology, 36(1), 59-65.
137. Klonoff-Cohen, H. S., Edelstein, S. L., Lefkowitz, E. S., Srinivasen, I. P., Kaegi, D., Chang, J. C. and Wiley, K. J. (1995). Effect of passive smoking and tobacco exposure through breast milk on sudden infant death syndrome. JAMA, 273(10), 795-798.

138. Martinez, F. D., Wright, A. L., Taussig, L. M. and Group Health Medical Associates. (1994). The effect of paternal smoking on the birthweight of newborns whose mothers did not smoke. American Journal of Public Health, 84(9), 1489-1491.
139. Berenson, A. B., Stiglich, N. J., Wilkinson, G. S. and Anderson, G. D. (1991). Drug abuse and other risk factors for physical abuse in pregnancy among white non-Hispanic, black, and Hispanic women. American Journal of Obstetrics and Gynecology, 164(6, Pt. 1), 1491-1499; Berenson, A. B., San Miguel, V. V. and Wilkinson, G. S. (1992). Violence and its relationship to substance use in adolescent pregnancy. Journal of Adolescent Health, 13(6), 470-474; McFarlane, J., Parker, B., Soeken, K. and Bullock, L. (1992). Assessing for abuse during pregnancy: Severity and frequency of injuries and associated entry into prenatal care, JAMA, 267(23), 3176-3178; Gomberg, E. S. L. and Nirenberg, T. D. (Eds.). (1993). Women and substance abuse. Norwood, NJ: Ablex Publishing; Amaro, H., Fried, L. E., Cabral, H. and Zuckerman, B. (1990). Violence during pregnancy and substance use. American Journal of Public Health, 80(5), 575-579; Kantor, G. K. and Straus, M. A. (1989). Substance abuse as a precipitant of wife abuse victimizations. American Journal of Drug and Alcohol Abuse, 15(2), 173-189; Parker, B., McFarlane, J. and Soeken, K. (1994). Abuse during pregnancy: Effects on maternal complications and birth weight in adult and teenage women. Obstetrics and Gynecology, 84(3), 323-328.
140. Center on Addiction and Substance Abuse at Columbia University. (1996).
141. U.S. General Accounting Office. (1990). Drug-exposed infants: A generation at risk. Washington, DC: U.S. General Accounting Office.
142. Phibbs, C. S., Bateman, D. A. and Schwartz, R. M. (1991). Neonatal costs of maternal cocaine use. Journal of the American Medical Association, 266(11), 1521-1526.
143. Joyce, T., Racine, A. D., McCalla, S. and Wehbeh, H. (1995). The impact of prenatal exposure to cocaine on newborn costs and length of stay. Health Services Research, 30(2), 341-358.
144. Lillie-Blanton, M. (1991). Meeting the needs of boarder babies and children of substance abusers in the District of Columbia: Mobilizing public and private sector resources for the challenge. Baltimore, MD, Washington, DC: Health Program Alliance, Johns Hopkins University, School of Hygiene and Public Health, and Center for Applied Research and Urban Policy, University of the District of Columbia; U. S. House of Representatives, Committee on Ways and Means, Subcommittee on Human Resources. (1990). The enemy within: Crack-cocaine and America's families. Washington, DC: U.S. Government Printing Office; Phibbs, C. S., Bateman, D. A. and Schwartz, R. M. (1991). Neonatal costs of maternal cocaine use. Journal of the American Medical Association, 266(11), 1521-1526.
145. Lillie-Blanton, M. (1991). Meeting the needs of boarder babies and children of substance abusers in the District of Columbia: Mobilizing public and private sector resources for the challenge. [Baltimore, MD, Washington, DC]: Health Program Alliance, Johns

Hopkins University, School of Hygiene and Public Health, and Center for Applied Research and Urban Policy, University of the District of Columbia.

146. U.S. General Accounting Office. (1990). Drug-exposed infants: A generation at risk. Washington, DC: U.S. General Accounting Office.

147. U.S. General Accounting Office. (1990). Drug-exposed infants: A generation at risk. Washington, DC: U.S. General Accounting Office.

148. U.S. General Accounting Office. (1994). Foster Care: Parental drug abuse has alarming impact on young children. Washington, DC: U.S. General Accounting Office.

149. The substance abuse-related cost of foster care is based on a study of kinship foster care cases active as of September 30, 1991, which found that substance abuse was a reason for placement in 77 percent of cases (New York City Human Resources Administration. (1992, August 31). Preliminary report on kinship foster family profile. Memorandum from George Gabel, Director, Office of Management Analysis to Robert L. Little, Executive Deputy Commissioner, p. 10). It was assumed that the substance abuse factor is the same for placement of children in kinship foster care as for placement with non-related foster parents. Therefore, the 77 percent was applied to the total foster cost. The total budget for foster care in New York City for FY 1994 was \$774 million (New York City Human Resources Administration, Program Operations, Data Analysis and Research, unpublished data). There were 43,484 children in foster care in FY 1995 (New York City Mayor's Office of Operations. (1995). Mayor's management report, Fiscal 1995, Volume II: Agency and citywide indicators. New York: Author, p. 102).

150. The cost is based on the number of children in foster care in the U.S. (approximately 400,000) and an estimated \$10,000 annual cost per child for foster care.

151. Rice, D. P., Kelman, S., Miller, L. S. (1991). Estimates of economic costs of alcohol and drug abuse and mental illness, 1985 and 1988. Public Health Reports, 106(3), 280-291; Rice, D. (1993). The economic cost of alcohol abuse and alcohol dependence: 1990. Alcohol Health and Research World, 17(1), 10-11.

152. DiFranza, J. R. and Lew, R. A. (1995). Effect of maternal cigarette smoking on pregnancy complications and sudden infant death syndrome. Journal of Family Practice, 40(4), 385-394.

153. Abel, E. L. and Sokol, R. J. (1991). A revised conservative estimate of the incidence of FAS and its economic impact. Alcoholism: Clinical and Experimental Research, 15(3), 514-524; Abel, E. L. and Sokol, R. J. (1987). Incidence of fetal alcohol syndrome and economic impact of FAS-related anomalies. Drug and Alcohol Dependence, 19(1), 51-70.

CHAPTER V.

REFERENCES

1. Nirenberg, T. D. and Gomberg, E. S. L. (1993). Prevention of alcohol and drug problems among women. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 339-359). Norwood, NJ: Ablex Publishing.
2. Marlatt, G. A. (1984). Sex differences in psychosocial alcohol research: Implications for prevention. In Women and alcohol: Health related issues: Proceedings of a conference May 23-25, 1984, NIDA research monograph 16 (pp. 260-271.). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; Beckman, L. J. (1994). Treatment needs of women with alcohol problems. Alcohol Health and Research World, 18(3), 206-211; Abbey, A., Ross, L. T., McDuffie, D. (1994). Alcohol's role in sexual assault. In R. R. Watson (Ed.), Addictive behaviors in women (pp.97-124). Totowa, NJ: Humana Press; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; Wilsnack, S. C. (1984). Drinking, sexuality and sexual dysfunction in women. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 189-227). New York: Guilford Press; Norris, J. (1994). Alcohol and female sexuality: A look at expectancies and risks. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 197-201). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146; George, W. H., Gourmic, S. J. and McAfee, M. P. (1988). Perceptions of postdrinking female sexuality: Effects of gender, beverage choice and drink payment. Journal of Applied Psychology, 18(5), 1295-1317; Frieze, I. H. and Schafer, P. C. (1984). Alcohol use and marital violence: Female and male differences in reactions to alcohol. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 260-279). New York: Guilford Press.
3. Barth, R. P., Pietrzak, J. and Ramler, M. (1993). Families living with drugs and HIV: Intervention and treatment strategies. New York: Guilford Press; Smith, P. H. and Chescheir, N. (1995). Battered substance-abusing women: A vulnerable and underserved population in North Carolina: Final report submitted to North Carolina Governor's Institute for Alcohol and Substance Abuse. Raleigh, NC: North Carolina Governor's Institute for Alcohol and Substance Abuse.
4. Novacek, J., Raskin, R. and Hogan, R. (1991). Why do adolescents use drugs?: Age, sex, and user differences. Journal of Youth and Adolescence, 20(5), 475-492.

5. Wilsnack, S. C. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.). The psychology of women's health: Progress and challenges in research and application. Washington, DC: American Psychological Press.
6. Gomberg, E. S. L. (1994). Risk factors for drinking over a woman's life span. Alcohol Health and Research World, 18(3), 220-227; Windle, M. and Barnes, G. M. (1988). Similarities and differences in correlates of alcohol consumption and problem behaviors among male and female adolescents. International Journal of Addictions, 23(7), 707-728; Marlatt, G. A. (1984). Sex differences in psychosocial alcohol research: Implications for prevention. In Women and alcohol: Health related issues: Proceedings of a conference May 23-25, 1984, NIDA research monograph 16 (pp. 260-271.). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; Liu, X. and Kaplan, H. B. (1996). Gender-related differences in circumstances surrounding initiation and escalation of alcohol and other substance use/abuse. Deviant Behavior: An Interdisciplinary Journal, 17(1), 71-106.
7. Swan, N. (1995). Early childhood behavior and temperament predict later substance use. NIDA Notes, 10(1), 13-18; Johnson, R. E. and Marcos, A. C. (1988). Correlates of adolescent drug use by gender and geographic location. American Journal of Drug Abuse, 14(1), 51-63; Diem, E. C., McKay, L. C. and Jamieson, J. L. (1994). Female adolescent alcohol, cigarette and marijuana use: Similarities and differences in patterns of use. International Journal of the Addictions, 29(8), 987-997; Gomberg, E. S. L. (1994). Risk factors for drinking over a woman's life span. Alcohol Health and Research World, 18(3), 220-227; Pope, S. K., Smith, P. D., et al. (1994). Gender differences in rural adolescent drinking patterns. Journal of Adolescent Health, 15(5), 359-365; Thompson, K. M. and Wilsnack, R. W. (1984). Drinking and drinking related problems among female adolescents: Patterns and influences. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 37-65). New York: Guilford Press; Kandel, D. B. and Davies, M. (1992). Progression to regular marijuana involvement: Phenomenology and risk factors for near-daily use. In M. Glantz and R. Pickens (Eds.), Vulnerability to drug abuse (pp. 211-253). Washington, DC: American Psychological Association.
8. Pope, S. K., Smith, P. D., et al. (1994). Gender differences in rural adolescent drinking patterns. Journal of Adolescent Health, 15(5), 359-365.
9. Kandel, D. B. and Yamaguchi, K. (1993). From beer to crack: Developmental patterns of drug involvement. American Journal of Public Health, 83(6), 851-855; Kandel, D. B., Kazuo, Y. and Chen, K. (1992). Stages of progression in drug involvement from adolescence to adulthood: Further evidence for the gateway theory. Journal of Studies on Alcohol, 53(5), 447-457; Mezzich, A. C., Moss, H., Tarter, R. E., Wolfenstein, M., Hsieh, Y. and Mauss, R. (1994). Gender differences in the pattern and progression of substance use in conduct-disordered adolescents. American Journal on Addictions, 3(4), 289-295.

10. U. S. Surgeon General. (1994). Preventing tobacco use among young people. Washington, DC: U.S. Department of Health and Human Services, Office on Smoking and Health.
11. Centers for Disease Control. (1995). Indicators of nicotine addiction among women: United States, 1991-1992. Morbidity and Mortality Weekly Report, 44(6), 103-105.
12. Breslau, N. and Peterson, E. L. (1996). Smoking cessation in young adults: Age at initiation of cigarette smoking and other suspected influences. American Journal of Public Health, 86(2), 214-220.
13. Gilpin, E. A., Lee, L., Evans, N. and Pierce, J. P. (1994). Smoking initiation rates in adults and minors: United States, 1944-1988. American Journal of Epidemiology, 140(6), 535-543.
14. Camp, D. E., Klesges, R. C. and Relyea, G. (1993). The relationship between body weight concerns and adolescent smoking. Health Psychology, 12(1), 24-32.
15. Patton, G. C., Hibbert, M., Rosier, M. J., Carlin, J. B., Caust, J. and Bowes, G. (1996). Is smoking associated with depression and anxiety in teenagers? American Journal of Public Health, 86(2), 225-230; French, S. A., Perry, C. L., Leon, G. R. and Fulkerson, J. A. (1994). Weight concerns, dieting behavior, and smoking initiation among adolescents: A prospective study. American Journal of Public Health, 84(11), 1818-1820; Clayton, S. (1991). Gender differences in psychosocial determinants of adolescent smoking. Journal of School Health, 61(3), 115-120; Novacek, J., Raskin, R. and Hogan, R. (1991). Why do adolescents use drugs?: Age, sex, and user differences. Journal of Youth and Adolescence, 20(5), 475-492; Surgeon General (1988). The health consequences of smoking: Nicotine addiction. Washington, DC: U.S. Department of Health and Human Services, Public Health Service, Office on Smoking and Health.
16. Novacek, J., Raskin, R. and Hogan, R. (1991). Why do adolescents use drugs?: Age, sex, and user differences. Journal of Youth and Adolescence, 20(5), 475-492; Watson, R. R. (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press.
17. Schifano, F., Forza, G. and Gallimberti, L. (1994). Smoking habit and psychological distress in adolescent female students. American Journal on Addictions, 3(2), 100-105.
18. Watson, R. R. (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press.
19. Kessler, L. (1989). Women's magazines' coverage of smoking related health hazards. Journalism Quarterly, 66(2), 316-323; Warner, K. E. (1985). Cigarette advertising and media coverage of smoking and health. New England Journal of Medicine, 312(6), 384-388.
20. Kessler, L. (1989). Women's magazines' coverage of smoking related health hazards. Journalism Quarterly, 66(2), 316-323.

21. Beck, K. H. and Summons, T. G. (1987). Adolescent gender differences in alcohol beliefs and behaviors. Journal of Alcohol and Drug Education, 33(1), 31-43.
22. Pascale, P. J., Trucksis, F. E. and Sylvester, J. (1985). Regional trends and sex differences of drug use and attitudes of high school students in northeast Ohio 1977-1983. Journal of Drug Education, 15(3), 241-250.
23. DiFranza, J. R., Savageau, J. A. and Aisquith, B. F. (1996). Youth access to tobacco: The effects of age, gender, vending machine locks, and "It's the Law" programs. American Journal of Public Health, 86(2), 221-224.
24. Blume, S. B. (1992). Alcohol and other drug problems in women. In J. H. Lowinson, P. Ruiz, R. B. Millman and J. G. Langrod (Eds.), Substance abuse: A comprehensive textbook (pp. 794-807). Baltimore, Hong Kong, London, Munich, Philadelphia, Sydney, Tokyo: Williams and Wilkins.
25. Wechsler, H., Dowdall, G., Davenport, A. and Castillo, S. (1995). Correlates of college student binge drinking. American Journal of Public Health, 85(7), 921-926; Wechsler, H. and Isaac, N. (1992). "Binge" drinking at Massachusetts colleges. Journal of the American Medical Association, 267(21), 2929-2931; Wechsler, H., Davenport, A., Dowdall, G., Moeykens, B. and Castillo, S. (1994). Health and behavioral consequences of binge drinking in college. Journal of the American Medical Association, 272(21), 1672-1677.
26. Mercer, P. W. and Khavari, K. A. (1990). Are women drinking more like men? An empirical examination of the convergence hypothesis. Alcoholism: Clinical and Experimental Research, 3, 461-466.
27. Wechsler, H., Kuh, G. and Davenport, A. (In Press). Fraternities, sororities and binge drinking: Results from a national study of American colleges. NASPA.
28. Lo, C. C. and Globetti, G. (1995). Addiction, 90, 1547-1551.
29. Celis, W. C. (1994, February 16). Drinking by college women raises new concern. New York Times, p. A18.
30. Morrissey, E. R. (1984). Of women, by women, or for women? Selected issues in the primary prevention of drinking problems. In National Institute on Alcohol Abuse and Alcoholism Women and alcohol: Health-related issues: Proceedings from a conference, May 23-25, 1984. NIAAA Research Monograph No. 16 (pp. 226-271). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism.
31. Morrissey, E. R. (1984). Of women, by women, or for women? Selected issues in the primary prevention of drinking problems. In National Institute on Alcohol Abuse and Alcoholism Women and alcohol: Health-related issues: Proceedings from a conference, May 23-25, 1984. NIAAA Research Monograph No. 16 (pp. 226-271). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism. But other studies have found that heavy drinking

rates generally don't differ between employed women and homemakers. See Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association.

32. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; LaRosa, J. H. (1990). Executive women and health: Perceptions and practices. American Journal of Public Health, 80(12), 1450-1454.

33. Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.

34. Springhouse Corporation. (1994). Illustrated manual of nursing practice. 2nd ed. Springhouse, PA: Springhouse Corporation; Carnegie Corporation of New York (1994). Starting points: Meeting the needs of our youngest children. New York: Carnegie Corporation of New York; Trachtenberg, A. I. and Fleming, M. F. (1994). Diagnosis and treatment of drug abuse in family practice. Kansas City, Missouri: American Academy of Family Physicians

35. Centers for Disease Control and Prevention. (1995). Sociodemographic and behavioral characteristics associated with alcohol consumption during pregnancy: United States, 1988. Morbidity and Mortality Weekly Report, 44(13), 261-264.

36. Watson, R. R. (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press; Lange, W. R., White, N. and Robinson, N. (1992). Medical complications of substance abuse. Substance Abuse, 92(3), 205-214; Mello, N. K., Mendelson, J. H. and Teoh, S. K. (1989). Neuroendocrine consequences of alcohol abuse in women. Annals of the New York Academy of Sciences, 562, 211-240.

37. Clarren, S. K. (1986). Prepared statement in placing infants at risk: Parental addiction and disease. Hearing before the Select Committee on Children, Youth and Families, George Miller, Chairman, U.S. House of Representatives, Ninety-Ninth Congress, Second Session, held in Washington, DC on May 21, 1986. Washington, DC: U.S. House of Representatives; Clarren, S. K. (1988). Fetal alcohol syndrome: Diagnosis, treatment, and mechanisms of teratogenesis. In R. Bellisario and G. J. Mizejewski (Eds.), Transplacental disorders: Perinatal detection, treatment, and management (including pediatric AIDS) (pp. 37-55). Albany, NY: Alan R. Liss; Abel, E.L. and Sokol, R.J. (1987). Incidence of fetal alcohol syndrome and economic impact of FAS-related anomalies. Alcohol and Drug Dependence 19, 51-70; Blume, S. B. (1984). Women and alcohol: Public policy issues. In Women and alcohol: Health-related issues, NIAAA research monograph 16 (pp. 294-311). Rockville, MD: National Institute on alcohol Abuse and Alcoholism

38. Abel, E. L. (1988). Fetal alcohol syndrome in families. Neurotoxicology and Teratology, 10(1), 1-2.
39. Day, N. L. and Richardson, G. A. (1991). Prenatal alcohol exposure: A continuum of effects. Seminars in Perinatology, 15(4), 271-279; Abel, E. L. and Sokol, R. J. (1987). Incidence of fetal alcohol syndrome and economic impact of FAS-related anomalies. Alcohol and Drug Dependence 19, 51-70.
40. Streissguth, A. P. (1994). Fetal alcohol syndrome: Understanding the problem: Understanding the solution: What Indian communities can do. American Indian Culture and Research Journal, 18(3), 45-83; Clarren, S. K. (1988). Fetal alcohol syndrome: Diagnosis, treatment, and mechanisms of teratogenesis. In R. Bellisario and G. J. Mizejewski (Eds.), Transplacental disorders: Perinatal detection, treatment, and management (including pediatric AIDS) (pp. 37-55). Albany, NY: Alan R. Liss; Little, B. B., Snell, L. M., Rosenfeld, C. R., Gilstrap, L. C. and Gant, N. F. (1990). Failure to recognize fetal alcohol syndrome in newborn infants. American Journal of Diseases of Children, 144(10), 1142-1146; Clarren, S. K. (1990). Maternal alcohol ingestion: Effects on the developing fetus. Current Pediatric Therapy, 13, 706-707; Stratton, K., Howe, C., Battaglia, F. and Institute of Medicine. (1995). Fetal alcohol syndrome: Diagnosis, epidemiology, prevention, and treatment. Washington, DC: National Academy Press.
41. Clarren, S. K. (1988). Fetal alcohol syndrome: Diagnosis, treatment, and mechanisms of teratogenesis. In R. Bellisario and G. J. Mizejewski (Eds.), Transplacental disorders: Perinatal detection, treatment, and management (including pediatric AIDS) (pp. 37-55). Albany, NY: Alan R. Liss.
42. Morrissey, E. R. (1984). Of women, by women, or for women? Selected issues in the primary prevention of drinking problems. In National Institute on Alcohol Abuse and Alcoholism Women and alcohol: Health-related issues: Proceedings from a conference, May 23-25, 1984. NIAAA Research Monograph No. 16 (pp. 226-271). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; Blume, S. B. (1984). Women and alcohol: Public policy issues. In Women and alcohol: Health-related issues, NIAAA research monograph 16 (pp. 294-311). Rockville, MD: National Institute on alcohol Abuse and Alcoholism; R. C. Engs (Ed.). Women: Alcohol and other drugs. Dubuque, IA: Kendall/Hunt Publishing.
43. Hankin, J. R., Firestone, I. J., Sloan, J. J., Ager, J. W., Sokol, R. J. and Matier, S. S. (1996). Heeding the alcoholic beverage warning label during pregnancy: Multiparae versus nulliparae. Journal of Studies on Alcohol, 57(2), 171-177.
44. Hankin, J. R., Sloan, J. J., Firestone, I. J., Ager, J. W., Sokol, R. J. and Martier, S. S. (1993). A time series analysis of the impact of the alcohol warning label on antenatal drinking. Alcoholism: Clinical and Experimental Research, 17(2), 284-289.

45. Barrett, M. E., Wong, F. Y. and McKay, D. R. R. (1993). Self-reported alcohol use among women of childbearing age and their knowledge of alcohol warning labels and signs. Archives of Family Medicine, 2(12), 1260-1264.
46. Weiner, L., Morse, B. A. and Garrido, P. (1989). FAS/FAE: Focusing prevention on women at risk. International Journal of the Addictions, 24(5), 385-395; Hankin, J. R., Firestone, I. J., Sloan, J. J., Ager, J. W., Sokol, R. J. and Matrier, S. S. (1996). Heeding the alcoholic beverage warning label during pregnancy: Multiparae versus nulliparae. Journal of Studies on Alcohol, 57(2), 171-177; Hankin, J. R., Sloan, J. J., Firestone, I. J., Ager, J. W., Sokol, R. J. and Martier, S. S. (1993). A time series analysis of the impact of the alcohol warning label on antenatal drinking. Alcoholism: Clinical and Experimental Research, 17(2), 284-289.
47. Gilchrist, L. D. and Gillmore, M. R. (1992). Methodological issues in prevention research on drug use and pregnancy. In M. M. Kilbey and K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 1-17). Rockville, MD: National Institute on Drug Abuse; Barth, R. P., Pietrzak, J. and Ramler, M. (1993). Families living with drugs and HIV: Intervention and treatment strategies. New York: Guilford Press; Grinspoon, L. and Bakalar, J. B. (1990). Harvard Medical School Mental Health Review: Alcohol abuse and dependence. Cambridge, MA: President and Fellows of Harvard College.
48. Little, R. E. and Wendt, J. K. (1993). The effects of maternal drinking in the reproductive period: An epidemiological review. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 191-213). Norwood, NJ: Ablex Publishing; Weiner, L. and Larsson, G. (1987). Clinical prevention of fetal alcohol effects--a reality: Evidence for the effectiveness of intervention. Alcohol Health and Research World, 11(4), 60-63, 92-94; Jacobson, J. L., Jacobson, S. W. and Sokol, R. J. (1994). Effects of prenatal exposure to alcohol, smoking and illicit drugs on postpartum somatic growth. Alcoholism: Clinical and Experimental Research, 18(2), 317-323; Blume, S. B. (1984). Women and alcohol: Public policy issues. In Women and alcohol: Health-related issues. NIAAA Research Monograph 16 (pp. 294-311). Rockville, MD: National Institute on alcohol Abuse and Alcoholism; Li, C. Q., Windsor, R. A., Perkins, L., Goldenberg, R. L. and Lowe, J. B. (1993). The impact on infant birth weight and gestational age of cotinine-validated smoking reduction during pregnancy. JAMA, 269(12), 1519-1524; Center for the Future of Children (1995). The future of children: Low birth weight. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children; U.S. General Accounting Office. (1990). Drug-exposed infants: A generation at risk. Washington, DC: U.S. General Accounting Office; Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321; Nair, B. S. and Watson, R. R. (1991). Cocaine and the pregnant woman. Journal of Reproductive Medicine, 36(12), 862-867; Chasnoff, I. J., Griffith, D. R., MacGregor, S., Dirkes, K. and Burns, K. A. (1989). Temporal patterns of cocaine use in pregnancy: Perinatal outcome. JAMA, 261(12), 1741-1744.

49. Chasnoff, I. J. (1992). Cocaine, pregnancy, and the growing child. Current Problems in Pediatrics, 22(7), 302-321; Racine, A., Joyce, T. and Anderson, R. (1993). Association between prenatal care and birthweight among women exposed to cocaine in New York City. JAMA, 270(13), 1581-1586; Feldman, J. G., Minkoff, H. L., McCalla, S. and Salwen, M. (1992). A cohort study of the impact of perinatal drug use on prematurity in an inner-city population. American Journal of Public Health, 82(5), 726-728; MacGregor, S. N., Keith, L. G., Bachicha, J. A. and Chasnoff, I. J. (1989). Cocaine abuse during pregnancy: Correlation between prenatal care and perinatal outcome. Obstetrics and Gynecology, 74(6), 882-885; Robins, L. N., Mills, J. L., Krulewitch, C. and Herman, A. A. (1993). Effects of in utero exposure to street drugs. American Journal of Public Health, 83(Suppl.), 9-32; Feig, L. (1990). Drug exposed infants and children: Service needs and policy questions. Washington, DC: U.S. Department of Health and Human Services.
50. Barth, R. P., Pietrzak, J. and Ramler, M. (1993). Families living with drugs and HIV: Intervention and treatment strategies. New York: Guilford Press.
51. Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Obstetrics and Gynecology Clinics of North America, 17(4), 905-925; Center for the Future of Children. (1995). The future of children: low birth weight. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children; National Institutes of Health, National Cancer Institute (1994). Tobacco and the clinician: Interventions for medical and dental practice. Smoking and tobacco control monograph 5. Bethesda, MD: National Cancer Institute.
52. Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Obstetrics and Gynecology Clinics of North America, 17(4), 905-925; Watson, R. R. (Ed). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press; Ershoff, D. H., Mullen, P. D. and Quinn, V. P. (1989). A randomized trial of a serialized self-help smoking cessation program for pregnant women in an HMO. American Journal of Public Health, 79(2), 182-187; Burling, T. A., Bigelow, G. E., Robinson, J. C. and Mead, A. M. (1991). Smoking during pregnancy: Reduction via objective assessment and directive advice. Behavior Therapy, 22(1), 31-40; Price, J. H., Desmond, S. M., Roberts, S. M., Krol, R. A., Losh, D. P. and Snyder, F. F. (1991). Comparison of three antismoking interventions among pregnant women in an urban setting: A randomized trial. Psychological Reports, 68(1-2), 595-604; Windsor, R. A., Lowe, J. B., Perkins, L. L., Smith-Yoder, D., Artz, L., Crawford, M., Amburgy, K. and Boyd, N. R. (1993). Health education for pregnant smokers: Its behavioral impact and cost benefit. American Journal of Public Health, 83(2), 201-206; McCrady, B. S. and Raytek, H. (1993). Women and substance abuse: Treatment and outcome modalities. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 314-338). Norwood, NJ: Ablex Publishing.
53. Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Obstetrics and Gynecology Clinics of North America, 17(4), 905-925.

54. Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Obstetrics and Gynecology Clinics of North America, 17(4), 905-925; Floyd, R. L., Rimer, B. K., Giovino, G. A., Mullen, P. D. and Sullivan, S. E. (1993). A review of smoking in pregnancy: Effects on pregnancy outcomes and cessation efforts. Annual Review of Public Health, 14, 379-411.
55. Chasnoff, I. J. (1992). Drug use in pregnancy: Does treatment work? Project report. Unpublished paper.
56. Kogan, M. D., Kotelchuck, M., Alexander, G. R. and Johnson, W. E. (1994). Racial disparities in reported prenatal care advice from health care providers. American Journal of Public Health, 84(1), 82-88.
57. Bingol, N., Schuster, C., Fuchs, M., Iosub, S., Turner, G., Stone, R. K. and Gromisch, D. S. (1987). The influence of socioeconomic factors on the occurrence of fetal alcohol syndrome. Advances in Alcoholism and Substance Abuse, 6(4), 105-118; Ramirez, A. G. and Gallion, K. J. (1993). Nicotine dependence among blacks and Hispanics. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 350-364). New York: Oxford University Press.
58. Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing Company; R. R. Watson (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press; Merrick, J. C. (1993). Maternal substance abuse during pregnancy. Journal of Legal Medicine, 14(1), 57-71.
59. S. S. Brown (Ed.) (1992). Including children and pregnant women in health care reform: Summary of two workshops. Washington, DC: National Academy Press.
60. Barth, R. P., Pietrzak, J. and Ramler, M. (1993). Families living with drugs and HIV: Intervention and treatment strategies. New York: Guilford Press; Ackatz, L., and Jones, E. D. (1992, January). Availability of substance abuse treatment programs for pregnant women: Results from three national surveys. Chicago, IL: National Committee to Prevent Child Abuse; Coalition on Alcohol and Drug Dependent Women and Their Children. ([1990]). Treatment committee survey--1990 [of pregnant and post-partum demonstration projects]. [Washington, DC]: National Council on Alcoholism and Drug Dependence, Coalition on Alcohol and Drug Dependent Women and Their Children; Finkelstein, N. (1994). Treatment issues for alcohol- and drug-dependent pregnant and parenting women. Health and Social Work, 19(1), 7-15; D. R. Gerstein and H. J. Harwood (Eds.). (1990). Treating drug problems: A study of the evolution, effectiveness, and financing of public and private drug treatment systems. Washington, DC: National Academy Press; Klein, D., Zahnd, E., Holtby, S. and Barth, R. (1994). Perinatal needs assessment executive summary. Berkeley, CA: California Health and Welfare Agency, Department of Alcohol and Drug Programs, Office of Perinatal Substance Abuse; Legal Action Center. (1994). Best practices: Models of care for pregnant alcoholic and drug dependent women and their children. New York: Legal Action Center; Morain, C. (1994). Still a long way to go, baby. American Medical News,

37(25), 7-12; Skolnick, A. (1990). Drug screening in prenatal care demands objective medical criteria, support services. JAMA, 264(3), 309-10; U.S. General Accounting Office. (1991). ADMS Block Grant: Women's set-aside does not assure drug treatment for pregnant women. Washington, DC: U.S. General Accounting Office; U.S. House of Representatives, Committee of Ways and Means, Subcommittee on Human Resources. (1990). The enemy within: Crack-cocaine and America's families. Washington, DC: U.S. Government Printing Office; Wallace, B. C. (1991). Chemical dependency treatment for the pregnant crack addict: Beyond the criminal-sanctions perspective. Psychology of Addictive Behavior, 5(1), 23-35; Beckman, L. J. (1994). Treatment needs of women with alcohol problems. Alcohol Health and Research World, 18(3), 206-211; Finnegan, L. P., Hagan, T. and Kaltenbach, K. A. (1991). Scientific foundation of clinical practice: Opiate use in pregnant women. Bulletin of the New York Academy of Medicine, 67(3), 223-239; Kaltenbach, K. A. and Finnegan, L. P. (1992). Studies of prenatal drug exposure and environmental research issues: The benefits of integrating within a treatment program. In M. M. Kilbey and K. Ashgar (Eds.), Methodological issues in epidemiological, prevention and treatment research on drug-exposed women and their children, NIDA Research Monograph 117 (pp. 259-270). Rockville, MD: National Institute on Drug Abuse.

61. Blume, S. B. (1988). Alcohol/drug dependent women: New insights into their special problems, treatment, recovery. Minneapolis, MN: Johnson Institute; Hill, S. Y. (1984). Physiological effects of alcohol in women. In Women and alcohol: Health-related issues, NIAAA Research Monograph 16 (pp. 199-214). Rockville, MD: National Institute on alcohol Abuse and Alcoholism.

62. Wilsnack, S. C. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application. Washington, DC: American Psychological Press; Marcenko, M. O., Spence, M. and Rohweder, C. (1994). Psychosocial characteristics of pregnant women with and without a history of substance abuse. Health and Social Work, 19(1), 17-22.

63. Council on Scientific Affairs, American Medical Association. (1996). Alcoholism in the elderly. JAMA, 275(10), 797-801.

64. Council on Scientific Affairs, American Medical Association. (1996). Alcoholism in the elderly. JAMA, 275(10), 797-801.

65. McCrady, B. S. and Raytek, H. (1993). Women and substance abuse: Treatment and outcome modalities. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 314-338). Norwood, NJ: Ablex Publishing; Kagle, J. D. (1987). Women who drink: Changing images, changing realities. Journal of Social Work Education, 23(3), 21-28; R. R. Watson (Ed.). (1994) Addictive behaviors in women. Totowa, NJ: Humana Press; Vanicelli, M. (1984). Treatment outcome of alcoholic women: The state of the art. Bulletin of the Society of Psychologists in Addictive Behavior, 3(3), 155-162; Murray, J. B. (1989). Psychologists and alcoholic women. Psychology Reports, 64(2), 627-644; Toneatto, A., Sobell, L. C. and Sobell, M. B. (1992). Gender issues in the treatment of

abusers of alcohol, nicotine, and other drugs. Journal of Substance Abuse, 4(2), 209-218; Hesselbrock, M. N. and Hesselbrock, V. M. (1993). Depression and antisocial personality disorder in alcoholism; Gender comparison. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 142-161). Norwood, NJ: Ablex Publishing; N. van den Bergh (Ed.). (1991). Feminist perspectives on addictions. New York: Springer Publishing Company; Quinby, P. M. and Graham, A. V. (1993). Substance abuse among women. Primary Care, 20(1), 131-140.

66. Toneatto, A., Sobell, L. C. and Sobell, M. B. (1992). Gender issues in the treatment of abusers of alcohol, nicotine, and other drugs. Journal of Substance Abuse, 4(2), 209-218; McCrady, B. S. and Raytek, H. (1993). Women and substance abuse: Treatment and outcome modalities. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 314-338). Norwood, NJ: Ablex Publishing; Hesselbrock, M. N. and Hesselbrock, V. M. (1993). Depression and antisocial personality disorder in alcoholism: Gender comparison. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 142-161). Norwood, NJ: Ablex Publishing; Nixon, S. J. and Glenn, S. W. (1995). Cognitive psychosocial performance and recovery in female alcoholics. In M. Galanter (Ed.), Alcoholism and women (pp. 287-307). New York: Plenum Press; Kagle, J. D. (1987). Women who drink: Changing images, changing realities. Journal of Social Work Education, 23(3), 21-28; Watson, R. R. (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press; Vanicelli, M. (1984). Treatment outcome of alcoholic women: The state of the art. Bulletin of the Society of Psychologists in Addictive Behavior, 3(3), 155-162; Murray, J. B. (1989). Psychologists and alcoholic women. Psychology Reports, 64(2), 627-644.

67. Jarvis, T. J. (1992). Implications of gender for alcohol treatment research: A quantitative and qualitative review. British Journal of Addiction, 87(9), 1249-1261.

68. Schmidt, L. and Weisner, C. (1995). The emergence of problem-drinking women as a special population in need of treatment. In M. Galanter (Ed.), Alcoholism and women (pp. 309-334). New York: Plenum Press. See also Williams, G. D., Grant, B. F., Harford, T. C. and Noble, J. (1989). Population projections using DSM-III criteria: Alcohol abuse and dependence, 1990-2000. Alcohol, Health and Research World, 13(4), 366-370; Grant, B. F., Harford, T. C., Dawson, D. A., Chou, P., Dufour, M. and Pickering, R. (1994). Prevalence of DSM-IV Alcohol Abuse and Dependence, United States, 1992. Alcohol, Health and Research World, 18(3), 243-248; Smith, A. R. (1986). Alcoholism and gender: Patterns of diagnosis and response. Journal of Drug Issues, 16(3), 407-420.

69. Schmidt, L. and Weisner, C. (1995). The emergence of problem-drinking women as a special population in need of treatment. In M. Galanter (Ed.), Alcoholism and women (pp. 309-334). New York: Plenum Press; Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C. B., Hughes, M., Eshleman, S., Wittchen, H. and Kendler, K. S. (1994). Lifetime and 12-month Prevalence of DSM-III-R Psychiatric Disorders in the United States: Results from the National Comorbidity Survey. Archives of General Psychiatry, 51(5), 8-19.

70. National Association of State Alcohol and Drug Abuse Directors. (1991). Survey on state alcohol and drug agency use of FY 1989 federal and state funds. Washington, DC: National Association of State Alcohol and Drug Abuse Directors.
71. Miller, G. (1989). Addicted infants and their mothers. Zero to Three, 9(5), 20-23.
72. U.S. Select Committee on Children, Youth, and Families. (1989). Hearing before the Select Committee on Children, Youth and Families: House of Representatives, One Hundred First Congress, hearing held in Washington, DC, April 27, 1989. Washington, DC: U.S. Government Printing Office; Miller, G. (1989). Addicted infants and their mothers. Zero to Three, 9(5), 20-23; U. S. House of Representatives, Committee on Ways and Means, Subcommittee on Human Resources. (1990). The enemy within: Crack-cocaine and America's families. Washington, DC: U.S. Government Printing Office.
73. California Elected Women's Association for Education and Research. (1994). Women and substance abuse. Sacramento, CA: California Elected Women's Association for Education and Research.
74. Seiden, A. M. (1992). Measures of pregnant, drug-abusing women for treatment research. In M. M. Kilbey and K. Ashgar (Eds.), Methodological issues in epidemiological, prevention and treatment research on drug-exposed women and their children, NIDA Research Monograph 117 (pp. 194-211). Rockville, MD: National Institute on Drug Abuse; U.S. General Accounting Office. (1990). Drug-exposed infants: A generation at risk. Washington, DC: U.S. General Accounting Office.
75. Legal Action Center. (1994). Dispelling the myth: Legal issues of treatment programs serving pregnant addicts. New York: Legal Action Center.
76. Wilsnack, S. C. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application. Washington, DC: American Psychological Press; Roman, P. M. (1988). Biological features of women's alcohol use: A review. Public Health Reports, 103(6), 628-637.
77. Beckman, L. J. (1994). Treatment needs of women with alcohol problems. Alcohol Health and Research World, 18(3), 206-211.
78. Blume, S. B. (1988). Alcohol/drug dependent women: New insights into their special problems, treatment, recovery. Minneapolis, MN: Johnson Institute; Blume, S. B. (1984). Women and alcohol: Public policy issues. In Women and alcohol: Health-related issues, NIAAA Research Monograph 16 (pp. 294-311). Rockville, MD: National Institute on alcohol Abuse and Alcoholism.
79. Waller, P. F. and Blow, F. C. (1995). Women, alcohol and driving. In M. Galanter (Ed.), Alcoholism and women (pp. 103-123). New York: Plenum Press; Dunham, R. G.

- (1986). Noticing alcoholism in the elderly and women: A nationwide examination of referral behavior. Journal of Drug Issues, 16(3), 397-406; Blume, S. B. (1984). Women and alcohol: Public policy issues. In Women and alcohol: Health-related issues, NIAAA Research Monograph 16 (pp. 294-311). Rockville, MD: National Institute on alcohol Abuse and Alcoholism.
80. Levy, M. L., Reichman, W., Herrington, S. and Young, D. (1980). Alcoholic women in industry: Some empirical data. International Journal of Group Tensions, 10(1-4), 120-129.
81. Blum, T. C., Roman, P. M. and Harwood, E. M. (1995). Employed women with alcohol problems who seek help from employee assistance programs: Descriptions and comparisons. In M. Galanter (Ed.), Alcoholism and women (pp. 125-156). New York: Plenum Press.
82. Toray, T., Coughlin, C., Vuchinich, S. and Patricelli, P. (1991). Gender differences associated with adolescent substance abuse comparisons and implications for treatment. Family Relations, 40(3), 338-344.
83. Bergmann, P. E., Smith, M. B. and Hoffmann, N. G. (1995). Adolescent treatment: Implications for assessment, practice guidelines, and outcome management. Pediatric Clinics of North America, 42(2), 453-472.
84. Dunham, R. G. (1986). Noticing alcoholism in the elderly and women: A nationwide examination of referral behavior. Journal of Drug Issues, 16(3), 397-406.
85. Dunham, R. G. (1986). Noticing alcoholism in the elderly and women: A nationwide examination of referral behavior. Journal of Drug Issues, 16(3), 397-406.
86. Gomberg, E. S. L. and Nirenberg, T. D. (1993). Antecedents and consequences. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 118-141). Norwood, NJ: Ablex Publishing; Gomberg, E. S. L. (1993). Women and alcohol: Use and abuse. Journal of Nervous and Mental Disease, 181(4), 211-219.
87. Gomberg, E. S. L. (1991). Women and alcohol: Psychosocial aspects. In D. J. Pittman and H. R. White (Eds.), Society, culture and drinking patterns reexamined (pp. 263-84). New Brunswick, NJ: Alcohol Research Documentation, Inc., Rutgers Center of Alcohol Studies; Blume, S. B. (1987). Women's health: Issues in mental health, alcoholism and substance abuse: Alcoholism and women's health. Public Health Reports, 102(Suppl.), 38-42.
88. Blume, S. B. (1987). Women's health: Issues in mental health, alcoholism and substance abuse: Alcoholism and women's health. Public Health Reports, 102(Suppl.), 38-42.
89. Sindelar, J. L. (1982). Differential use of medical care by sex. Journal of Political Economy, 90(5), 1003-10019; Verbrugge, L. M. and Wingard, D. L. (1987). Sex differentials in health and mortality. Women and Health, 12(2), 103-145; Cyr, M. G. and

Moulton, A. W. (1990). Substance abuse in women. Health Maintenance Strategies, 17(4), 905-925; U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. (1996). National ambulatory medical care survey: 1994 summary. Hyattsville, MD: National Center for Health Statistics.

90. Blume, S. B. (1987). Women's health: Issues in mental health, alcoholism and substance abuse: Alcoholism and women's health. Public Health Reports, 102(Suppl.), 38-42; Gearhart, J. G., Beebe, D. K., Milhorn, H. T. and Meeks, G. R. (1991). Alcoholism in women. American Family Physician, 44(3), 907-913; National Institutes of Health, National Cancer Institute (1994). Tobacco and the clinician: Interventions for medical and dental practice. Smoking and tobacco control monograph 5. Bethesda, MD: National Cancer Institute; Cyr, M. G. and Moulton, A. W. (1993). The physician's role in prevention, detection, and treatment of alcohol abuse in women. Psychiatric Annals, 23(8), 454-462.

91. Blume, S. B. (1996). Chemical dependency problems in ob/gyn practice. Primary Care Update for Ob/Gyns, 31(1), 1-8; Halliday, A., Bush, B., Cleary, P., Aronson, M. and Delbanco, T. (1986). Alcohol abuse in women seeking gynecologic care. Obstetrics and Gynecology, 68(3), 322-326.

92. Beckman, L. J. (1994). Treatment needs of women with alcohol problems. Alcohol Health and Research World, 18(3), 206-211; Wilsnack, S. C. and Wilsnack, R. W. (1990). Women and substance abuse: Research directions for the 1990s. Psychology of Addictive Behaviors, 4(1), 46-49; Brown, E. R., Wyn, R., Cumberland, W. G., Yu, H., Abel, E., Gelberg, L. and Ng, L. (1995). Women's health-related behaviors and use of clinical preventive services: A report to the Commonwealth Fund. Los Angeles: UCLA Center for Health Policy Research; Cyr, M. G. and Moulton, A. W. (1993). The physician's role in prevention, detection, and treatment of alcohol abuse in women. Psychiatric Annals, 23(8), 454-462.

93. Dawson, N. V., Dadheech, G., Speroff, T., Smith, R. L. and Schubert, D. S. P. (1992). The effect of patient gender on the prevalence and recognition of alcoholism on a general medicine inpatient service. Journal of General Internal Medicine, 7(1), 38-45.

94. Moore, R. D., Bone, L. R., Geller, G., Mamon, J. A., Stokes, E. J. and Levine, D. M. (1989). Prevalence, detection, and treatment of alcoholism in hospitalized patients. JAMA, 261(3), 403-407.

95. McCrady, B. S. and Raytek, H. (1993). Women and substance abuse: Treatment and outcome modalities. In E. S. L. Gombert and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 314-338). Norwood, NJ: Ablex Publishing; Wilsnack, S. C. and Wilsnack, R. W. (1990). Women and substance abuse: Research directions for the 1990s. Psychology of Addictive Behaviors, 4(1), 46-49; Quinby, P. and Graham, A. V. (1993). Substance abuse among women. Primary Care, 20(1), 131-140; Quinby, P. M. and Graham, A. V. (1993). Substance abuse among women. Primary Care, 20(1), 131-140.

96. N. van den Bergh (Ed.). (1991). Feminist perspectives on addictions. New York: Springer Publishing Company; Smith, A. R. (1986). Alcoholism and gender: Patterns of diagnosis and response. Journal of Drug Issues, 16(3), 407-420; McCrady, B. S. and Raytek, H. (1993). Women and substance abuse: Treatment and outcome modalities. In E. S. L. Gombert and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 314-338). Norwood, NJ: Ablex Publishing; Gearhart, J. G., Beebe, D. K., Milhorn, H. T. and Meeks, G. R. (1991). Alcoholism in women. American Family Physician, 44(3), 907-913.
97. Gearhart, J. G., Beebe, D. K., Milhorn, H. T. and Meeks, G. R. (1991). Alcoholism in women. American Family Physician, 44(3), 907-913; Reed, B. G. (1987). Developing women-sensitive drug dependence treatment services: Why so difficult? Journal of Psychoactive Drugs, 19(2), 151-164; Kagle, J. D. (1987). Women who drink: Changing images, changing realities. Journal of Social Work Education, 23(3), 21-28.
98. Hughes, T. L. (1990). Evaluating research on chemical dependency among women: A women's health perspective. Family and Community Health, 13(3), 35-46; Blume, S. B. (1992). Alcohol and other drug problems in women. In J. H. Lowinson, P. Ruiz, R. B. Millman and J. G. Langrod (Eds.), Substance abuse: A comprehensive textbook (pp. 794-807). Baltimore, Hong Kong, London, Munich, Philadelphia, Sydney, Tokyo: Williams and Wilkins; Cyr, M. G. and Moulton, A. W. (1993). The physician's role in prevention, detection, and treatment of alcohol abuse in women. Psychiatric Annals, 23(8), 454-462.
99. Wilke, D. (1994). Women and alcoholism: How a male-as-norm bias affects research, assessment, and treatment. Health and Social Work, 19(1), 29-35; Hughes, T. L. (1990). Evaluating research on chemical dependency among women: A women's health perspective. Family and Community Health, 13(3), 35-46.
100. Trachtenberg, A. I. and Fleming, M. F. (1994). Diagnosis and treatment of drug abuse in family practice. Kansas City, Missouri: American Academy of Family Physicians; Council on Scientific Affairs, American Medical Association. (1996). Alcoholism in the elderly; JAMA, 275(10), 797-801; American Medical Association (1995). Alcoholism in the elderly: Diagnosis, treatment and prevention guidelines for primary care physicians. Chicago, IL: American Medical Association; Dufour, M. C., Archer, L. and Gordis, E. (1992). Alcohol and the elderly. Clinics in Geriatric Medicine, 8(1), 127-141; Curtis, J. R., Geller, G., Stokes, E. J., Levine, D. M. and Moore, R. D. (1989). Characteristics, diagnosis and treatment of alcoholism in elderly patients. Journal of the American Geriatrics Society, 37(4), 310-316.
101. Dufour, M. C., Archer, L. and Gordis, E. (1992). Alcohol and the elderly. Clinics in Geriatric Medicine, 8(1), 127-141; Curtis, J. R., Geller, G., Stokes, E. J., Levine, D. M. and Moore, R. D. (1989). Characteristics, diagnosis and treatment of alcoholism in elderly patients. Journal of the American Geriatrics Society, 37(4), 310-316.

102. Gomberg, E. S. L. (1995). Older women and alcohol use and abuse. In M. Galanter (Ed.), Alcoholism and Women, (pp. 61-79). New York: Plenum Press; Gomberg, E. S. (1994). Risk factors for drinking over a woman's life span. Alcohol, Health and Research World, 18(3), 220-227; Gomberg, E. S. L. (1989). Alcoholism in women: use of other drugs. Alcoholism; Clinical and experimental research, 13, 338; Celentano, D. D. and McQueen, D. V. (1984). Multiple substance abuse among women with alcohol-related problems. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 97-116). New York: Guilford Press; Council on Scientific Affairs, American Medical Association. (1996). Alcoholism in the elderly; JAMA, 275(10), 797-801.
103. Gearhart, J. G., Beebe, D. K., Milhorn, H. T. and Meeks, G. R. (1991). Alcoholism in women. American Family Physician, 44(3), 907-913; Wilcox, D. O. and Yates, W. R. (1993). Gender and psychiatric comorbidity in substance-abusing individuals. American Journal on Addictions, 2(3), 202-206; Cyr, M. G. and Moulton, A. W. (1993). The physician's role in prevention, detection, and treatment of alcohol abuse in women. Psychiatric Annals, 23(8), 454-462; Council on Scientific Affairs, American Medical Association. (1996). Alcoholism in the elderly. JAMA, 275(10), 797-801.
104. Cyr, M. G. and Moulton, A. W. (1990). Substance abuse in women. Obstetrics and Gynecology Clinics of North America, 17(4), 905-925; Gomberg, E. S. L. and Lisansky, J. M. (1984). Antecedents of alcohol problems in women. In S. C. Wilsnack and L. J. Beckman (eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 233-259). New York: Guilford Press; Leckman, A. L., Umland, B. E. and Blay, M. (1984). Alcoholism in the families of family practice outpatients. Journal of Family Practice, 19(2), 205-207.
105. Blume, S. B. (1985). Women and alcohol. In T. E. Bratter and G. G. Forrest (Eds.) Alcoholism and substance abuse: Strategies for clinical intervention (pp. 623-638). New York: The Free Press.
106. Council on Scientific Affairs, American Medical Association. (1996). Alcoholism in the elderly. JAMA, 275(10), 797-801.
107. National Institute on Alcohol Abuse and Alcoholism. (1995). The physicians' guide to helping patients with alcohol problems. [Rockville, MD]: National Institute on Alcohol Abuse and Alcoholism.
108. Wilsnack, S. C. (1991). Barriers to treatment for alcoholic women. Addiction and Recovery, 11(4), 10-12.
109. Schmidt, L. and Weisner, C. (1995). The emergence of problem-drinking women as a special population in need of treatment. In M. Galanter (Ed.), Alcoholism and women (pp. 309-334). New York: Plenum Press; Blume, S. B. (1991). Women, alcohol, and drugs. In

- N. S. Miller (Ed.), Comprehensive handbook of drug and alcohol addiction (pp. 147-177). New York: Marcel Dekker.
110. Breitbart, V., Chavkin, W. and Wise, P. H. (1994). Accessibility of drug treatment for pregnant women: A survey of programs in five cities. American Journal of Public Health, 84(10), 1658-1661.
111. R. R. Watson (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press; J. Hill (Ed.). (1994). Vulnerable women and visionary programs: Safety net programs for drug-involved women and their children. Washington, DC: National Public Health and Hospital Institute; Chavkin, W., Paone, D., Friedmann, P. and Wilets, I. (1993). Reframing the debate: Toward effective treatment for inner city drug-abusing mothers. Bulletin of the New York Academy of Medicine, 70(1), 50-68.
112. Hughes, P. H., Coletti, S. D., Neri, R. L., Urmann, C. F., Sicilian, D. M., Stahl, S. S. and Anthony, J. C. (1994). Retention of cocaine abusing women in residential treatment. In L. S. Harris (Ed.), Problems of drug dependence, 1993: Proceeding of the 55th Annual Scientific Meeting, The College on Problems of Drug Dependencies, Inc. volume II: Abstracts, NIDA Research Monograph 141 (pp. 4). Rockville, MD: National Institute on Drug Abuse; Hughes, P. H., Coletti, S. D., Neri, R. L., Urmann, C. F., Stahl, S., Sicilian, D. M. and Anthony, J. C. (1995). Retaining cocaine-abusing women in a therapeutic community: The effect of a child live-in program. American Journal of Public Health, 85(8), 1149-1152.
113. Beckman, L. J. and Amaro, H. (1986). Personal and social difficulties faced by women and men entering alcoholism treatment. Journal of Studies on Alcohol, 47(2), 135-145; Murray, J. B. (1989). Psychologists and alcoholic women. Psychology Reports, 64(2), 627-644.
114. McCrady, B. S. and Raytek, H. (1993). Women and substance abuse: Treatment and outcome modalities. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 314-338). Norwood, NJ: Ablex Publishing; O'Connor, L. E., Berry, J. W., Inaba, D., Weiss, J. and Morrison, A. (1994). Shame, guilt, and depression in men and women in recovery from addiction. Journal of Substance Abuse Treatment, 11(6), 503-510; Quinby, P. M. and Graham, A. V. (1993). Substance abuse among women. Primary Care, 20(1), 131-140.
115. Blum, T. C., Roman, P. M. and Harwood, E. M. (1995). Employed women with alcohol problems who seek help from employee assistance programs: Descriptions and comparisons. In M. Galanter (Ed.), Alcoholism and women (pp. 125-156). New York: Plenum Press.
116. Blum, T. C., Roman, P. M. and Harwood, E. M. (1995). Employed women with alcohol problems who seek help from employee assistance programs: Descriptions and

- comparisons. In M. Galanter (Ed.), Alcoholism and women (pp. 125-156). New York: Plenum Press.
117. Blum, T. C., Roman, P. M. and Harwood, E. M. (1995). Employed women with alcohol problems who seek help from employee assistance programs: Descriptions and comparisons. In M. Galanter (Ed.), Alcoholism and women (pp. 125-156). New York: Plenum Press.
118. Waite-O'Brien, N., Clinical Director, Betty Ford Center at Eisenhower. (1996, January). Personal Communication.
119. Blume, S. B. (1991). Women, alcohol, and drugs. In N. S. Miller (Ed.), Comprehensive handbook of drug and alcohol addiction (pp. 147-177). New York: Marcel Dekker; Quinby, P. M. and Graham, A. V. (1993). Substance abuse among women. Primary Care, 20(1), 131-140; Gomberg, E. S. L. (1993). Women and alcohol: Use and abuse. Journal of Nervous and Mental Disease, 181(4), 211-219; Smith, A. R. (1986). Alcoholism and gender: Patterns of diagnosis and response. Journal of Drug Issues, 16(3), 407-420; Beckman, L. J. and Amaro, H. (1986). Personal and social difficulties faced by women and men entering alcoholism treatment. Journal of Studies on Alcohol, 47(2), 135-145.
120. Smith, A. R. (1986). Alcoholism and gender: Patterns of diagnosis and response. Journal of Drug Issues, 16(3), 407-420.
121. Davis, K. (1995). The federal budget and women's health. American Journal of Public Health, 85(8), 1051-1053; D. R. Gerstein and H. J. Harwood (Eds.). (1990). Treating drug problems: A study of the evolution, effectiveness, and financing of public and private drug treatment systems. Washington, DC: National Academy Press; Inciardi, J. A., Lockwood, D. and Pottieger, A. E. (1993). Women and crack-cocaine. New York: Macmillan Publishing Company; Klein, D., Zahnd, E., Holtby, S. and Barth, R. (1994). Perinatal needs assessment executive summary. Berkeley, CA: California Health and Welfare Agency, Department of Alcohol and Drug Programs, Office of Perinatal Substance Abuse; Amaro, H., Beckman, L. J. and Mays, V. M. (1987). A comparison of black and white women entering alcoholism treatment. Journal of Studies on Alcohol, 48(3), 220-228; Schliebner, C. T. (1994). Gender-sensitive therapy: An alternative for women in substance abuse treatment. Journal of Substance Abuse Treatment, 11(6), 511-515; Beckman, L. J. (1994). Treatment needs of women with alcohol problems. Alcohol Health and Research World, 18(3), 206-211.
122. Wells, D. V. B. and Jackson, J. F. (1992). HIV and chemically dependent women: Recommendations for appropriate health care and treatment services. International Journal of the Addictions, 27(5), 571-585.
123. Breitbart, V., Chavkin, W., and Wise, P. H. (1994). Accessibility of drug treatment for pregnant women: A survey of programs in five cities. American Journal of Public Health, 84(10), 1658-61; Gehshan, S. and Southern Regional Project on Infant Mortality.

(1993). A step toward recovery: Improving access to substance abuse treatment for pregnant and parenting women: Results from a regional study. Washington, DC: Southern Regional Project on Infant Mortality; Barth, R. P., Pietrzak, J. and Ramler, M. (1993). Families living with drugs and HIV: Intervention and treatment strategies. New York: Guilford Press.

124. R. R. Watson (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press; Bollerud, K. (1990). A model for the treatment of trauma-related syndromes among chemically dependent inpatient women. Journal of Substance Abuse Treatment, 7(2), 83-87; Beckman, L. J. (1994). Treatment needs of women with alcohol problems. Alcohol Health and Research World, 18(3), 206-211; Katz, J. L. (1990). Eating disorders: A primer for the substance abuse specialist: 2. Theories of etiology, treatment approaches, and considerations during comorbidity with substance abuse. Journal of Substance Abuse Treatment, 7(4), 211-217; Root, M. P. P. (1989). Treatment failures: The role of sexual victimization in women's addictive behavior. American Journal of Orthopsychiatry, 59(4), 542-549; Young, E. B. (1995). The role of incest issues in relapse and recovery. In A. B. Washton (Ed.), Psychotherapy and substance abuse: A practitioners handbook (pp. 451-469). New York: Guilford Press; Brady, K. T., Killen, T., Saladin, M. E., Dansky, B. and Becker, S. (1994). Comorbid substance abuse and posttraumatic stress disorder. American Journal on Addictions, 3(2), 160-164; Fullilove, M. T., Fullilove, R. E., Smith, M., Winkler, K., Michael, C., Panzer, P. G. and Wallace, R. (1993). Violence, trauma, and post-traumatic stress disorder among women drug users. Journal of Traumatic Stress, 6(4), 533-543; Gomberg, E. S. L. (1988). Alcoholic women in treatment: The question of stigma and age. Alcohol and Alcoholism, 23(6), 507-514; O'Connor, L. E., Berry, J. W., Inaba, D., Weiss, J. and Morrison, A. (1994). Shame, guilt, and depression in men and women in recovery from addiction. Journal of Substance Abuse Treatment, 11(6), 503-510.

125. Harvey, E. M., Rawson, R. A. and Obert, J. L. (1994). History of sexual assault and the treatment of substance abuse disorders. Journal of Psychoactive Drugs, 26(4), 361-367.

126. Harrison, P. A., Hoffmann, N. G. and Edwall, G. E. (1989). Differential drug use patterns among sexually abused adolescent girls in treatment for chemical dependency. International Journal of the Addictions, 24(6), 499-514; Harrison, P. A., Hoffman, N. G. and Edwall, G. E. (1989). Sexual abuse correlates: Similarities between male and female adolescents in chemical dependency treatment. Journal of Adolescent Research, 4(3), 385-399.

127. Harrison, P. A., Hoffman, N. G. and Edwall, G. E. (1989). Sexual abuse correlates: Similarities between male and female adolescents in chemical dependency treatment. Journal of Adolescent Research, 4(3), 385-399.

128. Miller, B. A., Downs, W. R. and Gondoli, D. M. (1989). Spousal violence among alcoholic women as compared to a random household sample of women. Journal of Studies on Alcohol, 50(6), 533-540.

129. Finnegan, L. P., Hagan, T. and Kaltenbach, K. A. (1991). Scientific foundation of clinical practice: Opiate use in pregnant women. Bulletin of the New York Academy of Medicine, 67(3), 223-239; Mitchell, J. L. (1993). Pregnant, substance-using women. Rockville, MD: U.S. Department of Health and Human Services, Center for Substance Abuse Treatment; Barth, R. P., Pietrzak, J. and Ramler, M. (1993). Families living with drugs and HIV: Intervention and treatment strategies. New York: Guilford Press; Kaltenbach, K. A. and Finnegan, L. P. (1992). Studies of prenatal drug exposure and environmental research issues: The benefits of integrating within a treatment program. In M. M. Kilbey and K. Ashgar (Eds.), Methodological issues in epidemiological, prevention and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 259-270). Rockville, MD: National Institute on Drug Abuse.
130. Finnegan, L. P., Hagan, T. and Kaltenbach, K. A. (1991). Scientific foundation of clinical practice: Opiate use in pregnant women. Bulletin of the New York Academy of Medicine, 67(3), 223-239; Beckman, L. J. (1994). Treatment needs of women with alcohol problems. Alcohol Health and Research World, 18(3), 206-211.
131. R. R. Watson (Ed.). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press; Marcenko, M. O., Spence, M. and Rohweder, C. (1994). Psychosocial characteristics of pregnant women with and without a history of substance abuse. Health and Social Work, 19(1), 17-22.
132. Silverman, K., Chutuape, M. A. D., Svikis, D. S., Bigelow, G. E. and Stitzer, M. L. (1995). Drug and Alcohol Dependence, 40(2), 115-123; Finnegan, L. P., Hagan, T. and Kaltenbach, K. A. (1991). Scientific foundation of clinical practice: Opiate use in pregnant women. Bulletin of the New York Academy of Medicine, 67(3), 223-239.
133. Finnegan, L. P., Hagan, T. and Kaltenbach, K. A. (1991). Scientific foundation of clinical practice: Opiate use in pregnant women. Bulletin of the New York Academy of Medicine, 67(3), 223-239; Chavkin, W., Paone, D., Friedmann, P. and Wilets, I. (1993). Reframing the debate: Toward effective treatment for inner city drug-abusing mothers. Bulletin of the New York Academy of Medicine, 70(1), 50-68; Blume, S. B. (1991). Women, alcohol, and drugs. In N. S. Miller (Ed.), Comprehensive handbook of drug and alcohol addiction (pp. 147-177). New York: Marcel Dekker; Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: TAB Books; Mathias, R. (1994). NIDA's Perinatal-20 projects: Laying the foundation for the treatment of drug-abusing pregnant and parenting women. NIDA Notes, 9(4), 6-7; Marcenko, M. O., Spence, M. and Rohweder, C. (1994). Psychosocial characteristics of pregnant women with and without a history of substance abuse. Health and Social Work, 19(1), 17-22.
134. Comtois, K. A. and Ries, R. K. (1995). Sex differences in dually diagnosed severely mentally ill clients in dual diagnosis outpatient treatment. American Journal on Addictions, 4(3), 245-253; Burman, S. (1992). A model for women's alcohol/drug treatment. Alcoholism Treatment Quarterly, 9(2), 87-99; Luthar, S. S. and Walsh, K. G. (1995). Treatment needs of drug-addicted mothers: Integrating parenting psychotherapy interventions. Journal of

- Substance Abuse Treatment, 12(5), 341-348; Schliebner, C. T. (1994). Gender-sensitive therapy: An alternative for women in substance abuse treatment. Journal of Substance Abuse Treatment, 11(6), 511-515; Kaltenbach, K. A. and Finnegan, L. P. (1992). Studies of prenatal drug exposure and environmental research issues: The benefits of integrating within a treatment program. In M. M. Kilbey and K. Ashgar (Eds.), Methodological issues in epidemiological, prevention and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 259-270). Rockville, MD: National Institute on Drug Abuse.
135. Burman, S. (1992). A model for women's alcohol/drug treatment. Alcoholism Treatment Quarterly, 9(2), 87-99; N. van den Bergh (Ed.). (1991). Feminist perspectives on addictions. New York: Springer Publishing Company; Quinby, P. M. and Graham, A. V. (1993). Substance abuse among women. Primary Care, 20(1), 131-140.
136. Blume, S. B. (1988). Alcohol/drug dependent women: New insights into their special problems, treatment, recovery. Minneapolis, MN: Johnson Institute; Comtois, K. A. and Ries, R. K. (1995). Sex differences in dually diagnosed severely mentally ill clients in dual diagnosis outpatient treatment. American Journal on Addictions, 4(3), 245-253; Wilke, D. (1994). Women and alcoholism: How a male-as-norm bias affects research, assessment, and treatment. Health and Social Work, 19(1), 29-35.
137. Chappel, J. N. (1993). Long-term recovery from alcoholism. Psychiatric Clinics of North America, 16(1), 177-187.
138. Schmidt, L. and Weisner, C. (1995). The emergence of problem-drinking women as a special population in need of treatment. In M. Galanter (Ed.), Alcoholism and women (pp. 309-334). New York: Plenum Press; Berenson, D. (1991). Powerlessness: Liberating or enslaving? Responding to the feminist critique of the Twelve Steps. In C. Bepko (Ed.), Feminism and addiction (pp. 67-84). New York: Haworth Press; Abbott, A. A. (1994). A feminist approach to substance abuse treatment and service delivery. Social Work in Health Care, 19(3/4), 67-83.
139. Chappel, J. N. (1993). Long-term recovery from alcoholism. Psychiatric Clinics of North America, 16(1), 177-187; Blume, S. B. (1991). Women, Alcohol, and drugs. In N. S. Miller, (Ed.), Comprehensive handbook of drug and alcohol addiction (pp. 147-177). New York: Marcel Dekker.
140. Chappel, J. N. (1993). Long-term recovery from alcoholism. Psychiatric Clinics of North America, 16(1), 177-187; U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Treatment. (1994). Practical approaches in the treatment of women who abuse alcohol and other drugs. Rockville, MD: U.S. Department of Health and Human Services.
141. McCrady, B. S. and Raytek, H. (1993). Women and substance abuse: Treatment and outcome modalities. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and

- substance abuse (pp. 314-338). Norwood, NJ: Ablex Publishing; Vanicelli, M. (1984). Treatment outcome of alcoholic women: The state of the art. Bulletin of the Society of Psychologists in Addictive Behavior, 3(3), 155-162; Halliday, A. and Bush, B. (1987). Women and alcohol abuse. In H. N. Barnes, M. D. Aronson and T. L. Delbanco (Eds.), Alcoholism: A guide for the primary care physician (pp. 176-180). New York: Springer Verlag; Murray, J. B. (1989). Psychologists and alcoholic women. Psychology Reports, 64(2), 627-644.
142. Dahlgren, L. and Willander, A. (1989). Are special treatment facilities for female alcoholics needed?: A controlled 2-year follow-up study from a specialized female unit (EWA) versus a mixed male/female treatment facility. Alcoholism: Clinical and Experimental Research, 13(4), 499-504.
143. New Standards, Inc. (1993). Betty Ford Center Treatment Outcomes November 1993. St. Paul, MN: New Standards, Inc.
144. Waite-O'Brien, N., Clinical Director, Betty Ford Center at Eisenhower. (1996). Personal communication.
145. R. R. Watson (Ed). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press.
146. Blake, S. M., Klepp, K., Pechacek, T. F., Folsom, A. R., Leupker, R. V., Jacobs, D. R. and Mittelmark, M. B. (1989). Differences in smoking cessation strategies between men and women. Addictive Behaviors, 14(4), 409-418; R. R. Watson (Ed). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press.
147. Blake, S. M., Klepp, K., Pechacek, T. F., Folsom, A. R., Leupker, R. V., Jacobs, D. R. and Mittelmark, M. B. (1989). Differences in smoking cessation strategies between men and women. Addictive Behaviors, 14(4), 409-418.
148. McBride, C. M., Pirie, P. L. and Curry, S. J. (1992). Postpartum relapse to smoking: A prospective study. Health Education Research, 7(3), 381-390. Center for the Future of Children (1995). The future of children: Low birth weight. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children; Fingerhut, L. A., Kleinman, J. C. and Kendrick, J. S. (1990). Smoking before, during, and after pregnancy. American Journal of Public Health, 80(5), 541-544.
149. R. R. Watson (Ed). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press.
150. Pirie, P. L., Murray, D. M. and Luepker, R. V. (1991). Gender differences in cigarette smoking and quitting in a cohort of young adults. American Journal of Public Health, 81(3), 324-327.

151. Berman, B. A. and Gritz, E. R. (1991). Women and smoking: Current trends and issues for the 1990s. Journal of Substance Abuse, 3(2), 221-238.
152. McBride, C. M., Pirie, P. L. and Curry, S. J. (1992). Postpartum relapse to smoking: A prospective study. Health Education Research, 7(3), 381-390.
153. Berman, B. A. and Gritz, E. R. (1993). Women and smoking: Toward the year 2000. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 258-285). Norwood NJ: Ablex Publishing.
154. Hall, S. M., Ginsberg, D. and Jones, R. T. (1986). Smoking cessation and weight gain. Journal of Consulting and Clinical Psychology, 54(3), 342-346; Gritz, E. R. and St. Jeor, S. T. (1992). Task force 3: Implications with respect to intervention and prevention. Health Psychology, 11(Suppl.), 17-25; Berman, B. A. and Gritz, E. R. (1993). Women and smoking: Toward the year 2000. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 258-285). Norwood NJ: Ablex Publishing.
155. Glassman, A. H., Helzer, J. E., Covey, L. S., Cottler, L. B., Stetner, F., Tipp, J. E. and Johnson, J. (1990). Smoking, smoking cessation, and major depression. JAMA, 264(12), 1546-1549; Anda, R. F., Williamson, D. F., Escobedo, L. G., Mast, E. E., Giovino, G. A. and Remington, P. L. (1990). Depression and the dynamics of smoking. JAMA, 264(12), 1541-1545; Fisher, E. B., Lichtenstein, E. and Haire-Joshu, D. (1993). Multiple determinants of tobacco use and cessation. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 59-88). New York: Oxford University Press.
156. Anda, R. F., Williamson, D. F., Escobedo, L. G., Mast, E. E., Giovino, G. A. and Remington, P. L. (1990). Depression and the dynamics of smoking. JAMA, 264(12), 1541-1545; Fisher, E. B., Lichtenstein, E. and Haire-Joshu, D. (1993). Multiple determinants of tobacco use and cessation. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 59-88). New York: Oxford University Press; Glassman, A. H., Helzer, J. E., Covey, L. S., Cottler, L. B., Stetner, F., Tipp, J. E. and Johnson, J. (1990). Smoking, smoking cessation, and major depression. JAMA, 264(12), 1546-1549.
157. Surgeon General (1988). The health consequences of smoking: Nicotine addiction. Washington, DC: U.S. Department of Health and Human Services, Public Health Service.
158. Berman, B. A. and Gritz, E. R. (1993). Women and smoking: Toward the year 2000. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 258-285). Norwood NJ: Ablex Publishing; Fisher, E. B., Lichtenstein, E. and Haire-Joshu, D. (1993). Multiple determinants of tobacco use and cessation. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 59-88). New York: Oxford University Press; Surgeon General (1988). The health consequences of smoking: Nicotine addiction. Washington, DC: U.S. Department of Health and Human Services, Public Health Service.

159. Gritz, E. E. and St. Jeor, S. T. (1992). Task force 3: Implications with respect to intervention and prevention. Health Psychology, 11(Suppl.), 17-25; Berman, B. A. and Gritz, E. R. (1991). Women and smoking: Current trends and issues for the 1990s. Journal of Substance Abuse, 3(2), 221-238; R. R. Watson (Ed). (1994). Addictive behaviors in women. Totowa, NJ: Humana Press.
160. Marcus, B. H., Albrecht, A. E., Niaura, R. S., Taylor, E. R., Simkin, L. R., Feder, S. I., Abrams, D. B. and Thompson, P. D. (1995). Exercise enhances the maintenance of smoking cessation in women. Addictive Behaviors, 20(1), 87-92.
161. Streater, J. A., Sargent, R. G. and Ward, D. S. (1989). A study of factors associated with weight change in women who attempt smoking cessation. Addictive Behaviors, 14(5), 523-530.
162. Berman, B. A. and Gritz, E. R. (1993). Women and smoking: Toward the year 2000. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 258-285). Norwood NJ: Ablex Publishing.
163. Pirie, P. L., McBride, C. M., Hellerstedt, W., Jeffery, R. W., Hatsukami, D., Allen, S. and Lando, H. (1992). Smoking cessation in women concerned about weight. American Journal of Public Health, 82(9), 1238-1243.
164. Horger, E. O., Brown, S. B. and Condon, C. M. (1990). Cocaine in pregnancy: Confronting the problem. Journal of the South Carolina Medical Association, 86(10), 527-531.
165. Chavkin, W., Walker, N. A. and Paone, D. (1990). Drug-using families and child protection: Results of a study and implications for change. University of Pittsburgh Law Review, 54(1), 295-324; Connolly, W. B. and Marshall, A. B. (1991). Drug addiction, pregnancy, and childbirth: Legal issues for the medical and social services communities. Clinics in Perinatology, 18(1), 147-186; English, A. (1990). Prenatal drug exposure and pediatric AIDS: New issues for children's attorneys. Clearinghouse Review, 24(5), 452-459; Weber, E. M. (1992). Alcohol- and drug-dependent pregnant women: Laws and public policies that promote and inhibit research and the delivery of services. In M. M. Kilbey and K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 349-365). Rockville, MD: National Institute on Drug Abuse; Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished; Lombardo, C. (1991). Mothers' prenatal drug use: Sufficient grounds for juvenile court jurisdiction. Journal of Juvenile Law, 12, 116-119; Horger, E. O., Brown, S. B. and Condon, C. M. (1990). Cocaine in pregnancy: Confronting the problem. Journal of the South Carolina Medical Association, 86(10), 527-531.
166. Jos, P. H., Marshall, M. F. and Perlmutter, M. (1995). The Charleston policy on cocaine use during pregnancy: A cautionary tale. Journal of Law, Medicine and Ethics,

- 23(2), 120-128; Hansen, M. (1992). Courts side with moms in drug cases. ABA Journal, 78, 18; Hawk, M. A. N. (1994). How social policies make matters worse: The case of maternal substance abuse. Journal of Drug Issues, 24(3), 517-526; Chavkin, W., Breitbart, V. and Wise, P. H. (1994). Finding common ground: The necessity of an integrated agenda for women's and children's health. Journal of Law, Medicine and Ethics, 22(3), 262-269.
167. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
168. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
169. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
170. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
171. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
172. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
173. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
174. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
175. Marshall, A. B. (1995). 1994 legislative update. Families, Addiction Research and Education Update, 1.
176. Wiese, D., Daro, D. and National Center on Child Abuse Prevention Research. (1995). Current trends in child abuse reporting and fatalities: The results of the 1994 annual fifty state survey. Chicago, IL: National Committee to Prevent Child Abuse; Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General; U.S. General Accounting Office. (1995). Child welfare: Complex needs strain capacity to provide services. Washington, DC: U.S. General Accounting Office; Jost, K. (1991). Foster care crisis. CO Researcher, 1(20), 707-727.
177. U.S. General Accounting Office. (1995). Child welfare: Complex needs strain capacity to provide services. Washington, DC: U.S. General Accounting Office; U.S. General Accounting Office. (1995). Child welfare: Complex needs strain capacity to provide services. Washington, DC: U.S. General Accounting Office; Jost, K. (1991). Foster care crisis. CO Researcher, 1(20), 707-727.

178. Berrick, J. D. and Lawrence-Karski, R. (1995). Emerging issues in child welfare: A state survey of child welfare administrators identifies issues of concern. Public Welfare, 4-11.
179. U.S. General Accounting Office. (1995). Child welfare: Complex needs strain capacity to provide services. Washington, DC: U.S. General Accounting Office; Berrick, J. D. and Lawrence-Karski, R. (1995). Emerging issues in child welfare: A state survey of child welfare administrators identifies issues of concern. Public Welfare, 4-11; Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.
180. Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.
181. Task Force on Family Violence. (1993). Behind closed doors: The city's response to family violence. Report of the Task Force on Family Violence. New York: Author.
182. Garcia, S. A. (1993). Maternal drug abuse: Laws and ethics as agents of just balances and therapeutic interventions. International Journal of the Addictions, 28(13), 1311-1339.
183. Horowitz, R. (1990). Perinatal substance abuse. Children Today, 19(4), 8-12.
184. Kusserow, R. P. (June 1990). Crack babies. Washington, DC: U.S. Department of Health and Human Services, Office of the Inspector General.
185. Garcia, S. A. (1992). Perinatal drug and alcohol abuse: Rights, laws, and responsibilities. Legal Medicine, 109-164.
186. Bays, J. (1990). Substance abuse and child abuse: Impact of addiction on the child. Pediatric Clinics of North America, 37(4), 881-904; Besharov, D. J. (1989). The children of crack: Will we protect them? Public Welfare, 47(4), 6-11; Jost, K. (1991). Foster care crisis. CO Researcher, 1(20), 707-727.
187. Sexton, J. (1996, March 12). Officials seek wider powers to seize children in drug homes. New York Times, p. B1.
188. Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
189. Allen, P. M. and Sandler, M. (1993). Critical components of obstetric management of chemically dependent women. Clinical Obstetrics and Gynecology, 36(2), 347-360; Chavkin, W. (1991). Mandatory treatment for drug use during pregnancy. JAMA, 266(11), 1556-1561.

190. Luthar, S. S. and Walsh, K. G. (1995). Treatment needs of drug-addicted mothers: Integrating parenting psychotherapy interventions. Journal of Substance Abuse Treatment, 12(5), 341-348; Seiden, A. M. (1992). Measures of pregnant, drug-abusing women for treatment research. In M. M. Kilbey and K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 194-211). Rockville, MD: National Institute on Drug Abuse.
191. U.S. General Accounting Office. (1995). Child welfare: Complex needs strain capacity to provide services. Washington, DC: U.S. General Accounting Office; Jost, K. (1991). Foster care crisis. CO Researcher, 1(20), 707-727.
192. C. G. Leukefeld and F. M. Tims (Eds.). (1988). Compulsory treatment of drug abuse: Research and clinical practice. NIDA Research Monograph 86. Washington, DC: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, National Institute on Drug Abuse; Smith, L. (1992). Help seeking in alcohol-dependent females. Alcohol and Alcoholism, 27(1), 3-9. Chavkin, W. (1991). Mandatory treatment for drug use during pregnancy. JAMA, 266(11), 1556-1561.
193. Chavkin, W. (1991). Mandatory treatment for drug use during pregnancy. JAMA, 266(11), 1556-1561; Barth, R. P., Pietrzak, J. and Ramler, M. (1993). Families living with drugs and HIV: Intervention and treatment strategies. New York: Guilford Press; Moss, K. L. (1991). Forced drug or alcohol treatment for pregnant and postpartum women: Part of the solution or part of the problem? New England Journal on Criminal and Civil Confinement, 17(1), 1-16.
194. Chavkin, W. (1991). Mandatory treatment for drug use during pregnancy. JAMA, 266(11), 1556-1561.
195. Chavkin, W. (1991). Mandatory treatment for drug use during pregnancy. JAMA, 266(11), 1556-1561; Garcia, S. A. (1993). Maternal drug abuse: Laws and ethics as agents of just balances and therapeutic interventions. International Journal of the Addictions, 28(13), 1311-1339; (1987, February 1). Jail by mistake for female alcoholics. New York Times, p. A42.
196. Chavkin, W. (1991). Mandatory treatment for drug use during pregnancy. JAMA, 266(11), 1556-1561; (1987, February 1). Jail by mistake for female alcoholics. New York Times, p. A42.
197. Garcia, S. A. and Keilitz, I. (1991). Involuntary civil commitment of drug-dependent persons with special reference to pregnant women. Mental and Physical Disability Law Register, 15(4), 418-437.
198. California Elected Women's Association for Education and Research. (1994). Women

and substance abuse. Sacramento, CA: California Elected Women's Association for Education and Research.

199. Besharov, D. J. (1996, January 13). How to save children. New York Times, p. A23.
200. Chasnoff, I. J. (1992). Drug use in pregnancy: Does treatment work? Project report. Unpublished paper.
201. Carroll, K. M., Chang, G., Behr, H., Clinton, B. and Kosten, T. R. (1995). Improving treatment outcome in pregnant, methadone-maintained women: Results from a randomized clinical trial. American Journal on Addictions, 4(1), 56-81; Chang, G., Carroll, K. M., Behr, H. M. and Kosten, T. R. (1992). Improving treatment outcome in pregnant opiate-dependent women. Journal of Substance Abuse Treatment, 9(4), 327-330.
202. Chang, G., Carroll, K. M., Behr, H. M. and Kosten, T. R. (1992). Improving treatment outcome in pregnant opiate-dependent women. Journal of Substance Abuse Treatment, 9(4), 327-330.
203. Carroll, K. M., Chang, G., Behr, H., Clinton, B. and Kosten, T. R. (1995). Improving treatment outcome in pregnant, methadone-maintained women: Results from a randomized clinical trial. American Journal on Addictions, 4(1), 56-81.
204. Seiden, A. M. (1992). Measures of pregnant, drug-abusing women for treatment research. In M. M. Kilbey and K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 194-211). Rockville, MD: National Institute on Drug Abuse; Allen, P. M. and Sandler, M. (1993). Critical components of obstetric management of chemically dependent women. Clinical Obstetrics and Gynecology, 36(2), 347-360; Chavkin, W. (1991). Mandatory treatment for drug use during pregnancy. JAMA, 266(11), 1556-1561; Chavkin, W., Paone, D., Friedmann, P. and Wilets, I. (1993). Reframing the debate: Toward effective treatment for inner city drug-abusing mothers. Bulletin of the New York Academy of Medicine, 70(1), 50-68.
205. Gfroerer, J. C. and Office of Applied Studies. (1995). Preliminary estimates from the 1994 National Household Survey on Drug Abuse. Rockville, MD: Office of Applied Studies; Klein, D., Zahnd, E., Holtby, S. and Barth, R. (1994). Perinatal needs assessment executive summary. Berkeley, CA: California Health and Welfare Agency, Department of Alcohol and Drug Programs, Office of Perinatal Substance Abuse; Law, M. and Tang, J. L. (1995). An analysis of the effectiveness of interventions intended to help people to stop smoking. Archives of Internal Medicine, 155, 1933-1941; Hall, J. A., Henggeler, S. W., Felice, M. E., Reynoso, T., Williams, N. M. and Sheets, R. (1993). Adolescent substance use during pregnancy. Journal of Pediatric Psychology, 18(2), 265-271; Gilchrist, L. D., Gillmore, M. R. and Lohr, M. J. (1990). Drug use among pregnant adolescents. Journal of Consulting and Clinical Psychology, 58(4), 402-407; Colmorgen, G. H. C., Johnson, C., Zazzarino, M. A.

- and Durinzi, K. (1992). Routine urine drug screening at the first prenatal visit. American Journal of Obstetrics and Gynecology, 166(2), 588-590.
206. Bayer, R. (1994). Ethical challenges posed by zidovudine treatment to reduce vertical transmission of HIV. New England Journal of Medicine, 331(18), 1223-5; Centers for Disease Control and Prevention. (1994). Zidovudine for the prevention of HIV transmission from mother to infant. Morbidity and Mortality Weekly Report, 43(16), 285-7.
207. Chavkin, W., Paone, D., Friedmann, P. and Wilets, I. (1993). Reframing the debate: Toward effective treatment for inner city drug-abusing mothers. Bulletin of the New York Academy of Medicine, 70(1), 50-68.
208. Raskin, V. D. (1993). Psychiatric aspects of substance use disorders in childbearing populations. Psychiatric Clinics of North America, 16(1), 157-165; Quinby, P. M. and Graham, A. V. (1993). Substance abuse among women. Primary Care, 20(1), 131-140; Gomberg, E. S. L. (1987). Shame and guilt issues among women alcoholics. Alcoholism Treatment Quarterly, 4(2), 139-155.
209. American Hospital Association (1991). The role of hospitals in caring for pregnant substance-abusing women. Chicago, IL: American Hospital Association; American Medical Association Board of Trustees. (1990). Legal interventions during pregnancy: Court-ordered medical treatments and legal penalties for potentially harmful behavior by pregnant women. JAMA, 264(20), 2663-2670; Chavkin, W. (1990). Drug addiction and pregnancy: Policy crossroads. American Journal of Public Health, 80(4), 483-487.
210. U.S. General Accounting Office. (1991). ADMS Block Grant: Women's set-aside does not assure drug treatment for pregnant women. Washington, DC: U.S. General Accounting Office; U.S. General Accounting Office. (1990). Drug-exposed infants: A generation at risk. Washington, DC: U.S. General Accounting Office; Chasnoff, I. J. (1992). Drug use in pregnancy: Does treatment work? Project report. Unpublished paper; Gehshan, S. and Southern Regional Project on Infant Mortality. (1993). A step toward recovery: Improving access to substance abuse treatment for pregnant and parenting women: Results from a regional study. Washington, DC: Southern Regional Project on Infant Mortality; Weber, E. M. (1992). Alcohol- and drug-dependent pregnant women: Laws and public policies that promote and inhibit research and the delivery of services. In M. M. Kilbey, K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 349-365). Rockville, MD: National Institute on Drug Abuse; Poland, M. L., Dombrowski, M. P., Ager, J. W. and Sokol, R. J. (1993). Punishing pregnant drug users: Enhancing the flight from care. Drug and Alcohol Dependence, 31, 199-203.
211. Chavkin, W. (1990). Drug addiction and pregnancy: Policy crossroads. American Journal of Public Health, 80(4), 483-487; Kaltenbach, K. A. and Finnegan, L. P. (1992). Studies of prenatal drug exposure and environmental research issues: The benefits of integrating within a treatment program. In M. M. Kilbey and K. Ashgar (Eds.),

- Methodological issues in epidemiological, prevention and treatment research on drug-exposed women and their children. NIDA research monograph 117 (pp. 259-270). Rockville, MD: National Institute on Drug Abuse; Sokol, R. J., Ager, J. W. and Martier, S. S. (1992). Methodological issues in obtaining and managing substance abuse information from prenatal patients. In M. M. Kilbey and K. Ashgar (Eds.), Methodological issues in epidemiological, prevention and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 80-95). Rockville, MD: National Institute on Drug Abuse.
212. Chavkin, W. (1996, February). Personal communication.
213. Chavkin, W. (1996, February). Personal communication.
214. Birchfield, M., Scully, J. and Handler, A. (1995). Perinatal screening for illicit drugs: Policies in hospitals in a large metropolitan area. Journal of Perinatology, 15(3), 208-214; Weber, E. M. (1992). Alcohol- and drug-dependent pregnant women: Laws and public policies that promote and inhibit research and the delivery of services. In M. M. Kilbey and K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 349-365). Rockville, MD: National Institute on Drug Abuse.
215. Weber, E. M. (1992). Alcohol- and drug-dependent pregnant women: Laws and public policies that promote and inhibit research and the delivery of services. In M. M. Kilbey and K. Asghar (Eds.), Methodological issues in epidemiological, prevention, and treatment research on drug-exposed women and their children. NIDA Research Monograph 117 (pp. 349-365). Rockville, MD: National Institute on Drug Abuse.
216. Birchfield, M., Scully, J. and Handler, A. (1995). Perinatal screening for illicit drugs: Policies in hospitals in a large metropolitan area. Journal of Perinatology, 15(3), 208-214.
217. Horowitz, R. (1990). Perinatal substance abuse. Children Today, 19(4), 8-12; Cadwalader, Wickersham and Taft. (1995). The law's response to maternal and prenatal substance abuse, unpublished.
218. Mann, T., Battaglin, J., Cooper, S. and Mahan, C. S. (1992). Some of my patients use drugs: Pregnancy/substance abuse and the physician. Journal of the Florida Medical Association, 79(1), 41-45.
219. Mann, T., Battaglin, J., Cooper, S. and Mahan, C. S. (1992). Some of my patients use drugs: Pregnancy/substance abuse and the physician. Journal of the Florida Medical Association, 79(1), 41-45.
220. Wilson, B. L. (1991). Treatment for two. Public Welfare, 49(4), 31-36.
221. American Bar Association, Center on Children and the Law, Young Lawyers Division. (1990). Drug exposed infants and their families: Coordinating responses of the legal, medical and child protection system. Washington, DC: American Bar Association, Center on

Children and the Law, Young Lawyers Division; Kaltenbach, K. A. and Finnegan, L. P. (1992). Studies of prenatal drug exposure and environmental research issues: The benefits of integrating within a treatment program. In M. M. Kilbey and K. Ashgar (Eds.), Methodological issues in epidemiological, prevention and treatment research on drug-exposed women and their children, NIDA Research Monograph 117 (pp. 259-270). Rockville, MD: National Institute on Drug Abuse; Coles, C. D. (1992). Discussion: Measurement issues in the study of effects of substance abuse in pregnancy. In M. M. Kilbey and K. Ashgar (Eds.), Methodological issues in epidemiological, prevention and treatment research on drug-exposed women and their children, NIDA Research Monograph 117 (pp. 248-258). National Institute on Drug Abuse; Birchfield, M., Scully, J. and Handler, A. (1995). Perinatal screening for illicit drugs: Policies in hospitals in a large metropolitan area. Journal of Perinatology, 15(3), 208-214.

222. Birchfield, M., Scully, J. and Handler, A. (1995). Perinatal screening for illicit drugs: Policies in hospitals in a large metropolitan area. Journal of Perinatology, 15(3), 208-214.

223. Birchfield, M., Scully, J. and Handler, A. (1995). Perinatal screening for illicit drugs: Policies in hospitals in a large metropolitan area. Journal of Perinatology, 15(3), 208-214.

224. Wiley, K., Gibbs, B., Kahn, S., Karlman, R., Tse, R. and Perez-Woods, R. (1991). Prevalence of illicit drug use among prenatal patients and predictive validity of nurses' judgments. Journal of Perinatology, 11(4), 330-335; Chasnoff, I. J., Landress, H. J. and Barrett, M. E. (1990). The prevalence of illicit-drug or alcohol use during pregnancy and discrepancies in mandatory reporting in Pinellas County, Florida. New England Journal of Medicine, 322(17), 1202-1206.

CHAPTER VI.

REFERENCES

1. Blume, S. B. (1991). Women, alcohol and drugs. In N. S. Miller (Ed.), Comprehensive handbook of drug and alcohol addiction (pp. 147-177). New York: Marcel Dekker.
2. Center for the Future of Children. (1995). The future of children: low birth weight. Los Altos, CA: David and Lucile Packard Foundation, Center for the Future of Children.
3. American Psychological Association (1996). Research Agenda for psychosocial and behavioral factors in women's health. Washington, DC: American Psychological Association.
4. Chavkin, W. (1994). Women and clinical research. Journal of the American Medical Womens Association, 49(4), 99-100.
5. In addition, studies that have tried to measure how a woman's hormone levels across the menstrual cycle may affect her metabolism of alcohol have produced inconsistent but provocative findings. While hormone changes do appear to affect her metabolism of alcohol, it is unclear whether a given amount of alcohol is more likely to make a woman feel drunk at any a particular time in her menstrual cycle. See Tivis, L. J. and Galaver, J. S. (1994). Alcohol, hormones and health in postmenopausal women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 185-188). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Hill, S. Y. (1984). Physiological effects of alcohol in women. In Women and Alcohol: Health-related issues, NIAAA Research Monograph 16 (pp. 199-214). Rockville, MD: National Institute on alcohol Abuse and Alcoholism; Lammers, S. M. M., Mainzer, D. E. H. and Breteler, M. H. M. (1995). Do alcohol pharmacokinetics in women vary due to the menstrual cycle? Addiction, 90(1), 23-30; Brot, M. D., Britton, K. T. and Koob, G. F. (1995). The estrous cycle, neurosteroids and the antianxiety effects of alcohol. In W. A. Hunt and S. Zakhari (Eds.), Stress, gender and alcohol-seeking behavior (pp. 345-364). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Sutker, P. B., Goist, K. C. and King, A. R. (1987). Acute alcohol intoxication in women: Relationship to dose and menstrual cycle phase. Alcoholism: Clinical and Experimental Research, 11(1), 74-79; Brick, J., Nathan, P. E., Westrick, E., Frankenstein, W. and Shapiro, A. (1980). The effects of menstrual cycle on blood alcohol and behavior. Journal of Studies on Alcohol, 47(6), 472-477; Roman, P. M. (1988). Biological features of women's alcohol use: A review. Public Health Reports, 103(6), 628-637.
6. Wilsnack, S. C. and Wilsnack, R. W. (1995). Drinking and problem drinking in U. S. women. In M. Galanter (Ed.), Alcoholism and Women (pp. 29-60). New York: Plenum Press.

7. Risch, H. A., Howe, G. R., Jain, M., Burch, J. D., Holowaty, E. J. and Miller, A. B. (1993). Are female smokers at higher risk of lung cancer than male smokers? American Journal of Epidemiology, 138(5), 281-293.
8. Wilsnack, S. C. and Wilsnack, R. W. (1995). Drinking and problem drinking in U. S. women. In M. Galanter (Ed.), Alcoholism and Women (pp. 29-60). New York: Plenum Press; Gomberg, E. S. L. (1994). Risk factors for drinking over a woman's life span. Alcohol Health and Research World, 18(3), 220-227; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.
9. Wilsnack, R. W. and Wilsnack, S. C. (1992). Women, work and alcohol: Failures of simple theories. Alcoholism: Clinical and Experimental Research, 16(2), 172-179.
10. Johnston, L. D., O'Malley, P. M. and Bachman, J. G. (1993). National survey results on drug use from the Monitoring the Future study, 1975-1992, volume II. Rockville, MD: National Institute on Drug Abuse.
11. R. J. Battjes, Z. Sloboda and W. C. Grace (Eds.) (1994). The context of HIV risk among drug users and their sexual partners. NIDA Research Monograph 143. Rockville, MD: National Institute on Drug Abuse.
12. Gardner, P. and Hudson, B. L. (1996). Advance report of final mortality statistics, 1993. Monthly Vital Statistics Report, 44(7), 1-84; Tivis, L. J. and Galaver, J. S. (1994). Alcohol, hormones and health in postmenopausal women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 185-188). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Barry, P. P. (1986). Gender as a factor in treating the elderly. In B. A. Ray and M. C. Braude (Eds.), Women and drugs: A new era for research: NIDA Research Monograph 65 (pp. 65-69). Rockville, MD: National Institute on Drug Abuse.
13. Heath, D. B. (1991). Women and alcohol: Cross-cultural perspectives. Journal of Substance Abuse, 3(2), 175-185; Braude, M. C. (1986). Drugs and drug interactions in elderly women. In B. A. Ray and M. C. Braude (Eds.), Women and drugs: A new era for research. NIDA Research Monograph 65 (pp. 58-64). Rockville, MD: National Institute on Drug Abuse; Barry, P. P. (1986). Gender as a factor in treating the elderly. In B. A. Ray and M. C. Braude (Eds.), Women and drugs: A new era for research: NIDA Research Monograph 65 (pp. 65-69). Rockville, MD: National Institute on Drug Abuse.
14. Gomberg, E. S. L. (1995). Older women and alcohol use and abuse. In M. Galanter (Ed.), Alcoholism and Women, (pp. 61-79). New York: Plenum Press.
15. Surgeon General. (1990). The health benefits of smoking cessation. Rockville, MD: Office on Smoking and Health; Li, C. Q., Windsor, R. A., Perkins, L., Goldenberg, R. L.

- and Lowe, J. B. (1993). The impact on infant birth weight and gestational age of cotinine-validated smoking reduction during pregnancy. JAMA, 269(12), 1519-1524.
16. Finnegan, L. P., Kaltenbach, K., Kandall, S. R., Lester, B., Mayes, L. C. and Paltrow, L. M. (1995). The conflicts for parenting drug dependent women: What does the research show us? Unpublished paper; Brooks-Gunn, J., McCarton, C. and Hawley, T. (1994). Effect of in utero drug exposure on children's development. Archives of Pediatrics and Adolescent Medicine, 148(1), 33-39; American Public Health Association. (1991). Policy statement 9020: Illicit drug use by pregnant women. American Journal of Public Health, 81(2), 253-254; Streissguth, A. P. (1991). The 1990 Betty Ford lecture: What every community should know about drinking during pregnancy and the lifelong consequences for society. Substance Abuse, 12(3), 114-127; Gomberg, E. S. L. (1986). Women: Alcohol and other drugs. In B. Segal (Ed.) Perspectives on drug use in the United States (pp. 75-109). New York: Haworth Press.
17. Mathias, R. (1993). What does the future hold for drug-exposed babies? NIDA Notes, 8(3), 1, 4-5, 7; Finnegan, L. P., Kaltenbach, K., Kandall, S. R., Lester, B., Mayes, L. C. and Paltrow, L. M. (1995). The conflicts for parenting drug dependent women: What does the research show us? Unpublished paper.
18. Mathias, R. (1993). What does the future hold for drug-exposed babies? NIDA Notes, 8(3), 1, 4-5, 7.
19. Streissguth, A. P. (1991). The 1990 Betty Ford lecture: What every community should know about drinking during pregnancy and the lifelong consequences for society. Substance Abuse, 12(3), 114-127; Gomberg, E. S. L. (1986). Women: Alcohol and other drugs. In B. Segal (Ed.) Perspectives on drug use in the United States (pp. 75-109). New York: Haworth Press.
20. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; Lex, B. W. (1995). Alcohol and other psychoactive substance dependence on women and men. In M. V. Seeman (Ed.), Gender and psychopathology (pp. 311-358). Washington, DC: American Psychiatric Press; Gomberg, E. S. L. (1989). Alcoholism in women: Use of other drugs. Alcoholism: Clinical and Experimental Research, 13, 338.
21. Gomberg, E. S. L. (1989). Alcoholism in women: Use of other drugs. Alcoholism: Clinical and Experimental Research, 13, 338.
22. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association.

23. American Psychological Association (1996). Research Agenda for psychosocial and behavioral factors in women's health. Washington, DC: American Psychological Association; Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A.L. Stanton and S.J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association.
24. Davis, E., Fennoy, I., Laraque, D., Kanem, N., Brown, G. and Mitchell, J. (1992). Autism and developmental abnormalities in children with perinatal cocaine exposure. Journal of the National Medical Association, 84(4), 315-319.
25. Helzer, J. E. and Pryzbeck, T. R. (1994). Psychiatric disorders in the addicted patient: The co-occurrence of alcoholism with other psychiatric disorders. In N. S. Miller (Ed.), Principles of addiction medicine (pp. 1-5). Chevy Chase, MD: American Society of Addiction Medicine; Anthenelli, R. M. and Schuckit, M. A. (1994). Psychiatric disorders in the addicted patient: Affective and anxiety disorders. In N. S. Miller, (Ed.), Principles of addiction medicine (pp. 1-7). Chevy Chase, MD: American Society of Addiction Medicine; Drake, R. E., Alterman, A. I. and Rosenberg, S. R. (1993). Detection of substance use disorders in severely mentally ill patients. Community Mental Health Journal, 29(2), 175-192.
26. American Psychological Association (1996). Research agenda for psychosocial and behavioral factors in women's health. Washington, DC: American Psychological Association; Wilsnack, S. C. and Wilsnack, R. W. (1995). Drinking and problem drinking in U. S. women. In M. Galanter (Ed.), Alcoholism and Women (pp. 29-60). New York: Plenum Press; Heath, D. B. (1991). Women and alcohol: Cross-cultural perspectives. Journal of Substance Abuse, 3(2), 175-185; Rebach, H. (1992). Alcohol and drug use among American minorities. In J. E. Tremble, C. S. Balek and S. J. Niemcryn (Eds.), Ethnic and multicultural drug abuse (pp. 23-57). Binghamton, NY: Haworth.
27. Caetano, R. C. (1994). Drinking and alcohol-related problems among minority women. In D. M. Welsh (Ed.), Alcohol Health and Research World: Women and Alcohol, 18(3), (pp. 233-241). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press.
28. Royce, J. M., Hymowitz, N., Corbett, K., Hartwell, T. D. and Orlandi, M. A. (1993). Smoking cessation factors among African Americans and whites. American Journal of Public Health, 83(2), 220-226; Solomon, L. J. and Flynn, B. S. (1993). Women who smoke. In C. T. Orleans and J. Slade (Eds.), Nicotine addiction: Principles and management (pp. 339-349). New York: Oxford University Press.
29. Royce, J. M., Hymowitz, N., Corbett, K., Hartwell, T. D. and Orlandi, M. A. (1993). Smoking cessation factors among African Americans and whites. American Journal of Public Health, 83(2), 220-226.

30. Weibel-Orlando, J. (1984). Women and alcohol: Special populations and cross-cultural variations. In Women and alcohol: Health-related issues: Proceedings of a conference, May 23-25, 1984; NIAAA Research Monograph 16 (pp. 161-187). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; Leland, J. (1984). Alcohol use and abuse in ethnic minority women. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and intervention (pp. 66-96). New York: Guilford Press; Ja, D. Y. and Aoki, B. (1993). Substance abuse treatment: Cultural barriers in the Asian-American community. Journal of Psychoactive Drugs, 25(1), 61-79.
31. Weibel-Orlando, J. (1984). Women and alcohol: Special populations and cross-cultural variations. In Women and alcohol: Health-related issues: Proceedings of a conference, May 23-25, 1984; NIAAA Research Monograph 16 (pp. 161-187). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism.
32. Weibel-Orlando, J. (1984). Women and alcohol: Special populations and cross-cultural variations. In Women and alcohol: Health-related issues: Proceedings of a conference, May 23-25, 1984; NIAAA Research Monograph 16 (pp. 161-187). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism.
33. American Psychological Association (1996). Research agenda for psychosocial and behavioral factors in women's health. Washington, DC: American Psychological Association; McCrady, B. S. and Raytek, H. (1993). Women and substance abuse: Treatment and outcome modalities. In E. S. L. Gombert and T. D. Nirenberg (Eds.), Women and substance abuse (pp. 314-338). Norwood, NJ: Ablex Publishing; Wilke, D. (1994). Women and alcoholism: How a male-as-norm bias affects research assessment and treatment. Health and Social Work, 19(1), 29-35; D. J. Pittman and H. R. White (Eds.). (1991). Society, culture and drinking patterns reexamined. New Brunswick, NJ: Alcohol Research Documentation, Inc., Rutgers Center of Alcohol Studies.
34. Mathias, R. (1994). NIDA's Perinatal-20 projects: Laying the foundation for the treatment of drug-abusing pregnant and parenting women. NIDA Notes, 9(4), 6-7.
35. Friedman, A. S., Granick, S., Bransfield, S. and Kreishner, C. (1995). Gender differences in early life risk factors for substance use/abuse: A study of an African-American sample. American Journal of Drug and Alcohol Abuse, 21(4), 511-531; Kendler, K. S., Neale, M. C., Kessler, R. C., Heath, A. C., Corey, L. A. and Eaves, L. J. (1996). Childhood parental loss and alcoholism in women: A causal analysis using a twin-family design. Psychological Medicine, 26(1), 79-95.
36. Wilsnack, S. C. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health: Progress and challenges in research and application. Washington, DC: American Psychological Press; Gombert, E. S. L. (1994). Risk factors for drinking over a woman's life span. Alcohol Health and Research World, 18(3), 220-227; Windle, M. and Barnes, G. M. (1988). Similarities and differences in correlates of alcohol consumption and problem behaviors among male and female

- adolescents. International Journal of Addictions, 23(7), 707-728; Marlatt, G. A. (1984). Sex differences in psychosocial alcohol research: Implications for prevention. In Women and alcohol: Health related issues: Proceedings of a conference May 23-25, 1984, NIDA Research Monograph 16 (pp. 260-271.). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism.
37. American Psychological Association (1996). Research agenda for psychosocial and behavioral factors in women's health. Washington, DC: American Psychological Association.
38. Brunswick, A. F. and Banaszak-Holl, J. (1996). HIV risk behavior and the Health Belief Model: An empirical test in an African American community sample. Journal of Community Psychology 24, 44-65.
39. Center on Addiction and Substance Abuse at Columbia University. (1994). Cigarettes, alcohol, marijuana: Gateways to illicit drug use. New York: CASA. However, as mentioned in Chapter 2, there are some indications that more black women than white women begin smoking after age 21.
40. Heath, D. B. (1991). Women and alcohol: Cross-cultural perspectives. Journal of Substance Abuse, 3(2), 175-18; Fillmore, K. M. (1984). "When angels fall": Women's drinking as cultural preoccupation and as reality. In S. C. Wilsnack and L. J. Beckman (Eds.), Alcohol problems in women: Antecedents, consequences and interventions (pp. 7-36). New York: Guilford Press; Penniman, L. J. and Agnew, J. (1989). Women, work and alcohol. Occupational Medicine, 4(2), 263-273.
41. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association.
42. Blume, S. B. (1985). Women and alcohol. In T. E. Bratter and G. G. Forrest (Eds.) Alcoholism and substance abuse: Strategies for clinical intervention (pp. 623-638). New York: The Free Press.
43. Council on Scientific Affairs, American Medical Association. (1996). Alcoholism in the elderly. JAMA, 275(10), 797-801.
44. Hardesty, L. and Greif, G. L. (1994). Common themes in a group for female IV drug users who are HIV positive. Journal of Psychoactive Drugs, 26(3), 289-293.
45. Blume, S. B. (1990). Chemical dependency in women: Important issues. American Journal of Drug and Alcohol Abuse, 16(3-4), 297-307.
46. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association.

47. Donovan, J. E., Jessor, R. and Jessor, L. Problem drinking in adolescence and young adulthood: A follow-up study. Journal of Studies on Alcohol, 44(1), 109-137; 1993 National Household Survey on Drug Abuse.
48. Blume, S. B. (1991). Women, alcohol and drugs. In N. S. Miller (Ed.), Comprehensive handbook of drugs and alcohol addiction (pp. 147-177). New York: Marcel Dekker.
49. Blume, S. B. (1991). Women, alcohol and drugs. In N. S. Miller (Ed.), Comprehensive handbook of drugs and alcohol addiction (pp. 147-177). New York: Marcel Dekker.
50. Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: Tab Books.
51. Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: Tab Books.
52. Gomberg, E. S. L. and Nirenberg, T. D. (1993). Antecedents and consequences. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp.118-142). Norwood, NJ: Ablex Publishing.
53. Gomberg, E. S. L. and Nirenberg, T. D. (1993). Antecedents and consequences. In E. S. L. Gomberg and T. D. Nirenberg (Eds.), Women and substance abuse (pp.118-142). Norwood, NJ: Ablex Publishing.
54. Wilsnack, S. (1995). Alcohol use and alcohol problems in women. In A. L. Stanton and S. J. Gallant (Eds.), The psychology of women's health (pp. 381-443). Washington, DC: American Psychological Association; Sandmaier, M. (1992). The invisible alcoholics: Women and alcohol. Blue Ridge Summit, PA: Tab Books.
55. Blume, S. B. (1991). Sexuality and stigma: The alcoholic woman. Alcohol Health and Research World, 15(2), 139-146.
56. Blume, S. B. (1992). Alcohol and other drug problems in women. In J. H. Lowinson, P. Ruiz, R. B. Millman and J. G. Langrod (Eds.), Substance abuse: A comprehensive textbook (pp. 794-807). Baltimore, Hong Kong, London, Munich, Philadelphia, Sydney, Tokyo: Williams and Wilkins.



U.S. Department of Education
Office of Educational Research and Improvement (OERI)
Educational Resources Information Center (ERIC)



REPRODUCTION RELEASE

(Specific Document)

I. DOCUMENT IDENTIFICATION:

Title: SUBSTANCE ABUSE and THE AMERICAN WOMAN	
Author(s):	
Corporate Source: The NATIONAL CENTER ON ADDICTION + SUBSTANCE ABUSE	Publication Date: JUNE 1996

II. REPRODUCTION RELEASE:

AT COLUMBIA UNIVERSITY

In order to disseminate as widely as possible timely and significant materials of interest to the educational community, documents announced in the monthly abstract journal of the ERIC system, *Resources in Education* (RIE), are usually made available to users in microfiche, reproduced paper copy, and electronic/optical media, and sold through the ERIC Document Reproduction Service (EDRS) or other ERIC vendors. Credit is given to the source of each document, and, if reproduction release is granted, one of the following notices is affixed to the document.

If permission is granted to reproduce and disseminate the identified document, please CHECK ONE of the following two options and sign at the bottom of the page.



Check here

For Level 1 Release:
Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical) and paper copy.

The sample sticker shown below will be affixed to all Level 1 documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY

_____ *Sample* _____

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Level 1

The sample sticker shown below will be affixed to all Level 2 documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN OTHER THAN PAPER COPY HAS BEEN GRANTED BY

_____ *Sample* _____

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Level 2



Check here

For Level 2 Release:
Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical), but *not* in paper copy.

Documents will be processed as indicated provided reproduction quality permits. If permission to reproduce is granted, but neither box is checked, documents will be processed at Level 1.

"I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce and disseminate this document as indicated above. Reproduction from the ERIC microfiche or electronic/optical media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete inquiries."

Sign here → please

Signature: <i>David G Man</i>	Printed Name/Position/Title: DAVID MAN LIBRARIAN	
Organization/Address: <i>The National Center on Addiction + Substance Abuse at Columbia University 152 West 57th Street New York, NY 10019-3310</i>	Telephone: 212 841 5255	FAX: 212-841-5200
	E-Mail Address:	Date: 9/9/1996.



III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):

If permission to reproduce is not granted to ERIC, or, if you wish ERIC to cite the availability of the document from another source, please provide the following information regarding the availability of the document. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents that cannot be made available through EDRS.)

Publisher/Distributor:
Address:
Price:

IV. REFERRAL OF ERIC TO COPYRIGHT/REPRODUCTION RIGHTS HOLDER:

If the right to grant reproduction release is held by someone other than the addressee, please provide the appropriate name and address:

Name:
Address:

V. WHERE TO SEND THIS FORM:

Send this form to the following ERIC Clearinghouse:

However, if solicited by the ERIC Facility, or if making an unsolicited contribution to ERIC, return this form (and the document being contributed) to:

ERIC Processing and Reference Facility

1100 West Street, 2d Floor
Laurel, Maryland 20707-3598

Telephone: 301-497-4080

Toll Free: 800-799-3742

FAX: 301-953-0263

e-mail: ericfac@inet.ed.gov

WWW: <http://ericfac.piccard.csc.com>