

DOCUMENT RESUME

ED 403 220

SP 037 074

AUTHOR Nichols, Sandra; Wilson, Janet  
 TITLE Healthy Students, Healthy Schools, Healthy Missouri. Recommendations for Improving School Health Education and Reducing Health Risks for Missouri Students.  
 INSTITUTION Missouri State Dept. of Elementary and Secondary Education, Jefferson City.  
 SPONS AGENCY Centers for Disease Control (DHHS), Atlanta, GA. Div. of Adolescent and School Health.  
 PUB DATE 96  
 CONTRACT U87/CCU709009-04  
 NOTE 19p.  
 PUB TYPE Reports - Research/Technical (143)

EDRS PRICE MF01/PC01 Plus Postage.  
 DESCRIPTORS \*Health Education; \*Health Promotion; High Schools; High School Students; \*Inservice Teacher Education; Profiles; \*State Surveys; Student Behavior; \*Student Characteristics; Teacher Qualifications  
 IDENTIFIERS \*Health Behavior; \*Missouri


ABSTRACT

Findings from the 1994 Missouri School Health Education Profile and the 1995 Missouri Youth Risk Behavior Survey, and the executive summary and key findings of the 1996 School Health Education Profile are here combined. The 1994 profile is based on questionnaires completed by 252 principals and 244 teachers. The 1995 survey reports on information from 4,900 students, and the 1996 profile contains data from 249 principals and 250 teachers. Key findings from the three studies are reported in five categories: influences on health behaviors; support for school health education; amount of classroom instruction; instructional content; and teacher preparation. In each category the report indicates very briefly what research says, status in Missouri, and implications. Among the recommendations for action are the following: (1) increase the involvement of parents, peers, and members of the community; (2) continue providing support for teachers to participate in professional development opportunities; (3) improve the content of school health instruction by using curricula that are developmentally appropriate, skills based, and culturally sensitive; and (4) increase the amount of health education available for grades 11 and 12 through appropriately integrated instruction and reinforcement activities, such as peer education. The 1996 study found improvements in areas recommended in the earlier report, but a decrease in the percentage of secondary schools requiring comprehensive health education and in the percentage of schools using trained peer educators. (MAH)

\*\*\*\*\*  
 \* Reproductions supplied by EDRS are the best that can be made \*  
 \* from the original document. \*  
 \*\*\*\*\*

*Recommendations for*  
**Improving School Health Education**  
*and Reducing Health Risks for*  
**Missouri Students**

ED 403 220



► *Findings from the*  
**1994 Missouri School Health**  
**Education Profile and**  
**1995 Missouri Youth Risk**  
**Behavior Survey**

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.

- Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE AND  
DISSEMINATE THIS MATERIAL  
HAS BEEN GRANTED BY

*R. E. Bartman*

TO THE EDUCATIONAL RESOURCES  
INFORMATION CENTER (ERIC)

**August 1996**  
**Missouri Department of**  
**Elementary and Secondary Education**  
**Robert E. Bartman, Commissioner of Education**

SP037074

### **ACKNOWLEDGMENTS**

This report was prepared by:

Sandra Nichols

Health and Physical Education Consultant

Janet Wilson

Supervisor, HIV/AIDS Prevention Education

The Missouri Department of Elementary and Secondary Education would like to extend sincere appreciation to Bill Datema, HIV/AIDS Prevention Education Supervisor from 1992 through 1995, who was responsible for administering the surveys. Because of his efforts and the cooperation of numerous administrators, teachers and students who completed the surveys, sufficient data were collected to assure that the results are representative of schools and students in Missouri.

The Department also acknowledges the Centers for Disease Control and Prevention's Division of Adolescent and School Health and WESTAT, Inc. for the expertise and support they provided with data collection, analysis and reporting. A special thanks is extended to Nancy Speiker and Tracey Searles for their technical assistance.

This document was printed with funds from a cooperative agreement between the Missouri Department of Elementary and Secondary Education and the Division of Adolescent and School Health, U.S. Centers for Disease Control and Prevention (No. U87/CCU709009-04.)

# Foreword

It has become obvious that educators cannot ignore the impact of new health risks on students. Today, many teachers and schools must deal regularly with problems such as teen pregnancy, drug and alcohol abuse, poor nutrition, child abuse, and emotional disturbances. When students bring such problems to school with them, educators must respond to both the personal and educational consequences.

There is widespread agreement that better education and prevention are the keys to reducing health risks for young people. We hope this report provides a useful snapshot of health education programs in Missouri's public schools and—more importantly—suggests steps we can take to reduce health risks for all young people in Missouri.

Robert E. Bartman  
Commissioner of Education

# About this Report

- *Summaries of the 1994 School Health Education Profile and the 1995 Youth Risk Behavior Survey*
- *Implications*
- *Recommendations*

Since 1988, the Missouri Department of Elementary and Secondary Education has maintained a cooperative agreement with the U.S. Centers for Disease Control and Prevention (CDC) for providing assistance to schools in improving comprehensive health education programs, including HIV-prevention education. Two activities of the agreement are to periodically collect information about the status of secondary school health and HIV-prevention education programs and to assess the degree to which high school students are engaging in various health risk behaviors.

The School Health Education Profile and the Youth Risk Behavior Survey were designed and validated by the CDC in collaboration with state and local education personnel. The School Health Education Profile surveys randomly selected secondary principals and health education teachers. Students in the 9th through 12th grades from randomly selected high



schools completed the Youth Risk Behavior Survey. Participation in both surveys was voluntary and anonymous. Responses were confidential; no school, principal, teacher or student was identified when results were compiled. Sufficient data were collected to assure that the results are representative of schools and students in Missouri.

Not all of the data from the two surveys are being reported. Summaries of the 1994 School Health Education Profile and the 1995 Youth Risk Behavior Survey, implications of the key findings from the surveys and recommendations for action are also included in this report.

The intent of the report is to generate thoughtful consideration of how secondary school health and HIV-prevention education can be improved to ensure that all students have the opportunity to learn how to live healthfully.

# Introduction

*“Too many students are engaging in behaviors which put them at risk of immediate or future health problems—or death.”*

**M**any students in Missouri secondary schools are engaging in health-enhancing behaviors such as regular vigorous physical activity and eating a diet that includes fruits and vegetables. Unfortunately, too many students also are engaging in behaviors which put them at risk of immediate and future health problems or death.

Of concern for immediate health problems are behaviors which result in suicide, physical injury or death due to violence and car crashes, unintended pregnancy, infection with HIV (the AIDS virus) and infection with other sexually transmitted diseases (STDs). Some students also are establishing behaviors such as regular alcohol and tobacco use, eating foods high in fat, and not participating in regular physical activity which puts them at risk for cardiovascular disease later in life. The use of alcohol and other drugs also contributes to other health risk behaviors such as violence, drinking and driving and early and unprotected sexual activity.

The goal of school health education is to help students acquire the knowledge and skills needed to be healthy,

active and productive throughout life. Schools, however, cannot be solely responsible for educating students about their health. Schools, parents, peers, churches and community organizations must work together to ensure students have multiple opportunities to learn how to live safely and healthfully.

Research has shown that programs with demonstrated effectiveness in improving student health knowledge, skills and behaviors appear to have several elements in common:

- Influences on health behaviors such as peer interaction and media messages are addressed
- Support for school health education is provided
- An adequate amount of classroom instruction is provided
- The content of classroom instruction is comprehensive
- Teacher preparation is thorough and effective

## THIS REPORT ADDRESSES:

- What the research says about effective programs
- The status of Missouri secondary schools, as indicated by the findings of the School Health Education Profile and the Youth Risk Behavior Survey
- Implications

# How are we doing?

## *Influences on health behaviors*

### WHAT THE RESEARCH SAYS

Effective programs involve students in decisions about programs and in presenting positive messages to their peers. Activities are included to educate about misperceptions of peer social norms and to provide reinforcement that not all students engage in unhealthy behaviors. Effective programs also help students realistically recognize their vulnerability to health risk behaviors. Effective programs challenge media messages that promote unhealthy behaviors and replace them with positive messages.

### STATUS IN MISSOURI

One-half of the secondary schools did not use trained peer educators to help teach about health. The most common method of involving students in health education activities, in addition to classroom instruction, was through school newspaper articles. Forty-one percent of lead health education teachers taught about perceptions of vulnerability and social norms related to HIV/AIDS risk behaviors. No data were collected about how schools addressed media influences.

### IMPLICATIONS

Because some schools were not involving students in peer education or other activities to reinforce health instruction, opportunities may have been missed for students to assume leadership and take responsibility for educating themselves and others. Also, teachers need to help students change inaccurate perceptions and to understand that specific behaviors make them vulnerable to health risks.

## *Support for school health education*

### WHAT THE RESEARCH SAYS

Support is provided by school administrators, parents, community health departments, churches and other youth-serving organizations. Programs that have someone designated to coordinate efforts within the school and with the community are more effective. Additionally, effective programs are supported by school policies which reinforce classroom instruction.

### STATUS IN MISSOURI

Most secondary principals reported that inservice training and professional development for teachers was supported by the school or district. Almost one-third reported that no one was designated to coordinate health education in the school. The majority of principals reported that parents had provided either positive or no feedback about the school health education program. The majority of schools did not have an active advisory council. Of those that did, almost two-thirds did not have parents represented, and fewer had community organizations represented. Almost a third of the lead health education teachers reported that they did not involve parents in health instruction.

### IMPLICATIONS

Although the vast majority of schools and districts provided support for teachers to participate in professional development, about one-fourth of the teachers had not attended an inservice program during the previous two years. Health education is a dynamic field requiring constant updating and refinement. To be effective, teachers need to participate in professional development regularly, and someone in the school needs to coordinate efforts to keep the curriculum current and aligned with other grade levels.

Community members, especially parents, did not appear to be very involved in health education programs. What is taught in school must be reinforced outside the classroom if healthy behaviors are to be promoted and risky behaviors prevented.

# *Amount of classroom instruction*

## **WHAT THE RESEARCH SAYS**

Studies show that students' health knowledge can be increased with only a few hours of instruction. However, to achieve improvements in self-reported behaviors, more time is necessary (approximately 40 to 50 hours per year in several consecutive school years).

## **STATUS IN MISSOURI**

The majority of Missouri secondary schools were providing more than one-half year of health education which should result in more than 40 hours of instruction. However, only about 10 percent of senior high schools were scheduling health education for 11th and 12th grade students.

## **IMPLICATIONS**

Secondary schools may be providing an adequate amount of instruction for students, except for those in the 11th and 12th grades. Because of the percentage of students in the 11th and 12th grade who reported engaging in some high-risk behaviors, lack of adequate instruction in these grades is a serious concern.

# *Content of classroom instruction*

## **WHAT THE RESEARCH SAYS**

Providing only factual information about a variety of health topics is not enough to influence student behavior. Effective programs provide repeated opportunities for students to learn, at different developmental stages, health-enhancing skills such as communication (e.g., refusal, resistance, conflict resolution), decision-making and problem-solving, goal setting and stress management. The content of effective programs also is sensitive to cultures and community values.

## **STATUS IN MISSOURI**

Secondary students were receiving a great deal of health information but fewer opportunities to learn skills. This conclusion is evidenced by the high percentage of teachers who taught knowledge in a variety of health topics, while a much smaller percentage also taught skills.

## **IMPLICATIONS**

Some high school students may not possess the skills needed to help them make healthy choices. Skills-based instruction, along with factual and necessary information, should help students be better prepared to make responsible decisions about their health.

# *Teacher preparation*

## **WHAT THE RESEARCH SAYS**

Teachers receiving adequate preparation through preservice and inservice programs are more likely to be motivated to teach health education well and to fully implement a comprehensive curriculum.

## **STATUS IN MISSOURI**

The majority of lead health education teachers were experienced physical education teachers who had appropriate credentials to teach health. While most had attended inservice on various health topics during the previous two years, many more expressed interest in attending inservice training, particularly on preventing violence, suicide, HIV infection and pregnancy, and alcohol, tobacco and other drug use. However, one-fourth of the teachers had not attended an inservice during the previous two years.

## **IMPLICATIONS**

The teachers' interest in receiving training related to critical health issues suggested a recognition that students were engaging in health risk behaviors. Their concerns were reinforced by data from the Youth Risk Behavior Survey. Universities, the state department of education, state and local health departments and community-based organizations must collaborate to meet the need for providing teacher inservice statewide.



# Recommendations for Action

Based on the findings of the two surveys and what research says about effective programs, the following recommendations should be considered:

1. Increase the involvement of parents, peers and members of the community by activating a comprehensive school health advisory committee, which can assist in identifying ways to improve the health education program.
2. Continue providing support for teachers to participate in professional development opportunities and encourage them to participate regularly to update and refine their skills.
3. Improve the content of school health instruction by using curricula that are comprehensive, skills-based, developmentally appropriate and culturally sensitive. Lack of skills-based instruction may be due, in part, to schools relying on textbooks which traditionally focus on dissemination of information. When possible, schools should rely less on textbooks and more on curricula that have been shown to be effective in affecting students' knowledge, skills and behaviors. When textbooks are used, skills activities should be incorporated with the instruction.
4. Increase the amount of health education available for 11th and 12th graders through appropriately integrated instruction and reinforcement activities such as peer education.
5. Use the "Healthy Active Living" Curriculum Framework to incorporate the "Show-Me Standards" into the district's curriculum and to assist with developing performance assessments.

6. Review the "Show Me Goals 2000 State Plan" for suggested initiatives to improve:

- Curriculum and Assessment
- Professional Development
- Health and Human Services Programs
- Parent and Community Involvement

Note: the "Show-Me Standards," Goals 200 State Plan, and the Healthy Active Living Curriculum Framework are published by the Missouri Department of Elementary and Secondary Education, Division of Instruction, P.O. Box 480, Jefferson City, MO 65102

## REFERENCES

- Connell, D., Turner, R., & Mason, E. "Summary of Findings of the School Health Education Evaluation: Health Promotion Effectiveness, Implementation, and Costs." *Journal of School Health*, 1985; 55 (8): 316-321.
- Kirby, D., Short, L., Collins, J., Rugg, D., Kolbe, L., Howard, M., Miller, B., Sonenstein, F., & Zabin, L. "School-Based Programs to Reduce Sexual Risk Behaviors: A Review of Effectiveness." *Public Health Reports*, 1994; 109 (3): 339-359.
- Marx, E., & Northrop D. *Educating for Health: A Guide to Implementing a Comprehensive Approach to School Health Education*. Newton, MA: Education Development, Inc. 1995.
- Cortese, P. Middleton, K., editors *The Comprehensive School Health Challenge: Promoting Health Through Education*, Volumes One and Two. Santa Cruz, CA: ETR, Associates, 1994.

# 1994 Missouri School Health Education Profile


## *Executive summary*

The 1994 School Health Education Profile Survey was conducted as a requirement for a cooperative agreement between the Missouri Department of Elementary and Secondary Education and the federal Centers for Disease Control and Prevention (CDC). The cooperative agreement program was titled "State and Local Comprehensive School Health Programs to Prevent Important Health Problems and Improve Educational Outcomes."

In the spring of 1994, questionnaires developed by the CDC Division of Adolescent and School Health were sent to the principal and the lead health education teacher in 345 secondary schools in the state. Usable

questionnaires were received from 252 principals and 244 teachers. The principals and lead health education teachers who responded are representative of secondary school principals and lead health education teachers in Missouri.

Survey results were compiled in the following categories: (1) overall results for all schools, (2) senior high school results for schools comprised primarily of grades 9-12, (3) middle school results for schools comprised primarily of grades 6-8, and (4) junior/senior high schools results for schools comprised primarily of grades 7-12. The results may be used to develop policies and improve programs for school health education, including HIV/AIDS prevention education.



# Key findings

## INFLUENCES ON HEALTH BEHAVIOR

- 50 percent of all schools did not use trained peer educators.
- 83 percent of all schools offered guest presentations and assemblies in addition to classroom instruction. 37 percent offered intramural fitness activities, and 31 percent offered school newspaper articles.

## SUPPORT FOR SCHOOL HEALTH EDUCATION

- Only 11 percent of all principals reported that the school or district provided NO support for inservice training or staff development in health education.
- 29 percent of all schools had no one designated to coordinate health education.
- 54 percent of all schools had no active health education advisory council. Of those that did, teachers and school administrators were the most common members (42 and 36 percent respectively). 30 percent of schools had parents represented on the council, 21 percent had students represented, 9 percent had public health departments represented, and 7 percent had churches represented.
- 39 percent of the lead health education teachers involved parents in health instruction through homework assignments. 30 percent did not involve parents at all.
- 45 percent of principals reported that feedback from parents about health education in the school was mainly positive during the past year. 48 percent said parents provided no feedback.
- 92 percent of schools had a smoke-free policy, 93 percent had a drug-free policy, and 53 percent had a violence-free policy.

## AMOUNT OF INSTRUCTION

- 86 percent of senior high schools required students to take one-half year or more of health education. 50 percent of all senior high schools scheduled health education at the 10th grade, while only 10 percent

scheduled it for 11th grade students and 9 percent for 12th graders.

- 61 percent of middle schools required students to take less than one-half year or more of health education. 86 percent did not require students to repeat health education if failed.

## INSTRUCTIONAL CONTENT

- 96 percent of lead health education teachers attempted to increase students' knowledge about alcohol, tobacco and other drug use prevention. 77 percent attempted to help students improve skills to avoid alcohol and other drugs.
- 91 percent of lead health education teachers attempted to increase students' knowledge about tobacco use, and 63 percent tried to improve students' skills to avoid tobacco.
- 86 percent of lead health education teachers attempted to increase students' knowledge about HIV prevention, and 59 percent tried to improve students' skills to avoid HIV infection.

## TEACHER PREPARATION

- 62 percent of all lead health education teachers' major emphasis of professional preparation was in physical education. 90 percent of all the lead teachers were certified or endorsed to teach health education.
- 65 percent of lead health education teachers had been teaching health education for more than five years. 26 percent of all lead teachers had attended no inservice training during the prior two years.
- More than 40 percent of lead health education teachers had received four or more hours of inservice training in alcohol and other drug use prevention and HIV prevention during the prior two years. More than 40 percent also indicated they would like to attend inservice on these topics.
- Less than 15 percent of lead health education teachers had attended inservice on violence and suicide prevention during the prior two years. More than 40 percent indicated that they would like to attend inservice on these topics.

# 1995 Missouri Youth Risk Behavior Survey

## *Executive summary*

The 84-item, multiple-choice Youth Risk Behavior Survey (YRBS) was administered to 4,900 students in 23 public high schools during the spring of 1995. The school response rate was 79 percent, and the student response rate was 80 percent. Survey procedures were designed to protect the privacy and confidentiality of all participating students. Participation was voluntary.

The students who participated in the survey are representative of students in Missouri. The results may be used to make important inferences concerning the health-risk behaviors of all Missouri public high school students in grades 9 through 12.

The YRBS was developed by the Division of Adolescent and School Health, National Center for Chronic Disease Prevention and Health Promotion, at the Centers for Disease Control and Prevention. Representatives from 71 state and local departments of education and 19 other federal agencies also assisted with the design of the survey. Its purpose is to monitor priority health-risk behaviors that contribute to the leading causes of mortality, morbidity and loss of quality of life among youth and adults in the United States. These behaviors fall into six categories:

- Tobacco use
- Alcohol and other drug use
- Dietary behaviors
- Physical activity
- Behaviors that result in unintentional and intentional injuries
- Sexual behaviors that result in HIV infection, other sexually transmitted diseases (STDs) and unintended pregnancies



# Key findings

## ALCOHOL, TOBACCO AND OTHER DRUG USE

- 75 percent had tried smoking cigarettes; 30 percent smoked two or more cigarettes on the days they smoked.
- 83 percent had tried alcohol, 40 percent before the age of 13.
- 40 percent drank five or more drinks in a row sometime during the past 30 days. (44 percent male, 36 percent female; 34 percent of 9th graders, 37 percent 10th graders, 47 percent 11th graders, and 44 percent 12th graders)
- 26 percent had someone offer, sell or give them illegal drugs on school property during the past 12 months.

## BEHAVIORS THAT RESULT IN UNINTENTIONAL AND INTENTIONAL INJURIES

- 45 percent had ridden in a vehicle during the past 30 days with someone who had been drinking alcohol.
- 30 percent of 11th graders and 31 percent of 12th graders had driven a car during the past 30 days when they had been drinking alcohol.
- 15 percent were in a physical fight on school property within the past 12 months.
- 8 percent had been threatened or injured with a weapon on school property
- 20 percent had made a plan about how they would commit suicide, and 9 percent attempted.

## SEXUAL BEHAVIORS THAT RESULT IN HIV INFECTION, OTHER SEXUALLY TRANSMITTED DISEASES AND UNINTENDED PREGNANCIES

- 54 percent had engaged in sexual intercourse during their lifetime (56 percent of males, 52 percent of females; 41 percent of 9th graders, 48 percent of 10th graders, 62 percent of 11th graders, and 69 percent of 12th graders).


- 13 percent had sexual intercourse for the first time before the age of 13.
- 39 percent had engaged in sexual intercourse during the previous three months, of which 52 percent used a condom the last time. (60 percent of 9th graders, 56 percent 10th graders, 47 percent 11th graders, and 48 percent 12th graders)
- 64 percent had discussed AIDS or HIV infection with their parents or other adults in their family.
- 86 percent had been taught about HIV infection or AIDS in school.

## DIETARY AND PHYSICAL ACTIVITY BEHAVIORS

- 60 percent ate fruit, 57 percent drank fruit juice, 50 percent ate cooked vegetables, and 29 percent ate green salad one or more times on the day before the survey.
- 65 percent ate french fries or potato chips, 59 percent ate hamburger, hot dog or sausage, and 64 percent ate cookies, doughnuts, pie or cake one or more times the day before the survey.
- 43 percent were trying to lose weight; 51 percent exercised to lose weight or keep from gaining weight during the past 30 days.
- 63 percent had exercised or participated in sports activities that made them sweat or breathe hard for at least 20 minutes on 3 or more of the past 7 days.
- 53 percent did stretching exercises and 50 percent did exercises to strengthen or tone muscles on 3 or more days of the past 7 days.
- 34 percent walked or bicycled for at least 30 minutes at a time on 3 or more of the past 7 days.
- 47 percent attended physical education class one or more days during an average school week (73 percent of 9th graders, 43 percent of 10th graders, 32 percent of 11th graders, and 32 percent of 12th graders).

# Summary of 1995 Youth Risk Behavior Survey findings

PERCENT OF PARTICIPANTS WHO . . .





Missouri Department of Elementary and Secondary Education  
Division of Instruction  
P.O. Box 480  
Jefferson City, MO 65102-0480



# 1996 School Health Education Profile

---

▶ *Executive Summary and Key Findings*

---

HEALTHY  
Healthy Students  
MISSOURI  
Healthy Schools

**October 1996**

**Missouri Department of Elementary and Secondary Education  
Robert E. Bartman, Commissioner of Education**



# 1996 Missouri School Health Education Profile

## *Executive summary*

The School Health Education Profile is a survey conducted in the spring of even-numbered years, as part of a cooperative agreement between the Missouri Department of Elementary and Secondary Education and the federal Centers for Disease Control and Prevention (CDC). The first survey was conducted in Missouri in 1994, and results were reported in a document titled *Recommendations for Improving School Health Education and Reducing Health Risks for Missouri Students* (August 1996). This report highlights the findings of the second (1996) School Health Education Profile survey.

During the spring of 1996, questionnaires prepared by the CDC Division of Adolescent and School Health were sent to the principal and a designated lead health education teacher in 340 secondary schools in the state. Usable questionnaires were returned by 249 principals and 250 teachers.

The responses are representative of secondary principals and health education teachers in Missouri public schools, and results may be used to develop policies and improve programs for school health education, including HIV/AIDS prevention education.

Survey results were compiled in the following categories: (1) overall results for all schools; (2) for schools composed primarily of grades 9-12; (3) for schools composed primarily of grades 6-8; and (4) for schools composed primarily of grades 7-12. Not all data are reported in this summary. Key findings representing significant changes from the 1994 survey results are reported and discussed.

## *Key findings from 1996*

### INFLUENCES ON HEALTH BEHAVIOR

- The percent of principals who reported that their schools did not use trained peer educators increased to 68 percent compared to 50 percent in 1994.
- 75 percent of the teachers reported they include analysis of media messages in their health education instruction. (No data reported in 1994.)

### SUPPORT FOR SCHOOL HEALTH EDUCATION

- School or district support for inservice training or staff development in health education increased, with 84 percent providing substitutes (58 percent in 1994) and 74 percent reimbursing expenses (58 percent in 1994).
- Schools with someone designated to coordinate health education increased to 90 percent from 71 percent in 1994.
- Teachers' attempts to involve parents in health instruction increased, with 57 percent including parents in homework assignments (39 percent in 1994) and 49 percent sending educational materials home to parents (28 percent in 1994).
- Schools with an active health education advisory council decreased to 41 percent from 46 percent in 1994. However, of those with a council, the composition changed considerably as shown by these percentages:

|   | <u>1994</u> | <u>1996</u> |
|---|-------------|-------------|
| Students                                  | 21%         | 67%         |
| Parents                                   | 30%         | 92%         |
| Teachers                                  | 42%         | 98%         |
| School administrators                     | 36%         | 97%         |
| Food service staff                        | 11%         | 32%         |
| School nurses                             | 24%         | 88%         |
| Counselors                                | 26%         | 73%         |
| School board members                      | 15%         | 49%         |
| Public health dept. staff                 | 9%          | 46%         |
| Business community                        | 8%          | 45%         |
| Medical community                         | 12%         | 59%         |
| Mental health community                   | 3%          | 18%         |
| Churches or other religious organizations | 7%          | 36%         |
| Community-based organizations             | 9%          | 39%         |
| Law enforcement organizations             | NA          | 35%         |
| Other                                     | 7%          | 22%         |

## AMOUNT OF INSTRUCTION

- Required health education for students in grades 11 and 12 more than doubled to 24 and 22 percent respectively from 10 and 9 percent in 1994.
- Health instruction for middle school students increased with 78 percent of the schools requiring one-half year or more, compared to 66 percent in 1994.

## INSTRUCTIONAL CONTENT

- A very high percentage of teachers continued attempts to increase student knowledge on a variety of health topics.
- A high percentage of teachers reported teaching a variety of skills in health education courses (e.g., decision-making, 98 percent; resisting social pressure for unhealthy behaviors, 97 percent; goal setting, 89 percent; stress management, 88 percent; communication, 87 percent; non-violent conflict resolution, 79 percent; and analysis of media messages, 75 percent).
- Significantly fewer teachers taught skills in 1994 (e.g., 69 percent taught decision-making and 53 percent taught communication skills to avoid HIV risk behaviors).

## TEACHER PREPARATION

- 40 percent of the teachers' major emphasis of professional preparation was in health and physical education. 90 percent were certified to teach health education. 61 percent had been teaching health education for more than five years. 49 percent had been in the teaching profession for 15 years or more.
- The most common inservices attended by teachers in the past two years were on CPR (51 percent) and alcohol and other drug-use prevention (41 percent).
- 34 percent of the teachers received four or more hours of inservice on HIV prevention during the past two years compared to 44 percent in 1994.
- Teachers' interest in receiving inservice increased overall from 1994, especially in the following topics:

|   | <u>1994</u> | <u>1996</u> |
|---|-------------|-------------|
| HIV Prevention                              | 48%         | 67%         |
| Suicide Prevention                          | 44%         | 62%         |
| Conflict Resolution/<br>Violence Prevention | 49%         | 56%         |
| Alcohol and other<br>Drug-Use Prevention    | 42%         | 54%         |

## Discussion

### SIGNIFICANT CHANGES FROM 1994

1. There was an increase in support for comprehensive health education as indicated by more schools identifying health education coordinators and through more involvement of parents, community members and students on advisory councils. Continued support is critical in order to have effective programs.
2. There was an increase in skills instruction. Research indicates that providing only factual information about health-related topics is not enough to influence student behavior. Therefore, it is essential for schools to continue to provide opportunities for students to learn health-enhancing skills which prepare them to make responsible decisions about their health.
3. There was an increase in teachers' desire for inservice in HIV, suicide, violence and alcohol-use prevention. This may indicate that teachers are concerned about the number of students who are engaging in health-risk behaviors. (The incidence of such behavior is well documented in the 1995 Missouri Youth Risk Behavior Survey.) Opportunities for teacher training and inservice should be continued.
4. Although there was an increase in required health education for 11th and 12th graders, the overall percent of secondary schools requiring comprehensive health education decreased. The Missouri School Improvement Program requires districts to provide comprehensive health instruction, including, specifically, drug and alcohol abuse prevention education and AIDS prevention education to all students at every grade level.
5. There was a decrease in the percentage of schools using trained peer educators. Research indicates that effective programs involve students in decisions about programs and allows them to present positive messages to their peers.

## Conclusion

Secondary school personnel are to be commended for the strides which have been made in improving comprehensive health education. It is hoped that this report will encourage continued efforts to improve the health of Missouri students and enhance programs that reduce health risks for all young people.


## ACKNOWLEDGMENTS

Sincere appreciation is extended to the secondary principals and lead health education teachers who completed the 1996 School Health Education Profile questionnaires.

Special thanks to Nancy Speicher and Candi Hitchcock of Westat, Inc., for their assistance with data tabulation and analysis.

This report was prepared by:  
Sandy Nichols, Health and Physical Education Consultant  
Janet Wilson, HIV/AIDS Prevention Education Supervisor

This document was printed with funding provided through a cooperative agreement (U87/CCU709009-04) between the Missouri Department of Elementary and Secondary Education and the Division of Adolescent and School Health of the U.S. Centers for Disease Control and Prevention.



Missouri Department of Elementary and Secondary Education  
Division of Instruction  
P.O. Box 480  
Jefferson City, MO 65102-0480



**U.S. DEPARTMENT OF EDUCATION**  
*Office of Educational Research and Improvement (OERI)*  
*Educational Resources Information Center (ERIC)*



## NOTICE

### REPRODUCTION BASIS

This document is covered by a signed "Reproduction Release (Blanket)" form (on file within the ERIC system), encompassing all or classes of documents from its source organization and, therefore, does not require a "Specific Document" Release form.

This document is Federally-funded, or carries its own permission to reproduce, or is otherwise in the public domain and, therefore, may be reproduced by ERIC without a signed Reproduction Release form (either "Specific Document" or "Blanket").