

## DOCUMENT RESUME

ED 440 332

CG 029 874

AUTHOR Weinstein, Erica S.  
TITLE School-Based Eating Disorder Prevention Programs for Pre-Adolescents and Adolescents: A Review of Recent Literature.  
PUB DATE 2000-04-00  
NOTE 17p.  
PUB TYPE Information Analyses (070)  
EDRS PRICE MF01/PC01 Plus Postage.  
DESCRIPTORS Adolescent Development; Early Adolescents; \*Eating Disorders; \*Females; Health Promotion; Late Adolescents; \*Literature Reviews; Prevention; Program Effectiveness; Secondary Education  
IDENTIFIERS \*School Based Services; Search Engines

## ABSTRACT

This paper presents a review of 25 sources on school-based eating disorder prevention programs for pre-adolescents and adolescents. The sources used to collect the information include Search ERIC database, PsycINFO, InterScience, and Expanded Academic. A review of the literature concluded that the most effective method of implementing a school-based eating disorder program has been to use a comprehensive curriculum integrating primary and secondary prevention techniques. A small group context was the format found to be most constructive for girls with eating disorders due to its high relational characteristics. The research suggests that a comprehensive program should address such essential components as developmental education, life skills training, empowerment, media literacy training, and self-esteem enhancement. It encourages longitudinal research since change occurs rapidly during the developmental stage of adolescents, and aspects of the prevention program that are effective at one time, may not be at another time. (Contains 25 references.) (JDM)

CG

ED 440 332

Running head: SCHOOL-BASED EATING DISORDER PREVENTION PROGRAMS

School-Based Eating Disorder Prevention Programs for Pre-Adolescents and Adolescents

A Review of Recent Literature

Erica S. Weinstein

Loyola College in MD

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

"PERMISSION TO REPRODUCE THIS MATERIAL HAS BEEN GRANTED BY

E. WEINSTEIN

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)."

BEST COPY AVAILABLE

## Abstract

In this review of the literature, 25 sources on the topic of school-based eating disorder prevention programs for pre-adolescents and adolescents were examined. The search engines used to collect sources included the Search ERIC database, PsycINFO, InterScience, and Expanded Academic. Research conducted by Neumark-Sztainer was well represented throughout the literature on eating disorders in youth. The culminating research concluded that the most effective method of implementing a school-based eating disorder program has been to use a comprehensive curriculum integrating primary and secondary prevention techniques. Due to its high relational characteristics, the small group context was the format that was found to be most constructive for girls with eating disorders. In the research, it was concluded that a comprehensive program should address such essential components as developmental education, life skills training, empowerment, media literacy training, and self-esteem enhancement. In order for this field of study to advance, longitudinal studies have been suggested.

## School-Based Eating Disorder Prevention Programs for Pre-Adolescents and Adolescents

The review of the literature on school-based eating disorder prevention programs for pre-adolescents and adolescents was culminated from 25 sources including two articles from the microfiche collection, one dissertation abstract, and the rest scholarly journals. Eating disorders have been a relatively new subject of discussion in the schools; therefore, the publication dates spanned the 1990s. The search engines used were ERIC, Expanded Academic, InterScience, and PsycINFO. For the purpose of this review, the term eating disorders was defined as anorexia nervosa, bulimia nervosa, and dieting as a means of extreme weight control.

The significance of this review relied upon the statistical data which showed that the numbers of school aged children suffering from eating disorders and eating disordered behavior have been growing. In addition, the age of children exhibiting symptoms has become younger and younger. In order to deter children from developing eating disorders, primary prevention programming was suggested to begin in the elementary grades. The review was divided into four subsections. The first part discussed various types of prevention programming and the differences between the efficacy of primary and secondary prevention. The second section reviewed the role of comprehensiveness in prevention programming. The third portion focused on the myriad variables contributing to disordered eating. Finally, the fourth segment synthesized the first three by reviewing components of successful school-based prevention programs.

### Types of Prevention Programming and the Efficacy of Primary and Secondary Prevention

The three identified types of prevention programming are primary, secondary, and tertiary. Primary preventative efforts have focused on the identification, reduction, and elimination of risk factors contributing to a disorder (Huon, Braganza, Brown, Ritchie, &

Roncolato, 1998). In a review of six school-based prevention programs, Carter, Stewart, Dunn, and Fairburn (1997) found that primary prevention strategies typically included “education about the nature and consequences of eating disorders; discussion about the adverse effects of dieting and other methods of weight control; and skills-training for resisting social pressures to diet” (p.168). Primary and secondary prevention methods are very similar, but their target populations have differed; there was substantial debate in the literature as to which strategy has been more effective in the school system.

According to Huon et al. (1998), secondary prevention programming has occurred when the focus is on “reducing chronicity through early identification and intervention” (p.456). Taking these definitions into account, the target population has been the key difference. Primary prevention has targeted low-risk groups of students (general school population), while secondary prevention has aimed to reach pre-adolescents and adolescents at high-risk for developing, or those whom have already been exhibiting symptoms of an eating disorder.

One of the conclusions in this field of study, was that prevention efforts aimed at low-risk populations have been more successful than those administered to high-risk subjects (Nakamura, Hoshino, Watanabe, Honda, Niwa, & Yamamoto, 1999). This was due to studies that indicated once in effect, disordered eating behaviors have been extremely difficult for pre-adolescents and adolescents to change, especially in regards to negative self-perception (Neumark-Sztainer, Butler, & Palti, 1995; Shisslak, Crago, Renger, & Clark-Wagner, 1998). Considering that poor self-image and distorted body image take years to manifest as destructive concepts, it is logical to expect that cognitive-behavioral change takes time, and considerable effort is required on the part of the eating disordered person in order to “unlearn” the undesired behavior. From this

perspective, it was the primary prevention that targeted low-risk students that seemed most effective.

However, other studies concluded that the opposite was true. Killen et al. (1993) as cited in Graber and Brooks-Gunn (1996) argued that primary prevention for eating disorders has been inappropriate. The high-risk students who have already displayed signs of disordered eating behaviors are the ones who should be targeted by secondary preventative measures. In addition, Varnando (1998) stated that targeting youth at the highest risk for developing eating disorders resulted in more effective prevention efforts. One of the reasons given for the ineffectiveness of including the entire school population in prevention efforts was the multiple demands made on school personnel's time by individual students' and group needs (Chally, 1998). Murray, Touyz, & Beaumont (1990) and Watts & Ellis (1992) as cited in Chally (1998) discovered that in spite of trying to prevent eating disordered behavior, deviant eating behavior was sometimes learned through discussion. From this perspective, it was the secondary prevention programming aimed at high-risk and/or already affected students that proved most effective.

Tertiary prevention has been the strategy applied least in the schools. Tertiary methods have involved treating the behavior that has already been manifested. Gabel and Kearney (1998) defined tertiary prevention as "focusing on reducing the impairment that may result from an established disorder" (p.33). It was evident why this prevention method has been used least in the schools; in most cases, a student who has been entrenched in an eating disorder was in need of more intensive therapy than what a school counselor can have provide (Cappuzzi, 1996). Frequently, these students have been given referrals to appropriate organizations and professionals outside of the school system.

It is vital to note that tertiary prevention, the most well developed area of research, has been the least likely method to be used in the schools. Yet, the school environment has been where the most appropriate target population for employing primary and secondary prevention exists (Paxton, 1996). Primary and secondary prevention research is still in the preliminary stages. Considering that the school setting has been the optimal place for prevention strategies to affect the highest number of pre-adolescents and adolescents, much more research on the implementation and evaluation of primary and secondary prevention programming has been needed.

The literature indicated that the most commonly recommended forms of prevention for eating disordered behavior has been a primary approach or a combination of primary and secondary, even though there has been a paucity of research describing developmentally appropriate and theoretically based primary prevention strategies (Rosen, 1998). This conclusion was contradictory. How has it been that the studies encouraged primary prevention as a means to deter disordered eating habits in pre-adolescents and adolescents, when there has been so little research to support this claim?

#### The Role of Comprehensiveness in Prevention Programming

In the majority of the literature the importance of comprehensiveness, or an all-encompassing approach was highly stressed as a component of successful prevention programs (Neumark-Sztainer, 1996; Azzarto, 1997; Rosen, 1998; Stipek, de la Sota, & Weishaupt, 1999). Due to the multidimensional factors that have contributed to the development of eating disorders, a holistic programming strategy was suggested. Pre-adolescents' and adolescents' behaviors, attitudes, and beliefs have been affected by several different variables: cultural, social, psychological, and biological to name a few (Steiner & Lock, 1998). Therefore, attention must

be paid to not only what occurs in the classroom, but also transactions in the entire school, between the school and the family, and in the wider community as a whole (Dixey, 1996).

Due to the expansiveness of programming, comprehensiveness also has ensured that the greatest number of students has been reached. Neumark-Sztainer (1996) provided a framework for a comprehensive program based on an ecological model for health promotion.

The components of the program included staff training; a module for preventing eating disturbances and obesity for junior high students; formal and informal integration of material into existing curricula for all students; individual counseling and small group work for high-risk adolescents; a referral system within the school and between the school and community health services; opportunities for healthy eating at school; modifications in physical education and sport activities to encourage opportunities for physical activity for all students and policies preventing extreme weight fluctuations in the interest of improving team or individual competitiveness; and outreach activities in the school and the community by students as part of the curriculum and by staff and interested parents (p.68).

From this example of an effective comprehensive prevention program, it was evident that both primary and secondary strategies needed to be employed. The spectrum of opportunities for teaching and learning in a comprehensive program should be broad and well represented in the school system.

Another essential component of comprehensive prevention programming has been the inclusion of evaluation methods. As reflected in the literature, this crucial step has often been omitted in several of the studies due to the difficulty of conducting longitudinal studies with school populations (Childress, Brewerton, Hodges, & Jarrell, 1993; Neumark-Sztainer, 1995;



Chally, 1998). Even though some of the studies have employed immediate evaluative procedures such as the pre-test/post-test method, only one longitudinal study has been performed. Long-term follow up has been suggested in the literature, but not implemented (Moreno & Thelen, 1993; Steiner & Lock, 1998). The lack of use of this evaluative tool has been problematic in the field of study of preventing eating disorders because the efficacy of the prevention programs has not been able to be measured over time.

### Variables Contributing to Disordered Eating

The varied factors that have contributed to eating disordered behavior were heavily represented in the research. In the literature, the three categorizations of factors that have contributed to disordered eating were environmental, personal, and behavioral. Perry, Story, and Lytle (1997), listed the following models under each category. Environmental factors included access to foods, role models, social support, social norms, and parenting styles. Knowledge, self-efficacy, functional meanings, attitudes, self-image, and locus of control were stated as personal factors. The behavioral factors were listed as dietary skills, decision-making skills, coping responses, intentions, reinforcements, and behavioral repertoire. In addition, age and gender have been strongly correlated with the incidences of disordered eating, perceptions of body image, and methods of weight regulation (Vervae, van Heeringen, & Jannes, 1998). For the purpose of this review, the personal factor of self-esteem and the social-environmental factor of the role of the media have been expounded upon. It has been indicated that the two are interrelated and comprise a large portion of the reason why so many young people seem obsessed with appearances.

Self-esteem has been identified as a multi-layered construct of the personality, which has lent itself to much scrutiny. A prevalent question in the research was which comes first, low

self-esteem or the eating disorder (Shisslak et al., 1998)? In this particular study, a causal relationship was not determined because there was no method of measurement. The question could have been answered if the researchers had conducted a longitudinal study measuring self-esteem and eating disordered behavior at baseline and then again at a later date. As mentioned previously, the absence of long-term and follow-up studies has stunted the development of more effective prevention programming for students suffering with eating disorders.

There have been a few approaches to enhancement of self-esteem (for girls) that could be used in a prevention program or in a health education curriculum. Participation in sports, media literacy training, and body enhancement were suggested as possible topics to be included in an intervention program (Shisslak et al.). A different viewpoint was offered in an article focusing on group work and empowerment. Friedman (1998) argued that the nature of prevention programming has been didactic and therefore disempowering, which has left girls in vulnerable positions. It was suggested that girls take a pro-active stance by leading and co-facilitating groups, which in turn was reported to empower them and increase their self-esteem.

Closely linked to the development of self-esteem, or lack thereof, has been the role of the media in young people's perceptions of what is normal, acceptable, and desirable in terms of body shape and size. In one cross-sectional study, high school students were asked the principal sources of information from which they had learned about food, nutrition, risk of being overweight, and dieting to reduce weight. The majority of the information was gleaned from newspapers (78%) and television (75.7%) (Brook & Tepper, 1997). It has been found that over-exposure to the media has been linked to the likelihood of whether or not a young adolescent will develop an eating disorder (Shisslak et al., 1999). These conclusions should send a social signal that something has been dreadfully wrong with the images that have been portrayed in the media.

In order to decrease the amount of influence the media has wielded over the pre-adolescent and adolescent audience, Stipek et al., (1999) suggested implementing a curriculum based prevention program which “helps students understand how outside entities that do not necessarily have their best interest in mind...attempt to influence their behavior” (p.450). In addition, teaching students coping skills to deal with media influenced pressures such as the desire to diet in order to remain thin was encouraged (Santonastaso, Zanetti, Ferrara, Olivotto, Magnavita, & Favaro, 1999). The glamorization of excessive thinness has been slowly, but progressively eating away at pre-adolescents and adolescents in the schools.

#### Components of Successful School-Based Eating Disorder Prevention Programs

Even though there has not been an abundance of successful prevention programs represented in the literature, the ones discussed shared certain components and formats. First, the small group format was found to prove most encouraging and helpful for pre-adolescent and adolescent girls (Paxton, 1996; Azzarto, 1997; Friedman, 1998). This was explained by the relational context which support groups have often provided. Pirvan (1995) postulated that working in a supportive group in the school with some adults, as well as with peers, led to the formation of an alternative and protective subculture. The relational context of the group has often provided girls at high-risk for development of, and those who have already been suffering with eating disorders with a space that has been safe and empowering.

Empowerment was cited as a crucial component for the efficacy of eating disorder prevention programming. One way in which adolescents experienced empowerment was when they were included in the decision-making and implementation processes. Often, power has been viewed as a construct that is associated with the dominance of the male culture. Many girls whom exhibited eating disordered behavior described a feeling of powerlessness when it comes

to food (Dixey, 1996). Therefore, when were responsible for part of the planning and decision-making processes within the group, they asserted themselves; they felt empowered. The strength they experienced by having been involved in the process may have led to an increase in self-esteem, while simultaneously have advanced them in social and leadership skill development.

Leadership skill development was a component of another well represented modality prevalent in the literature for the implementation of eating disorder prevention programs. Life skills training has been proven to work effectively with adolescents who exhibit any sort of high-risk behavior (Huon et al., 1998; Stipek et al., 1999). The comprehensiveness of teaching life skills in the schools has been the factor that has applied so well with prevention programming. Life skills training is not a one time course, mini-lecture, or special educational program; rather, it has been a continual method of enforcing healthy decision-making for students that have hopefully lasted a lifetime. Stipek et al. (1999) stated, “it is a way of educating the whole child by embedding lessons related to physical, mental, emotional, and social health within the everyday curriculum and children’s everyday experiences, in and out of the classroom” (p.451). By having focused on the whole child, instead of merely having concentrated on the disordered eating behavior, chances were greater that the student was able to integrate his learning experiences into making health promoting decisions regarding high-risk behavior.

In cohesion with having learned life skills, comes the developmental appropriateness of prevention programming. The research indicated that developmental education in preventative strategies for eating disorders has often been omitted. In an earlier study conducted by Graber, it was found that girls with persistent deviant eating behaviors in young and mid adolescence went through puberty earlier than other girls. It is no coincidence that the normative age at which girls have started to diet and become concerned with body image has occurred simultaneously with

the on-set of pubescent changes. It has been suggested that girls be taught that certain bodily developments, such as breast and hip growth, are normal and necessary transitional biological functions for their age group. A well-balanced, comprehensive curriculum that stressed normative developmental changes could assist girls in accepting and appreciating their bodies as they grow and mature into young women (Graber & Brooks-Gunn, 1996).

### Summary

In short, the research revealed that the most effective method of employing a school-based eating disorder prevention program or curriculum has been to follow a comprehensive plan that integrates primary and secondary prevention strategies. It has been necessary to incorporate the myriad factors that have contributed to disordered eating within the broader categories of environmental, personal, and behavioral. The most effective format for girls with eating disorders has been the group setting, as it has provided an environment rich in relational context. A few of the target areas that should be addressed in the program were self-esteem enhancement, media literacy training, empowerment, life skills training, and developmental education.

### Conclusions

Much of the content in the literature was repetitive. Many of the studies reached similar conclusions that there has not been sufficient enough research, specifically longitudinal studies to ensure that the prevention measures were sound and reliable. This conclusion led the writer to believe that the school-based eating disorder prevention-programming field of study is stagnant. Even though several studies have claimed that preventing eating disorders in pre-adolescents and adolescents is paramount, only one longitudinal study has been conducted. Cross-sectional studies were well represented in the literature and do not need to be replicated for further advancement in the field.

In order to broaden insights in the field of school-based eating disorder prevention programming, several longitudinal studies are suggested. In order to track the efficacy of the educational efforts of prevention programs over time, long-term studies must be conducted. It has seemed rather unintelligent to spend time, energy, and money on programs that have had no long-term evaluative effect. It has been well documented that change occurs quite rapidly during the developmental stage of adolescence. A post-test that has been administered 6 months after a prevention program may show significant improvement in adolescents' knowledge and attitudes toward a healthy lifestyle. However, a year, or even another 6 months later, the results of that study would have a good chance of being completely different. Until researchers decide to implement longitudinal studies tracking the efficacy of school-based eating disorder prevention programs, the field will continue to produce similar results and findings. This cyclical ineffectiveness certainly has not helped those students who have suffered.

## References

- Azzarto, J. (1997). A young women's support group: Prevention of a different kind. Health and Social Work, 22, 299-305.
- Brook, U. & Tepper, I. (1997). High school students' attitudes and knowledge of food consumption and body image: Implications for school based education. Patient Education and Counseling, 30, 283-288.
- Capuzzi, D. (1996). The secret and all-consuming obsessions: Anorexia and bulimia. In D.R. Gross (Ed.), Youth at risk: A prevention resource for counselors, teachers, and parents. (pp. 153-181). Alexandria, VA: American Counseling Association.
- Carter, J., Stewart, D., Dunn, V., & Fairburn, C. (1997). Primary prevention of eating disorders: Might it do more harm than good? International Journal of Eating Disorders, 22, 167-172.
- Chally, P. (1998). An eating disorders prevention program. Journal of Child and Adolescent Psychiatric Nursing, 11 (2), 51-60.
- Childress, A., Brewerton, T., Hodges, E., Jarrell, M. (1993). The kids' eating disorder survey (KEDS): A study of middle school students. Journal of the American Academy of Child and Adolescent Psychiatry, 32, 843-850.
- Dixey, R. (1996). Healthy eating in schools and "eating disorders"— are "healthy eating" messages part of the problem or part of the solution? Nutrition and Health, 11, 49-58.
- Friedman, S. (1998). Girls in the 90s: A gender-based model for eating disorder prevention. Patient Education and Counseling, 33, 217-224.
- Gabel, K., & Kearney, K. (1998). Promoting reasonable perspectives of body weight: Issues for school counselors. Professional School Counseling, 1 (5), 32-35.

- Graber, J., Brooks-Gunn, J. (1996). Prevention of eating problems and disorders: Including parents. Eating Disorders: The Journal of Treatment and Prevention, 4 (4), 348-363.
- Huon, G., Braganza, C., Brown, L., Ritchie, J., & Roncolato, W. (1998). Reflections on prevention in dieting-induced disorders. International Journal of Eating Disorders, 23, 455-458.
- Moreno, A., & Thelen, M. (1993). A preliminary prevention program for eating disorders in a junior high school population. Journal of Youth and Adolescence, 22, 109-124.
- Neumark-Sztainer, D., Butler, R., & Palti, H. (1995). Eating disturbances among adolescent girls: Evaluation of a school-based primary prevention program. Journal of Nutrition Education, 27 (1), 24-31.
- Neumark-Sztainer, D. (1996). School-based programs for preventing eating disturbances. Journal of School Health, 66 (2), 64-71.
- Paxton, S. (1996). Prevention implications of peer influences on body image dissatisfaction and disturbed eating in adolescent girls. Eating Disorders: The Journal of Treatment and Prevention, 4 (4), 334-337.
- Perry, C., Story, M., & Lytle, L. (1997). Promoting healthy dietary behaviors. Enhancing Children's Wellness (pp. 214-243).
- Piran, N. (1995). Prevention: Can early lessons lead to a delineation of an alternative model? A critical look at prevention with schoolchildren. Eating Disorders: The Journal of Treatment and Prevention, 3 (1), 28-36.
- Rosen, D., & Neumark-Sztainer, D. (1998). Review of options for primary prevention of eating disturbances among adolescents. Journal of Adolescent Health, 23, 354-363.



Santonastaso, P., Zanetti, T., Ferrara, S., Olivitto, M., Magnavita, N., & Favaro, A. (1999). A preventive intervention program in adolescent schoolgirls: A longitudinal study. Psychotherapy and Psychosomatics, *68* (1), 46-50.

Shisslak, C., Crago, M., Renger, R., & Clark-Wagner, A. (1998). Self-esteem and the prevention of eating disorders. Eating Disorders: The Journal of Treatment and Prevention, *6* (2), 105-118.

Steiner, H., & Lock, J. (1998). Anorexia nervosa and bulimia nervosa in children and adolescents: A review of the past 10 years. Journal of the American Academy of Child and Adolescent Psychiatry, *37*, 352-359.

Stipek, D., de la Sota, A., & Weishaupt, L. (1999). Life lessons: An embedded classroom approach to preventing high-risk behaviors among preadolescents. The Elementary School Journal, *99*, 433-352.

Varnado, P. (1998). Development and implementation of the body logic program for adolescents: A primary prevention program for eating disorders (Doctoral dissertation, The Louisiana State University and Agricultural and Mechanical College, 1998). Dissertation Abstracts International, *58*, (9-B) 5145.

Vervaeke, M., van Heeringen, C., & Jannes, C. (1998). Weight concerns and eating patterns in schoolboys and girls. Eating Disorders: The Journal of Treatment and Prevention, *6* (1), 41-50.

Whisenant, S., & Smith, B. (1995). Eating disorders: Current nutrition therapy and perceived needs in dietetics education and research. Journal of the American Dietetic Association, *95*, 1109-1112.



**U.S. Department of Education**  
 Office of Educational Research and Improvement  
 (OERI)  
 National Library of Education (NLE)  
 Educational Resources Information Center (ERIC)



## Reproduction Release

(Specific Document)

### I. DOCUMENT IDENTIFICATION:

Title: <i>School-Based Eating Disorder Prevention Programs for Pre-Adolescents and Adolescents: A Review of Recent Literature</i>	
Author(s): <i>Erica Sharyn Weinstein</i>	
Corporate Source:	Publication Date: <i>April 2000</i>

### II. REPRODUCTION RELEASE:


In order to disseminate as widely as possible timely and significant materials of interest to the educational community, documents announced in the monthly abstract journal of the ERIC system, Resources in Education (RIE), are usually made available to users in microfiche, reproduced paper copy, and electronic media, and sold through the ERIC Document Reproduction Service (EDRS). Credit is given to the source of each document, and, if reproduction release is granted, one of the following notices is affixed to the document.

If permission is granted to reproduce and disseminate the identified document, please CHECK ONE of the following three options and sign in the indicated space following.

The sample sticker shown below will be affixed to all Level 1 documents	The sample sticker shown below will be affixed to all Level 2A documents	The sample sticker shown below will be affixed to all Level 2B documents
PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY _____ _____ TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)	PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN MICROFICHE, AND IN ELECTRONIC MEDIA FOR ERIC COLLECTION SUBSCRIBERS ONLY, HAS BEEN GRANTED BY _____ _____ TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)	PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN MICROFICHE ONLY HAS BEEN GRANTED BY _____ _____ TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)
Level 1	Level 2A	Level 2B
Check here for Level 1 release, permitting reproduction and dissemination in microfiche or other ERIC archival media (e.g. electronic) and paper copy.	Check here for Level 2A release, permitting reproduction and dissemination in microfiche and in electronic media for ERIC archival collection subscribers only	Check here for Level 2B release, permitting reproduction and dissemination in microfiche only

Documents will be processed as indicated provided reproduction quality permits.  
If permission to reproduce is granted, but no box is checked, documents will be processed at Level 1.

*I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce and disseminate this document as indicated above. Reproduction from the ERIC microfiche, or electronic media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete inquiries.*

Signature: 	Printed Name/Position/Title: Erica S. Weinstein / graduate student	
Organization/Address: 1631 S. Hanover St. Baltimore MD 21230	Telephone: 410 837-2952	Fax:
	E-mail Address: esw71@hotmail.com	Date: April 25, 2000

**III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):**

If permission to reproduce is not granted to ERIC, or, if you wish ERIC to cite the availability of the document from another source, please provide the following information regarding the availability of the document. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents that cannot be made available through EDRS.)

Publisher/Distributor:
Address:
Price:

**IV. REFERRAL OF ERIC TO COPYRIGHT/REPRODUCTION RIGHTS HOLDER:**

If the right to grant this reproduction release is held by someone other than the addressee, please provide the appropriate name and address:

Name:
Address:

**V. WHERE TO SEND THIS FORM:**

<http://ericfac.piccard.csc.com/reprod.html>

4/20/00



Send this form to the following ERIC Clearinghouse:

However, if solicited by the ERIC Facility, or if making an unsolicited contribution to ERIC, return this form (and the document being contributed) to:

**ERIC Processing and Reference Facility**  
**4483-A Forbes Boulevard**  
**Lanham, Maryland 20706**  
**Telephone: 301-552-4200**  
**Toll Free: 800-799-3742**  
**e-mail: [ericfac@inet.ed.gov](mailto:ericfac@inet.ed.gov)**  
**WWW: <http://ericfac.piccard.csc.com>**

EFF-088 (Rev. 9/97)