

DOCUMENT RESUME

ED 459 905

PS 029 191

AUTHOR Croan, Tara; Hatcher, Juliet; Jager, Justin; Long, Melissa; O'Hare, William; Wertheimer, Richard

TITLE The Right Start State Trends: Conditions of Babies and Their Families across the Nation (1990-1998). Child Trends/KIDS COUNT Special Report.

INSTITUTION Child Trends, Inc., Washington, DC.

SPONS AGENCY Annie E. Casey Foundation, Baltimore, MD.

PUB DATE 2001-00-00

NOTE 275p.

AVAILABLE FROM Annie E. Casey Foundation, Attn: KIDS COUNT Special Report, 701 St. Paul Street, Baltimore, MD 21202. Tel: 410-223-2890; Fax: 410-547-6624; Web site: <http://www.childtrends.org>. For full text: <http://www.kidscount.org>.

PUB TYPE Numerical/Quantitative Data (110) -- Reports - Descriptive (141)

EDRS PRICE MF01/PC11 Plus Postage.

DESCRIPTORS Birth Weight; Births to Single Women; *Child Health; Early Experience; Early Parenthood; Educational Attainment; Family (Sociological Unit); *Infants; Parent Background; Poverty; Premature Infants; Prenatal Care; Racial Differences; Smoking; *Social Indicators; *Trend Analysis; *Well Being

IDENTIFIERS Ethnic Differences; *Indicators; *United States

ABSTRACT

This Kids Count special report examines how children born in each state fared on selected measures of a "right start." This statistical portrait is based on trends in eight indicators of well-being: (1) percent of total births to teens; (2) percent of repeat teen births; (3) percent of total births to unmarried women; (4) percent of total births to mothers with less than 12 years of education; (5) percent of total births to mothers receiving late or no prenatal care; (6) percent of total births to mothers who smoked during pregnancy; (7) percent of low birth-weight births; and (8) percent of preterm births. Following an introduction detailing the importance of each indicator, the report is comprised of profiles for each individual state. The data are provided for each year from 1990 to 1998, and data for the United States as a whole are presented to provide a point of comparison. Findings indicate that the most notable signs of progress in the 1990s were a drop in the percentage of mothers receiving late or no prenatal care and a drop in the percentage of mothers reporting smoking during pregnancy. Modest progress was evident in a decline in repeat teen births and a decline in the share of births to mothers with less than 12 years of education. The United States lost ground in the percentage of births that were low birth weight, the percentage of infants born prematurely, and the percentage of total births to unmarried women. Substantial differences were identified by race and Hispanic origin for each indicator except smoking during pregnancy. The report's three appendices contain state rankings by indicator for 1998; definitions, data sources, and reporting issues; and the primary contacts for state Kids Count projects. (KB)

Reproductions supplied by EDRS are the best that can be made
from the original document.

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

This document has been reproduced as
received from the person or organization
originating it.

Minor changes have been made to
improve reproduction quality.

Points of view or opinions stated in this
document do not necessarily represent
official OERI position or policy.

PERMISSION TO REPRODUCE AND
DISSEMINATE THIS MATERIAL HAS
BEEN GRANTED BY

Carol A. Emig

TO THE EDUCATIONAL RESOURCES
INFORMATION CENTER (ERIC)

1

THE RIGHT START STATE TRENDS

CONDITIONS OF BABIES AND THEIR FAMILIES ACROSS THE NATION (1990-1998)

BEST COPY AVAILABLE

PS 029191

KIDS COUNT, a project of the Annie E. Casey Foundation, is a national and state-by-state effort to track the status of children in the United States. By providing policymakers and citizens with benchmarks of child well-being, KIDS COUNT seeks to enrich local, state, and national discussions concerning ways to secure better futures for all children. At the national level, the principal activity of the initiative is the publication of the annual *KIDS COUNT Data Book*, which uses the best available data to measure the educational, social, economic, and physical well-being of children. The Foundation also funds a nationwide network of state-level KIDS COUNT projects that provide a more detailed community-by-community picture of the condition of children.

Child Trends is a nonprofit, nonpartisan research organization dedicated to studying children, youth, and families through research, data collection, and data analyses. Child Trends conducts basic and applied research and evaluation studies; gathers and analyzes data on the major indicators of children's well-being; develops and tests new measures of child and family well-being; and communicates key research findings to policymakers and the public.

Child Trends/KIDS COUNT Special Report

THE RIGHT START STATE TRENDS

CONDITIONS OF BABIES AND THEIR FAMILIES ACROSS THE NATION (1990-1998)

Tara Croan, Juliet Hatcher, Justin Jager, Melissa Long, William O'Hare, Richard Wertheimer

The Annie E. Casey Foundation

2000 6

2000 7

Acknowledgments

This *Child Trends/KIDS COUNT Special Report* was produced and distributed with the help of numerous people. We greatly appreciate the efforts of Stephanie J. Ventura, M.A., of the National Center for Health Statistics, who educated us in the intricacies of producing reliable tabulations of birth statistics, alerted us to key connections among the measures we tabulated, and reviewed the text and tables included in this volume. Any remaining errors are the responsibility of the authors.

We thank numerous representatives of state departments of health for advising us on statistical reporting changes that affect the interpretations of trends in birth statistics.

We owe special thanks to Carol Emig, Director for Public Information and Policy at Child Trends, for reviewing the text, making valuable suggestions on how to present the material most effectively, and coordinating the publication and distribution process. We also acknowledge Eugenie Devine, Polly Dement, and others at Hager Sharp for their skillful dissemination of this report to the media.

The authors wish to thank the Annie E. Casey Foundation for providing the funding for this volume.

Finally, we would like to thank the state KIDS COUNT projects listed at the end of this report for their efforts to bring these and other data on children and families to the attention of national, state, and local leaders across the country.

Permission to copy, disseminate, or otherwise use information from this *Special Report* is granted as long as appropriate acknowledgment is given.

To obtain additional copies of this publication, call 410.223.2890 or write to the Annie E. Casey Foundation, Attn: *KIDS COUNT Special Report*, 701 St. Paul Street, Baltimore, MD 21202.

This report is also available on the Internet at www.childtrends.org and www.kidscount.org.

KIDS COUNT and Child Trends extend a special thanks to the National Center for Health Statistics (NCHS), the federal government's principal vital and health statistics agency and the source of the data presented in this publication. For several decades, NCHS has consistently gathered, disseminated, and explained health and vital statistics to the American public and to the research and policy communities. Its staff has consistently carried out this mission with exceptional attention to quality and accuracy. The agency performs a valuable and often overlooked public service, for which we are most grateful.

4	Foreword
6	Introduction
15	State Profiles
16	Profiles in alphabetical order
119	Resources
120	Appendix 1: States Ranked by Indicator, 1998
124	Appendix 2: Definitions, Data Sources, and Reporting Issues
129	Appendix 3: Primary Contacts for State KIDS COUNT Projects

Foreword

What constitutes “the right start” to life? Surely a healthy birth into a family ready and able to meet children’s needs is at the top of the list.

In *The Right Start State Trends*, Child Trends and KIDS COUNT report how children born in each state fared on eight selected measures of a “right start.” The data are provided for each year from 1990 to 1998 and data for the U.S. as a whole are presented to provide a point of comparison. These measures include some of the factors that medical science tells us promote a healthy birth, including early prenatal care and a tobacco-free pregnancy, as well as health conditions that place a baby at risk, such as prematurity or low birthweight. We also include measures of circumstances that we know from social science research potentially limit a child’s future prospects. These include birth to a teen mother, to an unmarried mother, and to a mother with limited education. A companion volume, *The Right Start City Trends*, reports the same data for the nation’s 50 largest cities.

The U.S. made progress on some of these measures during the 1990s, despite some continued barriers to health care, persistent racial gaps in birth outcomes, and a teen birth rate which, while declining, still far exceeds rates in other industrialized nations. The most notable signs of progress were:

- A drop in the percentage of mothers in the U.S. who reported getting late or no prenatal care, from 6 percent in 1990 to 4 percent in 1998.
 - A drop in the percentage of mothers who reported smoking during pregnancy, from 18 percent in 1990 to 13 percent in 1998. Smoking during pregnancy is associated with low birthweight and infant mortality, as well as negative consequences for child health and development.
- Modest progress occurred in two other measures of a right start, specifically:
- A decline in “repeat teen births” – the percentage of teen births that were to young women who had already borne a child earlier in their adolescence – from 24 percent in 1990 to 22 percent in 1998. A single birth to a teen presents serious concerns for the future of both the child and the mother. A second birth only compounds these concerns.
 - A decline in the share of births to mothers with less than 12 years of education, from 24 percent in 1990 to 22 percent in 1998. Research has consistently shown that a mother’s educational level is associated with

several child outcomes, including infant mortality and a child’s educational attainment.

Finally, the U.S. lost ground on three other measures of a right start:

- The percentage of births that were low-birthweight (less than 5.5 lbs.) increased from 7 percent in 1990 to 8 percent in 1998. Low-birthweight babies are at higher risk of death, disability, and other illness than are babies born at normal weight. The increase in low-birthweight births is partly attributable to the increase in births of twins and higher-order multiple births.
- The percentage of babies that were born preterm, or before the 37th week of pregnancy, increased from 11 percent in 1990 to 12 percent in 1998. Many preterm babies are also low-birthweight and face the same health risks.
- An increase in the percentage of total births to unmarried women, from 28 percent in 1990 to 33 percent in 1998. Most of this increase occurred in the 1990-1994 period, and this figure has been relatively constant the past few years. Children born to unmarried mothers face a greater risk of poor birth outcomes and of poverty than do children born to married mothers.

The Right Start City Trends, the companion to this volume, paints a bleaker picture for babies born in the nation’s largest cities. On every measure presented here, except smoking during pregnancy, babies born in urban America fare much worse than those born in the rest of the nation. This contrast compels us to direct our attention and resources to the inner-city neighborhoods that have less to offer children in the way of a healthy and promising start.

Toward this end, the Annie E. Casey Foundation centers its work on the proposition that children do best when their families do well, and families do better when they live in supportive neighborhoods. The Foundation will continue its work with partners in cities across America to rebuild neighborhoods that support families raising children. Child Trends supports this effort through its work with KIDS COUNT and through its commitment to rigorous, nonpartisan research and data on children and families.


Improving birth outcomes and changing the circumstances of pregnancy clearly require a mix of effective public policies, improved services, and

Foreword

responsible personal choices. As this report illustrates, for each measure of a healthy and promising start to life, there is a diverse group of states that outperforms the nation as a whole, setting a standard for other states to meet. Further progress is well within our reach.

Douglas W. Nelson
President
The Annie E. Casey Foundation

Kristin A. Moore
President
Child Trends



INTRODUCTION

16

17

Introduction

In 1999, the Annie E. Casey Foundation published *The Right Start: Conditions of Babies and Their Families in America's Largest Cities*, which provided 1997 city-level data for ten measures that characterize the well-being of newborns in the United States. Similar data for states were presented in the Appendix of that report.

In an effort to increase the usefulness of these indicators, we have expanded the original *Right Start Data Book* in several ways. First, we updated the data through 1998, the most recent year available. Second, we have provided annual data for each year back to 1990. Third, we have provided a one-page narrative description of how each state fared during the 1990-1998 period. Fourth, we are publishing a separate companion book focused on city-level data.

The remainder of this introduction focuses on the data contained in *THE Right Start: State Trends: Conditions of Babies and Their Families Across the Nation (1990-1998)*. There is a similar introduction in the city volume which focuses on the data contained in that publication. In addition, we present the value of each indicator for the top 50 cities, five additional cities, and the 50-city average in a companion City volume titled *The Right Start City Trends: Conditions of Babies and Their Families in America's Largest Cities (1990-1998)*.

States were selected as the focus of one of the *Right Start* Data Books because KIDS COUNT Data Books have traditionally looked at states. States are also used as a unit of analysis because they are a prime locus of policy-making for many of the programs that make a difference in birth outcomes. State decision-making power has increased significantly during the "devolution" era of the 1990s. Finally, states are large enough to provide reliable measures but small enough to reflect significant national differences.

To place our state-level results in context, we also present the value of each indicator for the U.S. as a whole. In addition, we present the value of each indicator for the largest 50 cities in the United States in the companion City volume cited above.

We selected indicators that describe the well-being of infants at birth because conditions at birth often reflect the forces that will shape a young person's life. Indicators such as lack of timely prenatal care and smoking during pregnancy reflect social conditions prior to birth that can affect the health of an infant. Other measures included, such as birthweight and gestation period, reflect health status at the time of birth. Finally, we included three characteristics of the mother (marital status, age, and

education) that are likely to reflect conditions a newborn might experience early in life.

We utilize birth certificate data compiled and provided by the National Center for Health Statistics (NCHS) because they provide one of the few sets of systematic measures reflecting child well-being that are available consistently for all states. From the birth certificate data we were able to construct eight measures, expressed as proportions or percents, which reflect some dimension of well-being:

- 1) Teen Births
- 2) Repeat Teen Births
- 3) Births to Unmarried Women
- 4) Low Maternal Education
- 5) Late or No Prenatal Care
- 6) Smoking During Pregnancy
- 7) Low-Birthweight Births
- 8) Preterm Births

Findings

This compilation of nine years of data has revealed a number of interesting patterns for the 50 states plus the District of Columbia and for the nation as a whole. In particular, there are broad differences among the states for a number of indicators. State-level socioeconomic, demographic, and economic indicators might offer some explanations for why certain states consistently performed better on many of the indicators, while others typically did worse.

It is important to recognize that the statewide numbers presented here may mask important variations within a state. Studies suggest that negative birth outcomes are part of a constellation of measures that point toward particular neighborhoods with concentrated poverty and diminished opportunity.¹

While these measures can hardly capture the full range of forces shaping the lives of newborns, the indicators used in this series reflect several important dimensions of a newborn's life. Moreover, these indicators are consistently measured across all of the states and over time, permitting legitimate comparisons. Since many of the conditions related to a birth are linked to

Introduction

later developmental problems, the data illuminate future prospects for children in these states.

The key indicators of birth outcomes used here are all taken from data compiled by NCHS and reflect the official data for each indicator. While these measures are not derived from samples, many are based on relatively small numbers of births and therefore may exhibit a degree of random fluctuation from year to year. Since small differences among states may reflect random fluctuations rather than “real” distinctions in the well-being of children, we urge readers to focus on those differences and changes over time that are relatively large.

In the following pages we describe each of the measures in more detail, explain why each measure was selected as an indicator of well-being, and discuss how the measure is related to broad, long-term outcomes.

Teen Births

Teenage childbearing is problematic because it is associated with diminished opportunities for both the child and the young mother. Teen births are particularly troublesome because most of these mothers are unmarried, and a large segment have not completed high school. For example, nationally, only 21 percent of teenagers who gave birth in 1998 were married, and only 38 percent had completed 12 or more years of education.^{2,3}

If a large share of the births in a state is occurring to teenagers, it means that a significant number or proportion of children are starting life with a parent who is unlikely to have the resources needed to provide for a child. Most teenage mothers are not settled in a job or career, and many young fathers are not able to help. Data from the Census Bureau indicate that only 10 percent of mothers ages 15 to 17 received child-support payments in 1997.⁴ Data from the March 2000 Current Population Survey show that only 58 percent of all males ages 16 to 19 had any earned income in 1999 and that the average annual income for those who worked was slightly less than \$6,000.⁵

Teenage mothers are also more likely to receive late or no prenatal care, although timely prenatal care by teens did increase over the 1990s.⁶ They also are more likely to smoke during pregnancy than older mothers. Moreover, unlike other age groups, the percentage of teen mothers who smoke has increased since 1994.⁷

Children born to teenage mothers are less likely to obtain the emotional and financial resources they need to develop into independent, productive, well-adjusted adults. Thus, babies born to teens reflect a group of children who will have to overcome high odds to thrive.

In 1998, 13 percent of all births in the U.S. occurred to teenagers. Massachusetts had the lowest percentage of total births to teenagers (7 percent), while Mississippi had the highest (20 percent). Nationally, the share of births to teenagers stayed nearly constant during the 1990s (13 percent in both 1990 and 1998; see Table 1), although birth rates for teens dropped sharply during the decade.^{8, 9, 10} It is important to note that the percentage of total births to teenagers is influenced by the fertility of older women (above age 20) as well as by the childbearing patterns of teens.

Repeat Teen Births

Most teen mothers are ill equipped to provide for one child, and a second one severely compounds that challenge. Therefore, children born to a teenage mother who already has one or more children are unlikely to receive the kinds of support that children need to thrive. Moreover, a high rate of repeat teen births signals a problem with pregnancy prevention programs and offers a key opportunity for policy or program intervention. Twenty-two percent of all births to teens were repeat births in 1998. The percentage of teen births to young women who were already mothers ranged from a low of 12 percent in North Dakota to a high of 24 percent in Mississippi. Throughout the U.S., the percentage of teen births that were repeat births decreased from 24 percent in 1990 to 22 percent in 1998 (see Table 1).

Births to Unmarried Women

Research shows that children growing up with a single mother “are more likely to drop out of school, to give birth out of wedlock, to divorce or separate, and to be dependent on welfare.”¹¹ Numerous recent studies document the importance of fathers in the lives of their children. “Children develop best when they are provided with the opportunity to have warm, intimate, continuous, and enduring relationships with both their fathers and their mothers.”¹² Even if a marriage fails, children born into a married couple family have advantages over those born to unmarried women. In 1998, the poverty rate for single-parent families headed by a never-married mother was 55 percent, compared to 35 percent for families headed by a

Introduction

divorced or separated mother. Also, in 1998, the infant mortality rate of children born to an unmarried mother was almost twice that of children born to married mothers (10.2 compared to 5.7 deaths per 1,000 live births).¹³

Moreover, the likelihood of a child receiving a child-support award reflects the marital status of parents at the time of birth. Data from 1997 indicate that among never-married single parents, only 47 percent had a child-support award in place, compared to 70 percent of divorced single parents. But it should be noted that many custodial parents with child-support awards in place never receive the money that they are due. Only 22 percent of never-married single mothers actually received child-support payments in 1997, compared to 47 percent of divorced single parents.¹⁴

Finally, unmarried mothers are more likely to receive inadequate prenatal care than are their married counterparts.¹⁵

Thirty-three percent of all births in the U.S. occurred to unmarried women in 1998. The percentage of total births to unmarried women ranged from a low of 17 percent in Utah to a high of 45 percent in Mississippi. National data show that this proportion increased during the early 1990s, but was stable from 1995 to 1998. The percentage of total births to unmarried women increased from 28 percent in 1990 to 33 percent in 1994 and in 1998 (see Table 1 for 1990 and 1998 data).¹⁶

Low Maternal Education

Research has consistently shown that the education level of a child's mother is a good predictor of many child outcomes.¹⁷ Consequently, children born to women who have not graduated from high school face tough odds. The infant mortality rate for births to women with less than 12 years of education was 9.1 deaths per 1,000 live births in 1998, compared to 6.3 for women with at least a high school education.¹⁸ Women who do not get a good formal education are often less likely to provide the kind of educational and intellectual stimulation that their children need. In addition, parents with less education are less likely to be effective advocates for their children when they enter school or encounter problems with other institutions or public systems.

Finally, mothers with less than 12 years of education are more likely to smoke during pregnancy and to receive inadequate prenatal care.¹⁹

Nationwide in 1998, 22 percent of births occurred to women with low levels of education (less than 12 years). The percentage of total births to mothers

with low levels of education ranged from a low of 10 percent in North Dakota to a high of 33 percent in Texas. Nationally, the percentage of total births to mothers with low levels of education declined slightly during the 1990s, going from 24 percent in 1990 to 22 percent in 1998 (see Table 1).

Late or No Prenatal Care

Mothers who receive timely prenatal care are less likely to have babies with health problems. Failure to obtain early prenatal care may reflect a mother's indifference to her pregnancy or it may reflect a lack of available health care. Either situation is cause for concern. A woman who makes sure that she gets proper prenatal care is also likely to make sure that she does other things to protect her newborn. Failure to find timely prenatal care may also reflect the fact that a woman is in a precarious situation where many other kinds of resources are simply not available.

Nationwide, 4 percent of all births in 1998 occurred to mothers who received late or no prenatal care. Within the U.S., the share of births occurring to women who lacked timely prenatal care in 1998 ranged from a low of 2 percent in Rhode Island to a high of 9 percent in New Mexico.²⁰ During the 1990s, the U.S. as a whole improved on this statistic. The share of mothers who received late or no prenatal care fell from 6 percent in 1990 to 4 percent in 1998 (see Table 1).

Smoking During Pregnancy

Babies born to mothers who smoked during pregnancy are more likely to have health problems. "Smoking during pregnancy is associated with adverse outcomes, including low-birthweight, intrauterine growth retardation and infant mortality as well as negative consequences for child health and development."²¹ Moreover, smoking during pregnancy may be symptomatic of other conditions that reflect an unhealthy approach to pregnancy and childbearing.

Data on smoking during pregnancy from birth certificates in California, New York, Indiana, and South Dakota were either not available or not compatible with NCHS standards. Therefore, these data are missing for these states, and the figure shown under the U.S. Average reflects only the 46 states, plus New York City and the District of Columbia where these data were collected.

Introduction

Thirteen percent of all births in the U.S. were to mothers who smoked during pregnancy in 1998. Among the 47 states that collect these data, the rates ranged from a low of 7 percent in the Texas to a high of 26 percent in West Virginia. Nationally, the share of babies born to mothers who smoked during pregnancy has been declining, dropping from 18 percent in 1990 to 13 percent in 1998 (see Table 1).

Low-Birthweight Births (Less than 5.5 pounds)

While most American children get off to a healthy start, babies weighing less than 2,500 grams (about 5.5 pounds) at birth have a high probability of experiencing developmental problems, suffering from serious illnesses, and dying in the first year of life.^{22, 23} Therefore, the percentage of low-birthweight births reflects a group of children who are more likely to have health problems as they move through the growth stages than are children born at a normal weight.

Nationally, 298,208 babies were born weighing less than 5.5 pounds in 1998, accounting for 8 percent of all births. The relatively high rate of low-birthweight births in the U.S. raises a number of troubling issues. Research shows that women who do not receive adequate early prenatal care are more likely to give birth to a low-birthweight baby and that mothers who lack health insurance are less likely to seek and obtain prenatal care. According to Census Bureau data for 1999, more than one-fifth (21 percent) of all male and female Hispanics and more than one-fifth (21 percent) of all male and female black non-Hispanics did not have health insurance. People living in poverty, high school dropouts, and young adults (ages 18 to 24) are among those that are the least likely to have health insurance.²⁴ Among women of childbearing age (15 to 44), 25 percent of those living in central cities lacked health insurance, compared to 16 percent of those living in the suburbs. Among Hispanic women of childbearing age living in central cities, 41 percent lacked health insurance.²⁵

Nearly 8 percent of the total births in 1998 in the U.S. were low-birthweight births. In 1998, Oregon had the lowest rate of low-birthweight births at 5 percent, and Louisiana and Mississippi were tied with the highest rate of 10 percent. During the 1990s, the share of babies born weighing less than 5.5 pounds increased from 7 percent of all births in 1990 to nearly 8 percent in 1998 (see Table 1).

Preterm Births (Less Than 37 Completed Weeks of Gestation)

Babies born preterm often suffer related health and development problems; preterm births are often also of low-birthweight. A shortened gestation period is also related to an increased risk of infant death (death in the first year of life).²⁶

Nationwide, 12 percent of births in 1998 were preterm births. New Hampshire had the lowest percentage of births that were preterm in 1998 at 8 percent, while Mississippi had the highest rate at 16 percent. Nationally, the share of babies who were preterm increased from 11 percent in 1990 to 12 percent in 1998 (see Table 1).

Race and Hispanic Origin

For every measure that characterizes the well-being of U.S. newborns in this volume, there are substantial differences by race and Hispanic origin. With the exception of smoking during pregnancy, births to (non-Hispanic) whites have a lower value for each measure than births to (non-Hispanic) blacks (see Table 2). Values for Hispanics often, but not always, fall in between the values for whites and blacks. In 1998, for example, 9 percent of U.S. births to whites were to teen mothers, compared with 22 percent of births to blacks and 17 percent of births to Hispanics. Given these racial and ethnic differences, it is not surprising to note that differences seen among states are often related to differences in racial and ethnic composition in the states. States in which black and/or Hispanic births are a large percentage of total births tend to have higher values for most measures than states in which most births are to whites.

At the same time, it is important to recognize that many of the differences between whites and minorities reflect differences in things such as educational attainment, income, and the availability of high-quality services. For example, 9 percent of U.S. births to women with less than 12 years of education are low-birthweight, compared with only 6 percent of births to women with at least 16 years of education. Since 30 percent of white births are to women with at least 16 years of education, compared with 11 percent for blacks, part of the apparent racial difference in low-birthweight births is due to differences in the educational attainment of white and black mothers.

Introduction

Birth outcomes are often examined by race and Hispanic origin because they are easily available. Questions about race and Hispanic origin status are included on births certificates in all 50 states and the District of Columbia, while many important measures of socioeconomic status (e.g., family income) are not. Consequently, while racial and ethnic differences are easily calculated it is not possible to tabulate the measures of well-being used in this publication by critical measures such as the mother's poverty status.

In short, the racial and ethnic differences in birth outcomes are real, but it is important to recognize that many of those differences actually reflect differences in things like income, education and availability of effective services.

27

28

TABLE 1

U.S. Birth Outcomes: 1990 and 1998

Indicator	Percent		Number of states with statistically significant changes from 1990 to 1998	
	1990	1998	Increase	No change Decrease
Percent of total births to teens	13	13	13	14 24
Percent of teen births to women who were already mothers**	24	22	0	10 40
Percent of total births to unmarried women*	28	33	49	0 1
Percent of total births to mothers with less than 12 years of education*	24	22	9	5 35
Percent of total births to mothers receiving late or no prenatal care	6	4	2	5 44
Percent of total births to mothers who smoked during pregnancy***	18	13	0	1 45
Percent low-birthweight births (less than 5.5 pounds)	7	8	45	5 1
Percent preterm births (less than 37 completed weeks of gestation)	11	12	47	2 2

SOURCES: For 1990 data: "Advance Report of Final Natality Statistics, 1990." Monthly Vital Statistics Report, Vol. 41, No. 9, suppl. Hyattsville, MD: National Center for Health Statistics. For 1998 data: Ventura, S.J., Martin, J.A., Curtin, S.C., Mathews, J.J., Park, M.M. (2000), "Births: Final Data for 1998," *National Vital Statistics Reports*, Vol. 48, No. 3. Hyattsville, MD: National Center for Health Statistics.

*Not all states are included in the 1990 U.S. average for this indicator because data were not reliable for every state.

**Not all states are included in the 1998 U.S. average for this indicator because data were not reliable for every state.

***Not all states are included in the 1990 and 1998 averages for this indicator because data were not collected for every state.

TABLE 2
Percentage of U.S. births with selected characteristics by race and Hispanic origin, 1998

Race/ethnicity	Teen births	Repeat teen births	Births to unmarried women	Low maternal education	Late or no prenatal care	Smoking during pregnancy	Low-birthweight births	Preterm births
White non-Hispanic	9	18	22	13	2	16	7	10
Black non-Hispanic	22	27	69	27	7	10	13	18
Hispanic	17	24	42	49	6	4	6	11

SOURCES: Ventura, S.J., Martin, J.A., Curtin, S.C., Mathews, J.J., and Park, M.M. (2000), "Births: Final Data for 1998," *National Vital Statistics Reports*, Vol. 48, No. 3. Hyattsville, MD: National Center for Health Statistics. Curtin, S.C. and Martin, J.A. (2000), "Births: Preliminary Data for 1999," *National Vital Statistics Reports*, Vol. 48, No. 14. Hyattsville, MD: National Center for Health Statistics.

Introduction

- ¹ Coulton, C., and Pandey, S. (1992), "Geographic Concentration of Poverty and Risk to Children in Urban Neighborhoods," *American Behavioral Scientist*, Vol. 35, No.3. O'Campo, P., Xue, X., Wang, M., and Caughey, M.O. (1997), "Neighborhood Risk Factors for Low Birthweight in Baltimore: A Multilevel Analysis," *American Journal of Public Health*, Vol. 87, No. 7.
- ² Curtin, S.C., and Martin, J.A. (2000), "Births: Preliminary Data for 1999," *National Vital Statistics Reports*, Vol. 48, No. 14. Hyattsville, MD: National Center for Health Statistics.
- ³ Ventura, S.J., Martin, J.A., Curtin, S.C., Mathews, J.J., Park, M.M. (2000), "Births: Final Data for 1998," *National Vital Statistics Reports*, Vol. 48, No. 3. Hyattsville, MD: National Center for Health Statistics.
- ⁴ U.S. Census Bureau (2000), "Child Support 1997," October 13, 2000, <<http://www.census.gov/hhes/www/childsupport/97tables/tab4w.html>>.
- ⁵ Child Trends (2001), Special tabulations of the March 2000 Current Population Survey.
- ⁶ Ventura, S.J., Curtin, S.C., and Mathews, T.J. (2000), "Variations in Teenage Birth Rates, 1991-98: National and State Trends," *National Vital Statistics Reports*, Vol. 48, No. 6. Hyattsville, MD: National Center for Health Statistics.
- ⁷ Ibid.
- ⁸ Curtin, S.C., and Martin, J.A. (2000), "Births: Preliminary Data for 1999," *National Vital Statistics Reports*, Vol. 48, No. 14. Hyattsville, MD: National Center for Health Statistics.
- ⁹ Ventura, S.J., Martin, J.A., Curtin, S.C., Mathews, J.J., Park, M.M. (2000), "Births: Final Data for 1998," *National Vital Statistics Reports*, Vol. 48, No. 3. Hyattsville, MD: National Center for Health Statistics.
- ¹⁰ Ventura, S.J., Curtin, S.C., and Mathews, T.J. (2000), "Variations in Teenage Birth Rates, 1991-98: National and State Trends," *National Vital Statistics Reports*, Vol. 48, No. 6. Hyattsville, MD: National Center for Health Statistics.
- ¹¹ Garfinkel, I. and McLanahan, S.S. (1986), *Single Mothers and Their Children*, The Urban Institute, Washington, DC, pp. 1-2.
- ¹² Popenoe, David (1996), *Life Without Father: Compelling New Evidence that Fatherhood and Marriage Are Indispensable for the Good of Children and Society*, The Free Press, New York, NY.
- ¹³ Mathews, T.J., Curtin, S.C., and MacDorman, M.F. (2000), "Infant Mortality Statistics from the 1998 Period Linked Birth/Infant Death Data Set," *National Vital Statistics Reports*, Vol. 48, No. 12. Hyattsville, MD: National Center for Health Statistics.

- ¹⁴ Grail, Timothy. U.S. Census Bureau (2000), "Child Support for Custodial Mothers and Fathers," October 2000, <<http://www.census.gov/prod/2000pubs/p60-212.pdf>>.
- ¹⁵ Ventura, S.J. (1995), "Births to Unmarried Mothers: United States, 1980-1992," *Vital and Health Statistics*, Vol. 21, No. 53. Hyattsville, MD: National Center for Health Statistics.
- ¹⁶ Ventura, S.J., Bachrach, C.A. (2000), "Nonmarital Childbearing in the United States, 1940-1999," *National Vital Statistics Reports*, Vol. 48, No. 16. Hyattsville, Maryland: National Center for Health Statistics.
- ¹⁷ Haveman, R.B. and Wolfe, B. (1995), "The Determinants of Children's Attainments: A Review of Methods and Findings," *Journal of Economic Literature*, Vol. 33, No. 4.
- ¹⁸ Mathews, T.J., Curtin, S.C., and MacDorman, M.F. (2000), "Infant Mortality Statistics from the 1998 Period Linked Birth/Infant Death Data Set," *National Vital Statistics Reports*, Vol. 48, No. 12. Hyattsville, MD: National Center for Health Statistics.
- ¹⁹ Ventura, S.J., Martin, J.A., Curtin, S.C., Mathews, J.J., Park, M.M. (2000), "Births: Final Data for 1998," *National Vital Statistics Reports*, Vol. 48, No. 3. Hyattsville, MD: National Center for Health Statistics.
- ²⁰ Ibid.
- ²¹ Mathews, T.J. (1998), "Smoking During Pregnancy, 1990-96," *National Vital Statistics Reports*, Vol. 47, No. 10. Hyattsville, MD: National Center for Health Statistics, p. 1.
- ²² Mathews, T.J., Curtin, S.C., and MacDorman, M.F. (2000), "Infant Mortality Statistics from the 1998 Period Linked Birth/Infant Death Data Set," *National Vital Statistics Reports*, Vol. 48, No. 12. Hyattsville, MD: National Center for Health Statistics.
- ²³ MacDorman, M.F. and Atkinson, J.O. (1999), *National Vital Statistics Reports*, "Infant Mortality Statistics from the 1997 Period Linked Birth/Infant Death Data Set," Vol. 47, No. 23. Hyattsville, MD: National Center for Health Statistics.
- ²⁴ Mills, R.J., U.S. Census Bureau (2000), "Health Insurance Coverage," September 2000, <<http://www.census.gov/prod/2000pubs/p60-211.pdf>>.
- ²⁵ Annie E. Casey Foundation (2000), Special tabulations of the March 2000 Current Population Survey.
- ²⁶ MacDorman, M.F. and Atkinson, J.O. (1999), *National Vital Statistics Reports*, "Infant Mortality Statistics from the 1997 Period Linked Birth/Infant Death Data Set," Vol. 47, No. 23. Hyattsville, MD: National Center for Health Statistics.

STATE PROFILES

34

35

The yearly number of births in Alabama fell from 63,487 in 1990 to 62,074 in 1998. During this time, nearly two-thirds of total births were to whites, and nearly one-third were to blacks.*

During the 1990-1998 period, Alabama experienced changes in all eight measures of a healthy start to life, generally mirroring the trends in the national average. The more significant changes include drops in the percentages of births to women who smoked during pregnancy and to women who received late or no prenatal care, as well as an increase in the share of births to unmarried women. When compared with the 1998 U.S. average, the children of Alabama got off to a less healthy start to life on six of eight measures, including the proportion of births to teenage mothers (see figure).

More specifically:


- **Teen births.** Despite a slight overall drop in the share of births to teens, teen births consistently accounted for a substantially greater share of births in Alabama than in the U.S. overall. In 1998, 17 percent of Alabama births were to teens, compared with a national average of 13 percent.
- **Repeat teen births.** The percentage of teen births to young women who were already mothers fluctuated during the 1990s but underwent an overall decline, mirroring the trend in the U.S. average. Repeat teen births made up 23 percent of Alabama teen births in 1998, compared with 26 percent in 1990.
- **Births to unmarried women.** Births to unmarried women increased from 30 percent of total births in 1990 to 34 percent in 1993 through 1998. The nation as a whole experienced a similar upward trend.
- **Low maternal education.** The share of Alabama births to women with less than 12 years of education fell from 26 percent in 1990 and 1991 to 23 percent in 1998. This change paralleled the drop in the U.S. average.
- **Late or no prenatal care.** Alabama saw a reduction in the percentage of births to women who received late or no prenatal care, from 6 percent in 1990 to 4 percent in 1993, remaining stable thereafter. Alabama matched the national average for this indicator in both 1990 and 1998.
- **Smoking during pregnancy.** There was a notable drop in the proportion of Alabama births to women who smoked during pregnancy,

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

from 17 percent in 1990 to 13 percent in 1998. Alabama generally remained below the national average throughout the nine-year period.

- **Low-birthweight births.** The share of Alabama babies born at a low birthweight increased from 8 percent in 1990 to 9 percent in 1993 through 1998. Throughout this time, low-birthweight births were more common in Alabama than in the nation as a whole.
- **Preterm births.** Preterm births remained notably more prevalent in Alabama than in the U.S. as a whole throughout the 1990s. In 1998, preterm births accounted for 14 percent of Alabama births, compared with 12 percent of births in the nation overall.

In summary, Alabama saw changes in all eight measures of a healthy start to life, generally reflecting changes in the U.S. averages. Improvements include reductions in the percentages of births to women who smoked during pregnancy and to women who received late or no prenatal care. However, Alabama saw increases in the percentages of preterm and low-birthweight births, and in the share of births to unmarried women—three indicators for which the state was already above the respective national averages in 1990. In addition, when compared with the U.S. average, the proportion of births to teens remained high throughout the 1990-1998 period.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	18.2	18.5	18.2	17.9	18.6	18.5	18.4	17.6	17.1
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	26.2	26.4	27.1	24.2	22.1	22.2	23.2	22.6	23.0
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	30.1	31.9	32.6	33.5	34.5	34.5	33.7	33.9	34.1
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	26.0	26.1	25.8	25.4	25.0	24.0	23.9	23.5	23.3
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	6.1	5.3	4.7	4.2	3.9	3.8	3.8	3.7	3.9
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	16.5	16.4	15.1	14.6	13.7	13.4	13.0	12.5	12.6
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	8.4	8.7	8.5	8.7	9.0	9.0	9.3	9.2	9.3
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	13.1	13.2	13.2	13.8	14.1	13.5	13.9	14.3	14.3
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	40,778	40,296	39,774	39,508	39,132	39,073	39,326	40,203
	Black non-Hispanic	21,830	21,616	21,488	21,067	20,615	19,830	19,536	20,021
	Hispanic	344	400	444	509	579	758	936	1,056
	Other	535	498	554	622	613	668	690	505
	Total	63,487	62,810	62,260	61,706	60,939	60,488	60,914	62,074

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

During the 1990s, there was a marked overall drop in the total number of yearly births in Alaska, from 11,902 in 1990 to 9,926 in 1998. At the same time, the small percentage of births to Hispanics doubled, from 3 percent in 1990-1992 to 6 percent in 1995 through 1998.* There were 2,381 births to American Indians, Eskimos, and Aleuts in 1998—nearly one-quarter of Alaska's births.**

During this period, Alaska experienced a marked rise in the share of births to unmarried women and a notable overall drop in the share of births to women who smoked during pregnancy. The percentage of births to women with less than 12 years of education remained well below the national average throughout the 1990s (see figure).

More specifically:


- **Teen births.** Throughout the 1990s, the proportion of Alaska births to teens accounted for a smaller share of births in Alaska than in the nation as a whole. In 1998, 11 percent of Alaska births were to teenagers, compared with 13 percent in the U.S. overall.
- **Repeat teen births.** The small fluctuations during the 1990s in the percentage of Alaska teen births to young women who were already mothers were not significant. At 20 percent in 1998, Alaska was similar to the national average on this indicator.
- **Births to unmarried women.** Alaska saw a notable increase in the share of births to unmarried women, from 26 percent in 1990 to 31 percent in 1996 through 1998. The U.S. as a whole saw a similar upward trend.
- **Low maternal education.** Births to women with less than 12 years of education consistently made up a substantially smaller share of births in Alaska than in the nation as a whole. Thirteen percent of Alaska births in 1998 were to women with less than 12 years of education, compared with a national average of 22 percent.
- **Late or no prenatal care.** The percentage of Alaska births to mothers who received late or no prenatal care fell from 4 percent in 1990 to 3 percent in 1992-1996, and then rose to a high of 5 percent by 1998. Alaska went from below the U.S. average in 1990 to above it by 1998.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** These births account for 85 percent of the "Other" category. Asians and Pacific Islanders account for the rest.

- **Smoking during pregnancy.** The share of births to women who smoked during pregnancy fluctuated during the 1990s but decreased overall, from a high of 23 percent in 1992 to a low of 18 percent in 1998. Smoking during pregnancy remained considerably more common in Alaska than in the nation overall throughout the 1990-1998 period.
- **Low-birthweight births.** Despite a rise in the percentage of Alaska babies born at a low birthweight—from 5 percent in 1990 through 1993 to 6 percent in 1997 and 1998—low-birthweight births were consistently less prevalent in Alaska than in the U.S. as a whole.
- **Preterm births.** The proportion of births that were preterm in Alaska fluctuated between 8 percent and 10 percent during the 1990s. Alaska remained below the national average on this measure throughout this time.

In summary, Alaska saw an increase in the percentage of births to unmarried women and an overall decrease in the share of births to women who smoked during pregnancy. While the proportion of births to mothers with less than 12 years of education remained well below the U.S. average during the 1990s, the share of births to women who smoked during pregnancy was considerably greater than the national average throughout this time.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens									
Alaska	9.7	10.5	10.9	10.8	11.5	11.2	11.1	11.2	11.2
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers									
Alaska	21.1	21.8	20.5	21.5	19.7	17.9	18.5	19.3	20.1
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women									
Alaska	26.2	26.9	27.4	28.0	29.3	29.9	31.0	30.6	31.1
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education									
Alaska	14.7	14.6	15.1	14.8	14.9	13.4	13.9	14.2	13.2
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care									
Alaska	3.8	3.6	3.0	2.8	2.8	3.3	3.3	4.4	4.5
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy									
Alaska	21.9	23.2	23.3	21.6	21.6	19.2	20.5	19.3	18.4
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)									
Alaska	4.8	4.7	4.9	4.9	5.5	5.3	5.5	5.9	6.0
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)									
Alaska	9.0	8.6	9.3	9.3	9.1	8.0	10.1	10.2	10.0
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother									
White non-Hispanic	7,779	7,561	7,590	7,106	6,897	6,581	6,300	6,110	6,148
Black non-Hispanic	529	507	531	570	461	421	387	444	386
Hispanic	328	367	373	439	580	574	651	609	593
Other	3,266	3,251	3,232	2,958	2,740	2,668	2,699	2,784	2,799
Total	11,902	11,686	11,726	11,073	10,678	10,244	10,037	9,947	9,926

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

There was a marked increase in the yearly number of births in Arizona during the 1990s, from 68,995 in 1990 to 78,243 in 1998. While births to both whites and blacks decreased from 1990 to 1998, births to Hispanics increased dramatically.* Hispanics accounted for 38 percent of Arizona births in 1998, compared with 29 percent in 1990.

During this nine-year span, Arizona experienced a sharp reduction in the percentage of births to mothers who smoked during pregnancy and a substantial increase in the proportion of births to unmarried women. The share of births to women who received late or no prenatal care declined in the 1990s, but was still higher than the national average (see figure).

More specifically:

- **Teen births.** Throughout the 1990-1998 period, teenagers accounted for a larger proportion of births in Arizona than in the nation as a whole. In 1998, 15 percent of Arizona births were to teens, compared with a national average of 13 percent.
- **Repeat teen births.** The percentage of teen births to young women who already had a child fluctuated somewhat during the 1990s, but underwent an overall decrease, from 25 percent in 1990 through 1993 to 22 percent in 1998. Arizona was on par with the U.S. average in 1998.
- **Births to unmarried women.** Arizona saw a substantial increase in the proportion of births to unmarried women during the 1990s, from 33 percent in 1990 to 38 percent in 1998. Throughout this time, the percentage of Arizona births to unmarried women remained above the national average, which exhibited a parallel upward trend.
- **Low maternal education.** Births to women with less than 12 years of education consistently made up a larger share of births in Arizona than in the U.S. as a whole. In 1998, 30 percent of Arizona births were to women with less than 12 years of education, compared with the U.S. average of 22 percent.


- **Late or no prenatal care.** Arizona experienced a reduction in the percentage of births that were to women who received late or no prenatal care during the 1990s, from 10 percent in 1990 to 7 percent in 1998. The U.S. as a whole also saw a decline on this measure, and

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

Arizona remained above the national average throughout the 1990-1998 period.

- **Smoking during pregnancy.** The percentage of Arizona births to mothers who smoked during pregnancy fell from 15 percent in 1990 to less than 8 percent in 1998. This dramatic decline outpaced the downward trend for the nation as a whole. Throughout the 1990s, the proportion of Arizona births to women who smoked during pregnancy was lower than the national average.
- **Low-birthweight births.** The share of Arizona babies born at a low birthweight remained just below the national average throughout the 1990-1998 period. In 1998, 7 percent of births in Arizona were low-birthweight, compared with 8 percent in the U.S. as a whole.
- **Preterm births.** The proportion of Arizona births that were preterm fluctuated between 10 percent and 11 percent during the 1990s. Throughout this time, Arizona was near the U.S. average for this indicator.

In summary, Arizona experienced a striking reduction in the percentage of births to women who smoked during pregnancy, and was below the national average on this indicator throughout the 1990s. However, the proportion of births to women who received late or no prenatal care remained substantially higher than the U.S. average throughout the 1990-1998 period. An increase in the share of births to unmarried women, already above the national average in 1990, is also noteworthy.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	14.2	14.8	15.0	15.1	15.3	15.1	14.9	14.9	15.1	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Arizona.	25.2	24.7	25.0	24.5	22.5	20.8	21.8	22.7	22.0	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Arizona	32.7	35.1	36.2	37.9	38.3	38.2	38.8	37.6	38.4	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Arizona	28.8	30.6	33.3	30.1	29.3	29.2	29.5	29.3	29.5	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Arizona	10.1	9.3	8.1	8.5	7.4	8.2	7.4	6.6	7.2	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Arizona	15.0	14.3	12.6	12.4	12.2	11.5	10.3	8.7	7.6	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Arizona	6.4	6.4	6.4	6.7	6.8	6.8	6.7	6.9	6.8	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Arizona	10.2	10.2	10.6	11.1	11.3	11.3	10.3	11.2	11.4	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Percent preterm births (less than 37 completed weeks of gestation)	White non-Hispanic	39,775	38,214	37,790	36,517	36,597	38,474	38,602	37,860	38,621
	Black non-Hispanic	2,527	2,482	2,399	2,272	2,322	2,166	2,249	2,375	2,465
	Hispanic	19,663	20,552	21,862	22,579	24,135	25,504	27,901	28,472	29,682
	Other	7,030	6,861	6,778	7,688	7,792	6,319	6,570	6,992	7,475
	Total	68,995	68,109	68,829	69,056	70,846	72,463	75,322	75,699	78,243
Number of births by Race and Ethnicity of mother	Arizona**									

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



The yearly number of births in Arkansas dropped from 36,457 in 1990 to 34,289 in 1993, and then rose to 36,865 by 1998. In 1998 whites accounted for 72 percent of total births, while blacks accounted for 22 percent. Hispanic births quadrupled in the 1990s and accounted for 5 percent of total births in 1998.*

Over the 1990-1998 period, Arkansas experienced a decrease in the percentage of births to women who smoked during pregnancy (see figure). However, there was a sizable increase in the share of births to unmarried women. Moreover, when compared with the national average, the children of Arkansas got off to a less healthy start to life on all eight measures in 1998.


More specifically:

- **Teen births.** Teen births consistently accounted for a substantially greater share of births in Arkansas than in the U.S. as a whole. In 1998, 19 percent of Arkansas births were to teenage mothers, compared with 13 percent of births in the nation overall.
- **Repeat teen births.** The percentage of Arkansas teen births to young women who were already mothers fluctuated during the 1990s, dropping from a high of 26 percent in 1991 to a low of 21 percent in 1995, and then rising to 24 percent in 1997 and 1998. Repeat teen births made up a larger proportion of births in Arkansas than in the U.S. as a whole in 1998.
- **Births to unmarried women.** Arkansas saw a considerable increase in the percentage of births to unmarried women during the 1990s, mirroring the trend in the U.S. average. Unmarried women accounted for 35 percent of Arkansas births in 1998, compared with 29 percent in 1990.
- **Low maternal education.** Births to mothers with less than 12 years of education became a smaller share of Arkansas births during the 1990s, falling from 26 percent in 1990 and 1991 to 23 percent in 1995 through 1998. The nation as a whole experienced a similar downward trend.
- **Late or no prenatal care.** The percentage of births to women who received late or no prenatal care fluctuated during the 1990-1998 period, undergoing an overall decrease from 7 percent in 1990 to 5 percent in

1998. Throughout this time, Arkansas was above the national average for this indicator.

- **Smoking during pregnancy.** The proportion of Arkansas births to women who smoked during pregnancy dropped from 23 percent in 1990 to 19 percent in 1995-1998. There was a stronger downward trend in the national average, and smoking during pregnancy remained strikingly more common in Arkansas than in the U.S. as a whole.
- **Low-birthweight births.** The share of Arkansas births that were low-birthweight remained just above the U.S. average throughout the 1990-1998 period. In 1998, low-birthweight babies accounted for 9 percent of births in Arkansas, compared with 8 percent in the U.S. as a whole.
- **Preterm births.** Preterm births were slightly more prevalent in Arkansas than in the nation overall throughout the 1990s. In 1998, 13 percent of Arkansas births were preterm, compared with a national average of 12 percent.

In summary, Arkansas' children got off to a less healthy start to life on all eight measures, when compared with the 1998 U.S. average. In addition, there was a marked rise in the proportion of births to unmarried women. However, Arkansas saw notable reductions in the proportions of births to women with less than 12 years of education and to women who smoked during pregnancy. There were also declines in the percentages of teen births that were repeat births and births to mothers who received late or no prenatal care.



* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	19.7	19.9	19.4	19.4	20.1	19.6	19.8	19.2	18.6
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	25.9	26.5	26.0	22.9	22.8	20.7	23.1	23.7	23.5
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	29.4	29.9	31.0	31.7	32.6	32.9	33.9	34.2	35.0
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	25.8	25.9	25.0	24.0	23.6	23.4	22.9	23.0	22.8
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	7.0	6.8	6.5	5.7	6.1	6.3	6.8	5.8	5.1
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	22.6	22.0	21.1	20.5	20.0	19.3	19.3	18.6	19.0
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	8.2	8.2	8.2	8.2	8.2	8.2	8.5	8.4	8.9
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	12.5	12.8	12.2	12.6	12.7	12.0	12.2	12.5	13.3
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	27,055	26,425	25,828	25,403	25,594	26,572	26,506	26,545
	Black non-Hispanic	8,492	8,288	8,133	7,822	7,806	7,648	7,799	7,970
	Hispanic	414	379	472	579	782	1,004	1,542	1,724
	Other	496	387	387	485	536	561	613	626
	Total	36,457	35,479	34,820	34,289	34,718	35,175	36,371	36,478

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

There was a substantial decrease in the yearly number of births in California, from 612,628 in 1990 to 521,661 in 1998. This drop was due to marked reductions in births to both whites and blacks. Births to Hispanics rose slightly during this period.* Hispanics accounted for almost 48 percent of California births in 1998, compared with 40 percent in 1990.

During the 1990s, California experienced a sharp drop in the percentage of births to mothers who received late or no prenatal care. However, the share of births to California mothers with less than 12 years of education remained above the U.S. average throughout this time (see figure).

More specifically:

- **Teen births.** Teen births accounted for 12 percent of California births in 1990 through 1997, and slightly more than 11 percent in 1998. Throughout this time, California remained below the national average, where a steady 13 percent of births were to teen mothers.
- **Repeat teen births.** During the 1990s, California saw an overall drop in the proportion of teen births to young women who were already mothers, from 23 percent in 1990 and 1991 to 21 percent in 1998. California remained close to the U.S. average on this indicator throughout the nine-year period.
- **Births to unmarried women.** The trends in the proportion of births to unmarried women in the 1990s reflect, at least in part, changes in coding practices in the state of California.** In 1998, California was on par with the national average of 33 percent.
- **Low maternal education.** From 1990 to 1998, both California and the nation as a whole experienced a decline in the proportion of births to women with less than 12 years of education. However, California was substantially above the U.S. average throughout this time. In 1998, births to women with less than 12 years of education accounted for 31 percent of California births, compared with 22 percent of births in the U.S. overall.
- **Late or no prenatal care.** The percentage of births to mothers who received late or no prenatal care in California dropped markedly over the 1990-1998 period, from 7 percent in 1990 to less than 4 percent in


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** In 1995 California changed the method by which it inferred marital status to take into account naming conventions of Hispanics. In 1997, California began asking for mother's marital status directly. For more details, see Appendix 2.

1996 through 1998. This change outpaced the downward trend for the U.S. as a whole.

- **Smoking during pregnancy.** Data on smoking during pregnancy were not available for California births.
- **Low-birthweight births.** During the 1990s, the share of California babies born at a low birthweight remained relatively stable and below the national average. In 1998, 6 percent of California births were low-birthweight, compared with almost 8 percent of births in the nation as a whole.
- **Preterm births.** Preterm births accounted for 10 percent of California births during the 1990-1998 period. Throughout this time, California was slightly below the U.S. average on this measure.

In summary, California saw declines over the 1990-1998 period in the share of teen births that were repeat births, the proportion of births to women receiving late or no prenatal care, and the percentage of births to mothers with less than 12 years of education. Furthermore, the percentages of teen, low-birthweight, and preterm births in California were below the respective national averages in 1998. The share of births to women with less than 12 years of education, however, remained well above the national average throughout the 1990s.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	11.6	11.8	11.8	12.0	12.3	12.4	12.0	11.7	11.4
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	22.6	23.0	23.8	23.3	22.1	21.4	21.8	21.8	21.2
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	31.6	33.5	34.3	35.3	35.7	32.1	31.4	32.8	32.8
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	34.0	35.1	35.4	35.0	34.8	33.9	33.5	32.2	31.0
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	7.0	6.4	5.3	5.0	4.9	5.2	3.8	3.7	3.6
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	5.8	5.8	5.9	6.0	6.2	6.1	6.1	6.2	6.2
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	9.8	9.8	9.7	10.0	10.0	10.0	9.9	10.1	10.3
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	255,913	242,274	228,252	213,326	204,476	196,695	185,659	177,863
	Black non-Hispanic	48,004	46,756	45,510	43,497	41,510	39,284	37,133	36,109
	Hispanic	245,592	258,005	263,525	262,313	257,750	254,001	254,895	248,948
	Other	63,119	63,042	64,443	66,188	64,194	62,065	61,746	61,920
	Total	612,628	610,077	601,730	585,324	567,930	552,045	539,433	524,840

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**See text, footnote ** in text, and Appendix 2.

***N.A. = Not Available. See Appendix 2 for more details.

****The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Colorado increased from 53,525 in 1990 to 59,577 in 1998. This rise was due to a substantial increase in births to Hispanics, who accounted for almost 25 percent of Colorado births in 1998, compared with 17 percent in 1990.*

During the 1990-1998 period, the children of Colorado got off to a healthier start to life on several measures, when compared with the national average. In addition, there were two striking changes during this time—a dramatic drop in the proportion of births to mothers who smoked during pregnancy and a sizable increase in the percentage of births to unmarried women (see figure).


More specifically:

- **Teen births.** Teen births made up a slightly smaller share of births in Colorado than in the U.S. as a whole throughout the 1990s. Births to teens accounted for between 11 percent and 12 percent of Colorado births over the 1990-1998 period, compared with the steady national average of 13 percent.
- **Repeat teen births.** The percentage of Colorado teen births to young women who already had a child fluctuated somewhat during the 1990s, but underwent an overall decline from 22 percent in 1990 to 20 percent in 1998. Colorado was below the U.S. average on this measure throughout this time.
- **Births to unmarried women.** Over the 1990-1998 period, the proportion of Colorado births to unmarried women increased from 21 percent in 1990 to almost 26 percent in 1998. This overall increase from 1990 to 1998 paralleled a similar trend for the U.S. as a whole, although births to unmarried women were less common in Colorado than in the U.S. as a whole.
- **Low maternal education.** Between 18 percent and 19 percent of Colorado births during the 1990s were to women with less than 12 years of education. Colorado was consistently below the national average on this indicator throughout the nine-year period.
- **Late or no prenatal care.** The percentage of births to women who received late or no prenatal care remained stable at 5 percent from 1990 to 1995, and then decreased slightly to 4 percent thereafter. In 1998, Colorado was just above the U.S. average on this measure.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** During the 1990s, Colorado experienced a remarkable drop in the percentage of births to mothers who smoked during pregnancy, from 18 percent in 1990 to 11 percent in 1998. This change outpaced the downward trend for the nation as a whole, leaving Colorado below the U.S. average throughout most of the nine-year period.
- **Low-birthweight births.** The proportion of babies born at a low birthweight in Colorado was just above the U.S. average throughout the 1990s. In 1998, low-birthweight births accounted for 9 percent of Colorado births, compared with 8 percent of births in the nation overall.
- **Preterm births.** Preterm births made up a slightly smaller share of births in Colorado than in the nation overall throughout the 1990s. In 1998, 11 percent of Colorado births were preterm, compared with 12 percent of births in the U.S. as a whole.

In summary, when compared with the U.S. average, Colorado's children got off to a healthier start to life on a number of indicators during the 1990s. Moreover, the state experienced a dramatic reduction in the proportion of births to women who smoked during pregnancy. However, there was also a notable increase in the percentage of births to unmarried women during this time.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	11.3	11.8	12.0	11.9	12.3	12.1	11.9	11.9	12.1	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	21.8	22.3	22.7	20.3	18.5	17.8	19.5	18.9	20.3	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	21.2	23.6	23.8	24.8	25.0	24.9	24.8	25.2	25.6	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	17.8	18.8	18.2	18.0	18.0	18.1	18.7	18.9	19.4	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	5.0	4.8	4.9	5.1	4.6	5.1	4.5	3.9	4.3	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	18.1	17.8	15.5	14.9	13.3	12.7	12.6	10.9	11.0	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	8.0	8.2	8.5	8.4	8.5	8.4	8.8	8.8	8.6	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	9.6	10.1	10.3	10.2	10.4	10.7	10.7	11.0	11.2	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	39,791	39,487	39,512	39,156	38,930	38,142	38,667	38,685	39,936
	Black non-Hispanic	2,832	2,820	2,919	2,849	2,679	2,518	2,452	2,487	2,761
	Hispanic	9,278	9,803	10,303	10,268	10,667	11,523	12,375	13,217	14,654
	Other	1,624	1,703	1,801	1,749	1,795	2,149	2,313	2,144	2,226
	Total	53,525	53,813	54,535	54,022	54,071	54,332	55,807	56,533	59,577

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Connecticut

The yearly number of births in Connecticut decreased from 50,123 in 1990 to 43,820 in 1998. At the same time, births to Hispanics increased, from 5,318 (11 percent of total births) in 1990 to 6,224 (14 percent of total births) in 1998.*

In 1998, the children of Connecticut got off to a healthier start to life on six of the eight measures, when compared with the U.S. average. Moreover, the state experienced drops between 1990 and 1998 in the percentage of births to women who smoked during pregnancy and in the proportion of teen births to young women who were already mothers. Finally, the share of births to women with less than 12 years of education remained well below the national average throughout the 1990s (see figure).

More specifically:


- **Teen births.** Teen births were strikingly less common in Connecticut than in the U.S. as a whole throughout the 1990-1998 period. In both 1990 and 1998, teenagers accounted for 8 percent of births in Connecticut, compared with 13 percent in the U.S. overall.
- **Repeat teen births.** The proportion of teen births to young women who already had a child fell from a high of 27 percent in 1992 to 21 percent in 1998. The nation as a whole saw a similar downward trend.
- **Births to unmarried women.** Overall, unmarried women accounted for an increasing share of total births in Connecticut during the 1990-1998 period, paralleling the trend for the U.S. as a whole.** Among Connecticut births in 1998, 31 percent were to unmarried women, compared with 27 percent in 1990.
- **Low maternal education.** The proportion of Connecticut births to women with less than 12 years of education dropped from 14 percent in 1990 to 11 percent in 1996 through 1998. In 1998, births to mothers with less than 12 years of education were only half as common in Connecticut as in the U.S. as a whole.
- **Late or no prenatal care.** During the 1990-1998 period, births to women who received late or no prenatal care accounted for a smaller share of births in Connecticut than in the nation as a whole. In 1998, 3 percent of Connecticut births were to mothers who received late or no prenatal care, compared with 4 percent of births in the nation overall.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** Connecticut began determining marital status by direct question in June 1998. Births to unmarried women may be somewhat overstated in all years. See Appendix 2 for more details.

- **Smoking during pregnancy.** Connecticut experienced a reduction in the percentage of births to mothers who smoked during pregnancy, from 13 percent in 1990 to 9 percent in 1998. This change mirrored the downward trend in the U.S. as a whole, and Connecticut remained below the national average on this indicator throughout the nine-year period.
- **Low-birthweight births.** There was a slight overall increase in the proportion of births that were low-birthweight in Connecticut, from 7 percent in 1990 to 8 percent in 1998. Connecticut was on par with the national average for this indicator throughout the 1990-1998 period.
- **Preterm births.** Preterm births were somewhat less prevalent in Connecticut than in the U.S. as a whole throughout the 1990s. In 1998, preterm babies made up 10 percent of births in Connecticut, compared with 12 percent in the U.S. overall.

In summary, when compared with the national average, Connecticut's children got off to a healthier start to life on six of the eight indicators in 1998. The state also saw reductions in the shares of teen births that were repeat births, births to mothers who smoked during pregnancy, and births to mothers with less than 12 years of education. At the same time, there was a rise in the proportion of births to unmarried women.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	8.2	8.2	8.0	8.0	8.5	8.6	8.2	8.3	8.3	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	24.8	24.1	27.0	23.1	19.0	20.0	19.7	20.0	20.7	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	26.6	28.0	28.7	29.8	30.5	30.6	31.3	32.7	31.2	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	13.8	13.0	12.7	12.2	11.8	11.4	11.1	11.3	10.9	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	3.3	2.9	2.6	2.5	2.2	2.5	2.3	1.9	3.0	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	13.4	13.2	12.9	11.4	10.8	10.4	9.9	9.7	9.4	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	6.6	6.9	6.9	6.9	6.9	7.1	7.2	7.3	7.8	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	8.9	9.3	9.1	9.6	9.2	9.7	9.5	10.2	10.3	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	34,207	33,737	33,228	32,664	31,684	30,867	30,617	28,433	28,845
	Black non-Hispanic	5,199	5,451	5,349	5,199	4,836	4,974	4,845	4,896	4,990
	Hispanic	5,318	5,234	5,476	5,381	5,297	5,505	5,651	5,709	6,224
	Other	5,399	4,144	3,520	3,456	3,838	2,988	3,356	4,071	3,761
	Total	50,123	48,566	47,573	46,700	45,655	44,334	44,469	43,109	43,820

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**See footnote ** in text and Appendix 2.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



There was a slight reduction in the yearly number of births in Delaware from 11,113 in 1990 to 10,578 in 1998. All of this decline was accounted for by a decline in births to whites. At the same time, the Hispanic share of total births more than doubled, from 3 percent of births in 1990 to 7 percent of births in 1998.*

Delaware experienced a substantial increase in the share of births to unmarried women (see figure) and a marked decrease, followed by a partially offsetting increase, in the percentage of births to women who smoked during pregnancy. However, Delaware did not differ greatly from the U.S. average for most measures of a healthy start to life in 1998.

More specifically:


- **Teen births.** Teen births made up between 12 percent and 14 percent of Delaware births during the 1990s. In 1998, Delaware was not significantly different than the U.S. average of 13 percent.
- **Repeat teen births.** The percentage of Delaware teen births to young women who were already mothers fluctuated during the 1990s, undergoing an overall drop from 26 percent in 1990 to 22 percent in 1998. This change—most of which occurred between 1992 and 1993—paralleled the overall downward trend for the nation as a whole.
- **Births to unmarried women.** There was a dramatic increase in the share of Delaware births to unmarried women during the 1990s, from 29 percent in 1990 to 37 percent in 1998. This change outpaced the rise in the U.S. as a whole and widened the gap between Delaware and the national average.
- **Low maternal education.** The percentage of births to women with less than 12 years of education fluctuated but rose overall from 16 percent in 1990 to 18 percent in 1998. Despite this increase—and a small decrease in the U.S. average—births to women with less than 12 years of education continued to make up a smaller share of births in Delaware than in the nation as a whole in 1998.
- **Late or no prenatal care.** The share of births in Delaware to women who received late or no prenatal care fell from a high of 5 percent in 1991 to a low of 3 percent in 1995, and then rose to 4 percent in 1998. The U.S. as a whole also saw an overall decline in this measure, and Delaware was even with the U.S. average in 1998.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** The percentage of Delaware births to women who smoked during pregnancy dropped from 20 percent in 1990 to a low of 13 percent in 1995, and then rose to 15 percent in 1998. There was a steadier, sizable downward trend in the national average during the 1990s. Smoking during pregnancy was somewhat more common in Delaware than in the U.S. as a whole in 1998.

- **Low-birthweight births.** Low-birthweight births accounted for 7 to 9 percent of Delaware births during the 1990-1998 period. The state was just above the national average on this indicator in 1998.
- **Preterm births.** Preterm births became a larger proportion of Delaware births over the 1990-1998 period, rising from a low of 10 percent in 1993 to a high of 13 percent in 1998. In 1998, preterm births were somewhat more prevalent in Delaware than in the nation as a whole.

In summary, Delaware saw a considerable increase in the percentage of births to unmarried women, a sharp drop and then partial rebound in the proportion of births to women who smoked during pregnancy, and a decrease in the share of teen births that were repeat births. In general, Delaware did not differ greatly from the national average for most indicators of a healthy beginning in 1998.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	11.9	12.2	12.4	12.5	13.2	13.2	13.7	13.4	13.1	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	26.4	26.1	27.8	23.4	21.1	22.2	20.2	21.1	22.4	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	29.0	31.8	32.6	33.8	34.7	34.9	35.5	36.0	37.1	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	16.2	17.1	18.7	17.8	17.8	17.8	17.7	17.2	18.3	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	4.5	5.2	4.7	3.8	3.6	2.8	3.2	3.3	3.6	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	19.6	18.8	16.4	15.0	14.0	13.2	14.0	14.2	14.9	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	7.6	7.9	7.6	7.8	7.4	8.4	8.5	8.7	8.4	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	11.1	12.2	10.8	10.3	10.6	11.9	12.0	12.3	12.6	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	8,055	7,725	7,504	7,510	7,324	7,134	6,899	6,864	6,937
	Black non-Hispanic	2,564	2,831	2,502	2,363	2,315	2,326	2,348	2,462	2,604
	Hispanic	323	407	426	472	526	585	660	637	753
	Other	171	227	224	223	246	221	248	290	284
	Total	11,113	11,190	10,656	10,568	10,411	10,266	10,155	10,253	10,578

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

District of Columbia

The yearly number of births in the District of Columbia dropped markedly during the 1990-1998 period, from 11,850 in 1990 to 7,686 in 1998. Births to blacks accounted for the vast majority of total births throughout this period, including about three of every four births in 1998.* The drop in births was due primarily to a consistent decline in births to blacks since 1991, but all groups experienced declines.

During this same time period, the District saw improvements on all eight measures of a healthy start to life. The most striking changes were drops in the percentages of births to mothers who smoked during pregnancy and to mothers who received late or no prenatal care (see figure).

More specifically:

- **Teen births.** The proportion of births to teenagers declined from 18 percent in 1990 to 15 percent in 1998. This change brought the District closer to the national average, which remained steady at 13 percent during the 1990s.**
- **Repeat teen births.** The proportion of teen births to young women who were already mothers fluctuated during the nine-year period, from as high as 35 percent in 1993 to as low as 28 percent in 1998. The District remained substantially above the U.S. average throughout the 1990s.**
- **Births to unmarried women.** The percentage of births to unmarried women rose from 65 percent in 1990 to a high of 69 percent in 1994, and then dropped to 63 percent in 1998. Throughout this time, births to unmarried women were markedly more common in the District than in the nation as a whole.**
- **Low maternal education.** Births to women with less than 12 years of education became a smaller share of total births during the 1990s, falling from 31 percent in 1990 to 25 percent in 1998. This change narrowed** the gap between the District and the national average on this indicator.
- **Late or no prenatal care.** There was a striking reduction in the percentage of total births to women who received late or no prenatal care, from 16 percent in 1990 to 10 percent in 1998. This change mirrored the trend in the national average. However, the District remained well above the national average on this measure in 1998.**

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** To compare the District of Columbia with other large cities, see *The Right Start: City Trends*.


- **Smoking during pregnancy.** The District saw a dramatic decline in smoking during pregnancy that surpassed the downward trend in the U.S. as a whole. The percentage of total births in the District to mothers who smoked during pregnancy fell from 16 percent in 1990 to 5 percent in 1998. The District was well below the national average on this indicator throughout the 1990-1998 period.**

- **Low-birthweight births.** The share of low-birthweight babies declined from 15 percent in 1990 to 13 percent in 1998. However, the District remained substantially above the U.S. average throughout the 1990s.**

- **Preterm births.** There was a slight overall decline in the percentage of births that were preterm, from 21 percent in 1990 to 18 percent in 1998. Throughout the 1990-1998 period, preterm births remained notably more common in the District than in the nation as a whole, where 12 percent of births were preterm in 1998.**

In summary, the District of Columbia experienced improvements in all eight indicators during the 1990s, including especially striking reductions in the percentages of births to mothers who smoked during pregnancy and to mothers who received late or no prenatal care. Other sizable changes include drops in the percentages of births to teens and to women with less than 12 years of education.

Despite these important improvements, the District's children got off to a less healthy start to life on seven of eight indicators in 1998, when compared with the U.S. average. When making this comparison, however, it should be recalled that, unlike the 50 states, the District of Columbia is restricted to the city of Washington and contains no suburban or rural areas.**



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	17.8	17.4	16.3	17.4	16.4	16.3	16.7	15.6	15.3
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	31.0	33.0	34.5	34.9	32.1	30.1	31.0	31.0	28.0
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	64.9	66.3	66.9	67.8	68.8	65.8	66.1	63.6	62.9
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	31.2	31.4	30.1	31.9	31.5	28.9	30.1	26.5	25.2
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	15.8	16.3	13.9	15.8	15.4	14.9	11.8	12.3	10.2
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	16.3	13.2	13.0	10.2	9.7	8.3	7.0	5.5	4.8
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	15.1	15.4	14.3	14.6	14.2	13.4	14.3	13.4	13.1
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	20.7	20.5	18.7	20.3	19.9	18.5	18.4	18.3	18.5
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	1,640	1,401	1,319	1,355	1,260	1,354	1,274	1,314
	Black non-Hispanic	9,030	9,185	8,521	8,154	7,618	6,736	6,131	5,435
	Hispanic	890	865	893	930	850	685	777	694
	Other	290	325	227	190	202	239	235	282
	Total	11,850	11,776	10,960	10,629	9,930	9,014	8,390	7,927

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



The yearly number of births in Florida declined from 199,339 in 1990 to 188,723 in 1995, followed by an increase to 195,637 by 1998. The net decrease was due to a large reduction in births to whites that was partially offset by an increase in births to Hispanics. Hispanics accounted for 20 percent of Florida births in 1998, compared with 14 percent in 1990.*

During the 1990s, Florida experienced several improvements on measures of a healthy start to life, including considerable reductions in the percentages of births to mothers who smoked during pregnancy, teen births to young women who were already mothers, and births to women receiving late or no prenatal care (see figure). However, there was also a marked rise in the share of births to unmarried women.


More specifically:

- **Teen births.** Births to teens accounted for between 13 percent and 14 percent of Florida births during the 1990s. Teen births were slightly more common in Florida than in the U.S. as a whole, where a steady 13 percent of births were to teen mothers.
- **Repeat teen births.** There was an overall decline in the percentage of teen births to young women who were already mothers, from a high of 28 percent in 1991 to 22 percent in 1998. Throughout this time, Florida was just above the national average on this measure.
- **Births to unmarried women.** The share of Florida births to unmarried women increased from 32 percent in 1990 to 37 percent in 1998. This change mirrored the upward trend for the nation as a whole, where births to unmarried women were less common than in Florida throughout the 1990s.
- **Low maternal education.** During the 1990s, the percentage of births to women with less than 12 years of education declined steadily, from 25 percent in 1990 to 21 percent in 1997 and 1998. This decrease brought Florida from slightly above the national average in 1990 to just below the average in the latter part of the 1990s.
- **Late or no prenatal care.** The proportion of births to mothers who received late or no prenatal care in Florida was cut in half, from 7 percent in 1990 to 3.5 percent in 1998. Florida was below the U.S. average for this measure from 1992 through 1998.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Births to mothers who smoked during pregnancy became a substantially smaller share of Florida births over the 1990s, decreasing from 18 percent in 1990 to 11 percent in 1998. Florida was just below the U.S. average on this indicator throughout most of the nine-year period.
- **Low-birthweight births.** Low-birthweight babies accounted for between 7 percent and 8 percent of births in Florida during the 1990-1998 period, running slightly higher than in the nation as a whole.
- **Preterm births.** Throughout the 1990s, preterm births made up a slightly larger share of births in Florida than in the U.S. as a whole. In 1998, 13 percent of births in Florida were preterm, compared with the national average of 12 percent.

In summary, during the 1990s, Florida experienced sizable declines in the proportions of births to women who smoked during pregnancy, births to mothers who received late or no prenatal care, and repeat teen births. Over the same time, the state also saw a notable decrease in the proportion of births to mothers with less than 12 years of education. However, there was also a rise in the share of births to unmarried women.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	13.9	13.9	13.5	13.4	13.7	13.7	13.4	13.4	13.2
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	26.6	27.8	27.7	25.0	22.8	22.2	22.6	22.5	22.5
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	31.7	33.0	34.2	35.0	35.7	35.8	35.9	36.0	36.6
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	25.0	24.4	24.0	23.2	22.9	22.3	21.8	21.4	21.3
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	7.1	5.9	4.6	4.0	3.6	3.4	3.3	3.2	3.5
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	18.3	17.0	16.3	15.1	13.8	12.9	12.2	11.4	11.2
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	7.4	7.4	7.4	7.5	7.7	7.7	7.9	8.0	8.1
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	11.6	11.7	11.5	11.7	11.9	11.8	11.9	12.4	12.6
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother									
White non-Hispanic	123,514	118,154	115,131	114,016	111,500	108,831	107,900	107,757	107,754
Black non-Hispanic	45,362	44,681	43,954	43,470	42,299	41,191	41,305	42,546	43,483
Hispanic	27,570	28,183	29,367	31,550	32,895	34,509	35,699	37,369	39,540
Other	2,893	2,983	3,261	3,501	3,960	4,192	4,488	4,711	4,860
Total	199,339	194,001	191,713	192,537	190,654	188,723	189,392	192,383	195,637

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



The yearly number of births in Georgia increased from 112,666 in 1990 to 122,368 in 1998. The proportion of births to Hispanics more than tripled during this period, from 2 percent in 1990 to almost 7 percent in 1998.* Georgia experienced improvements over the nine-year span, including marked reductions in the proportions of teen births to young women who were already mothers, births to mothers who smoked during pregnancy, and births to mothers who received late or no prenatal care (see figure).


More specifically:

- **Teen births.** Teen births decreased from 17 percent of total births in 1990 to 15 percent in 1998. Despite this decrease, Georgia was still higher than the national average of 13 percent in 1998.
- **Repeat teen births.** The percentage of Georgia teen births to young women who were already mothers fluctuated during the 1990s, ranging between 23 percent and 29 percent of teen births. Between 1990 and 1998, Georgia experienced an overall drop in the share of teen births that were repeat births, but repeat births were still more common in Georgia than in the U.S. overall in 1998.
- **Births to unmarried women.** During the 1990-1998 period, both Georgia and the nation as a whole experienced an increase in the proportion of births to unmarried women, with Georgia consistently above the national average on this measure. In 1998, 36 percent of births in Georgia were to unmarried women, compared with the U.S. average of 33 percent.
- **Low maternal education.** Georgia saw an overall decrease in the share of births to women with less than 12 years of education during the 1990s, from 25 percent in the early 1990s to 23 percent in 1995 through 1998. Births to women with less than 12 years of education were slightly more common in Georgia than in the nation overall throughout this time.
- **Late or no prenatal care.** The percentage of Georgia births to mothers who received late or no prenatal care fell from 7 percent in 1990 to 3 percent in 1995, and remained relatively stable thereafter. This remarkable drop from 1990 to 1998 outpaced the downward trend for the nation as a whole and left Georgia below the U.S. average from 1993 through 1998.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Georgia witnessed a striking drop in the proportion of births to women who smoked during pregnancy during the 1990s, from more than 15 percent in 1990 to 10 percent in 1996 through 1998. The U.S. as a whole saw a similar decline on this measure from 1990 to 1998, although Georgia remained below the national average throughout the period.
- **Low-birthweight births.** The share of Georgia births that were low-birthweight remained stable and slightly above the national average throughout the 1990-1998 period. In 1998, approximately 9 percent of Georgia births were low-birthweight, compared with nearly 8 percent of births in the U.S. overall.
- **Preterm births.** Preterm births accounted for 13 percent of Georgia births in 1990 and 1991 and 12 percent of births thereafter. The percentage of preterm births in Georgia was above the U.S. average from 1990-1995. In 1996-1998, Georgia matched the U.S. average on this measure.

In summary, Georgia experienced large declines during the 1990s in the percentages of births to mothers who smoked during pregnancy, births to mothers who received late or no prenatal care, and teen births that were repeat births. Despite these improvements, Georgia's record was worse than the national average on five of the eight indicators in 1998. Moreover, there was a notable rise during the 1990s in the proportion of births to unmarried women.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	16.7	16.7	16.2	16.1	16.2	16.3	15.9	15.5	15.0	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	28.1	28.4	29.1	26.8	24.2	22.9	23.7	23.9	24.1	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	32.8	34.6	35.0	35.8	35.5	35.2	35.0	35.4	36.2	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	25.3	25.8	25.0	24.5	23.7	23.3	23.3	23.2	23.0	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	6.6	5.9	5.4	4.5	3.8	3.2	3.2	3.1	2.8	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	15.5	14.1	13.4	12.5	11.8	11.2	10.4	9.9	9.7	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	8.7	8.6	8.5	8.7	8.6	8.8	8.5	8.8	8.5	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	12.8	12.7	12.5	12.5	11.8	11.7	11.4	11.6	11.6	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	68,005	65,513	65,794	65,281	65,309	66,497	66,346	67,874	69,495
	Black non-Hispanic	40,412	40,173	40,279	39,707	38,852	38,217	38,167	39,404	40,913
	Hispanic	2,263	2,587	2,932	3,467	4,363	5,067	6,266	7,164	8,239
	Other	1,986	2,015	2,111	2,167	2,487	2,501	3,264	3,779	3,721
	Total	112,666	110,288	111,116	110,622	111,011	112,282	114,043	118,221	122,368

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

There was a drop in the yearly number of births in Hawaii from 20,489 in 1990 to 17,583 in 1998. This decline was most evident among whites, who accounted for 3,529 births in 1998, compared with 5,472 in 1990.* In 1998, births to "Other" races (which were nearly all to Asians and Pacific Islanders) were 64 percent of total births.

During this nine-year span, Hawaii experienced striking drops in the percentages of teen births to young women who were already mothers, births to mothers receiving late or no prenatal care, and births to mothers who smoked during pregnancy. Furthermore, when compared with the 1998 national average, the children of Hawaii got off to a significantly healthier start to life on seven out of eight measures, including the percentage of births to mothers with less than 12 years of education (see figure). There was, however, a marked increase in the share of births to unmarried women during the 1990s.

More specifically:

- **Teen births.** During the 1990s, the proportion of births to teens in Hawaii remained relatively stable and lower than the national average. In 1998, 11 percent of births in Hawaii were to teenagers, compared with 13 percent in the U.S. as a whole.
- **Repeat teen births.** The percentage of teen births to young women who were already mothers fluctuated during the 1990-1998 period but underwent an overall decrease, from a high of 22 percent in 1993 to a low of 17 percent in 1998. This change mirrored the downward trend in the U.S. overall, and Hawaii remained below the national average for this indicator throughout the nine-year period.
- **Births to unmarried women.** There was a substantial rise in the share of births to unmarried women in Hawaii that outpaced the upward trend in the U.S. as a whole. Births to unmarried women made up 32 percent of Hawaii births in 1998, compared with 25 percent in 1990.
- **Low maternal education.** Throughout the 1990-1998 period, births to women with less than 12 years of education remained strikingly less common in Hawaii than in the nation as a whole. In 1998, 11 percent of births in Hawaii were to mothers with less than 12 years of education, compared with a U.S. average of 22 percent.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.


- **Late or no prenatal care.** The share of births to women who received late or no prenatal care fluctuated during the 1990-1998 period, undergoing an overall decrease from 6 percent in 1990 through 1993 to 3 percent in 1998. The U.S. as a whole experienced a similar but steadier downward trend.

- **Smoking during pregnancy.** There was a sharp drop in the proportion of births to women who smoked during pregnancy, from 12 percent in 1990 to 8 percent in 1997 and 1998. The nation as a whole also experienced a sizable downward trend, but Hawaii remained below the U.S. average for this indicator throughout the 1990-1998 period.

- **Low-birthweight births.** The share of babies born at a low birthweight in Hawaii remained on par with the national average throughout the 1990s. In 1998, low-birthweight babies accounted for 8 percent of total births both in Hawaii and in the U.S. as a whole.

- **Preterm births.** Preterm babies accounted for 10 percent of Hawaii births in 1990 through 1992 and 11 percent of births thereafter. Preterm births were slightly less prevalent in Hawaii than in the nation overall in 1998.

In summary, during the 1990s, Hawaii saw marked drops in the percentages of teen births that were repeat births, births to mothers who received late or no prenatal care, and births to mothers who smoked during pregnancy. In addition, Hawaii's children got off to a significantly healthier start to life on all but one indicator in 1998, when compared with the U.S. average. Throughout the nine-year span, the percentage of Hawaii births to women with less than 12 years of education remained well below the national average. However, there was a large increase in the share of births to unmarried women in Hawaii during the 1990s.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	Hawaii 10.5	10.6	10.0	10.1	10.5	10.1	10.4	10.1	10.7
	U.S. Average 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	Hawaii 20.5	20.8	20.8	21.6	17.6	18.5	18.4	17.2	16.6
	U.S. Average 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	Hawaii 24.8	26.1	26.2	27.2	28.3	29.2	30.3	29.9	31.5
	U.S. Average 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	Hawaii 11.8	11.8	11.2	10.8	11.2	10.8	11.0	10.4	11.2
	U.S. Average 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	Hawaii 5.9	5.7	5.7	6.0	3.1	3.6	3.5	4.6	3.1
	U.S. Average 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	Hawaii 12.5	12.1	11.4	9.7	8.6	8.4	8.6	8.0	8.5
	U.S. Average 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	Hawaii 7.1	6.8	7.2	6.8	7.2	7.0	7.3	7.2	7.5
	U.S. Average 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	Hawaii 10.0	10.3	10.3	10.5	11.2	10.8	10.7	10.6	11.1
	U.S. Average 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	5,472	5,224	5,084	4,858	4,728	4,311	4,118	3,766
	Black non-Hispanic	651	597	654	602	601	540	489	550
	Hispanic	2,437	2,230	2,241	2,192	2,176	2,029	2,164	2,147
	Other	11,929	11,871	11,885	11,941	12,012	11,715	11,630	10,930
	Total	20,489	19,922	19,864	19,593	19,517	18,595	18,401	17,393

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



The yearly number of births in Idaho rose from 16,433 in 1990 to 19,391 in 1998. This increase was evident across all racial-ethnic groups but was most evident among Hispanics. Hispanics accounted for 13 percent of Idaho births in 1998, compared with just 9 percent in 1990.*

During the 1990s, Idaho experienced an increase in the share of births to unmarried women (see figure) and a drop in the share of births to women who received late or no prenatal care. Overall, the changes in measures of a healthy start to life in Idaho were generally small and tended to mirror the trends in the U.S. average.

More specifically:

- **Teen births.** The percentage of Idaho births to teens was similar in 1990 and 1998 despite a rise and fall in this percentage during the nine-year period. In 1998, the proportion of Idaho births to teens was on par with the national average at 13 percent.
- **Repeat teen births.** The percentage of teen births to young women who were already mothers was similar in 1990 and 1998. This percentage, however, dipped from 20 percent in 1990 to a low of 17 percent in 1995, and then rose to 19 percent by 1998.
- **Births to unmarried women.** During the 1990s, Idaho experienced an increase in the proportion of births to unmarried women, from 17 percent in 1990 to 22 percent in 1998. This change paralleled the rise in the U.S. average, but births to unmarried women remained substantially less common in Idaho than in the nation as a whole throughout the 1990-1998 period.
- **Low maternal education.** There was a slight overall drop in the share of total births to women with less than 12 years of education, from a high of more than 20 percent in 1991 to 18 percent in 1998. The U.S. as a whole saw a similar small downward trend, and women with less than 12 years of education continued to account for a somewhat smaller percentage of births in Idaho than in the nation each year throughout the 1990s.
- **Late or no prenatal care.** The percentage of Idaho births to women who received late or no prenatal care fell from 6 percent in 1990 to 4 percent in 1994, remaining relatively stable thereafter. This change mirrored the downward trend for the U.S. as a whole.


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Idaho saw a decrease in the proportion of births to women who smoked during pregnancy, from 16 percent in 1990 and 1991 to 13 percent by 1998. However, there was a larger drop in the U.S. average, and Idaho went from below the U.S. average in 1990 to simply matching it by 1998.

- **Low-birthweight births.** During the 1990-1998 period, the percentage of Idaho births that were low-birthweight remained relatively stable and below the national average. In 1998, 6 percent of Idaho babies were born at a low birthweight, compared with a national average of 8 percent.

- **Preterm births.** The percentage of births that were preterm fluctuated somewhat during the 1990s but underwent an overall increase, from 9 percent in 1990 to 10 percent in 1998. Nonetheless, Idaho remained below the U.S. average throughout this time.

In summary, Idaho experienced five significant changes in measures of a healthy start, often paralleling the trends in the national average. These include a rise in the percentage of births to unmarried women and a decrease in the share of births to women who received late or no prenatal care. Throughout the 1990-1998 period, the share of Idaho births to unmarried women and the percentages of low-birthweight and preterm births remained substantially lower than the U.S. average.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	12.3	13.3	13.0	13.4	13.1	14.0	13.5	12.7	12.8	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	20.2	21.1	21.0	20.5	17.7	16.5	18.2	20.0	19.1	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	16.7	17.4	18.3	18.7	18.7	19.9	21.3	20.7	22.0	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	19.5	20.4	19.9	19.4	19.0	18.7	19.3	19.0	18.1	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	5.9	5.5	5.2	4.5	4.5	4.1	4.5	4.2	4.4	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	15.9	16.2	15.7	14.7	14.1	13.7	14.3	13.2	13.1	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	5.7	5.8	5.5	5.3	5.5	5.9	5.8	6.3	6.0	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	8.7	8.5	8.6	8.1	8.7	9.4	8.9	9.5	9.6	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	14,521	14,792	15,065	15,044	14,972	15,375	15,614	15,387	16,091
	Black non-Hispanic	42	52	55	44	59	71	72	63	80
	Hispanic	1,434	1,535	1,758	1,850	1,978	2,040	2,180	2,365	2,428
	Other	436	442	484	502	517	549	759	767	792
	Total	16,433	16,821	17,362	17,440	17,526	18,035	18,625	18,582	19,391

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Illinois declined from 195,790 in 1990 to 182,588 in 1998. At the same time, Hispanics accounted for a growing share of Illinois births. While one-eighth of all births were to Hispanics in 1990, nearly one-fifth were to Hispanics in 1998.*

Over the 1990-1998 period, Illinois experienced a marked decline in the proportion of births to mothers who smoked during pregnancy (see figure) and a reduction in the percentage of teen births to young women who were already mothers. In 1998, Illinois was close to the U.S. average on most measures of a healthy start to life.

More specifically:


- **Teen births.** The share of Illinois births to teenagers was close to the U.S. average throughout the 1990-1998 period. In 1998, 12 percent of Illinois births were to teens, compared with 13 percent for the nation as a whole.
- **Repeat teen births.** Illinois saw a decline in the proportion of teen births to young women who already had a child over the 1990-1998 period, from a high of 28 percent in 1992 to 23 percent in 1998. This change mirrored the downward trend in the nation as a whole, where 22 percent of teen births were repeat births in 1998.
- **Births to unmarried women.** There was a small overall increase in the proportion of births to unmarried women in Illinois, from 32 percent in 1990 to 34 percent in 1998. The U.S. as a whole experienced a similar upward trend during the 1990s.
- **Low maternal education.** The percentage of births to women with less than 12 years of education remained relatively stable during the 1990s. In 1998, women with less than 12 years of education accounted for 22 percent of Illinois births, similar to the U.S. average.
- **Late or no prenatal care.** Both Illinois and the nation overall experienced a decline in the share of births to women receiving late or no prenatal care from 1990 to 1998. Women who received late or no prenatal care accounted for 4 percent of Illinois births in 1998, compared with 5 percent in 1990.
- **Smoking during pregnancy.** Over the 1990-1998 period, there was a marked decline in smoking during pregnancy in Illinois that paralleled

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

the downward trend in the U.S. average. The proportion of births to women who smoked during pregnancy fell from 17 percent in 1990 to 12 percent in 1998. Throughout this time, maternal smoking remained slightly less common in Illinois than in the U.S. as a whole.

- **Low-birthweight births.** Babies born at a low birthweight accounted for 8 percent of Illinois births during the 1990-1998 period. Throughout most of this time, low-birthweight births made up a slightly larger share of births in Illinois than in the U.S. overall.
- **Preterm births.** Between 11 percent and 12 percent of Illinois babies were preterm during the 1990s. In 1998, the percentage of preterm births in Illinois was just above the national average.

In summary, during the 1990-1998 period, Illinois experienced four significant improvements in measures of a healthy start to life. The percentage of teen births that were repeat births declined, as did the percentage of births to teens, births to women who received late or no prenatal care, and births to women who smoked during pregnancy. However, there were small increases in the proportion of births to unmarried women and in the shares of both preterm and low-birthweight births. For most measures of a healthy start to life, Illinois was close to the U.S. average in 1998.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	13.1	13.0	12.9	12.8	13.0	12.9	12.7	12.5	12.4
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	26.8	27.1	27.9	26.6	25.3	24.5	23.9	23.6	23.4
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	31.7	32.6	33.4	34.1	34.3	33.8	33.7	33.4	34.1
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	22.0	22.2	22.4	22.5	22.6	22.9	22.4	22.0	22.0
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	5.3	5.3	5.0	4.6	4.4	4.4	4.2	4.0	3.9
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	16.6	15.9	15.6	14.6	13.3	12.9	12.5	12.1	11.8
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	7.6	7.8	7.7	8.1	7.9	7.9	8.0	7.9	8.0
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	11.4	11.7	11.8	11.9	11.8	11.8	11.6	12.1	12.1
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	122,854	119,739	115,760	113,842	112,066	110,180	107,706	105,487
	Black non-Hispanic	41,808	41,996	42,346	42,573	40,798	37,308	36,014	35,256
	Hispanic	24,245	25,727	27,333	28,610	30,350	32,166	32,849	33,171
	Other	6,883	6,769	5,957	5,763	6,043	6,158	6,611	6,889
	Total	195,790	194,231	191,396	190,788	189,257	185,812	183,180	180,803

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Indiana decreased slightly from 86,214 in 1990 to 85,122 in 1998. The number of births to whites decreased, while the number of births to Hispanics rose steadily. The proportion of Hispanic births doubled during this period, from 2 percent in 1990 to 4 percent in 1998.

Indiana experienced significant changes in seven measures of a healthy start to life during the 1990-1998 period. The most striking change was a substantial increase in the percentage of births to unmarried women (see figure). Indiana was close to the U.S. average on all seven available indicators in 1998.


More specifically:

- **Teen births.** Teen births were slightly more common in Indiana than in the U.S. as a whole throughout the 1990-1998 period. During the 1990s, between 14 percent and 15 percent of births in Indiana were to teenagers, compared with the steady national average of 13 percent. The percentage of total births to teens was slightly lower in 1998 than in 1990.
- **Repeat teen births.** The percentage of Indiana teen births to young women who already had a child remained stable from 1990 through 1993 at 23 percent and ranged between 19 percent and 21 percent thereafter. In 1998, Indiana was in line with the U.S. average.
- **Births to unmarried women.** There was a dramatic increase in the proportion of Indiana births to unmarried women during the 1990s, from 26 percent in 1990 to 34 percent in 1998. This sharp rise outpaced the upward trend for the nation as a whole, bringing Indiana slightly above the U.S. average on this indicator by 1998.
- **Low maternal education.** Births to mothers with less than 12 years of education made up a slightly smaller share of births in Indiana than in the U.S. overall throughout the 1990-1998 period. In 1998, births to women with less than 12 years of education accounted for 21 percent of births in Indiana, compared with 22 percent in the nation as a whole.
- **Late or no prenatal care.** The percentage of Indiana births to women who received late or no prenatal care declined between 1990 and 1998. In 1998, 4 percent of births in Indiana and in the nation as a whole were to mothers receiving late or no prenatal care.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Data on smoking during pregnancy were not available for Indiana births.
- **Low-birthweight births.** The percentage of low-birthweight births in Indiana increased from 7 percent in 1990 to 8 percent in 1998, mirroring the slight upward trend for the nation as a whole.
- **Preterm births.** Preterm births became a larger proportion of Indiana births during the 1990s, rising from 10 percent in 1990 to 12 percent in 1998. In 1998, Indiana was on par with the national average for this indicator.

In summary, Indiana was similar to the U.S. as a whole on the majority of indicators of a healthy start to life in 1998. However, the proportions of births to unmarried women and to teenagers were just above the respective national averages in 1998. The most dramatic change in Indiana over the 1990-1998 period was a marked increase in the percentage of births to unmarried women. Indiana also witnessed smaller increases in the proportions of both preterm and low-birthweight births, as well as slight declines in the shares of births to teens and teen births that were repeat births.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	14.5	14.4	14.1	14.1	14.5	14.7	14.5	14.1	13.8	
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	22.5	22.8	23.4	22.9	20.2	19.1	21.4	20.4	21.3	
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	26.2	28.3	29.5	30.8	31.5	31.9	32.3	32.6	33.5	
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	21.2	21.3	20.9	20.5	20.1	20.2	20.3	20.4	20.5	
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	5.6	5.7	4.9	4.8	3.7	3.6	3.8	3.9	4.0	
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	6.6	6.7	6.7	7.0	6.8	7.5	7.6	7.7	7.9	
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	9.9	10.1	10.2	10.7	10.5	10.7	10.7	11.0	11.5	
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	73,933	73,528	71,956	71,507	70,172	70,525	70,539	69,840	70,642
	Black non-Hispanic	9,386	9,453	9,392	9,341	8,941	8,674	8,759	8,749	9,222
	Hispanic	1,856	1,841	1,941	2,131	2,324	2,546	2,950	3,409	3,770
	Other	1,039	885	851	970	1,158	1,090	1,265	1,438	1,488
	Total	86,214	85,707	84,140	83,949	82,595	82,835	83,513	83,436	85,122

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**N.A. = Not Available. See Appendix 2 for more details.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Iowa decreased from 39,409 in 1990 to 37,282 in 1998. At the same time, the relatively small number of births to Hispanics nearly tripled, from 623 (2 percent of all births) in 1990 to 1,739 (5 percent of all births) in 1998.* In addition, throughout this period, births to blacks accounted for around 3 percent of all births, while births to whites fell from 94 percent in 1990 to 89 percent in 1998.

During this nine-year span, Iowa saw significant changes in five measures of a healthy start to life. These changes, including an increase in the share of births to unmarried women and a decline in the share of births to women who smoked during pregnancy, tended to mirror changes in the national average. When compared with the 1998 U.S. average, Iowa's children got off to a healthier start to life on seven of eight indicators, including the percentage of births to women with less than 12 years of education (see figure).


More specifically:

- **Teen births.** Teen births made up a smaller share of births in Iowa than in the U.S. as a whole throughout the 1990-1998 period. In 1998, 11 percent of Iowa births were to teens, compared with a U.S. average of 13 percent.
- **Repeat teen births.** The percentage of teen births to young women who were already mothers dropped from 19 percent in 1990 to 16 percent in 1995, and then rose to 18 percent by 1997 and 1998. During this time, the share of Iowa teen births that were repeat births was consistently below the national average.
- **Births to unmarried women.** Iowa saw an increase in the proportion of births to unmarried women, from 21 percent in 1990 to 27 percent in 1998. There was a similar upward trend in the U.S. average. Throughout the 1990s, unmarried women continued to account for a smaller share of births in Iowa than in the nation as a whole.
- **Low maternal education.** Births to women with less than 12 years of education made up a strikingly smaller proportion of births in Iowa than in the U.S. overall during the 1990s. In 1998, 14 percent of Iowa births were to women with less than 12 years of education, compared with 22 percent in the U.S. overall.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Late or no prenatal care.** The percentage of Iowa births to women who received late or no prenatal care remained stable and well below the national average throughout the 1990-1998 period. In 1998, births to women who received late or no prenatal care made up 2 percent of births in Iowa, compared with 4 percent in the nation as a whole.
- **Smoking during pregnancy.** Although Iowa saw a decline in the share of births to women who smoked during pregnancy, from 23 percent in 1990 to 19 percent in 1998, the U.S. as a whole experienced a bigger decline. Iowa remained above the national average throughout the nine-year span.
- **Low-birthweight births.** Despite a slight increase in the proportion of Iowa births that were low-birthweight from 5 percent in 1990 to 6 percent in 1998, low-birthweight births remained somewhat less common in Iowa than in the U.S. as a whole throughout this period.
- **Preterm births.** The percentage of Iowa births that were preterm increased from 9 percent in the early 1990s to 11 percent in 1998. Throughout this time, however, preterm births were somewhat less prevalent in Iowa than in the nation as a whole.

In summary, Iowa saw increases during the 1990s in the share of births to unmarried women and in the percentages of preterm and low-birthweight births. The percentage of births to women who smoked during pregnancy declined, but remained above the U.S. average in 1998. However, throughout the 1990s, when compared with the U.S. average, the children of Iowa got off to a healthier start to life on all but one measure.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	Iowa 10.2	10.5	10.2	10.7	10.9	11.0	11.0	10.7	10.6
	<i>U.S. Average</i> 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	Iowa 19.3	20.6	20.0	18.3	17.0	15.5	17.4	18.0	17.9
	<i>U.S. Average</i> 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	Iowa 21.0	22.2	23.5	24.6	24.8	25.2	26.3	26.2	27.2
	<i>U.S. Average</i> 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	Iowa 12.8	13.2	13.0	13.3	13.1	12.5	13.2	13.6	13.6
	<i>U.S. Average</i> 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	Iowa 2.3	2.4	2.4	2.3	2.4	2.4	2.5	2.5	2.4
	<i>U.S. Average</i> 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	Iowa 22.6	21.6	20.5	19.2	18.6	18.6	18.9	18.3	18.9
	<i>U.S. Average</i> 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	Iowa 5.4	5.7	5.7	5.7	5.9	6.0	6.4	6.4	6.4
	<i>U.S. Average</i> 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	Iowa 8.7	8.7	9.2	9.4	9.7	9.6	10.0	10.4	10.7
	<i>U.S. Average</i> 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	37,080	36,371	35,724	35,004	34,177	33,463	33,311	32,685
	Black non-Hispanic	1,038	1,148	1,167	1,084	1,047	963	981	1,034
	Hispanic	623	732	853	967	1,060	1,279	1,481	1,739
	Other	668	738	725	771	795	1,105	1,366	1,321
	Total	39,409	38,989	38,469	37,826	37,079	36,810	37,139	36,659

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



The yearly number of births in Kansas was about the same in 1990 and 1998. In 1990 there were 39,020 births, while, in 1998, there were 38,422. Births to Hispanics doubled during this time from 5 percent of births to 10 percent.*

During this nine-year span, Kansas saw a decrease in the share of teen births to young women who were already mothers and an increase in the share of births to unmarried women (see figure). The percentage of births to unmarried Kansas women and to Kansas women with less than 12 years of education were substantially below the national average.

More specifically:

- **Teen births.** Teenagers accounted for a similar share of Kansas births in 1990 and 1998, despite a slight rise and fall in the percentage of teen births during this time. In 1998, Kansas matched the U.S. average of 13 percent for this indicator.
- **Repeat teen births.** The percentage of teen births to young women who were already mothers fell from 25 percent in 1990 to 21 percent in 1994, and remained relatively stable thereafter. In 1998, the share of Kansas teen births that were repeat births matched the national average.
- **Births to unmarried women.** Kansas saw a sizable increase in the proportion of births to unmarried women, from 22 percent in 1990 to 28 percent in 1998. This change mirrored the upward trend for the U.S. as a whole. However, the percentage of Kansas births to unmarried women remained below the national average throughout the nine-year period.
- **Low maternal education.** The percentage of births to women with less than 12 years of education ranged between 17 percent and 18 percent of Kansas births during the 1990-1998 period. In 1998, Kansas was below the national average of 22 percent.
- **Late or no prenatal care.** The share of births to women who received late or no prenatal care decreased from 4 percent in 1990 to 3 percent by 1995, and then remained stable through 1998. Kansas was below the U.S. average for this indicator throughout the 1990-1998 period.


- **Smoking during pregnancy.** Kansas experienced a slight drop in the percentage of births to women who smoked during pregnancy during the 1990s, from a high of 15 percent in 1991 to 13 percent in 1993-1998. Because there was a stronger downward trend for the U.S. as a whole,

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

Kansas moved from below the national average for this indicator in 1990 to matching it by 1998.

- **Low-birthweight births.** Despite a slight increase in the percentage of Kansas babies born at a low birthweight from 6 percent in 1990 to 7 percent in 1996 through 1998, Kansas remained just below the U.S. average on this measure throughout the 1990-1998 period.
- **Preterm births.** There was an overall increase in the share of Kansas births that were preterm, from 9 percent in 1990 to 11 percent in 1998. Since the nation as a whole experienced a similar increase, preterm births remained slightly less prevalent in Kansas than nationally throughout the nine-year period.

In summary, during the 1990-1998 period, Kansas experienced a marked decrease in the percentage of teen births that were repeat births and a sizable increase in the percentage of births to unmarried women. Although Kansas did not stray far from the national average for most indicators of a healthy beginning during the 1990s, the percentages of births to unmarried women and to women with less than 12 years of education were substantially below the U.S. average throughout this time.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	Kansas 12.3	12.2	12.4	12.9	12.9	13.1	13.1	12.9	12.6
	U.S. Average 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	Kansas 24.7	23.6	23.7	22.0	21.1	20.6	20.4	21.0	21.2
	U.S. Average 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	Kansas 21.5	23.1	24.3	25.9	26.0	25.9	26.9	27.6	27.8
	U.S. Average 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	Kansas 17.0	17.5	17.8	17.5	16.9	17.4	18.1	18.1	18.1
	U.S. Average 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	Kansas 3.9	3.7	3.2	3.2	3.1	2.7	2.6	2.7	2.8
	U.S. Average 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	Kansas 14.3	14.7	13.8	13.2	12.9	12.6	12.9	13.4	13.1
	U.S. Average 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	Kansas 6.2	6.2	6.4	6.6	6.5	6.4	6.9	6.9	7.0
	U.S. Average 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	Kansas 9.5	9.7	9.6	10.1	9.8	9.9	10.2	10.4	10.8
	U.S. Average 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	32,629	31,607	31,283	30,262	30,194	30,010	29,288	30,013
	Black non-Hispanic	3,318	3,142	3,287	3,204	3,102	2,857	2,740	2,759
	Hispanic	1,954	2,088	2,311	2,417	2,694	2,828	3,225	3,553
	Other	1,119	1,002	1,146	1,523	1,389	1,506	1,398	1,540
	Total	39,020	37,839	38,027	37,406	37,379	37,201	36,651	37,289

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



The yearly numbers of births in Kentucky were nearly the same in 1990 and 1998. In 1990 there were 54,362 births; in 1998, there were 54,329 births. Throughout this time, nearly 90 percent of total births were to whites and 9 percent were to blacks.*

During the 1990-1998 period, Kentucky saw improvements in five of eight measures of a healthy start to life, including a substantial drop in the share of births to women who received late or no prenatal care. At the same time, however, there was a sizable increase in the proportion of births to unmarried women, and the share of Kentucky births to women who smoked during pregnancy remained well above the U.S. average (see figure).

More specifically:

- **Teen births.** The percentage of Kentucky births to teens ranged between 17 and 18 percent during the 1990-1996 period, and then fell to 15 percent in 1998. Despite this decrease, teenagers accounted for a notably larger share of births in Kentucky than in the U.S. as a whole throughout the nine-year period.
- **Repeat teen births.** The proportion of Kentucky teen births to young women who were already mothers fell from a high of 23 percent in 1991 to a low of 19 percent in 1995, and then rose to 21 percent in 1997 and 1998. In 1998, Kentucky was in line with the U.S. average.
- **Births to unmarried women.** Births to unmarried women became a considerably larger share of Kentucky births, rising from 24 percent in 1990 to 30 percent in 1996 through 1998. This change mirrored the upward trend in the national average for this measure.
- **Low maternal education.** There was a drop in the percentage of Kentucky births to women with less than 12 years of education, from 26 percent in 1990 to 22 percent in 1998. This change closed the gap between Kentucky and the national average.
- **Late or no prenatal care.** Kentucky not only experienced a marked decline in the share of births to women who received late or no prenatal care, but also remained below the U.S. average for this measure throughout the 1990s. Births to women who received late or no prenatal care accounted for 2 percent of Kentucky births in 1998, compared with 4 percent of births in the U.S. as a whole.


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Despite a drop in the percentage of births to women who smoked during pregnancy from 29 percent in 1990 to 25 percent in 1994 through 1998, smoking during pregnancy remained strikingly more common in Kentucky than in the U.S. overall throughout the 1990s. In 1998, births to women who smoked during pregnancy accounted for 25 percent of births in Kentucky, compared with 13 percent in the nation as a whole.

- **Low-birthweight births.** The percentage of Kentucky babies born at a low birthweight remained close to the U.S. average throughout the 1990-1998 period. In 1998, 8 percent of births were low-birthweight both in Kentucky and in the U.S. as a whole.

- **Preterm births.** There was an increase in the share of Kentucky births that were preterm, from 10 percent in 1990 to 12 percent in 1998. Since the U.S. as a whole experienced a smaller upward trend, Kentucky was above the U.S. average by 1998.

In summary, Kentucky experienced improvements in five of eight measures of a healthy start to life, including a notable decrease in the percentage of births to women who receives late or no prenatal care. However, the state also saw marked increases in the share of births to unmarried women and the percentage of births that were preterm. Kentucky remained below the national average for the percentage of births to women who received late or no prenatal care during the 1990s, but was well above the U.S. average for the percentage of births to women who smoked during pregnancy.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens									
Kentucky	17.5	17.4	16.5	16.8	17.2	17.2	17.0	16.3	15.4
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers									
Kentucky	22.4	23.3	22.7	22.0	19.6	19.4	19.9	21.1	20.9
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women									
Kentucky	23.6	25.4	26.3	27.2	27.6	28.5	29.8	29.5	30.1
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education									
Kentucky	26.3	26.2	25.2	24.3	23.7	23.2	22.9	22.6	22.0
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care									
Kentucky	4.8	4.6	4.2	3.6	3.0	2.9	2.9	2.6	2.5
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy									
Kentucky	28.5	27.9	27.0	26.4	24.6	24.7	24.5	24.7	24.8
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)									
Kentucky	7.1	7.2	6.8	7.1	7.7	7.6	7.9	7.8	8.1
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)									
Kentucky	10.5	11.1	10.6	11.3	11.6	11.6	11.7	11.8	12.5
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother									
White non-Hispanic	48,574	48,447	47,840	47,266	47,179	46,634	46,727	47,230	48,095
Black non-Hispanic	5,065	5,255	5,167	4,822	4,876	4,759	4,830	4,630	4,845
Hispanic	274	231	372	401	466	493	555	727	751
Other	449	393	461	511	462	491	594	616	638
Total	54,362	54,326	53,840	53,000	52,983	52,377	52,706	53,203	54,329

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

106

107

The yearly number of births in Louisiana declined from 72,192 in 1990 to 66,888 in 1998. This decrease was reflected in reductions in births to both whites and blacks, who together accounted for most of Louisiana births throughout the nine-year period. In 1998, 55 percent of Louisiana births were to whites and 41 percent were to blacks.*

In 1998, Louisiana lagged behind the nation as a whole on six of the eight measures of a healthy start to life. Nevertheless, the state saw improvements during the 1990-1998 period in four measures, including a substantial drop in the percentage of births to women who smoked during pregnancy. At the same time, there was also a substantial rise in the proportion of births to unmarried women (see figure).

More specifically:


- **Teen births.** The percentage of births to teens in Louisiana remained between 17 and 19 percent during the 1990-1998 period, consistently above the national average. In 1998, 18 percent of Louisiana births were to teen mothers, compared with the U.S. average of 13 percent.
- **Repeat teen births.** The proportion of Louisiana teen births to young women who were already mothers fell from a high of 27 percent in 1992 to 23 percent in 1998. Despite this substantial decline, the percentage of Louisiana teen births that were repeat births remained slightly above the U.S. average throughout the 1990-1998 period.
- **Births to unmarried women.** During the 1990s, there was a sizable increase in the percentage of Louisiana births to unmarried women, from 37 percent in 1990 to 45 percent in 1998. This rise surpassed the upward trend for the nation as a whole, and Louisiana remained well above the U.S. average on this measure throughout the nine-year period.
- **Low maternal education.** Births to women with less than 12 years of education became a smaller share of Louisiana births over the 1990-1998 period, dropping from 27 percent in 1990 through 1993 to 25 percent in 1998. Louisiana was consistently above the national average on this measure throughout this time.
- **Late or no prenatal care.** The proportion of births to mothers receiving late or no prenatal care decreased during the 1990s, from 6 percent in the early 1990s to 4 percent in 1994 through 1998. This

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

overall change from 1990 to 1998 mirrored the downward trend for the U.S. as a whole.

- **Smoking during pregnancy.** Louisiana experienced a striking drop in the percentage of births to women who smoked during pregnancy over the 1990s, from 15 percent in 1990 to 10 percent in 1998. Smoking during pregnancy remained less common in Louisiana than in the nation as a whole throughout the nine-year period.
- **Low-birthweight births.** Over the 1990-1998 period, the proportion of Louisiana babies born at a low birthweight increased slightly and was consistently above the U.S. average. In 1998, 10 percent of births in Louisiana were low-birthweight, compared with 8 percent in the nation as a whole.
- **Preterm births.** Preterm births made up a larger share of births in Louisiana than in the U.S. overall throughout the 1990s. In 1998, 15 percent of births in Louisiana were preterm, compared with the national average of 12 percent.

In summary, when compared with the 1998 U.S. average, Louisiana's children got off to a less healthy start to life on six indicators. However, the percentage of births to mothers who smoked during pregnancy remained below the national average, while continuing to fall throughout the 1990-1998 period. Louisiana also saw reductions in the proportion of births to mothers who received late or no prenatal care and in the share of teen births that were repeat births. At the same time, however, there was a notable rise in the percentage of births to unmarried women.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	17.6	17.6	18.1	18.7	19.2	19.1	18.9	18.6	18.4	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	25.9	26.1	26.7	25.4	25.2	22.9	23.0	23.2	22.7	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	36.8	38.4	40.2	42.0	42.6	42.4	43.4	43.9	44.9	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	26.6	27.5	27.4	27.4	26.5	25.3	25.1	24.5	24.5	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	5.9	6.3	5.8	4.8	4.2	4.0	4.2	4.1	3.9	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	15.1	13.8	12.4	11.6	11.5	11.1	10.7	10.3	10.3	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	9.2	9.4	9.4	9.3	9.6	9.7	9.9	10.2	10.1	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	13.9	14.2	14.3	14.2	14.2	14.0	13.7	14.5	14.7	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	40,547	39,788	38,900	37,658	36,808	36,448	36,184	36,445	36,756
	Black non-Hispanic	29,688	30,347	29,751	29,633	28,432	26,784	26,438	27,097	27,385
	Hispanic	941	927	977	1,001	1,413	1,158	1,267	1,263	1,327
	Other	1,016	1,131	1,079	1,110	1,164	1,251	1,315	1,220	1,420
	Total	72,192	72,193	70,707	69,402	67,817	65,641	65,204	66,025	66,888

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Maine fell by about 20 percent from 17,359 in 1990 to 13,733 in 1998. Whites accounted for well over 90 percent of Maine births throughout this time.*

When compared with the national average, Maine's children got off to a healthier start to life on seven of eight indicators in 1998. Moreover, Maine saw a striking decrease during the 1990s in the already small percentage of teen births to young women who were already mothers (see figure). However, the proportion of births to unmarried women increased markedly during this time.


More specifically:

- **Teen births.** Throughout the 1990-1998 period, births to teens remained less common in Maine than in the nation as a whole. In 1998, teenagers accounted for 10 percent of births in Maine, compared with 13 percent in the U.S. overall.
- **Repeat teen births.** The proportion of Maine teen births to young women who were already mothers underwent a sizable decrease, from 20 percent in 1990 and 1991 to 15 percent in 1995, and then remained relatively stable thereafter. Since this drop outpaced the downward trend in the U.S. as a whole, Maine remained below the national average for this measure throughout the 1990-1998 period.
- **Births to unmarried women.** Maine saw a dramatic increase in the share of births to unmarried mothers, from 23 percent in 1990 to 31 percent in 1998. Although this change outpaced the increase in the U.S. average, births to unmarried women continued to account for a smaller proportion of births in Maine than in the U.S. as a whole in 1998.
- **Low maternal education.** Already well below the national average, the percentage of Maine births to women with less than 12 years of education fell from 14 percent in 1990 to 11 percent in 1998.
- **Late or no prenatal care.** Throughout the 1990-1998 period, births to women who received late or no prenatal care accounted for a smaller share of births in Maine than in the nation overall. In 1998, 2 percent of births in Maine were to women who received late or no prenatal care, compared with 4 percent in the U.S. as a whole.
- **Smoking during pregnancy.** The percentage of Maine births to women who smoked during pregnancy fell from 22 percent in 1990 and 1991 to

18 percent in 1995, and then rose to 20 percent in 1998. Smoking during pregnancy remained more common in Maine than in the U.S. as a whole throughout the nine-year period.

- **Low-birthweight births.** Despite a slight increase in the share of Maine births that were low-birthweight, Maine remained below the U.S. average for this measure throughout the 1990-1998 period. In 1998, low-birthweight babies made up 6 percent of births in Maine, compared with 8 percent in the nation as a whole.
- **Preterm births.** The percentage of births that were preterm remained smaller in Maine than in the U.S. overall throughout the 1990s. In 1998, 9 percent of Maine births were preterm, compared with the U.S. average of 12 percent.

In summary, when compared with the national average in 1998, the children of Maine got off to a healthier start to life on all indicators except maternal smoking. Further, Maine saw a number of improvements during the 1990-1998 period, including decreases in the already small percentages of births to mothers with less than 12 years of education and teen births that were repeat births. At the same time, however, there was a marked increase in the share of Maine births to unmarried women as well as an increase in the percentage of preterm births.



* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	Maine 10.8	10.9	10.2	10.1	10.2	10.3	9.7	10.1	9.8	
	U.S. Average 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	Maine 19.8	19.9	19.5	18.4	16.8	14.7	15.2	16.4	14.9	
	U.S. Average 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	Maine 22.6	25.0	25.3	27.0	28.2	27.8	28.7	29.7	30.6	
	U.S. Average 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	Maine 14.5	14.2	13.4	13.0	11.7	11.8	11.7	11.7	10.9	
	U.S. Average 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	Maine 2.5	3.3	1.8	1.7	1.4	1.7	1.7	1.7	1.7	
	U.S. Average 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	Maine 21.5	21.7	20.7	20.6	18.8	18.2	19.3	19.2	19.6	
	U.S. Average 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	Maine 5.1	5.4	5.0	5.4	5.7	6.1	5.9	5.9	5.8	
	U.S. Average 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	Maine 7.6	7.3	7.5	8.1	8.8	9.1	8.8	9.2	9.1	
	U.S. Average 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	16,081	15,602	15,033	14,340	13,819	13,248	12,951	12,656	12,674
	Black non-Hispanic	83	71	75	54	69	66	80	68	78
	Hispanic	127	107	101	107	107	112	115	121	131
	Other	1,068	973	848	564	446	470	628	824	850
	Total	17,359	16,753	16,057	15,065	14,441	13,896	13,774	13,669	13,733

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Maryland

The yearly number of births in Maryland declined from 80,245 in 1990 to 71,972 in 1998. In 1998, births to whites accounted for 57 percent of total births, while births to blacks accounted for nearly a third. The Hispanic share was 5 percent of total births.*


When compared with the national average in 1998, Maryland's children got off to a healthier start to life on five of the eight indicators, including the percentage of births to mothers with less than 12 years of education (see figure). Moreover, there were marked declines during the 1990-1998 period in the share of teen births to young women who were already mothers and in the share of all births to mothers who smoked during pregnancy. At the same time, the proportion of Maryland births to unmarried women increased. More specifically:

- **Teen births.** The percentage of Maryland births to teens remained relatively stable and below the U.S. average throughout the 1990-1998 period. In 1998, teenagers accounted for 10 percent of births in Maryland, compared with 13 percent in the nation as a whole.
- **Repeat teen births.** The proportion of Maryland teen births to young women who were already mothers dropped from 26 percent in 1990 and 1991 to 19 percent by 1995 and then remained relatively stable thereafter. This change brought Maryland from above the national average for this measure in 1990 to below it by 1995.
- **Births to unmarried women.** Maryland saw an overall increase in the share of births to unmarried women, from 30 percent in 1990 to 34 percent in 1998. This change mirrored the rise in the U.S. average.
- **Low maternal education.** Already well below the national average in 1990, the percentage of Maryland births to women with less than 12 years of education fell from 16 percent in 1990 to 13 percent in 1996 through 1998.
- **Late or no prenatal care.** Throughout the 1990-1998 period, births to women who received late or no prenatal care remained less common in Maryland than in the nation as a whole. In 1998, births to women who received late or no prenatal care accounted for 3 percent of births in Maryland, compared with 4 percent in the nation overall.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Maryland experienced a marked drop in the percentage of births to women who smoked during pregnancy, from 16 percent in 1990 to 10 percent in 1998. Moreover, smoking during pregnancy remained less common in Maryland than in the U.S. as a whole throughout the nine-year period.
- **Low-birthweight births.** There was a slight increase in the share of Maryland babies born at a low birthweight, from 8 percent in 1990 to 9 percent by 1997-1998. The percentage of Maryland births that were low-birthweight remained just above the national average throughout the 1990-1998 period.
- **Preterm births.** The share of Maryland births that were preterm rose from 11 percent in 1990 to 13 percent in 1998. During this time, preterm births were slightly more common in Maryland than in the U.S. as a whole.

In summary, the children of Maryland got off to a healthier start to life on five of eight indicators, when compared with the national average in 1998. Furthermore, during the 1990-1998 period, the state saw notable declines in the share of teen births that were repeat births and the percentage of births to women who smoked during pregnancy. There were smaller declines in the percentage of births to women with less than 12 years of education, and births to women who received late or no prenatal care. However, there were increases in the percentages of preterm and low-birthweight births and in the share of births to unmarried women.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens									
Maryland	10.5	10.4	9.8	10.0	10.3	10.3	10.3	10.3	10.2
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers									
Maryland	25.7	25.9	25.3	23.2	21.9	18.9	19.3	20.2	20.3
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women									
Maryland	29.6	30.7	30.5	32.5	33.7	33.3	33.5	33.5	34.4
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education									
Maryland	15.5	15.1	14.4	14.6	14.0	13.6	12.9	12.8	13.0
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care									
Maryland	3.9	3.7	3.7	3.8	3.5	3.0	2.7	2.6	3.0
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy									
Maryland	15.8	15.1	13.7	12.8	11.7	11.8	11.3	10.2	9.8
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)									
Maryland	7.8	8.1	8.3	8.5	8.5	8.5	8.6	8.8	8.7
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)									
Maryland	11.2	11.7	11.5	12.1	12.2	12.7	12.3	12.7	12.8
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother									
White non-Hispanic	49,463	47,821	46,492	44,519	44,019	43,477	42,214	40,890	40,893
Black non-Hispanic	23,716	23,883	24,549	23,967	23,641	22,348	22,497	22,490	23,741
Hispanic	2,523	2,477	2,980	2,996	3,090	3,155	3,234	3,274	3,580
Other	4,543	5,003	3,794	3,506	3,221	3,416	3,588	3,561	3,758
Total	80,245	79,184	77,815	74,988	73,971	72,396	71,533	70,215	71,972

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Massachusetts

There was a marked decrease in the yearly number of births in Massachusetts, from 92,654 in 1990 to 81,411 in 1998. Births to whites accounted for the vast majority of Massachusetts births, including three of every four births in 1998.* The decline in total births was due to reductions in births to whites and blacks. Births to Hispanics rose slightly and accounted for 11 percent of the total in 1998.

When compared with the U.S. average, the children of Massachusetts got off to a healthier start to life on all eight indicators in 1998. Furthermore, Massachusetts experienced a notable drop in the proportion of births to women with less than 12 years of education, as well as a substantial reduction in the percentage of teen births that were to young women who already had a child (see figure).

More specifically:

- **Teen births.** Teen births were notably less common in Massachusetts than in the nation as a whole during the 1990s. Throughout this period, births to teens accounted for between 7 and 8 percent of Massachusetts births, compared with the national average of 13 percent.
- **Repeat teen births.** During the 1990s, Massachusetts experienced a sizable reduction in the percentage of teen births to young women who were already mothers, from 21 percent in 1990 to less than 17 percent in 1998. This change outpaced the downward trend of the U.S. overall, leaving Massachusetts well below the national average in 1998.
- **Births to unmarried women.** The share of births to unmarried women in Massachusetts increased from 25 percent in 1990 to 27 percent in 1994, and then remained steady at 26 percent through 1998. The U.S. as a whole saw a sharper overall rise in this indicator from 1990 to 1998.

- **Low maternal education.** There was a reduction in the proportion of Massachusetts births to women with less than 12 years of education, from 16 percent in 1990 and 1991 to 13 percent in 1995 through 1998. Massachusetts was markedly below the U.S. average for this indicator throughout the nine-year period.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** A more accurate method for calculating this statistic was introduced in 1996. This introduces a temporary discontinuity in the trends in prenatal care receipt. See Appendix 2 for more details.


- **Late or no prenatal care.** Births to mothers who received late or no prenatal care made up a substantially smaller share of births in Massachusetts than in the nation as a whole throughout the 1990s. In 1998, births to women receiving late or no prenatal care accounted for 2 percent of births in Massachusetts, compared with 4 percent in the U.S. overall.**

- **Smoking during pregnancy.** The abrupt change in the percentage of births to mothers who smoked during pregnancy from 1992 to 1993 probably reflects the effects of state anti-smoking legislation passed in 1992. There was a decline in maternal smoking thereafter, from more than 16 percent in 1993 to less than 12 percent in 1998.

- **Low-birthweight births.** From 1990 to 1998, there was a slight rise in the proportion of Massachusetts births that were low-birthweight, from 6 percent in 1990 through 1996 to 7 percent thereafter. Despite this increase, low-birthweight births remained less prevalent in Massachusetts than in the U.S. as a whole.

- **Preterm births.** Preterm births became a larger share of Massachusetts births over the 1990-1998 period, rising from 8 percent in 1990 to 10 percent in 1998. Nevertheless, preterm births consistently accounted for a smaller proportion of births in Massachusetts than in the nation as a whole.

In summary, when compared with the national average, the children of Massachusetts got off to a healthier start to life on all eight measures. In addition, the state witnessed marked declines from 1990 to 1998 in the share of teen births that were repeat births and in the proportion of births to women with less than 12 years of education.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	8.0	8.0	7.7	7.8	7.8	7.5	7.3	7.4	7.2
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	21.4	21.0	21.5	20.8	20.3	18.0	18.1	18.0	16.5
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	24.7	25.9	25.9	26.4	26.6	25.6	25.5	25.9	26.1
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	16.1	16.0	15.3	14.6	14.2	13.1	13.2	13.4	13.4
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	2.8	2.5	2.1	2.0	1.9	1.9	3.0	2.7	2.4
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	25.3	24.5	23.3	16.5	15.0	13.6	13.2	12.4	11.7
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	5.9	5.9	6.0	6.2	6.4	6.3	6.4	7.0	6.9
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	7.7	7.7	7.7	7.8	8.0	8.3	8.9	9.5	9.8
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	72,261	68,390	67,564	65,659	64,821	63,067	61,446	62,073
	Black non-Hispanic	7,585	7,233	7,141	6,860	6,667	6,272	5,886	6,322
	Hispanic	8,425	8,485	8,522	8,252	8,435	8,109	7,732	8,235
	Other	4,383	4,097	4,004	3,897	3,864	4,200	4,606	4,475
	Total	92,654	88,205	87,231	84,668	83,787	81,648	80,276	80,364

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**See footnote ** in text and Appendix 2.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



There was a substantial decline in the yearly number of births in Michigan, from 153,700 in 1990 to 133,666 in 1998. In 1998, births to whites accounted for 70 percent of total births, births to blacks accounted for 18 percent, and births to Hispanics accounted for 4 percent.*

Over the 1990-1998 period, the percentage of Michigan births to women who smoked during pregnancy declined markedly, yet remained well above the U.S. average (see figure). Michigan saw smaller improvements in a number of other measures of a healthy start to life, including a drop in the share of teen births to young women who already had a child.

More specifically:

- **Teen births.** Teen births became a slightly smaller share of Michigan births during the 1990s. While teenagers accounted for 13 percent of births in 1990, they accounted for 12 percent in 1998.
- **Repeat teen births.** The share of Michigan teen births to young women who already had a child fluctuated during the 1990s but fell overall, from 25 percent in 1990 to 21 percent in 1998. The U.S. as a whole saw a similar downward trend.
- **Births to unmarried women.** The percentage of births to unmarried women is not displayed for 1990-1993 because the number of births to unmarried women in Michigan was underreported during these years.** From 1994 through 1998, the proportion of Michigan births to unmarried women ranged between 33 percent and 35 percent, remaining just above the national average throughout.
- **Low maternal education.** Births to women with less than 12 years of education became a smaller proportion of Michigan births during the 1990s, falling from 20 percent in 1990 through 1992 to 18 percent in 1997 and 1998. Births to women with less than 12 years of education remained less common in Michigan than in the U.S. overall.
- **Late or no prenatal care.** The percentage of Michigan births to mothers who received late or no prenatal care decreased from 4 percent in 1990 to 3 percent in 1998. Michigan remained somewhat below the national average for this indicator throughout the 1990-1998 period.
- **Smoking during pregnancy.** Michigan experienced a substantial decline in the percentage of births to mothers who smoked during


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** See Appendix 2 for more details.

pregnancy, from 23 percent in 1990 to 17 percent in 1998. Yet, throughout the 1990s, smoking during pregnancy remained markedly more common in Michigan than in the U.S. as a whole.

- **Low-birthweight births.** Low-birthweight births accounted for 8 percent of Michigan births throughout the 1990-1998 period. In 1998, Michigan was close to the U.S. average for this indicator.
- **Preterm births.** The percentage of Michigan births that were preterm remained in line with the U.S. average throughout the 1990s. In 1998, preterm births accounted for 11 percent of births in Michigan and 12 percent of births in the U.S. as a whole.

In summary, between 1990 and 1998, Michigan experienced improvements in five of seven available measures of a healthy start to life, including reductions in the shares of all births to mothers who smoked during pregnancy and of teen births to young women who were already mothers. However, the percentage of births to mothers who smoked during pregnancy remained considerably higher than the national average throughout the nine-year period.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	13.5	13.2	13.0	12.6	12.6	12.5	12.2	11.7	11.6	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	24.8	25.9	26.3	24.5	22.5	20.4	21.3	21.5	21.3	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	N.R.	N.R.	N.R.	N.R.	35.0	34.3	33.8	33.2	33.9	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	19.9	20.0	19.8	19.6	18.5	18.3	17.8	17.5	17.5	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	4.1	4.0	3.7	3.6	3.7	3.3	3.0	3.0	3.4	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	22.6	22.3	21.2	19.9	18.6	18.3	17.6	17.6	17.0	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	7.6	7.8	7.5	7.6	7.8	7.7	7.7	7.7	7.8	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	10.7	11.2	10.8	11.2	11.2	10.9	10.8	11.3	11.5	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	109,182	106,234	102,987	99,426	97,473	95,715	94,990	93,875	92,972
	Black non-Hispanic	31,440	31,261	29,463	27,950	26,876	24,679	23,792	23,957	24,007
	Hispanic	4,326	4,397	4,302	4,389	4,454	4,781	5,050	5,454	5,945
	Other	8,752	8,306	7,337	8,090	9,225	9,467	9,555	10,428	10,742
	Total	153,700	150,198	144,089	139,855	138,028	134,642	133,387	133,714	133,666

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**N.R. = Not Reliable. See text and Appendix 2 for more details.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Minnesota

During the 1990s, the yearly number of births in Minnesota decreased, from 68,013 in 1990 to 65,202 in 1998. This decline was due to a substantial reduction in births to whites, as births to both blacks and Hispanics increased over this time. Births to whites fell from 86 percent of births in 1990 to 78 percent in 1998.

Minnesota experienced two striking changes in measures of a healthy start to life during the 1990s: a drop in the proportion of births to mothers who smoked during pregnancy and a rise in the percentage of births to unmarried women. Throughout this time, when compared with the U.S. average, Minnesota's children got off to a healthier start to life on all eight indicators, including the share of births to women with less than 12 years of education (see figure).

More specifically:


- **Teen births.** Teen births were markedly less common in Minnesota than in the U.S. as a whole throughout the 1990-1998 period. In 1998, teen births accounted for 9 percent of births in Minnesota, compared with 13 percent in the nation overall.
- **Repeat teen births.** During the 1990s, between 18 percent and 22 percent of Minnesota teen births were to young women who already had a child. The percentage of Minnesota teen births that were repeat births remained below the U.S. average throughout this time.
- **Births to unmarried women.** There was a marked rise in the share of Minnesota births to unmarried women, from 21 percent in 1990 to almost 26 percent in 1998. A similar upward trend for the U.S. overall left Minnesota below the national average throughout this period.
- **Low maternal education.** The proportion of Minnesota births to mothers with less than 12 years of education remained relatively stable and dramatically below the U.S. average throughout the 1990s. In 1998, 11 percent of births in Minnesota were to women with less than 12 years of education, compared with the national average of 22 percent.
- **Late or no prenatal care.** Over the 1990-1998 period, there was a slight decrease in the percentage of births to women who received late or no prenatal care, from 4 percent in 1990 and 1991 to 3 percent thereafter. Although the nation as a whole experienced a stronger

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

downward trend from 1990 to 1998, Minnesota remained below the U.S. average throughout this time.

- **Smoking during pregnancy.** Births to mothers who smoked during pregnancy became a notably smaller share of Minnesota births over the 1990s, falling from 16 percent in 1990 to 12 percent in 1998. Despite a more pronounced drop in the U.S. average, smoking during pregnancy remained less common in Minnesota than in the nation as a whole throughout the 1990-1998 period.
- **Low-birthweight births.** Throughout the 1990s, low-birthweight births made up a smaller share of births in Minnesota than in the U.S. as a whole. In 1998, 6 percent of births were low-birthweight in Minnesota, compared with the national average of 8 percent.
- **Preterm births.** From 1990 to 1998, the percentage of preterm births in Minnesota rose slightly, from 8 percent in 1990 to 9 percent throughout the remainder of the period. Preterm births were less common in Minnesota than in the U.S. as a whole throughout this time.

In summary, throughout the 1990-1998 period, the children of Minnesota got off to a healthier start to life on all eight measures, when compared with the national average. Furthermore, the state experienced a sizable reduction in the proportion of births to women who smoked during pregnancy over this time. Another noteworthy change was an increase in the share of births to unmarried women.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	8.0	8.1	8.1	8.2	8.5	8.4	8.6	8.8	8.6	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	19.6	21.5	20.9	20.5	19.4	17.8	18.9	18.0	18.3	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	20.9	22.3	23.0	23.4	24.0	23.9	24.8	25.0	25.6	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	10.4	10.5	10.3	10.5	10.1	10.2	10.3	10.4	10.6	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	3.6	3.8	3.5	3.1	3.0	3.0	3.4	3.1	2.9	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	15.8	15.4	15.2	14.4	13.8	13.4	13.1	13.0	12.4	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	5.1	5.3	5.2	5.5	5.7	5.9	5.8	5.9	5.8	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	8.1	8.6	8.5	8.9	8.8	8.9	9.0	9.0	9.2	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	58,476	57,059	52,515	50,870	52,141	49,834	50,726	51,234	51,089
	Black non-Hispanic	2,211	2,368	2,132	2,163	2,554	2,705	3,049	3,284	3,594
	Hispanic	1,050	1,262	1,377	1,560	1,661	1,915	2,324	2,625	2,967
	Other	6,276	6,380	9,583	10,055	7,949	8,809	7,601	7,356	7,552
	Total	68,013	67,069	65,607	64,648	64,305	63,263	63,700	64,499	65,202

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Mississippi

The yearly number of births in Mississippi decreased from 43,563 in 1990 to 40,987 in 1996, and then rose to 42,939 births in 1998. Throughout this time, slightly more than half of Mississippi births were to whites and slightly less than half were to blacks.*

When compared with the national average in 1998, Mississippi's children got off to a significantly less healthy start to life on six of eight indicators, including the percentage of births to teens (see figure). Moreover, there was a marked rise during the 1990-1998 period in the already large share of births to unmarried women. On the other hand, Mississippi experienced a number of improvements, including a substantial drop in the percentage of teen births to young women who already had a child.


More specifically:

- **Teen births.** Throughout the 1990s, births to teens made up a substantially greater proportion of births in Mississippi than in the nation as a whole. In 1998, 20 percent of births in Mississippi were to teenagers, compared with 13 percent in the U.S. overall.
- **Repeat teen births.** The share of Mississippi teen births to young women who were already mothers decreased from a high of 29 percent in 1992 to 24 percent by 1995, and then remained relatively stable through 1998. The share of Mississippi teen births that were repeat births remained above the U.S. average throughout the 1990-1998 period.
- **Births to unmarried women.** The percentage of births to unmarried women in Mississippi rose from 40 percent in 1990 to 45 percent in 1994 and remained stable thereafter. Throughout the 1990-1998 period, births to unmarried women were substantially more common in Mississippi than in the nation as a whole.
- **Low maternal education.** Births to women with less than 12 years of education became a smaller share of Mississippi births during the 1990s, falling from 30 percent in 1990 to 26 percent in 1997 and 1998. Mississippi remained above the U.S. average for this indicator throughout the 1990s.
- **Late or no prenatal care.** The proportion of Mississippi births to women who received late or no prenatal care fell from a high of 5 percent in 1993 to 4 percent in 1998. Since the nation as a whole saw a

stronger downward trend, Mississippi went from being below the U.S. average for this indicator in 1990 to even with it in 1998.

- **Smoking during pregnancy.** The percentage of Mississippi births to women who smoked during pregnancy fell from 16 percent in 1990 to 13 percent in 1995 and remained stable thereafter. Mississippi went from below the national average in 1990 to even with it in 1998.
- **Low-birthweight.** Low-birthweight births were more prevalent in Mississippi than in the U.S. as a whole during the 1990-1998 period. In 1998, 10 percent of births in Mississippi were low-birthweight, compared with 8 percent in the U.S. overall.
- **Preterm births.** Preterm births were substantially more common in Mississippi than in the nation as a whole throughout the 1990s. In 1998, preterm births accounted for 16 percent of Mississippi births, compared with a national average of 12 percent.

In summary, when compared with the U.S. average, the children of Mississippi got off to a significantly less healthy start to life on six of eight measures in 1998. Furthermore, the state experienced an increase during the 1990-1998 period in the already high percentage of births to unmarried women. However, there were also a number of improvements. The share of teen births that were repeat births declined, as did the percentages of births to mothers who smoked during pregnancy, births to mothers who received late or no prenatal care, and births to mothers with less than 12 years of education.



* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	21.3	21.7	21.4	21.7	22.1	22.2	21.3	20.7	20.0	
<i>U.S. Average</i>	<i>12.8</i>	<i>12.9</i>	<i>12.7</i>	<i>12.8</i>	<i>13.1</i>	<i>13.1</i>	<i>12.9</i>	<i>12.7</i>	<i>12.5</i>	
Percent of teen births to women who were already mothers	28.4	28.5	29.1	27.2	25.0	23.6	24.6	25.3	24.4	
<i>U.S. Average</i>	<i>24.3</i>	<i>24.6</i>	<i>25.0</i>	<i>23.4</i>	<i>21.8</i>	<i>21.0</i>	<i>21.4</i>	<i>21.6</i>	<i>21.5</i>	
Percent of total births to unmarried women	40.5	42.4	42.9	44.4	45.4	45.3	45.0	45.4	45.4	
<i>U.S. Average</i>	<i>28.0</i>	<i>29.5</i>	<i>30.1</i>	<i>31.0</i>	<i>32.6</i>	<i>32.2</i>	<i>32.4</i>	<i>32.4</i>	<i>32.8</i>	
Percent of total births to mothers with less than 12 years of education	29.6	29.2	28.6	28.1	27.4	27.2	26.6	26.1	26.2	
<i>U.S. Average</i>	<i>23.8</i>	<i>23.9</i>	<i>23.6</i>	<i>23.3</i>	<i>22.9</i>	<i>22.6</i>	<i>22.4</i>	<i>22.1</i>	<i>21.9</i>	
Percent of total births to mothers receiving late or no prenatal care	4.8	4.9	5.0	5.4	5.0	4.8	4.2	4.1	4.0	
<i>U.S. Average</i>	<i>6.1</i>	<i>5.8</i>	<i>5.2</i>	<i>4.8</i>	<i>4.4</i>	<i>4.2</i>	<i>4.0</i>	<i>3.9</i>	<i>3.9</i>	
Percent of total births to mothers who smoked during pregnancy	15.8	15.4	14.9	14.3	14.3	13.0	12.8	12.6	13.0	
<i>U.S. Average</i>	<i>18.4</i>	<i>17.8</i>	<i>16.9</i>	<i>15.8</i>	<i>14.5</i>	<i>13.9</i>	<i>13.6</i>	<i>13.2</i>	<i>12.9</i>	
Percent low-birthweight births (less than 5.5 lbs)	9.6	9.7	9.9	10.1	9.9	9.8	9.9	10.1	10.1	
<i>U.S. Average</i>	<i>7.0</i>	<i>7.1</i>	<i>7.1</i>	<i>7.2</i>	<i>7.3</i>	<i>7.3</i>	<i>7.4</i>	<i>7.5</i>	<i>7.6</i>	
Percent preterm births (less than 37 completed weeks of gestation)	15.1	15.4	15.4	15.9	15.6	15.4	15.3	15.6	16.0	
<i>U.S. Average</i>	<i>10.6</i>	<i>10.8</i>	<i>10.7</i>	<i>11.0</i>	<i>11.0</i>	<i>11.0</i>	<i>11.0</i>	<i>11.4</i>	<i>11.6</i>	
Number of births by Race and Ethnicity of mother	White non-Hispanic	22,161	21,771	21,554	21,091	21,347	21,321	21,156	22,523	
	Black non-Hispanic	20,748	20,815	20,515	20,417	19,953	19,241	18,958	19,343	
	Hispanic	116	137	141	141	182	220	278	328	403
	Other	538	481	471	500	472	562	595	581	670
	Total	43,563	43,204	42,681	42,149	41,954	41,344	40,987	41,533	42,939

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Missouri decreased from 79,260 in 1990 to 75,358 in 1998. While this decrease was evident among births to both blacks and whites, births to Hispanics increased during the 1990s. Births to Hispanics accounted for almost 3 percent of Missouri births in 1998, compared with 1 percent 1990.*

During the 1990s, Missouri experienced a number of changes in measures of a healthy start to life. The most striking changes were a sharp decrease in the percentage of births to mothers who smoked during pregnancy (see figure) and a sizable increase in the proportion of births to unmarried mothers.

More specifically:


- **Teen births.** Teen births were more common in Missouri than in the U.S. as a whole throughout the 1990-1998 period. During this time, between 14 percent and 15 percent of Missouri births were to teens, compared with the steady national average of 13 percent.
- **Repeat teen births.** The share of teen births to young women who were already mothers increased from 24 percent in 1990 to 26 percent in 1992, and then decreased to 21 percent in 1998. Missouri was close to the national average for this measure throughout this period.
- **Births to unmarried women.** Over the 1990-1998 period, there was a marked increase in the proportion of Missouri births to unmarried mothers, from 29 percent in 1990 to 34 percent in 1998. This change mirrored the upward trend for the U.S. overall, with Missouri just above the national average in both 1990 and 1998.
- **Low maternal education.** Births to women with less than 12 years of education made up a slightly smaller share of births in Missouri than in the nation as a whole throughout the 1990-1998 period. In 1998, 20 percent of Missouri births were to women with less than 12 years of education, compared with the U.S. average of 22 percent.
- **Late or no prenatal care.** The percentage of Missouri births to mothers receiving late or no prenatal care decreased from 5 percent in 1990 to 3 percent in 1994, and remained stable through 1998. Missouri remained below the national average for this indicator throughout the nine-year period.

* Estimates for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Both Missouri and the nation as a whole experienced a striking reduction in maternal smoking during the 1990s. However, smoking during pregnancy remained more common in Missouri than in the U.S. as a whole throughout this period. In 1998, 19 percent of Missouri births were to women who smoked during pregnancy, compared with 13 percent of births in the U.S. overall.
- **Low-birthweight births.** The proportion of Missouri births that were low-birthweight remained similar to the national average throughout the 1990-1998 period. In 1998, low-birthweight births accounted for 8 percent of births in both Missouri and the nation as a whole.

- **Preterm births.** During the 1990s, preterm births accounted for between 11 percent and 12 percent of births in both Missouri and the U.S. overall.

In summary, Missouri experienced a substantial reduction in the percentage of births to mothers who smoked during pregnancy over the 1990-1998 period. However, this percentage remained markedly above the U.S. average throughout this time. Missouri also saw notable declines in the shares of teen births that were repeat births and births to mothers receiving late or no prenatal care. During the same time, there was a marked increase in the percentage of births to unmarried women.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	14.4	14.4	14.5	14.2	14.7	14.4	14.1	13.9	13.8	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	23.8	25.4	25.8	23.8	21.8	21.0	20.6	20.3	20.9	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	28.6	30.2	31.5	32.4	32.5	32.1	33.2	33.1	34.1	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	21.1	21.0	20.9	20.4	19.7	19.2	19.3	19.3	19.6	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	4.8	4.5	4.2	4.0	3.2	3.0	2.9	2.8	2.9	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	24.8	24.2	23.4	22.1	20.7	20.1	19.6	19.6	19.1	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	7.1	7.5	7.3	7.5	7.6	7.6	7.5	7.7	7.8	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	10.7	11.4	11.2	11.5	11.5	11.5	11.3	11.6	12.0	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	63,574	62,887	60,901	59,912	59,289	59,423	59,757	59,788	60,554
	Black non-Hispanic	13,538	13,646	13,282	12,974	11,813	10,990	11,089	10,999	11,367
	Hispanic	970	1,070	1,018	1,150	1,224	1,288	1,516	1,758	1,970
	Other	1,178	1,074	1,100	1,217	1,217	1,327	1,470	1,492	1,467
	Total	79,260	78,677	76,301	75,253	73,543	73,028	73,832	74,037	75,358

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Montana declined from 11,613 in 1990 to 10,795 in 1998. Births to whites accounted for more than 80 percent of all Montana births throughout this time.* Births to American Indians (included in Other) accounted for 11 percent of Montana births in 1998.

When compared with the U.S. average, the children of Montana got off to a significantly healthier start to life on five of eight measures in 1998.

Moreover, Montana saw marked drops during the 1990s in the already small percentages of births to mothers who received late or no prenatal care and of teen births to young women who already had a child (see figure). However, there were sizable increases in the shares of preterm births and births to unmarried women.

More specifically:

- **Teen births.** Teenagers accounted for a relatively stable share of Montana births during the 1990-1998 period. In 1998, the percentage of births to teenagers in Montana was 12 percent, on par with the national average.
- **Repeat teen births.** The percentage of Montana teen births to young women who were already mothers fell from 20 percent in 1990 to 16 percent in 1993, and then remained relatively stable thereafter. Montana remained below the national average for this indicator throughout the 1990-1998 period.
- **Births to unmarried women.** During the 1990s, Montana saw a sizable overall rise in the share of births to unmarried women, mirroring the trend in the U.S. average. Unmarried women accounted for 30 percent of Montana births in 1998, compared with 24 percent in 1990.
- **Low maternal education.** Births to women with less than 12 years of education were markedly less common in Montana than in the U.S. as a whole throughout the 1990-1998 period. In 1998, these births made up 15 percent of total births in Montana, compared with 22 percent in the U.S. overall.
- **Late or no prenatal care.** Montana saw a drop during the 1990s in the share of births to women who received late or no prenatal care, from 5 percent in 1990 to 3 percent by 1994, and remaining relatively stable thereafter. Throughout the 1990-1998 period, Montana was below the U.S. average for this indicator.


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Smoking during pregnancy remained substantially more common in Montana than in the U.S. as a whole throughout the 1990s. In 1998, births to women who smoked during pregnancy accounted for 19 percent of births in Montana, compared with 13 percent in the nation overall.

- **Low-birthweight births.** There was a small increase in the proportion of Montana babies born at a low birthweight, from 6 percent in 1990 through 1995 to 7 percent by 1998. Throughout this time, low-birthweight births remained slightly less prevalent in Montana than in the nation as a whole.

- **Preterm births.** Montana experienced an increase in the share of births that were preterm, from 8 percent in 1990 to 11 percent in 1998. In 1998, Montana was on par with the national average for this measure.

In summary, when compared with the national average in 1998, Montana's children got off to a significantly healthier start to life on five of eight measures. In addition, there were sizable reductions over the 1990-1998 period in the already small shares of births to women who received late or no prenatal care and of teen births that were repeat births. On the other hand, the proportions of preterm births and births to unmarried women rose markedly during this time, and the percentage of births to women who smoked during pregnancy remained substantially above the U.S. average throughout the nine-year span.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	11.5	11.5	11.9	12.5	12.1	12.6	12.4	12.2	12.3	
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	20.2	19.7	18.7	15.9	16.4	15.9	15.7	14.6	15.7	
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	23.7	25.2	26.4	27.3	25.5	26.5	27.9	28.7	29.9	
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	16.4	16.0	16.5	15.7	14.7	14.6	14.4	14.7	15.3	
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	5.1	4.4	4.2	3.7	3.4	3.5	3.2	3.6	3.2	
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	20.5	19.1	20.2	19.2	18.2	18.6	17.9	18.2	18.5	
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	6.2	5.6	6.0	6.0	6.2	5.8	6.4	6.3	7.0	
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	8.5	8.7	8.7	9.3	9.7	9.5	9.7	10.2	11.0	
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	9,373	9,376	9,458	9,445	9,172	9,226	8,894	8,818	8,951
	Black non-Hispanic	35	32	36	34	24	33	31	18	38
	Hispanic	292	260	189	190	250	282	298	302	336
	Other	1,913	1,845	1,789	1,696	1,621	1,601	1,633	1,711	1,470
	Total	11,613	11,513	11,472	11,365	11,067	11,142	10,856	10,849	10,795

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



Between 1990 and 1998, the yearly number of births in Nebraska decreased from 24,380 to 23,534. While births to whites decreased substantially during this time, the percentage of births to Hispanics tripled, from 3 percent in 1990 to 9 percent in 1998.*

During the 1990s, Nebraska had a reduction in the proportion of births to mothers who smoked during pregnancy and a rise in the percentage of births to unmarried women. Throughout this time, when compared with the U.S. average, Nebraska's children got off to a healthier start to life on most indicators, including the share of births to mothers with less than 12 years of education (see figure).

More specifically:

- **Teen births.** Teen births remained less common in Nebraska than in the nation as a whole throughout the 1990s. While births to teenagers accounted for between 10 percent and 11 percent of Nebraska births over the 1990-1998 period, they made up a steady 13 percent of births in the U.S. overall.
- **Repeat teen births.** There was a marked decrease in the proportion of Nebraska teen births to young women who were already mothers, from 22 percent in 1990 and 1991 to 18 percent in 1998. Furthermore, Nebraska remained below the U.S. average on this measure throughout the nine-year period.
- **Births to unmarried women.** Like the U.S. as a whole, Nebraska experienced a sizable rise in the percentage of births to unmarried women during the 1990s, from 21 percent in 1990 to 26 percent in 1997 and 1998. However, births to unmarried women were less common in Nebraska than in the U.S. as a whole.


- **Low maternal education.** During the 1990s, the share of births to women with less than 12 years of education increased from 11 percent in 1990 to almost 15 percent in 1998. The U.S. as a whole, in contrast, experienced a decline on this measure over the nine-year period. Despite these trends, Nebraska remained substantially below the national average throughout the 1990s.

- **Late or no prenatal care.** Births to mothers who received late or no prenatal care accounted for a steady 3 percent of Nebraska births throughout the 1990s. At the same time, the nation as a whole

experienced a notable decline on this measure. Nonetheless, the share of Nebraska births to mothers who received late or no prenatal care remained below the U.S. average throughout the 1990-1998 period.

- **Smoking during pregnancy.** Both Nebraska and the U.S. overall experienced a sharp decline in maternal smoking from 1990 to 1998. In Nebraska, births to mothers who smoked during pregnancy accounted for 16 percent of births in 1998, compared with 21 percent in 1990. However, smoking during pregnancy remained more common in Nebraska than in the nation as a whole throughout this time.
- **Low-birthweight births.** The percentage of Nebraska births that were low-birthweight rose from 5 percent in 1990 to 7 percent in 1997 and 1998. Despite this increase, low-birthweight births were less prevalent in Nebraska than in the U.S. overall throughout the 1990s.
- **Preterm births.** Over the 1990-1998 period, Nebraska saw a rise in the share of preterm births, but remained below the U.S. average in 1998. Eleven percent of Nebraska births were preterm in 1998, compared with 12 percent of births in the U.S. as a whole.

In summary, when compared with the 1998 U.S. average, the children of Nebraska got off to a healthier start to life on all but one indicator: the proportion of births to mothers who smoked during pregnancy. Even on this measure, though, Nebraska experienced a striking reduction during the 1990s. The state also saw a marked decline in the percentage of teen births that were repeat births. During the same period, there were notable increases in the percentages of births to unmarried women and to mothers with less than 12 years of education.



* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	9.8	9.8	9.9	10.1	11.0	10.0	10.6	10.5	10.6
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	21.7	22.0	20.0	19.9	18.6	17.3	16.8	18.5	18.3
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	20.7	21.6	22.6	23.5	24.8	24.3	24.8	25.8	26.2
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	11.5	12.0	12.3	12.3	12.8	13.1	13.5	14.3	14.5
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	3.3	3.2	3.2	2.8	3.0	2.9	2.6	2.9	3.2
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	20.8	19.7	19.9	18.9	18.5	17.5	16.5	16.6	16.3
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	5.3	5.6	5.6	5.9	6.1	6.3	6.3	7.0	6.5
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	8.5	8.9	8.8	9.6	9.5	10.4	10.0	10.8	10.7
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	21,332	20,767	20,013	19,621	19,399	19,050	18,817	18,753
	Black non-Hispanic	1,376	1,334	1,306	1,251	1,265	1,193	1,236	1,227
	Hispanic	802	954	1,105	1,261	1,382	1,615	1,866	1,978
	Other	870	962	973	1,091	1,110	1,130	1,177	1,288
	Total	24,380	24,017	23,397	23,224	23,156	23,243	23,286	23,319

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

There was a considerable rise in the yearly number of births in Nevada, from 21,599 in 1990 to 28,699 in 1998. The proportion of Nevada births to Hispanics doubled during this period, from 15 percent in 1990 to 30 percent in 1998, while the percentage of births to whites decreased markedly.*

Nevada experienced a dramatic drop over the 1990-1998 period in the percentage of births to mothers who smoked during pregnancy. At the same time, there was a substantial increase in the proportion of births to mothers with less than 12 years of education (see figure).

More specifically:

- **Teen births.** The proportion of Nevada births to teenagers ranged from 12 percent to 14 percent during the 1990s. In 1998, teen births accounted for 13 percent of births in Nevada, close to the U.S. average.
- **Repeat teen births.** Teen births to young women who were already mothers made up between 21 percent and 23 percent of Nevada teen births over the 1990-1998 period. In 1998, the percentage of Nevada teen births that were repeat births was not significantly different from the U.S. average.
- **Births to unmarried women.** The percentage of births to unmarried women increased fairly rapidly between 1990 and 1996. The apparent decline in 1997 is almost entirely due to a change in reporting procedures.** In 1997 and 1998, about 35 percent of Nevada births were to unmarried women, slightly higher than the national average.
- **Low maternal education.** During the 1990s, there was a sharp increase in the share of Nevada births to mothers with less than 12 years of education, contrasting with a decline on this measure in the U.S. overall. In 1998, 28 percent of Nevada births were to mothers with less than 12 years of education, compared with the national average of 22 percent.
- **Late or no prenatal care.** The proportion of births to mothers who received late or no prenatal care fluctuated during the 1990s, but was slightly lower in 1998 than in 1990. The U.S. as a whole experienced a more pronounced drop on this measure, widening the gap between Nevada and the national average over this period. In 1998, 7 percent of births in Nevada were to mothers who received late or no prenatal care, compared with 4 percent of births in the nation overall.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.


** In 1997, Nevada began asking for mother's marital status directly. For more details, see Appendix 2.

- **Smoking during pregnancy.** Nevada witnessed a remarkable decline in the percentage of births to mothers who smoked during pregnancy, from 20 percent in 1990 to 13 percent in 1998. Nevada was in line with the national average on this measure in 1998.

- **Low-birthweight births.** Low-birthweight babies made up a relatively constant share of births in Nevada and the U.S. as a whole over the 1990-1998 period. In 1998, 8 percent of births in both Nevada and the nation overall were low-birthweight.

- **Preterm births.** Preterm births became a larger proportion of Nevada births over the 1990-1998 period, rising from 11 percent in 1990 to 13 percent in 1998. Preterm births were slightly more prevalent in Nevada than in the nation as a whole throughout the nine-year period.

In summary, Nevada saw a marked increase in the proportion of births to mothers with less than 12 years of education over the 1990-1998 period. Moreover, the percentages of Nevada births to mothers with less than 12 years of education and to mothers who received late or no prenatal care were considerably higher than the U.S. average in 1998. On the other hand, the state experienced a striking drop during the 1990s in the percentage of births to mothers who smoked during pregnancy.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	Nevada 12.6	12.9	12.4	13.3	13.3	13.7	13.3	13.5	13.1
	U.S. Average 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	Nevada 23.1	22.4	21.6	22.1	20.6	21.4	20.8	21.7	22.8
	U.S. Average 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	Nevada** 25.4	31.9	33.3	34.0	35.0	38.8	39.3	35.5	35.0
	U.S. Average 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	Nevada 23.0	25.4	25.8	26.8	26.4	26.7	25.8	27.8	28.3
	U.S. Average 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	Nevada 8.1	10.1	7.8	7.7	7.6	7.9	6.0	6.3	7.0
	U.S. Average 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	Nevada 20.1	20.8	19.6	18.5	17.8	16.1	13.3	13.6	12.8
	U.S. Average 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	Nevada 7.2	7.2	7.1	7.4	7.6	7.4	7.5	7.6	7.6
	U.S. Average 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	Nevada 11.1	11.6	11.6	12.1	12.0	11.4	11.7	12.3	12.9
	U.S. Average 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	15,378	15,143	14,901	14,739	15,231	15,386	15,121	15,603
	Black non-Hispanic	1,887	2,043	2,127	1,958	2,042	1,863	1,941	2,064
	Hispanic	3,261	3,679	4,116	4,413	5,227	6,124	6,966	7,713
	Other	1,073	1,161	1,230	1,293	1,411	1,610	1,832	2,013
	Total	21,599	22,026	22,374	22,403	23,911	25,056	26,125	26,911

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.
 **See text, footnote ** in text, and Appendix 2 for more details.
 **The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in New Hampshire fell from 17,569 in 1990 to 14,429 in 1998. Births to non-Hispanic whites accounted for 92 percent of total births in 1998.

When compared with the national average in 1998, New Hampshire's children got off to a healthier start to life on seven of eight indicators, including the percentage of births to mothers with less than 12 years of education (see figure). Moreover, the state saw a number of improvements during the 1990-1998 period, including a drop in the percentage of births to mothers who smoked during pregnancy. At the same time, there was a sharp rise in the proportion of births to unmarried women.


More specifically:

- **Teen births.** Throughout the 1990s, teen births accounted for a much smaller share of births in New Hampshire than in the U.S. as a whole. In 1998, 8 percent of births in New Hampshire were to teens, compared with 13 percent in the U.S. overall.
- **Repeat teen births.** The percentage of New Hampshire teen births to young women who were already mothers fell, from 18 percent in 1990 to 13 percent in 1996. Although the percentage then rose to 15 percent by 1998, New Hampshire remained below the national average for this indicator throughout the nine-year period.
- **Births to unmarried women.** The increase in the share of New Hampshire births to unmarried women outpaced the rise in the U.S. average. Unmarried women made up 24 percent of New Hampshire births in 1997 and 1998, compared with 17 percent in 1990. Throughout this time, however, births to unmarried women remained considerably less common in New Hampshire than in the U.S. overall.
- **Low maternal education.** The proportion of births in New Hampshire to women with less than 12 years of education in New Hampshire decreased from 12 percent in 1990 to 10 percent in 1998. Throughout this time, births to women with less than 12 years of education were only half as common in New Hampshire as in the nation as a whole.
- **Late or no prenatal care.** Throughout the 1990s, the share of New Hampshire births to women who received late or no prenatal care remained substantially below the U.S. average. In 1998, births to

women who received late or no prenatal care made up 2 percent of births in New Hampshire, compared with 4 percent in the U.S. as a whole.

- **Smoking during pregnancy.** The percentage of New Hampshire births to women who smoked during pregnancy declined from 21 percent in 1990 and 1991 to 17 percent in 1996-1998. Throughout this time, however, births to women who smoked during pregnancy remained more common in New Hampshire than in the U.S. as a whole.
- **Low-birthweight births.** Low-birthweight births were consistently less prevalent in New Hampshire than in the nation as a whole during the 1990s. In 1998, 6 percent of births in New Hampshire were low-birthweight, compared with a national average of 8 percent.
- **Preterm births.** Throughout the 1990-1998 period, the share of New Hampshire births that were preterm remained below the U.S. average. In 1998, preterm births accounted for 8 percent of births in New Hampshire, compared with 12 percent in the nation as a whole.

In summary, when compared with the U.S. average in 1998, the children of New Hampshire got off to a healthier start to life on all measures except maternal smoking. Further, there were several improvements during the 1990s, including drops in the shares of births to women who smoked during pregnancy and of teen births that were repeat births. However, the state also saw a substantial rise in the proportion of births to unmarried women and increases in the percentages of preterm and low-birthweight births.



* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** Because Hispanic origin of births was not collected until 1993 and was not reliably reported until 1994, the racial/ethnic distribution of births is available only from 1994 and later.

Trends In Key Indicators, 1990-1998

Indicator**	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	7.2	7.1	6.7	6.8	7.0	7.6	7.4	7.8	7.7
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	18.2	18.0	18.5	17.3	15.9	16.3	12.6	14.1	14.7
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	16.9	18.3	19.2	20.6	22.1	22.2	23.4	23.8	24.1
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	12.4	12.1	11.7	11.0	11.1	11.1	10.5	10.9	10.3
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	2.5	2.5	2.1	1.6	1.9	1.8	1.5	1.8	1.9
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	20.8	20.7	19.6	18.5	18.2	17.5	16.8	17.4	16.6
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	4.9	4.9	5.3	5.0	5.1	5.5	4.8	5.8	5.7
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	6.8	7.2	7.3	7.7	7.5	8.2	7.0	8.2	8.1
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	N.A.	N.A.	N.A.	N.R.	13,848	13,659	13,350	13,332
	Black non-Hispanic	N.A.	N.A.	N.A.	N.R.	87	75	98	116
	Hispanic	N.A.	N.A.	N.A.	N.R.	187	214	231	256
	Other	N.A.	N.A.	N.A.	N.R.	984	717	542	639
	Total	17,569	16,341	15,990	15,436	15,106	14,665	14,520	14,313

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**N.A. = Not Available. See footnote ** in text.

***N.R. = Not Reliable. See footnote ** in text.

****The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



The yearly number of births in New Jersey declined from 122,289 in 1990 to 114,550 in 1998. At the same time, the number of births to Hispanic mothers increased. While Hispanic births accounted for 14 percent of all births in 1990, they accounted for 18 percent in 1998.*

During the 1990-1998 period, New Jersey saw a number of relatively small changes in measures of a healthy start to life, including an overall drop in the share of teen births to young women who were already mothers and a rise in the share of all births to unmarried women. Throughout this time, the percentages of New Jersey births to teens (see figure) and to women with less than 12 years of education were substantially below the respective national averages.

More specifically:

- **Teen births.** During the 1990s, teen births accounted for a much smaller share of births in New Jersey than in the U.S. as a whole. Throughout this time, 8 percent of births in New Jersey were to teens, compared with the U.S. average of 13 percent.
- **Repeat teen births.** The percentage of teen births to young women who were already mothers fluctuated during the 1990s but dropped overall, from 24 percent in 1990 to 20 percent by 1994, and remained relatively stable thereafter. The nation as a whole saw a similar overall downward trend.
- **Births to unmarried women.** The share of New Jersey births to unmarried women increased from 24 percent in 1990 to 28 percent in 1994 through 1998. This change mirrored the increase in the national average.
- **Low maternal education.** During the 1990-1998 period, births to women with less than 12 years of education remained markedly less common in New Jersey than in the U.S. as a whole. In 1998, women with less than 12 years of education accounted for 14 percent of births in New Jersey, compared with 22 percent in the U.S. overall.
- **Late or no prenatal care.** Births to mothers who received late or no prenatal care made up a relatively steady share of New Jersey births during the 1990-1998 period, despite a downward trend in the national average. In 1998, 5 percent of births in New Jersey were to mothers who received late or no prenatal care, compared with 4 percent in the U.S. as a whole.


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** The percentage of births to mothers who smoked during pregnancy in New Jersey decreased from 13 percent in 1990 to a low of 8 percent in 1994, and then rose to 11 percent by 1998. Despite a considerably stronger downward trend in the national average, New Jersey remained below the national average for this indicator throughout the nine-year period.

- **Low-birthweight births.** The share of New Jersey births that were low-birthweight remained close to the U.S. average throughout the 1990-1998 period. At 8 percent in 1998, New Jersey was just above the U.S. average.

- **Preterm births.** During the 1990s, the proportion of New Jersey births that were preterm fluctuated somewhat but underwent an overall rise, from 11 percent in 1990 through 1993 to 12 percent in 1998. On par with the U.S. average for this indicator in 1990, New Jersey was just above the U.S. average in 1998.

In summary, New Jersey saw a number of small changes in measures of a healthy start to life over the 1990-1998 period. On the one hand, there were increases in the shares of preterm births and births to unmarried women. On the other hand, the percentage of teen births that were repeat births declined somewhat, as did the proportion of all births to mothers who smoked during pregnancy. Moreover, the percentages of New Jersey births to teenagers and to women with less than 12 years of education remained well below the respective national averages throughout the nine-year span.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	8.4	8.4	8.0	7.9	8.2	8.2	7.7	7.8	7.7
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	23.6	22.6	23.9	21.8	20.1	20.0	19.9	20.6	20.3
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	24.3	26.3	26.4	27.1	28.1	27.6	28.0	28.0	28.3
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	15.2	15.3	14.9	14.6	14.6	14.6	14.3	14.0	13.9
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	4.5	4.6	4.6	4.4	4.5	4.2	4.6	4.7	4.6
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	13.2	11.9	10.3	9.4	8.5	8.9	12.0	12.1	11.2
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	7.0	7.4	7.2	7.6	7.6	7.6	7.7	7.9	8.0
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	10.7	10.8	10.8	11.1	11.5	10.5	11.5	12.2	12.5
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	78,423	76,192	74,348	72,747	71,802	69,375	67,216	66,244
	Black non-Hispanic	22,130	22,697	22,103	21,384	21,546	19,518	19,227	19,302
	Hispanic	16,972	17,500	17,609	17,650	18,083	18,835	19,451	19,635
	Other	4,764	5,017	5,849	5,905	6,070	7,100	8,412	8,174
	Total	122,289	121,406	119,909	117,686	117,501	114,828	114,306	113,279

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.



The yearly number of births in New Mexico remained steady during the 1990s. There were 27,402 births in 1990 and 27,318 births in 1998. The percentage of New Mexico births to Hispanics rose during this time, from 45 percent to 50 percent.*

During the 1990s, New Mexico saw two important improvements: a considerable decrease in the share of teen births to young women who were already mothers and a marked drop in the share of births to women receiving late or no prenatal care (see figure). At the same time, however, there was a dramatic increase in the proportion of births to unmarried women.

More specifically:

- **Teen births.** The percentage of teen births in New Mexico rose from 16 percent in 1990 to 18 percent in 1993, and then remained stable through 1998. In contrast, births to teenagers made up a steady 13 percent of births in the U.S. as a whole throughout the 1990s.
- **Repeat teen births.** Teen births to young women who were already mothers accounted for 23 percent of New Mexico teen births in the early 1990s and 20 percent in 1994 through 1998. This overall decline from 1990 to 1998 was similar to the downward trend in the U.S. as a whole, with New Mexico just below the U.S. average throughout this time.
- **Births to unmarried women.** Already above the national average in 1990, the percentage of New Mexico births to unmarried women rose sharply, outpacing a similar increase in the national average. Births to unmarried women accounted for 44 percent of New Mexico births in 1998, compared with 35 percent in 1990.
- **Low maternal education.** During the 1990s, births to mothers with less than 12 years of education ranged between 27 percent and 29 percent of New Mexico births. Throughout this period, New Mexico was higher than the U.S. average, which declined from 24 percent of total births in 1990 to 22 percent in 1998.
- **Late or no prenatal care.** Both New Mexico and the U.S. as a whole experienced a noteworthy decline in the percentage of births to women receiving late or no prenatal care during the 1990s. However, New Mexico was twice as high as the national average on this indicator in 1998, when 9 percent of New Mexico births were to mothers who


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

received late or no prenatal care, compared with the national average of 4 percent.

- **Smoking during pregnancy.** Despite fluctuations during the 1990s, the proportion of New Mexico births to mothers who smoked during pregnancy was 11 percent in both 1990 and 1998. The U.S. as a whole, in contrast, saw a substantial decline in maternal smoking over this period. Nevertheless, smoking during pregnancy remained less common in New Mexico than in the nation overall throughout this time.
- **Low-birthweight births.** The share of low-birthweight births in New Mexico was between 7 percent and 8 percent during the 1990s. In 1998, 8 percent of births were low-birthweight in both New Mexico and the U.S. overall.
- **Preterm births.** Preterm births accounted for 11 percent of New Mexico births throughout the 1990-1998 period, remaining close to the national average on this indicator throughout the nine-year span.

In summary, New Mexico saw sizable drops during the 1990s in the percentages of teen births that were repeat births and births to mothers receiving late or no prenatal care. Also, the proportion of New Mexico mothers who smoked during pregnancy was below the national average.

Nevertheless, there was a striking increase in the share of births to unmarried women, as well as a small increase in the proportion of births to teens. Moreover, the percentages of births to teens, to unmarried women, to women with less than 12 years of education, and to women receiving late or no prenatal care were well above the respective national averages during this time.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	New Mexico 16.3	16.5	17.0	17.8	18.0	18.4	17.8	17.9	18.2
	U.S. Average 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	New Mexico 22.9	23.3	22.9	21.9	20.2	20.1	19.7	19.6	19.8
	U.S. Average 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	New Mexico 35.4	37.6	39.5	41.4	41.7	42.6	42.1	43.5	44.0
	U.S. Average 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	New Mexico 27.2	27.6	29.0	28.8	28.1	28.0	28.0	27.1	28.2
	U.S. Average 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	New Mexico 12.8	11.3	9.9	9.3	7.9	8.1	8.0	7.6	8.5
	U.S. Average 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	New Mexico 11.3	11.8	11.8	11.1	9.9	10.2	11.3	10.3	11.0
	U.S. Average 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5-lbs)	New Mexico 7.4	7.1	7.2	7.3	7.3	7.5	7.5	7.8	7.6
	U.S. Average 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	New Mexico 11.0	10.5	10.6	11.3	11.2	11.2	10.9	11.0	11.3
	U.S. Average 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	10,527	10,563	10,331	10,155	10,185	9,914	10,085	9,545
	Black non-Hispanic	512	550	489	525	505	473	439	468
	Hispanic	12,193	12,592	12,957	13,054	12,984	12,900	13,255	13,322
	Other	4,170	4,095	4,145	4,118	3,917	3,633	3,449	3,527
	Total	27,402	27,800	27,922	27,852	27,591	26,920	27,228	26,871

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in New York State (including New York City) declined from 297,576 in 1990 to 258,207 in 1998. This decline was most evident among whites, who accounted for 48 percent of New York births in 1998, compared with 56 percent in 1990.*

Over the 1990-1998 period, New York saw substantial decreases in the percentage of teen births to young women who were already mothers and in the proportion of total births to women who received late or no prenatal care (see figure). Moreover, teen births were less common in New York than in the U.S. as a whole throughout the nine-year span.

More specifically:

- **Teen births.** The proportion of New York births to teens remained markedly below the U.S. average throughout the 1990-1998 period. In 1998, teenagers accounted for 9 percent of births in New York, compared with 13 percent in the U.S. as a whole.
- **Repeat teen births.** During the 1990s, there was a drop in the percentage of New York teen births to young women who were already mothers, from a high of 23 percent in 1991 to 18 percent in 1998. Throughout this time, repeat teen births made up a smaller share of teen births in New York than in the nation as a whole.
- **Births to unmarried women.** Births to unmarried women in New York City were overstated during 1990 through 1996 (New York City accounts for nearly half of all New York State's births).** Changes in coding procedures in 1997 in New York City have resulted in more reliable data for the city and the state. In 1998, 35 percent of New York births were to unmarried women.
- **Low maternal education.** Between 1992, when data on maternal education first became available, and 1998, women with less than 12 years of education accounted for a smaller share of births in New York than in the nation as a whole. In 1998, 19 percent of New York births were to women with less than 12 years of education, compared with a national average of 22 percent.
- **Late or no prenatal care.** The proportion of births to New York mothers who received late or no prenatal care decreased from 8 percent in 1990 to 5 percent in 1998. This change mirrored the downward trend


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** See Appendix 2 and New York City pages in *The Right Start: City Trends*.

for the U.S. as a whole. However, the percentage of births to New York women who received late or no prenatal care remained somewhat above the U.S. average throughout the nine-year period.

- **Smoking during pregnancy.** Data on smoking during pregnancy were available only for New York City births, not for the balance of the state.***
- **Low-birthweight births.** Throughout the 1990-1998 period, 8 percent of New York births were low-birthweight. New York was close to the U.S. average for this indicator in 1998.
- **Preterm births.** Preterm births accounted for 11 percent of New York births throughout the 1990s. In 1998, preterm births were slightly less prevalent in New York than in the nation as a whole, where 12 percent of births were preterm.

In summary, during the 1990s, the state of New York experienced marked reductions in the percentages of teen births to young women who were already mothers and total births to women who received late or no prenatal care. While the proportion of births to women who received late or no prenatal care remained somewhat above the national average during the 1990-1998 period, the share of New York births to teens remained substantially below the national average throughout this time.



*** See Appendix 2 for more details.

Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	New York 9.1	9.2	9.0	9.3	9.5	9.3	9.2	8.8	8.8
	U.S. Average 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	New York 22.5	22.8	22.2	22.2	21.3	20.8	20.0	19.1	18.3
	U.S. Average 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	New York** 33.0	34.1	34.8	37.2	37.6	37.9	39.6	35.2	34.9
	U.S. Average 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	New York*** N.A.	N.A.	18.8	19.2	19.3	19.6	19.5	19.1	19.2
	U.S. Average 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	New York 8.2	7.4	6.8	6.4	6.1	5.2	5.4	5.0	4.8
	U.S. Average 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	New York*** N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.
	U.S. Average 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	New York 7.6	7.9	7.6	7.7	7.6	7.6	7.7	7.8	7.8
	U.S. Average 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	New York 10.7	10.9	10.6	10.7	10.6	10.7	10.7	11.0	10.9
	U.S. Average 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	165,220	159,146	154,439	147,516	144,340	121,349	119,291	117,641
	Black non-Hispanic	56,998	56,986	55,549	53,848	53,155	49,730	48,096	47,172
	Hispanic	53,140	53,694	53,047	51,847	53,216	54,193	52,684	50,720
	Other	22,218	22,807	24,852	29,181	27,681	46,097	43,892	41,705
	Total	297,576	292,633	287,887	282,392	278,392	271,369	263,963	257,238

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**See text and footnote ** in text.

***N.A. = Not Available. See Appendix 2 for more details.

****The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

North Carolina

The yearly number of births in North Carolina increased from 104,525 in 1990 to 111,688 in 1998. The percentage of North Carolina births to Hispanics more than tripled during this time, rising from 2 percent of all births in 1990 to 7 percent in 1998.*

North Carolina experienced improvements in five measures of a healthy start to life over the 1990-1998 period. Particularly striking were a marked decline in the proportion of births to mothers who smoked during pregnancy and a substantial drop in the share of births to women who received late or no prenatal care (see figure).

More specifically:


- **Teen births.** The proportion of births to teenagers in North Carolina decreased from 16 percent in 1990 and 1991 to 14 percent in 1997 and 1998. Throughout the 1990s, teen births remained more common in North Carolina than in the nation as a whole, where a steady 13 percent of births were to teen mothers.
- **Repeat teen births.** The percentage of teen births to young women who were already mothers fluctuated during the 1990s, but declined overall, from 25 percent in 1990 to 22 percent in 1996 through 1998. Since the U.S. as a whole experienced a similar decline from 1990 to 1998, North Carolina remained close to the national average throughout this time.
- **Births to unmarried women.** There was a marked increase in the proportion of North Carolina births to unmarried women, from 29 percent in 1990 to 33 percent in 1998. North Carolina matched the U.S. average in 1998.
- **Low maternal education.** The percentage of North Carolina births to women with less than 12 years of education ranged from a high of 23 percent in 1990 and 1991 to a low of 21 percent in 1994-1995. At 22 percent in 1998, North Carolina matched the national average.
- **Late or no prenatal care.** The share of North Carolina births to women who received late or no prenatal care fell from 6 percent in 1990 to 3 percent in 1998. North Carolina remained below the national average for this indicator throughout the nine-year period.
- **Smoking during pregnancy.** There was a marked reduction in the percentage of North Carolina births to women who smoked during pregnancy from 21 percent in 1990 to 15 percent in 1997 and 1998.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

Despite this change, smoking during pregnancy remained more common in North Carolina than in the nation overall throughout the period.

- **Low-birthweight births.** The percentage of North Carolina babies born at a low birthweight remained just above the national average throughout the 1990s. In 1998, 9 percent of births in North Carolina were low-birthweight, compared with 8 percent in the U.S. as a whole.
- **Preterm births.** Preterm births were somewhat more prevalent in North Carolina than in the nation as a whole during the 1990s. In 1998, preterm births accounted for 13 percent of births in North Carolina, compared with a national average of 12 percent.

In summary, North Carolina experienced improvements on five of eight measures of a healthy start to life during the 1990s. These improvements include a dramatic reduction in the percentage of births to mothers receiving late or no prenatal care, as well as sizable decreases in the proportions of births to mothers who smoked during pregnancy and births to teen mothers. During the same time, there was an increase in the share of births to unmarried women.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	16.2	16.2	15.4	15.3	15.5	15.2	15.0	14.4	14.0	
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	24.9	25.5	26.1	23.4	20.6	19.8	22.0	21.5	22.3	
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	29.4	31.6	31.3	32.1	31.9	31.4	32.0	32.2	32.8	
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	22.7	23.2	22.1	21.6	21.1	21.1	21.5	21.6	21.7	
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	5.6	5.2	4.2	4.0	3.8	3.3	3.3	3.2	2.9	
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	20.7	19.7	18.6	17.8	17.1	15.9	15.8	15.2	14.9	
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	8.0	8.4	8.4	8.6	8.7	8.7	8.7	8.8	8.8	
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	12.5	12.8	12.5	12.9	12.5	12.6	12.5	12.7	12.7	
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	69,512	67,561	68,482	66,506	67,272	67,262	68,481	69,102	71,294
	Black non-Hispanic	30,726	30,381	30,262	29,435	27,807	26,833	27,029	27,376	28,153
	Hispanic	1,754	1,833	2,379	2,560	3,135	4,244	5,433	6,906	8,104
	Other	2,533	2,587	2,844	2,856	3,206	3,253	3,527	3,631	4,137
	Total	104,525	102,362	103,967	101,357	101,420	101,592	104,470	107,015	111,688

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

North Dakota

The yearly number of births in North Dakota fell from 9,250 in 1990 to 7,932 in 1998. Births to whites accounted for 85 percent of total births in 1998.* Births to American Indians (included in Other) accounted for 9 percent of North Dakota births in 1998.

When compared with the U.S. average, North Dakota's children got off to a healthier start to life on seven of eight indicators in 1998. Moreover, the state saw a marked drop during the 1990s in the share of teen births to young women who were already mothers (see figure). At the same time, there was a large increase in the percentage of births to unmarried women.


More specifically:

- **Teen births.** Teenagers consistently accounted for a smaller share of births in North Dakota than in the nation as a whole during the 1990s. In 1998, 10 percent of North Dakota births were to teenagers, compared with the U.S. average of 13 percent.
- **Repeat teen births.** The percentage of teen births to young women who were already mothers fluctuated, but underwent an overall drop, from 17 percent in 1990 to 12 percent in 1998. North Dakota remained well below the national average for this indicator throughout the nine-year period.
- **Births to unmarried women.** The dramatic increase in the share of North Dakota births to unmarried women outpaced the upward trend in the U.S. average. Unmarried women accounted for 27 percent of North Dakota births in 1998, compared with just 18 percent in 1990. Despite this change, births to unmarried women remained less common in North Dakota than in the U.S. overall throughout the 1990s.
- **Low maternal education.** Throughout the 1990-1998 period, the percentage of North Dakota births to women with less than 12 years of education was far below the national average. In 1998, births to women with less than 12 years of education accounted for 10 percent of births in North Dakota, compared with 22 percent in the nation as a whole.
- **Late or no prenatal care.** Births to women who received late or no prenatal care made up a smaller proportion of births in North Dakota than in the nation as a whole throughout the 1990-1998 period. In 1998, 3 percent of births in North Dakota were to women who received late or no prenatal care, compared with 4 percent in the U.S. as a whole.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** During the 1990s, the share of births to women who smoked during pregnancy fluctuated, undergoing an overall decline from 22 percent in 1990 to 19 percent in 1998. Smoking during pregnancy remained more common in North Dakota than in the U.S. overall throughout the 1990s.
- **Low-birthweight births.** There was an increase in the share of North Dakota births that were low-birthweight, from a low of 5 percent in 1991 to 7 percent in 1998. Despite this increase, the percentage of North Dakota babies born at a low birthweight remained below the U.S. average throughout the period.
- **Preterm births.** The proportion of North Dakota births that were preterm increased from 8 percent in 1990 and 1991 to 11 percent in 1998. Preterm births were slightly less common in North Dakota than in the U.S. overall in 1998.

In summary, when compared with the national average in 1998, the children of North Dakota got off to a healthier start to life on all but one measure: smoking during pregnancy. Furthermore, North Dakota saw a substantial decrease over the 1990-1998 period in the share of teen births that were repeat births. On the other hand, there was a striking rise in the share of North Dakota births to unmarried women, as well as increases in the percentages of preterm and low-birthweight births.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	8.6	8.7	9.3	9.6	9.4	9.6	9.6	9.1	9.8
<i>U.S. Average</i>	<i>12.8</i>	<i>12.9</i>	<i>12.7</i>	<i>12.8</i>	<i>13.1</i>	<i>13.1</i>	<i>12.9</i>	<i>12.7</i>	<i>12.5</i>
Percent of teen births to women who were already mothers	17.3	15.1	17.7	14.2	15.2	13.4	14.4	15.2	12.3
<i>U.S. Average</i>	<i>24.3</i>	<i>24.6</i>	<i>25.0</i>	<i>23.4</i>	<i>21.8</i>	<i>21.0</i>	<i>21.4</i>	<i>21.6</i>	<i>21.5</i>
Percent of total births to unmarried women	18.4	22.0	22.6	23.0	23.0	23.5	25.1	26.0	27.0
<i>U.S. Average</i>	<i>28.0</i>	<i>29.5</i>	<i>30.1</i>	<i>31.0</i>	<i>32.6</i>	<i>32.2</i>	<i>32.4</i>	<i>32.4</i>	<i>32.8</i>
Percent of total births to mothers with less than 12 years of education	9.2	10.1	9.6	9.8	9.3	9.5	9.0	8.7	10.0
<i>U.S. Average</i>	<i>23.8</i>	<i>23.9</i>	<i>23.6</i>	<i>23.3</i>	<i>22.9</i>	<i>22.6</i>	<i>22.4</i>	<i>22.1</i>	<i>21.9</i>
Percent of total births to mothers receiving late or no prenatal care	3.1	2.8	2.8	2.4	2.0	2.3	2.0	2.2	2.5
<i>U.S. Average</i>	<i>6.1</i>	<i>5.8</i>	<i>5.2</i>	<i>4.8</i>	<i>4.4</i>	<i>4.2</i>	<i>4.0</i>	<i>3.9</i>	<i>3.9</i>
Percent of total births to mothers who smoked during pregnancy	21.9	22.3	20.6	19.7	19.5	17.9	18.4	20.1	19.5
<i>U.S. Average</i>	<i>18.4</i>	<i>17.8</i>	<i>16.9</i>	<i>15.8</i>	<i>14.5</i>	<i>13.9</i>	<i>13.6</i>	<i>13.2</i>	<i>12.9</i>
Percent low-birthweight births (less than 5.5 lbs)	5.5	4.8	5.1	5.3	5.4	5.3	5.7	6.2	6.5
<i>U.S. Average</i>	<i>7.0</i>	<i>7.1</i>	<i>7.1</i>	<i>7.2</i>	<i>7.3</i>	<i>7.3</i>	<i>7.4</i>	<i>7.5</i>	<i>7.6</i>
Percent preterm births (less than 37 completed weeks of gestation)	8.3	8.3	9.0	9.1	9.3	10.1	9.3	10.2	10.6
<i>U.S. Average</i>	<i>10.6</i>	<i>10.8</i>	<i>10.7</i>	<i>11.0</i>	<i>11.0</i>	<i>11.0</i>	<i>11.0</i>	<i>11.4</i>	<i>11.6</i>
Number of births by Race and Ethnicity of mother	White non-Hispanic	8,160	7,734	7,657	7,544	7,513	7,375	7,100	6,707
	Black non-Hispanic	80	84	73	88	67	68	80	82
	Hispanic	123	110	119	122	116	147	170	152
	Other	887	959	962	936	888	989	1,003	991
	Total	9,250	8,887	8,811	8,690	8,584	8,347	8,353	7,932

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Ohio decreased from 166,913 in 1990 to 152,794 in 1998. In 1998, births to whites accounted for 81 percent of total births, while births to blacks accounted for 15 percent.

Over the 1990-1998 period, Ohio experienced a marked drop in the percentage of births to mothers who smoked during pregnancy, although this percentage remained well above the national average throughout the nine-year span (see figure). There was also a decrease in the proportion of teen births to young women who were already mothers and an increase in the share of births to unmarried women.

More specifically:

- **Teen births.** Teen births accounted for 14 percent of Ohio births in 1990 through 1995 and 13 percent of births thereafter. In 1998, Ohio was close to the U.S. average of 13 percent.
- **Repeat teen births.** The proportion of Ohio teen births to young women who already had a child fluctuated but underwent an overall drop, from 24 percent in 1990 to 21 percent by 1998. This change mirrored the overall decline in the national average.
- **Births to unmarried women.** The share of Ohio births to unmarried women rose from 29 percent in 1990 to 34 percent in 1997 and 1998. The nation as a whole saw a parallel upward trend.
- **Low maternal education.** Births to mothers with less than 12 years of education accounted for a smaller proportion of births in Ohio than in the U.S. overall throughout the 1990-1998 period. In 1998, 18 percent of births in Ohio were to mothers with less than 12 years of education, compared with 22 percent in the U.S. as a whole.
- **Late or no prenatal care.** It is not possible to interpret the trends in this indicator because of incomplete reporting of this information in Ohio.** In 1997 and 1998, however, the number of births to mothers who reported receiving late or no prenatal care was around 25 percent higher than in 1996.

- **Smoking during pregnancy.** Ohio experienced a substantial decrease in the percentage of births to mothers who smoked during pregnancy from 25 percent in 1990 to 19 percent in 1998. Despite this decrease,


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** See Appendix 2 for more details.

smoking during pregnancy remained considerably more common in Ohio than in the nation as a whole.

- **Low-birthweight births.** In both Ohio and the U.S. overall, low-birthweight babies accounted for 7 percent of births in 1990 and 8 percent in 1998.
- **Preterm births.** The percentage of preterm births in Ohio was similar to the national average throughout the 1990-1998 period. In 1998, preterm births accounted for 12 percent of births in both Ohio and the nation as a whole.

In summary, Ohio was close to the national average on most measures of a healthy start to life in 1998. Although the percentage of Ohio births to mothers who smoked during pregnancy remained higher than the national average, the state saw a sizable drop in this measure over the 1990-1998 period. In addition, there was a reduction in the share of Ohio teen births that were repeat births. However, the rise in the percentage of births to unmarried women is noteworthy.



177

176

Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	13.8	14.0	13.6	13.6	13.7	13.7	13.3	13.3	13.0
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	23.8	24.8	25.1	23.5	21.2	20.3	21.0	20.9	21.3
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	28.9	30.7	31.6	33.0	32.9	33.0	33.1	33.9	34.0
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	19.4	19.9	19.2	19.0	18.1	17.7	17.5	17.8	17.9
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	4.1	4.1	3.9	3.5	3.5	3.5	3.4	4.3	4.2
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	24.9	24.3	23.2	22.2	21.3	20.2	19.5	19.7	19.3
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	7.1	7.5	7.4	7.5	7.5	7.6	7.5	7.7	7.7
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	10.6	11.0	10.7	11.3	11.1	11.2	11.0	11.5	11.7
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	136,756	135,107	131,703	128,656	127,425	126,215	124,295	123,800
	Black non-Hispanic	25,809	26,227	25,880	25,321	23,505	22,702	21,876	22,286
	Hispanic	2,427	2,554	2,583	2,655	2,717	2,801	3,004	3,432
	Other	1,921	1,907	2,081	2,161	2,297	2,346	2,517	2,992
	Total	166,913	165,795	162,247	158,793	155,944	154,064	151,692	152,033

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**See text and Appendix 2.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Oklahoma

The yearly number of births in Oklahoma increased from 47,649 in 1990 to 49,461 in 1998. The number of births to Hispanics nearly doubled from 1991, when data for Hispanic births are first available to 1998, while the number of births to both whites and blacks decreased. Whites were the majority of Oklahoma births throughout this time, accounting for 71 percent of births in 1998.* Births to American Indians (included in Other) accounted for one in ten Oklahoma births in 1998.

Over the 1990-1998 period, Oklahoma experienced a marked decrease in the proportion of births to mothers who received late or no prenatal care but also a considerable rise in the percentage of births to unmarried women (see figure).

More specifically:

- **Teen births.** Throughout the 1990-1998 period, births to teenagers accounted for 16 to 17 percent of Oklahoma births, compared with the national average of 13 percent.
- **Repeat teen births.** Data on repeat teen births in 1997 and 1998 are not shown due to considerable underreporting of birth order.** In 1996, 20 percent of teen births were to young women who were already mothers. This is not significantly different from the U.S. average.
- **Births to unmarried women.** There was a steady rise in the percentage of Oklahoma births to unmarried women from 25 percent in 1990 to 33 percent in 1998. This increase brought Oklahoma from below the U.S. average in 1990 to matching the average in 1998.
- **Low maternal education.** During the 1990s, between 23 percent and 24 percent of births in Oklahoma were to women with less than 12 years of education. The state was just above the U.S. average of 22 percent in 1998.
- **Late or no prenatal care.** There was a marked decline in the share of births to Oklahoma mothers receiving late or no prenatal care that narrowed the gap between Oklahoma and the U.S. average during the 1990s. In 1998, births to mothers receiving late or no prenatal care accounted for 5 percent of Oklahoma births, compared with 4 percent in the U.S. overall.


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** If it is assumed that most of the births with birth order unknown are actually first-order births, the resulting percentage of repeat teen births for Oklahoma in 1997 and 1998 is comparable to that in most other states. See Appendix 2 for more details.

- **Smoking during pregnancy.** Oklahoma witnessed a slight reduction in maternal smoking from 1991, when data first became available, to 1998. However, the U.S. as a whole experienced a sharper drop in this measure, leaving Oklahoma above the national average throughout the nine-year period. In 1998, births to mothers who smoked during pregnancy accounted for 19 percent of Oklahoma births, compared with 13 percent of births in the nation overall.
- **Low-birthweight births.** Babies born at a low birthweight accounted for 7 percent of Oklahoma births throughout the 1990-1998 period. Oklahoma virtually matched the U.S. average throughout this time.
- **Preterm births.** Preterm births became a slightly larger share of Oklahoma births, rising from 10 percent in 1990 to 11 percent in 1998. Oklahoma was on par with the U.S. average in 1998.

In summary, Oklahoma witnessed three important improvements during the 1990-1998 period. There was a sizable reduction in the proportion of births to women who received late or no prenatal care, and smaller declines in the percentages of births to mothers who smoked during pregnancy and births to mothers with less than 12 years of education.

However, there was a substantial increase in the share of births to unmarried women and a rise in the percentage of births that were preterm. Moreover, the percentages of births to teens, to women who received late or no prenatal care, and to women who smoked during pregnancy were above the respective national averages in 1998.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	16.2	17.2	16.8	17.2	17.1	17.1	17.2	17.0	16.3
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers	22.7	24.4	24.1	23.0	21.0	20.2	20.0	N.R.	N.R.
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women	25.2	27.1	28.4	29.1	29.8	30.5	30.9	32.4	33.2
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education	23.6	23.7	23.4	23.5	23.3	23.3	22.9	23.3	23.0
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care	8.3	7.3	6.4	7.0	6.1	4.9	4.6	4.8	5.1
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy	N.A.	19.6	18.2	18.0	18.2	17.5	17.4	17.8	18.6
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)	6.6	6.6	6.7	6.7	7.0	7.0	7.4	7.3	7.2
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)	10.1	10.5	10.0	10.5	10.6	10.9	11.4	11.4	11.5
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother	White non-Hispanic	N.A.	35,684	35,320	34,074	33,595	33,727	34,750	35,059
	Black non-Hispanic	N.A.	5,135	5,124	4,896	4,734	4,469	4,459	4,688
	Hispanic	N.A.	1,862	2,045	2,133	2,260	2,356	2,881	3,226
	Other	N.A.	5,114	5,068	5,140	5,114	5,120	5,081	5,617
	Total	47,649	47,795	47,557	46,243	45,703	45,672	46,193	48,269

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**N.R. = Not Reliable. See text, footnote ** in text, and Appendix 2.

***N.A. = Not Available. See Appendix 2 for more details.

****The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Oregon

The yearly number of births in Oregon rose from 42,891 in 1990 to 45,273 in 1998. The yearly number of births to Hispanics more than doubled over this period, accounting for more than 14 percent of Oregon births in 1998, compared with 7 percent in 1990.

Oregon witnessed a remarkable reduction in the percentage of births to mothers who smoked during pregnancy over the 1990-1998 period. Furthermore, when compared with the national average, babies born at a low birthweight consistently accounted for a small share of Oregon births throughout the 1990s (see figure). At the same time, however, there was a substantial increase in the proportion of births to unmarried women.

More specifically:


- **Teen births.** Teen births accounted for 12 percent of Oregon births in 1990 through 1993 and 13 percent of births thereafter. Oregon matched the national average in the latter part of the 1990s, as a steady 13 percent of births in the nation as a whole were to teens throughout the nine-year period.
- **Repeat teen births.** There was an overall decline during the 1990s in the share of Oregon teen births to young women who were already mothers, from 21 percent in 1990 to 19 percent in 1994-1998. Oregon remained below the U.S. average throughout this time.
- **Births to unmarried women.** Oregon experienced a sizable increase in the proportion of births to unmarried women, from 26 percent in 1990 to 30 percent in 1998. The nation as a whole also saw an upward trend on this measure, with Oregon consistently below the U.S. average throughout the nine-year period.
- **Low maternal education.** Births to mothers with less than 12 years of education made up a slightly smaller share of births in Oregon than in the U.S. overall during the 1990s. While births to women with less than 12 years of education accounted for 21 percent to 22 percent of Oregon births during this time, they accounted for 22 percent to 24 percent of births in the nation overall.
- **Late or no prenatal care.** There was a small decline in the proportion of Oregon births to mothers who received late or no prenatal care, from 5 percent in 1990 and 1991 to 4 percent where it remained through to 1998. In 1998, Oregon was in line with the national average.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Oregon experienced a dramatic decline in maternal smoking over the 1990s. While births to women who smoked during pregnancy accounted for 22 percent of births in 1990, they accounted for 15 percent of births in 1998. Despite this change, Oregon remained above the U.S. average throughout the nine-year period.

- **Low-birthweight births.** Throughout the 1990s, babies born at a low birthweight were markedly less common in Oregon than in the U.S. as a whole. In 1998, low-birthweight babies accounted for 5 percent of Oregon births, compared with a national average of 8 percent.
- **Preterm births.** Preterm births became a slightly larger share of Oregon births, increasing from 8 percent in 1990 to 9 percent in 1998. Preterm babies continued to account for a considerably smaller percentage of births in Oregon than in the nation overall throughout this time.

In summary, when compared with the national average, the children of Oregon got off to a significantly healthier start to life on five of eight indicators. Although the state was above the national average for the percentage of births to mothers who smoked during pregnancy in 1998, Oregon experienced a substantial drop on this measure over the 1990-1998 period. The rise in the proportion of births to unmarried women is also noteworthy.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	12.0	12.3	12.4	12.4	12.8	13.0	13.2	12.5	12.5
<i>U.S. Average</i>	<i>12.8</i>	<i>12.9</i>	<i>12.7</i>	<i>12.8</i>	<i>13.1</i>	<i>13.1</i>	<i>12.9</i>	<i>12.7</i>	<i>12.5</i>
Percent of teen births to women who were already mothers	20.7	22.3	21.4	20.0	18.5	19.1	18.8	18.6	19.1
<i>U.S. Average</i>	<i>24.3</i>	<i>24.6</i>	<i>25.0</i>	<i>23.4</i>	<i>21.8</i>	<i>21.0</i>	<i>21.4</i>	<i>21.6</i>	<i>21.5</i>
Percent of total births to unmarried women	25.7	26.6	27.0	28.2	28.7	28.9	29.7	28.8	29.7
<i>U.S. Average</i>	<i>28.0</i>	<i>29.5</i>	<i>30.1</i>	<i>31.0</i>	<i>32.6</i>	<i>32.2</i>	<i>32.4</i>	<i>32.4</i>	<i>32.8</i>
Percent of total births to mothers with less than 12 years of education	21.6	21.9	21.5	21.3	21.6	21.3	21.6	20.5	20.6
<i>U.S. Average</i>	<i>23.8</i>	<i>23.9</i>	<i>23.6</i>	<i>23.3</i>	<i>22.9</i>	<i>22.6</i>	<i>22.4</i>	<i>22.1</i>	<i>21.9</i>
Percent of total births to mothers receiving late or no prenatal care	5.4	4.6	4.0	3.8	4.3	4.3	3.9	3.6	3.8
<i>U.S. Average</i>	<i>6.1</i>	<i>5.8</i>	<i>5.2</i>	<i>4.8</i>	<i>4.4</i>	<i>4.2</i>	<i>4.0</i>	<i>3.9</i>	<i>3.9</i>
Percent of total births to mothers who smoked during pregnancy	22.3	21.4	20.5	18.9	18.2	17.9	17.8	16.2	15.2
<i>U.S. Average</i>	<i>18.4</i>	<i>17.8</i>	<i>16.9</i>	<i>15.8</i>	<i>14.5</i>	<i>13.9</i>	<i>13.6</i>	<i>13.2</i>	<i>12.9</i>
Percent low-birthweight births (less than 5.5 lbs)	5.0	4.9	5.2	5.2	5.3	5.5	5.3	5.5	5.4
<i>U.S. Average</i>	<i>7.0</i>	<i>7.1</i>	<i>7.1</i>	<i>7.2</i>	<i>7.3</i>	<i>7.3</i>	<i>7.4</i>	<i>7.5</i>	<i>7.6</i>
Percent preterm births (less than 37 completed weeks of gestation)	8.0	8.3	8.2	8.6	8.6	8.9	8.4	9.1	9.1
<i>U.S. Average</i>	<i>10.6</i>	<i>10.8</i>	<i>10.7</i>	<i>11.0</i>	<i>11.0</i>	<i>11.0</i>	<i>11.0</i>	<i>11.4</i>	<i>11.6</i>
Number of births by Race and Ethnicity of mother	White non-Hispanic	37,045	36,324	35,577	34,726	34,499	34,781	34,990	35,138
	Black non-Hispanic	912	956	943	890	931	860	873	914
	Hispanic	2,974	3,285	3,561	4,002	4,357	5,002	5,458	6,501
	Other	1,960	1,934	1,954	1,958	2,050	2,168	2,337	2,672
	Total	42,891	42,499	42,035	41,576	41,837	42,811	43,658	43,809

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Pennsylvania

The yearly number of births in Pennsylvania decreased substantially from 171,961 in 1990 to 145,899 in 1998. While this decrease was evident among both blacks and whites, the number of births to Hispanics increased slightly over this period, and accounted for 5 percent of all Pennsylvania births in 1998.*

Pennsylvania experienced improvements on five of eight measures of a healthy start to life during the 1990s. The share of births to mothers with less than 12 years of education declined from 1990 to 1998 and was consistently below the national average throughout this time (see figure). During the same time period, there was a sizable increase in the percentage of births to unmarried women.


More specifically:

- **Teen births.** Births to teenagers accounted for 11 percent of Pennsylvania births in 1990 through 1996, and 10 percent in 1997 and 1998. These values remained below the U.S. as a whole, where a steady 13 percent of births were to teens throughout the 1990-1998 period.
- **Repeat teen births.** The percentage of Pennsylvania teen births to young women who were already mothers decreased overall, from 24 percent in 1990 and 1991 to 20 percent in 1998. Pennsylvania was just below the U.S. average in both 1990 and 1998.
- **Births to unmarried women.** Births to unmarried women became a markedly larger proportion of Pennsylvania births, rising from 29 percent in 1990 to 33 percent in 1997 and 1998. The U.S. as a whole experienced a similar rise on this measure, and Pennsylvania was close to the national average throughout this period.
- **Low maternal education.** Births to women with less than 12 years of education made up a substantially smaller share of births in Pennsylvania than in the U.S. as a whole throughout the 1990s. During this time, 15 percent to 17 percent of Pennsylvania births were to mothers with less than 12 years of education, compared with 22 percent to 24 percent of births in the nation overall.
- **Late or no prenatal care.** The proportion of Pennsylvania births to women receiving late or no prenatal care decreased from 5 percent in the early 1990s to 4 percent from 1994 through 1998. The U.S. as a whole also experienced a decline on this measure from 1990 to 1998.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Pennsylvania saw a notable reduction in the percentage of births to women who smoked during pregnancy, from 21 percent in 1990 through 1992 to 17 percent in 1998. However, Pennsylvania was above the national average throughout this time.
- **Low-birthweight births.** Both Pennsylvania and the U.S. overall witnessed a slight rise in the share of babies born at a low birthweight during the 1990s, from 7 percent in 1990 to 8 percent in 1998.
- **Preterm births.** Preterm births accounted for 10 percent of Pennsylvania births in 1990-1992 and around 11 percent thereafter. Throughout this time, Pennsylvania was just below the U.S. average, which also rose slightly from 1990 to 1998.

In summary, Pennsylvania experienced improvements in five measures of a healthy start to life over the 1990-1998 period, including declines in the percentages of births to mothers who smoked during pregnancy, teen births that were repeat births, and births to women receiving late or no prenatal care. However, when compared with the 1998 U.S. average, the proportion of births to mothers who smoked during pregnancy remained high. There was also a notable increase in the proportion of births to unmarried women during this time.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens									
Pennsylvania	10.9	10.8	10.5	10.5	10.8	10.8	10.6	10.4	10.3
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers									
Pennsylvania	23.6	24.0	24.5	23.6	22.1	20.8	20.4	20.6	20.4
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women									
Pennsylvania	28.6	30.4	31.6	32.2	32.8	32.4	32.3	32.8	32.8
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education									
Pennsylvania	16.4	16.7	16.3	16.2	16.0	15.5	15.4	15.3	15.1
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care									
Pennsylvania	5.4	5.4	5.2	4.7	4.4	3.9	3.6	4.4	3.5
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy									
Pennsylvania	20.9	20.7	20.5	19.9	19.3	18.6	18.1	17.9	17.5
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)									
Pennsylvania	7.1	7.3	7.2	7.4	7.4	7.4	7.5	7.6	7.6
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)									
Pennsylvania	10.3	10.4	10.3	10.6	10.7	10.6	10.5	10.7	10.9
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother									
White non-Hispanic	138,116	134,279	130,265	126,991	124,230	120,544	117,509	113,297	114,265
Black non-Hispanic	25,221	25,296	24,987	24,013	22,956	21,062	20,237	19,947	20,272
Hispanic	5,636	5,956	5,949	6,302	6,288	6,572	6,737	6,623	6,897
Other	2,988	3,320	3,424	3,456	3,597	3,672	3,855	4,357	4,465
Total	171,961	168,851	164,625	160,762	157,071	151,850	148,338	144,224	145,899

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Rhode Island

The yearly number of births in Rhode Island declined from 15,195 in 1990 to 12,599 in 1998. At the same time, the share of total births to Hispanics grew from 10 percent in 1990 to 15 percent in 1998.

Over the 1990-1998 period, there was a marked drop in the share of births to women who smoked during pregnancy in Rhode Island. Moreover, when compared with the national average in 1998, the children of Rhode Island got off to a healthier start to life on five of eight measures, including the proportion of births to women who received late or no prenatal care (see figure). However, the share of births to unmarried women climbed above the national average during the 1990s.

More specifically:

- **Teen births.** During the 1990-1998 period, teen births consistently made up a somewhat smaller share of births in Rhode Island than in the U.S. as a whole. In 1998, 11 percent of births in Rhode Island were to teenagers, compared with 13 percent in the U.S. overall.
- **Repeat teen births.** Although the percentage of teen births to young women who were already mothers fluctuated, this percentage was similar in 1990 and 1998. In 1998, Rhode Island was below the U.S. average, as 19 percent of teen births in Rhode Island were repeat teen births, compared with 22 percent in the nation overall.
- **Births to unmarried women.** Rhode Island saw a striking rise in the proportion of births to unmarried women, from 26 percent in 1990 to 34 percent in 1998. The share of Rhode Island births to unmarried women went from below the national average in 1990 to above it by 1998.
- **Low maternal education.** There was a drop in the percentage of births to women with less than 12 years of education in Rhode Island, from a high of 20 percent in 1991 to 15 percent in 1998. Throughout this time, births to women with less than 12 years of education were notably less common in Rhode Island than in the U.S. as a whole.
- **Late or no prenatal care.** Throughout the 1990-1998 period, Rhode Island was consistently below the U.S. average in the proportion of births to women who received late or no prenatal care. In 1998, 2

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.


** Hispanic origin was not reported or unknown for 13 percent of total births in 1998—up from only 5 percent in 1990. Consequently, the distribution of births by Hispanic origin is less reliable during the late 1990s. Over 90 percent of births with unknown Hispanic origin in 1998 were to white women.

percent of Rhode Island births were to women who received late or no prenatal care, compared with a national average of 4 percent.

- **Smoking during pregnancy.** There was a sizable drop in the share of Rhode Island births to women who smoked during pregnancy, from 22 percent in 1990 to 15 percent in 1998. However, smoking during pregnancy remained more common in Rhode Island than in the U.S. overall throughout the 1990s.
- **Low-birthweight births.** The share of births that were low-birthweight in Rhode Island rose from 6 percent in the early 1990s to 8 percent in 1998. This change brought Rhode Island from below the national average in 1990 to matching it by 1998.

- **Preterm births.** The percentage of Rhode Island births that were preterm fluctuated during the 1990s, increasing from a low of 9 percent in 1991 to 11 percent in 1998. There was a similar rise in the national average, and preterm births remained somewhat less prevalent in Rhode Island than in the U.S. as a whole throughout the 1990-1998 period.

In summary, Rhode Island saw a considerable increase during the 1990s in the percentage of births to unmarried women. On the other hand, there was a sharp drop in the share of births to women who smoked during pregnancy. Furthermore, when compared with the national average in 1998, Rhode Island's children got off to a healthier start to life on five of eight measures, including the proportion of births to women who received late or no prenatal care.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens									
Rhode Island	10.5	9.8	9.8	10.4	10.6	10.1	10.4	10.7	10.5
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers									
Rhode Island	21.9	26.4	22.5	22.7	21.1	20.3	21.2	19.9	19.2
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women									
Rhode Island	26.3	27.6	29.6	31.7	32.1	31.1	33.3	33.1	33.9
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education									
Rhode Island	19.2	19.7	19.3	18.3	17.7	16.5	16.5	16.4	15.5
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care									
Rhode Island	2.0	1.7	2.0	1.6	1.6	1.3	1.5	1.7	1.5
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy									
Rhode Island	22.4	21.5	20.4	19.0	18.5	16.6	16.4	16.2	15.2
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)									
Rhode Island	6.2	6.0	6.3	6.5	6.5	6.8	6.9	7.4	7.6
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)									
Rhode Island	9.6	8.8	9.6	9.5	9.9	9.5	9.5	10.0	10.7
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother									
White non-Hispanic	11,300	11,044	10,408	9,624	9,231	8,256	8,210	7,742	7,743
Black non-Hispanic	1,019	1,022	1,017	925	946	683	750	777	828
Hispanic	1,560	1,400	1,553	1,594	1,623	1,554	1,667	1,780	1,865
Other	1,316	1,268	1,522	1,833	1,666	2,283	2,025	2,156	2,163
Total	15,195	14,734	14,500	13,976	13,466	12,776	12,652	12,455	12,599

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**See footnote ** in text.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

South Carolina

The yearly number of births in South Carolina decreased from 58,610 in 1990 to 50,926 in 1995, and then rose to 53,877 in 1998. White births accounted for 61 percent of total births in 1998, while black births accounted for 35 percent.*

During the 1990-1998 period, there were improvements in five of eight measures of a healthy start to life, including a dramatic reduction in the percentage of births to women who received late or no prenatal care (see figure). However, when compared with the national average in 1998, the children of South Carolina got off to a significantly less healthy start to life on six of eight measures.

More specifically:


- **Teen births.** Despite a slight overall decrease in the percentage of births to teens, teenagers continued to account for a markedly larger share of births in South Carolina than in the U.S. as a whole throughout the 1990s. In 1998, 16 percent of births in South Carolina were to teen mothers, compared with 13 percent in the U.S. overall.
- **Repeat teen births.** Births to young women who already had a child became a smaller share of all teen births in South Carolina during the 1990s, falling from a high of 27 percent in 1992 to a low of 19 percent in 1995. The percentage then rose to 21 percent in 1998, similar to the 1998 national average.
- **Births to unmarried women.** There was a sizable increase in the proportion of births to unmarried women in South Carolina that mirrored the rise in the U.S. average. Unmarried women accounted for 39 percent of South Carolina births in 1998, compared with 33 percent in 1990. South Carolina remained above the U.S. average for this indicator throughout the nine-year period.
- **Low maternal education.** The share of South Carolina births to women with less than 12 years of education was similar to the national average throughout the 1990s. In 1998, these births made up 22 percent of total births both in South Carolina and in the nation as a whole.
- **Late or no prenatal care.** The percentage of births to women who received late or no prenatal care fell by nearly half, from 8 percent in 1990 and 1991 to 4 percent in 1998. This change closed the gap between South Carolina and the national average.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** During the 1990s, there was a large drop in the share of births to women who smoked during pregnancy in South Carolina, mirroring the downward trend in the U.S. average. Births to women who smoked during pregnancy made up 14 percent of South Carolina births in 1998, compared with 19 percent in 1990.

- **Low-birthweight births.** Low-birthweight births remained more prevalent in South Carolina than in the U.S. as a whole throughout the 1990-1998 period. In 1998, 10 percent of births in South Carolina were low-birthweight, compared with 8 percent in the U.S. overall.
- **Preterm births.** Preterm births were consistently more common in South Carolina than in the nation as a whole during the 1990s. In 1998, preterm births accounted for 14 percent of births in South Carolina, compared with 12 percent in the nation overall.

In summary, South Carolina saw improvements on five of eight measures of a healthy start to life during the 1990-1998 period. Particularly striking were drops in the percentages of births to women who smoked during pregnancy and births to women who received late or no prenatal care. At the same time, however, there was an increase in the proportion of births to unmarried women. Moreover, the percentages of teen, low-birthweight, and preterm births remained higher than the respective national averages in 1998.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens	17.1	17.0	16.6	16.2	17.0	17.3	16.8	16.4	16.0
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
South Carolina.	26.0	26.4	26.9	24.4	20.6	19.4	20.5	20.8	21.3
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
South Carolina	32.7	34.7	35.5	36.0	36.8	37.4	37.3	38.0	38.8
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
South Carolina	24.2	24.6	24.0	23.1	22.5	22.1	22.1	22.1	22.0
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
South Carolina	7.9	8.1	6.9	6.3	5.6	4.8	4.7	4.7	4.2
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
South Carolina	19.5	18.0	17.1	16.5	15.0	14.8	14.3	13.9	13.7
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
South Carolina	8.7	9.2	9.0	9.3	9.2	9.3	9.2	9.2	9.5
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
South Carolina	12.3	12.9	12.8	12.6	12.4	12.5	12.3	12.6	13.6
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
White non-Hispanic	35,055	34,039	33,379	32,116	31,397	31,127	31,427	31,971	32,885
Black non-Hispanic	22,336	22,404	21,535	20,482	19,380	18,393	17,973	18,382	18,840
Hispanic	571	599	625	617	643	763	973	1,129	1,307
Other	648	530	653	620	623	643	744	732	845
Total	58,610	57,572	56,192	53,835	52,043	50,926	51,117	52,214	53,877

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in South Dakota was similar in 1990 and 1998. There were 10,999 births in 1990 and 10,288 births in 1998. Throughout this time, births to whites accounted for the vast majority of South Dakota births, including 80 percent in 1998.* In 1998, American Indians (included in Other) accounted for 17 percent of South Dakota births.

Over the 1990-1998 period, South Dakota saw a sizable drop in the share of teen births to young women who were already mothers and a substantial increase in the share of all births to unmarried women (see figure). When compared with the 1998 U.S. average, the children of South Dakota got off to a significantly healthier start to life on five of the seven available measures.

More specifically:


- **Teen births.** Teenagers accounted for between 11 percent and 12 percent of births in South Dakota during the 1990-1998 period. At 12 percent in 1998, South Dakota was not significantly different from the national average.
- **Repeat teen births.** The percentage of South Dakota teen births to young women who were already mothers fluctuated during the 1990s, undergoing a marked overall drop that outpaced a similar trend in the U.S. average. Repeat teen births made up 17 percent of South Dakota teen births in 1998, compared with the U.S. average of 22 percent.
- **Births to unmarried women.** The proportion of births to unmarried women in South Dakota increased from 23 percent in 1990 to 32 percent in 1998, outpacing the increase in the national average. By 1998, the proportion of births to unmarried women in South Dakota virtually matched the U.S. average.
- **Low maternal education.** Throughout the 1990-1998 period, births to women with less than 12 years of education made up a substantially smaller share of total births in South Dakota than in the U.S. as a whole. In 1998, 16 percent of South Dakota births were to women with less than 12 years of education, compared with the U.S. average of 22 percent.
- **Late or no prenatal care.** South Dakota experienced a marked decrease in the percentage of births to mothers who received late or no prenatal care, mirroring the downward trend in the U.S. average. Births

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

to mothers who received late or no prenatal care dropped from 5 percent in 1990-1993 to 3 percent by 1998.

- **Smoking during pregnancy.** Data on smoking during pregnancy were not available for South Dakota births.
- **Low-birthweight births.** Throughout the 1990-1998 period, South Dakota was below the national average in the percentage of babies born at a low birthweight. In 1998, 6 percent of births in South Dakota were low-birthweight, compared with 8 percent in the nation as a whole.
- **Preterm births.** There was an increase in the percentage of South Dakota births that were preterm, from 9 percent in 1990-1995 to 11 percent in 1998. Despite this change, preterm births remained slightly less prevalent in South Dakota than in the U.S. overall in 1998.

In summary, over the 1990-1998 period, South Dakota experienced marked improvements in two important measures of a healthy beginning to life: the proportion of teen births that were repeat births and the share of all births to women who received late or no prenatal care. Furthermore, when compared with the 1998 U.S. average, the children of South Dakota got off to a significantly healthier start to life on five of seven available indicators. However, there were also rises in the percentages of births to unmarried women and births that were preterm during the nine-year period.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	10.8	11.0	11.4	11.1	11.4	11.4	11.5	12.2	12.0	
<i>U.S. Average</i>	<i>12.8</i>	<i>12.9</i>	<i>12.7</i>	<i>12.8</i>	<i>13.1</i>	<i>13.1</i>	<i>12.9</i>	<i>12.7</i>	<i>12.5</i>	
Percent of teen births to women who were already mothers	23.7	19.8	20.5	19.3	16.6	19.0	16.6	16.0	17.2	
<i>U.S. Average</i>	<i>24.3</i>	<i>24.6</i>	<i>25.0</i>	<i>23.4</i>	<i>21.8</i>	<i>21.0</i>	<i>21.4</i>	<i>21.6</i>	<i>21.5</i>	
Percent of total births to unmarried women	22.9	24.8	26.6	27.7	27.7	28.0	29.5	31.1	32.0	
<i>U.S. Average</i>	<i>28.0</i>	<i>29.5</i>	<i>30.1</i>	<i>31.0</i>	<i>32.6</i>	<i>32.2</i>	<i>32.4</i>	<i>32.4</i>	<i>32.8</i>	
Percent of total births to mothers with less than 12 years of education	15.2	15.9	16.0	15.1	15.6	15.0	15.4	15.8	15.5	
<i>U.S. Average</i>	<i>23.8</i>	<i>23.9</i>	<i>23.6</i>	<i>23.3</i>	<i>22.9</i>	<i>22.6</i>	<i>22.4</i>	<i>22.1</i>	<i>21.9</i>	
Percent of total births to mothers receiving late or no prenatal care	4.9	4.7	5.0	4.8	3.9	3.6	3.5	3.4	3.2	
<i>U.S. Average</i>	<i>6.1</i>	<i>5.8</i>	<i>5.2</i>	<i>4.8</i>	<i>4.4</i>	<i>4.2</i>	<i>4.0</i>	<i>3.9</i>	<i>3.9</i>	
Percent of total births to mothers who smoked during pregnancy	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	
<i>U.S. Average</i>	<i>18.4</i>	<i>17.8</i>	<i>16.9</i>	<i>15.8</i>	<i>14.5</i>	<i>13.9</i>	<i>13.6</i>	<i>13.2</i>	<i>12.9</i>	
Percent low-birthweight births (less than 5.5 lbs)	5.1	5.4	5.2	5.5	5.9	5.6	5.8	5.5	5.8	
<i>U.S. Average</i>	<i>7.0</i>	<i>7.1</i>	<i>7.1</i>	<i>7.2</i>	<i>7.3</i>	<i>7.3</i>	<i>7.4</i>	<i>7.5</i>	<i>7.6</i>	
Percent preterm births (less than 37 completed weeks of gestation)	8.9	8.8	9.0	8.8	9.3	9.5	10.0	10.2	10.8	
<i>U.S. Average</i>	<i>10.6</i>	<i>10.8</i>	<i>10.7</i>	<i>11.0</i>	<i>11.0</i>	<i>11.0</i>	<i>11.0</i>	<i>11.4</i>	<i>11.6</i>	
Number of births by Race and Ethnicity of mother	White non-Hispanic	9,076	9,030	9,008	8,721	8,661	8,590	8,520	8,284	8,252
	Black non-Hispanic	68	54	74	76	76	80	89	82	
	Hispanic	113	97	109	122	124	146	141	153	
	Other	1,742	1,765	1,827	1,800	1,646	1,672	1,727	1,801	
	Total	10,999	10,946	11,018	10,719	10,507	10,475	10,173	10,288	

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**N.A. = Not Available. See Appendix 2 for more details.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Tennessee decreased from 74,962 in 1990 to 73,017 in 1993 before increasing to 77,396 by 1998. While the number of births to both whites and Hispanics increased from 1990 to 1998, births to blacks decreased slightly. * During the 1990s, whites accounted for about three out of every four births in Tennessee.

During the 1990s, Tennessee saw improvements on five of eight measures of a healthy start to life, including a sizable decline in the percentage of births to mothers who smoked during pregnancy (see figure). However, there was also a sharp increase in the share of births to unmarried women.

More specifically:


- **Teen births.** Teen births became a smaller share of Tennessee births, decreasing from 18 percent in 1990 and 1991 to 16 percent in 1997 and 1998. However, Tennessee remained substantially above the steady U.S. average of 13 percent throughout the 1990s.
- **Repeat teen births.** The proportion of teen births to young women who were already mothers in Tennessee fell from a high of 26 percent in 1991 to 22 percent in 1996 through 1998. Tennessee was on par with the national average in 1998.
- **Births to unmarried women.** Over the 1990-1998 period, there was a marked rise in the percentage of Tennessee births to unmarried women that paralleled a similar trend in the nation as a whole. While births to unmarried women accounted for 30 percent of Tennessee births in 1990, they accounted for 35 percent in 1998.
- **Low maternal education.** Tennessee experienced a sizable reduction in the share of births to women with less than 12 years of education, from 26 percent in 1990 and 1991 to 22 percent in 1995 through 1998. This decrease closed the gap between Tennessee and the national average over the nine-year period.
- **Late or no prenatal care.** Births to mothers who received late or no prenatal care accounted for 5 percent of Tennessee births in 1990 and 4 percent of births from 1991 through 1998. Births to women receiving late or no prenatal care made up a slightly smaller percentage of births in Tennessee than in the nation as a whole in 1998.
- **Smoking during pregnancy.** There was a striking decline in the proportion of Tennessee births to mothers who smoked during

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

pregnancy over the 1990s, from 22 percent in 1990 to 17 percent in 1998. Despite this change, Tennessee remained above the national average throughout the nine-year period.

- **Low-birthweight births.** Low-birthweight births accounted for 8 percent of Tennessee births in 1990 and 9 percent of births thereafter. Babies born at a low birthweight made up a consistently larger share of births in Tennessee than in the nation as a whole throughout this time.
- **Preterm births.** Preterm births were more common in Tennessee than in the U.S. overall throughout the 1990s. In 1998, 14 percent of babies in Tennessee were preterm, compared with 12 percent in the U.S. overall.

In summary, when compared with the U.S. average, the children of Tennessee got off to a significantly less healthy start to life on five of eight indicators in 1998. Moreover, there was a substantial increase in the share of births to unmarried women over the 1990-1998 period. At the same time, however, Tennessee saw a number of improvements—including sizable reductions in the percentages of teen births to young women who were already mothers, births to mothers with less than 12 years of education, and births to mothers who smoked during pregnancy.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	Tennessee 17.6	17.7	16.9	16.9	17.4	16.9	16.8	16.3	15.9	
	U.S. Average 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	Tennessee 24.8	26.0	25.9	24.3	22.6	21.4	22.0	21.8	22.0	
	U.S. Average 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	Tennessee 30.2	32.2	32.7	33.6	33.4	33.1	33.4	34.1	34.9	
	U.S. Average 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	Tennessee 25.9	26.1	25.3	24.0	23.4	22.4	22.3	22.1	22.1	
	U.S. Average 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	Tennessee 4.7	4.4	4.0	3.6	3.8	3.6	3.7	3.9	3.6	
	U.S. Average 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	Tennessee 21.9	21.4	20.8	19.5	18.2	18.1	17.6	17.4	16.8	
	U.S. Average 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	Tennessee 8.2	8.8	8.5	8.8	8.8	8.7	8.8	8.8	9.1	
	U.S. Average 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	Tennessee 12.6	13.0	12.6	13.4	13.3	13.0	12.6	13.1	13.6	
	U.S. Average 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	56,178	55,525	54,700	53,896	54,783	54,875	55,120	55,288	57,345
	Black non-Hispanic	17,582	17,797	17,490	17,550	16,576	16,130	15,985	16,297	16,853
	Hispanic	444	481	595	694	845	1,111	1,425	1,721	1,997
	Other	758	707	829	877	987	1,224	1,224	1,172	1,201
	Total	74,962	74,510	73,614	73,017	73,191	73,173	73,754	74,478	77,396

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Texas increased from 316,423 in 1990 to 342,283 in 1998. This rise was due to a 30 percent increase in the number of births to Hispanics, who accounted for 44 percent of Texas births in 1998, compared with 37 percent in 1990.* The number of births to both whites and blacks decreased during this time.

During the 1990s, Texas experienced sizable declines in the share of teen births to young women who were already mothers and in the percentages of all births to mothers who smoked during pregnancy and to mothers who received late or no prenatal care. However, the percentage of Texas births to women with less than 12 years of education remained markedly above the national average throughout the 1990-1998 period (see figure).

More specifically:

- **Teen births.** Teen births were more common in Texas than in the nation as a whole throughout the 1990-1998 period. During this time, between 16 percent and 17 percent of Texas births were to teens, compared with the steady U.S. average of 13 percent.
- **Repeat teen births.** Texas experienced a reduction in the proportion of teen births to young women who were already mothers, from 27 percent in 1990 to around 24 percent in 1996 through 1998. Despite this progress, Texas remained above the national average on this indicator throughout this time.
- **Births to unmarried women.** Births to unmarried women were substantially underreported during 1990-1993 in Texas.** Births to unmarried women made up 29 percent of total births in 1994 and increased to 31 percent by 1998. Texas was slightly below the national average in 1998.
- **Low maternal education.** Births to women with less than 12 years of education made up a substantially larger share of births in Texas than in the U.S. as a whole throughout the 1990-1998 period. In 1998, 33 percent of Texas births were to mothers with less than 12 years of education, compared with 22 percent of births in the nation overall.
- **Late or no prenatal care.** The percentage of Texas births to women who received late or no prenatal care fell by nearly half, from 10 percent


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

** Trends can therefore be examined meaningfully only since 1994. See Appendix 2 for more details.

in 1990 and 1991 to 5 percent by 1998. This decline narrowed the gap between Texas and the national average on this indicator.

- **Smoking during pregnancy.** Both Texas and the nation as a whole saw a marked drop in the proportion of births to women who smoked during pregnancy over the 1990-1998 period, with Texas well below the U.S. average throughout this time. In 1998, 7 percent of births in Texas were to women who smoked during pregnancy, compared with the U.S. average of 13 percent.
- **Low-birthweight births.** Low-birthweight births accounted for 7 percent of Texas births during the 1990s. Texas was close to the national average throughout this time.
- **Preterm births.** During the 1990s, the percentage of preterm births ranged between 11 percent and 12 percent of Texas births. Texas was just above the national average in 1998.

In summary, Texas saw important declines over the 1990-1998 period in the percentage of teen births that were repeat births and in the shares of all births to mothers who received late or no prenatal care and to mothers who smoked during pregnancy. In addition, births to mothers who smoked during pregnancy made up a substantially smaller proportion of births in Texas than in the U.S. overall throughout the 1990s. However, when compared with the national average, Texas' children got off to a less healthy start to life on five of eight indicators in 1998.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens									
Texas	15.6	16.0	15.9	16.1	16.5	16.6	16.2	16.1	16.1
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers									
Texas	26.8	25.9	26.2	24.2	23.4	23.2	23.9	24.4	24.1
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women									
Texas**	N.R.	N.R.	N.R.	N.R.	28.9	30.0	30.4	30.7	31.5
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education									
Texas	33.5	34.2	34.4	34.2	34.1	34.1	33.9	33.6	33.1
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care									
Texas	10.0	10.1	9.2	8.4	6.1	5.7	5.4	5.1	5.3
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy									
Texas	10.4	9.6	9.3	8.6	8.4	7.9	7.7	7.4	7.2
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)									
Texas	6.9	7.1	7.0	7.1	7.0	7.1	7.2	7.3	7.4
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)									
Texas	11.2	11.4	11.0	11.4	11.6	11.4	11.7	12.0	12.4
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother									
White non-Hispanic	150,304	146,199	143,915	141,086	139,105	137,816	138,925	137,580	139,980
Black non-Hispanic	43,121	42,876	42,786	41,417	39,907	38,434	38,497	38,928	39,631
Hispanic	115,845	121,234	126,357	131,293	133,125	137,131	142,831	146,580	151,487
Other	7,153	7,437	7,787	8,275	8,977	9,372	10,153	10,886	11,185
Total	316,423	317,746	320,845	322,071	321,114	322,753	330,406	333,974	342,283

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**N.R. = Not Reliable. See text, footnote ** in text, and Appendix 2 for more details.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Utah rose from 36,277 in 1990 to 45,165 in 1998. During this time, the share of births to Hispanics nearly doubled, from 6 percent in 1990 to 11 percent in 1998.*

When compared with the 1998 national average, the children of Utah got off to a healthier start to life on seven of eight indicators, including the share of births to unmarried women (see figure). However, the share of births to unmarried women rose from 1990-1998, as did the percentages of births to mothers with less than 12 years of education, births to women receiving late or no prenatal care, and both low-birthweight and preterm births.

More specifically:

- **Teen births.** The share of Utah births to teenagers remained notably lower than the U.S. average during the 1990-1998 period. In 1998, teenagers accounted for 10 percent of births in Utah, compared with a U.S. average of 13 percent.
- **Repeat teen births.** The percentage of teen births in Utah to young women who were already mothers fell from 21 percent in 1990 to a low of 16 percent in 1994, and then rose to 18 percent by 1998. The national average also showed an overall decrease, and Utah was below the national average for this indicator throughout the nine-year period.
- **Births to unmarried women.** There was an increase in the share of Utah births to unmarried women, from 14 percent in 1990 to 17 percent in 1998. However, births to unmarried women remained substantially less common in Utah than in the U.S. as a whole, where unmarried women accounted for 33 percent of births in 1998.
- **Low maternal education.** The proportion of Utah births to women with less than 12 years of education rose slightly from 14 percent in 1990 to 15 percent in 1998. However, the 1998 percentage is still much lower than the national figure of 22 percent.
- **Late or no prenatal care.** The share of Utah births to mothers who received late or no prenatal care decreased between 1990 and 1993 and then steadily increased thereafter—undergoing an overall rise from 3 percent in 1990 to 4 percent in 1998. This change brought Utah from well below the national average in 1990 to even with it by 1998.


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Utah saw a reduction in the already small percentage of births to women who smoked during pregnancy, from 11 percent in 1990 and 1991 to 9 percent in 1994-1998. Although the U.S. as a whole saw a stronger downward trend in this measure, Utah remained below the U.S. average throughout the nine-year period.

- **Low-birthweight births.** The share of Utah births with a low birthweight rose from less than 6 percent in 1990 to 7 percent in 1996-1998. However, low-birthweight births remained less prevalent in Utah than in the U.S. as a whole in 1998.

- **Preterm births.** There was a small overall rise in the share of Utah births that were preterm, from 9 percent in 1990 to 10 percent in 1998. The U.S. as a whole saw a similar increase over the nine-year span, and Utah was below the U.S. average for this indicator in 1998.

In summary, Utah experienced small changes in most measures of a healthy start to life during the 1990-1998 period. When compared with the 1998 national average, Utah's children got off to a healthier start to life on seven of eight indicators. These included the share of births to women who smoked during pregnancy and the percentage of teen births that were repeat births. Although the percentage of Utah births to unmarried women increased from 1990 to 1998, births to unmarried women made up a consistently smaller proportion of births in Utah than in the U.S. overall throughout this time.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	Utah 10.3	10.8	10.5	10.8	10.7	10.8	10.7	10.6	9.8	
	U.S. Average 12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	Utah 20.8	20.4	19.6	18.3	16.1	16.2	17.1	18.7	18.3	
	U.S. Average 24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	Utah 13.5	14.4	15.1	15.5	15.7	15.7	16.2	16.6	17.1	
	U.S. Average 28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	Utah 13.5	13.9	13.8	13.6	13.3	13.5	14.5	14.4	14.6	
	U.S. Average 23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	Utah 3.0	3.0	2.5	2.4	2.8	3.0	3.3	3.5	4.1	
	U.S. Average 6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	Utah 10.8	10.7	10.1	9.6	9.3	9.2	9.3	8.7	8.7	
	U.S. Average 18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	Utah 5.7	6.0	5.6	5.9	5.9	6.3	6.6	6.6	6.7	
	U.S. Average 7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	Utah 8.7	8.6	8.5	8.9	8.9	9.2	9.5	9.9	10.2	
	U.S. Average 10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	32,450	32,050	33,062	32,834	33,714	34,496	36,006	36,534	37,982
	Black non-Hispanic	214	170	208	202	201	252	242	263	263
	Hispanic	2,043	2,175	2,282	2,456	2,704	3,110	3,944	4,298	4,879
	Other	1,570	1,638	1,648	1,635	1,660	1,740	1,885	1,985	2,041
	Total	36,277	36,033	37,200	37,127	38,279	39,577	42,087	43,059	45,165

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Vermont

The yearly number of births in Vermont declined during the 1990s, from 8,273 in 1990 to 6,582 in 1998. Throughout this time, the majority of births were to whites. In 1998, births to whites accounted for 96 percent of total births.*

During the 1990-1998 period, Vermont experienced a substantial increase in the percentage of births to unmarried women and a drop in the share of births to women who smoked during pregnancy. The percentage of births to women with less than 12 years of education was much lower than the U.S. average throughout the 1990s (see figure).

More specifically:


- **Teen births.** Throughout the 1990-1998 period, the percentage of births to teens was markedly lower in Vermont than in the U.S. as a whole. In 1998, teenagers accounted for 8 percent of births in Vermont, compared with 13 percent in the U.S. overall.
- **Repeat teen births.** The share of teen births to young women who were already mothers fluctuated during the 1990s, from a high of 21 percent in 1992 to a low of 14 percent in 1997, but was below the U.S. average throughout this time. In 1998, 17 percent of teen births in Vermont were repeat births, compared with 22 percent in the nation as a whole.
- **Births to unmarried women.** The proportion of births to unmarried women increased dramatically, from 20 percent in 1990 to 28 percent in 1998. Births to unmarried women were less common in Vermont than in the U.S. overall throughout the 1990s.
- **Low maternal education.** Throughout the 1990-1998 period, the percentage of births in Vermont to women with less than 12 years of education was about half that in the U.S. as a whole. In 1998, 11 percent of births in Vermont were to women with less than 12 years of education, compared with 22 percent in the U.S. overall.
- **Late or no prenatal care.** The percentage of Vermont births to women who received late or no prenatal care fell from 3 percent in 1990 to 2 percent in 1993 through 1998. Throughout this time, Vermont was well below the national average for this indicator.
- **Smoking during pregnancy.** Vermont experienced an overall drop in the percentage of births to mothers who smoked during pregnancy, from 23 percent in 1990 to 18 percent in 1998. However, births to mothers

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

who smoked during pregnancy remained more common in Vermont than in the nation overall throughout the 1990s.

- **Low-birthweight births.** The share of Vermont births that were low-birthweight rose from 5 percent in 1990 to 7 percent in 1998. Despite this increase, babies born at a low birthweight continued to account for a smaller share of births in Vermont than in the U.S. as a whole throughout the period.
- **Preterm births.** The proportion of Vermont births that were preterm fluctuated somewhat but underwent an overall increase, from 7 percent in 1990 to 9 percent in 1998. However, Vermont remained below the U.S. average throughout the nine-year period.

The children of Vermont got off to a healthier start to life on seven of eight indicators in 1998, when compared with the national average. Vermont also experienced a notable decrease during the 1990s in the percentage of births to mothers who smoked during pregnancy, although the state remained above the national average throughout the nine-year span. A considerable rise in the share of births to unmarried women during this time is also noteworthy.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	8.5	9.3	8.5	8.7	8.5	8.2	8.9	8.4	7.9	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	17.8	16.6	21.2	18.1	16.9	19.3	14.1	13.5	16.6	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	20.1	22.7	23.4	24.2	25.3	24.9	26.4	26.1	28.0	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	12.0	13.1	12.4	11.5	12.0	10.9	11.6	10.7	10.6	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	3.3	2.7	2.7	2.4	2.1	1.9	2.0	2.0	2.0	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	22.6	21.1	18.9	18.9	17.9	17.0	19.0	17.6	17.9	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	5.3	5.7	5.6	5.7	6.0	5.4	6.2	6.3	6.5	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	7.1	8.1	8.0	7.8	8.3	8.2	8.5	7.6	9.4	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	7,204	6,918	6,672	6,940	6,758	6,268	6,305	6,297	
	Black non-Hispanic	17	23	32	28	25	33	13	22	20
	Hispanic	30	31	30	25	38	27	39	22	37
	Other	1,022	993	1,003	464	556	455	358	258	228
	Total	8,273	7,965	7,737	7,457	7,377	6,783	6,767	6,607	6,582

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Virginia declined from 99,352 in 1990 to 94,351 in 1998. In 1998, births to whites accounted for nearly two-thirds of all Virginia births, while births to blacks accounted for 23 percent. Hispanics accounted for 6 percent of Virginia births in 1998.*

When compared with the 1998 national average, the children of Virginia got off to a healthier start to life on six of eight measures, including the share of births to women with less than 12 years of education (see figure). The state also experienced a number of improvements during the 1990s, the most striking of which was a substantial reduction in the percentage of births to mothers who smoked during pregnancy.


More specifically:

- **Teen births.** Teen births accounted for a smaller proportion of births in Virginia than in the U.S. as a whole throughout the 1990-1998 period. In 1998, 11 percent of births in Virginia were to teen mothers, compared with the national average of 13 percent.
- **Repeat teen births.** During the 1990s, the share of teen births that were to young women who already had a child fluctuated somewhat between 18 percent and 24 percent of Virginia teen births. Both Virginia and the nation as a whole experienced an overall decline on this measure from 1990 to 1998, with Virginia below the national average throughout this time.
- **Births to unmarried women.** There was a sizable increase in the percentage of Virginia births to unmarried women, from 26 percent in 1990 to 30 percent in 1998. Nevertheless, Virginia was below the national average for this measure throughout the 1990-1998 period.
- **Low maternal education.** Births to women with less than 12 years of education decreased from 18 percent of Virginia births in 1990 to 15 percent in 1998. Virginia was substantially below the U.S. average for this measure throughout this period.
- **Late or no prenatal care.** Already well below the U.S. average in 1990, the share of Virginia births to women receiving late or no prenatal care continued to decrease over the 1990-1998 period. In 1998, births to mothers receiving late or no prenatal care accounted for 3 percent of births in Virginia, compared with 4 percent in the U.S. overall.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** The percentage of births to mothers who smoked during pregnancy decreased from 16 percent in 1990 to 10 percent in 1998. Moreover, smoking during pregnancy remained less common in Virginia than in the U.S. overall throughout this period.
- **Low-birthweight births.** Both Virginia and the nation as a whole witnessed a slight increase in the percentage of babies born at a low birthweight during the 1990s, from 7 percent in 1990 to 8 percent in 1998.
- **Preterm births.** Preterm births accounted for between 11 percent and 12 percent of Virginia births during the 1990-1998 period. Throughout this time, Virginia was slightly above the U.S. average, which remained steady at 11 percent.

In summary, Virginia saw improvements on five of eight measures of a healthy start to life over the 1990-1998 period, including considerable declines in the percentages of births to women who smoked during pregnancy and to women who received late or no prenatal care. Furthermore, the proportion of births to women with less than 12 years of education remained remarkably lower than the national average throughout the 1990s. However, the state also saw a marked increase in the percentage of births to unmarried women during this nine-year span.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	11.7	11.5	11.0	10.9	11.3	11.4	11.0	11.0	10.8	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	23.0	23.5	23.9	21.2	19.5	18.4	19.1	20.2	20.5	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	26.0	27.9	28.3	29.0	29.2	29.3	28.8	29.3	29.8	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	18.1	18.4	17.4	16.8	16.3	16.0	15.6	15.7	15.2	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	4.6	4.3	3.7	3.4	3.5	3.2	3.3	3.2	3.3	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	15.9	15.3	14.9	14.6	13.9	13.0	11.8	11.3	10.4	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	7.2	7.2	7.4	7.3	7.5	7.7	7.7	7.7	7.9	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	11.0	11.1	11.4	11.6	11.5	11.9	11.9	12.1	12.3	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	68,944	66,858	66,370	64,359	64,824	62,660	62,192	61,428	62,119
	Black non-Hispanic	23,770	23,733	23,737	23,170	22,176	21,209	20,808	20,956	21,913
	Hispanic	3,459	3,749	3,890	4,135	4,609	4,841	5,150	5,374	5,806
	Other	3,179	3,030	3,201	3,280	3,430	3,868	4,204	4,104	4,513
	Total	99,352	97,370	97,198	94,944	95,039	92,578	92,354	91,862	94,351

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Washington was similar in 1990 and 1998. In 1990 there were 79,251 births, while in 1998 there were 79,663. During this time, however, births to whites decreased, from 63,707 in 1990 to 57,214 in 1998, while births to Hispanics rose, from 5,700 in 1990 to 10,074 in 1998. Asian/Pacific Islanders (included in Other) accounted for 7 percent of Washington births in 1998.

When compared with the national average in 1998, Washington's children got off to a healthier start to life on seven of eight indicators, including the percentage of low-birthweight births (see figure). Moreover, the state saw decreases during the 1990s in the shares of births to women who smoked during pregnancy and to women who received late or no prenatal care. At the same time, however, there was an increase in the share of births to unmarried women.

More specifically:


- **Teen births.** Throughout the 1990s, the proportion of Washington births to teens remained somewhat below the national average. In 1998, 11 percent of births in Washington were to teenagers, compared with 13 percent in the U.S. as a whole.
- **Repeat teen births.** The percentage of Washington teen births to young women who were already mothers fluctuated during the 1990s, undergoing an overall decline from a high of 22 percent in 1992 to 18 percent in 1998. The share of Washington teen births that were repeat births remained below the U.S. average throughout the 1990-1998 period.
- **Births to unmarried women.** Washington saw an increase in the share of births to unmarried women, from 24 percent in 1990 to 28 percent in 1998. Births to unmarried women remained less common in Washington than in the nation as a whole throughout the nine-year span.
- **Low maternal education.** Between 1992, when data first became available, and 1998, births to women with less than 12 years of education made up a smaller share of births in Washington than in the U.S. as a whole. In 1998, 18 percent of births in Washington were to women with less than 12 years of education, compared with 22 percent in the U.S. overall.
- **Late or no prenatal care.** The percentage of births in Washington to women who received late or no prenatal care fell from 5 percent in 1990

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

to 3 percent in 1998. Washington remained below the U.S. average for this indicator throughout the 1990-1998 period.

- **Smoking during pregnancy.** The share of Washington births to women who smoked during pregnancy dropped from 20 percent in 1990 through 1992 to less than 15 percent in 1997 and 1998. Smoking during pregnancy remained somewhat more common in Washington than in the nation overall throughout the 1990s.
- **Low-birthweight births.** Low-birthweight births were consistently less prevalent in Washington than in the U.S. as a whole during the 1990-1998 period. In 1998, 6 percent of births in Washington were low-birthweight, compared with 8 percent in the U.S. overall.
- **Preterm births.** There was a small increase in the share of Washington births that were preterm, from 8 percent in 1990 through 1994 to 9 percent in 1998. However, the percentage of preterm births in Washington remained below the national average throughout the 1990-1998 period.

In summary, over the 1990-1998 period, the changes in measures of a healthy start to life in Washington generally mirrored the trends in the nation as a whole. The state saw drops in the percentages of births to women who smoked during pregnancy, births to women who received late or no prenatal care, and teen births that were repeat births. On the other hand, the share of births to unmarried women and the percentage of births that were preterm increased. When compared with the U.S. average, Washington's children got off to a healthier start to life on seven of eight indicators in 1998.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	10.8	10.8	10.6	11.0	11.1	11.5	11.2	11.0	10.9	
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	20.7	20.7	21.5	19.3	18.3	17.8	18.7	18.5	17.7	
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	23.7	24.9	25.3	26.3	26.0	26.7	27.3	27.1	27.9	
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	N.A.	N.A.	17.7	18.6	18.5	18.7	18.3	18.1	18.1	
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	4.8	4.3	4.0	3.7	3.4	3.5	3.6	3.4	3.2	
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	20.2	19.5	19.9	18.3	17.0	16.2	16.0	14.6	14.6	
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	5.3	5.1	5.3	5.2	5.3	5.5	5.6	5.6	5.7	
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	8.4	8.4	8.3	8.4	8.4	8.5	8.6	8.9	9.5	
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	63,707	62,567	61,224	59,636	57,511	56,839	56,950	56,617	57,214
	Black non-Hispanic	2,759	2,887	2,993	3,036	2,880	2,757	2,886	2,968	2,888
	Hispanic	5,700	6,492	7,182	7,708	8,108	8,502	8,930	9,336	10,074
	Other	7,085	7,765	8,051	8,265	8,859	9,130	9,179	9,269	9,487
	Total	79,251	79,711	79,450	78,645	77,358	77,945	78,190	79,663	

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**N.A. = Not Available. See Appendix 2 for more details.

***The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

West Virginia

The yearly number of births in West Virginia decreased from 22,585 in 1990 to 20,747 in 1998. Whites accounted for 95 percent of all births in 1998, while blacks accounted for 4 percent. These percentages remained constant throughout the 1990-1998 period.*

During this nine-year span, West Virginia saw a substantial drop in the already small share of births to women who received late or no prenatal care. At the same time, however, there were marked increases in the percentages of preterm births and births to unmarried women. Moreover, the proportion of births to women who smoked during pregnancy remained strikingly higher than the national average throughout this period (see figure).

More specifically:

- **Teen births.** Teen births became a somewhat smaller share of West Virginia births during the 1990s, falling overall from 18 percent in 1990 to 16 percent in 1998. Despite this change, the percentage of births to teens in West Virginia remained notably above the national average throughout the 1990s.
- **Repeat teen births.** The proportion of West Virginia teen births to young women who already had a child remained somewhat below the U.S. average throughout the 1990-1998 period. In 1998, 19 percent of teen births in West Virginia were repeat births, compared with 22 percent in the U.S. as a whole.
- **Births to unmarried women.** There was a substantial increase in the share of West Virginia births to unmarried women, from 25 percent in 1990 to 32 percent in 1998. The nation as a whole experienced a somewhat weaker upward trend, and the percentage of West Virginia births to unmarried women went from below the national average in 1990 to even with it in 1998.
- **Low maternal education.** West Virginia saw a sizable decline in the share of births to women with less than 12 years of education, from more than 25 percent in 1990 and 1991 to less than 21 percent in 1998. This change brought West Virginia from just above the U.S. average in 1990 to just below it by 1998.
- **Late or no prenatal care.** The percentage of West Virginia births to women who received late or no prenatal care fell by more than half, from 6 percent in 1990 to less than 3 percent by 1998. Moreover, West


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

Virginia remained below the U.S. average for this measure throughout the nine-year span.

- **Smoking during pregnancy.** Births to women who smoked during pregnancy were twice as common in West Virginia than in the U.S. as a whole in 1998. In 1998, 26 percent of West Virginia births were to mothers who smoked during pregnancy, compared with a national average of 13 percent.
- **Low-birthweight births.** The percentage of West Virginia babies born at a low birthweight remained close to the U.S. average throughout the 1990-1998 period. At 8 percent in 1998, West Virginia was just above the national average.
- **Preterm births.** Preterm births became a larger share of West Virginia births during the 1990s, rising from 10 percent in 1990 to 12 percent in 1997 and 1998. This change brought West Virginia from below the U.S. average in 1990 to above it by 1998.

In summary, West Virginia experienced a dramatic reduction in the percentage of births to women who received late or no prenatal care and remained below the U.S. average for this indicator throughout the 1990-1998 period. The state also saw notable drops during this time in the shares of births to teens and to women with less than 12 years of education. However, there were increases in the proportion of births to unmarried women and in the percentages of preterm and low-birthweight births. Throughout the nine-year period, smoking during pregnancy was substantially more common in West Virginia than in the nation as a whole.

Smoking During Pregnancy



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	17.8	17.6	17.2	17.6	17.4	17.2	16.7	15.9	15.7	
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	20.4	19.4	20.8	18.2	18.3	17.7	18.2	18.9	18.6	
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	25.4	26.8	27.7	29.0	30.2	30.5	31.3	31.3	32.4	
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	25.5	25.6	24.8	23.9	22.6	22.0	21.5	21.0	20.6	
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	5.7	4.9	4.2	3.7	3.4	3.0	2.7	3.1	2.6	
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	27.8	26.8	27.0	26.5	26.1	24.6	25.5	25.3	25.6	
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	7.1	6.8	7.2	7.2	7.5	7.9	8.0	8.3	8.0	
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	9.9	10.1	10.3	10.7	10.7	11.2	11.4	12.4	12.4	
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	21,545	21,525	21,185	20,771	20,377	20,162	19,757	19,778	19,728
	Black non-Hispanic	859	801	812	815	794	806	738	744	757
	Hispanic	73	73	75	81	106	90	125	82	93
	Other	108	109	98	125	98	104	130	126	169
	Total	22,585	22,508	22,170	21,792	21,375	21,162	20,750	20,730	20,747

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

Wisconsin

The yearly number of births in Wisconsin decreased from 72,895 in 1990 to 67,450 in 1998. The number of births to Hispanics nearly doubled during this time, accounting for 5 percent of Wisconsin births in 1998, compared with 3 percent in 1990. Concurrently, the number of births to whites and blacks decreased.*

When compared with the national average in 1998, Wisconsin's children got off to a significantly healthier start to life on six of eight indicators, including the share of births to women with less than 12 years of education (see figure). The proportion of births to women who smoked during pregnancy, however, was well above the national average throughout the 1990-1998 period. During this nine-year span, the changes that Wisconsin saw in measures of a healthy start to life tended to mirror changes in the U.S. average.


More specifically:

- **Teen births.** Throughout the 1990-1998 period, teen births made up a smaller share of births in Wisconsin than in the nation as a whole. In 1998, 10 percent of Wisconsin births were to teens, compared with a national average of 13 percent.
- **Repeat teen births.** The percentage of Wisconsin teen births to young women who were already mothers fluctuated, undergoing an overall drop from 25 percent in 1990 to 21 percent in 1998. Wisconsin was even with the U.S. average for this indicator in 1998.
- **Births to unmarried women.** Wisconsin experienced a rise in the share of births to unmarried women, from 24 percent in 1990 to 28 percent in 1998. However, births to unmarried women remained less common in Wisconsin than in the nation as a whole throughout the nine-year period.
- **Low maternal education.** Births to women with less than 12 years of education made up a smaller share of births in Wisconsin than in the U.S. as a whole throughout the 1990-1998 period. In 1998, these births accounted for 16 percent of births in Wisconsin, compared with 22 percent in the U.S. overall.
- **Late or no prenatal care.** There was a small decrease in the percentage of births to women who received late or no prenatal care in Wisconsin, from 4 percent in 1990 to 3 percent by 1998. In 1998, this percentage was just below the national average of 4 percent.

* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

- **Smoking during pregnancy.** Wisconsin experienced a drop in the share of births to women who smoked during pregnancy, from 23 percent in 1990 to 18 percent in 1998. However, smoking during pregnancy remained markedly more common in Wisconsin than in the nation as a whole throughout the nine-year period.
- **Low-birthweight births.** The percentage of Wisconsin babies born at a low birthweight remained below the national average throughout the 1990-1998 period. In 1998, 7 percent of births in Wisconsin were low-birthweight, compared with 8 percent in the nation overall.
- **Preterm births.** The share of Wisconsin births that were preterm increased from 9 percent in 1990 to 11 percent in 1998. However, preterm births remained somewhat less prevalent in Wisconsin than in the U.S. as a whole in 1998.

In summary, Wisconsin saw marked drops during the 1990-1998 period in the share of births to women who smoked during pregnancy and in the percentage of teen births that were repeat births. Moreover, the proportion of births to women with less than 12 years of education was dramatically lower than the U.S. average throughout this nine-year period. On the other hand, there were sizable increases in the percentages of preterm births and births to unmarried women, and the share of births to women who smoked during pregnancy remained well above the national average throughout the 1990s.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998	
Percent of total births to teens	10.2	10.3	10.2	10.4	10.3	10.5	10.6	10.7	10.5	
<i>U.S. Average</i>	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5	
Percent of teen births to women who were already mothers	25.2	25.5	25.7	24.6	21.8	20.2	20.7	19.6	21.1	
<i>U.S. Average</i>	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5	
Percent of total births to unmarried women	24.2	25.3	26.1	27.1	27.2	27.4	27.4	28.1	28.5	
<i>U.S. Average</i>	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8	
Percent of total births to mothers with less than 12 years of education	16.9	17.1	17.0	16.9	16.6	16.3	16.1	16.1	15.8	
<i>U.S. Average</i>	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9	
Percent of total births to mothers receiving late or no prenatal care	3.8	3.8	3.6	3.6	3.5	3.4	3.3	3.2	3.4	
<i>U.S. Average</i>	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9	
Percent of total births to mothers who smoked during pregnancy	22.9	22.6	21.7	20.8	19.6	18.9	18.1	17.8	17.8	
<i>U.S. Average</i>	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9	
Percent low-birthweight births (less than 5.5 lbs)	5.9	6.1	5.9	6.1	6.4	6.0	6.3	6.4	6.5	
<i>U.S. Average</i>	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6	
Percent preterm births (less than 37 completed weeks of gestation)	9.2	9.6	9.2	9.6	10.2	9.7	9.7	10.3	10.5	
<i>U.S. Average</i>	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6	
Number of births by Race and Ethnicity of mother	White non-Hispanic	61,496	60,374	58,558	57,720	56,419	55,365	54,863	54,292	54,636
	Black non-Hispanic	7,108	7,154	7,265	7,128	6,814	6,476	6,386	6,366	6,490
	Hispanic	1,866	1,950	2,132	2,175	2,401	2,856	3,139	3,207	3,641
	Other	2,425	2,593	2,715	2,744	2,648	2,782	2,718	2,692	2,683
	Total	72,895	72,071	70,670	69,767	68,282	67,479	67,106	66,557	67,450

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

The yearly number of births in Wyoming decreased from 6,985 in 1990 to 6,252 in 1998. In 1998, births to whites accounted for 85 percent of Wyoming births.*

Wyoming saw a remarkable increase over the 1990-1998 period in the share of births to unmarried women (see figure). Moreover, throughout this time, smoking during pregnancy was dramatically more common in Wyoming than in the U.S. as a whole. On the other hand, the share of births to women with less than 12 years of education was below the U.S. average in 1998.

More specifically:

- **Teen births.** The percentage of Wyoming births to teens fluctuated during the 1990s but increased overall, from 14 percent in 1990 to 16 percent in 1998. The relatively stable U.S. average was considerably lower at 13 percent.
- **Repeat teen births.** The share of Wyoming teen births to young women who were already mothers fluctuated during the 1990-1998 period but remained below the national average throughout the nine-year span. In 1998, 18 percent of teen births in Wyoming were repeat births, compared with 22 percent in the U.S. as a whole.
- **Births to unmarried women.** The proportion of Wyoming births to unmarried women rose from 20 percent in 1990 to 30 percent in 1998. While births to unmarried women remained somewhat less common in Wyoming than in the U.S. as a whole in 1998, this change narrowed the gap between Wyoming and the U.S. average.
- **Low maternal education.** Throughout the 1990-1998 period, births to women with less than 12 years of education made up a smaller share of births in Wyoming than in the U.S. as a whole. In 1998, these births accounted for 17 percent of births in Wyoming, compared with 22 percent in the U.S. overall.
- **Late or no prenatal care.** The percentage of Wyoming births to women who received late or no prenatal care dropped slightly from 4 percent during 1990 through 1992 to less than 3 percent in 1993. However, by 1998 the percentage had returned to 4 percent, matching the U.S. average in 1998.
- **Smoking during pregnancy.** The proportion of births in Wyoming to women who smoked during pregnancy fell from 24 percent in 1990 to a


* Data for whites and blacks exclude Hispanics. Persons of Hispanic origin may be of any race.

low of 20 percent in 1995, and then rose to 23 percent in 1998.

Throughout this time, smoking during pregnancy was substantially more common in Wyoming than in the U.S. as a whole, where 13 percent of births were to mothers who smoked during pregnancy in 1998.

- **Low-birthweight births.** The share of Wyoming babies born at a low birthweight fluctuated but increased overall from 7 percent in 1990 to 9 percent by 1998. In 1998, low-birthweight births were more prevalent in Wyoming than in the U.S. as a whole.
- **Preterm births.** There was an overall increase during the 1990-1998 period in the percentage of Wyoming births that were preterm, despite some fluctuations during this time. The proportion of births that were preterm rose from 10 percent in 1990 to 12 percent in 1998, matching the 1998 national average.

In summary, over the 1990-1998 period, Wyoming saw a dramatic increase in the share of births to unmarried women, and smaller rises in the percentages of preterm, low-birthweight, and teen births. In 1998, the proportions of births to mothers who smoked during pregnancy and to teenage mothers were substantially higher than the respective national averages, while the percentages of teen births that were repeat births and of all births to women with less than 12 years of education were relatively low.



Trends In Key Indicators, 1990-1998

Indicator*	1990	1991	1992	1993	1994	1995	1996	1997	1998
Percent of total births to teens									
Wyoming	13.6	14.0	13.2	14.0	14.5	15.2	14.3	14.2	16.2
U.S. Average	12.8	12.9	12.7	12.8	13.1	13.1	12.9	12.7	12.5
Percent of teen births to women who were already mothers									
Wyoming	18.9	20.9	17.1	16.3	16.7	19.3	14.5	15.9	17.9
U.S. Average	24.3	24.6	25.0	23.4	21.8	21.0	21.4	21.6	21.5
Percent of total births to unmarried women									
Wyoming	19.8	23.1	24.0	25.8	27.5	26.4	27.0	27.4	29.6
U.S. Average	28.0	29.5	30.1	31.0	32.6	32.2	32.4	32.4	32.8
Percent of total births to mothers with less than 12 years of education									
Wyoming	17.7	18.2	16.7	16.9	16.3	16.1	15.6	16.3	17.2
U.S. Average	23.8	23.9	23.6	23.3	22.9	22.6	22.4	22.1	21.9
Percent of total births to mothers receiving late or no prenatal care									
Wyoming	3.8	4.1	4.3	2.9	3.6	3.8	3.7	3.7	4.1
U.S. Average	6.1	5.8	5.2	4.8	4.4	4.2	4.0	3.9	3.9
Percent of total births to mothers who smoked during pregnancy									
Wyoming	24.3	24.3	23.6	21.5	21.8	19.8	20.5	20.5	22.5
U.S. Average	18.4	17.8	16.9	15.8	14.5	13.9	13.6	13.2	12.9
Percent low-birthweight births (less than 5.5 lbs)									
Wyoming	7.4	7.0	7.3	7.3	8.8	7.4	8.4	9.0	8.9
U.S. Average	7.0	7.1	7.1	7.2	7.3	7.3	7.4	7.5	7.6
Percent preterm births (less than 37 completed weeks of gestation)									
Wyoming	10.3	10.1	9.9	10.8	11.4	10.7	10.8	11.4	11.8
U.S. Average	10.6	10.8	10.7	11.0	11.0	11.0	11.0	11.4	11.6
Number of births by Race and Ethnicity of mother									
White non-Hispanic	6,079	5,826	5,852	5,713	5,656	5,368	5,441	5,527	5,319
Black non-Hispanic	64	59	61	60	57	69	48	58	52
Hispanic	511	497	482	488	460	548	523	538	584
Other	331	321	328	294	255	276	274	264	297
Total	6,985	6,703	6,723	6,555	6,428	6,261	6,286	6,387	6,252

*For a complete definition and a list of sources for each of the nine indicators, please see Appendix 2.

**The Other category consists of Asian-Pacific Islanders, American Indians, and whites and blacks where Hispanic origin is unknown.

RESOURCES

240

Appendix 1: States Ranked by Indicator, 1998

Percent of total births to teens

Rank	State	Percent	Rank	State	Percent
	The Nation	12.5	26	Illinois	12.4
1	Massachusetts	7.2	27	Oregon	12.5
2	New Hampshire	7.7	28	Kansas	12.6
2	New Jersey	7.7	29	Idaho	12.8
4	Vermont	7.9	30	Ohio	13.0
5	Connecticut	8.3	31	Nevada	13.1
6	Minnesota	8.6	31	Delaware	13.1
7	New York	8.8	33	Florida	13.2
8	Maine	9.8	34	Indiana	13.8
8	Utah	9.8	34	Missouri	13.8
8	North Dakota	9.8	36	North Carolina	14.0
11	Maryland	10.2	37	Georgia	15.0
12	Pennsylvania	10.3	38	Arizona	15.1
13	Wisconsin	10.5	39	District of Columbia	15.3
13	Rhode Island	10.5	40	Kentucky	15.4
15	Nebraska	10.6	41	West Virginia	15.7
15	Iowa	10.6	42	Tennessee	15.9
17	Hawaii	10.7	43	South Carolina	16.0
18	Virginia	10.8	44	Texas	16.1
19	Washington	10.9	45	Wyoming	16.2
20	Alaska	11.2	46	Oklahoma	16.3
21	California	11.4	47	Alabama	17.1
22	Michigan	11.6	48	New Mexico	18.2
23	South Dakota	12.0	49	Louisiana	18.4
24	Colorado	12.1	50	Arkansas	18.6
25	Montana	12.3	51	Mississippi	20.0

Percent of teen births to women who were already mothers

Rank	State	Percent	Rank	State	Percent
	The Nation	21.5	26	Virginia	20.5
1	North Dakota	12.3	27	Connecticut	20.7
2	New Hampshire	14.7	28	Missouri	20.9
3	Maine	14.9	28	Kentucky	20.9
4	Montana	15.7	30	Wisconsin	21.1
5	Massachusetts	16.5	31	Kansas	21.2
6	Vermont	16.6	31	California	21.2
6	Hawaii	16.6	33	Indiana	21.3
8	South Dakota	17.2	33	Michigan	21.3
9	Washington	17.7	33	Ohio	21.3
10	Iowa	17.9	33	South Carolina	21.3
10	Wyoming	17.9	37	Arizona	22.0
12	New York	18.3	37	Tennessee	22.0
12	Utah	18.3	39	North Carolina	22.3
12	Nebraska	18.3	40	Delaware	22.4
12	Minnesota	18.3	41	Florida	22.5
16	West Virginia	18.6	42	Louisiana	22.7
17	Oregon	19.1	43	Nevada	22.8
17	Idaho	19.1	44	Alabama	23.0
19	Rhode Island	19.2	45	Illinois	23.4
20	New Mexico	19.8	46	Arkansas	23.5
21	Alaska	20.1	47	Georgia	24.1
22	Maryland	20.3	47	Texas	24.1
22	New Jersey	20.3	49	Mississippi	24.4
22	Colorado	20.3	50	District of Columbia	28.0
25	Pennsylvania	20.4		Oklahoma*	N.R.

* N.R. - Not Reliable



Percent of total births to unmarried women

Rank	State	Percent	Rank	State	Percent
	The Nation	32.8	26	West Virginia	32.4
1	Utah	17.1	27	California	32.8
2	Idaho	22.0	27	North Carolina	32.8
3	New Hampshire	24.1	27	Pennsylvania	32.8
4	Colorado	25.6	30	Oklahoma	33.2
4	Minnesota	25.6	31	Indiana	33.5
6	Massachusetts	26.1	32	Rhode Island	33.9
7	Nebraska	26.2	32	Michigan	33.9
8	North Dakota	27.0	34	Ohio	34.0
9	Iowa	27.2	35	Missouri	34.1
10	Kansas	27.8	35	Alabama	34.1
11	Washington	27.9	35	Illinois	34.1
12	Vermont	28.0	38	Maryland	34.4
13	New Jersey	28.3	39	Tennessee	34.9
14	Wisconsin	28.5	39	New York	34.9
15	Wyoming	29.6	41	Nevada	35.0
16	Oregon	29.7	41	Arkansas	35.0
17	Virginia	29.8	43	Georgia	36.2
18	Montana	29.9	44	Florida	36.6
19	Kentucky	30.1	45	Delaware	37.1
20	Maine	30.6	46	Arizona	38.4
21	Alaska	31.1	47	South Carolina	38.8
22	Connecticut	31.2	48	New Mexico	44.0
23	Texas	31.5	49	Louisiana	44.9
23	Hawaii	31.5	50	Mississippi	45.4
25	South Dakota	32.0	51	District of Columbia	62.9

Percent of total births to mothers with less than 12 years of education

Rank	State	Percent	Rank	State	Percent
	The Nation	21.9	24	Kansas	18.1
1	North Dakota	10.0	27	Delaware	18.3
2	New Hampshire	10.3	28	New York	19.2
3	Vermont	10.6	29	Colorado	19.4
3	Minnesota	10.6	30	Missouri	19.6
5	Connecticut	10.9	31	Indiana	20.5
5	Maine	10.9	32	West Virginia	20.6
7	Hawaii	11.2	32	Oregon	20.6
8	Maryland	13.0	34	Florida	21.3
9	Alaska	13.2	35	North Carolina	21.7
10	Massachusetts	13.4	36	South Carolina	22.0
11	Iowa	13.6	36	Illinois	22.0
12	New Jersey	13.9	36	Kentucky	22.0
13	Nebraska	14.5	39	Tennessee	22.1
14	Utah	14.6	40	Arkansas	22.8
15	Pennsylvania	15.1	41	Georgia	23.0
16	Virginia	15.2	41	Oklahoma	23.0
17	Montana	15.3	43	Alabama	23.3
18	Rhode Island	15.5	44	Louisiana	24.5
18	South Dakota	15.5	45	District of Columbia	25.2
20	Wisconsin	15.8	46	Mississippi	26.2
21	Wyoming	17.2	47	New Mexico	28.2
22	Michigan	17.5	48	Nevada	28.3
23	Ohio	17.9	49	Arizona	29.5
24	Washington	18.1	50	California	31.0
24	Idaho	18.1	51	Texas	33.1

Percent of total births to mothers receiving late or no prenatal care

Rank	State	Percent	Rank	State	Percent
	The Nation	3.9	25	Pennsylvania	3.5
1	Rhode Island	1.5	27	Delaware	3.6
2	Maine	1.7	27	Tennessee	3.6
3	New Hampshire	1.9	27	California	3.6
4	Vermont	2.0	30	Oregon	3.8
5	Iowa	2.4	31	Alabama	3.9
5	Massachusetts	2.4	31	Illinois	3.9
7	Kentucky	2.5	31	Louisiana	3.9
7	North Dakota	2.5	34	Mississippi	4.0
9	West Virginia	2.6	34	Indiana	4.0
10	Georgia	2.8	36	Utah	4.1
10	Kansas	2.8	36	Wyoming	4.1
12	Missouri	2.9	38	South Carolina	4.2
12	Minnesota	2.9	38	Ohio	4.2
12	North Carolina	2.9	40	Colorado	4.3
15	Maryland	3.0	41	Idaho	4.4
15	Connecticut	3.0	42	Alaska	4.5
17	Hawaii	3.1	43	New Jersey	4.6
18	Washington	3.2	44	New York	4.8
18	Nebraska	3.2	45	Oklahoma	5.1
18	Montana	3.2	45	Arkansas	5.1
18	South Dakota	3.2	47	Texas	5.3
22	Virginia	3.3	48	Nevada	7.0
23	Michigan	3.4	49	Arizona	7.2
23	Wisconsin	3.4	50	New Mexico	8.5
25	Florida	3.5	51	District of Columbia	10.2

Percent of total births to mothers who smoked during pregnancy

Rank	State	Percent	Rank	State	Percent
	The Nation*	12.9	25	Delaware	14.9
1	District of Columbia	4.8	27	Oregon	15.2
2	Texas	7.2	27	Rhode Island	15.2
3	Arizona	7.6	29	Nebraska	16.3
4	Hawaii	8.5	30	New Hampshire	16.6
5	Utah	8.7	31	Tennessee	16.8
6	Connecticut	9.4	32	Michigan	17.0
7	Georgia	9.7	33	Pennsylvania	17.5
8	Maryland	9.8	34	Wisconsin	17.8
9	Louisiana	10.3	35	Vermont	17.9
10	Virginia	10.4	36	Alaska	18.4
11	Colorado	11.0	37	Montana	18.5
11	New Mexico	11.0	38	Oklahoma	18.6
13	Florida	11.2	39	Iowa	18.9
13	New Jersey	11.2	40	Arkansas	19.0
15	Massachusetts	11.7	41	Missouri	19.1
16	Illinois	11.8	42	Ohio	19.3
17	Minnesota	12.4	43	North Dakota	19.5
18	Alabama	12.6	44	Maine	19.6
19	Nevada	12.8	45	Wyoming	22.5
20	Mississippi	13.0	46	Kentucky	24.8
21	Idaho	13.1	47	West Virginia	25.6
21	Kansas	13.1		California**	N.A.
23	South Carolina	13.7		Indiana**	N.A.
24	Washington	14.6		New York**	N.A.
25	North Carolina	14.9		South Dakota**	N.A.

* California, Indiana, New York, and South Dakota are excluded from the national average.

** N.A. - Not Available

245

246

Percent low-birthweight births (less than 5.5 pounds)

Rank	State	Percent	Rank	State	Percent
	The Nation	7.6	23	Pennsylvania	7.6
1	Oregon	5.4	27	Ohio	7.7
2	New Hampshire	5.7	28	Connecticut	7.8
2	Washington	5.7	28	Michigan	7.8
4	Minnesota	5.8	28	Missouri	7.8
4	South Dakota	5.8	28	New York	7.8
4	Maine	5.8	32	Indiana	7.9
7	Alaska	6.0	32	Virginia	7.9
7	Idaho	6.0	34	Illinois	8.0
9	California	6.2	34	New Jersey	8.0
10	Iowa	6.4	34	West Virginia	8.0
11	Vermont	6.5	37	Florida	8.1
11	Nebraska	6.5	37	Kentucky	8.1
11	Wisconsin	6.5	39	Delaware	8.4
11	North Dakota	6.5	40	Georgia	8.5
15	Utah	6.7	41	Colorado	8.6
16	Arizona	6.8	42	Maryland	8.7
17	Massachusetts	6.9	43	North Carolina	8.8
18	Montana	7.0	44	Arkansas	8.9
18	Kansas	7.0	44	Wyoming	8.9
20	Oklahoma	7.2	46	Tennessee	9.1
21	Texas	7.4	47	Alabama	9.3
22	Hawaii	7.5	48	South Carolina	9.5
23	Rhode Island	7.6	49	Mississippi	10.1
23	Nevada	7.6	49	Louisiana	10.1
23	New Mexico	7.6	51	District of Columbia	13.1

Percent preterm births (less than 37 completed weeks of gestation)

Rank	State	Percent	Rank	State	Percent
	The Nation	11.6	26	Arizona	11.4
1	New Hampshire	8.1	27	Oklahoma	11.5
2	Oregon	9.1	27	Michigan	11.5
2	Maine	9.1	27	Indiana	11.5
4	Minnesota	9.2	30	Georgia	11.6
5	Vermont	9.4	31	Ohio	11.7
6	Washington	9.5	32	Wyoming	11.8
7	Idaho	9.6	33	Missouri	12.0
8	Massachusetts	9.8	34	Illinois	12.1
9	Alaska	10.0	35	Virginia	12.3
10	Utah	10.2	36	Texas	12.4
11	Connecticut	10.3	36	West Virginia	12.4
11	California	10.3	38	Kentucky	12.5
13	Wisconsin	10.5	38	New Jersey	12.5
14	North Dakota	10.6	40	Florida	12.6
15	Rhode Island	10.7	40	Delaware	12.6
15	Nebraska	10.7	42	North Carolina	12.7
15	Iowa	10.7	43	Maryland	12.8
18	Kansas	10.8	44	Nevada	12.9
18	South Dakota	10.8	45	Arkansas	13.3
20	Pennsylvania	10.9	46	South Carolina	13.6
20	New York	10.9	46	Tennessee	13.6
22	Montana	11.0	48	Alabama	14.3
23	Hawaii	11.1	49	Louisiana	14.7
24	Colorado	11.2	50	Mississippi	16.0
25	New Mexico	11.3	51	District of Columbia	18.5

247

248

Appendix 2: Definitions, Data Sources, and Reporting Issues

Introduction

This Appendix provides definitions for the measures in this report and for technical terms in the narrative. It also addresses situations where there have been changes or problems in the way data have been collected, coded or reported by state departments of health. Such changes can result in substantial year-to-year changes in statistics that are due, at least in part, to the change in the methodology used to collect and report the data rather than to actual changes in behavior. All data are reported by the mother's place of residence, not the place where the infant was born.

All percentages in the *text* are rounded to the nearest whole percent; all percentages in the state *tables* are rounded to the nearest tenth of a percent. Thus, 9.46 rounds to 9 percent in the text and 9.5 percent in the tables

Definitions and data sources

Births to unmarried women (percent of total births to unmarried women) is the percentage of all births occurring to women who were unmarried at the time of the birth. In 1998, marital status was obtained from a direct question on the birth certificate in 48 states and the District of Columbia. In Michigan and New York, marital status was inferred from other information on the birth certificate. See the next section for more details.

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics. Data for Nevada for 1995 and 1996 only are from the Technical Notes in Ventura, S.J., Martin, J.A., Curtin S.C., Mathews T.J. (1998), "Births: Final data for 1997," *National Vital Statistics Reports*; Vol 47 no 18. Hyattsville, Maryland: National Center for Health Statistics.

Gap means the difference (measured in percentage points) between a percentage reported for a state and the comparable percentage for the U.S. as a whole. For example, in 1998, 14 percent of births in Alabama were preterm, compared with 12 percent for the U.S. as a whole. In this example, the gap is 2 percentage points.

Late or no prenatal care (percent of total births to mothers receiving late or no prenatal care) is the percentage of births that occurred to mothers who reported receiving prenatal care only in the third trimester of their pregnancy, or reported receiving no prenatal care. Birth certificates that did not report information about prenatal care were not included in this

calculation. In three states, there were reporting issues in some years. See the next section for more details.

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics.

Low-birthweight births (percent low-birthweight births) is the percentage of live births weighing less than 2,500 grams (5.5 pounds). Births of unknown weight were not included in these calculations.

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics.

Low maternal education (percent of total births to mothers with less than 12 years of education) is the percentage of women who had completed fewer than 12 years of education at the time of the birth. Birth certificates on which maternal education was not reported were not included in this calculation. Data on maternal education were not available for two states in the early 1990s. See the next section for more details.

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics.

Number of births by race and ethnicity of mother contains information about race/Hispanic origin of the mother. On birth certificates, as on most federal data collection forms, the question regarding whether a person is Hispanic is separate from the question asking whether a person is white, black, Asian or Pacific Islander, or Native American. Thus, people are asked to select a racial group and to indicate whether they are of Hispanic origin. A birth to a woman who reported that she was Hispanic and white would usually be included in figures for both of these groups. In order to create mutually exclusive categories, Hispanics were removed from the black and white racial categories in the tabulations presented here. This allows more meaningful comparisons between minorities and the group people typically think of when we say "white."

It should also be noted that these figures represent the race of the mother, not the race of the child. This is important because increasing numbers of children are born to parents of different races.

Births for which Hispanic origin was not reported are included in the "Other" category.

In 1990, Oklahoma did not collect data on Hispanic origin. New Hampshire did not report Hispanic origin until 1993 and did not collect this information reliably until 1994. See the next section for more details.

Appendix 2: Definitions, Data Sources, and Reporting Issues

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics.

Preterm births (percent preterm births) is the percentage of babies born with a gestational age of less than 37 completed weeks. Birth certificates that did not report gestational age were not included in this tabulation.

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics.

Repeat teen births (percent of teen births to women who were already mothers) is the percentage of births that were second or higher order births to mothers who were under the age of 20 at the time of the birth. Birth certificates that did not contain information on birth order were not included in this calculation.

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics.

Significant means *statistically* significant. The number of births reported for an area is essentially a complete count and is not subject to sampling error. (This is also true for births classified by various characteristics.) Nevertheless, when the number of births is used for analytic purposes, the number of events that actually occurred can be thought of as one in a large series of results that *could have* occurred under the same circumstances. When considered in this way, the number of births is subject to random variation.¹

In this volume, the difference in two percentages is considered statistically significant if the probability that the difference is due to random variation (in the sense of the previous paragraph) is less than 5 percent. All differences explicitly referred to in the text of this volume are statistically significant *whether or not* the term "significant" is present in the sentence.

Smoking during pregnancy (percent of total births to mothers who smoked during pregnancy) is the percentage of women who smoked during pregnancy. In 1998, data for smoking were not collected in NCHS' standard format in California, and not collected at all in New York (except New York City), Indiana, and South Dakota. See the next section for more details. Birth certificates on which information on smoking during pregnancy was not reported were not included in this calculation.

¹ For more information, see Technical Notes in Ventura, S.J., Martin, J.A., Curtin S.C., Mathews T.J., and Park M.M. (2000), "Births: Final data for 1998," *National Vital Statistics Reports*, Vol. 48, no. 3. Hyattsville, Maryland: National Center for Health Statistics.

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics.

Teen births (percent of total births to teens) was calculated by dividing the number of births to women under age 20 by the total number of births to women of all ages. It should be noted that this is not the same as a teen birth rate, which measures the risk that a teenager will give birth. The percentage of total births to women under age 20 is affected by the fertility of women over age 20, as well as the fertility of teenagers.

Sources: 1990-1998 Natality Data Set CD Series 21, numbers 2-9 and 11, National Center for Health Statistics.

Trend, as used in this report, means a general upward or downward change in a percentage over a specific multi-year time period. For example, the decrease in the percentage of Arizona births to mothers who received late or no prenatal care between 1990 and 1998 is a downward trend.

U.S. Average is obtained by dividing total U.S. births with a given characteristic by total U.S. births (as reported by NCHS). For Smoking During Pregnancy, the average is based only on those states with reported data. Data were not available for the following states: California, New York, Indiana, and South Dakota.

Reporting issues

Births to unmarried women.² As of 2000, "births to unmarried women are identified by a question on the birth certificates of all but two States. . . . In the two States (Michigan and New York) that use inferential procedures to compile birth statistics by marital status . . . , a birth is inferred as nonmarital if either of these factors is present: a paternity acknowledgment³ was received or the father's name is missing."

However, in 1990, the first year included in this volume, six states used inferential procedures.⁴ Thus, during the 1990-1998 period covered by this volume, four states—California (1997), Connecticut (1998), Nevada (1997),

² Unless otherwise indicated, this discussion and all direct quotations are drawn from Ventura, S.J., and Bachrach, C.A. (2000), "Nonmarital Childbearing in the United States, 1940-99," *National Vital Statistics Reports*, vol. 48, no. 16. Hyattsville, Maryland: National Center for Health Statistics.

³ A paternity acknowledgment is an acknowledgment from a man who is not married to a child's mother that he is the child's father.

⁴ Ventura, S.J., and Martin, J.A. (1993), "Advance Report of Final Natality Statistics, 1990," *Monthly Vital Statistics Reports*, vol. 41, no. 9 (supplement). Hyattsville, Maryland: National Center for Health Statistics.

Appendix 2: Definitions, Data Sources, and Reporting Issues

and Texas (1994)—switched from using inferential procedures to using a direct question to identify births to unmarried women.⁵ At each of these transition points, there is the potential for the percentage of births to unmarried women to change because of the change in the method of identifying births to unmarried women.

In addition, among those states using inferential procedures, in some states the inferential procedures changed during the 1990-1998 period. For example, in the counts of births to unmarried women submitted by Michigan to NCHS, births with paternity acknowledgments were counted as births to unmarried women beginning in 1994. Thus, there is an undercount of about 25 percent for births to unmarried women in Michigan between 1990 and 1993.

Table A2.1 summarizes the reporting transitions made by each of the affected states by providing the date of each change and the likely effect of the change in reporting as estimated by NCHS. In some cases, these changes have substantial effects on the percentage of births to unmarried women that are included in the tables in this volume. In such cases, trend analysis is either impossible or must be limited to an abbreviated period. All such cases are indicated with footnotes in both the text and the table.

Late or no prenatal care. There were several states in which the percentage of women who received late or no prenatal care changed substantially between two consecutive years. In these situations, we consulted with both NCHS staff and staff of the appropriate state department of health to check into the possibility that reporting problems were responsible, at least in part, for the changes. Table A2.2 summarizes all instances in which reporting problems were documented.

Low maternal education. In 1990 and 1991, Washington and New York State (exclusive of New York City) did not require reporting of educational attainment.

Race and ethnicity. Oklahoma did not ask about Hispanic origin until 1991. New Hampshire began asking for Hispanic origin in 1993, but did not collect this information reliably until 1994. Thus, data on race and ethnicity of births are not included in the table in 1990 for Oklahoma and 1990-1993 for New Hampshire.

In Rhode Island, Hispanic origin was not reported or unknown for 13 percent of total births in 1998—up from only 5 percent in 1990. Consequently, the distribution of births by Hispanic origin is less reliable

⁵ With the exception of Connecticut, these changes took place at the beginning of the year cited. The date of Connecticut's change was June 15.

during the late 1990s than in the early 1990s. Over 90 percent of births with unknown Hispanic origin in 1998 were to white women.

Smoking during pregnancy. Indiana, New York State, and South Dakota did not ask a question on smoking during pregnancy on their birth certificate. California asked it in a form that was not compatible with the standard recommended by NCHS. Oklahoma began reporting smoking data on the birth certificate in 1991, and New York City, a registration area separate from that of New York State, began reporting in 1994.⁶

Repeat teen births. In 1997 and 1998, the number of teen births in which birth order was unknown or not reported was extremely high in Oklahoma. Consequently, the statistics on the percentage of teen births to women who were already mothers are not reliable in those years and are not reported in this volume. Generally, most births with birth order unknown are first births. If it is assumed that most of the births with birth order unknown are actually first-order births, the resulting percentage of repeat teen births for Oklahoma is comparable to that in most other states.

⁶ This paragraph is a direct quotation from Mathews T. J. (1998), "Smoking During Pregnancy, 1990-96," *National Vital Statistics Reports*, vol. 47, no. 10. Hyattsville, Maryland: National Center for Health Statistics.

TABLE A2.1.

Summary of reporting changes for births to unmarried women, 1990-1998

State	Year of change ⁷	Nature of change	Estimated effect of change
California	1995	Changed methodology for inferring mother's marital status by taking into account the naming conventions of Hispanic mothers—especially the use of hyphenated surnames. If the child was given a double surname of the mother's and father's surnames, regardless of sequence, the mother's marital status was coded as "married."	Nonmarital births to Hispanic women in California fell about 17 percent from 1994 to 1995, but nonmarital births declined for other race and ethnic origin groups as well (down 6 percent for non-Hispanic women as a group). This indicates that the drop in Hispanic nonmarital births was not solely the result of the methodological change.
	1997	Began determining mother's marital status by direct question.	Nonmarital births to women aged 15-24 increased, while nonmarital births to older women decreased. These two changes were compensating, so that the overall levels of nonmarital births were only modestly higher in 1997 compared with 1996.
Connecticut	1998 (June 15)	Began determining mother's marital status by direct question as of June 15, 1998. Previously, inferred marital status by comparison of surnames of the parents and child.	The proportion of births to unmarried women was higher (33 percent) in the first six months of 1998 than in the last six months (29 percent). Thus, it appears that the inferential procedures used prior to June 15, 1998, resulted in some overestimation of the number of births to unmarried women.
Michigan	1994	Changed methodology for inferring mother's marital status by taking into account the number of births with paternity acknowledgments.	NCHS estimates that there was underreporting of births to unmarried women of about 25 percent for the 1990-1993 period. Thus, there is substantial discontinuity in the nonmarital birth data between 1993 and 1994.
Nevada	1997	Began determining mother's marital status by direct question in 1997. Previously, inferred marital status by comparing the surnames of the mother, father, and child.	The proportion of births to unmarried women based on the direct question is somewhat lower than the proportion based on the inferential procedures.
New York	1997	New York City changed its method of inferring marital status and began assuming mother is unmarried if the father's name is missing from the birth certificate, or if a paternity affidavit was filed. This change made New York City's methodology consistent with the rest of the state.	Births to unmarried women in New York City (and, thus, New York State) were overstated during 1990 through 1996. (New York City accounts for nearly half of all New York State's births.) The 1997 changes in coding procedures in New York City have resulted in more reliable data for the state.
Texas	1994	Began determining mother's marital status by direct question.	The number of births to unmarried women was underreported during the years 1990-1993. Thus, there is a considerable discontinuity in the data for Texas between 1993 and 1994, when the reported proportion of births to unmarried women increased from 17 to 29 percent statewide.

Sources: Ventura, S.J., Martin, J.A., Curtin S.C., Mathews T.J., and Park M.M. (2000). "Births: Final data for 1998." *National Vital Statistics Reports*, Vol. 48, no. 3. Hyattsville, Maryland: National Center for Health Statistics. Ventura, S.J., Bachrach, C.A. (2000). "Nonmarital childbearing in the United States, 1940-1999." *National Vital Statistics Reports*, Vol. 48, No. 16. Hyattsville, MD: National Center for Health Statistics.

⁷ All changes occurred at the beginning of the year unless otherwise indicated.

Appendix 2: Definitions, Data Sources, and Reporting Issues

TABLE A2.2

Reporting problems with respect to the percentage of women receiving late or no prenatal care

State	Year(s) in which problem occurred	Nature of problem	Implications of problem
Massachusetts	1996	Massachusetts began asking for the exact date of the first prenatal care visit instead of the month prenatal care began.	This change produced more accurate data on the percentage of women receiving late or no prenatal care beginning in 1996. However, the apparent increase in this percentage between 1995 and 1996 may reflect, at least in part, the change in the wording of the question.
Ohio	1997-1998	According to NCHS data for Ohio, the percentage of births for which no prenatal care was reported nearly doubled between 1996 and 1998. ⁸ Over three-quarters of the reported statewide increase in the percentage of births with no prenatal care occurred in Cleveland and Columbus. In Cleveland, this percentage tripled between 1996 and 1998; in Columbus, this percentage increased nine fold. According to a representative of the Ohio Department of Health, this increase may be associated with incomplete recording of information on prenatal care by several hospitals in Cleveland and Columbus.	Since the increase in the percentage of births in which no prenatal care was recorded may be due to reporting problems, the percentage of women receiving late or no prenatal care in 1997 and 1998 may be overstated and should be viewed with caution.

⁸ Source: unpublished tabulation provided by Stephanie Ventura, National Center for Health Statistics.

Appendix 3: Primary Contacts for State KIDS COUNT Projects

The Annie E. Casey Foundation funds a nationwide network of KIDS COUNT projects that provide a more detailed community-by-community picture of the condition of children.

Alabama	VOICES for Alabama's Children	Arkansas	Arkansas Advocates for Children & Families
Linda Tilly <i>Executive Director</i>	PO Box 4576 Montgomery, AL 36103 (334) 213-2410 (334) 213-2413 (fax) ltilly@alavoices.org www.alavoices.org	Julie Robbins <i>Director of Communications & Development</i>	523 S Louisiana Suite 700 Little Rock, AR 72201-4531 (501) 371-9678 (501) 371-9681 (fax) julesrob@swbell.net www.aradvocates.org

Alaska	KIDS COUNT Alaska	California	Children Now
Norm Dinges <i>Project Director</i>	University of Alaska-Anchorage Institute of Social and Economic Research 3211 Providence Dr. Anchorage, AK 99508 (907) 786-7744 (907) 786-7739 (fax) afngd@uaa.alaska.edu www.kidscount.alaska.edu	Amy Dominguez- Arms <i>Director of Policy</i>	1212 Broadway Suite 530 Oakland, CA 94612 (510) 763-2444 (510) 763-1974 (fax) ada@childrennow.org www.childrennow.org

Arizona	Children's Action Alliance	Colorado	Colorado Children's Campaign
Dana Naimark <i>Assistant Director</i>	4001 N 3rd St. Suite 160 Phoenix, AZ 85012 (602) 266-0707 (602) 263-8792 (fax) dnaimark@azchildren.org www.azchildren.org	Kaye Boeke <i>KIDS COUNT Coordinator</i>	225 E 16th Ave. Suite B-300 Denver, CO 80203 (303) 839-1580 (303) 839-1354 (fax) kaye@coloradokids.org www.coloradokids.org

Appendix 3: Primary Contacts for State KIDS COUNT Projects

<p>Connecticut</p> <p>Connecticut Association for Human Services 110 Bartholomew Ave. Suite 4030 Hartford, CT 06106 (860) 951-2212 (860) 951-6511 (fax) www.caahs.org</p>	<p>Florida</p> <p>Susan Weitzel <i>Director</i></p> <p>Center for the Study of Children's Futures Florida Mental Health Institute University of South Florida 13301 Bruce B. Downs Blvd. Tampa, FL 33612 (813) 974-7411 (813) 974-8534 (fax) weitzel@hal.fmhi.usf.edu www.floridakidscount.org</p>	<p>Idaho</p> <p>Linda Jensen <i>KIDS COUNT Director</i></p> <p>Mountain States Group 1607 W Jefferson St. Boise, ID 83702 (208) 388-1014 (208) 331-0267 (fax) ljensen@mtstatesgroup.org www.idahokids.org</p>
<p>Delaware</p> <p>Teresa Schooley <i>KIDS COUNT Project Director</i></p> <p>University of Delaware 298K Graham Hall Newark, DE 19716 (302) 831-4966 (302) 831-4987 (fax) terrys@diamond.net.udel.edu www.dekidscount.org</p>	<p>Georgia</p> <p>Ann Grace Marchetti <i>KIDS COUNT Project Director</i></p> <p>Georgians For Children 300 W. Wieuca Rd., NW Suite 216 Atlanta, GA 30342 (404) 843-0017 (404) 843-0019 (fax) ann@georgians.com www.georgians.com</p>	<p>Illinois</p> <p>Julie Zasadny <i>Director of Communications</i></p> <p>Voices for Illinois Children 208 S LaSalle St. Suite 1490 Chicago, IL 60604 (312) 516-5551 (312) 456-0088 (fax) jzasadny@voices4kids.org www.voices4kids.org</p>
<p>District of Columbia</p> <p>Sonali Patel <i>Director of Public Policy</i></p> <p>DC Children's Trust Fund 2021 L St., NW Suite 205 Washington, DC 20036 (202) 624-5555 (202) 624-0396 (fax) spatel@dcctf.org www.dcchildrenstrustfund.org</p>	<p>Hawaii</p> <p>Marcia Hartsock <i>KIDS COUNT Project Director</i></p> <p>Center on the Family University of Hawaii – Manoa 2515 Campus Rd. Miller Hall 103 Honolulu, HI 96822 (808) 956-4136 (808) 956-4147 (fax) marciah@hawaii.edu uhfamily.hawaii.edu</p>	<p>Indiana</p> <p>Karen Ruprecht <i>Director of Programs</i></p> <p>Indiana Youth Institute 3901 N Meridian St. Suite 200 Indianapolis, IN 46208-4046 (317) 924-3657 (317) 924-1314 (fax) ruprecht@iyi.org www.yi.org</p>



<p>Iowa Child & Family Policy Center 218 Sixth Ave. Suite 1021 Des Moines, IA 50309 (515) 280-9027 (515) 244-8997 (fax) mcrawford@cfpciowa.org www.cfpciowa.org</p>	<p>Louisiana Shannon Johnson KIDS COUNT Coordinator Sjohnson@agendaforchildren.org www.agendaforchildren.org</p>	<p>Massachusetts Massachusetts Citizens for Children 14 Beacon St. Suite 706 Boston, MA 02108 (617) 969-3235 (617) 742-7808 (fax) barryhock@insidehealthcare.com www.masskids.org</p>
<p>Kansas Kansas Action for Children PO Box 463 Topeka, KS 66601 (785) 232-0550 (785) 232-0699 (fax) brunck@kac.org www.kac.org</p>	<p>Maine Lynn Davey KIDS COUNT Director ldavey@mekids.org www.mekids.org</p>	<p>Michigan Jane Zehnder-Merrell KIDS COUNT Project Director janez@mian.net www.mills.org</p>
<p>Kentucky Valerie Salley KIDS COUNT Coordinator vsalley@kyyouth.org www.kyyouth.org</p>	<p>Maryland Jennean Everett-Reynolds KIDS COUNT Project Director jenneanr@aol.com www.acy.org</p>	<p>Minnesota Diane Benjamin KIDS COUNT Director benjamin@cdf-mn.org www.cdf-mn.org</p>

Appendix 3: Primary Contacts for State KIDS COUNT Projects



<p>Mississippi Mississippi Forum on Children & Families, Inc. 737 N President St. Jackson, MS 39202 (601) 355-4911 (601) 355-4813 (fax) kccald@meta3.net www.mfcf.org</p> <p>K.C. Caldwell <i>KIDS COUNT Project Coordinator</i></p>	<p>Nebraska Voices for Children in Nebraska 7521 Main St. Suite 103 Omaha, NE 68127 (402) 597-3100 (402) 597-2705 (fax) voices@uswest.net</p> <p>Janet Johnston <i>Research Coordinator</i></p>	<p>New Jersey Association for Children of New Jersey 35 Halsey St. Newark, NJ 07102 (973) 643-3876 (973) 643-9153 (fax) eloisa@acnj.org www.acnj.org</p> <p>Eloisa Hernandez <i>KIDS COUNT Director</i></p>
<p>Missouri Citizens for Missouri's Children 621 E McCarty Suite D Jefferson City, MO 65101 (573) 634-4324 (573) 634-7540 (fax) cmewest@socket.net www.mokids.org</p> <p>Cande Iveson <i>KIDS COUNT Project Director</i></p>	<p>Nevada Center for Business and Economic Research UNLV 4505 S Maryland Pkwy. Box 456002 Las Vegas, NV 89154 (702) 895-3191 (702) 895-3606 (fax) kidscount@nevada.edu www.nscee.edu/cber</p> <p>Marlys Morton <i>KIDS COUNT Project Coordinator</i></p>	<p>New Mexico New Mexico Advocates for Children & Families 8505 Rancho Diego Place, NE Albuquerque, NM 87113 (505) 823-9324 (505) 244-9509 (fax) kodonnel@uswest.net www.nmadvocates.org</p> <p>Kelly O'Donnell <i>KIDS COUNT Coordinator</i></p>
<p>Montana University of Montana Bureau of Business and Economic Analysis Missoula, MT 59812-6840 (406) 243-5113 steve.seninger@business.umt.edu</p> <p>Steve Seninger <i>Director of Economic Analysis</i></p>	<p>New Hampshire Children's Alliance of New Hampshire 2 Greenwood Ave. Concord, NH 03301 (603) 225-2264 (603) 225-8264 (fax) eshemitz@childrennh.org www.childrennh.org</p> <p>Ellen Shemitz <i>President</i></p>	<p>New York State of New York Council on Children and Families 5 Empire State Plaza Suite 2810 Albany, NY 12223-1533 (518) 474-6294 (518) 473-2570 (fax) debbie.benson@ccf.state.ny.us www.capital.net/com/council</p> <p>Deborah Benson <i>Director of Policy Planning & Research</i></p>

Appendix 3: Primary Contacts for State KIDS COUNT Projects

North Carolina
North Carolina Child Advocacy Institute
 Joann Haggerty
Research & Data Director
 311 E Edenton St.
 Raleigh, NC 27601-1017
 (919) 834-6623
 (919) 829-7299 (fax)
 jhaggerty@intrex.net
 www.ncchild.org

Oklahoma
 Marlo Nash
KIDS COUNT Coordinator
 Oklahoma Institute for Child Advocacy
 420 NW 13th St.
 Suite 101
 Oklahoma City, OK 73103
 (405) 236-5437
 (405) 236-5439 (fax)
 marlonash@oica.org
 www.oica.org

Rhode Island
 Elizabeth Burke
Bryant Executive Director
 Rhode Island KIDS COUNT
 1 Union Station
 Providence, RI 02903
 (401) 351-9400
 (401) 351-1758 (fax)
 ebb@rikidscount.org
 www.rikidscount.org

North Dakota
North Dakota KIDS COUNT
 University of North Dakota
 PO Box 7090
 Gillette Hall
 Room 3
 Grand Forks, ND 58202-7090
 (701) 777-4086
 (701) 777-4257 (fax)
 ann_lochner@mail.und.nodak.edu

Oregon
 Tonia Hunt
Public Policy Director
 Children First for Oregon
 PO Box 14914
 Portland, OR 97214
 (503) 236-9754
 (503) 236-3048 (fax)
 tonia@cifo.org
 www.cifo.org

South Carolina
 Baron Holmes
KIDS COUNT Project Director
 South Carolina Budget & Control Board
 PO Box 12444
 Columbia, SC 29211
 (803) 734-2291
 (803) 734-1276 (fax)
 bholmes@ogc.state.sc.us
 www.ors.state.sc.us/kc.html

Ohio
 David Norris
Communications Manager
 Children's Defense Fund Ohio
 52 E Lynn St.
 Suite 400
 Columbus, OH 43215-3551
 (614) 221-2244
 (614) 221-2247 (fax)
 dnorris@cdfohio.org
 www.cdfohio.org

Pennsylvania
 Diane Ollivier
KIDS COUNT Director
 Pennsylvania Partnerships for Children
 20 N Market Sq.
 Suite 300
 Harrisburg, PA 17101-1632
 (717) 236-5680
 (717) 236-7745 (fax)
 dollivier@papartnerships.org
 www.papartnerships.org

South Dakota
 Carole Cochran
Project Coordinator
 Business Research Bureau
 University of South Dakota
 414 E Clark St.
 Vermillion, SD 57069
 (605) 677-5287
 (605) 677-5427 (fax)
 ccochran@usd.edu
 www.usd.edu/brbinfo/brb/kc

Appendix 3: Primary Contacts for State KIDS COUNT Projects

Tennessee	Tennessee Commission on Children & Youth	Virgin Islands	The Community Foundation of the Virgin Islands	Washington	Human Services Policy Center Institute for Public Policy & Management University of Washington
Pam Brown <i>KIDS COUNT Project Director</i>	Andrew Johnson Tower 9th Floor 710 James Robertson Pkwy. Nashville, TN 37243-0880 (615) 532-1571 (615) 741-5956 (fax) pbrown3@mail.state.tn.us www.state.tn.us/tecy	Dee Baecher-Brown <i>Executive Director</i>	PO Box 11790 38A Dronningens Gade St. Thomas, USVI 00801 (340) 774-6031 (340) 774-3852 (fax) dbrownctvi@worldnet.att.net	Richard Brandon <i>Executive Director</i>	Seattle, WA 98195-3060 (206) 543-8483 (206) 616-5769 (fax) brandon@u.washington.edu www.hspc.org
Texas	Center for Public Policy Priorities	Virginia	Action Alliance for Virginia's Children & Youth	West Virginia	West Virginia KIDS COUNT Fund
Pam Hormuth <i>KIDS COUNT Project Director</i>	900 Lydia St. Austin, TX 78702 (512) 320-0222 (512) 320-0227 (fax) hormuth@cphp.org www.cphp.org/kidscount	Lisa Wood <i>KIDS COUNT Project Director</i>	701 E Franklin St. Suite 807 Richmond, VA 23219 (804) 649-0184 (804) 649-0161 (fax) lisa@vakids.org www.vakids.org	Margie Hale <i>Executive Director</i>	1031 Quarrier St. Suite 313 Atlas Building Charleston, WV 25301 (304) 345-2101 (304) 345-2102 (fax) hh3174@handsnet.org www.wvkidscountfund.org
Utah	Utah Children	Vermont	Vermont Children's Forum	Wisconsin	Wisconsin Council on Children & Families
Terry Haven <i>KIDS COUNT Coordinator</i>	757 East South Temple St. Suite 250 Salt Lake City, UT 84102 (801) 364-1182 (801) 364-1186 (fax) terryh@utahchildren.net www.utahchildren.net	Carlen Finn <i>Executive Director</i>	PO Box 261 7 Court St. 4th Floor Montpelier, VT 05601 (802) 229-6377 (802) 229-4929 (fax) vtcyf@together.net	M. Martha Cranley <i>KIDS COUNT Coordinator</i>	16 N Carroll St. Suite 600 Madison, WI 53703 (608) 284-0580 (608) 284-0583 (fax) mcranley@weccf.org

269

270

Appendix 3: Primary Contacts for State KIDS COUNT Projects

Wyoming

Wyoming Children's Action

Alliance

2622 Pioneer Ave.

Cheyenne, WY 82001

(307) 635-2272

(307) 635-4680 (fax)

kemmons@trib.com

www.wykids.com

Kathy Emmons

Executive

Director

271

272

The Annie E. Casey Foundation is a private charitable organization dedicated to helping build better futures for disadvantaged children in the United States. It was established in 1948 by Jim Casey, one of the founders of the United Parcel Service, and his siblings, who named the Foundation in honor of their mother. The primary mission of the Foundation is to foster public policies, human-service reforms, and community supports that more effectively meet the needs of today's vulnerable children and families. In pursuit of this goal, the Foundation makes grants that help states, cities, and communities fashion more innovative, cost-effective responses to these needs.

Additional copies of this
report are available from:

**The Annie E. Casey
Foundation**

701 St. Paul Street

Baltimore, MD 21202

410.223.2890 Publications Line

410.547.6600

410.547.6624 Fax

www.kidscount.org

273

274



**The Annie E. Casey
Foundation**
701 St. Paul Street
Baltimore, MD 21202
410.547.6600
410.547.6624 Fax
www.kidscount.org



Child Trends
4301 Connecticut Avenue, NW
Suite 100
Washington, DC 20008
202.362.5580
202.362.5533 Fax
www.childtrends.org



U.S. Department of Education
Office of Educational Research and Improvement (OERI)
National Library of Education (NLE)
Educational Resources Information Center (ERIC)



NOTICE

Reproduction Basis



This document is covered by a signed "Reproduction Release (Blanket)" form (on file within the ERIC system), encompassing all or classes of documents from its source organization and, therefore, does not require a "Specific Document" Release form.



This document is Federally-funded, or carries its own permission to reproduce, or is otherwise in the public domain and, therefore, may be reproduced by ERIC without a signed Reproduction Release form (either "Specific Document" or "Blanket").

EFF-089 (3/2000)