

## DOCUMENT RESUME


ED 463 465

CG 031 602

AUTHOR Coridan, Cathi; O'Connell, Cara  
TITLE Meeting the Challenge: Ending Treatment Disparities for Women of Color. A Background Paper.  
INSTITUTION National Mental Health Association, Alexandria, VA.  
SPONS AGENCY Substance Abuse and Mental Health Services Administration (DHHS/PHS), Rockville, MD. Center for Mental Health Services.; National Association of State Mental Health Program Directors, Washington, DC.  
PUB DATE 2001-00-00  
NOTE 26p.; Part of the Targeted Technical Assistance Project of the National Association of State Mental Health Program Directors.  
AVAILABLE FROM National Mental Health Association, 1021 Prince St., Alexandria, VA 22314. Tel: 800-433-5959 (Toll Free); TTY: 800-433-5959 (Toll Free); Fax: 703-684-5968. For full text: [http://www.nmha.org/substance/women\\_disparities.cfm](http://www.nmha.org/substance/women_disparities.cfm).  
PUB TYPE Information Analyses (070)  
EDRS PRICE MF01/PC02 Plus Postage.  
DESCRIPTORS Community Action; Community Health Services; \*Comprehensive Programs; Counseling; Cross Cultural Training; Delivery Systems; \*Females; \*Health Needs; Intercultural Communication; \*Mental Health; \*Minority Groups; \*Substance Abuse  
IDENTIFIERS \*People of Color

## ABSTRACT

Since women are more likely than men to experience co-occurring mental health and substance abuse problems, they have a greater need for comprehensive treatment models. Additionally, women of color often have less access to routine medical care where early diagnosis and intervention can be done, so their mental health and substance abuse problems are often more developed and complicated. Culturally sensitive social service practice is believed to increase many aspects of care including mutual communication, utilization of services, treatment effectiveness, and consumer satisfaction. Following background information on the problem of mental health disorders and substance abuse addictions in minority women, a discussion is included on the stigma surrounding women of color who have mental health or substance abuse problems. Service delivery and treatment program issues are reviewed followed by action steps communities can take to help address these problems. (Contains 88 references.) (JDM)



*Meeting the Challenge:*  
**Ending Treatment  
 Disparities for  
 Women of Color**  
**A Background Paper**



U.S. DEPARTMENT OF EDUCATION  
 Office of Educational Research and Improvement  
 EDUCATIONAL RESOURCES INFORMATION  
 CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE AND  
 DISSEMINATE THIS MATERIAL HAS  
 BEEN GRANTED BY

K. BATTISTA -  
FRAZEE

TO THE EDUCATIONAL RESOURCES  
 INFORMATION CENTER (ERIC)

**BEST COPY AVAILABLE**

# *Meeting the Challenge:*

## Ending Treatment Disparities for Women of Color

### A Background Paper

**Cathi Coridan, MA**

NMHA Senior Director, Substance Abuse Programs and Policy

**Cara O'Connell**

NMHA Intern

Special thanks to NMHA's Working Group on Substance Abuse/Addiction and Paula Heim of the MHA of Hawaii for their input and feedback in the development of this document.

This document was developed by the National Mental Health Association as part of the Targeted Technical Assistance project of the National Association of State Mental Health Program Directors (NASMHPD) and the Division of State and Community Systems Development (Mental Health Block Grant) of the Center for Mental Health Services, Substance Abuse and Mental Health Services Administration, U.S. Department of Health and Human Services.



# Table of Contents

I. Introduction.....	1
II. Background Information.....	1
Mental Health Disorders In Minority Populations.....	1
Mental Health Disorders In Women and Women of Color .....	3
Substance Abuse/Addiction Issues for Women and Women of Color .....	4
Co-occurring Mental Health and Substance Abuse/Addiction Disorders .....	6
III. Stigma.....	7
Stigma and Women.....	7
Criminal Justice Issues .....	8
Related Health Issues.....	8
IV. Service Delivery and Treatment Program Issues.....	9
Societal and Environmental Risk Factors for Women of Color .....	9
The Critical Need to Address Diversity .....	9
Ethnic, Cultural and Language Issues .....	10
Need for a Safe Treatment Environment.....	10
Access to Child Care .....	11
Support Services for Children and Family.....	11
Economic Issues .....	12
Access to Safe Housing.....	12
Access to Transportation .....	12
Access to Health Care .....	13
V. Action Steps .....	13
Community Call to Action .....	14
VI. Bibliography.....	15

# I. Introduction

Research clearly shows that women are more likely than men to experience co-occurring mental health and substance abuse/addiction disorders (NIDA, 1998), thus women have an increased need for comprehensive treatment models. Because the majority of persons who suffer from addictive disorders, and hence those who seek treatment for substance abuse/addiction and co-occurring mental disorders are male and white, women of minority status are often unaware of, or unable to access, appropriate services. When women of color who have mental health problems, substance abuse/addiction and co-occurring disorders do seek help, they can encounter inadequate and irrelevant treatment programs with little or no sensitivities to specific gender, racial and cultural issues, such as childcare and family support services. Additionally, many treatment programs do not provide services that support a woman's ability to recover successfully in the community, such as appropriate job skills training, transportation and affordable housing.

The National Mental Health Association (NMHA) believes that it is essential for community leaders and mental health and substance abuse/addiction stakeholders and treatment providers to work across systems, races and cultures to develop pathways for women of color to access services reflective of, and responsive to, their needs and experiences. Cultural background plays a large role in how the symptoms of mental illness and substance abuse/addiction develop, are reported and interpreted, and consequently, if and how they are treated. Women of color are apt to have less access to routine medical care where early diagnosis and intervention can be done, so their mental health and substance abuse/addiction problems are often more developed and complicated, and their social supports more depleted when they do try to access treatment. Cultural identity and cultural sensitivity issues are complex and cannot be isolated from other

factors that may either draw a woman of color into treatment or cause her to leave treatment (Jones et al, 1997). Culturally sensitive social service practice is believed to increase mutual communication, utilization of services, compliance, treatment effectiveness and consumer satisfaction (Wright & Donoghue, 1997).

Siloed funding, fragmented services and lack of understanding of the treatment needs of women of color create barriers to recovery. NMHA challenges advocates, families and service providers in communities to listen to the voices of these women and provide service coordination and appropriate funding to meet their needs. This background paper provides research-based information, strategies and action steps for grant writing and other funding, advocacy, coalition building, community services assessments and program development.

Now is the time to open the doors for healing to women of color. This process takes commitment, persistence and leadership in each community, and must be guided at all stages by the voices of the women served, and not served, by the system. When these women can heal, and when their families can heal, only then can their communities heal.

## II. Background Information

### **Mental Health Disorders In Minority Populations:**

According to *Mental Health: A Report of the Surgeon General*, the U.S. mental health system is not well equipped to meet the needs of racial and ethnic minority populations, making these populations underserved. The report also states that if members of these groups succeed in accessing services, the treatment may be inappropriate to meet their needs. Without

culturally competent services, this problem will only worsen because of the anticipated growth of minority populations (DHHS, 2000).

Depression, in particular, has often been misdiagnosed in communities of color because of cultural barriers such as language, trust and values in the relationship between doctor and patient, and reliance on the support of family and the religious community rather than mental health professionals during periods of emotional distress. Many individuals in minority cultures "mask" depressive symptoms with other medical conditions, somatic complaints, substance abuse/addiction and other psychiatric illnesses. Socioeconomic factors that contribute to misdiagnosis and inadequate care include limited access to medical care, and a mistrust of medical health professionals based in part on historical higher-than-average institutionalization for members of minority groups with mental illness (NMHA, 2000-B).

The prevalence of mental disorders is estimated to be higher among African Americans than among whites, most likely due to socioeconomic differences. They are under-represented in some private outpatient populations, over-represented in inpatient populations and more likely than whites to use the emergency room for mental health treatment (DHHS, 2000). Numerous studies confirm that African Americans drop out of services at a significantly higher rate than whites and use fewer treatment sessions for mental health services. They enter treatment at a later, more advanced stage than whites, under-consume community mental health services of all kinds, are misdiagnosed more often than whites, and are more often diagnosed with a severe mental illness than whites (WICHE, 1996).

*According to Mental Health: A Report of the Surgeon General, the U.S. mental health system is not well equipped to meet the needs of racial and ethnic minority populations, making these populations underserved.*

The prevailing attitudes of a racial or ethnic group can greatly influence their access to and participation in services. According to an NMHA survey on attitudes and beliefs about depression in the African-American community, approximately 63 percent of African Americans believe that depression is a "personal weakness," compared to the overall survey average of 54 percent. Only 31 percent of African Americans

said they believed depression is a "health" problem, and nearly 30 percent of African Americans said they would "handle it" themselves if they were depressed, while nearly 20 percent said they would seek help for depression from friends and family. Only one in four African

Americans recognize that a change in eating habits and sleeping patterns are signs of depression; only 16 percent recognize irritability as a sign. Only one-third of African Americans said they would take medication for depression, if prescribed by a doctor, compared to 69 percent of the general population. Almost two-thirds of respondents said they believe prayer and faith alone will successfully treat depression "almost all of the time" or "some of the time" (NMHA, 2000-B).

Asian Americans have higher rates of depression than traditionally believed (NIMH, 2000). Many Asian Americans attach stigma to mental disorders because they think that revealing problems or dealing with problems by seeking professional help are signs of personal immaturity, weakness and a lack of self-discipline. They are apprehensive about the reaction of their families to the fact they are seeking help. Some research indicates Asian Americans often somatize mental health problems to avoid shame and maintain the honor of the family (Uba, 1994). These and other cultural factors may cause this group to seek

treatment from a community or religious leader rather than a mental health professional.

There are specific spiritual and cultural relationships that must be taken into consideration in working with the Hawaiian and Pacific Islander communities. Mental illness is often seen as a spiritual experience, a communication from a relative who has passed on, or a result of some misdeed. Substance abuse/addiction and suicide rates are increasing, due to generally lower socio-economic and educational achievement and lack of culturally sensitive service providers. Co-occurrence and exacerbation of physical health problems like hypertension, diabetes and obesity are similar to rates found in the American Indian population (Uba, 1994).

Within the American Indian (AI) population, there are over 400 separate tribes, each of which has its own cultural and social norms. Depression is also a significant problem in American Indian communities. Alcohol abuse and dependence are approximately double the rates found in other population groups, and suicides occur at "alarmingly high levels" (DHHS, 2000).

### **Mental Health Disorders In Women and Women of Color:**

Approximately 12 million women in the United States experience clinical depression each year (NIMH, 1999-A). About one in every eight women can expect to develop clinical depression during their lifetime, roughly twice the rate of men (NIMH, 1999-A/B). Depression occurs most frequently in women aged 25 to 44, prime child-bearing years, and is misdiagnosed approximately 30 to 50 percent of the time (NIMH, 1994, 1999-B). Research shows that

between one-third and one half of depressed people also suffer from some form of substance abuse/addiction or dependence (NIMH, 1999-C). Women who are depressed are more likely to become alcoholic than their male counterparts (McGrath et al., 1995).

Many factors in women may contribute to depression, such as developmental, reproductive, hormonal, genetic and other biological differences. Social factors may also lead to higher rates of clinical depression among women, including stress from work, family responsibilities, the roles and expectations of women and increased rates of sexual abuse and poverty (NIMH, 1995).

*Research suggests that the majority of women suffering from substance abuse/addiction and mental health problems have been victims of rape, incest or physical and emotional abuse.*

Research suggests that the majority of women suffering from substance abuse/addiction and mental health problems have been victims of rape, incest or physical and emotional abuse (Grella, 1996). The life-long psychological and emotional impact of sexual abuse has important implications for treatment and recovery in women with co-occurring disorders. Trauma survivors frequently experience chronic depression and anxiety. Post Traumatic Stress Disorder (PTSD) is the most common anxiety disorder in women and often develops as a result of sexual trauma and abuse. Women with PTSD commonly experience states of both hyper-arousal and dissociation, and substance abuse/addiction can become a form of self-medication that mitigates symptoms of fear, anxiety and depression in survivors of trauma.

Some evidence suggests that women and minorities are most likely to have depression and PTSD (Leshner, 1999), which makes understanding the link between PTSD and

substance abuse/addiction critical in the treatment of women. Substance use disorders have been linked with a host of other risk factors among women, including involvement in criminal acts to obtain drugs (e.g., larceny or prostitution) and participation in high-risk sexual behavior. Similarly, many interpersonal and social conflicts stem from substance use and its associated physical and mental health problems, including instability in relationships, unwanted or early pregnancy, failed educational pursuits and diminished educational achievement, impoverished occupational performance, jeopardized access to employment opportunities, restricted social integration and lifestyle behaviors that affect physical health and emotional well-being (NIDA, 1998).

Females with severe mental illness (SMI) frequently experience conflict and stress in their intimate relationships. Studies indicate higher rates of sexual activity, marriage, divorce, and parenthood among females with SMI compared to their male counterparts. Females with SMI are more likely to choose a spouse with mental illness, which can result in heightened marital conflict and exacerbated symptoms. It is important for treatment programs to address issues surrounding relationships and sexuality as they relate to the lives of women with mental illness.

### **Substance Abuse/Addiction Issues For Women and Women of Color:**

Substance abuse and addiction are among the most critical health and social problems facing the people of the United States today. The use,

misuse and abuse of alcohol and other drugs, including tobacco, are responsible for more than one in four preventable deaths each year, and costs to the economy was estimated to be about \$246 billion for 1992. While whites consistently use the majority of alcohol and drugs, there are consistent use that needs to be addressed among minorities, including minority women. The 1998 National Household Survey on Drug Abuse estimates past month use of any illicit drug by 4.5 percent of Hispanic/Latina females and 5.2 percent of African American females (SAMHSA, 1999).



Compared to other ethnic groups, Hispanics/ Latinas come in second highest in use of alcohol, binge drinking and heavy alcohol use. In the general Hispanic/Latino community, the main substance used is alcohol; the two main illegal drugs used are marijuana and cocaine. Hispanic/Latino families living in poverty are at greater risk for having problems with substance use, and tend to keep substance use problems a 'secret' within the family. Hispanic/Latina females use illegal drugs at a younger

age than males, and often get involved with alcohol and drug use as a way to be 'supportive' of their partner who is using alcohol and drugs. In at least one study, six out of 10 Hispanic/Latina pregnant adolescents report drinking before their third month of pregnancy, and half report smoking marijuana (SAMHSA/CMHS, 1999).

Alcohol is the primary drug of choice among American Indians. While they represent a small percentage of the total US population, their age-adjusted alcohol-related mortality rate is 5.3



times higher than that of the general US population (Walker, et al, 1996). American Indians use and abuse alcohol and other drugs in combination, at a younger age and at higher rates than other groups (SAMHSA/CSAT, 1999). Socioeconomic factors including high unemployment rates and low educational achievement have been used to explain the increased prevalence among this population. Researchers have also suggested that American Indians may be genetically predisposed to alcoholism (Beauvais, 1998).

AI women tend to die at higher rates than other women due to alcohol involved causes. They also have higher rates of co-occurring disorders than their male counterparts. Physical abuse plays a significant role in the development of substance abuse/addiction problems (SAMHSA/CSAT, 1999). Though there is very scant research on American Indians and mental health and substance abuse/addiction prevention and treatment issues, preliminary studies point to a high incidence of co-occurring mental health and substance abuse/addiction problems, including anxiety, depression and suicide (Walker, et al, 1996; WICHE, 1996).

Asian Americans and Pacific Islanders' alcohol and substance abuse/addiction rates vary considerably among subgroups and are influenced by socioeconomic variables including income, educational achievement, and family cohesiveness. Although alcohol and substance abuse/addiction rates are low among female Asian Americans and Pacific Islanders, they are at increased risk for developing co-occurring mental and physical disorders since they often serve as primary caretakers while supporting their families financially (CSAT, 1999). Often cultural expectations can serve as barriers to treatment. In certain social situations, use of

alcohol and mood-altering drugs is seen as culturally acceptable and a necessary aspect of bonding and affiliation (CSAP, 1998).

It is noted throughout the literature that there is severe lack of research on the treatment needs of women, the effects of new treatment on women, and clinical trials for the development of new medications on women (NIDA, 1998; Greenberger, 1998). What is known is that women develop alcohol and drug-related complications, such as liver and heart diseases much more quickly, and after consuming much less of the drug, than men (NIAAA, 2000).

*Minority women tend to initiate alcohol and other drug use at younger ages than non-minority women — a factor that has important implications for prevention efforts.*

Minority women tend to initiate alcohol and other drug use at younger ages than non-minority women — a factor that has important implications for prevention efforts. In addition, they often use drugs in groups of friends, creating a network of dependence

that can be difficult to disengage from. Socioeconomic factors have been linked to differences in substance abuse/addiction patterns within different minority groups.

Nicotine dependence is now the most common psychiatric diagnosis in the United States and the number one cause of preventable death and is rising among females (Bergen & Caporaso, 1999). Factors that have been associated with cigarette use include low socio-economic status, divorce or separation, and the presence of a mental illness (SAMHSA, 1997). Young women between the ages of 18-25 are at the greatest risk for smoking. Among all females, smoking rates are lowest among Hispanic/Latina and Asian/Pacific Islanders and highest in Caucasian and African American women.

Lung cancer is now the leading cause of cancer death for women. According to the recently

released *Women and Smoking: A Report of the Surgeon General - 2001*, socio-economic variables including level of education, income, and neighborhood factors put women of color, who are over-represented in lower SES groups, at significant risk for smoking. Smoking prevalence today is nearly three times higher for women who only have 9-11 years of education (32.9 percent) than among women with 16 or more years of education (11.2 percent) (HHS, 2001).

Additionally, women who smoke experience gender-specific health consequences, including increased risk of adverse reproductive outcomes. Studies indicate high rates of co-occurrence between nicotine dependence and depression, anxiety disorders, and other substance abuse/addiction problems (Breslau et al., 1991). This association is particularly significant for women, who are twice as likely to develop clinical depression and five times more likely to have an anxiety disorder than men. Smoking cessation efforts in women may be complicated by other factors such as post-cessation weight gain and acute increases in tension, anxiety, and irritability (HHS, 2001).

### **Co-occurring Mental Health and Substance Abuse/Addiction Disorders:**

Mental illness and substance abuse/addiction are problems that often occur together, exacerbating and complicating the symptoms and consequences of each disorder, and their treatment. A report from the U.S. Substance Abuse and Mental Health Services Administration (SAMHSA) indicates that in any twelve-month period 10 million people in this country will have a combination of at least one mental health and one substance abuse/addiction disorder.

The prevalence of substance abuse/addiction disorders in patients with severe mental illness is higher than in the general population. While people with schizophrenia do not constitute a large portion of the drug abusing population, an extraordinarily high percentage of people with this disorder abuse drugs (Leshner, 1999).

Studies in the last ten years have demonstrated co-occurrence rates from 29 percent to over 50 percent (Reiger, Farmer, Rae, et. al, 1990; Kessler, 1994). The 1998 National Household Survey on Drug Abuse shows that 52 percent of those with a lifetime history of alcohol abuse or dependence also had a lifetime mental disorder; 59 of those with a lifetime history of drug abuse or dependence had a lifetime mental disorder (SAMHSA, 1999). Three million individuals with co-occurring disorders have at least three disorders; one million have four or more (National Health Policy Forum, 1998).

*[I]n any twelve-month period 10 million people in this country will have a combination of at least one mental health and one substance abuse/addiction disorder.*

Some theories suggest that alcohol and drugs are used by a segment of substance abusing population to self-medicate symptoms of mental disorders such as anxiety and depression. Long term substance abuse/addiction, and the stressors of the accompanying lifestyle, i.e., school and work failure, relationship problems, criminal involvement, can also lead to the development of mental health problems.

Women with eating disorders are also at high risk for alcohol and substance abuse/addiction problems. While the disorders have different causes, the presence of one puts an individual at higher risk for developing the other.

Research shows that co-occurring mental health and substance abuse/addiction disorders are correlated with increased risk of relapse in recovery, violence, incarceration, depression,

suicide, homelessness, and HIV/AIDS, and other sexually transmitted and infectious diseases, such as Hepatitis C (Alexander, 1996; Institute of Medicine, 1999). Severely disabled women are especially vulnerable to assault which can exacerbate mental health and substance abuse/addiction problems.

Women of child-bearing age with mental health and substance abuse/addiction disorders face numerous obstacles

when attempting to access treatment. Few programs are designed to meet the needs of women with children, or who are pregnant, who have co-occurring disorders. Scientific evaluation of these programs to make them "evidence-based," and thus available for replication with grant monies, are even rarer (Grella, 1996).

Because of the prevalence of co-occurring disorders among women who have substance abuse/addiction and mental health disorders, it is critical that service providers be adequately cross-trained and that the gatekeepers of the service systems understand the dynamics of women with co-occurring disorders and assess for them at all points in the treatment process.

### III. Stigma

#### **Stigma and Women:**

The stigma issues surrounding women of color who have mental health and/or substance abuse/addiction disorders are staggering and complex. This is especially true for women who have been arrested and/or incarcerated, lost one or more jobs, had one or more unsuccessful treatment experiences, or perhaps lost custody of her children as a result of a

mental health or drug problem. Poverty, family violence, and lack of adequate housing and job skills add to stigma. Society's judgments may be even harsher toward the drug addicted pregnant woman or woman with children (Grella, 1996). Often the systems that are made to support those with mental health and substance abuse/addiction problems add to the stigma by blaming the consumer for her illness,

or restricting access to care and poorly coordinating care.

No group encounters more stigma, social alienation or difficulty getting basic needs met than persons with mental health and/or

substance abuse/addiction who are homeless, especially women with young children. Once homeless, persons with co-occurring disorders require more services and are more likely to remain homeless than other groups of homeless persons (Drake, Yovetich, Bebout, et. al, 1997).

Stigma often prevents people from learning about what services may be available in their community. For this reason, it is vital to increase public awareness about all the issues related to the mental health and treatment needs of women of color. Public education and media messages must be carefully crafted to help reduce stigma and increase treatment options for women of color.

Marketing and outreach for treatment programs need to be designed to attract the attention of the group that need them most. Studies show that special emphasis needs to be placed on outreach to and engagement of women of color at each step through the referral and treatment process (Jones, 1997; Kline, 1996). Outreach needs to be done repeatedly, in many settings, and by people that women of color can identify with and trust.

## Criminal Justice Issues:

Incarceration and experiences of trauma, abuse and degradation in the justice system increase stigma. Women of color are over-represented in the justice system, and many are there for minor criminal activity related to substance abuse/addiction, mental illness and co-occurring disorders (Peters & Bartoi, 1997, CSAT, 1999). Women arrested for all crimes have more frequent and more severe addictions than men (Grant & Campbell, 1998). Most of the women entering the criminal justice system are under 40, 80 percent are parents and only 11 percent were arrested for a violent offense (CSAT, 1999). In recent years, many women have received long sentences for non-violent, minor offenses because of repeat offender policies (CSAT, 1999).

Diversion programs for non-violent offenders that include treatment services for women with mental health and substance abuse/addiction problems would reduce the number of women in jails and prisons, and lessen the amount of trauma experienced by women in the criminal justice system.

Successful re-entry into the community from jail or prison is compromised by lack of effective, even basic, treatment programs (SAMHSA/CSAT, 1998). Often the justice system is used as the de-facto treatment program for women (and men) who cannot access treatment services in their communities, and done with little or no aftercare or transition planning. Comprehensive and integrated community-based service systems are needed to help women break the cycle of relapse/decompensation, arrest and incarceration (GAINS, 1999).



## Related Health Issues:

Women are the fastest growing population in the AIDS epidemic (AIA, 1996; Nyamathi, et al, 1997). Substance abusing women of childbearing age have been found in one study to be more than 100 times more likely to be HIV positive than all other women of childbearing age in the United States. According to the National Institute on Drug Abuse, infection with the Hepatitis C Virus (HCV) is a significant issue, intertwined with substance abuse/addiction and HIV-AIDS because injection drug use now accounts for at least 60 percent of HCV transmission in the United States (NIDA, 2000).

The issues regarding HIV/AIDS and HCV are complex and critical for women of color with mental health and substance abuse/addiction problems. They are at growing risk for infection due to relationships with infected partners, shared use of contaminated needles, and trading sex for drugs (AIA, 1996). The economic, psychological and medical consequences of HIV/AIDS are devastating, but the pressing need for drugs and money often take precedence over HIV prevention and health for women who live on the streets (Turner & Solomon, 1996). Their children, both born and unborn, are at increased risk for infection and for losing their mothers to early death (Adnopoz, et al, 1997).

The risk of HIV infection is even higher for persons with co-occurring mental health and substance abuse/addiction disorders. Providing ways for those who are HIV positive to address their physical health issues while in treatment is quite complicated but

crucial, since these women are also at risk for additional medical and mental health disorders, ranging from diabetes to dementia. And, they must confront difficult issues such as confidentiality and access to medication (SAMHSA/CSAT, 2000-B).

HIV/AIDS/HVC education and prevention is critical in all communities, and it must include clear messages targeted toward women of color.

## **IV. Service Delivery and Treatment Program Issues**

### **Societal and Environmental Risk Factors for Women of Color:**

Poverty and economic disadvantage are recognized as risk factors to both mental illness and substance abuse/addiction. According to the National Council of La Raza (NCLR), the poverty rates for Hispanic families, working Hispanic families, and Hispanic children remain disproportionately high (Soles, 2000). Poverty, overcrowding, illiteracy, unemployment, the breakdown of two-parent families, and environmental stressors associated with systemic and interpersonal racism have been repeatedly cited as probable causative factors for substance abuse/addiction in African Americans (SAMHSA/CSAT, 1999).

Institutionalized racism continues to support prejudice and discrimination even among those who are well trained and well-intentioned (Wright & Donoghue, 1997). These biases are known to permeate many systems of the health care and human services fields. Effective

programs in inner-city communities should be community-based, family-oriented and multi-generational, since these are the systems that provide the social supports in the African American, Latino, Asian and American Indian cultures.

There is little written specifically about the treatment and prevention of substance abuse and mental health problems of African American, Latina, Asian American and American Indian women, most likely because their data are not captured in the "traditional" settings. Information dissemination from service providers who are working with women of color to the rest of the field is critical to the development of effective programs in communities across the country. And there is an urgent need to find ways to share the stories and experiences of those women who are successfully healing and recovering.

### **The Critical Need to Address Diversity:**

The lack of attention to diversity between and within cultures and races and the absence of culturally competent staff create additional barriers to accessible and effective treatment for women of color. It is critical that treatment programs serving non-English-speaking women have staff who are bi-lingual, and who have a clear understanding of the acculturation process and can address the fears and the needs of immigrant women (Trepper, et. al, 1997).

Also, in addition to concerns and issues related to gender, race, ethnicity and sexual orientation, it is very important to be sensitive to the life experiences of women who have mental illness and/or are addicted, such as trading sex for drugs or having experiences of domestic and family violence.

*Poverty and economic disadvantage are recognized as risk factors to both mental illness and substance abuse/addiction.*

## Ethnic, Cultural and Language Issues:

The prevention literature consistently reports that ritual in families and communities is a very strong protective factor against drug use and other high-risk behavior (NACoA, 1998). Family relationships are the primary support systems in many minority communities, in ways and configurations quite different from the majority culture. Relationships with family and friends who are not drug users provide very valuable supports for women of color while in treatment (Jones, et al, 1997; Kline, 1996; Argeriou & Daley, 1997). The literature strongly supports the critical place of family, religion/spirituality, and culturally similar role modeling and mentoring as important factors in successful recovery for people of color.

Not being able to understand the language, or not feeling as if they can clearly communicate their needs can make women feel frustrated and unwanted, and can create enormous barriers to treatment admission and retention. Having family members, especially children, do translating for a woman while she is speaking of sensitive topics, such as mental illness, sexual history and drug-taking behavior violates family boundaries and cultural taboos, and significantly decreases the likelihood of getting accurate information.

It is crucial that treatment providers receive cultural competency training that promotes awareness and understanding of different racial and ethnic backgrounds, practices and taboos. Successful program participants have identified having treatment plans that included "cultural identity components," such as strong links to healthy immediate and extended family members and faith community, as helpful. Hiring staff that are proficient in a foreign

language or that are of the same ethnic group as their clientele is an effective method of improving treatment services.

## Need for Safe Treatment Environment:

Women with mental health, substance abuse/addiction and co-occurring disorders are more likely to have experienced childhood physical or sexual abuse (Rach-Beisel, Scott & Dixon, 1999). Women need to have treatment providers who ask about, and make it safe to discuss, incest, sexual and physical abuse and experiences on the street exchanging sex for drugs, without being or feeling re-victimized, intentionally or not. Not only do these women need focused attention to their own trauma issues for their recovery, they are also at increased risk for abusing their children (SAMHSA/CSAT, 2000-A). Important components of effective treatment services for women include teaching skills for safety and empowerment and access to services for sexuality and reproduction (Rach-Beisel, Scott & Dixon, 1999).

Studies reveal that women in coed treatment facilities often experience sexual harassment from their male counterparts (PROTOTYPES, 2000). Traditional treatment models where men and women are in recovery groups together, or where confrontation is a core treatment group value, may trigger memories of prior abusive situations, and be unintentionally damaging to persons who have PTSD (Kline, 1996). The treatment women receive is often so abrasive and blaming that it reinforces their distrust in service providers and decreases their willingness to utilize services (Wright & Donoghue, 1997). Availability of single-sex programs is one proven way for communities to provide safer environments for women to heal and recover.

*Family relationships are the primary support systems in many minority communities.*

## Access to Child Care:

By far, the largest barrier to accessing treatment for women of child-bearing age is leaving their children in the care of relatives or in the foster care system, for fear of losing permanent custody of their children (Lewis, et al, 1996; Kline, 1996). Until recently, there were few substance abuse/addiction and mental health treatment programs that would take children or pregnant women, as they pose legal liabilities and can be hard to fund. There is an interesting duality that children are the primary source of motivation for treatment for women, yet fears about the safety and care of their children while in treatment are also the primary reason for poor retention rates. One program in Florida found that women who were allowed to bring their children to the program were five times more likely to stay for a year, and stayed about three times longer than women not offered onsite child care and support treatment (CASA, 1996).

Even when residential programs that accommodate women and their children are

available, there is seldom childcare available in outpatient settings, which may be the only treatment option in a community. Aftercare and ongoing peer support also depend on being able to balance the demands of recovery and parenting, and little attention is paid to these issues in traditional treatment programs. Additionally, women need parenting education, family therapy, parallel treatment for their children, an opportunity for children to live-in with their mothers while in treatment, and "women only" groups for safety and support regarding sexuality, violence and trauma (PROTOTYPES, 2000).

*[C]hildren are the primary source of motivation for treatment for women, yet fears about the safety and care of their children while in treatment are also the primary reason for poor retention rates.*

## Support Services for Children and Family:

Research consistently shows that children of parents with substance abuse/addiction problems and parents with mental illness are at high risk for substance abuse/addiction and mental health problems, and need focused education and support systems (NACoA, 1998). Jail is often a reality for women with substance abuse/addiction problems and over seventy-five percent of all incarcerated women have children (PROTOTYPES, 2000). Support and education services to strengthen family systems and provide support for the children of the women in treatment are critical components of effective treatment programs. There is a strong need for the family to support an individual in the whole process of treatment, from the point of access, through the ups and downs of the treatment and recovery process (Kline, 1996).

The research literature and conventional wisdom both suggest that the family members of persons with substance abuse/addiction and mental health

problems who are in treatment should receive support services. School-aged and younger children are viewed as particularly benefiting from such structured programs (NACoA, 1999). Yet, in a survey of substance abuse/addiction treatment programs, the National Association for Children of Alcoholics reported that sixty-nine percent of the treatment providers that responded to the survey indicated that they do not offer services to the school-aged children of their clients (NACoA, 1999). The primary reason given was lack of funding for such services.

The long-term community benefits of providing support services include substance abuse prevention for the other members of the family. Community stakeholders who understand the need to strengthen the family to improve the outcomes of treatment for women should work to secure the funding to provide education and support services for these children.

### **Economic Issues:**

Women of color who are experiencing mental health problems or have serious substance abuse/addiction problems are less likely to be employed than their male counterparts, with unemployment rates estimated as high as 80 percent (CASA, 1996). This means that they have to rely on publicly funded treatment, increasing their chances of being turned away or put on a waiting list. Women of color may be single parents who are supporting their families with poorly paying jobs, no insurance benefits, who are unable to find funds to pay for treatment even when they can find openings in appropriate programs. Economic pressure can create tension among family members that causes them to discourage treatment for the mother, daughter, wife or girlfriend.

Traditional treatment programs have focused on the job-skill and job-readiness needs of men returning from treatment to the workforce.



Several studies indicate that women are less likely to participate in vocational training services and comprise less than one third of the clientele in these programs. Even when women do complete job skills training, they frequently report

earning salaries less than 56 percent of those earned by their male counterparts. Staff expectations and the quality of services delivered can appear to be lower for female participants. Therefore, the vocational needs of women, who have a higher probability of having few or no job skills, often do not get addressed. This creates barriers for these women to become self-supporting (Grella, 1996). Without changes in the economic and employment prospects for this group of women, any treatment or prevention efforts will have little long term effect (Cohen, 1999).

### **Access to Safe Housing:**

A very significant risk factor for relapse from recovery from mental health and substance abuse/addiction problems is a return to a recovering individual's former environment, possibly one filled with violence (Wright & Donoghue, 1997), and at least filled with cues and people that "tempt" a return to the former lifestyle, whether it be drug or alcohol abuse, non-compliance with medication or an abusive relationship. Safe, affordable housing with adequate space and facilities for children is a critical need in effective treatment for all women, but especially women of color.

### **Access to Transportation:**

Even in metropolitan areas, access to services may be severely restricted by a lack of efficient and reliable transportation. Many individuals seeking substance abuse/addiction treatment have had their driving privileges revoked, and where public transportation is available, it may not be reliable or time effective, especially when a woman in recovery is trying to balance return to the workplace with parenting and treatment needs (Trepper, et. al, 1997). Locating programs on bus lines, providing transportation vouchers and offering services in proximity to related programs such as AA, help keep women in recovery and motivated to continue their journey of healing.



## Access to Health Care:

Women with mental health and substance abuse/addiction problems have higher rates of co-occurring physical health problems, including sexually transmitted diseases, reproductive problems and general health concerns, and they need consistent access to quality health care (Grella, 1996). Primary care providers who treat women of color should be part of community advocacy and treatment teams, providing and receiving training and technical assistance on key issues in substance abuse/addiction and mental health treatment and recovery.

## V. Action Steps

Now is the time for action! This publication was developed to motivate community action. New and existing coalitions need to work together to increase awareness about the prevention, intervention and treatment needs of women of color, and work across systems in local communities to increase access to, and improve the quality of, prevention interventions and treatment services for women of color.

Addressing issues of gender and cultural competence is not easy — it makes people uncomfortable and can challenge long held personal beliefs and community norms. But action on this information is critical — understanding the issues discussed in this paper merely set the stage. Coalitions and community work groups will find both profound challenges and great

satisfaction in addressing the treatment needs of women of color.

**Be sure that appropriate services are available and known.** In some communities it will take time to design and fund programs that address the multiple treatment needs of women of color. Coordination of existing services and sharing information with other family-oriented service providers and community programs can strengthen and enhance the quality of care dramatically and quickly.

**Market services wisely.** Having treatment services available is not effective if these services are unknown in and to the community they serve. Information needs to be available in places where women in need and their families will see them, and formatted in ways that encourage access. Faith communities, youth organizations and other social service providers can be excellent resources for marketing outreach.

**Educate the public and the stakeholders.** Adopt an aggressive strategy to make the needs of these women known to the general public. Promote the many faces of recovery. Because public perception of women with mental health and addiction issues can be very negative, be

*“Having to go to various places for different services is really hard, especially when you have kids and you have to get on the bus and you’ve got to be somewhere at a certain time and you’re tired.*

*It made a real difference to go to my treatment program and have everything I needed. I had parenting training, family therapy, mental health therapy — in group and individually. I also had alcohol and drug sessions. My children played a big role in my recovery. And I am glad they have had this attention from the treatment center now. I hope that they will not follow in my footsteps.” (SAMHSA News, 1999)*

From a woman in recovery for 17 months, now clean and sober, caring for all of her children at home and serving as a consumer advisor to her treatment agency.

She was formerly homeless, addicted to cocaine and heroin, and facing the permanent loss of her five children and prison time for felony narcotics distribution.

sure that their stories and their strengths are emphasized.

**Do not assume that the stakeholders understand all of the needs and issues.** Many believe that the services being provided are adequate. They need to hear about specific gaps and barriers and be shown how more effective treatment is also cost-effective in other systems. Include stakeholders outside of the usual membership of mental health or substance abuse/addiction coalitions, such as local health clinics, community business leaders, faith community leaders, law enforcement and criminal justice personnel and community housing development organizations.

**Outreach is key.** The stigma that women who have mental health and drug abuse issues face is formidable. They need to be invited repeatedly by people whom they believe are credible and trustworthy. They need to see treatment and recovery as something that they can relate to — and succeed at!

**Address myths and misconceptions up front and often.** There are many misconceptions about women of color that suffer from addiction and/or mental illness. There are also many misconceptions within ethnic communities about mental health and substance abuse/addiction issues. All of these misconceptions need to be named and addressed directly, in multiple settings in the community.

The faces and voices of people of similar culture, race and background are critical. Hopeful messages to women and their families have to penetrate a great deal of noise and competing messages. There need to be role models and mentors in communities that convey a sense of leadership and possibility.

Involving consumers and family members is very important. They know what and where the major barriers to treatment and recovery are. They know which programs offer meaningful services — and which don't! People in recovery, their friends and

their families can make excellent and insightful outreach volunteers.

Prevention must be targeted to high-risk groups. Children of parents with mental health and substance abuse problems are known to be at increased risk for an array of psycho-social problems. It is critical to get culturally sensitive and appropriate prevention messages into the community, targeted to families, young children and adolescents who are at increased risk for mental illness or substance abuse/addiction.

### **Community Call to Action:**

It is unnecessary and inexcusable that women of color who suffer from mental health problems, substance abuse/addiction and co-occurring disorders are often unable to access appropriate treatment and prevention services in their communities. The stigma, fragmentation of services and system gridlock are unacceptable.

The information and stories in this background paper are only as useful as they serve to motivate communities to action. Community assessments, media campaigns and coalition meetings need to be outcome based. Policy and funding should respond to need, and providers must be motivated to work together to break old barriers and provide effective services.

The ultimate success of these efforts will be measured by the healing experiences of women who can share the joys and struggles of recovery and contribute to the health of their families and communities. Those in leadership roles must make a commitment to guarantee that sustained prevention and treatment services become more accessible to women of color, that stigma about mental illness and substance abuse/addiction are reduced, and community-based services are provided with gender, race and cultural sensitivity and competency.

## VI. Bibliography

Abandoned Infants Assistance Resource Center (AIA). *Women and Children with HIV/AIDS (factsheet)*. Berkeley, California: Author. 1996.

Acuff, C, Archambeault, J, Greenberg, B, et al. *Mental Health Care for People Living With or Affected by HIV AIDS: A Practical Guide*. Rockville, MD: Research Triangle Institute. 1999.

Alexander, MJ. *Women with co-occurring addictive and mental disorders: An emerging profile of vulnerability*. *Am. J. Orthopsychiatry*. 1996;66:61-70.

Argeriou, M & Daley, M. An examination of racial and ethnic differences within a sample of Hispanic, white (non-Hispanic), and African American Medicaid-eligible pregnant substance abusers. The MOTHERS project. *J Subst Abuse Treat*. 1997;14:489-498.

Bearman, D. Breaking the cycle of dependency: Dual diagnosis and AFDC families. *Journal of Psychoactive Drugs*. 1997;29:359-367.

Beauvais, F. American Indians and alcohol. *Alcohol and Health Research World*. 1998;22:253-259.

Bergen, A & Caporaso, N. Cigarette Smoking. *Journal of the National Cancer Institute*. 1999;91:1365-1375.

Breslau, N. Psychiatric sequelae of posttraumatic stress disorder in women. *Archives of General Psychiatry* 1997;54:81-87.

Breslau, N, Kilbey, M, Adreski, P. Nicotine dependence, major depression, and anxiety in young adults. *Arch Gen Psychiatry*. 1993;50:31-5.

Diaz, J. *Hispanic Children and AIDS factsheet*. National Council of La Raza. Washington, DC: 1993.

Drake, RE, McHugo, GJ, Clark, RE, et al. Assertive community treatment for patients with co-occurring severe mental illness and substance abuse disorders: A clinical trial. *Am. J. Orthopsychiatry*. 1998;68:201-215.

Drake, RE, Mercer-McFadden, C, Mueser, KT, et al. Review of integrated mental health and substance abuse treatment for patients with dual disorders. *Schizophrenia Bulletin*. 1998;24:589-608.

Drake, RE, Mueser, KT, Clark, RE & Wallach, MA. The course, treatment, and outcome of substance disorder in persons with severe mental illness. *Am J Orthopsychiatry*. 1996;66:42-51.

Drake, RE, Yovetich, NA, Bebout, RR, Harris, M & McHugo, GJ. Integrated treatment for dually diagnosed homeless adults. *J. Nervous and Mental Disease*. 1997;185:298-305.

Drug Strategies. *Keeping Score on Alcohol*. Washington, DC: Drug Strategies. 1999.

Entertainment Industries Council, Inc. *Spotlight on Depiction of Health and Social Issues*. Entertainment Industries Council, Inc., National Institute on Drug Abuse, The Robert Wood Johnson Foundation. 2000; 1.

GAINS Center. *The Courage to Change: A Guide for Communities to Create Integrated Services for People with Co-Occurring Disorders in the Justice System*. Delmar, NY: Author,1999.

Grant, B, Hasin, D, & Dawson, D. The relationship between DSM-IV alcohol use disorders and DSM-IV major depression: examination of the primary-secondary distinction in a general population sample. *J Affective Disorders*.1995;38:113-128.

Grant, D. *Delivering Culturally Competent Services to Women & Children Who Are Affected By Drugs*. National Abandoned Infants Assistance Resource Center. Berkeley, CA. 1997.

Grant, E.G & Campbell, J.C. Abuse, drugs, and crime: Pathways to vicious cycle. *Journal of Addictions Nursing*.1998;10:15-27.

Grella, C. Background and overview of mental health and substance abuse treatment systems: Meeting the needs of women who are pregnant or parenting. *Journal of Psychoactive Drugs*.1996;28:319-343.

Grella, C. Services for perinatal women with substance abuse and mental health disorders: The unmet need. *Journal of Psychoactive Drugs*.1997;29:67-78.

Greenberger, P. News from the Society for the Advancement of Women's Health Research: Targeting mental illness and substance abuse among women. *Journal of Women's Health*. 1998;7:407-410.

Hanson, M.S. HIV-risk behaviors and mental health characteristics among homeless or drug-recovering women and their closest sources of social support. *Nursing Research*. 1997;46:133-137.

Hassin, DS, & Nunes, EV Comorbidity of alcohol, drug and psychiatric disorders: epidemiology. In: Kranzler, HR, & Rounsaville, BJ, eds. *Dual Diagnosis and Treatment: Substance Abuse and Comorbid Medical and Psychiatric Disorders*. New York, NY: Marcel Dekker, Inc.,1998.

Jerrell, JM & Wilson, JL Ethnic differences in the treatment of dual mental and substance abuse disorders. A preliminary analysis. *J. Substance Abuse Treatment*. 1997;14:133-140.

Jerrell, JM & Ridgley, MS. Evaluating changes in symptoms and functioning of dually-diagnosed clients in specialized treatment. *Psych Services*. 1995(A);46:233-238.

Jones, J, Hutchins, F, Childress, C et al. *Cultural Identity As An Indicator Of Recovery From Substance Abuse: African American Women's Perspectives*. San Joaquin County, CA. 1997.

Kessler, RC, Nelson, CB, McGonagle, KA, et al. The epidemiology of co-occurring addictive and mental disorders: Implications for prevention and service utilization. *Am. J. Orthopsychiatry*. 1996;66:17-31.

Kessler, RC, McGonagle, KA, Zhao, S., et.al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States; results from the National Comorbidity Survey. *Arch. Gen. Psychiatry*. 1996; 51:8-19.

- Kline, A. Pathways into drug user treatment: the influence of gender and racial/ethnic identity. *Subst Use Misuse*. 1996;3:323-342.
- Lilenfeld, LR & Kaye, WH. The link between alcoholism and eating disorders. *Alcohol Health and Research World*. 1996;20:94-99.
- Leshner, A. Drug abuse and mental disorders: Comorbidity is reality. *NIDA Notes*. 1999. V.14 #4.
- Lewis-Idema, D, Leinman, J, Meyer, J, et al. *Health Care Access and Coverage for Women: Changing Times, Changing Issues?* New York, NY: The Commonwealth Fund, 1999.
- Lewis, R, Haller, D, Branch, D et al. Treatment for Drug Exposed Women and Their Children: Retention Issues involving Drug-Abusing Women In Treatment. Rockville, MD: *NIDA Research Monograph 166*. 1996.
- McGrath, E, Keita GP, Strickland BR, et al.: *Women and Depression: Risks Factors and Treatment issues*. Washington, DC: American Psychological Association, 1995.
- Merikangas, KR, Stevens, D, Fenton, B. Comorbidity of alcoholism and anxiety disorders: The role of family studies. *Alcohol Health and Research World*. 1996;20:100-106.
- Minkoff, K. Program components of a comprehensive integrated care system for seriously mentally ill patients with substance abuse disorders. In: *New Directions for Mental Health Services*. San Francisco: Jossey-Bass.1991.
- Mueser, KT, Drake, RE & Wallach, MA. Dual diagnosis: a review of etiological theories. *Addiction Behaviors*. 1998;23:727-734.
- National Association for Children of Alcoholics. *Children of Alcoholics: Important Facts*. Rockville, MD: Author. 1998.
- National Association for Children of Alcoholics. *Educational Support Services Needed for Children of Addicted Parents in Treatment*. Rockville, MD: Author. 1999.
- National Center on Addiction and Substance Abuse (CASA). *Substance Abuse and The American Woman*. New York, NY: Author. 1996.
- National Institute on Drug Abuse (NIDA). *Treatment of Drug-Dependent Individuals with Comorbid Mental Disorders*. *National Institutes of Drug Abuse*. Rockville, MD: Author. 1997.
- National Institute on Drug Abuse (NIDA). *Drug Addiction Research and the Health of Women*. National Institute on Drug Abuse. Rockville, MD: Author. 1998.
- National Institute of Mental Health (NIMH). *Depression: Treat It, Defeat It*. Bethesda, MD: Author. 1999-B.
- National Institute of Mental Health (NIMH). *Co-Occurrence of Depression with Medical, Psychiatric and Substance abuse/addiction Disorders*. Bethesda, MD: Author. 1999-C.

- National Institute of Mental Health (NIMH). *Strategic Plan for Reducing Health Disparities*. Bethesda, MD: Author. 2000.
- National Mental Health Association (NMHA). *Clinical Depression and Women (factsheet)*. Alexandria, VA: Author. 2000-A.
- National Mental Health Association (NMHA). *Depression and African Americans (factsheet)*. Alexandria, VA: Author. 2000-B
- National Mental Health Association (NMHA). *Depression: What Women Need To Know (factsheet)*. Alexandria, VA: Author. 2000-C.
- National Mental Health Association (NMHA). *Depression and the Latino Community: What You Need To Know*. Alexandria, VA: Author. 2000-D. (bi-lingual)
- Nunes, E & Quitkin, F. Treatment of Drug-Dependent Individuals With Comorbid Mental Disorders: Treatment of Depression in Drug-Dependent Patients, Effects on Mood and Drug Use. Rockville, MD: *NIDA Research Monograph 172*. 1997.
- Nyamathi, A, Flaskerud, J, Leake, B. HIV-Risk Behaviors and Mental Health Characteristics Among Homeless or Drug-Recovering Women and Their Closest Sources of Social Support. *Nursing Research*. 1997;46:133-137.
- Osher, FC & Drake, RE. Reversing the history of unmet needs: Approaches to care for persons with co-occurring addictive and mental disorders. *Am J Orthopsychiatry*. 1996;66:4-11.
- Osher, FC. A vision for the future: toward a service system responsive to those with co-occurring addictive and mental disorders. *Am J Orthopsychiatry*. 1996;66:71-76.
- Peters, R, Bartoi-Green, M. *Screening and Assessment of Co-Occurring Disorders in the Justice System*. Delmar, NY: GAINS Center. 1997.
- PROTOTYPES Systems Change Center. *Changing and Improving Services for Women and Children: Strategies Used and Lessons Learned*. Culver City, CA: Brown, VB, Ramirez-Fogg, E, Broady, J, et al. 2000.
- Rach-Beisel, J, Scott, J & Dixon, L. Co-occurring severe mental illness and substance use disorders: A review of recent research. *Psychiatric Services*. 1999;50:1427-1434.
- Regier, DA, Farmer, ME, Rae, DS, et al. Comorbidity of mental disorders with alcohol and other drug use: Results from the Epidemiological Catchment Area (ECA) study. *JAMA* 1990;264:2511-2518.
- SAMHSA. *Substance Use Among Women in the United States*. SAMHSA Office of Applied Studies. Rockville, MD: Author. 1997.
- SAMHSA. *Summary of Findings from the 1999 National Household Survey on Drug Abuse*. SAMHSA Office of Applied Studies. Rockville, MD: Author. 2000.

SAMHSA. *Summary of Findings from the 1998 National Household Survey on Drug Abuse*. SAMHSA Office of Applied Studies. Rockville, MD: Author. 1999.

SAMHSA/Center for Mental Health Services (CMHS). *Creating a Vision for the 21st Century: National Congress for Hispanic Mental Health, Summary of Congress Proceedings*. Rockville, MD: Author. 2000.

SAMHSA/CSAP Communications Team. (CSAP) *Following Specific Guidelines Will Help You Assess Cultural Competence in Program Design, Implementation and Management*. Rockville, MD: Author, 1994-A.

SAMHSA/Center for Substance Abuse Prevention (CSAP). *ATOD Resource Guide: African Americans*. Rockville, MD: Author. 1995.

SAMHSA/Center for Substance Abuse Prevention (CSAP). *Cultural Competency for Social Workers: A Guide for Alcoholic and Other Drug Abuse Prevention Professionals Working With Ethnic/Racial Communities*. Rockville, MD: Author. 1995.

SAMHSA/Center for Substance Abuse Prevention (CSAP). *Substance Abuse Resource Guide: Hispanic/Latino Americans*. Rockville, MD: Author. 1996.

SAMHSA/Center for Substance Abuse Prevention (CSAP). *Cultural Competence for Health Care Professionals Working With African-American Communities: Theory and Practice*. Rockville, MD: Author.1998.

SAMHSA/Center for Substance Abuse Prevention (CSAP). *CSAP Resource Guide: American Indians and Native Alaskans*. Rockville, MD: Author.1998.

SAMHSA/Center for Substance Abuse Prevention (CSAP). *Responding to Pacific Islanders: Culturally Competent Perspectives for Substance Abuse Prevention*. Rockville, MD: Author. 1998.

SAMHSA/Center for Substance Abuse Treatment (CSAT). *Assessment and Treatment of Patients with Coexisting Mental Illness and Alcohol and Other Drug Abuse. Treatment Improvement Protocol Series #9*. Rockville, MD: Author.1994.

SAMHSA/Center for Substance Abuse Treatment (CSAT). *Continuity of Offender Treatment for Substance Use Disorders From Institution to Community. Treatment Improvement Protocol Series#30*. Rockville, MD: Author. 1998.

SAMHSA/Center for Substance Abuse Treatment (CSAT). *Cultural Issues In Substance Abuse Treatment*. Rockville, MD: Author. 1999.

SAMHSA/Center for Substance Abuse Treatment (CSAT). *Substance Abuse Treatment for Persons with Child Abuse and Neglect Issues. Treatment Improvement Protocol Series #36*. Rockville, MD: Author. 2000-A

SAMHSA/Center for Substance Abuse Treatment (CSAT). *Substance Abuse Treatment for Persons with HIV/AIDS. Treatment Improvement Protocol Series #37*. Rockville, MD: Author. 2000B

SAMHSA. SAMHSA News Vol. VIII, No. 4. 1999.

Sciacca, K & Thompson, CM. *Program development and integrated treatment across systems: Mental illness, drug addiction and alcoholism.* J. Mental Health Administration. 1996;23:288-297.

Soles, C. *Drink Too Much, Risk Too Much: Alcohol Abuse HIV, and Latinos Factsheet.* National Council of La Raza. Washington, DC: 1999.

Soles, C. *Hepatitis C Factsheet.* National Council of La Raza. Washington, DC: 1999-B.

Soles, C. *Substance Abuse: A Real Situation That Places The Hispanic Community At Risk For HIV Infection And Other Diseases factsheet.* National Council of La Raza. Washington, DC: 2000.

Steinberg, D & Gehshan, S. *State Responses to Maternal Drug and Alcohol Use: An Update.* Washington, DC: National Conference of State Legislatures. 2000.

Swan, N. Gender affects relationships between drug abuse and psychiatric disorders. *NIDA Notes.* Vol. 12 #5. 1997.

Thompson, J. Commentary on the Alcohol Abuse in Urban Indian Adolescents and Women: A Longitudinal Study of Assessment and Risk Evaluation. *American Indian and Alaska Native Mental Health Research.* 1996;7:57-59.

Trepper, TS, Nelson, TS, McCollum, EE, et al. Improving substance abuse service delivery to Hispanic women through increased cultural competencies; a qualitative study. *J Subst Abuse Treat.* 1997;14:225-234.

U.S. Department of Health and Human Services (DHHS) *Women and Smoking: A Report of the Surgeon General - 2001.* Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health, 2001.

U.S. Department of Health and Human Services (DHHS). *Mental Health: A Report of the Surgeon General-Executive Summary.* Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health, 1999.

U.S. Department of Health and Human Services (DHHS). *Special Report to the U.S. Congress on Alcohol and Health.* Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, National Institute on Alcohol Abuse and Alcoholism, 2000.

Uba, L. *Asian Americans: Personality Patterns, Identity, and Mental Health.* NY:Guilford Press. 1994.

Walker, D. Response to Alcohol Abuse in Urban Indian Adolescents and Women: A Longitudinal Study of Assessment and Risk Evaluation. *American Indian and Alaska Native Mental Health Research.* 1996;7:94-97.



Walker, R.D. Alcohol abuse in urban Indian adolescents and women: Longitudinal study for assessment and risk evaluation. *American Indian and Alaska Native Mental Health Research Journal of the National Center*. 1996;7:1-47.

Weiss, RD, Najavits, LM. Overview of treatment modalities for dual diagnosis patients: Pharmacotherapy, psychotherapy, and 12-step programs. In: Kranzler, HR, & Rounsaville, BJ, eds. *Dual Diagnosis and Treatment: Substance Abuse and Comorbid Medical and Psychiatric Disorders*. New York, NY: Marcel Dekker, Inc., 1998.

Western Interstate Commission for the Higher Education (WICHE). *Cultural Competence Standards in Managed Mental Health Care for Four Underserved/Underrepresented Racial/Ethnic Groups*. Boulder, CO: Author. 1996.

Wright, T & Donoghue, K. Obstacles To Culturally Sensitive Services For Drug-Addicted Women And Their Children. In: *AIA Monograph on Cultural Competency*. Author: Berkeley, CA. 1997.

## **Web-Based Resources**

National Mental Health Association — [www.nmha.org](http://www.nmha.org)

Center for Mental Health Services/Knowledge Exchange Network — [www.mentalhealth.org](http://www.mentalhealth.org)

Substance Abuse and Mental Health Services Administration/  
National Clearinghouse on Alcohol and Drug Information — [www.health.org](http://www.health.org)


National Alliance for Hispanic Health — [www.hispanichealth.org](http://www.hispanichealth.org)

National Association for Children of Alcoholics — [www.nacoa.net](http://www.nacoa.net)

Office of Minority Health Resource Center — [www.omhrc.gov](http://www.omhrc.gov)

National Institute on Drug Abuse — [www.drugabuse.gov](http://www.drugabuse.gov)

National Institute of Mental Health — [www.nih.nimh.gov](http://www.nih.nimh.gov)



**1021 Prince Street  
Alexandria, VA 22314-2971  
703-684-7722 • 703-684-5968 (fax)  
800-969-NMHA (6642)  
TTY: 800-433-5959  
<http://www.nmha.org>**



**U.S. Department of Education**  
Office of Educational Research and Improvement (OERI)  
National Library of Education (NLE)  
Educational Resources Information Center (ERIC)



## REPRODUCTION RELEASE

(Specific Document)

### I. DOCUMENT IDENTIFICATION:

Title: <u>Ending Treatment Disparities For Women of color</u>	
Author(s):	
Corporate Source: <u>National Mental Health Association</u>	Publication Date: <u>6/01</u>

### II. REPRODUCTION RELEASE:

In order to disseminate as widely as possible timely and significant materials of interest to the educational community, documents announced in the monthly abstract journal of the ERIC system, *Resources in Education* (RIE), are usually made available to users in microfiche, reproduced paper copy, and electronic media, and sold through the ERIC Document Reproduction Service (EDRS). Credit is given to the source of each document, and, if reproduction release is granted, one of the following notices is affixed to each document.

If permission is granted to reproduce and disseminate the identified documents, please CHECK ONE of the following three options and sign at the bottom of the page.

The sample sticker shown below will be affixed to all Level 1 documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY

Kristin Battista-Frazee

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

**1**

Level 1

Check here for Level 1 release, permitting reproduction and dissemination in microfiche or other ERIC archival media (e.g., electronic) and paper copy.

The sample sticker shown below will be affixed to all Level 2A documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN MICROFICHE, AND IN ELECTRONIC MEDIA FOR ERIC COLLECTION SUBSCRIBERS ONLY, HAS BEEN GRANTED BY

\_\_\_\_\_

\_\_\_\_\_

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

**2A**

Level 2A

Check here for Level 2A release, permitting reproduction and dissemination in microfiche and in electronic media for ERIC archival collection subscribers only

The sample sticker shown below will be affixed to all Level 2B documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN MICROFICHE ONLY HAS BEEN GRANTED BY

\_\_\_\_\_

\_\_\_\_\_

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

**2B**

Level 2B

Check here for Level 2B release, permitting reproduction and dissemination in microfiche only

Documents will be processed as indicated provided reproduction quality permits.  
If permission to reproduce is granted, but no box is checked, documents will be processed at Level 1.

*I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce and disseminate these documents as indicated above. Reproduction from the ERIC microfiche or electronic media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete inquiries.*

**Sign here, → please**

Signature: <u>Kristin Battista-Frazee</u>	Printed Name/Position/Title: <u>Kristin Battista-Frazee, Dir market</u>	
Organization/Address: <u>National Mental Health Association 1021 Prince St. Alexandria, VA 22314</u>	Telephone: <u>703-637-4797</u>	FAX: <u>703-684-5968</u>
	E-Mail Address: <u>kbattista@nmha.org</u>	Date: <u>4/8/02</u>

### III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):

If permission to reproduce is not granted to ERIC, or, if you wish ERIC to cite the availability of these documents from another source, please provide the following information regarding the availability of these documents. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents that cannot be made available through EDRS.)

Publisher/Distributor:
Address:
Price:

### IV. REFERRAL OF ERIC TO COPYRIGHT/REPRODUCTION RIGHTS HOLDER:

If the right to grant this reproduction release is held by someone other than the addressee, please provide the appropriate name and address:

Name:
Address:

### V. WHERE TO SEND THIS FORM:

Send this form to the following ERIC Clearinghouse: **ERIC Counseling & Student Services**  
**University of North Carolina at Greensboro**  
**201 Ferguson Building**  
**PO Box 26171**  
**Greensboro, NC 27402-6171**