

AUTHOR Hirschman, Jay; Schmidt, Stefanie; McKinney, Patricia; Frost, Alberta

TITLE School Lunch Salad Bars: Nutrition Assistance Program Report Series. Special Nutrition Programs.

INSTITUTION Food and Nutrition Service (USDA), Alexandria, VA. Office of Analysis and Evaluation.

REPORT NO CN-02-SB

PUB DATE 2002-04-00

NOTE 42p.

AVAILABLE FROM USDA Food and Nutrition Service, 3101 Park Center Drive, Room 926, Alexandria, VA 22302. E-mail: OANEWEB@fns.usda.gov; Web site: <http://www.fns.usda.gov/oane>. For full text: <http://www.fns.usda.gov/oane/MENU/Published/CNP/FILES/saladbar.pdf>.

PUB TYPE Reports - Descriptive (141)

EDRS PRICE EDRS Price MF01/PC02 Plus Postage.

DESCRIPTORS Children; *Comparative Analysis; Elementary Schools; Elementary Secondary Education; Geographic Location; High Schools; Institutional Characteristics; *Lunch Programs; Mail Surveys; Middle Schools; *Nutrition; *Public Schools; Rural Schools; Suburban Schools; Telephone Surveys; Urban Schools

IDENTIFIERS *School Lunch Program

ABSTRACT

Noting that children's average daily intake of fruits and vegetables is well below recommended minimums, this report responds to a request from the Appropriations Committee Directives, Fiscal Year 2002 to compare fruits/vegetable availability in schools with and without salad bars. The comparison was made using data from the School Nutrition Dietary Assessment Study collected during the 1998-99 school year. Information was collected from School Food Authorities by telephone and from cafeteria managers in sample schools using a mail survey. Section 1 of the report describes salad bar availability and analyzes trends during the 1990s. Section 2 describes what fruits/vegetables are contained in salad bars. Section 3 compares the variety of fruits/vegetables in schools with and without salad bars. Section 4 describes characteristics of schools with and without salad bars. Section 5 notes data limitations, and Section 6 presents discussion and conclusions. The key findings are as follows: (1) salad bars are available at least once weekly in 21 percent of public schools, are most common in high schools, least common in elementary schools, and more common in more affluent schools than in less affluent schools; (2) a wide range of vegetables/fruits are available in salad bars; (3) schools with salad bars offer a wider variety of fruits/vegetables than other schools; and (4) salad bars are more common in rural and suburban schools than in urban schools. The report notes that determining the qualities of fruits/vegetables served or consumed would require additional data and cautions that differences associated with salad bars were not necessarily caused by schools adding salad bars. It is further noted that schools do not, on average, meet nutrition standards for fat, saturated fat, and sodium. The report's three appendices provide supplementary data tables, an exploratory comparison of food and nutrient characteristics of meals in schools with and without salad bars, and a discussion of the study's methodology. (KB)

Reproductions supplied by EDRS are the best that can be made
from the original document.

PS PSTR 6256

ED 479 405

Nutrition Assistance Program Report Series

The Office of Analysis, Nutrition and Evaluation

Special Nutrition Programs

Report No. CN-02-SB

SCHOOL LUNCH SALAD BARS

U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

This document has been reproduced as received from the person or organization originating it.

Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.



United States
Department of
Agriculture

Food and
Nutrition
Service

April 2002

BEST COPY AVAILABLE

PS 031475
ERIC
Full Text Provided by ERIC

“The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA’s TARGET Center at (202) 720-2600 (voice and TDD)”

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.



United States Food and
Department of Nutrition
Agriculture Service

April 2002
Special Nutrition Programs
Report No. CN-02-SB

School Lunch Salad Bars

This report is available on the Food and Nutrition Service web site:
<http://www.fns.usda.gov/oane>.

Suggested Citation:

U.S. Department of Agriculture, Food and Nutrition Service, Office of Analysis, Nutrition and Evaluation "School Lunch Salad Bars" *Nutrition Assistance Program Report Series*, NO. CN-02-SB, Alexandria, VA, 22302.

ACKNOWLEDGEMENTS

Several staff members in the Office of Analysis, Nutrition and Evaluation (OANE), Food and Nutrition Service, U.S. Department of Agriculture contributed to the analyses and production of this report: Jay Hirschman (Staff Director), Stefanie Schmidt and Patricia McKinney (program analysts). Alberta Frost (Office Director) also contributed much with her thoughtful review and comments on the drafts. Child Nutrition Program staff members Susan Fouts, Clare Miller and Lori French provided helpful review from the Program perspective.

OANE staff would like to thank Mary Kay Fox of Abt Associates for providing this project with detailed information on salad bar ingredients that were not available on the SNDA-II data files. We would also like to thank Virginia Webb, Coordinator, On Site Services, National Food Service Management Institute at The University of Mississippi for her valuable assistance in researching the costs of implementing a salad bar.

School Lunch Salad Bars

EXECUTIVE SUMMARY

This report fulfills a request to the Department of Agriculture (USDA) from the Appropriations Committee Directives, Fiscal Year 2002. The following language is contained in House Report 107-116:

School Lunch Salad Bars. –The Committee is concerned about school lunch nutrition, and in particular about increasing the consumption of fruits and vegetables among children. The Committee directs the Department to analyze data collected in the School Nutrition Dietary Assessment Study, Part II to compare the amount of fruit and vegetables available to children in schools with salad/fruit bars versus those without salad/fruit bars. The Committee requests a report on this analysis by April 1, 2002.

As requested, this report compares the availability of fruits and vegetables in schools with and without salad bars using data from the School Nutrition Dietary Assessment Study, Part II (SNDA-II), which were collected during the School Year (SY) 1998-99. SNDA-II data enables us to examine the choice and variety of foods offered at salad bars, but not the quantity in a typical serving or the amount consumed.

Our key findings are as follows:

Salad bar availability varies by grade level and free and reduced price eligibility status:

- Twenty-one percent of public schools offer a salad bar at least once per week.
- Salad bars are most common in high schools and least common in elementary schools. Forty-one percent of high schools, 26 percent of middle schools, and 14 percent of elementary schools offer a salad bar at least once per week.
- On a typical school day, 20 percent of public school children have access to a salad bar, that is, they are enrolled in a school where a salad bar is served.
- Free and reduced price approved children are less likely to be enrolled in a school which offers a salad bar than paid status children because salad bars were more commonly found in the more affluent public NSLP schools.

A wide range of vegetables and fruits are available in salad bars:

- Nearly all salad bars include at least one vegetable, with the most prevalent being lettuce, tomatoes, and other raw vegetables.

- Over one-half of salad bars include at least one type of fruit, with the most prevalent types being fresh fruit and canned fruit.

Schools with salad bars offer a wider variety of vegetables and fruits than other schools:

- At all grade levels, schools with salad bars are more likely to offer green salad, raw vegetables, fresh fruit, canned fruit, and dried fruit than schools without salad bars.
- Elementary schools with salad bars are more likely to offer fruit or vegetable juice. Middle schools with salad bars are more likely to offer legumes, and high schools with salad bars are more likely to offer legumes, cooked vegetables, or french fries (either baked or fried) than schools without salad bars.
- Middle schools with salad bars are more likely to serve baked french fries and less likely to serve fried french fries than middle schools without salad bars. Baked french fries have, on average, a lower percent of calories from fat than fried french fries.

The presence of a salad bar is related to school characteristics and NSLP participation:

- Schools offering a salad bar at least once per week have a lower percentage of students who are free and reduced price approved than schools without salad bars.
- In middle schools and high schools, National School Lunch Program (NSLP) participation rates for all students are higher for schools with a salad bar at least once per week than for schools without salad bars. Urban schools are less likely to have salad bars than rural or suburban schools.
- High schools with 500-999 students are more likely to have salad bars than larger or smaller schools.

Determining the quantities of fruits and vegetables served to or consumed by students would require additional data collection such as a third, expanded version of the FNS-sponsored School Nutrition Dietary Assessment series.

One overarching caveat for this report is that the differences associated with salad bars noted above have not necessarily been caused by schools adding salad bars. It is possible that schools with these pre-existing characteristics were more likely to add salad bars. For example, schools with pre-existing higher NSLP participation may have chosen to add salad bars so one cannot conclude from this report that adding salad bars caused the higher NSLP participation.

Table of Contents

| | Page |
|---|------|
| Executive Summary | iii |
| List of Tables and Figures | vii |
| Introduction | 1 |
| Background | 1 |
| Section I How Common Are Salad Bars? | 2 |
| A. What's the Availability of Salad Bars in Schools? | 2 |
| B. How Available Were Salad Bars to Students in School Year (SY) 1998-99? | 4 |
| C. Did Salad Bars Become More Prevalent During the 1990s? | 4 |
| Section II What Vegetables and Fruits Are Offered in Salad Bars? | 6 |
| A. Number of Fruits and Vegetables | 6 |
| B. Percent of Salad Bars Offering Different Types of Vegetables and Fruits | 6 |
| Section III Is There a Difference in The Variety of Fruits and Vegetables in Schools with Salad Bars and Schools without Salad Bars? | 8 |
| Section IV What Are The Characteristics of Schools Offering SaladBars? | 10 |
| A. Free and Reduced Price Eligibility and Participation in The National School Lunch Program..... | 10 |
| B. Urbanicity and School Size | 11 |
| C. Food and Nutrient Characteristics..... | 13 |
| Section V Limitations | 13 |
| Section VI Discussions and Conclusions | 13 |

Bibliography 15

Appendix A: Supplementary Tables 17

Appendix B: Exploratory Comparison of the Food and Nutrient Characteristics of Meals in Schools with and without Salad Bars 26

Appendix C: Methodology and Limitations 30

List of Tables and Figures

| Table | | Page |
|--------------|---|-------------|
| 1 | Percentage of Public NSLP Schools Offering Different Types of Salad Bars | 3 |
| 2 | Percentage of Public NSLP Schools Offering Entrée or Side Salad Bar: 1991-1992 School Year v. 1998-99 School Year | 5 |
| 3 | Mean Number of Fruits and Vegetables Offered in a Salad Bar in Average Public NSLP School with Salad Bar | 6 |
| 4 | Average Percent of Salad Bars in NSLP Public Schools which include Particular Types of Vegetables | 7 |
| 5 | Average Percent of Salad Bars in NSLP Public Schools which Include Particular Types of Fruits..... | 7 |
| 6 | Types of Fruits and Vegetables Served in NSLP Lunches | 9 |
| 7 | Percentage of Student Enrollment Approved for Free/Reduced Price NSLP Meals, 1998-99..... | 10 |
| 8 | National School Lunch Program Participation Rates in NSLP Public Schools During the Target Week..... | 11 |
| 9 | Availability of Salad Bars in Public NSLP Schools by School Size SY 1998-99 | 12 |

Figures

| | | |
|---|---|----|
| 1 | Free and Reduced Price Students Have Less Access to Salad Bars at Every School Level..... | 4 |
| 2 | Salad Bars are Less Available at Urban Middle and High School | 12 |

School Lunch Salad Bars

Introduction

This report fulfills a request to the Department of Agriculture (USDA) from the Appropriations Committee Directives, Fiscal Year 2002. The following language is contained in House Report 107-116:

School Lunch Salad Bars. –The Committee is concerned about school lunch nutrition, and in particular about increasing the consumption of fruits and vegetables among children. The Committee directs the Department to analyze data collected in the School Nutrition Dietary Assessment Study, Part II to compare the amount of fruit and vegetables available to children in schools with salad/fruit bars versus those without salad/fruit bars. The Committee requests a report on this analysis by April 1, 2002.

As requested, this report compares the availability of fruits and vegetables in schools with and without salad bars using data from the School Nutrition Dietary Assessment Study, Part II (SNDA-II) [Fox, et. al., 2001]. SNDA-II data enables us to examine the choice and variety of foods offered at salad bars, but not the quantity in a typical serving or the amount consumed.

Background

The *Dietary Guidelines for Americans* (U.S. Department of Agriculture and the Department of Health and Human Services, 2000) states: “Eating plenty of fruits and vegetables of different kinds, as part of [a] healthful eating pattern... may help protect you against many chronic diseases. Fruits and vegetables provide essential vitamins and minerals, fiber, and other substances that are important for good health. Most people, including children, eat fewer servings of fruits and vegetables than are recommended. To promote health, [one should] eat a variety of fruits and vegetables---at least 2 servings of fruits and 3 servings of vegetables---each day.” An FNS-sponsored report *Changes in Children’s Diets: 1989-1991 to 1994-1996* (Gleason and Sutor, 2001) found that only 14 percent of school-aged children met the target for fruits and only 17 percent met the target for vegetables. The children’s mean daily intakes of both fruits (1.4 servings) and vegetables (2.6 servings) were below the Food Guide Pyramid’s (U.S. Department of Agriculture, 1995) recommended minimum. Any form of fruits and/or vegetables---fresh, frozen, canned, cooked, dried---whether consumed alone or in a mixture count as Food Guide Pyramid servings. National School Lunch Program (NSLP) meals offer students the opportunity to select from a variety of fruits and vegetables.

The SNDA-II study produced national cross-sectional estimates of the nutrient composition of USDA meals served in elementary and secondary schools in SY 1998-99. The study focused exclusively on public schools, which account for roughly 90 percent of

all institutional NSLP participants. The study design included separate nationally representative probability samples of public School Food Authorities (SFAs), public elementary schools, middle schools, and high schools participating in the NSLP. Thus, study results are generalizable to *public* SFAs and *public* schools nationwide but not to the entire NSLP.¹

The report is organized as follows. Section I describes the availability of salad bars, including an examination of whether the availability of salad bars grew during the 1990s. Section II describes what fruits and vegetables are contained in salad bars. Section III compares the variety of fruits and vegetables in schools with and without salad bars. Section IV describes the characteristics of schools with and without salad bars. The reports limitations are presented in Section V, and Section VI concludes.

Definitions: What is a Salad Bar?

For the purposes of this report, a salad bar is a self-serve station where students can select two or more fruits and/or vegetables.² Some salad bars can provide an entire NSLP reimbursable meal because they also provide meat or meat alternates, breads /grains and milk. The salad bars that comprise an entire reimbursable meal can be either entrée salad bars or theme bars. Theme salad bars include potato bars, soup and salad bars, salad and sandwich bars, and potato and salad bars.³ Some salad bars do not offer all of the required meal components and therefore are typically combined with entrées and milk to make reimbursable lunches. Three categories of salad bars comprise part of reimbursable meals: side salad bars, self-serve fruit bars, and assorted self-serve raw vegetables.

Section I: How Common Are Salad Bars?

Section I. A. What's the Availability of Salad Bars in Schools?

About one-fifth (21 percent) of schools offer a salad bar at least once per week (Table 1).⁴ High schools are most likely and elementary schools are least likely to offer salad bars. Forty-one percent of high schools offered some type of salad bar at least once per week, compared to 26 percent of middle schools and 14 percent of elementary schools. A smaller share of schools offered some type of salad bar on a daily basis: 31 percent of

¹ In SY 1998/99 private schools only account for 7% of the NSLP schools and 3 percent of total enrollment.

² For one school, a potato bar was counted as a salad bar even though it had only one fruit or vegetable—the baked potato.

³ Taco salad bars were not included in our definition of a salad bar although they could include vegetables such as lettuce and tomato.

⁴ Our definition of a salad bar comes from data on what foods were served during one week between September 1998 and May 1999. Some schools may have had salad bars at other times during the school year but not during the reference week. Data from the SNDA-II survey of cafeteria managers suggest that many schools may have salad bars sometimes but did not have them during the reference week. Nine percent all schools (including 15 percent of high schools) had no salad bar during the reference week but had a cafeteria manger who reported that a salad bar was offered at least once per week.

high schools, 20 percent of middle schools, and 10 percent of elementary schools.⁵ The most common types of salad bars are entrée salad bars and side salad bars, with theme salad bars, fruit bars, and self-serve assorted raw vegetables offered by a relatively small share of schools.

| Percentage of Schools Offering | Elementary Schools | Middle Schools | High Schools | All Schools |
|--|---------------------|----------------|--------------|-------------|
| Salad Bars That Are Entire Reimbursable Meal | | | | |
| Daily Entrée Salad Bar | 4*** ⁺⁺ | 12** | 22 | 9 |
| Entrée Salad Bar at Least Once Per Week | 6*** ⁺⁺ | 18** | 31 | 12 |
| Daily Theme Salad Bar (Potato Bar or Combination Salad/Sandwich, Salad/Soup or Salad/Potato Bar) | <0.5 | 0 | 1 | <0.5 |
| Theme Salad Bar at Least Once Per Week | 2 | <0.5* | 3 | 2 |
| Salad Bars That are Part of Reimbursable Meal | | | | |
| Daily Side Salad Bar | 6 | 8 | 7 | 7 |
| Side Salad Bar at Least Once Per Week | 8 | 10 | 10 | 9 |
| Daily Self-Serve Fruit Bar | 2 | 1 | 1 | 2 |
| Self-Serve Fruit Bar at Least Once Per Week | 2 | 1 | 3 | 2 |
| Daily Self-Serve Assorted Raw Vegetables | 1 | 1 | 1 | 1 |
| Self-Serve Assorted Raw Vegetables at Least Once Per Week | 1 | 1 | 1 | 1 |
| All Types of Salad Bars | | | | |
| Daily Salad Bar of Any Type | 10*** ⁺⁺ | 20** | 31 | 16 |
| Any Type Of Salad Bar at Least Once Per Week | 14*** ⁺⁺ | 26** | 41 | 21 |
| <i>Sample Size (Number of Schools):</i> | | | | |
| <i>Unweighted</i> | 385 | 329 | 328 | 1042 |
| <i>Weighted</i> | 45,979 | 12,342 | 12,856 | 71,177 |

*Difference when compared to high schools is statistically significant at the .05 level

**Difference when compared to high schools is statistically significant at the .01 level


⁺⁺ Difference when compared to middle schools is statistically significant at the .01 level

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

⁵ All Schools column in all tables reflects all grade levels. Because elementary schools are more numerous than middle schools or high schools, elementary schools have the largest contribution to the all schools total. The analyses are not weighted by the student enrollment in each school.

Section I.B.: How Available Were Salad Bars to Students in School Year (SY) 1998-99?

Next, we turn to the share of *students* who have access to salad bars, that is, students who are enrolled in schools where salad bars are offered. On a typical day, 20 percent of students are enrolled in a school where a salad bar is offered. Access varies by both grade level and free and reduced price status. High school students are the most likely to have access to a salad bar and elementary school students are the least likely. Thirty-four percent of high school students, 20 percent of middle school students, and 10 percent of elementary school students have access to a salad bar on a typical day. Free and reduced price students are less likely to have access to a salad bar than paid students. On a typical day, 15 percent of free and reduced price students are enrolled in a school where a salad bar is offered, compared with 23 percent of paid students (Figure 1 and Appendix Table A2).



Section I.C: Did Salad Bars Become More Prevalent During the 1990s?

Given the USDA's growing emphasis on the nutritional content of school lunches, a natural question is whether salad bars have become more prevalent over time. We examine this issue by comparing the two rounds of SNDA data. The original SNDA was collected during the 1991-92 school year and SNDA-II was collected during the 1998-99 school year. We examine the proportion of schools that offer either an entrée or side salad bar, since those were the only tabulations that are available for the SNDA data. Note that because the SNDA survey included private schools while the SNDA-II survey did not, the statistics calculated using the two surveys are not directly comparable. Comparing the

data can likely tell us about the direction of change in the availability of salad bars between the two survey years, but not the magnitude of change.

It appears that the availability of salad bars in elementary schools grew during the 1990s (Table 2). The percentage of elementary schools that offered an entrée or side salad bar at least once per week was 7 percent in SNDA and 12 percent in SNDA-II. The availability of daily salad bars also appears to have grown. Four percent of elementary schools offered a daily entrée or side salad bar in 1991-92 compared to 9 percent in 1998-99. Despite positive growth, salad bars remain considerably less prevalent in elementary schools than in secondary schools.

In middle schools and high schools, it appears that the availability of entrée and side salad bars did not increase substantially between the 1991-92 and 1998-99 school years. The data from middle schools show decline in both daily salad bars and salad bars served at least once per week. The data from high schools show a decrease in daily salad bars and perhaps a small increase (37 vs. 38 percent) in the percent of schools that offered salad bars at least once per week.

| Percentage of Schools Offering | Elementary Schools | | Middle Schools | | High Schools | | All Schools | |
|---|--------------------|---------|----------------|---------|--------------|---------|-------------|---------|
| | 1991-92 | 1998-99 | 1991-92 | 1998-99 | 1991-92 | 1998-99 | 1991-92 | 1998-99 |
| Daily entrée or side salad bar | 4 | 9 | 22 | 19 | 33 | 29 | 12 | 15 |
| Entrée or side salad bar at least once per week | 7 | 12 | 29 | 25 | 37 | 38 | 15 | 19 |
| <i>Sample size (unweighted)</i> | 278 | 385 | 92 | 329 | 145 | 328 | 515 | 1042 |

Sources: 1991-92 School Year Weighted tabulations of menu data from SNDA (public and private schools combined) for one week between February 1992 and May 1992. Tabulations taken from p.77 of SNDA report. 1998-99 School Year: Weighted tabulations of menu data from SNDA-II (public schools only) for one week between September 1998 and May 1999.

Note: Published SNDA data are not available in enough detail to analyze whether the differences over time are statistically significant.

Section II: What Vegetables and Fruits are Offered in Salad Bars?

Section II.A: Number of Fruits and Vegetables.

The typical salad bar offers a wide variety of vegetables and fruits, 5.4 vegetables and 1.8 fruits (Table 3). Data on salad bar ingredients other than fruits and vegetables (such as meat and meat alternates, salad dressing, cheese, and croutons) are not available. This section does not describe the fruit and vegetable offerings that were not included as part of a salad bar, but were offered as part of the classic serving line. Section IV below presents a discussion of fruits and vegetables that includes non-salad bar menu items.

Salad bar offerings vary by grade level. Elementary school salad bars offer fewer vegetables on average (4.6) than either middle schools (5.7) or high schools (6.1). The lower number of total vegetables in elementary schools was due in part to the lower number of raw vegetables. Elementary schools offered an average of 2.9 raw vegetables compared to 3.4 in middle schools and 3.7 in high schools. The total number of fruits offered did not differ significantly among grade levels.

| | Elementary Schools | Middle Schools | High Schools | All Schools |
|---|-----------------------|-------------------|-----------------|----------------|
| Mean Number of Vegetables | 4.6** ⁺ | 5.7 | 6.1 | 5.4 |
| Mean Number of Raw Vegetables | 2.9** ⁺ | 3.4 | 3.7 | 3.3 |
| Mean Number of Fruits | 1.9 | 2 | 1.7 | 1.8 |
| <i>Sample Size (Number of Schools):</i> | | | | |
| <i>Weighted</i> | <i>54</i> | <i>86</i> | <i>117</i> | <i>257</i> |
| <i>Unweighted</i> | <i>6,423</i> | <i>3,222</i> | <i>5,267</i> | <i>14,914</i> |

*Difference when compared to high schools is statistically significant at the .05 level

**Difference when compared to high schools is statistically significant at the .01 level

⁺ Difference when compared to middle schools is statistically significant at the .05 level

Note: Only schools serving salad bars are included in the table.

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

Section II.B: Percent of Salad Bars Offering Different Types of Vegetables and Fruits

Nearly all salad bars (98 percent) include at least one vegetable. The most common salad bar offerings are lettuce, tomatoes, and other raw vegetables (Table 4). Ninety-two percent of salad bars include lettuce, 70 percent included tomatoes, and 90 percent include raw vegetables besides lettuce and tomatoes. The most common raw vegetables are broccoli, cauliflower, carrots, cucumber, and green peppers. Other common vegetable offerings are cooked vegetables and non-green salads (such as potato salad). (See Appendix Table A3 for more detail on individual vegetables.)

Over half of all salad bars (58 percent) offer at least one type of fruit (Table 5).

The most common fruit offerings were fresh and canned fruits. Forty-five percent of salad bars include fresh fruit and 35 percent included canned fruit. The most common individual fruit items are fresh apples, fresh bananas, fresh oranges, canned pineapple, and canned peaches. (See Appendix Table A4 for more detail on individual fruits.)

Elementary school salad bars are more likely to offer fresh fruit and less likely to offer canned fruit than high school salad bars.

| | Elementary School | Middle Schools | High Schools | All Schools |
|---|-------------------|----------------|--------------|-------------|
| Any Vegetable | 98 | 100 | 97 | 98 |
| Lettuce | 87 | 97 | 94 | 92 |
| Tomato | 62* | 72 | 77 | 70 |
| Any Raw Vegetable (excluding lettuce and tomato): | 89 | 92 | 89 | 90 |
| Any Cooked Vegetable | 8* | 14 | 20 | 13 |
| Any Non-green Salad | 8** | 13 | 25 | 14 |
| Any Legume | 0***++ | 8 | 11 | 5 |
| Vegetable Soup | 0 | <0.5 | 2 | 1 |
| Baked Potato | 10 | 4 | 4 | 6 |

#Note: Only schools serving salad bars at least once per week are included in this table. Only days in which salad bars were served are included in the analysis.

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

**Difference when compared to high schools is statistically significant at the .01 level.

* Difference when compared to high schools is statistically significant at the .05 level.

++ Difference when compared to middle schools is statistically significant at .01 level

| | Elementary Schools | Middle Schools | High Schools | All Schools |
|-------------------|--------------------|----------------|--------------|-------------|
| Any Type of Fruit | 65 | 49 | 57 | 58 |
| Fresh Fruit | 58** | 39 | 34 | 45 |
| Canned Fruit | 25** | 36 | 48 | 35 |
| Dried Fruit | 11 | 10 | 5 | 9 |
| Frozen Fruit | 7 | 4 | 3 | 5 |

Note: Only schools serving salad bars at least once per week are included in this table. Days in which salad bars were not served were excluded from the analysis.

Source: Weighted tabulations of menu data from SNDA-II **for one week between September 1998 and May 1999.

Section III: Is There a Difference in The Variety of Fruits and Vegetables in Schools With Salad Bars and Schools Without Salad Bars?

Next, this report addresses the issue of whether schools with salad bars serve a wider variety of fruits and vegetables than schools without salad bars. The major finding from Table 6 is that schools with salad bars offer a significantly wider range of fruit and vegetable categories than other schools. For all grade levels, schools with salad bars are more likely to serve green salads, raw vegetables, fresh fruit, canned fruit, and dried fruit. For middle and high school levels, schools with salad bars are also more likely to serve legumes. Elementary schools with salad bars are more likely to serve fruit and/or vegetable juice. High schools with salad bars are more likely to serve cooked vegetables.

For green salads and raw vegetables, the findings are quite striking. For example, in the average elementary school that had a salad bar at least once a week, 87 percent of daily menus included green salad and 81 percent included raw vegetables. In contrast, in the average elementary school with no salad bars green salads were served on 35 percent of daily menus and raw vegetables on only 15 percent of daily menus. The differences are also large for middle schools and high schools.

The findings for french fries are interesting. In middle schools with salad bars, menus are more likely to include baked french fries and less likely to include fried french fries. Baked french fries have a lower percentage of their calories from fat than fried french fries.⁶ In the average middle school that has a salad bar at least once per week, 7 percent of daily menus included fried french fries and 34 percent included baked french fries. In contrast, in the average middle school without a salad bar, fried french fries were served in 16 percent of the daily menus whereas baked french fries were served in 22 percent of the daily menus.

High schools with daily salad bars are more likely to serve french fries of either type (baked or fried) than high schools without salad bars. High schools with daily salad bars serve french fries in 57 percent of daily menus on average compared with 47 percent in high schools without salad bars.

⁶ In the SNDAIL data, the average serving of baked french fries has 41 percent of its calories from fat while the average serving of fried french fries has 47 percent of its calories from fat.

| Table 6 Types of Fruits and Vegetables Served in NSLP Lunches | | | | | | | | | | | | |
|--|--------------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|
| Percentage of Daily Menus Including Menu Item in Salad Bar or not in Salad Bar | | | | | | | | | | | | |
| | Elementary schools | | | Middle schools | | | High Schools | | | All Schools | | |
| | Daily Salad Bar | Salad Bar At Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar At Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar At Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar At Least Once Per Week | No Salad Bar |
| Types of Vegetables: | | | | | | | | | | | | |
| Cooked Vegetables | 39 | 39 | 42 | 47 | 48 | 50 | 65** | 61** | 45 | 50 | 49* | 44 |
| French Fries, Deep Fried | 4 | 3 | 3 | 5* | 7* | 16 | 17 | 17 | 17 | 9 | 9 | 7 |
| French Fries, Baked | 19 | 17 | 18 | 39** | 34** | 22 | 39 | 36 | 33 | 30** | 27** | 21 |
| Potatoes, other | 25 | 25 | 21 | 22 | 23 | 25 | 31 | 29 | 27 | 26* | 26* | 22 |
| Lettuce and/or Tomato [#] | 10 | 8 | 7 | 12 | 10 | 14 | 16 | 15 | 16 | 13 | 11 | 9 |
| Green Salad | 95** | 87** | 35 | 100** | 92** | 59 | 100** | 92** | 61 | 98** | 90** | 43 |
| Other Salads | 6 | 9 | 6 | 12** | 18** | 7 | 36** | 30** | 8 | 20** | 19** | 7 |
| Raw Vegetables | 98** | 81** | 15 | 92** | 84** | 20 | 97** | 86** | 15 | 97** | 83** | 16 |
| Legumes | 7 | 7 | 7 | 17** | 15** | 8 | 19** | 17** | 9 | 13** | 12** | 7 |
| Other Vegetables (Soups, casseroles) | 3 | 4 | 4 | 5 | 4 | 7 | 12 | 11 | 9 | 7 | 6 | 5 |
| Types of Fruit: | | | | | | | | | | | | |
| Canned Fruit | 77** | 72** | 57 | 72** | 72** | 55 | 75** | 74** | 53 | 75** | 73** | 56 |
| Fresh Fruit | 81** | 68** | 38 | 78** | 70** | 54 | 76** | 70** | 50 | 79** | 69** | 42 |
| Fruit/Vegetable Juice (100 percent) | 45** | 34** | 18 | 26 | 22 | 23 | 23 | 22 | 22 | 33** | 27** | 19 |
| Dried Fruit | 15** | 12** | 1 | 13** | 10** | 2 | 7* | 6* | 1 | 12** | 9** | 1 |
| Frozen Fruit | 10 | 9 | 8 | 9 | 8 | 6 | 7 | 6 | 4 | 9 | 8 | 7 |
| Combination of Fresh, Frozen, Canned and/or Dried | 1 | 1 | 1 | 2 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Sample Size (Number of Schools): | | | | | | | | | | | | |
| Unweighted | 38 | 54 | 331 | 61 | 86 | 243 | 86 | 117 | 211 | 185 | 257 | 785 |
| Weighted | 4,719 | 6,424 | 39,555 | 2,452 | 3,223 | 9,119 | 4,083 | 5,268 | 7,589 | 1,1254 | 14,914 | 56,262 |

[#]Note: Typically, the lettuce and/or tomato category includes lettuce and/or tomato that are offered as toppings to sandwiches. A salad bar was counted as having lettuce and/or tomato if it offered tomato but not lettuce. A salad bar was categorized as green salad if it offered lettuce. Chef's salads were categorized as green salads. The raw vegetables category excludes lettuce and tomato.

*Difference in those schools with and without salad bars is statistically significant at the .05 level

**Difference in those schools with and without salad bars is statistically significant at the .01 level

Section IV: What Are The Characteristics of Schools Offering Salad Bars?

This section describes selected characteristics of school food service programs in public NSLP schools that offered salad bars and other fruit/vegetable-based self-serve bars.

Section IV.A. Free and Reduced Price Eligibility and Participation in the National School Lunch Program

Participation in the NSLP is open to all students in participating schools. Students from low-income families are eligible to receive meals free of charge or at a reduced price (The maximum charge for a reduced price lunch is \$0.40 per day). In SY 1998-99, the year data were collected in SNDA II, one-third of students enrolled in public schools were approved for free meal benefits. Another eight percent were approved for reduced price meals. The percentage of students approved for free and reduced price meal benefits is higher in elementary schools (45 percent) than in middle schools (38 percent) or high schools (30 percent). (See SNDA-II report page15.)

Salad bars were more commonly found in the more affluent public NSLP schools (Table 7). Schools offering salad bars at least once a week had a significantly lower percentage of students approved for free and reduced price meals than those schools not offering salad bars. This was evident in elementary, middle and high schools. It is also true in high schools offering salad bars on a daily basis.

| | Schools Offering Daily Salad Bars | Schools Offering Salad Bars at Least Once per Week | Schools without Salad Bars |
|--------------------|-----------------------------------|--|----------------------------|
| Elementary Schools | 38 | 38* | 46 |
| Middle Schools | 34 | 32* | 40 |
| High Schools | 25** | 25** | 33 |
| All Schools | 32* | 32** | 43 |

*Difference in those schools with and without salad bars is statistically significant at the .05 level

**Difference in those schools with and without salad bars is statistically significant at the .01 level

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

On an average day during the target week for the study, approximately 60 percent of all students in public NSLP schools received an NSLP lunch. Participation varied by type of school, with participation being highest in elementary schools (67 percent) and lowest in high schools (39 percent). Participation also varied by meal benefits. Students approved to receive free lunches participated at a higher rate (80 percent overall) than either students approved to receive reduced price lunches (69 percent) or students who paid full price (48 percent). (See SNDA-II report, page15)

The participation rate for students paying full price in middle or high schools was significantly higher for schools offering a salad bar either daily or at least once per week than for schools without salad bars (Table 8). The participation rate was unrelated to availability of salad bars in elementary schools. The positive association between the presence of salad bar and school lunch participation does not necessarily imply that salad bars improve school lunch participation. Other factors could account for the relationship.⁷

| | Free and Reduced | | | Paid (Full Price) | | |
|--------------------|------------------|----------------------------------|--------------|-------------------|----------------------------------|--------------|
| | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar |
| Elementary Schools | 85 | 84 | 83 | 59 | 61 | 56 |
| Middle Schools | 79** | 78** | 71 | 48* | 48* | 37 |
| High Schools | 62 | 64* | 58 | 58** | 58** | 47 |
| All Schools | 75 | 75 | 78 | 48 | 49 | 49 |

*Difference in those schools with and without salad bars is statistically significant at the .05 level

**Difference in those schools with and without salad bars is statistically significant at the .01 level

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

Section IV.B: Urbanicity and School Size

Among public NSLP middle schools and high schools, urban schools are less likely to have salad bars than are suburban or rural schools (Figure 2 and Appendix Table A7). Among high schools, those in the 500-999 student range are most likely to have salad bars. (Table 9)

High school size and urbanicity are closely related. For example, urban areas are more likely to have high schools with 2000 or more students than rural or suburban areas. In order to disentangle the relationship between urbanicity and school size, we conducted a regression analysis of the determinants of whether a schools had a salad bar. In a multivariate context, both urbanicity and school size are important determinants of salad bar availability in high schools (See Appendix Tables A5 and A6.)

⁷For example, it is possible that schools that already had high participation rates have been the first to adopt salad bars.

**Figure 2. Salad Bars are Less Available at
Urban Middle and High Schools**

Percent of Schools with Salad Bar At Least Once Per Week
* = Statistically significant difference (.05 level) when compared to urban schools




Table 9
Availability of Salad Bars in Public NSLP Schools by School Size, SY 1998-99

| | High Schools | | | | All Schools | | | |
|--|---------------|------------------|--------------------|----------------|---------------|------------------|--------------------|----------------|
| | <500 students | 500-999 Students | 1000-1999 Students | 2000+ Students | <500 students | 500-999 Students | 1000-1999 Students | 2000+ Students |
| Daily Salad Bar | 18* | 48 | 24* | 33* | 12 | 20 | 17 | 27 |
| Salad Bar at Least Once Per Week | 29* | 63 | 32* | 33* | 18 | 25 | 24 | 28 |
| <i>Sample Size (Number of Schools)</i> | | | | | | | | |
| <i>Unweighted</i> | 72 | 81 | 112 | 52 | 392 | 384 | 177 | 55 |
| <i>Weighted</i> | 2587 | 4053 | 4145 | 1756 | 31,441 | 28,622 | 6,786 | 2,118 |

*Statistically significant difference (.05 level) when compared to schools with 500-999 students

**Statistically significant difference (.01 level) when compared to schools with 500-999 students

Note: Elementary schools and middle schools are not shown separately because sample sizes are too small to produce reliable estimates for some size categories.

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

Section IV.C: Food and Nutrient Characteristics:

One of the major findings of the SNDA-II study is that the average school lunch did not meet *Dietary Guidelines for Americans* recommendations for percent of calories from fat and saturated fat. Lunches served to students in SY 1998-99 provided more than one-third of the Recommended Dietary Allowances (RDAs) for targeted nutrients and calories except in secondary schools, where calories fell short of meeting the recommended level. (See SNDA-II report, page 76) In the context of this report, this begs the question “**Are Meals Served at Schools with Salad Bars More Nutritious?**” SNDA-II did not collect data on the specific quantities of different foods selected by students from self-serve salad bars. In the SNDA-II report assumptions were made about the foods selected to enable nutritional analysis, a reasonable approach given that salad bars were available at only about 20 percent of the schools. However, without data on the actual quantities selected from the salad bar, a direct comparison of the nutrient content of meals served at schools with and without salad bars cannot be considered reliable. As an *exploratory* consideration, Appendix B examines food and nutrients using the SNDA-II assumptions on selections from salad bars.

Section V: Limitations

This report is limited by the SNDA-II data, which provides information on foods offered and average meals served, but not on actual intake or plate waste. It is also limited to public schools, as the SNDA-II sample did not include private schools. The lack of data on selection of specific foods from self-serve salad bars restricts analysis intended to compare the food and nutrient content of average meals at schools with salad bars relative to other schools. A more detailed discussion of limitations is provided in Appendix C.

Section VI: Discussion and Conclusions

Using data from SY 1998-99, our analysis finds that salad bars are available at least once per week in 21 percent of public schools. The availability of salad bars is low in schools with high percentages of children approved for free and reduced price meals and schools with low rates of school lunch participation. Salad bars are less likely to be available in urban schools than in suburban schools and rural schools. Salad bars are most common in high schools and least common in elementary schools.

Access to salad bars is associated with access to a greater variety of fruit and vegetable offerings. Because of data limitations, we cannot determine whether access to salad bars is associated with increased *consumption* (intake) of fruits and vegetables.

One barrier to expanding access to salad bars could be the cost. Anecdotal reports indicate that many school systems believe that self-service salad bars greatly increase meal costs due to the need for substantial upfront equipment investment as well as ongoing costs related to increased food and labor costs. Studies are needed to validate or refute this common belief. Detailed information from one elementary school that recently implemented a salad bar indicated that the start-up costs were approximately \$7000.

Other issues schools must take into account when considering the adding a salad bar include additional labor costs for preparation of items (i.e. washing, cleaning, chopping produce, etc), replenishing items during service periods and cleaning the unit, monitoring and documenting temperatures for food safety purposes, monitoring student use of salad bar for proper use (related to food safety, serving portions for a reimbursable meal and plate waste) and additional time for students serving themselves.

The SNDA-II original report and an exploratory analysis in this report indicate that while at all grade levels, schools with and without salad bars are generally doing well in providing RDA nutrients, they do not on average meet the NSLP nutrition standards for fat, saturated fat and sodium. Continued improvement will be needed to achieve meals offered, served and consumed that are fully consistent with the principles of the *Dietary Guidelines for Americans*. An additional study would be needed to determine whether the availability of school lunch salad bars would actually increase school meal and total daily consumption of fruits and vegetables by students.

Bibliography

Burghardt, John, A. Gordon, N. Chapman, P. Gleason and T. Fraker. "School Nutrition and Dietary Assessment Study School Food Service, Meals Offered, and Dietary Intakes". Project Officer, Patricia McKinney and Leslie Christovich. U.S. Department of Agriculture. Food and Nutrition Service, Office of Analysis, Nutrition and Evaluation, Alexandria, VA, 1993.

Fox, Mary Kay, M.K. Crepinsek, P. Connor and M. Battaglia, "School Nutrition and Dietary Assessment Study-II Final Report", *Nutrition Assistance Report Series*, No. CN-01-SNDAIIFR, Project Officer, Patricia McKinney. U.S. Department of Agriculture. Food and Nutrition Service, Office of Analysis, Nutrition and Evaluation, Alexandria, VA, 2001.

Gleason, Phil and Carol Suitor, "Changes in Children's Diets: 1989-1991 to 1994-1996", *Nutrition Assistance Report Series*, No. CN-01-CD2, Project Officer, Ed Herzog. U.S. Department of Agriculture, Food and Nutrition Service, Office of Analysis, Nutrition and Evaluation, Alexandria, VA, 2001.

National Research Council, *Recommended Dietary Allowances*, 10th edition. Washington, DC: National Academy Press. (1989a)

National Research Council, *Diet and Health: Implications for Reducing Chronic Disease Risk*. Washington, DC: National Academy Press (1989b)

U.S. Department of Agriculture, *The Food Guide Pyramid*. USDA Home and Garden Bulletin 252, 1995.

U.S. Department of Agriculture and U.S. Department of Health and Human Services. *Nutrition and Your Health: Dietary Guidelines for Americans*, 5th edition. Washington, DC: U.S. Government Printing Office. 2000. [Note: NSLP standards are based on the 1990 version of the *Dietary Guidelines*].

Appendix A

| Table | | Page |
|-------|--|------|
| A1 | Nutrition Standards Used in Evaluating School Meals..... | 18 |
| A2 | On a Typical Day, Percentage of Students in Public NSLP Schools with Access to Salad Bar | 19 |
| A3 | Percent of Days in Which a Particular Vegetable Was Offered in an Average Public NSLP School with Salad Bar | 20 |
| A4 | Percent of Days in Which a Particular Fruit Was Offered in an Average Public NSLP School with Salad Bar | 21 |
| A5 | Linear Probability Model: Dependent Variable = Salad Bar served at least once per week..... | 22 |
| A6 | Linear Probability Model: Dependent Variable = Salad Bar served daily | 23 |
| A7 | Availability of Salad Bars in Public NSLP Schools by Urbanicity SY 1998-99 | 24 |
| A8 | Meal Planning Characteristics and Price of Full Price Lunch in Public NSLP Schools with and without Salad Bars, SY 1998-99 | 25 |

Appendix B

| | | |
|----|--|----|
| B1 | Exploratory Analysis-Mean Calorie and Nutrient Content of Average NSLP Lunches Served, SY 1998-99..... | 28 |
| B2 | Exploratory Analysis: Percentage Public Schools Meeting Nutrition Standards For Lunch by Salad Bar Availability, SY 1998-99..... | 29 |

Appendix C

| | | |
|----|---|----|
| C1 | Coding Rules for Fruits and Vegetables in Salad Bars..... | 31 |
|----|---|----|

APPENDIX A
SUPPLEMENTARY TABLES

| Table A1 | |
|--|---|
| Nutrition Standards Used in Evaluating School Meals | |
| NUTRIENT | STANDARD |
| Nutrition Standards Defined in NSLP and SBP Regulations | |
| Nutrients with established Recommended Dietary Allowances (RDAs) | |
| Calories, protein, vitamin A, vitamin C, calcium, and iron | Breakfast: One-fourth of the RDA Lunch: One-third of the RDA |
| Nutrients included in the <i>Dietary Guidelines for Americans</i>: | |
| Breakfast and Lunch: | |
| Total fat | ≤30 percent of total calories |
| Saturated fat | <10 percent of total calories |
| National Research Council <i>Diet and Health</i> –Based Recommendations | |
| Carbohydrate | Breakfast and Lunch: >55 percent of total calories |
| Cholesterol | Breakfast: ≤75 mg Lunch: ≤100 mg |
| Sodium | Breakfast: ≤600 mg Lunch: ≤800 mg |

Note: Recommendations for cholesterol and sodium are equivalent to one-third (lunch) and one-fourth (breakfast) of the recommended maximum daily intake.

BEST COPY AVAILABLE

Table A2
On a Typical Day, Percentage of Students in Public NSLP Schools
With Access To Salad Bar

| | Elementary School Students | Middle School Students | High Schools Students | All Grade Levels |
|---|-------------------------------|------------------------------|-----------------------------|---------------------|
| Students of All School Meal Eligibility Categories | 10 | 20 | 34 | 20 |
| Free and Reduced Price Students | 8 | 16 | 25 | 15 |
| Paid Students | 12 | 21 | 37 | 23 |

Source: Weighted tabulations of data from SNDA-II for one week between September 1998 and May 1999.

Table A3
Percent of Days in Which a Particular Vegetable Was Offered in an Average Public NSLP School with Salad Bar

| | Elementary Schools | Middle Schools | High Schools | All Schools |
|---|--------------------|----------------|--------------|-------------|
| Any Vegetable | 98 | 100 | 97 | 98 |
| Lettuce | 87+ | 97 | 94 | 92 |
| Tomato | 62* | 72 | 77 | 70 |
| Any Raw Vegetable (excluding lettuce and tomato): | 89 | 92 | 89 | 90 |
| Raw Broccoli | 54 | 61 | 51 | 55 |
| Raw Cabbage | 4 | 11 | 5 | 6 |
| Raw Carrots | 71 | 70 | 70 | 70 |
| Raw Cauliflower | 23** | 34 | 42 | 32 |
| Raw Celery | 41* | 34 | 25 | 34 |
| Raw Cucumber | 47***+ | 70 | 72 | 61 |
| Raw Green Pepper | 23** | 41 | 50 | 37 |
| Raw Mushrooms | 6* | 13 | 15 | 10 |
| Raw Onion | 1 | 5 | 12 | 5 |
| Raw Radishes | 10 | 15 | 14 | 13 |
| Raw Spinach | 0 | 0 | 3 | 1 |
| Raw Zucchini | 0 | <0.5 | 2 | 1 |
| Sprouts | 4 | <0.5 | 3 | 3 |
| Any Cooked Vegetable | 8* | 14 | 20 | 13 |
| Canned Beets | 1***+ | 9 | 15 | 8 |
| Canned Carrots | 0 | 1 | 1 | 1 |
| Frozen Peas | 7 | 5 | 6 | 6 |
| Any Non-green Salad | 8** | 13 | 25 | 14 |
| Carrot-Raisin Salad | 0 | 2 | 1 | 1 |
| Coleslaw or Cabbage Salad | 5 | 2 | 4 | 4 |
| Marinated Mushrooms | 0 | 0 | 3 | 1 |
| Potato Salad | 3 | 8 | 14 | 8 |
| Three Bean Salad | 0 | 4 | 2 | 2 |
| Any Legume | 0***+ | 8 | 11 | 5 |
| Canned Baked Beans | 0 | 3* | <0.5 | 1 |
| Canned Chick Peas | 0***+ | 4 | 10 | 5 |
| Canned Kidney Beans | 0 | <0.5 | 3 | 1 |
| Other: | | | | |
| Vegetable Soup | 0 | <0.5 | 2 | 1 |
| Baked Potato | 10 | 4 | 4 | 6 |
| <i>Sample Size (Number of Schools):</i> | | | | |
| <i>Unweighted</i> | 54 | 86 | 117 | 257 |
| <i>Weighted</i> | 6,423 | 3,222 | 5,267 | 14,914 |

*Difference when compared to high schools is statistically significant at the .05 level

**Difference when compared to high schools is statistically significant at the .01 level

+ Difference when compared to middle schools is statistically significant at the .05 level

** Difference when compared to middle schools is statistically significant at the .01 level

Note: Vegetables found in less than one percent of salad bars are not shown as individual vegetables, but are included in the category total. Examples of such vegetables include canned corn and canned okra, which are included in the any canned vegetable total, and red pepper and summer squash, which are included in the any fresh vegetable total.

| | Elementary Schools | Middle Schools | High Schools | All Schools |
|----------------------------------|-----------------------|-------------------|-----------------|----------------|
| Any Type of Fruit | 65 | 49 | 57 | 58 |
| Any Type of Fresh Fruit | 58** ⁺ | 39 | 34 | 45 |
| Apple | 38* | 30 | 23 | 31 |
| Banana | 20 | 18 | 9 | 16 |
| Cantaloupe | 0* ⁺ | 4 | 6 | 3 |
| Grapes | 13* | 16** | 4 | 11 |
| Honeydew | 1 | 1 | 2 | 1 |
| Kiwi | 5 | 6 | 2 | 4 |
| Oranges | 27* | 28 | 18 | 24 |
| Pear | 8 | 4 | <0.5 | 4 |
| Any Type of Canned Fruit | 25** | 36 | 48 | 35 |
| Canned Applesauce | 12 | 10 | 15 | 13 |
| Canned Apricots | 4 | 6 | 2 | 4 |
| Canned Fruit Cocktail | 12 | 13 | 13 | 13 |
| Canned Maraschino Cherries | 3 | 2 | 0 | 2 |
| Canned Peaches | 4** ⁺ | 13 | 22 | 12 |
| Canned Pears | 2* | 14 | 11 | 8 |
| Canned Pineapple | 17 | 23 | 26 | 22 |
| Any Type of Dried Fruit | 11 | 10 | 5 | 9 |
| Dates | 7* | 4* | <0.5 | 4 |
| Raisins | 4 | 4 | 4 | 4 |
| Any Type of Frozen Fruit | 7 | 4 | 3 | 5 |
| Frozen Blueberries | 3 | 2 | 1 | 2 |
| Frozen Cherries | 4 | 2 | 3 | 3 |
| Sample Size (Number of Schools): | | | | |
| <i>Unweighted</i> | 54 | 86 | 117 | 257 |
| <i>Weighted</i> | 6,423 | 3,222 | 5,267 | 14,914 |

*Difference when compared to high schools is statistically significant at the .05 level

**Difference when compared to high schools is statistically significant at the .01 level

⁺ Difference when compared to middle schools is statistically significant at the .05 level

Note: Fruits found in less than one percent of salad bars are not shown as individual fruits, but are included in the category totals. Examples of such fruits are fresh strawberries, fresh pineapple, and fresh grapefruit, which are counted as any type of fresh fruit in the category total.

Note: Only schools serving salad bars at least once per week are included in this table.

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

| Table A5 | | | |
|---|-------------------|-----------------------|---------------------|
| Linear Probability Model: | | | |
| Dependent Variable=Salad Bar served at least once per week | | | |
| Independent variables | Elementary | Middle Schools | High Schools |
| Intercept | .171** | .403** | .694** |
| Urban | -.0356 | -.217** | -.163* |
| Rural | .0400 | -.0708 | -.0851 |
| Suburban (omitted) | Omitted | Omitted | Omitted |
| 0-499 students | .0231 | -.0388 | -.289* |
| 500-999 students | Omitted | Omitted | Omitted |
| 1000-1999 students | -.0196 | -.0923 | -.292** |
| 2000+ students | -.208 | -.311 | -.212* |
| Free and reduced approval rate | -.00236* | -.00252* | -.00332* |
| Northeast | .00668 | -.0464 | -.113 |
| Mid-Atlantic | -.0146 | .0124 | .128 |
| Southeast | .0496 | .0191 | .0849 |
| Mid-West | Omitted | Omitted | Omitted |
| Southwest | .0779 | .0234 | .00513 |
| Mountain Plains | .114 | .267* | .280* |
| West | .196** | .128 | .0575 |
| <i>R-Squared</i> | .055 | .110 | .180 |
| <i>Sample size</i> | 365 | 301 | 309 |

*Statistically significant difference (.05 level)

**Statistically significant difference (.01 level)

| Table A6 | | | |
|--|-------------------|-----------------------|---------------------|
| Linear Probability Model: | | | |
| Dependent Variable=Salad Bar Served Daily | | | |
| Independent variables | Elementary | Middle Schools | High Schools |
| Intercept | .145** | .291** | .430** |
| Urban | -.0628 | -.172** | -.263** |
| Rural | .0382 | -.0400 | -.0117 |
| Suburban (omitted) | Omitted | Omitted | Omitted |
| 0-499 students | .00000802 | -.0000983 | -.276** |
| 500-999 students | Omitted | Omitted | Omitted |
| 1000-1999 students | -.00282 | -.0817 | -.182** |
| 2000+ students | -.149 | -.222 | -.00182 |
| Free and reduced approval rate | -.00204* | -.00102 | -.00270* |
| Northeast | -.00958 | -.121 | -.0821 |
| Mid-Atlantic | -.0373 | .0536 | .256** |
| Southeast | .0381 | .0232 | .202* |
| Mid-West | Omitted | Omitted | Omitted |
| Southwest | .105 | .0456 | .0801 |
| Mountain Plains | .0628 | .267 | .286** |
| West | .202** | .122 | .163 |
| <i>R-Squared</i> | .074 | .099 | .183 |
| <i>Sample size</i> | 365 | 301 | 309 |

*Statistically significant difference (.05 level)

**Statistically significant difference (.01 level)

| Table A7 Availability of Salad Bars in Public NSLP Schools by Urbanicity, SY 1998-99 | | | | | | | | | | | | |
|---|--------------------|----------|-------|----------------|----------|-------|--------------|----------|-------|-------------|----------|-------|
| | Elementary Schools | | | Middle Schools | | | High Schools | | | All Schools | | |
| | Urban | Suburban | Rural | Urban | Suburban | Rural | Urban | Suburban | Rural | Urban | Suburban | Rural |
| Daily Salad Bars | 6 | 12 | 12 | 7 | 26* | 21* | 10 | 40* | 32* | 6 | 20 | 17 |
| Salad Bars at Least Once Per Week | 10 | 15 | 16 | 11 | 33* | 26* | 22 | 49* | 40* | 12 | 25 | 22 |
| <i>Sample Size (Number of Schools):</i> | | | | | | | | | | | | |
| <i>Unweighted</i> | 97 | 180 | 106 | 92 | 174 | 56 | 75 | 169 | 81 | 264 | 523 | 243 |
| <i>Weighted</i> | 10999 | 22139 | 12745 | 3343 | 7095 | 1811 | 2258 | 6608 | 3617 | 16899 | 35842 | 18173 |

*Statistically significant difference (.05 level) when compared to urban schools.

**Statistically significant difference (.01 level) when compared to urban schools.

BEST COPY AVAILABLE

| Table A8 | | | | | | | | | | | | |
|---|--------------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|
| Meal Planning Characteristics and Price of Full Price Lunch in Public NSLP Schools with and without Salad Bars, SY 1998-99 | | | | | | | | | | | | |
| | Elementary Schools | | | Middle Schools | | | High Schools | | | All Schools | | |
| | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar |
| Who Plans Menu | | | | | | | | | | | | |
| District | 58 | 62 | 64 | 74 | 75 | 66 | 54 | 57 | 62 | 60 | 63 | 64 |
| Off-site kitchen | 10 | 10 | 8 | 6 | 7 | 5 | 1 | 1 | 2 | 6 | 6 | 6 |
| School | 22 | 20 | 19 | 7 | 8 | 14 | 35 | 32 | 26 | 24 | 22 | 19 |
| Combination of above | 10 | 8 | 6 | 12 | 9 | 12 | 9 | 10 | 9 | 10 | 9 | 7 |
| Who Buys Food | | | | | | | | | | | | |
| District | 41 | 40 | 53 | 64 | 58 | 62 | 41 | 43 | 51 | 46* | 45* | 54 |
| Off-site kitchen | 21* | 18* | 8 | 5 | 7 | 4 | <0.5 | 2 | 1 | 10 | 10 | 7 |
| School | 12 | 16 | 16 | 19 | 19 | 15 | 28 | 26 | 25 | 19 | 20 | 17 |
| Combination of above | 27 | 26 | 21 | 11 | 16 | 18 | 31* | 29 | 20 | 25 | 25 | 20 |
| Menu Planning System | | | | | | | | | | | | |
| NSMP/ANSM | 43* | 35 | 27 | 29 | 28 | 28 | 32 | 30 | 26 | 36* | 32 | 27 |
| Enhanced food | 10** | 13** | 31 | 26 | 24 | 30 | 23 | 26 | 30 | 18** | 20** | 31 |
| Traditional food | 37 | 41 | 39 | 39 | 40 | 39 | 43 | 42 | 38 | 40 | 41 | 39 |
| Price of Full-Price (Paid) Lunch | | | | | | | | | | | | |
| Price Paid | \$1.40* | \$1.35 | \$1.29 | \$1.49 | \$1.48 | \$1.43 | \$1.48 | \$1.46 | \$1.42 | \$1.45** | \$1.42** | \$1.33 |
| Sample Size (Number of Schools) | | | | | | | | | | | | |
| Unweighted | 38 | 54 | 330 | 61 | 86 | 242 | 86 | 117 | 211 | 185 | 257 | 783 |
| Weighted | 4,719 | 6,424 | 39,555 | 2,452 | 3,223 | 9,119 | 4,082 | 5,268 | 7,589 | 11,254 | 14,914 | 56,263 |

*Difference in those schools with and without salad bars is statistically significant at the .05 level

**Difference in those schools with and without salad bars is statistically significant at the .01 level

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

BEST COPY AVAILABLE

APPENDIX B

EXPLORATORY COMPARISON OF THE FOOD AND NUTRIENT CHARACTERISTICS OF MEALS IN SCHOOLS WITH AND WITHOUT SALAD BARS

This Appendix examines whether lunches offered in schools with salad bars and lunches offered in schools without salad bars differ in the nutritional content. The nutritional analysis includes salad bar and non-salad bar menu items, and all types of foods offered, including fruits and vegetables, meat and meat alternates, milk, and grains. *This analysis is considered exploratory because SNDA-II did not collect data on the specific quantities of different foods selected by students from self-serve salad bars. Therefore, assumptions were made about the foods selected from the salad bar. The results are sensitive to changes in these assumptions, and must be interpreted with caution.*

Food Energy (calories) and RDA Nutrients

Depending on the nutrient and the grade level, there are some differences in the nutrient content of lunches served in schools with and without salad bars (Appendix Table B1). Average calories served are higher for elementary schools with salad bars than those without salad bars. For all grade levels, lunches served in schools with salad bars average higher levels of vitamin A. In elementary schools and middle schools, with daily salad bars, lunches served in schools with salad bars average higher levels of vitamin C.

Next, this Appendix examines the percent of schools meeting the NSLP Nutrition Standards for Lunch (Appendix Table B2). Schools with salad bars are more likely to meet the nutrient standards for some types of nutrients and some grade levels. For all grade levels, schools with salad bars are significantly more likely to meet the RDA-based NSLP nutrient standards for Vitamin C with the exception of elementary schools with salad bars at least once per week. High schools with salad bars are more likely to meet the RDA-based NSLP standard for vitamin A. However, elementary schools with salad bars are less likely to meet the nutrient standard for iron.

Fat and Saturated Fat

In addition, there are some findings that suggest that schools with salad bars are more likely to meet the standards for saturated fat, which state that on average less than 10 percent of calories should come from saturated fat. Specifically, public high schools offering salad bars are more likely to average less than 10 percent of calories from saturated fat than high schools with no salad bars. Twenty-two percent of high schools with daily salad bars and 11 percent of high schools with no salad bars meet the saturated fat recommendation. Elementary schools with daily salad bars or salad bars at least once per week are more likely to meet the recommended levels of saturated fat than elementary schools with no salad bars. But an important caveat should be used in

interpreting the elementary school results. In particular, elementary schools with salad bars are more likely to meet the *Dietary Guidelines* for saturated fat, but they are also more likely to have relatively high levels of saturated fat.⁸

Sodium

In regards to sodium, very few elementary or middle schools met the recommended level and the amount of sodium in an average elementary schools with daily or at least once per week salad bars was even higher than for schools without a salad bar.

⁸ In particular, the 75th percentile of saturated fat level for elementary schools with daily salad bars, with salad bars at least once per week, and no salad bars are 13.4 grams, 13.4 grams, and 13.0 grams, respectively.

Table B1
Exploratory Analysis-Mean Calorie and Nutrient Content of Average NSLP Lunches Served, SY
1998-99

| | Elementary Schools | | | Middle Schools | | | High Schools | | | All Schools | | |
|--|--------------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|
| | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar |
| Nutrients with RDA-based standards | | | | | | | | | | | | |
| Calories | 721* | 721* | 689 | 731 | 722 | 708 | 719 | 728 | 741 | 722* | 724** | 699 |
| Vitamin A | 490 | 489* | 426 | 456** | 447** | 372 | 419* | 422** | 362 | 457* | 456** | 409 |
| Vitamin C | 38** | 34** | 26 | 34** | 31 | 27 | 30 | 30 | 29 | 34** | 32** | 27 |
| Protein | 29 | 29 | 29 | 30 | 30 | 29 | 30 | 31 | 31 | 30 | 30 | 29 |
| Iron | 4.5 | 4.5 | 4.4 | 4.7 | 4.6 | 4.6 | 4.7 | 4.7 | 4.8 | 4.6* | 4.6** | 4.5 |
| Calcium | 473 | 473 | 478 | 477 | 478 | 470 | 475 | 482 | 476 | 475 | 456 | 477 |
| Nutrients with NSLP Standards Based on Dietary Guidelines | | | | | | | | | | | | |
| ≤30 percent calories from fat | 34 | 34 | 33 | 34 | 34 | 34 | 34 | 34 | 35 | 34 | 34 | 33 |
| <10 percent calories saturated fat | 12.0 | 11.8 | 11.9 | 11.9 | 12.0 | 12.1 | 11.8* | 11.9* | 12.4 | 11.9 | 11.9 | 12.0 |
| Nutrients Based on NRC Recommendations | | | | | | | | | | | | |
| >55 percent calories carbohydrate | 52 | 51 | 51 | 51 | 50 | 50 | 50 | 50 | 49 | 51 | 51 | 51 |
| Sodium <800 mg | 1335* | 1308* | 1244 | 1410* | 1378 | 1334 | 1412 | 1405 | 1429 | 1379* | 1357* | 1283 |
| Cholesterol ≤100 mg | 61 | 63 | 65 | 65 | 66 | 67 | 67 | 68 | 70 | 64 | 65 | 66 |
| Sample Size: Number of Schools | | | | | | | | | | | | |
| <i>Unweighted</i> | 38 | 54 | 331 | 61 | 86 | 243 | 86 | 117 | 211 | 185 | 257 | 785 |
| <i>Weighted</i> | 4,719 | 6,424 | 39,555 | 2,452 | 3,223 | 9,119 | 4,082 | 5,268 | 7,589 | 11,254 | 14,914 | 5,263 |

*Difference in those schools with and without salad bars is statistically significant at the .05 level

**Difference in those schools with and without salad bars is statistically significant at the .01 level

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

BEST COPY AVAILABLE

Table B2
Exploratory Analysis: Percentage Public Schools Meeting Nutrition Standards
For Lunch by Salad Bar Availability, SY 1998-99

| | Elementary Schools | | | Middle Schools | | | High Schools | | | All Schools | | |
|--|--------------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|-----------------|----------------------------------|--------------|
| | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar | Daily Salad Bar | Salad Bar at Least Once Per Week | No Salad Bar |
| Nutrients with RDA-based standards | | | | | | | | | | | | |
| Calories | 61 | 66 | 65 | 28 | 23 | 20 | 11 | 13 | 17 | 36** | 38** | 51 |
| Vitamin A | 100 | 100 | 98 | 64 | 68 | 59 | 77** | 78** | 56 | 84 | 85 | 86 |
| Vitamin C | 89 | 89 | 85 | 94* | 91* | 81 | 81* | 81* | 69 | 87 | 87 | 82 |
| Protein | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Iron | 78** | 82* | 93 | 58 | 54 | 53 | 62 | 65 | 60 | 68** | 70** | 82 |
| Calcium | 100 | 100 | 100 | 90 | 88 | 86 | 80 | 84 | 85 | 91** | 92* | 96 |
| Nutrients with NSLP Standards Based on Dietary Guidelines | | | | | | | | | | | | |
| ≤30 percent calories from fat | 23 | 23 | 22 | 14 | 15 | 14 | 15 | 12 | 15 | 18 | 17 | 20 |
| <10 percent calories saturated fat | 27* | 25* | 14 | 19 | 18 | 11 | 22* | 18 | 11 | 23** | 21* | 13 |
| Nutrients Based on NRC Recommendations | | | | | | | | | | | | |
| >55 percent calories carbohydrate | 73 | 75 | 83 | 84 | 85 | 86 | 90 | 90 | 84 | 82 | 82 | 84 |
| Sodium <800 mg | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | <0.5 | 0 | <0.5 | 1 |
| Cholesterol ≤100 mg | 98 | 99 | 99 | 96 | 97 | 95 | 99 | 99 | 96 | 98 | 99 | 98 |
| Sample Size: Number of Schools | | | | | | | | | | | | |
| <i>Unweighted</i> | 38 | 54 | 331 | 61 | 86 | 243 | 86 | 117 | 211 | 185 | 257 | 785 |
| <i>Weighted</i> | 4,719 | 6,424 | 39,555 | 2,452 | 3,223 | 9,119 | 4,082 | 5,268 | 7,589 | 11,254 | 14,914 | 56,263 |

*Difference in those schools with and without salad bars is statistically significant at the .05 level

**Difference in those schools with and without salad bars is statistically significant at the .01 level

Source: Weighted tabulations of menu data from SNDA-II for one week between September 1998 and May 1999.

BEST COPY AVAILABLE

APPENDIX C

METHODOLOGY AND LIMITATIONS

Methodology

SNDA-II data were collected from SFA directors by telephone and from cafeteria managers in sampled schools using a self-administered mail survey. SFA directors provided information on such items as enrollment, number of students approved for free or reduced price, and various menu-planning practices. Cafeteria managers provided information on the number of servings and types of food that were actually served to students during a specified five-day period. Cafeteria managers also provided information on local food service operations such as the availability of a la carte items, types of meal service offered such as salad or other bars, and number of reimbursable meals served each day during the target week, by reimbursement category (free, reduced price or paid). The data collection instruments and a more detailed discussion of the SNDA-II methodology may be found in the original SNDA-II report, Appendix C.

The statistical analyses in this report were conducted by FNS staff using microdata from SNDA-II survey and the statistical software, SAS (SAS Institute, 2002).

Limitations to the Analysis of School Lunch Salad Bars

This report does not explore the number of servings of fruits and vegetables offered to students as part of the National School Lunch Program (NSLP). Program regulations require that all schools participating in the NSLP meet specified nutrition standards for reimbursable meals. Schools generally do so by using a food-based menu planning system that specifies the food items and quantities that must be offered or by using computerized software in planning menus in which milk must be offered along with an entrée and one side dish.⁹ More than two-thirds of all schools used a food-based menu planning system in SY 1998-99. The traditional food-based system requires that students be offered a minimum of two servings of fruit and/or vegetables, and an enhanced food-based system requires larger servings of fruits and vegetables and more servings of bread and grain products.

For most menu items, SNDA-II data provide information on *average meals served to or selected by* students participating in the NSLP. Schools provided information from production records of the types and quantities of foods available for students to select. For the most part, serving sizes specified on the menus were those required by current program regulations. If cafeteria managers were unable to provide information on portion sizes, default values were applied based on these program requirements.

⁹ Although schools using the computerized menu planning option do not have to offer fruits or vegetables as the side dish, they are required to offer menus that meet RDA standards for vitamins A and C---nutrients found most often in fruits and vegetables.

Cafeteria managers provided information on all foods offered on salad bars and other self-serve bars. For bars served on multiple days, managers were asked to provide information only for the first day the bar was offered. Although portions were available for some items (e.g. the average size of a potato used on a potato bar), portions (quantities) were not specified for most self-serve bars.

Because limited data on portions were collected, the nutritional analysis of salad bars was conducted using a set of assumptions about standard salad bar recipes (Table C-1). For example, if a salad bar included lettuce, tomato, broccoli, and salad dressing, the survey did not collect data on how much lettuce, tomato, broccoli, or salad dressing the average student selected or was served. Rather, the nutritional analyses were conducted assuming that the typical salad selected from that salad bar by a secondary school student included 3/4 cup lettuce, 3/8 cup tomato, 3/8 broccoli, and a modest amount of salad dressing.

| Table C-1 | |
|---|---|
| Coding Rules for Fruits and Vegetables in Salad Bars | |
| Elementary Schools | Secondary Schools |
| 3/8 cup vegetable, half of which is lettuce, plus 3/8 cup fruit | 1 cup vegetable, half of which is lettuce, plus 1/2 cup fruit |
| If no fruit, 3/4 cup vegetable, half of which is lettuce | If no fruit, 1 1/2 cup vegetable, half of which is lettuce |

SNDA-II studies what was offered at school lunches, not what students consumed. In addition, the survey collected detailed data on foods served as part of reimbursable meals. Only very limited data on a la carte items is available.

For salad bars, the survey collected detailed information on which individual fruits and vegetables were offered (e.g. peaches), and the number of fruits and vegetables within each category (e.g. canned fruit). These data were not included in the published SNDA-II report, but were obtained via a supplemental data file from Abt Associates. Therefore, we are able to tabulate the percent of salad bars that offered an individual fruit or vegetable.

For non-salad bar fruits and vegetables served as part of the reimbursable meal, the survey collected less detailed information. The nutrient analysis software food coding system only allowed data to be generated as to whether a school offered a fruit or vegetable within a broad category (e.g., canned fruit), not which individual fruits or vegetables were offered (e.g., peaches). Thus, the data allow us to compare whether a certain category of food for (e.g., canned fruit) is more commonly served in schools with salad bars compared to those without salad bars. But we cannot disaggregate those categories to compare the number of fruits or vegetables within a particular category. For example, one can examine whether schools with salad bars are more likely to serve canned fruit than other schools, but one cannot examine whether schools with salad bars serve a larger number of types of canned fruit (e.g., canned peaches and canned pears) than other schools.

When this report analyzes the percent of schools that serve a category of fruits and vegetables, the analysis differs from the published SNDA-II report in an important way. Published tables in the SNDA-II report about the prevalence of categories of fruits and vegetables in schools did not include fruits and vegetables included as part of an entree. For example, tomato sauce topping for pasta would not be included in our analysis of whether tomatoes and/or lettuce are more common in schools with salad bars, even though that tomato sauce would count as at least part of a serving of vegetables in the Food Guide Pyramid.

This report does measure the prevalence of fruits and vegetables in two kinds of entrees: salad bars and chef's salads. The analysis used supplemental data from Abt on the types of fruits and vegetables contained in salad bars. In the interest of treating chef's salads similarly to salad bars, the analysis also assumes that chef's salads contain lettuce and thus are green salads.



U.S. Department of Education
Office of Educational Research and Improvement (OERI)
National Library of Education (NLE)
Educational Resources Information Center (ERIC)



NOTICE

Reproduction Basis

- This document is covered by a signed "Reproduction Release (Blanket)" form (on file within the ERIC system), encompassing all or classes of documents from its source organization and, therefore, does not require a "Specific Document" Release form.
- This document is Federally-funded, or carries its own permission to reproduce, or is otherwise in the public domain and, therefore, may be reproduced by ERIC without a signed Reproduction Release form (either "Specific Document" or "Blanket").