

ED482340 2003-08-00 Depression and Disability in Children and Adolescents. ERIC Digest.

ERIC Development Team

www.eric.ed.gov

Table of Contents

If you're viewing this document online, you can click any of the topics below to link directly to that section.

Depression and Disability in Children and Adolescents. ERIC Digest....	2
MAJOR DEPRESSIVE DISORDER.....	2
DYSTHYMIC DISORDER.....	2
BIPOLAR DISORDER.....	2
OTHER DISABILITIES ASSOCIATED WITH DEPRESSIVE DISORDERS.....	3
THE LINK BETWEEN DEPRESSION AND SUICIDE.....	3
CAUSAL FACTORS RELATED TO DEPRESSION.....	3
DIAGNOSIS AND ASSESSMENT OF DEPRESSIVE ILLNESS IN YOUNG PEOPLE.....	4
TREATMENT OF DEPRESSIVE DISORDERS.....	4
SCHOOL AND CLASSROOM INTERVENTION.....	4
SUMMARY.....	5
RESOURCES.....	5



ERIC Identifier: ED482340

Publication Date: 2003-08-00

Author: Guetzloe, Eleanor

Source: ERIC Clearinghouse on Disabilities and Gifted Education Arlington VA.

Depression and Disability in Children and Adolescents. ERIC Digest.

THIS DIGEST WAS CREATED BY ERIC, THE EDUCATIONAL RESOURCES INFORMATION CENTER. FOR MORE INFORMATION ABOUT ERIC, CONTACT ACCESS ERIC 1-800-LET-ERIC

For many years, depression and other disorders of mood were thought to be afflictions of only adults. Within the past three decades, however, it has become evident that mood disorders are common among children and adolescents. Population studies reveal that between 10% and 15% of the child and adolescent population exhibit some symptoms of depression (U. S. Department of Health and Human Services [USDHHS], 2000).

In children and adolescents, the most frequently diagnosed mood disorders are major depressive disorder, dysthymic disorder, and bipolar disorder. This digest focuses on these three disorders as they are exhibited in childhood and adolescence—their symptoms, causal factors, and treatment.

MAJOR DEPRESSIVE DISORDER

Major depressive disorder is a serious condition characterized by one or more major depressive episodes. In children and adolescents, an episode lasts an average of seven to nine months (Birmaher et al., 1996a, 1996b). Depressed children are sad and lose interest in activities they used to enjoy. They feel unloved, pessimistic, or even hopeless; they think that life is not worth living; and they may think about or threaten suicide. They are often irritable, which may lead to disruptive or aggressive behavior. They may be indecisive, have problems concentrating, and lack energy or motivation. They may neglect appearance and hygiene, and their normal eating and sleeping patterns may be disturbed (USDHHS, 2000).

DYSTHYMIC DISORDER

Dysthymic disorder has fewer symptoms, but is more persistent. The child or adolescent is depressed for most of the day on most days, and symptoms may continue for several years, the average dysthymic period being approximately four years. Seventy percent of children and adolescents with dysthymia eventually experience an episode of major depression. When this combination of major depression and dysthymia occurs, the condition is referred to as double depression (USDHHS, 2000).

BIPOLAR DISORDER

In bipolar disorder, episodes of depression alternate with episodes of mania. The depressive episode usually comes first, with the first manic features becoming evident

months or even years later. Adolescents with mania feel energetic and confident; may have difficulty sleeping but do not tire; and talk a great deal, often speaking very loudly or rapidly. They may complain of racing thoughts. They may do schoolwork quickly and creatively, but in a chaotic, disorganized way. In the manic stage, they may have exaggerated or even delusional ideas about their capabilities and importance, become overconfident, and be uninhibited with others. They may engage in reckless behavior (e.g., fast driving or unsafe sex). Sexual preoccupations are increased and may be associated with promiscuous behavior (USDHHS, 2000).

OTHER DISABILITIES ASSOCIATED WITH DEPRESSIVE DISORDERS

Approximately two-thirds of children and adolescents with major depressive disorder also have another mental disorder, such as anxiety disorder, conduct disorder, oppositional defiant disorder, psychoactive substance abuse or dependence, or phobias (Anderson & McGee, 1994). Authorities have also noted that children with medical problems often face extreme and/or chronic stress, which places them at risk for depression. Estimates of depression among youngsters with medical problems range from 7% in general medical patients to 23% in orthopedic patients (Guetzloe, 1991). Depression has also been linked to a variety of other medical conditions, including endocrinopathies and metabolic disorders (e.g., diabetes and hypoglycemia), viral infections (e.g., influenza, viral hepatitis, and viral pneumonia), rheumatoid arthritis, cancer, central nervous system disorders, metal intoxications, and disabling diseases of all kinds. Some of these conditions may be temporary, but some may be diagnosed as primary disabilities in youngsters with health impairments.

THE LINK BETWEEN DEPRESSION AND SUICIDE

A number of studies have confirmed that children and adolescents with depression are at high risk for suicidal behavior (see Guetzloe, 1991). Because mood disorders substantially increase the risk of suicide, suicidal behavior is a matter of serious concern for parents, educators, and clinicians who deal with the mental health problems of children and adolescents. Over 90% of children and adolescents who commit suicide have a mental disorder (USDHHS, 2000).

CAUSAL FACTORS RELATED TO DEPRESSION

The precise causes of depression are not known. Research on adults with depression generally points to both biological and psychosocial factors, but there has been considerably less research on children and adolescents (Kendler, 1995).

* Family and genetic factors. Between 20% and 50% of depressed children and adolescents have a family history of depression. It is not clear whether the relationship between parent and childhood depression derives from genetic factors or if depressed parents create an environment in which children are more likely to develop mental

disorders (USDHHS, 2000).

* Biological factors. Biochemical and physiological correlates of depression have been studied by medical researchers, with results that generally point to a chemical imbalance in the brain as a causal factor (Birmaher et al., 1996a,1996b). Most of these studies have been conducted with adults, so the findings may not apply to children and adolescents (Guetzloe, 1991).

* Cognitive factors. For several decades there has been considerable interest in the relationship between a pessimistic mindset and a predisposition to depression. Pessimistic individuals generally react more passively, helplessly, and ineffectively to negative events than optimistic individuals. The specific origins of pessimistic mindset have not been established (USDHHS, 2000) but are topics of current research interest (Alloy et al., 2001; Garber & Flynn, 2001).

DIAGNOSIS AND ASSESSMENT OF DEPRESSIVE ILLNESS IN YOUNG PEOPLE

Recent research has focused on the development and validation of checklists and protocols to be used by mental health professionals along with clinical interviews and medical tests. An accurate diagnosis of depression is a complex task, extremely difficult for even highly skilled physicians and other clinicians. It requires a careful examination of physical, mental, emotional, environmental, and cultural factors related to the child or adolescent, his/her family, and the environment. Teachers, counselors, and other school personnel are not expected to diagnose depression in young people; the major roles of educators are to detect the symptoms of depression and make appropriate referrals.

TREATMENT OF DEPRESSIVE DISORDERS

Treatment approaches for children and adolescents include psychosocial interventions (e. g., cognitive behavior therapy) and medication, as well as traditional psychotherapy. Two forms of cognitive therapy (i.e., self-control therapy for prepubertal children and coping skills for adolescents) have been judged as probably effective (Kaslow & Thompson, 1998).

A number of medications are commonly prescribed for children and adolescents with depression, but many of these have not yet been subjected to sufficient study. Effective treatment requires intervention by both medical and mental health professionals, with support from all others who come in contact with the young person; and is therefore not within the purview of the school alone.

SCHOOL AND CLASSROOM INTERVENTION

The educator's most important contribution is the provision of a positive and supportive environment, components of which include satisfaction of basic needs, caring relationships with adults, and physical and psychological security. Any inclusion in a student's program that serves to enhance feelings of self-worth, self-control, and optimism has the potential for ameliorating feelings of depression. Aversive techniques (e. g., punishment and "get tough" approaches) should be avoided to the extent possible (Guetzloe, 1989, 1991).

Educators must use instructional strategies that are both positive and effective so that the student will achieve success and enjoy the learning process. Examples include direct instruction with positive reinforcement, thematic instructional units with varied levels of classroom assignments, learning strategies (e. g., mnemonic devices) and utilization of the principles of universal design for learning, which promote access to the general curriculum for students with learning problems. Some protective factors have been addressed in published curricula (e.g., preventing alienation, enhancing self-esteem, and learning self-control). Other interventions that have implications for school programs (e.g., phototherapy and exercise) have been found to have value in reducing symptoms of depression in adults (Brosse, Sheets, Lett, & Blumenthal, 2002; USDHHS, 2003), but have not yet been subjected to sufficient study with children and adolescents.

SUMMARY

Mood disorders, including major depression, dysthymia, and bipolar disorder, are now recognized as serious problems among children and adolescents. This brief discussion has focused on the symptoms of these disorders, their relationships to other mental and physical problems, their treatment, and appropriate school intervention.

RESOURCES

Alloy, L. B., Abramson, L. Y., Tashman, N., Berrebbi, D. S., Hogan, M. E., Whitehouse, W. G., Crossfield, A. G., & Morocco, A. (2001). Developmental origins of cognitive vulnerability to depression: Parenting, cognitive, and inferential feedback styles of the parents of individuals at high and low cognitive risk for depression. *Cognitive Therapy and Research*, 25, 397-423.

Anderson, J. C., & McGee, R. (1994). Comorbidity of depression in children and adolescents. In W. M. Reynolds & H. F. Johnson (Eds.), *Handbook of depression in children and adolescents* (pp. 581-601). New York: Plenum.

Birmaher, B., Ryan, N. D., Williamson, D. E., Brent, D. A., & Kaufman, J. (1996a). Childhood and adolescent depression: A review of the past 10 years. Part II. *Journal of the American Academy of Child and Adolescent Psychiatry*, 35, 1575-1583.

Birmaher, B., Ryan, N. D., Williamson, D. E., Brent, D. A., Kaufman, J., Dahl, R. E., Perel, J., & Nelson, B. (1996b). Childhood and adolescent depression: A review of the past 10 years. Part I. *Journal of the American Academy of Child and Adolescent*

Psychiatry, 35, 1427-1439.

Brosse, A. L., Sheets, E. S., Lett, H. S., & Blumenthal, J. A. (2002). Exercise and the treatment of clinical depression in adults: Recent findings and future directions. *Sports Medicine* 32(12),741-760.

Garber, A., & Flynn, C. A. (2001). Predictors of depressive cognitions in young adolescents. *Cognitive Therapy and Research*, 25, 353-376.

Guetzloe, E. C. (1991). *Depression and suicide: Special education students at risk*. Reston, VA: Council for Exceptional Children.

Guetzloe, E. C. (1989). *Youth suicide: What the educator should know*. Reston, VA: The Council for Exceptional Children.

Kaslow, N. J., & Thompson, M. P. (1998). Applying the criteria for empirically supported treatments to studies of psychosocial interventions for child and adolescent depression. *Journal of Clinical Child Psychology*, 27, 146-155.

Kendler, K. S. (1995). Genetic epidemiology in psychiatry. Taking both genes and environment seriously. *Archives of General Psychiatry*, 52, 895-899.

U. S. Department of Health and Human Services (USDHHS). (2000). *Mental health: A report of the Surgeon General*. Rockville, MD: U. S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health.

U. S. Department of Health and Human Services (2003). *Mood disorders*. Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, The Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health.
<http://www.mentalhealth.org/publications/allpubs/ken98-0049/default.asp>



Depression & Bipolar Support Alliance (DBSA)



www.DBSAAlliance.org



National Alliance for the Mentally Ill (NAMI)



<http://www.nami.org>



National Foundation for Depressive Illness, Inc.



<http://www.depression.org>



National Institute of Mental Health



<http://www.nimh.nih.gov>



National Mental Health Association (NMHA)



<http://www.nmha.org>

ERIC Digests are in the public domain and may be freely reproduced and disseminated, but please acknowledge your source. This digest was prepared with funding from the Institute of Education Sciences (IES), U.S. Department of Education, under Contract No. ED-99-C0-0026. The opinions expressed in this publication do not necessarily reflect the positions of IES or the Department of Education.

Title: Depression and Disability in Children and Adolescents. ERIC Digest.

Note: Digest number E648.

Document Type: Information Analyses---ERIC Information Analysis Products (IAPs) (071); Information Analyses---ERIC Digests (Selected) in Full Text (073);

Available From: ERIC Clearinghouse on Disabilities and Gifted Education, 1110 North Glebe Rd., Arlington, VA 22201-5704. Tel: 703-620-3660; Tel: 800-328-0272 (Toll Free); For full text: <http://www.ericcec.org>.

Descriptors: Adolescents, Biological Influences, Classroom Techniques, Clinical

Diagnosis, Cognitive Restructuring, Counseling Techniques, Depression (Psychology), Disability Identification, Elementary Secondary Education, Environmental Influences, Etiology, Genetics, Inclusive Schools, Symptoms (Individual Disorders), Teaching Methods

Identifiers: Bipolar Traits, ERIC Digests