

## Cost-Savings of School-Based Health Centers

1. Adams EK, Johnson V. An elementary SBHC: can it reduce Medicaid costs? *Pediatrics* 2000 Apr;105(4 Pt 1):780-8.
  2. Guo JJ, Jan R, Keller KN, McCracken AL, Pan W, Cluxton RJ. Impact of SBHCs on Children with Asthma. *J Adolesc Health*. 2005; 37(4): 266-274.
  3. Kaplan DW, Calonge BN, Guernsey BP, Hanrahan MB. Managed care and SBHCs. Use of health services. *Arch Pediatr Adolesc Med*. 1998 Jan;152(1):25-33.
  4. Key JD, Washington EC, Hulseley TC. Reduced emergency department utilization associated with SBHC enrollment. *J Adolesc Health*. 2002; 30:273-2.
  5. Santelli J, Kouzis A, Newcomer S. SBHCs and adolescent use of primary care and hospital care. *J Adolesc Health*. 1996; 19(4): 267-275.
  6. Tai, T, and Bame, SI. Cost-benefit analysis of childhood asthma management through school-based clinic programs. *J Community Health*. 2010; DOI 10.1007/s10900-010-9305-y
  7. Webber MP, Carpiniello KE, Oruwariye T, Yungtai L, Burton WB, Appel DK. Burden of asthma in elementary school children: Do SBHCs make a difference? *Arch Pediatr Adolesc Med*. 2003; 157: 125-129.
  8. Wade TJ and Guo JJ. Linking improvements in health-related quality of life to reductions in Medicaid costs among students who use school-based health centers. *AJPH*. 2010; 9 (100): 1611-1616.
- Research and evaluations have demonstrated that school-based health centers represent cost-effective investments of public resources:
- A study by Johns Hopkins University found that school-based health centers **reduced inappropriate emergency room use**, increased use of primary care, and resulted in **fewer hospitalizations** among regular users. (5)
  - A study of Medicaid-enrolled children served by a SBHC in Atlanta, Georgia found **significantly lower inpatient, nonemergency department transportation, drug, and emergency department Medicaid expenses** as compared to children without a SBHC. In 1996 the total yearly expense per individual for the SBHC was \$898.98, as compared to \$2360.46 for individuals without a SBHC. (1)
  - The number of **hospitalizations and emergency department visits decreased for children with SBHCs** in Cincinnati schools (2.4-fold and 33.5% respectively) – with an estimated savings of nearly \$1,000 per child. (2)
  - Students in New York City schools with SBHCs were **less likely to have been hospitalized for asthma** at least once in the past year (10.5%) compared to those in schools without SBHCs (17.1%). (7)
  - In South Carolina, prevention-oriented health care provided in a SBHC **decreased emergency department visit rates** by 41% to 57% -- 18% greater than the decrease in students who did not use the SBHC. (4)
  - Adolescents with access to SBHCs in Denver had **38% - 55% fewer after-hours care (emergent or urgent) visits** than those without school-based health center access. (3)
  - Students in Ohio who used an SBHC reported more positive self-perceptions of their health, which correlated with **lower Medicaid costs**. SBHC patients cost Medicaid an average of \$30.40 less than comparable, non-SBHC patients. (8)
  - A study that explored the cost-benefit of a nationwide SBHC program to manage childhood asthma estimated total savings for opportunity costs of work loss and premature death at \$23.13 billion. (6)

February, 2011