

monitoring the future occasional paper series

Paper 97

***DEMOGRAPHIC SUBGROUP TRENDS AMONG ADOLESCENTS
IN THE USE OF VARIOUS LICIT AND ILLICIT DRUGS
1975–2021***

Lloyd D. Johnston
Richard A. Miech
Patrick M. O'Malley
Jerald G. Bachman
John E. Schulenberg
Megan E. Patrick

Institute for Social Research
The University of Michigan
Ann Arbor, Michigan

**Demographic Subgroup Trends among Adolescents
in the Use of Various Licit and Illicit Drugs
1975–2021**

Monitoring the Future Occasional Paper 97

Lloyd D. Johnston, Ph.D.
Richard A. Miech, Ph.D.
Patrick M. O'Malley, Ph.D.
Jerald G. Bachman, Ph.D.
John E. Schulenberg, Ph.D.
Megan E. Patrick, Ph.D.

The University of Michigan
Institute for Social Research
Ann Arbor, MI

The Monitoring the Future project is supported by Grant Number R01DA001411 from the National Institute on Drug Abuse. The content of this occasional paper is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Drug Abuse or the National Institutes of Health.

©Published 2022

Public Domain Notice

All materials appearing in this volume are in the public domain and may be reproduced or copied, whether in print or in non-print media including derivatives, in any reasonable manner, without permission from the authors. To modify the material, please contact the Monitoring the Future Project at mtfinformation@umich.edu for verification of accuracy. Citation of the source is appreciated, including at least the following: Monitoring the Future, Institute for Social Research, University of Michigan.



Recommended Citation

Johnston, L. D., Miech, R. A., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Patrick, M. E. (2022). *Demographic subgroup trends among adolescents in the use of various licit and illicit drugs, 1975–2021* (Monitoring the Future Occasional Paper No. 97). Ann Arbor, MI: Institute for Social Research, The University of Michigan.

CONTENTS

Demographic Subgroup Trends among Adolescents in the Use of Various Licit and Illicit Drugs, 1975-2021

INTRODUCTION	i
LIST OF FIGURES	ii
LIST OF SUPPLEMENTAL TABLES	xv
FIGURES	1
DEFINITION OF BACKGROUND AND DEMOGRAPHIC SUBGROUPS [This information also appears as Appendix B of <i>Monitoring the Future National Survey Results on Drug Use, 1975–2021. Volume I: Secondary School Students</i>].....	469
TRENDS BY SUBGROUP: SUPPLEMENTAL TABLES FOR SECONDARY SCHOOL STUDENTS [These tables complement Appendix D of <i>Monitoring the Future National Survey Results on Drug Use, 1975-2021. Volume I: Secondary School Students</i>]	473
TABLES	474

INTRODUCTION

This occasional paper presents national demographic subgroup data for the 1975-2021 Monitoring the Future (MTF) national survey results on 8th, 10th, and 12th graders' use of drugs, alcohol, and tobacco. MTF is funded by the National Institute on Drug Abuse at the National Institutes of Health under a series of investigator-initiated, competitive research grants to the University of Michigan. The study covers all major classes of illicit and licit psychoactive drugs for an array of population subgroups. The 2020 subgroup data presented here accompany the [Monitoring the Future National Survey Results on Drug Use: 1975-2021: Overview, Key Findings on Adolescent Drug Use](#)¹ and the [Monitoring the Future National Survey Results on Drug Use, 1975-2021, forthcoming: Volume I, Secondary School Students](#).² Prior to 2014 subgroup data were available in tabular form only in Appendices B and D of the relevant year's *Volume I*.³ Since 2014, the MTF subgroup definitions and data have been presented in this series of occasional papers, in both tables and figures to facilitate the examination and interpretation of trend data.

The Overview of Key Findings presents trends in prevalence, perceived risk, disapproval, and perceived availability for most drugs under study and a brief description of subgroup differences. *Volume I* contains a description of MTF's design and purposes, as well as extended reporting on substance use of all kinds—licit and illicit—and a number of related factors, such as attitudes and beliefs about drugs, age of initiation, noncontinuation of use, perceived availability, relevant conditions in the social environment, history of daily marijuana use, use of drugs for the treatment of ADHD, and sources of prescription drugs used outside of medical supervision.

The trends offered here in tabular and graphic forms cover demographic subgroups based on:

- Gender
- College plans
- Region of the country
- Population density
- Education level of the parents (a proxy for socioeconomic level)
- Racial/ethnic identification

Detailed descriptions of the demographic categories are provided in the section starting on page 469 of this paper.

The graphs and tables in this occasional paper present trend data for 8th-, 10th-, and 12th-grade respondents separately. Data for 12th grade begins with 1975, the first year in which a nationally representative sample of high school seniors was surveyed. Data for 8th and 10th grades begin with 1991, when the study's nationally representative annual surveys were expanded to include surveys of those lower grade levels.

¹Johnston, L. D., Miech, R. A., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Patrick, M. E. (2022). *Monitoring the Future national survey results on drug use, 1975-2021: Overview, key findings on adolescent drug use*. Ann Arbor: Institute for Social Research, The University of Michigan.

²Miech, R. A., Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Patrick, M. E. (2022, forthcoming). *Monitoring the Future national survey results on drug use, 1975-2021: Volume I, Secondary school students*. Ann Arbor: Institute for Social Research, The University of Michigan.

³Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Miech, R. A. (2016 and preceding years). *Monitoring the Future national survey results on drug use, 1975-2015: Volume I, Secondary school students*. Ann Arbor: Institute for Social Research, The University of Michigan.

**LIST OF FIGURES GIVING
DEMOGRAPHIC SUBGROUP TRENDS**

NOTE: The lists of figures and tables provided below contain clickable links to all graphics and tables in the paper.

Any Illicit Drug: Trends in Annual Prevalence

- [By Gender \(Figure 1\)](#)..... 1
- [By College Plans \(Figure 2\)](#)..... 2
- [By Region \(Figure 3\)](#)..... 3
- [By Population Density \(Figure 4\)](#)..... 4
- [By Average Education Level of Parents \(Figure 5\)](#)..... 5
- [By Race/Ethnicity \(Figure 6\)](#)..... 6

Any Illicit Drug other than Marijuana: Trends in Annual Prevalence

- [By Gender \(Figure 7\)](#)..... 7
- [By College Plans \(Figure 8\)](#)..... 8
- [By Region \(Figure 9\)](#)..... 9
- [By Population Density \(Figure 10\)](#)..... 10
- [By Average Education Level of Parents \(Figure 11\)](#)..... 11
- [By Race/Ethnicity \(Figure 12\)](#)..... 12

Marijuana: Trends in Lifetime Prevalence

- [By Gender \(Figure 13\)](#)..... 13
- [By College Plans \(Figure 14\)](#)..... 14
- [By Region \(Figure 15\)](#)..... 15
- [By Population Density \(Figure 16\)](#)..... 16
- [By Average Education Level of Parents \(Figure 17\)](#)..... 17
- [By Race/Ethnicity \(Figure 18\)](#)..... 18

Marijuana: Trends in Annual Prevalence

- [By Gender \(Figure 19\)](#)..... 19
- [By College Plans \(Figure 20\)](#)..... 20
- [By Region \(Figure 21\)](#)..... 21
- [By Population Density \(Figure 22\)](#)..... 22
- [By Average Education Level of Parents \(Figure 23\)](#)..... 23
- [By Race/Ethnicity \(Figure 24\)](#)..... 24

Marijuana: Trends in 30-Day Prevalence

- [By Gender \(Figure 25\)](#)..... 25
- [By College Plans \(Figure 26\)](#)..... 26
- [By Region \(Figure 27\)](#)..... 27
- [By Population Density \(Figure 28\)](#)..... 28
- [By Average Education Level of Parents \(Figure 29\)](#)..... 29
- [By Race/Ethnicity \(Figure 30\)](#)..... 30

LIST OF FIGURES (continued)

Marijuana: Trends in 30-Day Prevalence of Daily Use	
By Gender (Figure 31).....	31
By College Plans (Figure 32).....	32
By Region (Figure 33).....	33
By Population Density (Figure 34).....	34
By Average Education Level of Parents (Figure 35).....	35
By Race/Ethnicity (Figure 36).....	36
Synthetic Marijuana: Trends in Annual Prevalence	
By Gender (Figure 37).....	37
By College Plans (Figure 38).....	38
By Region (Figure 39).....	39
By Population Density (Figure 40).....	40
By Average Education Level of Parents (Figure 41).....	41
By Race/Ethnicity (Figure 42).....	42
Inhalants: Trends in Annual Prevalence	
By Gender (Figure 43).....	43
By College Plans (Figure 44).....	44
By Region (Figure 45).....	45
By Population Density (Figure 46).....	46
By Average Education Level of Parents (Figure 47).....	47
By Race/Ethnicity (Figure 48).....	48
Hallucinogens: Trends in Annual Prevalence	
By Gender (Figure 49).....	49
By College Plans (Figure 50).....	50
By Region (Figure 51).....	51
By Population Density (Figure 52).....	52
By Average Education Level of Parents (Figure 53).....	53
By Race/Ethnicity (Figure 54).....	54
LSD: Trends in Annual Prevalence	
By Gender (Figure 55).....	55
By College Plans (Figure 56).....	56
By Region (Figure 57).....	57
By Population Density (Figure 58).....	58
By Average Education Level of Parents (Figure 59).....	59
By Race/Ethnicity (Figure 60).....	60
Hallucinogens other than LSD: Trends in Annual Prevalence	
By Gender (Figure 61).....	61
By College Plans (Figure 62).....	62
By Region (Figure 63).....	63
By Population Density (Figure 64).....	64
By Average Education Level of Parents (Figure 65).....	65
By Race/Ethnicity (Figure 66).....	66

LIST OF FIGURES (continued)

MDMA (Ecstasy, Molly): Trends in Annual Prevalence	
By Gender (Figure 67).....	67
By College Plans (Figure 68).....	68
By Region (Figure 69).....	69
By Population Density (Figure 70).....	70
By Average Education Level of Parents (Figure 71).....	71
By Race/Ethnicity (Figure 72).....	72
Salvia: Trends in Annual Prevalence	
By Gender (Figure 73).....	73
By College Plans (Figure 74).....	74
By Region (Figure 75).....	75
By Population Density (Figure 76).....	76
By Average Education Level of Parents (Figure 77).....	77
By Race/Ethnicity (Figure 78).....	78
Cocaine: Trends in Annual Prevalence	
By Gender (Figure 79).....	79
By College Plans (Figure 80).....	80
By Region (Figure 81).....	81
By Population Density (Figure 82).....	82
By Average Education Level of Parents (Figure 83).....	83
By Race/Ethnicity (Figure 84).....	84
Crack: Trends in Annual Prevalence	
By Gender (Figure 85).....	85
By College Plans (Figure 86).....	86
By Region (Figure 87).....	87
By Population Density (Figure 88).....	88
By Average Education Level of Parents (Figure 89).....	89
By Race/Ethnicity (Figure 90).....	90
Cocaine other than Crack: Trends in Annual Prevalence	
By Gender (Figure 91).....	91
By College Plans (Figure 92).....	92
By Region (Figure 93).....	93
By Population Density (Figure 94).....	94
By Average Education Level of Parents (Figure 95).....	95
By Race/Ethnicity (Figure 96).....	96
Heroin: Trends in Annual Prevalence	
By Gender (Figure 97).....	97
By College Plans (Figure 98).....	98
By Region (Figure 99).....	99
By Population Density (Figure 100).....	100
By Average Education Level of Parents (Figure 101).....	101
By Race/Ethnicity (Figure 102).....	102

LIST OF FIGURES (continued)

Heroin with a Needle: Trends in Annual Prevalence	
By Gender (Figure 103).....	103
By College Plans (Figure 104).....	104
By Region (Figure 105).....	105
By Population Density (Figure 106).....	106
By Average Education Level of Parents (Figure 107).....	107
By Race/Ethnicity (Figure 108).....	108
Heroin without a Needle: Trends in Annual Prevalence	
By Gender (Figure 109).....	109
By College Plans (Figure 110).....	110
By Region (Figure 111).....	111
By Population Density (Figure 112).....	112
By Average Education Level of Parents (Figure 113).....	113
By Race/Ethnicity (Figure 114).....	114
Narcotics other than Heroin: Trends in Annual Prevalence	
By Gender (Figure 115).....	115
By College Plans (Figure 116).....	116
By Region (Figure 117).....	117
By Population Density (Figure 118).....	118
By Average Education Level of Parents (Figure 119).....	119
By Race/Ethnicity (Figure 120).....	120
OxyContin: Trends in Annual Prevalence	
By Gender (Figure 121).....	121
By College Plans (Figure 122).....	122
By Region (Figure 123).....	123
By Population Density (Figure 124).....	124
By Average Education Level of Parents (Figure 125).....	125
By Race/Ethnicity (Figure 126).....	126
Vicodin: Trends in Annual Prevalence	
By Gender (Figure 127).....	127
By College Plans (Figure 128).....	128
By Region (Figure 129).....	129
By Population Density (Figure 130).....	130
By Average Education Level of Parents (Figure 131).....	131
By Race/Ethnicity (Figure 132).....	132
Amphetamines: Trends in Annual Prevalence	
By Gender (Figure 133).....	133
By College Plans (Figure 134).....	134
By Region (Figure 135).....	135
By Population Density (Figure 136).....	136
By Average Education Level of Parents (Figure 137).....	137
By Race/Ethnicity (Figure 138).....	138

LIST OF FIGURES (continued)

Ritalin: Trends in Annual Prevalence	
By Gender (Figure 139).....	139
By College Plans (Figure 140).....	140
By Region (Figure 141).....	141
By Population Density (Figure 142).....	142
By Average Education Level of Parents (Figure 143).....	143
By Race/Ethnicity (Figure 144).....	144
Adderall: Trends in Annual Prevalence	
By Gender (Figure 145).....	145
By College Plans (Figure 146).....	146
By Region (Figure 147).....	147
By Population Density (Figure 148).....	148
By Average Education Level of Parents (Figure 149).....	149
By Race/Ethnicity (Figure 150).....	150
Provigil: Trends in Annual Prevalence	
By Gender (Figure 151).....	151
By College Plans (Figure 152).....	152
By Region (Figure 153).....	153
By Population Density (Figure 154).....	154
By Average Education Level of Parents (Figure 155).....	155
By Race/Ethnicity (Figure 156).....	156
Methamphetamine: Trends in Annual Prevalence	
By Gender (Figure 157).....	157
By College Plans (Figure 158).....	158
By Region (Figure 159).....	159
By Population Density (Figure 160).....	160
By Average Education Level of Parents (Figure 161).....	161
By Race/Ethnicity (Figure 162).....	162
Crystal Methamphetamine (Ice): Trends in Annual Prevalence	
By Gender (Figure 163).....	163
By College Plans (Figure 164).....	164
By Region (Figure 165).....	165
By Population Density (Figure 166).....	166
By Average Education Level of Parents (Figure 167).....	167
By Race/Ethnicity (Figure 168).....	168
Bath Salts (Synthetic Stimulants): Trends in Annual Prevalence	
By Gender (Figure 169).....	169
By College Plans (Figure 170).....	170
By Region (Figure 171).....	171
By Population Density (Figure 172).....	172
By Average Education Level of Parents (Figure 173).....	173
By Race/Ethnicity (Figure 174).....	174

LIST OF FIGURES (continued)

Sedatives (Barbiturates): Trends in Annual Prevalence	
By Gender (Figure 175)	175
By College Plans (Figure 176).....	176
By Region (Figure 177)	177
By Population Density (Figure 178)	178
By Average Education Level of Parents (Figure 179).....	179
By Race/Ethnicity (Figure 180)	180
Tranquilizers: Trends in Annual Prevalence	
By Gender (Figure 181)	181
By College Plans (Figure 182).....	182
By Region (Figure 183)	183
By Population Density (Figure 184)	184
By Average Education Level of Parents (Figure 185).....	185
By Race/Ethnicity (Figure 186)	186
Any Prescription Drug: Trends in Annual Prevalence	
By Gender (Figure 187)	187
By College Plans (Figure 188).....	188
By Region (Figure 189)	189
By Population Density (Figure 190)	190
By Average Education Level of Parents (Figure 191).....	191
By Race/Ethnicity (Figure 192)	192
Over-the-Counter Cough/Cold Medicines: Trends in Annual Prevalence	
By Gender (Figure 193)	193
By College Plans (Figure 194).....	194
By Region (Figure 195)	195
By Population Density (Figure 196)	196
By Average Education Level of Parents (Figure 197).....	197
By Race/Ethnicity (Figure 198)	198
Rohypnol: Trends in Annual Prevalence	
By Gender (Figure 199)	199
By College Plans (Figure 200).....	200
By Region (Figure 201)	201
By Population Density (Figure 202)	202
By Average Education Level of Parents (Figure 203).....	203
By Race/Ethnicity (Figure 204)	204
Alcohol: Trends in 30-Day Prevalence	
By Gender (Figure 205)	205
By College Plans (Figure 206).....	206
By Region (Figure 207)	207
By Population Density (Figure 208)	208
By Average Education Level of Parents (Figure 209).....	209
By Race/Ethnicity (Figure 210)	210

LIST OF FIGURES (continued)

Alcohol: Trends in Daily Prevalence	
By Gender (Figure 211)	211
By College Plans (Figure 212).....	212
By Region (Figure 213)	213
By Population Density (Figure 214)	214
By Average Education Level of Parents (Figure 215).....	215
By Race/Ethnicity (Figure 216)	216
Alcohol: Trends in 30-Day Prevalence of Having Been Drunk	
By Gender (Figure 217)	217
By College Plans (Figure 218).....	218
By Region (Figure 219)	219
By Population Density (Figure 220)	220
By Average Education Level of Parents (Figure 221).....	221
By Race/Ethnicity (Figure 222)	222
Alcohol: Trends in 2-Week Prevalence of 5 or More Drinks in a Row	
By Gender (Figure 223)	223
By College Plans (Figure 224).....	224
By Region (Figure 225)	225
By Population Density (Figure 226)	226
By Average Education Level of Parents (Figure 227).....	227
By Race/Ethnicity (Figure 228)	228
Alcohol: Trends in 2-Week Prevalence of 10 or More Drinks in a Row	
By Gender (Figure 229)	229
By College Plans (Figure 230).....	230
By Region (Figure 231)	231
By Population Density (Figure 232)	232
By Average Education Level of Parents (Figure 233).....	233
By Race/Ethnicity (Figure 234)	234
Alcohol: Trends in 2-Week Prevalence of 15 or More Drinks in a Row	
By Gender (Figure 235)	235
By College Plans (Figure 236).....	236
By Region (Figure 237)	237
By Population Density (Figure 238)	238
By Average Education Level of Parents (Figure 239).....	239
By Race/Ethnicity (Figure 240)	240
Beer: Trends in 30-Day Prevalence	
By Gender (Figure 241)	241
By College Plans (Figure 242).....	242
By Region (Figure 243)	243
By Population Density (Figure 244)	244
By Average Education Level of Parents (Figure 245).....	245
By Race/Ethnicity (Figure 246)	246

LIST OF FIGURES (continued)

Beer: Trends in 2-Week Prevalence of 5 or More Drinks in a Row	
By Gender (Figure 247)	247
By College Plans (Figure 248).....	248
By Region (Figure 249)	249
By Population Density (Figure 250)	250
By Average Education Level of Parents (Figure 251)	251
By Race/Ethnicity (Figure 252)	252
Liquor: Trends in 30-Day Prevalence	
By Gender (Figure 253)	253
By College Plans (Figure 254).....	254
By Region (Figure 255)	255
By Population Density (Figure 256)	256
By Average Education Level of Parents (Figure 257)	257
By Race/Ethnicity (Figure 258)	258
Liquor: Trends in 2-Week Prevalence of 5 or More Drinks in a Row	
By Gender (Figure 259)	259
By College Plans (Figure 260).....	260
By Region (Figure 261)	261
By Population Density (Figure 262)	262
By Average Education Level of Parents (Figure 263)	263
By Race/Ethnicity (Figure 264)	264
Wine: Trends in 30-Day Prevalence	
By Gender (Figure 265)	265
By College Plans (Figure 266).....	266
By Region (Figure 267)	267
By Population Density (Figure 268)	268
By Average Education Level of Parents (Figure 269)	269
By Race/Ethnicity (Figure 270)	270
Wine: Trends in 2-Week Prevalence of 5 or More Drinks in a Row	
By Gender (Figure 271)	271
By College Plans (Figure 272).....	272
By Region (Figure 273)	273
By Population Density (Figure 274)	274
By Average Education Level of Parents (Figure 275)	275
By Race/Ethnicity (Figure 276)	276
Wine Coolers: Trends in 30-Day Prevalence	
By Gender (Figure 277)	277
By College Plans (Figure 278).....	278
By Region (Figure 279)	279
By Population Density (Figure 280)	280
By Average Education Level of Parents (Figure 281)	281
By Race/Ethnicity (Figure 282)	282

LIST OF FIGURES (continued)

Wine Coolers: Trends in 2-Week Prevalence of 5 or More Drinks in a Row	
By Gender (Figure 283)	283
By College Plans (Figure 284)	284
By Region (Figure 285)	285
By Population Density (Figure 286)	286
By Average Education Level of Parents (Figure 287)	287
By Race/Ethnicity (Figure 288)	288
Flavored Alcoholic Beverages: Trends in 30-Day Prevalence	
By Gender (Figure 289)	289
By College Plans (Figure 290)	290
By Region (Figure 291)	291
By Population Density (Figure 292)	292
By Average Education Level of Parents (Figure 293)	293
By Race/Ethnicity (Figure 294)	294
Alcoholic Beverages containing Caffeine: Trends in Annual Prevalence	
By Gender (Figure 295)	295
By College Plans (Figure 296)	296
By Region (Figure 297)	297
By Population Density (Figure 298)	298
By Average Education Level of Parents (Figure 299)	299
By Race/Ethnicity (Figure 300)	300
Cigarettes: Trends in 30-Day Prevalence	
By Gender (Figure 301)	301
By College Plans (Figure 302)	302
By Region (Figure 303)	303
By Population Density (Figure 304)	304
By Average Education Level of Parents (Figure 305)	305
By Race/Ethnicity (Figure 306)	306
Cigarettes: Trends in 30-Day Prevalence of Daily Use	
By Gender (Figure 307)	307
By College Plans (Figure 308)	308
By Region (Figure 309)	309
By Population Density (Figure 310)	310
By Average Education Level of Parents (Figure 311)	311
By Race/Ethnicity (Figure 312)	312
Cigarettes: Trends in 30-Day Prevalence of Use of Half Pack a Day or More	
By Gender (Figure 313)	313
By College Plans (Figure 314)	314
By Region (Figure 315)	315
By Population Density (Figure 316)	316
By Average Education Level of Parents (Figure 317)	317
By Race/Ethnicity (Figure 318)	318

LIST OF FIGURES (continued)

Tobacco using a Hookah: Trends in Annual Prevalence	
By Gender (Figure 319)	319
By College Plans (Figure 320)	320
By Region (Figure 321)	321
By Population Density (Figure 322)	322
By Average Education Level of Parents (Figure 323)	323
By Race/Ethnicity (Figure 324)	324
Small Cigars: Trends in Annual Prevalence	
By Gender (Figure 325)	325
By College Plans (Figure 326)	326
By Region (Figure 327)	327
By Population Density (Figure 328)	328
By Average Education Level of Parents (Figure 329)	329
By Race/Ethnicity (Figure 330)	330
Flavored Little Cigars: Trends in 30-Day Prevalence	
By Gender (Figure 331)	331
By College Plans (Figure 332)	332
By Region (Figure 333)	333
By Population Density (Figure 334)	334
By Average Education Level of Parents (Figure 335)	335
By Race/Ethnicity (Figure 336)	336
Regular Little Cigars: Trends in 30-Day Prevalence	
By Gender (Figure 337)	337
By College Plans (Figure 338)	338
By Region (Figure 339)	339
By Population Density (Figure 340)	340
By Average Education Level of Parents (Figure 341)	341
By Race/Ethnicity (Figure 342)	342
Large Cigars: Trends in 30-Day Prevalence	
By Gender (Figure 343)	343
By College Plans (Figure 344)	344
By Region (Figure 345)	345
By Population Density (Figure 346)	346
By Average Education Level of Parents (Figure 347)	347
By Race/Ethnicity (Figure 348)	348
Any Vaping: Trends in 30-Day Prevalence	
By Gender (Figure 349)	349
By College Plans (Figure 350)	350
By Region (Figure 351)	351
By Population Density (Figure 352)	352
By Average Education Level of Parents (Figure 353)	353
By Race/Ethnicity (Figure 354)	354

LIST OF FIGURES (continued)

Vaping Nicotine: Trends in 30-Day Prevalence	
By Gender (Figure 355).....	355
By College Plans (Figure 356).....	356
By Region (Figure 357).....	357
By Population Density (Figure 358).....	358
By Average Education Level of Parents (Figure 359).....	359
By Race/Ethnicity (Figure 360).....	360
Vaping Marijuana: Trends in 30-Day Prevalence	
By Gender (Figure 361).....	361
By College Plans (Figure 362).....	362
By Region (Figure 363).....	363
By Population Density (Figure 364).....	364
By Average Education Level of Parents (Figure 365).....	365
By Race/Ethnicity (Figure 366).....	366
Vaping Just Flavoring: Trends in 30-Day Prevalence	
By Gender (Figure 367).....	367
By College Plans (Figure 368).....	368
By Region (Figure 369).....	369
By Population Density (Figure 370).....	370
By Average Education Level of Parents (Figure 371).....	371
By Race/Ethnicity (Figure 372).....	372
Smokeless Tobacco: Trends in 30-Day Prevalence	
By Gender (Figure 373).....	373
By College Plans (Figure 374).....	374
By Region (Figure 375).....	375
By Population Density (Figure 376).....	376
By Average Education Level of Parents (Figure 377).....	377
By Race/Ethnicity (Figure 378).....	378
Smokeless Tobacco: Trends in 30-Day Prevalence of Daily Use	
By Gender (Figure 379).....	379
By College Plans (Figure 380).....	380
By Region (Figure 381).....	381
By Population Density (Figure 382).....	382
By Average Education Level of Parents (Figure 383).....	383
By Race/Ethnicity (Figure 384).....	384
Dissolvable Tobacco Products: Trends in Annual Prevalence	
By Gender (Figure 385).....	385
By College Plans (Figure 386).....	386
By Region (Figure 387).....	387
By Population Density (Figure 388).....	388
By Average Education Level of Parents (Figure 389).....	389
By Race/Ethnicity (Figure 390).....	390

LIST OF FIGURES (continued)

Snus: Trends in Annual Prevalence	
By Gender (Figure 391)	391
By College Plans (Figure 392).....	392
By Region (Figure 393)	393
By Population Density (Figure 394)	394
By Average Education Level of Parents (Figure 395).....	395
By Race/Ethnicity (Figure 396)	396
Steroids: Trends in Annual Prevalence	
By Gender (Figure 397)	397
By College Plans (Figure 398).....	398
By Region (Figure 399)	399
By Population Density (Figure 400)	400
By Average Education Level of Parents (Figure 401).....	401
By Race/Ethnicity (Figure 402)	402
Marijuana: Trends in Lifetime Prevalence of Daily Use for Month or More	
By Gender (Figure 403)	403
By College Plans (Figure 404).....	404
By Region (Figure 405)	405
By Population Density (Figure 406)	406
By Average Education Level of Parents (Figure 407).....	407
By Race/Ethnicity (Figure 408)	408
Nonprescription Diet Pills: Trends in Annual Prevalence	
By Gender (Figure 409)	409
By College Plans (Figure 410).....	410
By Region (Figure 411)	411
By Population Density (Figure 412)	412
By Average Education Level of Parents (Figure 413).....	413
By Race/Ethnicity (Figure 414)	414
Nonprescription Stay-Awake Pills: Trends in Annual Prevalence	
By Gender (Figure 415)	415
By College Plans (Figure 416).....	416
By Region (Figure 417)	417
By Population Density (Figure 418)	418
By Average Education Level of Parents (Figure 419).....	419
By Race/Ethnicity (Figure 420)	420
Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use	
By Gender (Figure 421)	421
By College Plans (Figure 422).....	422
By Region (Figure 423)	423
By Population Density (Figure 424)	424
By Average Education Level of Parents (Figure 425).....	425
By Race/Ethnicity (Figure 426)	426

LIST OF FIGURES (continued)

Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use	
By Gender (Figure 427).....	427
By College Plans (Figure 428).....	428
By Region (Figure 429).....	429
By Population Density (Figure 430).....	430
By Average Education Level of Parents (Figure 431).....	431
By Race/Ethnicity (Figure 432).....	432
Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use	
By Gender (Figure 433).....	433
By College Plans (Figure 434).....	434
By Region (Figure 435).....	435
By Population Density (Figure 436).....	436
By Average Education Level of Parents (Figure 437).....	437
By Race/Ethnicity (Figure 438).....	438
Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use	
By Gender (Figure 439).....	439
By College Plans (Figure 440).....	440
By Region (Figure 441).....	441
By Population Density (Figure 442).....	442
By Average Education Level of Parents (Figure 443).....	443
By Race/Ethnicity (Figure 444).....	444
Either Stimulant or Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use	
By Gender (Figure 445).....	445
By College Plans (Figure 446).....	446
By Region (Figure 447).....	447
By Population Density (Figure 448).....	448
By Average Education Level of Parents (Figure 449).....	449
By Race/Ethnicity (Figure 450).....	450
Either Stimulant or Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use	
By Gender (Figure 451).....	451
By College Plans (Figure 452).....	452
By Region (Figure 453).....	453
By Population Density (Figure 454).....	454
By Average Education Level of Parents (Figure 455).....	455
By Race/Ethnicity (Figure 456).....	456
Any Nicotine Use: Trends in 30-Day Prevalence by Gender	
By Gender (Figure 457).....	457
By College Plans (Figure 458).....	458
By Region (Figure 459).....	459
By Population Density (Figure 460).....	460
By Average Education Level of Parents (Figure 461).....	461
By Race/Ethnicity (Figure 462).....	462

LIST OF FIGURES (continued)

Any Nicotine Use other than Vaping: Trends in 30-Day Prevalence by Gender

By Gender (Figure 463).....	463
By College Plans (Figure 464).....	464
By Region (Figure 465).....	465
By Population Density (Figure 466).....	466
By Average Education Level of Parents (Figure 467).....	467
By Race/Ethnicity (Figure 468).....	468


**LIST OF SUPPLEMENTAL TABLES
DEMOGRAPHIC SUBGROUPS TRENDS**

Tables are listed by drug and prevalence period. For each drug and prevalence period, trend data is displayed separately for grades 8, 10, and 12 unless otherwise noted.

Drugs	Prevalence	
Any Illicit Drug	Annual	476
Any Illicit Drug other than Marijuana	Annual	484
Marijuana	Lifetime	494
	Annual	501
	30-Day	508
	30-Day of Daily Use.....	515
Synthetic Marijuana	Annual	522
Inhalants	Annual	525
Hallucinogens	Annual	532
LSD	Annual	539
Hallucinogens other than LSD	Annual	546
MDMA (Ecstasy, Molly)	Annual	553
Salvia	Annual	559
Cocaine	Annual	562
Crack	Annual	569
Other Cocaine	Annual	576
Heroin	Annual	583
Heroin with a Needle	Annual	590
Heroin without a Needle	Annual	596
Narcotics other than Heroin	Annual, Grade 12 only.....	602
OxyContin	Annual	606
Vicodin	Annual	609
Amphetamines	Annual	612
Ritalin	Annual	622
Adderall	Annual	625
Provigil	Annual, Grade 12 only.....	628
Methamphetamine	Annual	629
Crystal Methamphetamine (Ice)	Annual, Grade 12 only.....	632
Bath Salts (Synthetic Stimulants)	Annual	634
Sedatives (Barbiturates)	Annual, Grade 12 only.....	637
Tranquilizers	Annual	640
Any Prescription Drug without Medical Supervision	Annual, Grade 12 only.....	647
Over-the-Counter Cough/Cold Medicines	Annual	648
Rohypnol	Annual	651
Alcohol	30-Day	657
	30-Day of Daily Use.....	664
Been Drunk	30-Day	671
	2-Week of 5+ Drinks in a Row.....	677
	2-Week of 10+ Drinks in a Row.....	684
Beer	30-Day	687
	2-Week of 5+ Drinks in a Row.....	694
Liquor	30-Day, Grade 12 only	701
	2-Week of 5+ Drinks in a Row, Grade 12 only	704
Wine	30-Day, Grade 12 only	707

	2-Week of 5+ Drinks in a Row, Grade 12 only	710
Wine Coolers	30-Day	713
	2-Week of 5+ Drinks in a Row	718
Flavored Alcoholic Beverages	30-Day	721
Alcoholic Beverages containing Caffeine	Annual	724
Cigarettes	30-Day	727
	30-Day of Daily Use	734
	30-Day of Half Pack a Day	741
Tobacco Using Hookah	Annual, Grade 12 only	748
Small Cigars	Annual, Grade 12 only	749
Flavored Little Cigars or Cigarillos	30-Day	750
Regular Little Cigars or Cigarillos	30-Day	752
Large Cigars	30-Day	754
Any Vaping	30-Day	756
Vaping Nicotine	30-Day	757
Vaping Marijuana	30-Day	758
Vaping Just Flavoring	30-Day	759
Smokeless Tobacco	30-Day	760
	30-Day of Daily Use	767
Dissolvable Tobacco Products	Annual	774
Snus	Annual	777
Steroids	Annual	780
Marijuana	Lifetime	786
Nonprescription Diet Pills	Annual	789
Nonprescription Stay-Awake Pills	Annual	792
Stimulant-Type ADHD Drugs	Lifetime	795
	Current	798
Non-Stimulant-Type ADHD Drugs	Lifetime	801
	Current	804
Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs	Lifetime	807
	Current	810
Any Nicotine Use	30-Day	813
Any Nicotine Use other than Vaping	30-Day	814
Use of Marijuana under a Doctor's Orders	Lifetime	815
Powdered Alcohol	Annual	816
Energy Drinks	Daily	817
Energy Shots	Daily	820
Either Energy Drinks or Energy Shots	Daily	823
Approximate Weighted Ns by Subgroups in Grade 8		826
Approximate Weighted Ns by Subgroups in Grade 10		828
Approximate Weighted Ns by Subgroups in Grade 12		830

FIGURE 1
Any Illicit Drug:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2013, a revised set of questions on amphetamine use were introduced. Data for any illicit drug are affected by these changes.

FIGURE 2
Any Illicit Drug:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2013, a revised set of questions on amphetamine use were introduced. Data for any illicit drug are affected by these changes.

FIGURE 3
Any Illicit Drug:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2013, a revised set of questions on amphetamine use were introduced. Data for any illicit drug are affected by these changes.

FIGURE 4
Any Illicit Drug:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2013, a revised set of questions on amphetamine use were introduced. Data for any illicit drug are affected by these changes.

FIGURE 5
Any Illicit Drug:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2013, a revised set of questions on amphetamine use were introduced. Data for any illicit drug are affected by these changes.

**Parental education is an average score of mother's education and father's education.

FIGURE 6
Any Illicit Drug*: Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2013, a revised set of questions on amphetamine use were introduced. Data for any illicit drug are affected by these changes.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 7
Any Illicit Drug other than Marijuana:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, revised sets of questions on other hallucinogen and tranquilizer use were introduced. Beginning in 2013, a revised set of questions on amphetamines were introduced. Data for any illicit drug other than marijuana are affected by these changes.

FIGURE 8
Any Illicit Drug other than Marijuana:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, revised sets of questions on other hallucinogen and tranquilizer use were introduced. Beginning in 2013, a revised set of questions on amphetamines were introduced. Data for any illicit drug other than marijuana are affected by these changes.

FIGURE 9
Any Illicit Drug other than Marijuana:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, revised sets of questions on other hallucinogen and tranquilizer use were introduced. Beginning in 2013, a revised set of questions on amphetamines were introduced. Data for any illicit drug other than marijuana are affected by these changes.

FIGURE 10
Any Illicit Drug other than Marijuana:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, revised sets of questions on other hallucinogen and tranquilizer use were introduced. Beginning in 2013, a revised set of questions on amphetamines were introduced. Data for any illicit drug other than marijuana are affected by these changes.

FIGURE 11

Any Illicit Drug other than Marijuana:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, revised sets of questions on other hallucinogen and tranquilizer use were introduced. Beginning in 2013, a revised set of questions on amphetamines were introduced. Data for any illicit drug other than marijuana are affected by these changes.

**Parental education is an average score of mother's education and father's education.

FIGURE 12
Any Illicit Drug other than Marijuana:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, revised sets of questions on other hallucinogen and tranquilizer use were introduced. Beginning in 2013, a revised set of questions on amphetamines were introduced. Data for any illicit drug other than marijuana are affected by these changes.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 13
Marijuana: Trends in Lifetime Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 14
Marijuana: Trends in Lifetime Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 15
Marijuana: Trends in Lifetime Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 16
Marijuana: Trends in Lifetime Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 17
Marijuana: Trends in Lifetime Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 18
Marijuana: Trends in Lifetime Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 19
Marijuana: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 20
Marijuana: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 21
Marijuana: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 22
Marijuana: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 23
Marijuana: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 24
Marijuana: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 25
Marijuana: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 26
Marijuana: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 27
Marijuana: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 28
Marijuana: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 29
Marijuana: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 30
Marijuana: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 31
Marijuana: Trends in 30-Day Prevalence of Daily Use by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 32
Marijuana: Trends in 30-Day Prevalence of Daily Use by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 33
Marijuana: Trends in 30-Day Prevalence of Daily Use by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 34
Marijuana: Trends in 30-Day Prevalence of Daily Use by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 35
Marijuana: Trends 30-Day Prevalence of Daily Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 36
Marijuana: Trends in 30-Day Prevalence of Daily Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 37
Synthetic Marijuana: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 38
Synthetic Marijuana: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 39
Synthetic Marijuana: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 40
Synthetic Marijuana: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 41
Synthetic Marijuana: Trends Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 42
Synthetic Marijuana: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 43
Inhalants: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 44
Inhalants: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.


Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 45
Inhalants: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.
 Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 46
Inhalants: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 47
Inhalants: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


*Parental education is an average score of mother's education and father's education.

FIGURE 48
Inhalants: Trends in Annual Prevalence by Race/Ethnicity*



Source. The Monitoring the Future study, the University of Michigan.
 Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.
 *To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 49
Hallucinogens:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. Data for hallucinogens are affected by these changes.

FIGURE 50
Hallucinogens:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. Data for hallucinogens are affected by these changes.

FIGURE 51
Hallucinogens:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. Data for hallucinogens are affected by these changes.

FIGURE 52
Hallucinogens:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. Data for hallucinogens are affected by these changes.

FIGURE 53
Hallucinogens:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. Data for hallucinogens are affected by these changes.

**Parental education is an average score of mother's education and father's education.

FIGURE 54
Hallucinogens:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. Data for hallucinogens are affected by these changes.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 55
LSD: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 56
LSD: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 57
LSD: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 58
LSD: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 59
LSD: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 60
LSD: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 61
Hallucinogens other than LSD:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples.

FIGURE 62
Hallucinogens other than LSD:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples.

FIGURE 63
Hallucinogens other than LSD:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples.

FIGURE 64
Hallucinogens other than LSD:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples.

FIGURE 65
Hallucinogens other than LSD:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

str(*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples.

**Parental education is an average score of mother's education and father's education.

FIGURE 66
Hallucinogens other than LSD:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on other hallucinogen use was introduced. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 67
MDMA (Ecstasy, Molly)*: Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2014, the question was changed in some of the questionnaire forms to include molly in the list of examples. In 2015, it was changed in the remaining forms. Data is presented here for both versions of the question.

FIGURE 68
MDMA (Ecstasy, Molly)*: Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2014, the question was changed in some of the questionnaire forms to include molly in the list of examples. In 2015, it was changed in the remaining forms. Data is presented here for both versions of the question.

FIGURE 69
MDMA (Ecstasy, Molly)*: Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2014, the question was changed in some of the questionnaire forms to include molly in the list of examples. In 2015, it was changed in the remaining forms. Data is presented here for both versions of the question.

FIGURE 70
MDMA (Ecstasy, Molly)*: Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2014, the question was changed in some of the questionnaire forms to include molly in the list of examples. In 2015, it was changed in the remaining forms. Data is presented here for both versions of the question.

FIGURE 71
MDMA (Ecstasy, Molly)*: Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2014, the question was changed in some of the questionnaire forms to include molly in the list of examples. In 2015, it was changed in the remaining forms. Data is presented here for both versions of the question.

**Parental education is an average score of mother's education and father's education.

FIGURE 72
MDMA (Ecstasy, Molly)*: Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2014, the question was changed in some of the questionnaire forms to include molly in the list of examples. In 2015, it was changed in the remaining forms. Data is presented here for both versions of the question.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 73
Salvia: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 74
Salvia: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 75
Salvia: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 76
Salvia: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 77
Salvia: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 78
Salvia: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 79
Cocaine*: Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Cocaine in any form.

FIGURE 80
Cocaine*: Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Cocaine in any form.

FIGURE 81
Cocaine*: Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Cocaine in any form.

FIGURE 82
Cocaine*: Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Cocaine in any form.

FIGURE 83
Cocaine*: Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Cocaine in any form.

**Parental education is an average score of mother's education and father's education.

FIGURE 84
Cocaine*: Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Cocaine in any form.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 85
Crack: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 86
Crack: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 87
Crack: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 88
Crack: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 89
Crack: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 90
Crack: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 91
Cocaine other than Crack*: Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Almost entirely powder cocaine for many years.

FIGURE 92
Cocaine other than Crack*: Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Almost entirely powder cocaine for many years.

FIGURE 93
Cocaine other than Crack*: Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Almost entirely powder cocaine for many years.

FIGURE 94
Cocaine other than Crack*: Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Almost entirely powder cocaine for many years.

FIGURE 95
Cocaine other than Crack*: Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Almost entirely powder cocaine for many years.

**Parental education is an average score of mother's education and father's education.

FIGURE 96
Cocaine other than Crack*: Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Almost entirely powder cocaine for many years.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 97
Heroin: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 98
Heroin: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 99
Heroin: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 100
Heroin: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 101
Heroin: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 102
Heroin: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 103
Heroin with a Needle: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 104
Heroin with a Needle: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 105
Heroin with a Needle: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 106
Heroin with a Needle: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 107
Heroin with a Needle: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 108
Heroin with a Needle: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 109
Heroin without a Needle: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 110
Heroin without a Needle: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 111
Heroin without a Needle: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 112
Heroin without a Needle: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 113
Heroin without a Needle: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 114
Heroin without a Needle: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 115
Narcotics other than Heroin:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2002, a revised set of questions on use of narcotics other than heroin was introduced. Talwin, laudanum, and paregoric were replaced with Vicodin, OxyContin, and Percocet.

FIGURE 116
Narcotics other than Heroin:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2002, a revised set of questions on use of narcotics other than heroin was introduced. Talwin, laudanum, and paregoric were replaced with Vicodin, OxyContin, and Percocet.

FIGURE 117
Narcotics other than Heroin:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2002, a revised set of questions on use of narcotics other than heroin was introduced. Talwin, laudanum, and paregoric were replaced with Vicodin, OxyContin, and Percocet.

FIGURE 118
Narcotics other than Heroin:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2002, a revised set of questions on use of narcotics other than heroin was introduced. Talwin, laudanum, and paregoric were replaced with Vicodin, OxyContin, and Percocet.

FIGURE 119
Narcotics other than Heroin:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2002, a revised set of questions on use of narcotics other than heroin was introduced. Talwin, laudanum, and paregoric were replaced with Vicodin, OxyContin, and Percocet.

**Parental education is an average score of mother's education and father's education.

FIGURE 120
Narcotics other than Heroin:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2002, a revised set of questions on use of narcotics other than heroin was introduced. Talwin, laudanum, and paregoric were replaced with Vicodin, OxyContin, and Percocet.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 121
OxyContin: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 122
OxyContin: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 123
OxyContin: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 124
OxyContin: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 125
OxyContin: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 126
OxyContin: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 127
Vicodin: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 128
Vicodin: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 129
Vicodin: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 130
Vicodin: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 131
Vicodin: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 132
Vicodin: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 133
Amphetamines:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*After 1982, the question about amphetamine use was revised to further clarify that nonprescription stimulants should not be included in the answers.

Beginning in 2013, a revised set of questions on amphetamine use was introduced in which the phrase 'or other prescription stimulant drugs' was added.

FIGURE 134
Amphetamines:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*After 1982, the question about amphetamine use was revised to further clarify that nonprescription stimulants should not be included in the answers.

Beginning in 2013, a revised set of questions on amphetamine use was introduced in which the phrase 'or other prescription stimulant drugs' was added.

FIGURE 135
Amphetamines:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*After 1982, the question about amphetamine use was revised to further clarify that nonprescription stimulants should not be included in the answers.

Beginning in 2013, a revised set of questions on amphetamine use was introduced in which the phrase 'or other prescription stimulant drugs' was added.

FIGURE 136
Amphetamines:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*After 1982, the question about amphetamine use was revised to further clarify that nonprescription stimulants should not be included in the answers.

Beginning in 2013, a revised set of questions on amphetamine use was introduced in which the phrase 'or other prescription stimulant drugs' was added.

FIGURE 137
Amphetamines:* Trends in Annual Prevalence by Average Education Level of Parents**



Source. The Monitoring the Future study, the University of Michigan.


Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*After 1982, the question about amphetamine use was revised to further clarify that nonprescription stimulants should not be included in the answers.

Beginning in 2013, a revised set of questions on amphetamine use was introduced in which the phrase 'or other prescription stimulant drugs' was added.

**Parental education is an average score of mother's education and father's education.

FIGURE 138
Amphetamines:* Trends in Annual Prevalence by Race/Ethnicity**



Source. The Monitoring the Future study, the University of Michigan.


Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*After 1982, the question about amphetamine use was revised to further clarify that nonprescription stimulants should not be included in the answers.

Beginning in 2013, a revised set of questions on amphetamine use was introduced in which the phrase 'or other prescription stimulant drugs' was added.


**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 139
Ritalin: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.


FIGURE 140
Ritalin: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 141
Ritalin: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 142
Ritalin: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 143
Ritalin: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 144
Ritalin: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 145
Adderall: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 146
Adderall: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 147
Adderall: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 148
Adderall: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 149
Adderall: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 150
Adderall: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 151
Provigil:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2012.

FIGURE 152
Provigil:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2012.

FIGURE 153
Provigil:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2012.

FIGURE 154
Provigil:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2012.

FIGURE 155
Provigil:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2012.

**Parental education is an average score of mother's education and father's education.

FIGURE 156
Provigil:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2012.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 157
Methamphetamine: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 158
Methamphetamine: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 159
Methamphetamine: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.


Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 160
Methamphetamine: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.
 Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 161
Methamphetamine: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 162
Methamphetamine: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 163
Crystal Methamphetamine (Ice): Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 164
Crystal Methamphetamine (Ice): Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 165
Crystal Methamphetamine (Ice): Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 166
Crystal Methamphetamine (Ice): Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 167
Crystal Methamphetamine (Ice): Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 168
Crystal Methamphetamine (Ice): Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 169
Bath Salts (Synthetic Stimulants):* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2019.

FIGURE 170
Bath Salts (Synthetic Stimulants): Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2019.

FIGURE 171
Bath Salts (Synthetic Stimulants):* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2019.

FIGURE 172
Bath Salts (Synthetic Stimulants):* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2019.

FIGURE 173

Bath Salts (Synthetic Stimulants): Trends in Annual Prevalence by Average Education Level of Parents***




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

**Question discontinued in 2019.

FIGURE 174
Bath Salts (Synthetic Stimulants):* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued in 2019.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 175
Sedatives (Barbiturates): Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 176
Sedatives (Barbiturates): Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 177
Sedatives (Barbiturates): Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 178
Sedatives (Barbiturates): Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 179
Sedatives (Barbiturates): Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 180
Sedatives (Barbiturates): Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 181
Tranquilizers:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on tranquilizer use was introduced in which Miltown was replaced with Xanax.

FIGURE 182
Tranquilizers:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on tranquilizer use was introduced in which Miltown was replaced with Xanax.

FIGURE 183
Tranquilizers:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on tranquilizer use was introduced in which Miltown was replaced with Xanax.

FIGURE 184
Tranquilizers:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on tranquilizer use was introduced in which Miltown was replaced with Xanax.

FIGURE 185
Tranquilizers:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on tranquilizer use was introduced in which Miltown was replaced with Xanax.

**Parental education is an average score of mother's education and father's education.

FIGURE 186
Tranquilizers:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 2001, a revised set of questions on tranquilizer use was introduced in which Miltown was replaced with Xanax.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 187
Any Prescription Drug:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*The use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers without a doctor telling you to use them. Beginning in 2013, a revised set of questions on amphetamine use was introduced

FIGURE 188
Any Prescription Drug:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*The use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers without a doctor telling you to use them. Beginning in 2013, a revised set of questions on amphetamine use was introduced

FIGURE 189
Any Prescription Drug:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*The use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers without a doctor telling you to use them. Beginning in 2013, a revised set of questions on amphetamine use was introduced

FIGURE 190
Any Prescription Drug:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*The use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers without a doctor telling you to use them. Beginning in 2013, a revised set of questions on amphetamine use was introduced

FIGURE 191
Any Prescription Drug:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*The use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers without a doctor telling you to use them. Beginning in 2013, a revised set of questions on amphetamine use was introduced

**Parental education is an average score of mother's education and father's education.

FIGURE 192
Any Prescription Drug:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*The use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers without a doctor telling you to use them. Beginning in 2013, a revised set of questions on amphetamine use was introduced

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 193
Over-the-Counter Cough/Cold Medicines:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Taken to get high.

FIGURE 194
Over-the-Counter Cough/Cold Medicines:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Taken to get high.

FIGURE 195
Over-the-Counter Cough/Cold Medicines:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Taken to get high.

FIGURE 196
Over-the-Counter Cough/Cold Medicines:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Taken to get high.

FIGURE 197
Over-the-Counter Cough/Cold Medicines:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Taken to get high.

**Parental education is an average score of mother's education and father's education.

FIGURE 198
Over-the-Counter Cough/Cold Medicines:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Taken to get high.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 199
Rohypnol:* Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*For 12th grade only, the 2001 and 2002 data are not comparable due to changes in the questionnaire forms.

FIGURE 200
Rohypnol:* Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*For 12th grade only, the 2001 and 2002 data are not comparable due to changes in the questionnaire forms.

FIGURE 201
Rohypnol:* Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*For 12th grade only, the 2001 and 2002 data are not comparable due to changes in the questionnaire forms.

FIGURE 202
Rohypnol:* Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*For 12th grade only, the 2001 and 2002 data are not comparable due to changes in the questionnaire forms.

FIGURE 203
Rohypnol:* Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

str(*For 12th grade only, the 2001 and 2002 data are not comparable due to changes in the questionnaire forms.)

**Parental education is an average score of mother's education and father's education.

FIGURE 204
Rohypnol:* Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

***For 12th grade only, the 2001 and 2002 data are not comparable due to changes in the questionnaire forms.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 205
Alcohol:* Trends in 30-Day Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

FIGURE 206
Alcohol:* Trends in 30-Day Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

FIGURE 207
Alcohol:* Trends in 30-Day Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

FIGURE 208
Alcohol:* Trends in 30-Day Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

FIGURE 209
Alcohol:* Trends in 30-Day Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

**Parental education is an average score of mother's education and father's education.

FIGURE 210
Alcohol:* Trends in 30-Day Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 211
Alcohol:* Trends in Daily Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

FIGURE 212
Alcohol:* Trends in Daily Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

FIGURE 213
Alcohol:* Trends in Daily Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

FIGURE 214
Alcohol:* Trends in Daily Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

FIGURE 215
Alcohol:* Trends in Daily Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

**Parental education is an average score of mother's education and father's education.

FIGURE 216
Alcohol:* Trends in Daily Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Beginning in 1993, a revised set of questions on alcohol use was introduced that indicated that a drink meant more than a few sips.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 217
Alcohol: Trends in 30-Day Prevalence of Having Been Drunk by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 218
Alcohol: Trends in 30-Day Prevalence of Having Been Drunk by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 219
Alcohol: Trends in 30-Day Prevalence of Having Been Drunk by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 220
Alcohol: Trends in 30-Day Prevalence of Having Been Drunk by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 221

Alcohol: Trends in 30-Day Prevalence of Having Been Drunk by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 222
Alcohol: Trends in 30-Day Prevalence of Having Been Drunk by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 223
Binge Drinking: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 224
Binge Drinking: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 225
Binge Drinking: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 226
Binge Drinking: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 227

Binge Drinking: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 228
Binge Drinking: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 229
Binge Drinking: Trends in 2-Week Prevalence of 10 or More Drinks in a Row by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 230
Binge Drinking: Trends in 2-Week Prevalence of 10 or More Drinks in a Row by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 231
Binge Drinking: Trends in 2-Week Prevalence of 10 or More Drinks in a Row by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 232
Binge Drinking: Trends in 2-Week Prevalence of 10 or More Drinks in a Row by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 233

Binge Drinking: Trends in 2-Week Prevalence of 10 or More Drinks in a Row by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 234
Binge Drinking: Trends in 2-Week Prevalence of 10 or More Drinks in a Row by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 235
Binge Drinking: Trends in 2-Week Prevalence of 15 or More Drinks in a Row by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 236
Binge Drinking: Trends in 2-Week Prevalence of 15 or More Drinks in a Row by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 237
Binge Drinking: Trends in 2-Week Prevalence of 15 or More Drinks in a Row by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 238
Binge Drinking: Trends in 2-Week Prevalence of 15 or More Drinks in a Row by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 239

Binge Drinking: Trends in 2-Week Prevalence of 15 or More Drinks in a Row by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 240
Binge Drinking: Trends in 2-Week Prevalence of 15 or More Drinks in a Row by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 241
Beer: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 242
Beer: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 243
Beer: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 244
Beer: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 245
Beer: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 246
Beer: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 247
Beer: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 248
Beer: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 249
Beer: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 250
Beer: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 251

Beer: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 252
Beer: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 253
Liquor: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 254
Liquor: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 255
Liquor: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 256
Liquor: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 257
Liquor: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 258
Liquor: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 259
Liquor: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 260
Liquor: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 261
Liquor: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 262

Liquor: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 263

Liquor: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 264
Liquor: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 265
Wine:* Trends in 30-Day Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

FIGURE 266
Wine:* Trends in 30-Day Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

FIGURE 267
Wine:* Trends in 30-Day Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

FIGURE 268
Wine:* Trends in 30-Day Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

FIGURE 269
Wine:* Trends in 30-Day Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

**Parental education is an average score of mother's education and father's education.

FIGURE 270
Wine:* Trends in 30-Day Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 271
Wine:* Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

FIGURE 272

Wine:* Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

FIGURE 273
Wine:* Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

FIGURE 274

Wine:* Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

FIGURE 275

Wine:* Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Average Education Level of Parents**



Source. The Monitoring the Future study, the University of Michigan.


Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

**Parental education is an average score of mother's education and father's education.

FIGURE 276

Wine:* Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 277
Wine Coolers:* Trends in 30-Day Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued for 8th and 10th graders in 2004.

FIGURE 278
Wine Coolers:* Trends in 30-Day Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued for 8th and 10th graders in 2004.

FIGURE 279
Wine Coolers:* Trends in 30-Day Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued for 8th and 10th graders in 2004.

FIGURE 280
Wine Coolers:* Trends in 30-Day Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued for 8th and 10th graders in 2004.

FIGURE 281
Wine Coolers:* Trends in 30-Day Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued for 8th and 10th graders in 2004.

**Parental education is an average score of mother's education and father's education.

FIGURE 282
Wine Coolers:* Trends in 30-Day Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question discontinued for 8th and 10th graders in 2004.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 283
Wine Coolers: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 284

Wine Coolers: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 285


Wine Coolers: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 286
Wine Coolers: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 287

Wine Coolers: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 288

Wine Coolers: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 289
Flavored Alcoholic Beverages: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 290
Flavored Alcoholic Beverages: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 291
Flavored Alcoholic Beverages: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 292
Flavored Alcoholic Beverages: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 293
Flavored Alcoholic Beverages: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 294
Flavored Alcoholic Beverages: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 295
Alcoholic Beverages containing Caffeine: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 296
Alcoholic Beverages containing Caffeine: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 297
Alcoholic Beverages containing Caffeine: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 298
Alcoholic Beverages containing Caffeine: Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 299

Alcoholic Beverages containing Caffeine: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 300
Alcoholic Beverages containing Caffeine: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 301
Cigarettes: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 302
Cigarettes: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 303
Cigarettes: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 304
Cigarettes: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 305
Cigarettes: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 306
Cigarettes: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 307
Cigarettes: Trends in 30-Day Prevalence of Daily Use by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 308
Cigarettes: Trends in 30-Day Prevalence of Daily Use by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 309
Cigarettes: Trends in 30-Day Prevalence of Daily Use by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 310
Cigarettes: Trends in 30-Day Prevalence of Daily Use by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 311
Cigarettes: Trends in 30-Day Prevalence of Daily Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 312
Cigarettes: Trends in 30-Day Prevalence of Daily Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 313
Cigarettes: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 314
Cigarettes: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 315
Cigarettes: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 316
Cigarettes: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 317

Cigarettes: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 318
Cigarettes: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 319
Tobacco using a Hookah: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 320
Tobacco using a Hookah: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 321
Tobacco using a Hookah: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 322
Tobacco using a Hookah: Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 323

Tobacco using a Hookah: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 324
Tobacco using a Hookah: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 325
Small Cigars*: Trends in Annual Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 12 months, on how many occasions have you smoked small cigars?

FIGURE 326
Small Cigars*[†]: Trends in Annual Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 12 months, on how many occasions have you smoked small cigars?

FIGURE 327
Small Cigars*: Trends in Annual Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 12 months, on how many occasions have you smoked small cigars?

FIGURE 328
Small Cigars*: Trends in Annual Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 12 months, on how many occasions have you smoked small cigars?

FIGURE 329
Small Cigars*: Trends in Annual Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 12 months, on how many occasions have you smoked small cigars?

**Parental education is an average score of mother's education and father's education.

FIGURE 330
Small Cigars*: Trends in Annual Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 12 months, on how many occasions have you smoked small cigars?

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 331
Flavored Little Cigars*: Trends in 30-Day Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked flavored little cigars or cigarillos?

FIGURE 332
Flavored Little Cigars*: Trends in 30-Day Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked flavored little cigars or cigarillos?

FIGURE 333
Flavored Little Cigars*: Trends in 30-Day Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked flavored little cigars or cigarillos?

FIGURE 334
Flavored Little Cigars*: Trends in 30-Day Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked flavored little cigars or cigarillos?

FIGURE 335
Flavored Little Cigars*: Trends in 30-Day Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked flavored little cigars or cigarillos?

**Parental education is an average score of mother's education and father's education.

FIGURE 336
Flavored Little Cigars*: Trends in 30-Day Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked flavored little cigars or cigarillos?

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 337
Regular Little Cigars*: Trends in 30-Day Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked regular little cigars or cigarillos?

FIGURE 338
Regular Little Cigars*: Trends in 30-Day Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked regular little cigars or cigarillos?

FIGURE 339
Regular Little Cigars*: Trends in 30-Day Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked regular little cigars or cigarillos?

FIGURE 340
Regular Little Cigars*: Trends in 30-Day Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked regular little cigars or cigarillos?

FIGURE 341
Regular Little Cigars*: Trends in 30-Day Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked regular little cigars or cigarillos?

**Parental education is an average score of mother's education and father's education.

FIGURE 342
Regular Little Cigars*: Trends in 30-Day Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Question text is: During the last 30 days, on how many days (if any) have you smoked regular little cigars or cigarillos?

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 343
Large Cigars: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 344
Large Cigars: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 345
Large Cigars: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 346
Large Cigars: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 347
Large Cigars: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 348
Large Cigars: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 349
Any Vaping*: Trends in 30-Day Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2017, the surveys switched from asking about vaping in general to asking separately about vaping nicotine, marijuana, and vaping just flavoring. Beginning in 2017, data presented are based on these new questions showing vaping of any of those substances.

FIGURE 350
Any Vaping*: Trends in 30-Day Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2017, the surveys switched from asking about vaping in general to asking separately about vaping nicotine, marijuana, and vaping just flavoring. Beginning in 2017, data presented are based on these new questions showing vaping of any of those substances.

FIGURE 351
Any Vaping*: Trends in 30-Day Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2017, the surveys switched from asking about vaping in general to asking separately about vaping nicotine, marijuana, and vaping just flavoring. Beginning in 2017, data presented are based on these new questions showing vaping of any of those substances.

FIGURE 352
Any Vaping*: Trends in 30-Day Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2017, the surveys switched from asking about vaping in general to asking separately about vaping nicotine, marijuana, and vaping just flavoring. Beginning in 2017, data presented are based on these new questions showing vaping of any of those substances.

FIGURE 353
Any Vaping*: Trends in 30-Day Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2017, the surveys switched from asking about vaping in general to asking separately about vaping nicotine, marijuana, and vaping just flavoring. Beginning in 2017, data presented are based on these new questions showing vaping of any of those substances.

**Parental education is an average score of mother's education and father's education.

FIGURE 354
Any Vaping*: Trends in 30-Day Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*In 2017, the surveys switched from asking about vaping in general to asking separately about vaping nicotine, marijuana, and vaping just flavoring. Beginning in 2017, data presented are based on these new questions showing vaping of any of those substances.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 355
Vaping Nicotine: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 356
Vaping Nicotine: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 357
Vaping Nicotine: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 358
Vaping Nicotine: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 359
Vaping Nicotine: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


*Parental education is an average score of mother's education and father's education.

FIGURE 360
Vaping Nicotine: Trends in 30-Day Prevalence by Race/Ethnicity*



Source. The Monitoring the Future study, the University of Michigan.
 Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.
 *To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 361
Vaping Marijuana: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 362
Vaping Marijuana: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 363
Vaping Marijuana: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 364
Vaping Marijuana: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 365
Vaping Marijuana: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 366
Vaping Marijuana: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 367
Vaping Just Flavoring: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 368
Vaping Just Flavoring: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 369
Vaping Just Flavoring: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 370
Vaping Just Flavoring: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 371
Vaping Just Flavoring: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 372
Vaping Just Flavoring: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 373
Smokeless Tobacco: Trends in 30-Day Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 374
Smokeless Tobacco: Trends in 30-Day Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 375
Smokeless Tobacco: Trends in 30-Day Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 376
Smokeless Tobacco: Trends in 30-Day Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 377
Smokeless Tobacco: Trends in 30-Day Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 378
Smokeless Tobacco: Trends in 30-Day Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 379
Smokeless Tobacco: Trends in 30-Day Prevalence of Daily Use by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 380
Smokeless Tobacco: Trends in 30-Day Prevalence of Daily Use by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 381
Smokeless Tobacco: Trends in 30-Day Prevalence of Daily Use by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 382
Smokeless Tobacco: Trends in 30-Day Prevalence of Daily Use by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 383

Smokeless Tobacco: Trends in 30-Day Prevalence of Daily Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 384
Smokeless Tobacco: Trends in 30-Day Prevalence of Daily Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 385
Dissolvable Tobacco Products: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 386
Dissolvable Tobacco Products: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 387
Dissolvable Tobacco Products: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 388
Dissolvable Tobacco Products: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 389
Dissolvable Tobacco Products: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 390
Dissolvable Tobacco Products: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 391
Snus: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 392
Snus: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 393
Snus: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 394
Snus: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 395
Snus: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 396
Snus: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 397
Steroids: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 398
Steroids: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 399
Steroids: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 400
Steroids: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 401
Steroids: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 402
Steroids: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 403
Marijuana: Trends in Lifetime Prevalence of Daily Use for Month or More by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 404
Marijuana: Trends in Lifetime Prevalence of Daily Use for Month or More by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 405
Marijuana: Trends in Lifetime Prevalence of Daily Use for Month or More by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 406

Marijuana: Trends in Lifetime Prevalence of Daily Use for Month or More by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 407

Marijuana: Trends in Lifetime Prevalence of Daily Use for Month or More by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 408
Marijuana: Trends in Lifetime Prevalence of Daily Use for Month or More by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 409
Nonprescription Diet Pills: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 410
Nonprescription Diet Pills: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 411
Nonprescription Diet Pills: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 412
Nonprescription Diet Pills: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 413
Nonprescription Diet Pills: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 414
Nonprescription Diet Pills: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 415
Nonprescription Stay-Awake Pills: Trends in Annual Prevalence by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 416
Nonprescription Stay-Awake Pills: Trends in Annual Prevalence by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 417
Nonprescription Stay-Awake Pills: Trends in Annual Prevalence by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 418
Nonprescription Stay-Awake Pills: Trends in Annual Prevalence by Population Density



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 419
Nonprescription Stay-Awake Pills: Trends in Annual Prevalence by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 420
Nonprescription Stay-Awake Pills: Trends in Annual Prevalence by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 421
Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 422
Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 423
Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 424
Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 425

Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 426
Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 427
Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 428
Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 429
Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 430
Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 431

Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 432
Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 433
Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 434
Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 435
Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 436

Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 437

Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 438
Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.


FIGURE 439
Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Gender



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 440
Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by College Plans



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.


FIGURE 441
Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Region



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 442
Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 443

Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 444
Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 445

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 446

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 447

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 448

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 449

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 450

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Lifetime Prevalence of Legal Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 451

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 452

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 453

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 454

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

FIGURE 455

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Average Education Level of Parents*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Parental education is an average score of mother's education and father's education.

FIGURE 456

Either Stimulant-Type or Non-Stimulant-Type ADHD Drugs: Trends in Current Prevalence of Legal Use by Race/Ethnicity*




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 457
Any Nicotine Use*: Trends in 30-Day Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, smokeless tobacco, or vaping nicotine.

FIGURE 458
Any Nicotine Use*: Trends in 30-Day Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, smokeless tobacco, or vaping nicotine.

FIGURE 459
Any Nicotine Use*: Trends in 30-Day Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, smokeless tobacco, or vaping nicotine.

FIGURE 460
Any Nicotine Use*: Trends in 30-Day Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, smokeless tobacco, or vaping nicotine.

FIGURE 461
Any Nicotine Use*: Trends in 30-Day Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, smokeless tobacco, or vaping nicotine.

**Parental education is an average score of mother's education and father's education.

FIGURE 462
Any Nicotine Use*: Trends in 30-Day Prevalence by Race/Ethnicity**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, smokeless tobacco, or vaping nicotine.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

FIGURE 463
Any Nicotine Use other than Vaping*: Trends in 30-Day Prevalence by Gender




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, or smokeless tobacco.

FIGURE 464
Any Nicotine Use other than Vaping*: Trends in 30-Day Prevalence by College Plans




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, or smokeless tobacco.

FIGURE 465
Any Nicotine Use other than Vaping*: Trends in 30-Day Prevalence by Region




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, or smokeless tobacco.

FIGURE 466
Any Nicotine Use other than Vaping*: Trends in 30-Day Prevalence by Population Density




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, or smokeless tobacco.

FIGURE 467

Any Nicotine Use other than Vaping*: Trends in 30-Day Prevalence by Average Education Level of Parents**




Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, or smokeless tobacco.

**Parental education is an average score of mother's education and father's education.

FIGURE 468
Any Nicotine Use other than Vaping*: Trends in 30-Day Prevalence by Race/Ethnicity**



Source. The Monitoring the Future study, the University of Michigan.

Note. In year 2020 chart data imputed as average of 2019 and 2021 estimates. No 2020 estimates are available because the COVID-19 pandemic curtailed data collection in this year and MTF collected only about one-quarter the number of surveys it collects in a typical year.

*Includes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, or smokeless tobacco.

**To derive percentages for each racial/ethnic subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates.

DEFINITION OF BACKGROUND AND DEMOGRAPHIC SUBGROUPS

The following are brief definitions of the background and demographic subgroups explored in the Monitoring the Future (MTF) national survey of 8th, 10th, and 12th graders' attitudes toward and use of drugs (including alcohol and tobacco). A more thorough discussion of the findings can be found in Appendix B of the MTF Volume I monograph, [*Monitoring the Future National Survey Results on Drug Use, 1975-2021: Volume I, Secondary School Students*](#).⁴ (Note: All case counts provided in the tables are based on weighted Ns.)

Total: The total sample of respondents in a given year based on weighted cases (set to equal the total number of actual cases).

Gender: *Male and female.* Respondents are asked, "What is your sex?" Those with missing data on the question are omitted from the data presented by gender.

College Plans: Respondents are asked how likely it is that they will graduate from a four-year college program. College plans groupings are defined as follows:

None or under four years. Respondents who indicate they "definitely won't" or "probably won't" graduate from a four-year college program. (Note that, among those who do not expect to complete a four-year college program, a number still expect to get some postsecondary education.)

Complete four years. Respondents who indicate they "definitely will" or "probably will" graduate from a four-year college program.

Those not answering the college plans question are omitted from both groupings.

Region: Region of the country in which the respondent's school is located. There are four mutually exclusive regions in the US based on Census Bureau categories, defined as follows:

Northeast. Census classifications of New England and Middle Atlantic states consist of Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania.

Midwest. Census classifications of East North Central and West North Central states consist of Ohio, Indiana, Illinois, Michigan, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas.

South. Census classifications of South Atlantic, East South Central, and West South Central states consist of Delaware, Maryland, District of Columbia, Virginia, and West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas.

⁴Miech, R. A., Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Patrick, M. E. (2022, forthcoming). *Monitoring the Future national survey results on drug use, 1975-2021: Volume I, Secondary school students*. Ann Arbor: Institute for Social Research, The University of Michigan.

West. Census classifications of Mountain and Pacific states consist of Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, and California (Alaska and Hawaii are also included in this Census region, but are not included in the MTF study).

**Population
Density:**

Population density of the area in which the schools are located. There are three mutually exclusive groups into which schools have been placed in a given year based on population density. The 1975–1985 samples were based on the 1970 Census; in 1986, one-half of the sample was based on the 1970 Census and the other half was based on the 1980 Census. In 1987 through 1993, all samples were based on the 1980 Census; in 1994, half of the sample was based on the 1980 Census and half on the 1990 Census. Starting in 2006 until 2013, each first-year half-sample of schools comes from a sample design that utilizes 2000 Census counts as the measure of size for first-stage units. Counts from the 2010 Census were used for the samples beginning in 2014.

The three levels of population density were defined in terms of Standard Metropolitan Statistical Area (SMSA) designations through 1985, and then changed to the new Office of Management and Budget (OMB) classifications of Metropolitan Statistical Areas (MSAs).⁵ Except in the New England states, an MSA is a county or group of contiguous counties that contain at least one city of 50,000 inhabitants or more or twin cities with a combined population of at least 50,000. In the New England states, MSAs consisted of towns and cities instead of counties until 1994, after which New England Consolidated Metropolitan Areas (NECMAs) were used to define MSAs. Each MSA must include at least one central city, and the complete title of an MSA identifies the central city or cities. For the complete description of the criteria used in defining MSAs, see the OMB publication, *Metropolitan Statistical Areas, 1990* (NTIS-PB90-214420), Washington, D.C. Although MTF has updated the measures of size of the MSAs and non-MSAs following the 2000 and 2010 Censuses, the project has not altered MSA definitions since the introduction of its new sample design in 1994. Thus, MTF continues to utilize the MSAs as defined by OMB in June 1990.⁶ The population living in an MSA is designated as the metropolitan population. The levels of population density used in MTF include those described here:

Large MSA. These were the 12 largest SMSAs as of the 1970 Census and were used for the 1975–1985 samples: New York, Los Angeles, Chicago, Philadelphia, Detroit, San Francisco, Washington, Boston, Pittsburgh, St. Louis, Baltimore, and Cleveland. As of the 1980 Census, the Large MSA group consisted of the 16 largest MSAs in the nation. This new structure was used for the 1986–1994 samples. These 16 MSAs include all of those mentioned above except Cleveland, plus Dallas-Fort Worth, Houston, Nassau-Suffolk, Minneapolis-St. Paul, and Atlanta.

⁵ OMB utilizes several names for geographic areas, such as Primary Metropolitan Statistical Areas (PMSAs) which are component parts of Consolidated Metropolitan Statistical Areas (CMSAs). For example, in June 1990, the Ann Arbor MI PMSA and Detroit MI PMSA constituted the Detroit-Ann Arbor MI CMSA. For the sake of simplicity, this document utilizes MSA throughout.

⁶ For example, OMB currently defines the Detroit-Warren-Dearborn MSA as Wayne, Oakland, Macomb, Livingston, St. Clair, and Lapeer Counties, while MTF continues to define the Detroit MSA as Wayne, Oakland, Macomb, Livingston, St. Clair, Monroe, and Lapeer Counties, as OMB defined Detroit in June 1990.

A new sample design was developed based on the 1990 Census, beginning with the first-year half-A sample of schools chosen in 1994. In the 1990s sample, only the eight largest MSAs are represented with certainty at all three grade levels; 16 other large MSAs are divided into pairs, with half randomly assigned to both the 8th- and 12th-grade samples and the other half assigned to the 10th-grade sample. The eight largest MSAs are New York, Los Angeles-Long Beach, Chicago, Philadelphia PA-NJ, Detroit, Washington DC-MD-VA, Dallas-Ft. Worth, and Boston-Lawrence-Salem-Lowell-Brockton. The other 16 large MSAs are Houston, Atlanta, Seattle-Tacoma, Minneapolis-St. Paul MN-WI, St. Louis MO-IL, San Diego, Baltimore, Pittsburgh, Phoenix, Oakland, Cleveland, Miami, Newark, Denver, San Francisco, and Kansas City MO-KS.

Other MSAs. This category consists of all other, smaller MSAs, as defined by OMB, except those listed previously.

Non-MSAs. This category consists of all areas not designated as MSAs—in other words, they do not contain a town (or twin cities) of at least 50,000 inhabitants. The population living outside of MSAs constitutes the nonmetropolitan population.

**Parental
Education:**

This is an average of mother’s education and father’s education based on the respondents’ answers about the highest level of education achieved by each parent, using the following scale: (1) completed grade school or less, (2) some high school, (3) completed high school, (4) some college, (5) completed college, and (6) graduate or professional school after college. Missing data were allowed for one of the two parents. The respondent was instructed, “If you were raised mostly by foster parents, stepparents, or others, answer for them. For example, if you have both a stepfather and a natural father, answer for the one that was most important in raising you.”

**Race/
Ethnicity:**

From 1975 through 2004, respondents were asked, “How do you describe yourself?” and were presented with a list of various racial/ethnic categories. A general instruction told them to select the *one* best response for each question. In 2005, the instructions in half of the questionnaire forms were revised in order to be more consistent with the guidelines of the Office of Management and Budget for assessing race/ethnicity. In the changed forms, respondents were presented with a list of racial/ethnic categories and instructed to “select one or more responses.” An examination of the data showed that relatively few respondents (about 6% in 2005 and about 6.6% of the sample in 2014) selected more than one racial/ethnic category. Because some survey questions appear in only one or a few forms, there was some variation in the version of the race/ethnicity question upon which the 2005 data were based. Based on the analyses we have examined, we do not believe these different permutations make any appreciable difference in the results. In 2006 and thereafter the revised instruction was used in all forms. Those checking multiple racial/ethnic groups or one of the other specified groups are omitted from the reporting on race/ethnicity in this volume because of the small numbers of cases.

White/Caucasian. Consists of those respondents who describe themselves as White or Caucasian in 1975–2004. In 2005, the unchanged questionnaire forms were treated in a

similar manner. For the revised question in 2005 and for all forms in 2006 and beyond, those checking only White and no other racial/ethnic group were categorized as White.

African American. Consists of those respondents who in 1975–1990 describe themselves as Black or Afro-American or who, in 1991–2004, describe themselves as Black or African American. In 2005, the unchanged questionnaire forms were treated in a similar manner; for the revised question in 2005 and for all forms in 2006 and beyond, only those checking Black or African American and no other racial ethnic group were categorized as African American.

Hispanic. Consists of those respondents who in 1975–1990 describe themselves as Mexican American or Chicano, or Puerto Rican or other Latin American. After 1990, this group includes those respondents who describe themselves as Mexican American or Chicano, Cuban American, Puerto Rican American, or other Latin American. (The term “Puerto Rican American” was shortened to “Puerto Rican” after 1994.) In 2005, the unchanged questionnaire forms were treated in a similar manner; the changed forms

in 2005 and for all forms in 2006 and beyond, only those checking Mexican American or Chicano, Cuban American, Puerto Rican, or Other Hispanic or Latino and no other racial/ethnic group were categorized as Hispanic.

TRENDS BY SUBGROUP: SUPPLEMENTAL TABLES FOR SECONDARY SCHOOL STUDENTS

The tables in this section contain the numbers underlying the trend lines shown in the figures on subgroup trends. Trends in subgroup data for the 1975–2020 Monitoring the Future (MTF) results containing 8th, 10th, and 12th graders' use of drugs and alcohol are presented in tables below. The trends observed span major classes of drugs for population subgroups defined by gender, college plans, region, population density, parental education, and race/ethnicity. Appendix D of Volume I contains a portion of the information presented here.⁷

Due to the sheer quantity of information such trend tables generate for each prevalence measure (e.g., lifetime, annual, 30-day, daily), we have selected only the prevalence periods that seem most useful for understanding subgroup differences below. Thus, for most drugs, we include only annual prevalence; rates for additional prevalence periods are provided for alcohol, cigarettes, and smokeless tobacco because of their more frequent use. The numbers of cases that go with the data in the figures may be found in the complete set of tables that are included below.

The subgroups distinguished in these tables are used throughout Volume I and defined in the previous section. The reader should note that two-year moving averages are given for the three major racial/ethnic groups included in order to reduce fluctuations in the trends due to random sampling differences. A footnote in each table describes the procedure used to obtain the presented data. A question about race/ethnicity identification was changed in 2005 in half of the questionnaire forms; allowing respondents to select multiple categories of race/ethnicity rather than just one (see the previous section for more details). In 2006, the remaining questionnaire forms were changed in a like manner. However, we believe that this change has had little effect on the results because so few respondents selected more than one of the categories offered.

For nearly all drugs, we provide one table showing the subgroup trends (since 1991) for 8th graders, a second table of trends (since 1991) for 10th graders and a third table showing the longer-term trends (since 1975) for 12th graders. However, for two of the drugs—sedatives (barbiturates) and narcotics other than heroin—the 8th- and 10th-grade data have been omitted, as they have been throughout MTF publications because we are less certain about the validity of the answers provided by those younger students. Specifically, we believe that they often fail to omit nonprescription substances. Usage questions for a few other drugs are simply not asked of 8th and 10th graders; thus, only 12th-grade tables are presented.

Sample sizes should be taken into account when interpreting the importance of any changes observed. The numbers provided in the tables assume that all respondents in a given grade were asked about their use of the drug. Some of the drugs were not contained in all questionnaire forms, meaning that the subgroup and total numbers must be adjusted accordingly. The “Notes” section at the bottom of each table will indicate if only a fraction of the total sample received the question.

This paper also includes graphic presentations of the trends presented in the tables for the various demographic subgroups. These figures use color to help distinguish the various subgroups and improve readability.

⁷Miech, R. A., Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Patrick, M. E. (2022, forthcoming). *Monitoring the Future national survey results on drug use, 1975-2021: Volume I, Secondary school students*. Ann Arbor: Institute for Social Research, The University of Michigan.

List of 2021 Tables

Substance	Table Number			Time Period				
	8th Grade	10th Grade	12th Grade	Lifetime	Annual	30-Day	Daily	2-Weeks
Any illicit drug	1	2	3		X			
Any illicit drug other than marijuana	4	5	6		X			
Marijuana, Lifetime	7	8	9	X				
Marijuana, Annual	10	11	12		X			
Marijuana, 30-Day	13	14	15			X		
Marijuana, Daily	16	17	18				X	
Synthetic Marijuana	19	20	21		X			
Inhalants	22	23	24		X			
Hallucinogens	25	26	27		X			
LSD	28	29	30		X			
Hallucinogens other than LSD	31	32	33		X			
Ecstasy (MDMA)	34	35	36		X			
Salvia	37	38	39		X			
Cocaine	40	41	42		X			
Crack	43	44	45		X			
Other cocaine	46	47	48		X			
Heroin	49	50	51		X			
Heroin with a needle	52	53	54		X			
Heroin without a needle	55	56	57		X			
Narcotics other than heroin	—	—	58		X			
OxyContin	59	60	61		X			
Vicodin	62	63	64		X			
Amphetamines	65	66	67		X			
Ritalin	68	69	70		X			
Adderall	71	72	73		X			
Provigil	—	—	74		X			
Methamphetamine	75	76	77		X			
Crystal methamphetamine (ice)	—	—	78		X			
Bath Salts (Synthetic Stimulants)	79	80	81		X			
Sedatives (barbiturates)	—	—	82		X			
Tranquilizers	83	84	85		X			
Any Prescription Drug	—	—	86					
OTC cough/cold medicines	87	88	89		X			
Rohypnol	90	91	92		X			
Alcohol, 30-Day	93	94	95			X		
Alcohol, Daily	96	97	98				X	
Been drunk	99	100	101			X		
5+ drinks in a row	102	103	104					X
10+ drinks in a row	105	105	105					X
15+ drinks in a row	—	—	106					X
Beer	107	108	109			X		
5+ drinks in a row	110	111	112					X
Liquor	—	—	113			X		
5+ drinks in a row	—	—	114					X
Wine	—	—	115			X		
5+ drinks in a row	—	—	116					X
Wine coolers	117	118	119			X		
5+ drinks in a row	—	—	120					X
Flavored alcoholic beverages	121	122	123			X		
Alcoholic Beverages containing Caffeine	124	125	126		X			
Cigarettes	127	128	129			X		
Daily	130	131	132				X	
1/2 pack+/day	133	134	135				X	
Tobacco using a Hookah	—	—	136		X			
Small cigars	—	—	137		X			
Flavored little cigars or cigarillos	138	138	138			X		
Regular little cigars or cigarillos	139	139	139			X		
Large cigars	140	140	140			X		

Substance	Table Number			Time Period				
	8th Grade	10th Grade	12th Grade	Lifetime	Annual	30-Day	Daily	2-Weeks
Any Vaping	141	141	141			X		
Vaping nicotine	142	142	142			X		
Vaping marijuana	143	143	143			X		
Vaping just flavoring	144	144	144			X		
Smokeless tobacco	145	146	147			X		
Daily	148	149	150				X	
Dissolvable Tobacco Products	151	152	153		X			
Snus	154	155	156		X			
Steroids	157	158	159		X			
Marijuana, Ever Daily for Month or More	—	—	160				X	
Nonprescription Diet Pills	—	—	161		X			
Nonprescription Stay-Awake Pills	—	—	162		X			
Stimulant-Type ADHD Drugs, Lifetime	163	164	165	X				
Stimulant-Type ADHD Drugs, Current	166	167	168					
Non-Stimulant-Type ADHD Drugs, Lifetime	169	170	171	X				
Non-Stimulant-Type ADHD Drugs, Current	172	173	174					
Either Type ADHD Drugs, Lifetime	175	176	177	X				
Either Type ADHD Drugs, Current	178	179	180					
Any Nicotine Use	—	—	181			X		
Any Nicotine Use other than Vaping	—	—	182			X		
Marijuana, Under a Doctor's Orders			183	X				
Powdered Alcohol	184	184	184		X			
Energy Drinks	185	186	187				X	
Energy Shots	188	189	190				X	
Either Energy Drinks or Shots	191	192	193				X	
Weighted <i>N</i> s by subgroups	S1	S2	S3					

TABLE 1
ANY ILLICIT DRUG: ^{a,d} Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Approximate weighted N =	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	11.3	12.9	15.1	18.5	21.4	23.6	22.1	21.0	20.5	19.5	19.5	17.7	16.1	15.2	15.5
Gender:															
Male	11.7	11.9	15.2	19.4	22.3	23.6	22.6	21.3	21.3	19.7	21.3	19.2	16.4	15.0	15.2
Female	11.0	13.6	14.9	17.6	20.2	23.3	21.3	20.4	19.7	19.0	17.5	16.3	15.5	15.2	15.6
College Plans:															
None or under 4 years	22.8	25.6	30.7	34.6	38.4	40.3	39.6	41.3	39.9	38.9	38.5	36.8	34.7	33.6	35.2
Complete 4 years	9.5	10.9	12.8	16.3	19.1	21.0	19.9	18.4	18.0	17.1	17.2	15.7	14.0	13.2	13.3
Region:															
Northeast	9.3	10.6	11.5	16.6	17.9	20.3	20.2	16.0	18.2	16.6	18.1	13.8	13.1	13.0	12.3
Midwest	11.2	13.0	13.9	17.2	23.3	24.7	22.3	21.9	22.6	20.6	18.0	17.0	15.7	14.2	15.9
South	11.5	12.9	15.1	17.6	20.8	22.5	21.6	22.3	21.0	19.2	21.5	20.1	18.1	16.8	17.4
West	13.3	15.0	21.1	23.7	23.3	27.1	24.4	22.0	19.2	21.0	18.9	18.4	15.6	15.5	14.5
Population Density:															
Large MSA	10.5	12.0	13.1	16.2	15.2	23.4	20.5	19.8	19.1	18.1	16.4	15.9	14.3	14.0	14.9
Other MSA	12.1	14.4	17.3	21.5	23.7	24.9	22.6	21.4	19.5	18.8	21.5	18.3	16.2	15.9	16.2
Non-MSA	10.8	11.2	12.9	14.0	20.3	21.4	22.9	21.6	24.3	22.7	19.1	18.9	18.1	15.3	15.0
Parental Education: ^b															
1.0–2.0 (Low)	19.5	18.5	20.8	26.1	29.8	26.7	29.5	30.4	30.2	30.9	29.9	27.3	26.4	25.7	24.9
2.5–3.0	11.7	14.1	17.1	20.2	24.3	25.7	25.5	24.2	24.9	23.9	23.3	22.4	20.4	18.5	20.5
3.5–4.0	11.6	13.6	15.4	19.7	23.4	26.2	23.8	21.8	21.0	20.1	21.5	18.5	16.9	16.7	16.7
4.5–5.0	8.7	10.2	12.8	15.7	17.4	21.3	19.3	17.8	15.6	14.6	15.0	14.5	12.0	12.1	11.4
5.5–6.0 (High)	10.2	10.1	11.8	14.9	17.7	19.8	16.8	17.1	15.8	15.0	13.4	12.1	11.0	10.9	9.8
Race/Ethnicity (2-year average): ^c															
White	—	11.8	13.6	15.7	19.2	22.4	23.0	21.5	19.9	19.1	19.0	18.3	16.5	14.9	14.4
African American	—	7.9	9.3	13.0	15.8	17.5	18.1	18.1	18.6	18.3	16.7	15.1	14.6	14.6	15.6
Hispanic	—	18.1	20.6	24.6	26.7	26.9	26.5	26.7	27.4	25.1	24.3	24.8	22.8	20.9	18.8

Table continued on next page.

TABLE 1 (cont.)
ANY ILLICIT DRUG:^{a,d} Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	14.8	13.2	14.1	14.5	16.0	14.7	13.4‡	15.2	14.6	14.8	12.0	12.9	13.4	14.8	15.6	10.2	-4.7* sss
Gender:																	
Male	15.1	13.2	14.3	15.7	16.7	15.7	13.8‡	14.9	14.1	14.0	11.3	11.6	13.2	12.6	§	7.4	-5.2 sss
Female	14.2	12.9	13.7	13.1	15.4	13.3	12.7‡	14.9	14.5	15.3	12.5	13.6	13.3	16.7	§	11.4	-5.2 sss
College Plans:																	
None or under 4 years	33.1	31.9	34.5	30.7	33.6	32.1	30.1‡	31.3	29.0	27.5	25.4	24.7	25.8	27.0	§	15.9	-11.1 sss
Complete 4 years	13.0	11.4	12.3	13.0	14.6	13.2	12.1‡	14.0	13.3	13.6	10.7	11.5	11.6	13.0	§	9.1	-3.9 sss
Region:																	
Northeast	10.8	8.1	11.3	13.8	14.1	9.8	9.6‡	12.5	12.9	11.6	7.5	10.0	8.8	10.8	§	8.2	-2.6
Midwest	13.9	12.9	15.5	13.5	14.2	13.0	9.9‡	13.4	13.4	15.6	12.2	11.2	11.3	13.9	§	11.7	-2.2
South	17.7	15.2	15.6	15.9	18.4	15.7	15.8‡	17.3	14.7	14.9	13.1	13.2	13.2	15.4	§	9.6	-5.8 sss
West	13.9	13.3	12.4	13.5	15.5	18.6	15.6‡	15.5	16.7	16.1	13.0	16.0	19.2	17.6	§	11.0	-6.6 s
Population Density:																	
Large MSA	13.5	11.5	12.5	12.4	15.3	13.4	12.6‡	15.9	16.0	14.8	11.3	12.5	14.0	13.1	§	8.6	-4.5 s
Other MSA	15.6	13.8	15.1	15.7	17.2	16.1	14.3‡	14.8	14.1	15.7	11.8	13.5	13.5	15.5	§	11.2	-4.3 ss
Non-MSA	15.2	14.5	14.2	14.5	14.4	13.7	12.7‡	14.9	13.7	12.9	13.3	12.0	12.0	15.8	§	10.3	-5.5 s
Parental Education: ^b																	
1.0–2.0 (Low)	23.7	23.2	23.3	21.5	24.6	23.4	22.0‡	25.8	22.9	24.0	17.9	19.8	19.0	22.1	§	14.6	-7.5 s
2.5–3.0	18.6	16.8	19.2	19.0	20.6	20.1	18.4‡	22.1	19.6	18.8	15.1	17.5	18.0	23.2	§	15.4	-7.7 sss
3.5–4.0	16.5	14.9	15.3	15.8	17.1	16.4	15.4‡	17.9	16.4	16.1	13.5	13.5	15.7	16.3	§	13.2	-3.1
4.5–5.0	11.0	9.2	10.5	11.1	12.8	10.7	9.6‡	11.0	10.4	10.3	9.7	9.4	9.9	11.4	§	8.7	-2.7 s
5.5–6.0 (High)	9.9	8.1	9.6	9.9	10.5	8.7	7.9‡	9.0	10.5	10.0	7.8	8.7	7.5	9.1	§	7.1	-2.0
Race/Ethnicity (2-year average): ^c																	
White	14.1	13.1	13.0	13.5	14.1	13.8	12.1	11.1‡	11.5	11.5	10.4	9.8	10.3	10.9	§	12.6	+1.7
African American	14.9	12.7	12.3	14.2	15.7	15.0	13.5	14.9‡	14.9	14.7	14.3	13.9	13.1	13.8	§	11.9	-1.9
Hispanic	17.9	16.8	16.1	16.8	18.2	19.3	19.3	18.9‡	18.9	17.9	15.8	14.3	15.5	17.6	§	11.9	-5.8 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. † ‡ indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. *Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. The 2013 data are based on half the forms; N is one half of N indicated. Beginning in 2014 data are based on all forms. See the amphetamine notes for details of the changes.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aUse of any illicit drug includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of amphetamines or tranquilizers not under a doctor's orders. The use of narcotics other than heroin and sedatives (barbiturates) has been excluded because 8th and 10th graders appear to overreport their use (perhaps because they include the use of nonprescription drugs in their answers).

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

Any illicit drug use and any illicit drug use including inhalants were also impacted by this change.

TABLE 2
ANY ILLICIT DRUG: ^{a,d} Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Approximate weighted N =	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	21.4	20.4	24.7	30.0	33.3	37.5	38.5	35.0	35.9	36.4	37.2	34.8	32.0	31.1	29.8
Gender:															
Male	21.6	20.4	25.1	31.8	33.7	38.8	40.1	35.3	37.0	39.4	39.6	35.9	33.2	32.0	30.5
Female	21.1	20.1	24.0	28.0	32.5	36.3	36.8	34.7	34.6	33.5	35.0	33.7	30.8	30.2	28.9
College Plans:															
None or under 4 years	32.7	32.0	37.7	43.2	47.3	52.4	55.2	50.5	51.8	53.5	52.7	51.5	48.6	46.9	45.8
Complete 4 years	18.9	17.8	21.9	27.0	30.8	35.0	35.7	32.2	33.2	33.9	34.6	32.1	29.2	28.9	27.6
Region:															
Northeast	21.8	19.0	26.9	29.6	32.4	37.7	37.8	39.0	38.2	34.0	37.4	35.2	32.6	32.8	31.3
Midwest	21.7	20.7	22.4	28.5	32.1	37.6	37.7	32.0	35.2	34.8	35.9	33.7	28.8	28.8	29.1
South	19.2	17.9	23.3	29.2	33.2	37.9	38.7	35.1	34.8	36.0	36.7	33.9	34.0	32.9	31.1
West	23.7	25.5	28.9	34.4	36.1	36.8	40.2	34.5	36.0	41.6	40.7	37.7	32.3	29.8	26.9
Population Density:															
Large MSA	21.4	19.9	24.0	29.4	28.7	35.5	37.2	32.6	35.0	36.5	34.0	33.0	30.5	28.9	28.0
Other MSA	22.0	20.8	25.1	32.7	35.5	40.0	40.0	36.9	37.3	36.6	39.0	36.9	32.4	32.0	31.1
Non-MSA	20.4	20.1	24.4	24.7	30.7	35.1	37.2	34.5	33.9	35.8	37.4	32.8	33.5	32.3	29.5
Parental Education: ^b															
1.0–2.0 (Low)	25.5	24.8	29.2	32.6	38.2	39.5	38.3	36.6	42.2	42.4	39.0	39.4	38.8	37.7	33.2
2.5–3.0	23.0	21.3	25.4	31.1	37.1	39.1	40.8	39.1	39.4	39.2	41.6	39.4	36.4	34.0	34.3
3.5–4.0	21.2	20.6	24.9	30.5	34.7	40.1	41.6	35.6	35.4	39.5	38.2	35.5	33.3	33.6	30.6
4.5–5.0	19.4	18.7	22.5	28.1	30.9	35.5	36.3	31.9	32.8	32.6	35.1	31.9	27.8	27.7	27.6
5.5–6.0 (High)	21.1	18.5	23.6	27.2	26.6	33.6	33.7	31.5	34.6	31.3	32.7	29.1	27.5	26.2	25.9
Race/Ethnicity (2-year average): ^c															
White	—	22.4	23.7	27.9	32.6	36.5	39.3	38.2	36.4	36.9	37.6	37.6	35.0	32.2	31.0
African American	—	10.8	11.9	18.5	23.6	27.3	30.2	28.9	28.4	29.7	30.5	28.5	27.3	29.3	29.0
Hispanic	—	23.6	26.3	30.3	34.3	40.0	41.3	38.1	38.4	39.3	38.8	36.2	33.8	34.5	32.8

Table continued on next page.

TABLE 2 (cont.)
ANY ILLICIT DRUG: ^{a,d} Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	28.7	28.1	26.9	29.4	30.2	31.1	30.1‡	32.1	29.9	27.9	26.8	27.8	29.9	31.0	30.4	18.7	-12.4* sss
Gender:																	
Male	29.3	29.5	27.9	31.6	33.0	33.8	32.9‡	34.9	30.3	27.8	25.9	26.1	29.5	29.5	§	16.9	-12.6 sss
Female	28.1	26.7	25.6	27.3	27.3	28.2	27.5‡	29.4	29.4	27.6	27.1	28.9	30.2	32.1	§	18.9	-13.2 sss
College Plans:																	
None or under 4 years	46.2	45.7	47.0	48.5	48.8	50.4	46.2‡	46.2	44.4	44.3	41.0	39.1	43.2	42.1	§	30.3	-11.8 sss
Complete 4 years	26.4	25.7	24.2	27.3	28.0	28.9	28.5‡	30.6	28.4	26.1	25.2	26.1	28.0	28.8	§	15.6	-13.2 sss
Region:																	
Northeast	27.8	27.7	28.6	26.2	27.9	30.6	33.2‡	29.7	29.5	28.9	26.0	27.8	32.3	32.5	§	21.4	-11.1 sss
Midwest	28.2	28.1	25.8	26.2	28.1	29.0	26.3‡	29.9	26.6	23.9	25.5	27.5	27.4	28.8	§	20.1	-8.6 ss
South	29.7	26.7	27.1	31.6	31.0	31.4	30.3‡	33.6	31.1	28.4	27.3	26.1	28.9	30.1	§	19.3	-10.8 sss
West	29.0	30.2	26.2	32.6	33.3	33.2	31.6‡	34.1	31.8	30.2	27.8	30.4	32.2	33.9	§	14.3	-19.6 sss
Population Density:																	
Large MSA	27.3	27.9	26.4	29.0	28.9	29.5	30.8‡	31.0	30.0	28.2	26.4	29.9	31.6	31.5	§	14.4	-17.1 sss
Other MSA	28.6	28.3	28.1	30.7	31.9	32.6	30.2‡	33.8	30.5	28.2	26.6	27.4	30.7	32.1	§	19.8	-12.3 sss
Non-MSA	31.4	27.8	24.8	27.4	28.6	30.2	29.2‡	29.5	28.5	26.3	28.0	25.2	25.3	27.1	§	23.3	-3.8
Parental Education: ^b																	
1.0–2.0 (Low)	36.2	32.0	33.6	39.5	37.4	37.7	39.8‡	40.5	35.0	36.8	30.1	31.8	34.1	35.1	§	25.6	-9.4 s
2.5–3.0	33.7	33.0	31.7	32.8	36.1	36.3	34.5‡	40.6	34.8	34.0	31.9	31.9	34.1	38.8	§	22.7	-16.0 sss
3.5–4.0	30.6	30.2	29.2	30.3	31.9	33.2	31.8‡	32.1	33.6	31.8	28.7	30.4	31.8	33.7	§	21.8	-11.9 sss
4.5–5.0	24.6	24.7	22.3	26.5	24.6	27.2	27.5‡	29.2	25.5	24.5	24.1	24.7	26.9	26.8	§	16.4	-10.4
5.5–6.0 (High)	24.3	21.7	21.7	24.5	25.7	25.1	24.8‡	24.2	24.3	20.7	23.5	24.0	26.5	25.1	§	13.7	-11.4 sss
Race/Ethnicity (2-year average): ^c																	
White	30.2	29.4	28.0	27.9	29.2	30.0	29.7	29.2‡	29.0	27.2	26.6	27.1	27.7	28.2	§	23.8	-4.3 ss
African American	26.2	24.2	22.4	23.5	26.6	29.6	30.6	32.5‡	33.2	30.8	28.5	26.2	26.4	29.5	§	22.4	-7.2 s
Hispanic	28.5	27.3	30.1	33.1	34.0	33.5	34.3	36.3‡	35.8	32.8	29.8	28.8	31.0	32.7	§	25.3	-7.5

Source. The Monitoring the Future study, the University of Michigan.

Notes. ‡ indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of
 Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See
 Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. The 2013 data are based on half the forms; N is
 one half of N indicated. Beginning in 2014 data are based on all forms. See the amphetamine notes for details of the changes.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aUse of any illicit drug includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of amphetamines or tranquilizers
 not under a doctor's orders. The use of narcotics other than heroin and sedatives (barbiturates) has been excluded because 8th and 10th graders appear
 to overreport their use (perhaps because they include the use of nonprescription drugs in their answers).

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data
 that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, a

edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

Any illicit drug use and any illicit drug use including inhalants were also impacted by this change.

TABLE 3
ANY ILLICIT DRUG: ^{a,b,c} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	45.0	48.1	51.1	53.8	54.2	53.1	52.1	49.4	47.4	45.8	46.3	44.3	41.7	38.5	35.4	32.5
Gender:																
Male	49.0	52.6	55.4	58.6	58.1	56.0	53.6	51.8	49.7	48.0	48.3	45.7	43.2	40.6	37.7	34.3
Female	41.4	43.0	46.7	48.7	50.1	49.8	50.8	46.3	44.4	42.8	43.8	42.3	39.7	36.1	32.8	30.1
College Plans:																
None or under 4 years	—	50.6	54.3	55.5	56.8	56.5	55.8	53.4	50.8	50.3	50.1	48.6	46.7	42.0	40.9	37.8
Complete 4 years	—	44.3	46.8	50.5	50.5	49.7	48.6	45.5	43.7	41.4	43.1	41.2	39.0	36.5	32.6	29.6
Region:																
Northeast	51.5	55.3	56.8	61.6	62.9	58.9	58.8	55.1	53.8	54.7	53.2	49.7	45.8	41.2	36.0	36.4
Midwest	45.5	47.6	51.9	54.6	55.0	53.1	53.4	50.3	46.9	42.4	45.7	45.0	42.7	39.7	38.7	34.0
South	38.1	42.3	46.2	47.5	45.4	47.0	43.7	42.2	41.3	41.4	37.2	37.4	35.9	34.2	30.7	27.6
West	48.3	49.7	50.0	53.2	56.4	55.8	55.5	51.7	50.7	49.1	53.3	47.8	45.7	41.8	39.5	34.4
Population Density:																
Large MSA	54.5	54.6	56.3	60.3	61.3	59.9	57.8	54.8	52.0	49.7	49.9	48.0	43.9	39.1	32.9	32.6
Other MSA	45.0	47.8	52.4	54.5	55.2	53.8	52.1	49.8	49.6	46.7	46.5	45.1	42.7	40.5	36.3	33.5
Non-MSA	38.8	43.7	45.2	47.4	47.6	47.0	47.6	44.0	41.1	41.4	43.0	40.0	37.6	34.3	36.0	30.1
Parental Education: ^c																
1.0–2.0 (Low)	—	43.4	45.3	47.7	50.2	49.5	48.1	44.3	45.1	42.7	42.8	38.1	35.4	35.8	28.4	26.6
2.5–3.0	—	49.2	51.8	53.3	53.2	53.0	51.2	48.8	46.3	45.7	46.0	44.8	41.8	37.2	35.3	32.7
3.5–4.0	—	48.9	53.1	55.1	56.1	54.2	52.8	50.8	46.5	47.6	47.2	45.6	42.2	38.6	37.7	33.8
4.5–5.0	—	50.8	51.7	56.3	57.1	54.0	53.4	49.7	48.9	44.9	48.4	44.7	43.1	40.0	35.5	33.1
5.5–6.0 (High)	—	51.3	51.8	59.1	54.3	55.0	54.8	48.5	46.1	45.5	44.5	44.5	43.5	40.6	36.3	33.3
Race/Ethnicity (2-year average): ^d																
White	—	—	50.4	53.5	55.2	54.9	54.4	50.7	49.3	47.4	47.6	47.2	45.2	43.0	40.3	37.5
African American	—	—	40.8	42.8	41.5	40.5	39.0	36.4	38.5	37.8	35.9	33.3	28.9	25.0	21.3	17.0
Hispanic	—	—	49.9	49.5	48.4	48.1	46.8	42.7	42.0	43.1	43.9	42.8	38.9	35.4	30.1	26.4

Table continued on next page.

TABLE 3 (cont.)
ANY ILLICIT DRUG: ^{a,b,c} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	29.4	27.1	31.0	35.8	39.0	40.2	42.4	41.4	42.1	40.9	41.4	41.0	39.3	38.8	38.4	36.5
Gender:																
Male	32.1	29.0	33.5	38.6	41.5	43.4	44.1	45.2	45.0	43.4	43.8	43.5	41.3	41.4	42.1	37.5
Female	26.2	24.7	27.9	32.7	35.8	36.2	40.0	37.2	38.9	38.0	38.4	37.8	36.7	35.9	34.5	35.0
College Plans:																
None or under 4 years	33.9	33.5	34.9	40.8	44.1	46.2	48.8	47.3	47.9	45.1	46.2	46.2	46.8	44.4	46.5	46.1
Complete 4 years	27.1	24.4	29.2	33.6	36.7	37.8	40.1	39.1	40.3	38.8	39.6	39.3	36.6	36.8	36.1	33.8
Region:																
Northeast	31.9	28.7	36.1	39.4	41.7	44.6	47.7	46.2	42.5	46.2	47.4	45.7	43.7	43.9	43.2	42.0
Midwest	31.3	27.8	30.8	36.5	40.9	40.9	40.6	38.1	42.1	38.9	42.0	41.6	40.0	38.2	38.8	36.3
South	24.5	23.7	28.2	34.1	36.4	37.6	38.8	40.5	40.8	35.3	37.3	38.1	34.8	35.7	36.6	34.9
West	32.6	31.1	31.8	34.7	38.2	39.1	45.9	43.1	44.2	47.4	41.9	40.5	41.4	39.6	36.3	34.6
Population Density:																
Large MSA	28.6	26.8	32.9	36.4	41.7	41.3	42.1	42.0	42.4	41.1	43.9	41.9	35.7	36.3	38.5	38.2
Other MSA	33.0	27.3	31.7	37.8	39.0	42.3	44.2	42.1	43.3	42.6	41.0	42.4	42.7	42.4	39.8	36.7
Non-MSA	23.8	27.0	28.4	31.6	35.9	35.4	39.2	39.3	39.9	37.5	39.4	37.1	37.6	34.9	35.5	33.5
Parental Education: ^c																
1.0–2.0 (Low)	28.7	27.7	29.5	32.9	37.7	36.6	40.3	38.9	40.9	41.3	38.8	35.3	35.8	34.7	32.0	37.6
2.5–3.0	28.7	26.4	29.2	35.4	38.3	39.9	40.8	40.5	43.7	40.5	41.9	39.3	39.0	37.7	38.9	38.2
3.5–4.0	29.6	28.1	31.6	36.4	38.8	40.4	42.0	42.9	42.9	41.6	42.7	43.2	41.3	42.1	41.3	37.1
4.5–5.0	28.7	26.2	31.5	36.5	39.0	40.5	43.6	40.9	40.0	39.6	41.3	42.0	38.9	38.2	38.8	33.2
5.5–6.0 (High)	31.9	26.8	33.4	35.7	40.7	40.6	44.0	41.8	42.3	41.1	40.1	40.6	36.5	37.4	36.0	37.3
Race/Ethnicity (2-year average): ^d																
White	33.9	30.5	31.4	35.5	39.0	40.8	42.8	44.0	43.3	42.8	43.1	43.6	42.8	42.0	41.6	40.0
African American	14.7	14.5	16.6	23.5	29.6	32.4	33.0	32.3	32.8	32.7	31.7	30.4	28.3	27.7	29.0	30.0
Hispanic	29.4	30.3	28.8	31.2	35.5	38.0	41.2	41.9	42.5	44.8	41.8	39.0	35.8	34.4	34.5	34.2

Table continued on next page.

TABLE 3 (cont.)
ANY ILLICIT DRUG: ^{a,b,c} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	35.9	36.6	36.5	38.3	40.0	39.7‡	40.1	38.7	38.6	38.3	39.9	38.8	38.0	36.8	32.0	-6.0* sss
Gender:																
Male	38.4	38.8	39.9	41.4	43.2	43.5‡	43.2	39.8	39.7	39.2	40.2	39.7	37.5	§	29.5	-8.0 sss
Female	33.2	34.1	33.2	34.7	36.2	35.1‡	36.5	36.9	37.2	37.0	38.9	36.8	37.7	§	33.4	-4.2
College Plans:																
None or under 4 years	43.7	46.0	42.7	46.5	46.6	44.3‡	47.5	44.5	42.5	40.4	42.4	42.5	42.4	§	34.9	-7.5 s
Complete 4 years	34.0	34.2	34.9	36.1	38.2	38.2‡	38.1	36.7	37.3	37.1	39.0	37.2	36.3	§	30.6	-5.7 ss
Region:																
Northeast	41.9	40.0	40.5	44.1	42.8	40.9‡	43.7	43.0	43.3	40.7	42.9	42.8	39.1	§	38.1	-1.0
Midwest	37.3	35.1	36.5	36.3	37.9	35.3‡	35.4	38.3	35.8	37.0	41.1	40.1	37.1	§	31.9	-5.1 s
South	33.6	36.0	34.7	36.2	37.0	37.5‡	38.8	37.7	35.3	35.7	36.2	34.6	36.4	§	25.7	-10.7 sss
West	33.4	36.5	36.0	38.9	44.4	46.1‡	44.3	37.0	43.1	42.3	42.8	41.7	42.1	§	37.4	-4.7
Population Density:																
Large MSA	37.9	37.7	40.2	39.9	41.5	43.0‡	43.2	38.8	41.6	40.8	42.4	39.9	38.4	§	35.2	-3.2
Other MSA	35.1	35.7	35.8	39.4	41.3	40.3‡	40.9	41.2	38.3	38.1	39.2	38.1	38.4	§	32.0	-6.4 ss
Non-MSA	34.7	37.2	33.4	33.2	34.6	32.9‡	33.6	32.3	34.2	34.7	37.2	38.7	36.4	§	26.8	-9.6 ss
Parental Education: ^c																
1.0–2.0 (Low)	35.6	34.6	33.0	36.9	40.3	45.4‡	41.6	40.7	40.3	37.7	37.1	36.2	37.3	§	25.4	-11.9 sss
2.5–3.0	37.8	37.5	37.4	39.2	40.5	40.9‡	42.8	41.4	41.1	38.9	42.2	42.2	39.6	§	32.9	-6.6
3.5–4.0	35.0	35.9	38.4	40.6	41.7	40.8‡	41.8	39.1	41.0	40.3	41.6	41.2	42.1	§	35.8	-6.3 ss
4.5–5.0	34.9	36.5	35.4	37.9	38.5	38.5‡	38.1	36.5	35.4	37.4	38.9	37.4	33.0	§	31.5	-1.5
5.5–6.0 (High)	36.9	36.7	36.6	32.6	37.9	34.8‡	37.9	37.1	36.1	35.5	37.5	35.6	37.9	§	32.3	-5.7
Race/Ethnicity (2-year average): ^d																
White	38.5	38.7	38.7	38.8	40.2	40.3	39.3‡	38.9	38.5	38.5	39.2	39.5	38.1	§	33.5	-4.6 s
African American	28.7	28.6	30.4	33.1	34.8	35.0	37.7‡	39.6	36.3	35.6	38.0	38.8	38.3	§	36.6	-1.7
Hispanic	32.2	31.4	32.6	34.5	37.5	40.5	42.2‡	39.9	38.6	39.7	39.5	37.2	35.9	§	30.2	-5.8

Table continued on next page.

TABLE 3 (cont.)
ANY ILLICIT DRUG:^{a,b,c} Trends in Annual Prevalence of Use by Subgroups in Grade 12

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimates and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. The 2013 data are based on half the forms; *N* is one half of *N* indicated. Beginning in 2014 data are based on all forms. See the amphetamine notes for details of the changes.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aUse of any illicit drug includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of narcotics other than heroin, amphetamines, sedatives (barbiturates), methaqualone (excluded since 1990), or tranquilizers not under a doctor's orders.

^bBeginning in 1982 the question about amphetamine use was revised to get respondents to exclude the inappropriate reporting of nonprescription amphetamines. The prevalence-of-use rate dropped slightly as a result of this methodological change.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^dEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^eFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. *N* is one half of *N* indicated.

Any illicit drug use and any illicit drug use including inhalants were also impacted by this change.

TABLE 4
ANY ILLICIT DRUG OTHER THAN MARIJUANA: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^b	2002	2003	2004	2005
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	8.4	9.3	10.4	11.3	12.6	13.1	11.8	11.0	10.5	10.2†	10.8	8.8	8.8	7.9	8.1
Gender:															
Male	8.0	8.0	9.2	10.1	11.5	11.0	10.8	9.6	9.7	9.1†	10.0	8.1	7.9	7.0	6.6
Female	8.8	10.4	11.5	12.3	13.5	14.7	12.6	12.1	11.2	10.9†	11.2	9.3	9.4	8.8	9.3
College Plans:															
None or under 4 years	16.3	18.5	21.3	21.2	25.3	23.0	22.1	23.8	23.4	22.7†	21.5	19.7	20.0	18.0	18.2
Complete 4 years	7.2	8.0	8.9	9.9	10.9	11.6	10.6	9.4	9.0	8.7†	9.5	7.6	7.5	6.9	7.0
Region:															
Northeast	6.8	6.6	8.2	10.3	10.7	11.3	9.5	8.5	8.5	8.0†	9.5	5.8	7.1	5.9	5.8
Midwest	8.6	10.4	9.4	10.2	14.0	14.3	12.5	10.5	11.9	11.2†	9.9	8.7	8.7	7.7	8.1
South	8.6	9.7	11.0	11.7	12.5	12.6	11.8	12.5	11.2	10.3†	12.4	10.6	10.0	9.0	9.5
West	9.3	9.8	13.4	12.7	12.7	14.0	13.0	11.1	9.3	10.5†	10.1	8.4	7.9	8.2	7.7
Population Density:															
Large MSA	8.0	8.1	8.8	9.8	8.7	12.3	9.9	8.9	8.4	8.5†	9.7	7.4	7.1	6.9	7.1
Other MSA	8.6	10.4	11.8	12.5	13.5	14.1	12.2	11.2	10.7	10.1†	11.8	9.2	8.7	8.5	8.8
Non-MSA	8.6	8.9	9.8	9.8	13.2	12.1	13.0	12.8	12.8	12.3†	10.3	9.8	11.0	8.0	8.1
Parental Education: ^c															
1.0–2.0 (Low)	12.9	12.9	14.4	15.6	18.0	15.5	14.8	17.3	16.0	15.8†	14.3	13.0	14.1	15.2	14.3
2.5–3.0	8.5	10.1	11.8	12.4	14.2	13.9	12.9	12.2	12.1	12.2†	13.2	10.9	10.9	8.8	10.3
3.5–4.0	8.7	10.1	10.6	11.8	14.2	14.5	12.5	11.2	11.3	10.6†	11.7	9.0	9.0	8.4	9.0
4.5–5.0	7.1	7.5	9.1	9.5	9.7	12.0	10.6	9.4	8.5	7.7†	8.9	7.6	6.7	6.8	5.7
5.5–6.0 (High)	7.8	8.0	8.2	9.4	10.1	11.7	10.3	9.5	8.3	8.4†	8.0	6.5	6.8	5.9	5.3
Race/Ethnicity (2-year average): ^d															
White	—	9.0	10.0	10.8	12.6	13.9	13.5	12.5	11.5	11.1	10.6†	10.3 ^e	9.3	8.7	8.1
African American	—	4.9	5.0	5.9	5.7	5.3	4.7	4.0	4.1	3.8	3.9†	4.4 ^e	4.4	4.4	4.2
Hispanic	—	12.2	13.7	15.2	15.3	14.7	13.6	13.5	14.5	13.9	12.2†	11.9 ^e	10.8	10.4	9.9

Table continued on next page.

TABLE 4 (cont.)
ANY ILLICIT DRUG OTHER THAN MARIJUANA: ^a Trends in Annual Prevalence of Use
 by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021 change	
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700		
Total	7.7	7.0	7.4	7.0	7.1	6.4	5.5‡	6.3	6.4	6.3	5.4	5.8	6.1	6.5	7.7	4.6	-1.9* sss	
Gender:																		
Male	6.5	5.9	6.0	6.6	6.4	6.1	4.8‡	5.5	5.3	5.1	4.7	4.4	5.8	5.2	§	3.3	-1.9 sss	
Female	8.7	8.0	8.7	7.3	7.7	6.5	6.0‡	6.9	7.1	7.4	6.1	6.8	6.1	7.6	§	5.3	-2.3 sss	
College Plans:																		
None or under 4 years	17.7	18.1	20.3	18.2	16.4	17.2	13.0‡	14.6	14.7	11.0	11.5	10.5	11.6	12.5	§	7.2	-5.3 sss	
Complete 4 years	6.7	6.0	6.3	6.0	6.3	5.5	4.9‡	5.7	5.7	5.9	4.8	5.1	5.2	5.6	§	4.2	-1.4 ss	
Region:																		
Northeast	4.4	4.4	5.2	5.0	4.6	4.4	4.1‡	4.0	4.3	5.3	3.0	3.9	3.6	4.7	§	2.5	-2.2 ss	
Midwest	7.1	7.4	7.6	6.7	7.0	6.0	3.7‡	5.4	5.6	5.9	4.9	4.9	4.6	6.2	§	4.9	-1.3	
South	9.8	7.8	8.8	8.4	8.3	6.7	7.3‡	8.2	7.1	6.7	6.4	6.1	6.8	6.8	§	5.0	-1.7 ss	
West	7.4	6.9	6.7	6.3	7.0	7.8	5.4‡	5.9	7.5	6.6	5.9	7.3	8.2	7.7	§	5.2	-2.5 s	
Population Density:																		
Large MSA	6.6	6.1	6.8	5.5	5.7	5.7	5.1‡	5.7	5.7	5.8	4.9	6.0	7.1	5.9	§	3.9	-2.0 ss	
Other MSA	8.0	7.6	7.7	7.8	8.0	7.0	6.1‡	6.5	6.9	7.1	5.5	5.8	5.8	6.9	§	4.9	-2.0 ss	
Non-MSA	8.8	7.1	7.6	7.1	6.9	6.0	5.0‡	6.9	6.4	5.1	6.2	5.4	5.0	6.4	§	5.3	-1.1	
Parental Education: ^c																		
1.0–2.0 (Low)	12.3	11.9	13.6	12.4	11.5	10.7	8.3‡	10.6	9.4	9.8	8.5	9.7	8.3	9.0	§	6.4	-2.6	
2.5–3.0	9.4	8.3	9.7	8.7	8.8	9.1	7.2‡	7.1	7.6	7.0	6.1	6.8	7.4	9.3	§	5.1	-4.3 sss	
3.5–4.0	8.4	7.8	8.0	7.4	7.3	7.0	6.5‡	8.2	7.0	6.8	5.4	5.8	6.8	6.5	§	5.7	-0.8	
4.5–5.0	6.0	4.7	5.4	5.4	6.0	4.6	3.5‡	5.2	4.9	5.0	4.8	4.6	5.2	5.8	§	4.5	-1.3	
5.5–6.0 (High)	6.2	5.5	5.7	5.1	4.7	4.3	4.4‡	4.0	5.4	5.1	4.8	4.2	4.0	4.8	§	3.8	-1.1	
Race/Ethnicity (2-year average): ^d																		
White	8.1	7.6	7.6	7.5	7.2	6.7	5.6	4.9‡	5.4	5.7	5.2	4.8	5.0	5.5	§	6.5	+1.0	
African American	3.9	3.3	3.4	4.1	4.0	3.8	3.9	4.0‡	5.2	5.1	5.3	4.9	4.8	4.7	§	3.0	-1.8 s	
Hispanic	9.2	8.6	8.4	8.6	8.7	8.7	8.1	7.4‡	7.4	7.1	6.8	6.5	7.0	7.8	§	5.2	-2.5	

Table continued
on next page.

TABLE 4 (cont.)
**ANY ILLICIT DRUG OTHER THAN MARIJUANA:^a Trends in Annual Prevalence of Use
by Subgroups in Grade 8**

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. The 2013 data are based on half the forms; N is one half of N indicated. Beginning in 2014 data are based on all forms. See the amphetamine notes for details of the changes.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aUse of any illicit drug other than marijuana includes any use of LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of amphetamines or tranquilizers not under a doctor's orders. The use of narcotics other than heroin and sedatives (barbiturates) has been excluded because 8th and 10th graders appear to overreport their use (perhaps because they include the use of nonprescription drugs in their answers).

^bIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. For the tranquilizer list of examples, Miltown was replaced with Xanax. The 2001 data are based on the changed forms only; N is one half of N indicated. In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^dEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^eDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 5
ANY ILLICIT DRUG OTHER THAN MARIJUANA: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^o	2002	2003	2004	2005
Approximate weighted N =	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	12.2	12.3	13.9	15.2	17.5	18.4	18.2	16.6	16.7	16.7†	17.9	15.7	13.8	13.5	12.9
Gender:															
Male	11.2	11.1	13.4	14.1	15.8	17.2	17.2	15.6	15.9	16.7†	18.3	15.1	13.0	12.7	12.0
Female	13.1	13.2	14.3	16.0	18.9	19.6	19.1	17.5	17.3	16.6†	17.4	16.4	14.3	14.2	13.6
College Plans:															
None or under 4 years	19.6	20.2	23.1	24.0	27.5	29.5	29.6	27.8	27.3	27.7†	32.1	27.1	23.8	25.3	24.0
Complete 4 years	10.7	10.5	12.0	13.3	15.7	16.5	16.3	14.6	15.0	15.0†	15.5	14.0	12.1	11.9	11.4
Region:															
Northeast	10.6	9.6	12.8	13.7	14.1	17.2	16.0	17.2	18.2	14.7†	16.2	13.4	11.7	13.7	13.0
Midwest	13.2	12.9	12.8	14.8	19.0	20.0	16.2	14.4	16.1	15.8†	16.5	15.3	13.1	12.0	12.2
South	11.9	12.2	14.7	15.3	18.4	18.6	20.8	18.3	16.8	17.5†	19.5	16.8	15.7	14.8	13.9
West	12.7	14.1	15.6	17.2	17.2	17.4	18.7	15.8	15.7	18.5†	19.0	16.7	13.8	13.5	12.0
Population Density:															
Large MSA	11.8	11.4	12.2	13.1	13.5	16.8	16.3	14.6	15.0	17.2†	15.6	14.3	10.8	10.4	10.9
Other MSA	12.3	12.3	14.1	16.1	18.5	19.5	18.0	16.6	17.3	15.6†	17.4	16.5	14.2	15.1	13.2
Non-MSA	12.4	13.1	15.0	14.6	17.6	18.3	20.8	18.9	17.5	18.1†	21.5	16.1	17.2	14.5	14.7
Parental Education: ^c															
1.0–2.0 (Low)	14.4	16.6	18.1	17.1	20.8	22.7	19.1	21.5	19.2	20.4†	19.6	21.0	19.1	16.5	15.9
2.5–3.0	13.7	12.5	14.6	16.3	19.7	19.4	19.9	19.1	19.1	19.4†	20.3	18.3	16.7	15.6	14.4
3.5–4.0	12.1	12.7	14.8	15.9	18.3	19.9	19.8	16.4	16.5	17.4†	19.7	16.1	14.5	14.6	13.7
4.5–5.0	11.0	10.9	11.7	13.3	15.9	16.6	16.5	14.1	15.4	14.5†	15.6	13.7	11.2	11.6	11.5
5.5–6.0 (High)	11.6	10.7	12.2	12.8	13.4	15.4	15.4	14.4	15.6	14.5†	14.6	12.2	10.5	11.0	10.8
Race/Ethnicity (2-year average): ^d															
White	—	13.7	14.4	15.4	17.7	20.0	20.5	19.7	18.7	18.6	19.2†	18.9 ^e	17.2	15.7	14.9
African American	—	4.3	4.6	5.4	5.4	4.5	4.8	4.7	4.5	4.2	4.7†	5.7 ^e	4.7	4.6	4.8
Hispanic	—	11.8	13.7	16.1	16.9	18.8	19.1	17.5	17.9	17.8	15.8†	15.7 ^e	15.2	15.1	14.9

Table continued on next page.

TABLE 5 (cont.)
ANY ILLICIT DRUG OTHER THAN MARIJUANA: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months															2019-2021 change	
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	12.7	13.1	11.3	12.2	12.1	11.2	10.8‡	11.2	11.2	10.5	9.8	9.4	9.6	9.1	8.6	5.1	-4.0* sss
Gender:																	
Male	12.0	12.9	10.9	11.8	12.3	12.2	11.3‡	12.5	10.4	10.0	9.7	9.3	9.8	9.5	§	4.0	-5.5 sss
Female	13.1	13.2	11.5	12.3	11.8	10.0	10.4‡	9.9	11.8	10.7	9.7	9.1	9.2	8.6	§	5.4	-3.2 sss
College Plans:																	
None or under 4 years	24.7	25.0	22.9	25.4	24.5	24.1	24.5‡	17.2	21.9	19.7	18.4	17.7	18.2	13.9	§	8.3	-5.6 sss
Complete 4 years	11.1	11.5	9.8	10.7	10.7	9.7	9.4‡	10.5	10.1	9.5	8.8	8.3	8.3	8.1	§	4.2	-3.9 sss
Region:																	
Northeast	10.4	11.3	8.2	9.0	9.8	9.4	9.3‡	8.5	9.8	9.9	6.8	6.7	7.4	6.1	§	4.9	-1.2
Midwest	12.7	13.4	11.8	11.5	12.0	10.8	10.4‡	11.0	11.8	9.0	9.2	10.3	9.6	8.5	§	5.1	-3.4 sss
South	13.7	13.3	12.6	13.6	12.4	12.2	12.4‡	12.7	11.9	11.3	10.7	8.7	10.3	9.4	§	5.6	-3.9 ss
West	13.3	13.8	11.5	13.7	14.1	11.7	10.3‡	11.6	10.9	11.3	11.5	11.6	10.0	11.4	§	4.6	-6.7 sss
Population Density:																	
Large MSA	11.5	12.0	9.9	10.7	11.2	10.4	10.0‡	10.4	10.2	10.3	9.5	9.9	9.6	8.2	§	4.8	-3.3 sss
Other MSA	12.9	13.4	12.1	13.4	12.6	11.2	10.9‡	11.9	11.3	10.6	10.2	9.4	9.8	9.8	§	4.8	-5.0 sss
Non-MSA	14.1	14.0	11.8	11.7	12.7	12.6	11.9‡	10.6	12.7	10.6	9.2	8.7	9.0	8.8	§	6.4	-2.3 ss
Parental Education: ^c																	
1.0–2.0 (Low)	16.0	15.6	15.2	18.3	15.8	13.3	16.8‡	15.2	14.1	15.1	11.3	10.5	12.5	11.5	§	8.6	-2.9
2.5–3.0	14.9	15.2	13.2	13.6	14.8	13.1	11.7‡	13.5	12.4	12.8	11.1	11.2	10.9	11.4	§	6.3	-5.1 sss
3.5–4.0	13.4	14.2	12.7	13.0	12.9	12.0	10.9‡	11.6	13.2	12.3	11.0	9.5	9.8	9.7	§	5.9	-3.8 ss
4.5–5.0	10.7	11.6	9.0	9.9	10.2	9.6	9.7‡	9.9	9.2	8.9	9.0	8.5	8.2	7.3	§	4.0	-3.2 sss
5.5–6.0 (High)	11.1	9.9	8.6	10.6	9.4	9.2	9.1‡	9.1	9.0	7.4	8.3	8.5	9.1	7.4	§	3.8	-3.6 sss
Race/Ethnicity (2-year average): ^d																	
White	14.4	14.5	13.8	12.9	13.0	12.5	11.5	11.2‡	11.5	10.9	10.3	10.0	9.8	9.1	§	6.3	-2.8 sss
African American	4.9	4.2	4.1	4.3	5.0	5.5	5.8	6.5‡	8.4	7.8	7.8	6.3	5.2	5.4	§	5.7	+0.3
Hispanic	13.0	11.8	12.4	13.2	13.5	12.9	12.4‡	11.7	11.1	11.1	10.6	11.0	11.0	§	7.0	-4.0 ss	

Table continued
on next page.

TABLE 5 (cont.)
**ANY ILLICIT DRUG OTHER THAN MARIJUANA:^a Trends in Annual Prevalence of Use
by Subgroups in Grade 10**

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. The 2013 data are based on half the forms; *N* is one half of *N* indicated. Beginning in 2014 data are based on all forms. See the amphetamine notes for details of the changes.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aUse of any illicit drug other than marijuana includes any use of LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of amphetamines or tranquilizers not under a doctor's orders. The use of narcotics other than heroin and sedatives (barbiturates) has been excluded because 8th and 10th graders appear to overreport their use (perhaps because they include the use of nonprescription drugs in their answers).

^bIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. For the tranquilizer list of examples, Miltown was replaced with Xanax. The 2001 data are based on the changed forms only; *N* is one half of *N* indicated. In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^dEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^eDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 6
ANY ILLICIT DRUG OTHER THAN MARIJUANA: ^{a,b}
Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200	
Total	26.2	25.4	26.0	27.1	28.2	30.4	34.0	30.1	28.4	28.0	27.4	25.9	24.1	21.1	20.0	17.9	
Gender:																	
Male	25.9	25.7	26.3	27.9	29.4	30.2	32.8	31.0	28.9	28.2	27.9	26.2	24.3	22.2	21.0	19.2	
Female	26.2	24.4	25.3	25.7	26.3	30.0	34.3	28.3	27.3	26.9	26.2	24.8	23.3	19.3	18.5	16.0	
College Plans:																	
None or under 4 years	—	28.7	30.1	30.0	31.8	35.5	38.3	34.0	32.3	32.9	31.6	31.3	28.8	24.5	25.5	23.1	
Complete 4 years	—	20.9	20.8	22.7	23.5	25.5	30.1	26.0	24.7	23.3	24.1	22.2	21.3	19.0	17.2	15.2	
Region:																	
Northeast	26.0	26.1	27.8	30.7	32.0	32.1	38.0	33.5	31.2	33.8	32.9	29.5	25.5	20.2	19.2	17.1	
Midwest	29.2	26.1	27.7	26.8	27.6	30.9	36.1	31.1	28.6	26.1	25.9	25.1	22.7	20.3	21.5	18.0	
South	22.5	23.4	22.9	24.0	23.2	25.8	26.1	24.7	23.8	24.2	21.0	20.6	21.1	20.0	18.1	16.9	
West	28.2	26.6	26.0	28.8	33.3	35.2	38.7	32.7	33.0	31.3	33.0	31.6	29.5	24.8	22.3	20.4	
Population Density:																	
Large MSA	30.3	27.5	27.1	30.2	32.1	34.6	38.3	33.8	31.5	30.5	30.4	28.3	24.5	20.7	16.9	16.0	
Other MSA	26.3	25.8	26.8	27.3	28.7	30.1	33.3	30.0	29.7	27.8	26.9	26.4	24.5	22.7	20.9	18.5	
Non-MSA	23.4	23.3	24.2	24.2	24.7	27.5	31.4	27.0	24.4	26.2	25.5	23.1	23.0	18.4	21.1	18.4	
Parental Education: ^c																	
1.0–2.0 (Low)	—	23.2	23.2	24.7	25.2	28.2	29.2	25.7	25.6	27.3	25.8	23.2	21.5	19.7	18.2	15.2	
2.5–3.0	—	25.6	27.0	26.4	27.6	30.7	33.5	30.4	28.1	28.2	27.4	27.0	24.2	20.5	20.0	17.9	
3.5–4.0	—	26.1	26.2	27.8	29.2	30.7	34.7	30.9	28.6	29.3	28.9	26.6	24.7	20.5	21.4	19.1	
4.5–5.0	—	27.2	25.9	27.3	28.7	29.9	34.8	29.4	30.0	26.2	27.1	24.9	23.8	21.7	19.3	17.5	
5.5–6.0 (High)	—	25.6	24.8	28.6	30.4	30.8	36.7	31.3	29.0	26.2	23.8	23.8	24.9	22.0	19.6	17.2	
Race/Ethnicity (2-year average): ^d																	
White	—	—	26.6	27.7	28.8	30.6	34.5	32.1	31.2	30.2	29.6	28.2	26.6	24.4	22.5	21.0	
African American	—	—	14.2	13.4	13.0	13.8	13.2	14.5	15.2	12.9	12.0	12.1	11.1	10.3	8.6	6.5	
Hispanic	—	—	23.8	23.5	23.3	24.7	27.6	25.5	25.2	26.2	27.2	26.2	23.0	20.5	17.7	15.6	

Table continued on next page.

TABLE 6 (cont.)
ANY ILLICIT DRUG OTHER THAN MARIJUANA: ^{a,b}
Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^g	2002	2003	2004	2005	2006
Approximate weighted N =	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	16.2	14.9	17.1	18.0	19.4	19.8	20.7	20.2	20.7	20.4 [‡]	21.6	20.9	19.8	20.5	19.7	19.2
Gender:																
Male	17.0	15.5	17.8	18.5	20.7	21.7	21.7	21.7	22.5	21.5 [‡]	23.3	22.0	21.1	21.5	21.4	19.7
Female	14.8	13.8	15.8	16.9	17.3	16.8	18.8	18.0	18.5	18.6 [‡]	19.0	19.0	17.9	18.7	17.5	18.3
College Plans:																
None or under 4 years	20.1	19.5	19.8	22.9	23.9	24.2	25.8	26.5	24.4	24.7 [‡]	24.5	27.2	26.5	26.0	25.3	25.4
Complete 4 years	14.3	13.0	15.9	16.0	17.5	17.9	18.4	17.8	19.4	18.5 [‡]	19.9	19.0	17.4	18.4	18.0	17.4
Region:																
Northeast	15.6	14.7	18.6	17.2	20.2	22.9	24.1	20.7	19.5	21.7 [‡]	22.4	21.5	19.3	20.4	18.0	20.3
Midwest	17.4	15.5	16.4	20.1	19.1	19.2	18.9	19.8	20.9	18.9 [‡]	24.0	19.9	18.7	19.1	21.4	17.9
South	14.4	14.0	16.0	17.3	19.0	18.6	19.8	20.3	21.2	19.0 [‡]	18.6	21.3	19.4	20.8	20.5	19.7
West	17.9	15.8	18.5	17.3	19.9	19.2	20.9	20.0	20.8	23.2 [‡]	22.6	21.1	22.4	21.9	18.0	19.1
Population Density:																
Large MSA	14.2	13.5	15.1	16.7	20.2	18.9	18.6	19.0	17.6	19.4 [‡]	23.6	20.6	15.9	18.0	18.6	20.1
Other MSA	17.9	14.9	18.2	19.2	19.2	20.2	21.5	20.4	21.6	20.9 [‡]	21.2	21.6	22.0	22.3	20.6	18.7
Non-MSA	14.9	16.1	16.8	17.2	18.7	19.8	21.2	21.3	22.4	20.7 [‡]	20.3	20.2	20.9	19.7	19.3	18.9
Parental Education: ^c																
1.0–2.0 (Low)	17.4	14.9	15.6	17.8	19.4	16.9	19.9	20.0	22.2	20.0 [‡]	17.3	17.8	19.4	18.7	14.8	19.7
2.5–3.0	16.8	15.0	16.1	18.5	19.4	19.7	19.4	20.1	21.0	19.9 [‡]	20.9	20.3	19.6	20.0	20.5	19.4
3.5–4.0	16.3	15.0	17.8	17.6	19.2	19.8	20.4	21.0	21.9	20.5 [‡]	24.5	22.0	21.7	23.0	21.5	20.2
4.5–5.0	14.6	14.3	17.5	18.4	19.4	19.3	21.0	19.1	19.3	20.1 [‡]	20.1	21.2	18.5	19.7	19.7	18.6
5.5–6.0 (High)	14.9	14.3	17.6	16.5	18.3	20.2	21.7	18.9	19.4	20.3 [‡]	18.7	19.8	17.0	17.6	17.5	18.0
Race/Ethnicity (2-year average): ^d																
White	18.7	17.1	17.9	19.4	20.3	21.2	22.3	23.1	22.9	22.7	23.0 [‡]	24.1 ^f	23.0	22.8	22.7	22.0
African American	5.7	5.3	4.8	6.1	6.9	6.0	6.4	7.1	6.8	6.4	6.3 [‡]	6.0 ^f	6.3	6.5	7.1	7.2
Hispanic	15.8	15.1	15.6	16.5	17.9	19.7	18.9	17.5	18.5	21.2	18.2 [‡]	16.1 ^f	16.0	16.2	16.7	17.3

Table continued on next page.

TABLE 6 (cont.)
ANY ILLICIT DRUG OTHER THAN MARIJUANA: ^{a,b} Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months															2018-2019
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	18.5	18.3	17.0	17.3	17.6	17.0‡	17.8	15.9	15.2	14.3	13.3	12.4	11.5	11.4	7.2	-4.2* sss
Gender:																
Male	20.0	19.9	18.8	19.3	19.1	18.4‡	18.9	16.9	15.7	15.4	14.9	13.9	12.4	§	8.0	-4.5 sss
Female	16.7	16.2	14.9	15.0	15.5	14.9‡	15.9	14.2	14.1	12.5	11.3	9.7	9.7	§	6.0	-3.7 sss
College Plans:																
None or under 4 years	26.0	25.0	22.1	23.6	25.1	19.4‡	24.4	19.8	17.7	15.4	15.7	16.0	15.4	§	9.0	-6.4 sss
Complete 4 years	16.7	16.5	15.7	15.7	15.8	16.1‡	16.2	14.5	14.3	13.6	12.3	10.8	10.0	§	6.5	-3.5 sss
Region:																
Northeast	19.4	18.9	16.8	17.6	16.4	15.1‡	16.7	13.9	13.1	12.3	11.5	12.1	8.1	§	6.7	-1.4
Midwest	18.4	16.8	19.1	16.9	17.5	17.2‡	18.9	16.2	15.0	15.7	12.1	12.2	11.9	§	5.9	-6.0 sss
South	18.8	19.1	16.4	17.4	16.2	16.0‡	17.5	17.2	15.2	14.0	13.6	12.2	11.3	§	6.7	-4.6 sss
West	17.3	18.1	15.6	17.4	20.6	19.5‡	17.9	15.1	16.9	15.0	15.1	13.3	14.7	§	9.7	-5.0 s
Population Density:																
Large MSA	18.4	16.9	15.9	17.3	16.6	18.3‡	18.1	14.6	14.8	14.3	12.5	11.0	10.7	§	7.3	-3.3 ss
Other MSA	18.1	18.2	17.4	17.7	18.6	17.2‡	18.8	17.7	15.9	14.5	13.9	12.9	11.9	§	7.7	-4.2 sss
Non-MSA	19.5	19.8	17.4	16.3	16.7	14.7‡	15.0	13.0	14.1	13.8	13.4	14.0	11.8	§	6.0	-5.8 sss
Parental Education: ^c																
1.0–2.0 (Low)	16.7	16.4	14.7	14.9	16.0	16.5‡	15.9	14.7	16.5	13.0	10.7	11.4	10.5	§	5.7	-4.8 ss
2.5–3.0	19.7	17.7	17.3	17.7	16.6	18.0‡	18.6	16.8	14.4	13.2	14.6	12.3	11.7	§	7.0	-4.7 s
3.5–4.0	18.4	18.7	17.6	18.4	18.8	17.6‡	18.3	15.7	15.8	15.4	13.4	12.4	12.9	§	7.5	-5.4 sss
4.5–5.0	18.5	18.2	17.8	16.8	16.6	16.8‡	18.7	15.1	14.7	14.5	12.5	12.3	10.6	§	7.2	-3.4 ss
5.5–6.0 (High)	17.4	17.9	15.4	15.7	18.3	15.2‡	16.3	16.8	16.0	14.0	14.4	12.1	10.0	§	7.8	-2.2
Race/Ethnicity (2-year average): ^d																
White	21.6	21.5	20.5	19.9	20.0	19.6	18.7‡	17.6	16.1	15.5	14.8	13.9	12.5	§	8.6	-3.9 sss
African American	7.0	7.0	6.8	8.3	8.7	7.6	9.5‡	13.0	11.2	10.3	9.1	7.5	7.2	§	8.0	+0.8
Hispanic	15.8	13.7	13.6	12.8	12.8	13.6	14.7‡	15.1	14.0	13.7	12.7	10.8	10.1	§	8.7	-1.4

Table continued on next page.

TABLE 6 (cont.)
**ANY ILLICIT DRUG OTHER THAN MARIJUANA:^{a,b} Trends in Annual Prevalence of Use
by Subgroups in Grade 12**

Source. The Monitoring the Future study, the University of Michigan.

Notes. '‡' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. The 2013 data are based on half the forms; *N* is one half of *N* indicated. Beginning in 2014 data are based on all forms. See the amphetamine notes for details of the changes.

§ Insufficient data for 2020 estimate.

^aChanges across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aUse of any illicit drug other than marijuana includes any use of LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of narcotics other than heroin, amphetamines, sedatives (barbiturates), methaqualone (excluded since 1990), or tranquilizers not under a doctor's orders.

^bBeginning in 1982 the question about amphetamine use was revised to get respondents to exclude the inappropriate reporting of nonprescription amphetamines. The prevalence-of-use rate dropped slightly as a result of this methodological change.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^dEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^eIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens and shrooms was added to the list of examples. For the tranquilizer list of examples, Miltown was replaced with Xanax. The 2001 data are based on the changed forms only; *N* is one half of *N* indicated. In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^dDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 7
MARIJUANA: ^c Trends in Lifetime Prevalence of Use by Subgroups in Grade 8

	Percentage who ever used														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	10.2	11.2	12.6	16.7	19.9	23.1	22.6	22.2	22.0	20.3	20.4	19.2	17.5	16.3	16.5
Gender:															
Male	11.9	11.9	14.5	19.6	22.5	25.5	24.7	24.3	24.3	22.2	23.6	22.2	19.3	17.6	18.0
Female	8.3	10.3	10.8	13.7	17.2	20.5	20.2	19.6	19.8	18.1	17.2	16.2	15.6	14.9	14.8
College Plans:															
None or under 4 years	22.9	24.5	27.9	33.1	36.2	41.6	41.8	43.5	44.1	41.2	40.8	39.8	38.0	36.9	38.5
Complete 4 years	8.1	9.0	10.4	14.3	17.7	20.2	19.9	19.4	19.1	17.6	17.9	17.1	15.2	14.1	14.0
Region:															
Northeast	7.4	9.2	8.8	15.3	16.0	19.9	20.2	15.6	18.2	17.3	18.7	15.2	13.7	13.6	12.8
Midwest	9.9	9.4	11.5	15.6	21.7	23.1	21.3	23.7	24.6	21.0	18.3	17.9	16.8	15.4	17.6
South	10.8	11.8	12.5	14.9	19.4	22.1	22.7	23.9	23.0	20.5	22.4	22.1	20.0	18.0	18.5
West	12.4	15.1	18.7	22.7	22.4	27.4	25.9	23.1	20.6	21.5	20.7	19.5	17.5	16.8	15.0
Population Density:															
Large MSA	8.9	10.5	10.6	16.7	19.6	24.0	21.6	20.4	20.5	18.8	17.3	17.3	17.1	15.0	15.7
Other MSA	11.4	12.6	15.0	19.8	21.2	24.6	23.2	22.9	21.2	19.4	22.3	19.7	17.1	17.0	17.1
Non-MSA	9.3	9.4	10.1	11.0	18.1	19.1	22.4	22.7	25.6	23.8	20.3	20.7	18.9	16.7	16.3
Parental Education: ^a															
1.0–2.0 (Low)	20.6	19.9	18.6	24.9	28.7	26.6	31.6	30.6	35.1	32.7	31.8	31.2	31.1	25.7	25.3
2.5–3.0	11.9	13.6	14.6	18.1	23.0	25.9	25.7	26.0	28.3	25.0	25.3	25.5	22.2	21.7	23.0
3.5–4.0	10.1	10.4	13.3	16.8	21.4	25.2	25.0	23.9	21.7	20.7	22.0	19.8	18.3	17.0	17.5
4.5–5.0	6.6	7.9	10.0	13.8	16.3	20.1	19.8	17.9	16.2	14.8	15.3	14.9	12.7	12.7	11.7
5.5–6.0 (High)	6.7	7.3	8.7	14.1	14.9	18.8	15.5	16.8	15.2	14.6	12.8	11.8	11.3	10.5	9.8
Race (2-year average): ^b															
White	—	9.9	11.0	12.9	16.6	20.5	22.0	21.1	20.0	19.0	18.5	18.0	16.6	15.1	14.6
African American	—	7.4	9.3	13.2	17.1	19.7	21.7	23.0	23.9	23.8	22.2	20.2	19.7	19.9	20.5
Hispanic	—	19.0	20.3	23.3	26.3	27.2	28.6	29.6	29.7	27.1	26.2	27.4	26.1	23.7	20.5

Table continued on next page.

TABLE 7 (cont.)
MARIJUANA: ^c Trends in Lifetime Prevalence of Use by Subgroups in Grade 8

	Percentage who ever used																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	15.7	14.2	14.6	15.7	17.3	16.4	15.2	16.5	15.6	15.5	12.8	13.5	13.9	15.2	14.8	10.2	-5.0* sss
Gender:																	
Male	17.1	15.2	16.3	17.9	18.9	18.1	16.5	17.0	15.8	16.0	12.8	13.4	14.4	13.6	§	8.0	-5.6 sss
Female	14.0	12.8	12.7	13.4	15.8	14.1	13.6	15.6	14.7	14.6	12.8	13.3	13.3	16.4	§	11.1	-5.4 sss
College Plans:																	
None or under 4 years	35.4	35.3	35.0	32.2	36.8	35.5	32.9	34.3	29.2	30.0	27.8	26.3	26.9	27.0	§	18.4	-8.6 sss
Complete 4 years	13.6	12.2	12.7	14.3	15.7	14.8	13.8	15.2	14.4	14.1	11.5	12.0	12.1	13.5	§	8.7	-4.8 sss
Region:																	
Northeast	12.6	8.8	11.7	14.2	15.5	10.4	10.0	13.1	12.8	10.9	8.6	10.1	9.4	9.7	§	9.1	-0.6
Midwest	15.6	13.5	16.6	15.3	15.1	13.8	12.2	13.3	14.5	16.4	13.0	11.5	11.1	14.0	§	12.5	-1.4
South	18.4	16.6	15.8	17.2	19.9	17.8	17.6	19.4	15.8	15.9	14.0	13.7	13.5	16.7	§	9.0	-7.7 sss
West	13.6	14.0	12.9	14.9	16.6	21.5	18.3	17.4	18.1	16.7	13.7	17.3	20.5	17.8	§	10.7	-7.1 s
Population Density:																	
Large MSA	14.7	11.6	12.9	13.1	16.5	14.7	14.4	17.4	17.6	15.6	12.5	12.5	14.0	13.3	§	8.2	-5.1 s
Other MSA	16.2	15.3	15.9	17.1	18.3	17.8	15.8	15.0	14.7	15.9	12.4	14.6	14.0	15.9	§	11.6	-4.3 ss
Non-MSA	16.2	15.4	14.2	16.3	16.3	16.1	15.3	18.2	14.3	14.1	14.4	12.0	13.5	16.6	§	10.5	-6.1 s
Parental Education: ^a																	
1.0–2.0 (Low)	25.6	26.6	24.5	24.1	26.2	27.8	27.8	29.1	25.2	25.3	20.9	21.6	22.4	24.6	§	16.0	-8.6 ss
2.5–3.0	20.8	19.8	19.7	21.2	23.1	23.0	22.2	23.4	21.8	20.5	18.0	18.9	20.0	25.0	§	16.7	-8.3 sss
3.5–4.0	17.2	15.5	16.0	17.2	18.9	18.1	16.7	18.6	18.0	17.3	14.3	14.8	16.3	17.3	§	15.3	-2.0
4.5–5.0	11.1	9.2	11.0	11.4	13.3	11.7	10.6	11.6	10.4	9.6	10.0	9.2	9.0	10.2	§	8.0	-2.3 s
5.5–6.0 (High)	9.3	6.5	8.3	9.7	10.3	8.6	7.6	9.4	10.0	9.4	6.6	8.1	7.1	8.8	§	5.5	-3.3 s
Race (2-year average): ^b																	
White	14.1	12.9	12.5	13.3	14.4	14.3	12.7	11.7	11.4	10.9	10.0	9.5	9.9	10.3	§	11.3	+1.0
African American	19.5	16.4	15.9	18.1	19.9	18.9	17.2	18.9	18.8	17.6	17.2	17.0	15.6	16.7	§	15.1	-1.6
Hispanic	19.3	19.1	18.1	18.5	20.4	21.8	22.8	22.1	21.2	20.0	17.5	15.5	16.4	18.5	§	9.7	-8.8 sss

Source. The Monitoring the Future Study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '§' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 8
MARIJUANA: ^c Trends in Lifetime Prevalence of Use by Subgroups in Grade 10

	Percentage who ever used														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	23.4	21.4	24.4	30.4	34.1	39.8	42.3	39.6	40.9	40.3	40.1	38.7	36.4	35.1	34.1
Gender:															
Male	25.3	23.3	27.0	33.6	36.2	42.3	45.1	41.6	43.3	44.2	43.9	41.7	38.5	37.2	36.1
Female	21.4	19.3	21.7	26.8	31.6	37.3	39.3	37.6	38.4	36.3	36.4	35.8	34.1	32.9	31.8
College Plans:															
None or under 4 years	37.7	34.6	39.4	44.9	49.5	55.9	60.6	56.1	59.6	58.1	56.7	57.5	53.3	51.7	50.5
Complete 4 years	20.2	18.4	21.1	27.1	31.4	37.0	39.2	36.5	37.8	37.5	37.4	35.7	33.5	32.8	31.7
Region:															
Northeast	22.3	19.3	26.5	30.0	33.1	39.6	41.2	41.9	41.0	37.4	38.8	38.2	36.1	36.0	35.0
Midwest	23.2	20.8	21.7	27.4	30.8	38.0	41.3	35.7	40.3	38.0	38.8	36.6	32.3	32.1	33.4
South	21.3	19.4	23.1	29.8	34.9	41.3	41.9	39.8	39.6	39.8	39.6	38.0	38.9	37.4	35.6
West	27.4	27.4	29.4	36.7	38.4	39.4	45.7	41.4	43.6	46.9	45.2	43.2	37.8	34.8	31.5
Population Density:															
Large MSA	23.6	20.5	23.5	30.9	32.4	38.3	42.2	36.8	40.3	39.9	36.4	37.0	35.6	33.6	33.2
Other MSA	24.9	22.0	25.5	33.6	37.2	42.5	44.1	41.7	42.0	41.2	42.4	40.9	36.5	35.1	34.7
Non-MSA	20.6	21.1	23.2	24.0	30.0	36.2	39.1	38.8	39.2	38.9	39.8	36.4	37.2	37.0	33.9
Parental Education: ^a															
1.0–2.0 (Low)	30.3	28.7	32.0	35.8	39.3	40.8	46.2	43.2	49.0	46.1	44.0	45.9	43.1	45.6	40.1
2.5–3.0	26.2	22.9	26.0	32.8	38.0	42.3	45.7	45.2	45.2	44.3	45.7	44.8	41.9	40.3	40.0
3.5–4.0	23.4	21.4	24.6	30.8	35.5	42.4	44.8	40.1	40.9	44.0	40.6	38.1	37.9	37.6	34.9
4.5–5.0	20.3	18.8	20.4	27.0	31.7	37.4	38.9	35.0	36.8	35.7	37.3	35.2	31.2	30.0	30.4
5.5–6.0 (High)	20.0	17.7	21.9	25.9	26.8	34.7	35.4	33.8	36.2	32.7	33.8	31.4	29.7	27.5	28.2
Race (2-year average): ^b															
White	—	23.0	23.2	27.2	32.3	37.1	41.1	40.9	39.9	40.1	39.7	39.3	37.4	35.1	33.7
African American	—	16.2	16.2	22.1	28.6	34.3	38.1	37.1	37.3	39.5	40.1	38.1	36.9	38.0	38.3
Hispanic	—	26.9	28.8	33.5	37.5	42.2	46.5	46.6	46.5	46.0	44.9	41.9	39.4	39.9	38.8

Table continued on next page.

TABLE 8 (cont.)
MARIJUANA:^c Trends in Lifetime Prevalence of Use by Subgroups in Grade 10

	Percentage who ever used																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	31.8	31.0	29.9	32.3	33.4	34.5	33.8	35.8	33.7	31.1	29.7	30.7	32.6	34.0	33.3	22.0	-12.0* sss
Gender:																	
Male	32.9	33.5	31.7	35.8	36.9	37.8	37.1	38.9	35.2	31.7	29.2	29.9	32.4	32.9	§	20.5	-12.4 sss
Female	30.6	28.4	28.0	29.0	29.8	31.2	30.6	32.6	31.9	30.2	29.8	31.0	32.5	34.8	§	21.7	-13.2 sss
College Plans:																	
None or under 4 years	51.6	50.3	51.7	52.8	54.5	54.1	51.3	53.3	50.5	48.3	44.0	42.5	46.8	46.6	§	34.8	-11.8 sss
Complete 4 years	29.2	28.3	27.0	30.0	30.9	32.2	31.9	33.9	31.8	29.3	28.1	28.9	30.4	31.6	§	18.8	-12.8 sss
Region:																	
Northeast	30.6	30.6	31.5	29.4	30.8	33.7	35.7	32.7	31.8	31.3	28.2	30.1	34.1	35.2	§	23.1	-12.1 sss
Midwest	30.7	29.9	27.8	28.5	31.1	31.9	28.6	32.8	30.0	26.9	28.4	29.6	29.2	30.9	§	23.4	-7.5 s
South	33.0	30.7	31.1	35.0	35.2	35.7	34.5	38.6	34.6	31.9	30.3	29.6	32.0	34.2	§	22.4	-11.9 sss
West	32.3	32.7	29.0	34.9	35.4	36.4	36.7	37.8	37.4	33.9	31.4	33.9	35.8	36.1	§	19.4	-16.7 sss
Population Density:																	
Large MSA	29.6	31.0	30.1	31.4	31.2	32.5	33.8	34.1	33.3	30.7	29.2	31.2	34.8	33.4	§	17.1	-16.2 sss
Other MSA	31.7	30.9	30.2	33.4	35.3	36.5	34.1	37.5	34.4	31.3	29.1	30.9	33.4	35.7	§	23.8	-12.0 sss
Non-MSA	35.3	31.0	28.6	31.2	32.7	33.7	32.9	34.4	32.5	31.1	32.4	29.5	27.2	30.3	§	26.2	-4.1
Parental Education: ^a																	
1.0–2.0 (Low)	41.7	37.4	39.7	43.8	43.6	44.4	47.5	45.3	40.9	42.6	34.7	35.8	38.3	40.8	§	31.4	-9.5 s
2.5–3.0	39.1	37.8	35.8	37.2	40.4	41.6	39.8	43.7	41.2	38.8	37.0	36.4	39.2	43.1	§	27.2	-15.9 sss
3.5–4.0	33.0	32.8	32.3	33.9	35.5	36.9	35.6	38.1	38.0	35.3	31.7	33.1	34.9	36.9	§	26.4	-10.5 sss
4.5–5.0	26.7	26.3	24.0	28.0	26.6	28.7	30.4	31.5	27.9	26.2	26.1	27.4	28.3	28.8	§	19.3	-9.6 sss
5.5–6.0 (High)	25.0	22.2	21.8	25.1	27.0	25.9	26.2	26.5	25.4	22.3	24.4	24.8	26.1	26.2	§	15.0	-11.2 sss
Race (2-year average): ^b																	
White	32.4	31.2	29.6	29.7	31.3	32.1	31.8	31.7	31.6	29.8	28.9	28.7	28.8	29.8	§	27.7	-2.1
African American	36.0	32.9	30.0	29.7	32.6	37.0	38.3	38.7	38.3	35.6	33.0	31.7	32.4	34.9	§	32.6	-2.2
Hispanic	34.7	32.1	34.8	37.8	38.9	38.9	40.4	42.9	42.1	39.3	35.2	33.5	35.3	36.8	§	31.4	-5.4

Source. The Monitoring the Future Study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 9
MARIJUANA: ^c Trends in Lifetime Prevalence of Use by Subgroups in Grade 12

Percentage who ever used

	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	47.3	52.8	56.4	59.2	60.4	60.3	59.5	58.7	57.0	54.9	54.2	50.9	50.2	47.2	43.7	40.7
Gender:																
Male	52.7	58.9	61.9	64.4	65.0	64.4	62.5	61.5	59.9	57.9	56.6	53.3	52.0	49.8	46.6	43.6
Female	42.7	46.1	50.8	53.9	55.7	56.1	56.2	55.5	53.4	51.3	51.5	48.2	48.0	44.5	40.4	37.3
College Plans:																
None or under 4 years	—	55.3	59.6	61.4	62.9	64.3	63.5	63.8	61.2	60.7	59.1	56.0	57.0	53.6	51.5	48.0
Complete 4 years	—	48.7	52.0	55.5	56.8	56.8	55.9	54.0	52.2	49.7	50.2	47.5	46.4	44.0	39.7	37.2
Region:																
Northeast	56.3	60.7	62.5	66.7	69.8	67.4	67.8	64.6	63.7	64.0	62.2	56.8	55.7	49.6	45.4	44.5
Midwest	46.9	52.1	56.0	60.6	60.9	60.2	59.9	59.8	57.0	52.3	53.8	52.0	50.1	48.0	46.8	42.6
South	38.8	45.7	51.4	52.4	51.6	53.6	50.8	51.1	50.8	49.1	44.5	43.7	43.6	42.4	38.2	35.4
West	52.5	55.9	57.1	59.0	62.1	62.9	63.2	61.7	59.2	59.8	60.4	53.8	55.1	52.0	48.4	43.0
Population Density:																
Large MSA	58.1	60.1	62.5	66.2	68.5	67.9	65.9	64.0	62.3	59.3	59.2	55.8	53.2	47.8	41.4	40.8
Other MSA	48.1	52.3	57.7	60.2	62.0	61.0	59.6	59.6	58.8	56.0	54.6	51.5	52.0	49.7	45.0	42.5
Non-MSA	39.6	47.8	49.7	51.9	52.1	53.9	54.6	53.1	50.5	49.8	49.3	45.9	43.5	41.9	43.2	36.9
Parental Education: ^a																
1.0–2.0 (Low)	42.3	47.8	50.1	53.1	56.5	57.3	56.5	55.4	56.2	52.6	52.9	47.0	46.1	48.0	39.5	38.1
2.5–3.0	47.0	54.4	56.9	59.1	60.1	61.0	59.0	58.8	56.7	55.6	54.5	52.1	51.6	47.0	45.5	42.2
3.5–4.0	45.6	53.2	58.8	60.5	61.9	60.8	60.0	60.7	56.6	56.3	55.6	51.7	51.0	47.9	45.6	42.1
4.5–5.0	48.5	55.5	57.6	61.6	61.8	60.9	60.9	59.8	58.0	53.2	54.5	50.7	49.7	47.7	42.2	40.4
5.5–6.0 (High)	46.9	57.3	56.9	63.3	61.5	62.0	61.6	58.4	56.7	53.7	50.0	49.5	49.5	45.0	40.7	37.9
Race (2-year average): ^b																
White	—	—	55.1	58.7	60.6	61.3	61.3	60.7	58.8	56.5	55.5	53.8	52.2	51.1	48.3	45.4
African American	—	—	48.5	50.8	50.7	51.4	51.1	51.4	53.9	52.6	49.7	46.9	42.6	38.8	33.4	27.7
Hispanic	—	—	55.6	54.7	55.3	56.7	53.9	53.9	54.4	54.5	53.4	51.0	47.7	46.0	42.8	38.5

Table continued on next page.

TABLE 9 (cont.)
MARIJUANA: ^c Trends in Lifetime Prevalence of Use by Subgroups in Grade 12

Percentage who ever used

	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700
Total	36.7	32.6	35.3	38.2	41.7	44.9	49.6	49.1	49.7	48.8	49.0	47.8	46.1	45.7	44.8
Gender:															
Male	40.3	36.3	38.9	42.8	45.2	48.6	52.0	53.2	52.1	51.5	51.9	50.7	50.1	48.5	49.1
Female	32.8	28.6	31.2	33.7	37.7	40.7	46.6	44.6	47.0	45.7	45.7	44.7	42.0	42.6	40.2
College Plans:															
None or under 4 years	43.6	41.8	41.7	44.6	48.8	53.1	56.5	55.6	57.2	55.2	55.9	55.5	55.6	52.6	54.7
Complete 4 years	33.8	28.8	32.7	35.6	38.8	41.8	47.2	46.4	47.0	45.9	46.6	45.6	42.9	43.5	42.1
Region:															
Northeast	40.4	34.5	41.1	44.3	44.5	48.9	54.3	54.1	50.8	52.5	54.9	52.4	50.0	51.2	50.4
Midwest	39.3	32.1	34.5	37.3	43.3	45.4	47.0	45.1	48.1	46.9	48.2	48.2	48.2	45.6	44.0
South	31.2	29.9	32.9	36.1	39.2	41.9	46.3	48.3	48.8	43.6	44.9	44.6	40.9	42.0	43.1
West	39.3	36.8	35.7	37.7	41.5	45.3	54.6	51.0	52.3	56.1	51.2	48.3	48.6	46.5	43.4
Population Density:															
Large MSA	36.1	32.5	36.5	38.6	44.1	46.4	48.9	50.7	50.3	49.1	50.1	48.1	43.4	42.8	45.5
Other MSA	41.4	32.6	36.3	40.7	42.1	47.8	52.6	50.3	50.6	50.2	49.1	50.5	49.5	49.3	45.9
Non-MSA	29.4	32.7	32.7	33.2	38.3	38.2	44.5	44.5	47.4	45.9	47.5	42.2	43.5	42.2	41.8
Parental Education: ^a															
1.0–2.0 (Low)	38.0	36.6	34.6	35.4	39.8	42.1	46.0	45.6	50.5	51.9	48.8	42.8	44.5	43.1	39.6
2.5–3.0	36.8	32.9	34.8	38.5	41.8	44.8	49.7	50.2	51.5	50.4	50.5	49.4	47.8	47.3	47.8
3.5–4.0	36.8	33.2	36.7	39.3	41.7	45.0	49.3	50.5	51.2	49.3	50.4	50.1	47.1	49.0	47.5
4.5–5.0	35.5	30.0	34.3	38.3	40.7	45.5	49.7	48.0	46.2	45.8	47.9	47.1	45.4	43.3	43.4
5.5–6.0 (High)	38.0	29.6	34.9	37.6	43.2	44.2	51.1	47.8	47.6	47.2	45.1	45.4	41.9	42.8	40.7
Race (2-year average): ^b															
White	41.3	36.3	35.6	38.3	41.1	43.9	48.3	50.9	50.3	49.4	49.8	49.9	48.8	48.2	47.5
African American	24.9	23.0	24.1	29.4	34.7	39.0	41.3	42.7	45.1	45.7	43.5	41.7	40.0	39.1	39.0
Hispanic	39.9	40.5	36.8	36.6	39.8	43.5	48.6	50.2	51.1	55.0	52.0	47.6	44.2	42.7	42.8

Table continued on next page.

TABLE 9 (cont.)
MARIJUANA: ^c Trends in Lifetime Prevalence of Use by Subgroups in Grade 12

	Percentage who ever used															2019-2021 change	
Approximate weighted N =	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Total	42.3	41.8	42.6	42.0	43.8	45.5	45.2	45.5	44.4	44.7	44.5	45.0	43.6	43.7	43.7	38.6	-5.2* ss
Gender:																	
Male	43.7	44.8	44.8	45.4	47.1	48.7	49.2	49.1	46.2	46.2	45.5	45.7	43.5	43.7	§	36.4	-7.3 sss
Female	40.6	38.6	40.1	38.1	40.0	41.8	40.5	41.3	42.3	43.3	43.3	43.9	42.8	42.9	§	40.0	-2.9
College Plans:																	
None or under 4 years	53.3	49.8	54.2	50.4	54.7	52.6	53.6	54.1	52.0	51.0	48.2	49.0	48.9	49.3	§	43.0	-6.3 s
Complete 4 years	39.2	39.5	39.6	39.8	41.0	43.5	43.0	43.1	42.0	43.1	43.0	43.6	41.8	41.6	§	37.0	-4.5 s
Region:																	
Northeast	48.5	48.9	46.7	45.4	50.3	48.8	47.3	49.5	46.7	49.2	46.6	48.0	48.3	44.5	§	45.1	+0.7
Midwest	42.5	42.5	41.3	42.1	40.8	43.0	39.4	38.8	43.5	41.0	41.8	46.1	45.0	41.6	§	38.4	-3.2
South	41.0	40.0	42.0	40.2	41.6	42.8	43.5	45.0	43.2	41.3	42.0	42.0	38.6	42.4	§	31.8	-10.6 sss
West	38.9	37.9	42.0	41.7	45.1	50.1	52.1	50.0	45.1	50.5	50.0	46.8	47.3	49.1	§	44.4	-4.8
Population Density:																	
Large MSA	44.4	43.6	44.4	45.6	45.4	47.9	48.1	48.5	44.1	47.2	46.5	47.4	44.4	44.5	§	43.8	-0.8
Other MSA	42.3	41.2	41.2	40.4	45.1	46.3	45.4	46.5	46.6	44.5	44.9	44.1	42.9	43.2	§	36.9	-6.4 s
Non-MSA	39.5	40.2	43.8	40.8	38.2	40.3	40.1	38.5	39.2	40.8	40.8	43.2	43.8	43.6	§	33.8	-9.8 ss
Parental Education: ^a																	
1.0–2.0 (Low)	46.8	44.6	42.6	42.7	44.5	48.3	52.6	50.8	48.4	48.8	47.7	43.2	42.9	45.6	§	33.3	-12.3 ss
2.5–3.0	46.1	45.0	46.3	44.2	45.6	48.6	48.2	49.0	48.9	48.6	46.5	48.5	46.9	46.5	§	43.0	-3.5
3.5–4.0	43.3	41.5	42.9	44.0	45.6	47.3	46.3	47.9	45.5	46.8	47.4	47.1	46.3	47.8	§	41.9	-5.9 s
4.5–5.0	37.1	39.4	40.1	38.7	42.8	42.2	42.7	41.4	41.3	41.1	41.8	43.5	42.0	37.2	§	37.2	0.0
5.5–6.0 (High)	40.5	39.0	40.2	38.4	36.9	40.9	38.6	39.0	39.0	39.5	37.9	40.5	38.6	41.7	§	36.3	-5.4
Race (2-year average): ^b																	
White	45.3	43.3	43.4	43.3	43.3	44.8	45.2	44.0	43.7	43.8	43.6	44.3	44.2	42.7	§	39.5	-3.2
African American	38.9	37.7	38.0	40.0	41.0	41.8	42.1	43.8	43.7	41.1	41.8	43.7	43.7	43.0	§	41.7	-1.3
Hispanic	43.1	41.3	40.1	40.7	42.3	45.6	48.7	50.7	48.7	48.2	50.3	47.7	43.7	43.6	§	40.4	-3.1

Source. The Monitoring the Future Study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 10
MARIJUANA: ^c Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	6.2	7.2	9.2	13.0	15.8	18.3	17.7	16.9	16.5	15.6	15.4	14.6	12.8	11.8	12.2
Gender:															
Male	7.3	7.4	10.5	15.1	17.7	19.6	19.2	18.0	18.1	16.7	18.1	17.0	13.9	12.3	13.0
Female	5.1	6.9	8.0	10.9	13.7	16.9	16.1	15.3	14.9	14.3	12.8	12.4	11.5	11.2	11.4
College Plans:															
None or under 4 years	15.8	17.5	22.4	27.7	30.3	34.6	34.5	35.0	34.9	33.6	34.4	33.1	31.3	29.0	30.5
Complete 4 years	4.6	5.5	7.3	11.0	13.8	15.8	15.5	14.5	14.0	13.4	13.2	12.7	10.7	10.0	10.2
Region:															
Northeast	5.0	5.8	6.2	12.1	13.0	15.3	16.2	11.7	14.4	13.2	14.5	11.3	10.0	10.2	9.8
Midwest	5.9	6.0	8.0	12.0	17.5	18.6	17.0	18.1	18.5	16.6	14.1	14.3	12.3	10.9	12.8
South	6.1	7.3	9.0	11.4	14.7	17.1	17.2	17.9	16.7	15.2	16.8	16.3	14.4	12.9	13.5
West	7.8	10.3	14.8	18.1	18.4	22.5	20.6	18.2	15.6	16.9	15.5	15.3	12.9	12.3	11.4
Population Density:															
Large MSA	5.2	6.7	8.0	13.1	15.6	18.3	16.4	16.0	15.6	14.5	12.4	12.7	11.5	10.8	11.7
Other MSA	7.2	8.3	10.9	15.7	17.2	19.5	18.2	17.4	15.4	14.8	17.4	15.2	12.9	12.4	12.9
Non-MSA	5.3	5.7	7.2	8.0	13.7	15.8	18.0	16.9	19.7	18.5	15.3	16.1	14.1	11.9	11.6
Parental Education: ^a															
1.0–2.0 (Low)	13.2	12.7	13.6	18.7	23.0	20.2	24.8	25.0	25.8	26.2	26.0	24.2	22.5	19.7	18.2
2.5–3.0	7.0	7.7	10.7	14.5	17.9	20.6	20.3	20.0	20.8	19.4	19.5	18.7	16.5	15.4	17.4
3.5–4.0	6.2	7.0	9.7	13.2	17.2	20.2	19.5	17.7	16.3	15.9	16.7	15.4	13.7	12.8	12.8
4.5–5.0	3.7	5.4	7.4	10.9	12.7	16.2	15.7	13.7	11.7	10.8	11.1	11.4	8.9	9.0	8.9
5.5–6.0 (High)	4.6	5.2	6.4	11.0	13.0	14.7	12.1	12.7	12.4	11.5	9.4	9.7	8.0	7.7	7.1
Race/Ethnicity (2-year average): ^b															
White	—	6.4	7.8	10.0	13.5	16.7	17.8	16.7	15.4	14.9	14.7	14.5	13.0	11.5	11.1
African American	—	4.1	5.7	8.9	11.9	14.0	15.3	16.0	16.3	16.1	14.6	12.7	12.6	12.8	13.6
Hispanic	—	11.9	13.9	18.1	20.4	20.8	21.8	22.7	22.8	20.1	19.9	21.1	19.1	16.7	14.7

Table continued on next page.

TABLE 10 (cont.)
MARIJUANA: ^c Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021 change
<i>Approximate weighted N =</i>	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Total	11.7	10.3	10.9	11.8	13.7	12.5	11.4	12.7	11.7	11.8	9.4	10.1	10.5	11.8	11.4	7.1	-4.7* sss
Gender:																	
Male	12.7	11.2	12.2	13.5	15.0	14.0	12.2	12.6	11.9	11.9	9.3	9.6	10.6	10.2	§	5.2	-5.0 sss
Female	10.4	9.3	9.5	10.0	12.4	10.6	10.4	12.5	11.0	11.4	9.3	10.3	10.2	13.0	§	8.0	-5.0 sss
College Plans:																	
None or under 4 years	28.8	28.1	27.7	26.7	30.7	27.6	28.4	27.8	24.1	24.3	22.2	21.1	21.6	21.9	§	12.4	-9.5 sss
Complete 4 years	10.0	8.6	9.4	10.5	12.3	11.2	10.1	11.6	10.6	10.6	8.2	8.8	8.9	10.3	§	6.1	-4.2 sss
Region:																	
Northeast	9.3	6.4	8.3	11.9	12.2	7.8	7.9	10.1	10.3	8.7	6.0	7.8	7.4	8.2	§	6.9	-1.3
Midwest	11.0	9.8	12.6	11.1	12.1	10.7	8.5	10.5	10.7	13.1	10.0	8.7	8.4	10.9	§	7.8	-3.1 ss
South	13.8	11.9	11.5	12.5	15.2	13.3	13.2	14.5	11.5	11.6	9.9	9.8	9.7	12.0	§	6.3	-5.7 sss
West	10.6	10.5	10.3	11.2	13.7	16.8	14.1	13.7	14.0	12.9	10.2	13.6	16.0	14.6	§	7.9	-6.7 s
Population Density:																	
Large MSA	10.3	8.7	9.9	10.1	13.5	11.4	10.7	13.5	13.6	12.0	9.1	9.5	10.8	9.9	§	5.8	-4.1 s
Other MSA	12.5	10.8	11.8	12.8	14.3	14.0	12.3	11.7	10.8	12.2	8.9	10.9	10.7	12.6	§	8.0	-4.5 ss
Non-MSA	12.0	11.4	10.4	11.8	12.3	11.2	10.8	13.6	10.9	10.4	10.9	9.2	9.5	12.8	§	7.1	-5.7 s
Parental Education: ^a																	
1.0–2.0 (Low)	19.4	19.4	18.5	17.6	20.7	20.2	19.2	21.9	19.3	19.0	14.3	16.0	15.9	18.8	§	10.9	-7.9 s
2.5–3.0	15.1	13.8	14.6	15.8	18.4	17.5	16.3	18.3	16.7	15.8	13.0	14.7	15.1	19.5	§	12.3	-7.2 sss
3.5–4.0	13.1	11.3	11.8	13.0	14.7	14.0	13.0	15.1	12.9	12.9	10.7	10.6	12.5	13.4	§	10.5	-2.8
4.5–5.0	8.0	7.2	8.2	8.7	10.6	9.1	8.2	8.6	8.0	7.6	7.4	6.7	7.0	8.0	§	5.0	-3.0 ss
5.5–6.0 (High)	7.1	4.9	6.7	7.4	8.1	6.6	5.8	6.6	7.9	7.3	4.7	6.7	5.2	6.8	§	4.1	-2.7 s
Race/Ethnicity (2-year average): ^b																	
White	10.9	9.9	9.6	10.3	11.5	11.4	10.0	9.1	8.9	8.7	7.8	7.3	7.5	8.0	§	8.4	+0.5
African American	12.9	11.1	10.6	12.3	14.1	13.6	12.4	13.6	13.2	11.9	11.7	11.5	10.7	11.6	§	10.7	-0.9
Hispanic	14.3	13.7	13.2	13.6	15.3	16.9	17.2	16.8	16.0	15.0	12.8	11.4	12.4	14.3	§	8.2	-6.0 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 11
MARIJUANA: ^c Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	
Approximate weighted N =	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
Total	16.5	15.2	19.2	25.2	28.7	33.6	34.8	31.1	32.1	32.2	32.7	30.3	28.2	27.5	26.6	
Gender:																
Male	17.7	16.3	21.2	28.2	30.6	36.0	37.3	32.2	34.3	36.1	36.0	32.3	30.0	29.0	28.1	
Female	15.1	13.9	16.9	21.9	26.5	31.4	32.3	30.1	29.7	28.4	29.6	28.4	26.4	25.8	24.9	
College Plans:																
None or under 4 years	26.9	25.1	31.5	37.3	41.8	48.9	51.5	46.8	48.3	48.8	47.4	46.6	44.6	41.9	41.4	
Complete 4 years	14.2	13.0	16.5	22.4	26.4	31.0	32.0	28.2	29.3	29.7	30.3	27.7	25.5	25.4	24.6	
Region:																
Northeast	17.1	14.9	22.4	25.6	28.8	34.8	34.6	35.4	34.4	30.3	34.1	31.7	29.5	29.2	28.7	
Midwest	15.8	14.8	17.4	23.4	26.6	33.1	34.4	28.5	31.6	31.1	31.7	29.0	25.0	25.3	26.4	
South	14.5	12.5	16.4	23.8	28.4	33.9	34.4	30.7	30.9	31.4	31.2	28.9	29.6	29.0	27.4	
West	19.4	20.4	24.0	30.0	32.2	32.4	36.5	30.7	32.0	37.1	36.4	33.4	28.8	26.4	23.4	
Population Density:																
Large MSA	16.5	15.1	19.0	26.3	27.8	31.5	34.1	28.7	31.2	32.4	30.0	29.0	27.1	25.9	25.3	
Other MSA	17.3	15.9	19.8	28.2	31.2	36.2	36.6	33.1	33.6	32.8	35.2	32.4	28.7	27.8	28.0	
Non-MSA	14.9	13.9	18.2	18.5	24.8	30.9	32.5	30.2	30.0	31.1	30.9	27.6	29.0	29.0	25.5	
Parental Education: ^a																
1.0–2.0 (Low)	20.3	18.9	22.4	25.8	32.0	32.9	34.5	31.7	38.1	37.1	33.6	33.5	33.7	33.2	28.6	
2.5–3.0	17.8	16.0	19.7	26.3	31.8	35.6	36.8	35.3	35.4	34.9	37.1	35.1	32.2	30.6	31.1	
3.5–4.0	16.2	15.1	19.3	25.6	30.0	36.4	37.8	31.6	31.9	35.0	32.6	30.1	29.4	29.5	27.1	
4.5–5.0	14.9	14.1	17.6	23.8	27.0	31.7	33.1	28.3	28.8	28.9	31.4	27.9	24.3	24.3	24.5	
5.5–6.0 (High)	15.9	13.7	18.5	23.3	23.4	30.3	30.5	27.7	30.6	27.3	29.4	25.8	24.3	22.5	23.6	
Race/Ethnicity (2-year average): ^b																
White	—	17.0	18.0	22.6	27.7	32.0	35.3	34.2	32.5	32.6	32.9	32.7	30.6	28.2	27.3	
African American	—	7.6	8.7	15.3	20.9	25.7	28.4	26.9	26.3	27.6	28.7	26.5	25.1	27.0	27.2	
Hispanic	—	18.9	21.3	25.1	29.2	34.6	36.8	34.4	34.0	34.8	34.9	31.6	28.8	29.8	28.6	

Table continued on next page.

TABLE 11 (cont.)
MARIJUANA: ^c Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																2018-2019 change
<i>Approximate weighted N =</i>	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	25.2	24.6	23.9	26.7	27.5	28.8	28.0	29.8	27.3	25.4	23.9	25.5	27.5	28.8	28.0	17.3	-11.5* sss
Gender:																	
Male	26.3	26.8	25.5	29.6	30.7	31.8	31.0	32.7	28.3	25.9	23.3	24.2	27.2	27.6	§	16.2	-11.3 sss
Female	24.0	22.4	22.2	23.9	24.3	25.8	25.3	27.0	26.0	24.7	24.0	26.5	27.6	29.7	§	17.0	-12.7 sss
College Plans:																	
None or under 4 years	42.0	42.1	43.5	44.5	45.8	47.9	43.6	46.5	41.4	40.8	37.2	35.5	40.3	40.2	§	28.7	-11.5 sss
Complete 4 years	23.0	22.2	21.4	24.7	25.3	26.7	26.5	28.0	25.7	23.8	22.4	24.0	25.6	26.6	§	14.4	-12.3 sss
Region:																	
Northeast	24.9	25.0	26.3	24.1	26.2	29.3	31.4	28.7	27.4	26.7	24.1	25.9	29.8	30.7	§	19.7	-11.0 sss
Midwest	24.8	24.9	22.6	23.6	25.7	26.8	23.7	27.5	24.5	21.5	23.0	25.7	25.0	26.5	§	19.4	-7.2 s
South	25.5	22.7	24.0	28.7	27.8	28.6	27.9	31.1	27.6	25.8	23.8	23.5	26.2	27.9	§	17.6	-10.3 sss
West	25.3	26.2	23.4	29.4	30.3	30.9	30.2	31.3	29.4	27.7	24.6	27.9	30.2	31.4	§	13.1	-18.3 sss
Population Density:																	
Large MSA	24.0	24.5	24.2	26.6	26.6	27.4	28.6	28.7	27.5	25.6	24.0	27.0	29.1	29.1	§	13.0	-16.0 sss
Other MSA	25.1	24.9	24.7	27.8	29.1	30.6	28.1	31.5	28.0	25.9	23.2	25.2	28.5	30.1	§	18.6	-11.5 sss
Non-MSA	27.4	23.9	21.7	24.5	25.7	27.2	27.0	27.3	25.1	23.8	25.4	23.7	22.4	24.8	§	21.6	-3.2
Parental Education: ^a																	
1.0–2.0 (Low)	31.9	28.4	30.2	36.0	34.8	35.8	37.1	36.9	31.5	34.2	26.6	28.8	31.2	32.3	§	24.3	-8.0 s
2.5–3.0	29.6	29.4	28.3	29.9	33.0	34.0	33.0	35.7	32.0	31.1	29.1	29.1	32.0	36.3	§	21.1	-15.2 sss
3.5–4.0	26.6	26.3	26.0	27.3	29.3	31.0	29.7	31.8	30.6	29.0	26.1	28.7	29.2	31.6	§	20.4	-11.2 sss
4.5–5.0	21.8	21.2	19.8	24.2	22.2	24.6	25.5	26.3	23.3	21.9	20.9	23.0	24.7	24.9	§	15.5	-9.5
5.5–6.0 (High)	21.2	18.9	18.9	21.8	23.1	23.1	22.9	23.2	21.6	18.6	20.7	21.3	23.6	22.9	§	12.8	-10.2 sss
Race/Ethnicity (2-year average): ^b																	
White	26.6	25.7	24.6	25.1	26.5	27.4	27.5	27.3	26.7	24.8	24.2	24.6	25.3	26.0	§	23.8	-2.2
African American	24.5	22.3	20.6	22.0	25.0	28.1	29.1	30.5	30.6	28.1	25.4	23.8	24.8	27.7	§	24.7	-3.0
Hispanic	24.9	23.7	26.6	29.9	31.1	31.0	31.8	33.4	32.6	30.6	27.2	26.0	28.2	29.9	§	24.2	-5.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 12
MARIJUANA: ^c Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	40.0	44.5	47.6	50.2	50.8	48.8	46.1	44.3	42.3	40.0	40.6	38.8	36.3	33.1	29.6	27.0
Gender:																
Male	45.8	50.6	53.2	55.9	55.8	53.4	49.2	47.2	45.7	43.2	43.1	41.2	38.6	35.8	32.8	29.4
Female	34.9	37.8	42.0	44.3	45.7	44.1	42.5	40.8	38.4	36.0	37.8	36.0	33.8	30.3	26.3	24.2
College Plans:																
None or under 4 years	—	46.8	50.7	51.6	53.1	51.7	49.7	48.2	46.0	44.2	44.0	42.7	40.6	36.2	34.4	31.1
Complete 4 years	—	40.7	43.4	47.1	47.3	45.9	42.6	40.6	38.3	35.9	37.5	36.1	34.0	31.3	27.3	24.7
Region:																
Northeast	47.4	52.7	53.5	59.2	60.6	55.5	53.2	50.9	49.3	49.6	48.2	44.6	41.2	36.7	31.3	32.2
Midwest	40.1	44.0	48.1	51.6	52.2	48.9	46.8	45.6	42.0	36.4	40.8	40.2	37.4	34.3	33.0	28.7
South	32.4	37.9	42.5	42.7	41.2	42.0	38.0	36.7	36.1	35.6	31.0	31.7	30.2	28.7	25.0	21.4
West	44.1	45.8	46.8	49.1	51.9	51.7	49.6	45.5	44.8	43.2	46.2	41.2	39.6	35.6	32.3	28.3
Population Density:																
Large MSA	50.4	51.3	53.2	57.2	58.7	56.3	51.4	50.4	47.0	44.2	44.4	42.6	39.3	34.3	27.8	27.7
Other MSA	40.3	44.2	48.9	50.8	51.9	49.8	46.4	44.8	44.0	41.0	40.7	39.4	36.9	34.7	30.3	28.3
Non-MSA	32.9	39.8	41.2	43.3	43.3	41.9	41.6	38.5	36.5	35.3	37.3	34.7	32.2	29.0	30.0	23.5
Parental Education: ^a																
1.0–2.0 (Low)	35.2	38.9	41.0	42.5	46.0	43.7	41.8	38.9	39.7	35.7	37.1	33.4	30.7	30.7	23.3	21.0
2.5–3.0	39.2	46.1	48.2	50.3	50.0	49.0	45.3	44.5	42.2	40.1	40.6	38.8	36.3	31.1	29.6	26.9
3.5–4.0	38.5	44.9	49.5	51.4	52.7	49.8	47.0	46.5	42.2	41.4	41.0	40.1	36.8	33.4	31.4	27.6
4.5–5.0	40.6	46.8	49.3	53.2	53.7	50.5	47.6	45.9	43.5	39.6	43.2	39.9	37.5	35.1	29.7	28.5
5.5–6.0 (High)	38.7	47.5	48.6	55.2	51.2	52.0	48.5	45.7	43.7	39.9	37.9	38.9	38.6	35.9	30.7	29.4
Race/Ethnicity (2-year average): ^b																
White	—	—	46.8	50.1	51.8	51.2	49.1	47.1	44.6	42.0	41.6	41.4	39.7	37.6	34.5	31.6
African American	—	—	37.9	39.6	38.4	37.5	36.1	35.5	37.4	36.4	33.4	30.6	25.7	21.2	17.8	13.7
Hispanic	—	—	45.8	43.4	42.1	44.1	41.2	38.8	38.3	38.8	37.8	36.7	33.3	29.6	25.0	21.6

Table continued on next page.

TABLE 12 (cont.)
MARIJUANA: ^c Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	23.9	21.9	26.0	30.7	34.7	35.8	38.5	37.5	37.8	36.5	37.0	36.2	34.9	34.3	33.6	31.5
Gender:																
Male	27.2	24.4	29.0	35.1	38.1	39.4	40.9	41.7	41.4	39.2	40.1	39.9	37.8	37.4	37.6	32.7
Female	20.1	18.9	22.4	26.4	30.6	31.6	35.5	33.0	34.1	33.4	33.6	32.4	31.6	30.8	29.6	29.7
College Plans:																
None or under 4 years	27.6	27.5	29.1	34.4	39.0	41.7	44.6	43.0	43.2	40.3	41.5	40.9	42.1	39.0	41.3	39.9
Complete 4 years	22.0	19.4	24.4	29.1	32.6	33.4	36.4	35.2	35.9	34.6	35.3	34.7	32.3	32.6	31.5	29.1
Region:																
Northeast	28.2	23.9	31.2	36.0	37.7	40.0	43.5	43.0	39.0	42.3	43.8	41.9	40.5	40.1	39.6	38.1
Midwest	26.1	22.7	26.0	30.5	36.9	36.9	36.5	33.8	38.0	34.5	36.9	37.5	36.5	34.0	33.7	32.0
South	18.1	18.1	23.2	28.7	31.8	32.8	35.0	36.5	36.0	30.7	32.4	32.6	29.4	30.1	31.2	29.4
West	26.8	26.1	26.4	30.0	33.8	35.6	42.6	39.0	39.8	43.1	38.4	35.2	36.2	35.6	32.0	28.8
Population Density:																
Large MSA	24.3	22.6	29.1	32.0	37.5	37.2	38.3	38.4	38.7	37.1	39.2	37.7	32.3	32.3	34.5	33.4
Other MSA	27.5	22.1	26.2	32.7	34.9	38.6	40.5	38.8	39.1	38.1	36.7	38.0	38.1	37.9	35.0	31.9
Non-MSA	17.5	21.0	23.1	25.8	31.0	29.6	34.9	33.5	34.7	32.9	35.1	30.8	32.2	29.7	29.7	28.1
Parental Education: ^a																
1.0–2.0 (Low)	22.4	21.2	23.0	26.3	30.9	31.3	34.8	34.2	35.5	36.5	33.9	30.8	31.4	28.7	27.3	32.0
2.5–3.0	22.5	21.1	24.1	29.7	33.8	35.1	37.4	36.1	38.6	35.1	37.0	34.9	34.2	33.1	34.8	33.2
3.5–4.0	24.0	22.7	26.6	31.5	34.2	36.1	38.1	39.0	38.6	36.8	38.4	38.5	36.2	36.9	35.5	31.6
4.5–5.0	23.8	20.8	27.2	32.0	35.0	36.6	40.1	37.4	35.9	35.8	37.4	37.0	35.4	34.4	33.9	28.4
5.5–6.0 (High)	28.2	22.6	28.0	32.3	37.5	36.7	39.7	38.3	39.2	38.0	35.2	36.1	32.6	34.0	32.3	32.9
Race/Ethnicity (2-year average): ^b																
White	28.2	24.9	25.9	30.2	34.2	36.4	38.7	39.9	39.1	38.2	38.5	38.7	37.9	37.3	36.6	34.8
African American	11.4	11.5	14.2	20.7	26.8	30.2	30.4	30.0	30.4	30.0	29.0	27.3	26.3	25.5	26.3	27.7
Hispanic	23.6	24.7	23.5	25.7	29.7	32.3	36.4	37.2	37.8	40.5	37.6	34.6	31.1	29.5	29.6	28.7

Table continued on next page.

TABLE 12 (cont.)
MARIJUANA: ^c Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2018-2019 change
	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	31.7	32.4	32.8	34.8	36.4	36.4	36.4	35.1	34.9	35.6	37.1	35.9	35.7	35.2	30.5	-5.2* ss
Gender:																
Male	35.0	35.1	36.7	38.3	40.0	40.8	39.8	37.0	36.3	36.9	37.4	36.8	35.2	§	27.8	-7.3 sss
Female	28.3	29.5	28.7	30.7	32.1	31.3	32.4	33.0	33.5	34.0	36.2	34.1	35.5	§	32.3	-3.2
College Plans:																
None or under 4 years	38.6	41.0	38.0	42.7	41.5	41.5	42.1	40.8	38.7	37.8	39.6	39.5	39.8	§	33.9	-5.9 s
Complete 4 years	29.8	30.2	31.1	32.5	34.7	34.8	34.8	33.1	33.7	34.5	36.1	34.5	34.0	§	29.1	-4.9 ss
Region:																
Northeast	39.2	36.6	37.8	41.3	40.7	38.1	40.9	39.6	41.0	39.0	41.5	41.1	37.6	§	36.8	-0.7
Midwest	32.6	31.1	32.5	32.5	34.7	32.0	31.6	34.6	32.7	33.4	38.8	38.2	34.4	§	30.4	-4.0
South	29.0	31.0	30.6	32.4	33.1	33.9	34.4	33.6	31.2	32.8	32.7	30.7	33.9	§	24.0	-10.0 sss
West	29.1	32.8	32.1	35.8	40.0	43.0	40.9	34.3	38.9	40.1	39.6	38.9	39.8	§	36.1	-3.7
Population Density:																
Large MSA	33.9	34.8	37.2	36.8	38.7	40.4	38.9	35.8	38.5	38.1	40.2	37.0	36.4	§	33.6	-2.8
Other MSA	31.6	31.2	32.0	36.1	37.4	36.8	37.9	36.9	34.4	35.7	36.0	35.3	35.8	§	30.4	-5.4 s
Non-MSA	28.8	32.2	28.7	28.8	30.4	29.2	29.0	29.7	30.4	31.7	34.2	35.5	34.4	§	25.7	-8.7 ss
Parental Education: ^a																
1.0–2.0 (Low)	31.1	30.5	28.9	32.8	35.7	42.0	38.4	37.3	35.3	35.0	34.6	33.1	34.5	§	23.7	-10.8 sss
2.5–3.0	33.8	32.8	33.6	35.4	36.8	37.6	39.0	37.5	37.8	36.7	39.7	38.9	37.6	§	32.5	-5.1
3.5–4.0	30.6	32.0	34.5	36.8	38.1	37.5	38.3	35.9	36.8	37.1	39.0	38.6	39.4	§	34.5	-4.9 s
4.5–5.0	30.4	32.7	31.0	34.5	35.0	35.0	33.3	32.9	32.5	34.8	35.6	34.9	30.8	§	30.0	-0.8
5.5–6.0 (High)	32.8	32.8	33.6	29.9	34.5	32.2	33.8	32.9	32.1	32.6	34.1	32.8	35.8	§	29.7	-6.1
Race/Ethnicity (2-year average): ^b																
White	33.4	33.9	34.3	34.8	36.3	36.6	35.6	35.1	35.1	35.3	36.5	36.9	35.6	§	31.8	-3.7
African American	27.0	26.8	28.6	30.8	32.4	33.2	35.0	35.9	32.8	32.4	35.5	36.3	35.7	§	34.4	-1.2
Hispanic	27.2	27.3	28.8	31.6	34.4	37.3	39.2	37.1	35.6	36.9	36.8	34.2	33.4	§	29.4	-4.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 13
MARIJUANA: ^c Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
Approximate weighted N =	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	
Total	3.2	3.7	5.1	7.8	9.1	11.3	10.2	9.7	9.7	9.1	9.2	8.3	7.5	6.4	6.6	
Gender:																
Male	3.8	3.8	6.1	9.5	9.8	12.1	11.4	10.3	10.5	10.2	11.0	9.5	8.5	6.3	7.6	
Female	2.6	3.5	4.1	6.0	8.2	10.2	8.9	8.8	8.8	7.8	7.3	7.1	6.4	6.3	5.7	
College Plans:																
None or under 4 years	9.8	11.5	13.2	19.7	20.6	24.5	23.8	24.6	25.9	23.4	23.9	23.2	21.7	19.0	19.7	
Complete 4 years	2.1	2.5	3.9	6.1	7.5	9.1	8.4	7.8	7.7	7.4	7.5	6.8	6.0	4.9	5.2	
Region:																
Northeast	2.4	3.1	3.3	7.7	7.2	9.2	9.0	6.5	7.6	7.5	8.7	6.3	5.9	5.3	5.6	
Midwest	3.3	3.0	3.7	7.3	10.1	11.8	10.4	10.8	11.7	9.5	7.9	8.0	7.3	5.6	6.6	
South	3.3	3.7	5.0	6.6	8.5	10.3	9.7	10.2	10.1	8.9	10.0	9.4	8.7	7.0	7.6	
West	3.9	5.6	9.2	10.9	10.6	13.9	11.9	10.5	8.4	10.4	9.8	8.6	7.0	7.0	5.9	
Population Density:																
Large MSA	2.8	3.3	4.1	7.6	9.1	11.3	8.7	9.3	8.6	7.5	7.1	6.7	6.5	6.6	6.3	
Other MSA	3.7	4.4	6.5	9.8	9.9	12.0	10.7	9.8	9.4	9.1	10.5	8.8	7.5	6.4	7.4	
Non-MSA	2.8	3.1	3.2	4.3	7.7	9.7	10.8	10.2	11.6	11.0	9.1	9.4	9.0	6.1	5.7	
Parental Education: ^a																
1.0–2.0 (Low)	7.4	7.2	7.2	12.4	13.5	12.5	15.5	15.7	17.5	18.8	16.4	13.5	13.7	11.3	10.6	
2.5–3.0	3.9	3.9	5.5	9.0	10.4	13.4	11.6	11.4	12.1	11.9	11.9	11.3	10.0	8.3	9.6	
3.5–4.0	3.1	3.5	5.7	7.8	9.9	12.0	11.0	10.6	9.1	8.8	9.6	8.7	7.6	7.0	7.3	
4.5–5.0	1.7	2.9	3.6	6.0	7.1	9.3	9.0	7.1	6.7	5.3	6.5	6.2	5.0	4.4	4.2	
5.5–6.0 (High)	2.3	2.4	4.2	5.8	7.4	9.2	6.7	7.3	7.5	5.8	5.3	5.4	4.8	3.6	3.3	
Race (2-year average): ^b																
White	—	3.3	4.1	5.6	7.8	10.0	10.6	9.5	8.7	8.4	8.4	8.3	7.4	6.2	5.8	
African American	—	2.0	2.9	5.0	6.6	8.0	9.0	9.1	9.7	9.3	8.1	7.4	7.3	7.7	8.2	
Hispanic	—	6.4	8.3	12.1	12.9	12.5	13.1	13.5	14.3	12.7	12.6	12.6	10.3	9.3	8.3	

Table continued on next page.

TABLE 13 (cont.)
MARIJUANA: ^c Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	6.5	5.7	5.8	6.5	8.0	7.2	6.5	7.0	6.5	6.5	5.4	5.5	5.6	6.6	6.5	4.1	-2.5* sss
Gender:																	
Male	6.7	6.2	6.6	7.5	9.2	8.5	7.0	6.7	6.9	6.6	5.7	5.3	5.6	5.9	§	2.7	-3.1 sss
Female	6.0	4.9	4.8	5.3	6.8	5.7	6.0	7.2	5.9	6.2	5.0	5.5	5.3	7.4	§	4.9	-2.4 ss
College Plans:																	
None or under 4 years	20.1	18.3	19.1	18.9	21.4	19.4	20.0	19.7	17.6	16.4	15.4	13.1	14.2	13.5	§	7.2	-6.3 ss
Complete 4 years	5.1	4.5	4.5	5.4	6.8	6.2	5.4	6.2	5.6	5.6	4.5	4.6	4.3	5.7	§	3.4	-2.2 sss
Region:																	
Northeast	5.2	3.4	4.2	6.6	7.0	4.3	4.7	5.4	6.2	5.3	3.2	3.8	4.0	4.3	§	4.3	0.0
Midwest	5.6	5.3	6.8	6.2	6.8	5.7	4.4	5.8	6.0	6.6	5.8	4.3	4.5	6.4	§	5.4	-0.9
South	7.9	6.5	6.0	6.8	8.8	7.7	7.5	8.5	6.1	6.7	5.8	5.3	5.3	6.5	§	3.2	-3.3 sss
West	5.9	5.9	5.6	6.0	8.5	10.2	8.2	7.0	7.7	6.8	6.0	7.9	8.4	8.7	§	4.1	-4.6 s
Population Density:																	
Large MSA	5.3	4.3	4.9	5.5	7.7	6.8	5.8	7.2	7.5	7.1	4.9	4.4	5.8	5.5	§	3.4	-2.0
Other MSA	7.4	6.1	6.3	7.2	8.3	8.3	7.0	6.7	6.0	6.5	5.1	6.4	5.6	7.3	§	4.6	-2.6 sss
Non-MSA	6.3	6.7	5.8	6.1	7.4	5.5	6.5	7.5	6.1	5.7	7.1	4.9	5.3	6.9	§	4.0	-2.9
Parental Education: ^a																	
1.0–2.0 (Low)	12.0	10.4	11.1	10.3	12.3	12.2	11.4	13.6	12.4	11.9	8.9	9.5	8.6	11.3	§	7.3	-4.0
2.5–3.0	8.9	7.3	8.4	8.6	10.5	10.1	9.9	10.9	8.8	8.5	7.6	7.6	8.5	11.7	§	6.9	-4.8 s
3.5–4.0	7.0	6.7	5.6	7.1	8.8	7.7	7.2	8.1	7.1	6.7	5.9	6.1	6.3	6.9	§	5.0	-1.9
4.5–5.0	3.7	3.6	3.8	5.1	6.1	5.5	4.1	4.3	4.1	3.6	4.0	3.2	3.5	3.9	§	2.9	-1.0
5.5–6.0 (High)	3.4	2.3	3.6	4.0	4.3	3.4	3.1	3.4	4.1	4.1	2.6	3.5	2.4	3.6	§	2.4	-1.2
Race (2-year average): ^b																	
White	5.8	5.4	5.0	5.4	6.5	6.5	5.3	4.7	4.7	4.6	4.1	3.9	4.0	4.2	§	4.3	+0.1
African American	7.5	6.4	6.1	6.7	7.7	8.1	7.6	8.2	7.7	7.1	7.2	6.5	5.5	6.1	§	5.5	-0.7
Hispanic	8.1	7.6	7.1	7.3	8.8	10.1	10.4	10.0	8.9	8.3	7.3	6.5	6.8	8.3	§	4.2	-4.0 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 14
MARIJUANA: ^c Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>
Total	8.7	8.1	10.9	15.8	17.2	20.4	20.5	18.7	19.4	19.7	19.8	17.8	17.0	15.9	15.2
Gender:															
Male	10.1	9.0	13.1	18.6	19.1	22.3	23.0	20.3	21.8	23.3	22.7	19.3	19.0	17.4	16.7
Female	7.3	7.1	8.6	12.8	15.0	18.6	17.9	17.2	17.0	16.2	16.8	16.4	15.0	14.2	13.4
College Plans:															
None or under 4 years	15.0	14.3	20.4	25.8	27.9	35.2	34.7	32.6	33.8	33.7	35.2	31.5	31.3	27.5	27.0
Complete 4 years	7.4	6.7	8.8	13.6	15.2	17.9	18.1	16.1	17.0	17.6	17.3	15.6	14.7	14.2	13.4
Region:															
Northeast	10.1	7.7	13.4	16.9	17.1	21.9	21.6	22.7	22.0	19.4	21.3	19.4	18.3	16.5	17.6
Midwest	8.3	7.6	10.5	15.3	16.8	20.7	19.8	16.6	19.6	19.8	19.1	16.7	14.2	14.8	14.3
South	7.2	6.6	9.1	14.8	17.2	20.3	20.1	18.7	18.1	18.2	18.0	16.6	18.5	17.1	15.7
West	10.3	11.7	12.1	17.3	17.6	18.9	21.1	17.2	18.8	22.8	23.0	20.2	17.0	14.7	12.9
Population Density:															
Large MSA	9.3	8.0	11.2	17.2	16.9	18.4	19.8	17.4	19.0	20.6	18.5	16.8	15.4	13.8	14.6
Other MSA	9.1	8.6	11.2	17.5	18.5	22.2	21.7	19.7	20.3	19.7	21.0	19.1	17.3	16.7	15.6
Non-MSA	7.6	7.2	10.1	11.4	14.7	19.3	19.2	18.4	18.1	18.8	18.7	16.5	18.8	16.9	15.0
Parental Education: ^a															
1.0–2.0 (Low)	11.4	11.2	13.3	15.8	20.6	21.4	20.9	20.4	25.0	23.4	20.6	21.6	21.6	19.5	15.8
2.5–3.0	9.5	8.0	11.3	16.3	19.3	22.9	21.7	22.1	22.0	21.0	22.8	20.4	20.3	17.9	18.1
3.5–4.0	8.1	8.3	10.9	15.8	18.0	22.1	22.5	18.3	19.3	21.2	20.8	18.1	17.8	17.3	15.9
4.5–5.0	7.9	7.4	10.0	15.0	15.6	18.1	19.2	16.4	16.6	17.9	17.5	15.2	13.4	13.2	12.7
5.5–6.0 (High)	8.9	7.5	9.4	15.5	13.3	17.4	17.1	16.2	18.2	16.7	17.2	15.5	13.8	12.7	13.5
Race (2-year average): ^b															
White	—	9.0	9.8	13.4	16.8	19.3	21.2	20.3	19.8	20.2	20.2	19.8	18.2	16.5	15.7
African American	—	3.6	4.9	9.8	13.8	15.9	16.5	15.3	14.6	15.8	16.7	15.2	15.1	16.3	15.3
Hispanic	—	10.4	12.4	15.6	17.7	19.1	21.3	21.4	20.6	20.5	20.5	18.2	17.0	17.7	16.0

Table continued on next page.

TABLE 14 (cont.)
MARIJUANA: ^c Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>																	
Total	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	-8.3* sss
Gender:																	
Male	14.2	14.2	13.8	15.9	16.7	17.6	17.0	18.0	16.6	14.8	14.0	15.7	16.7	18.4	16.6	10.1	
Female	15.7	15.8	15.2	18.7	20.1	20.8	19.8	20.6	17.4	15.6	14.2	15.3	17.3	18.1	§	9.7	-8.4 sss
College Plans:	12.6	12.5	12.3	13.2	13.3	14.5	14.4	15.3	15.7	13.7	13.7	15.8	15.8	18.3	§	9.4	-8.9 sss
None or under 4 years	28.1	27.6	28.4	28.0	34.0	34.2	30.3	31.0	29.3	27.9	24.8	24.7	28.4	29.1	§	18.4	-10.7 sss
Complete 4 years	12.4	12.3	11.9	14.5	14.6	15.7	15.7	16.5	15.1	13.4	12.8	14.4	14.8	16.2	§	7.9	-8.3 sss
Region:																	
Northeast	14.6	14.3	16.6	15.3	15.5	19.0	19.1	17.3	17.5	14.7	14.3	17.2	20.4	19.7	§	12.8	-7.0 ss
Midwest	13.6	14.7	12.5	13.3	15.4	16.4	14.4	16.9	14.4	12.6	14.0	15.6	14.9	17.0	§	9.9	-7.1 ss
South	14.6	12.8	14.0	17.3	16.7	16.7	17.2	19.0	17.0	15.3	14.3	13.8	15.4	17.8	§	10.2	-7.6 sss
West	14.1	15.3	12.9	17.2	19.0	19.2	18.0	18.2	17.3	16.4	13.4	17.7	17.5	20.0	§	8.1	-11.9 sss
Population Density:																	
Large MSA	14.2	14.2	13.9	16.1	16.0	17.0	17.3	17.2	16.7	14.8	14.5	17.4	18.1	18.3	§	7.6	-10.7 sss
Other MSA	13.6	13.5	14.4	16.4	18.0	18.9	17.2	19.1	17.1	15.4	13.7	15.4	17.6	19.8	§	10.6	-9.2 sss
Non-MSA	15.7	15.5	12.4	14.8	15.1	15.8	16.3	16.3	14.9	13.4	14.2	13.9	12.4	14.6	§	13.0	-1.6
Parental Education: ^a																	
1.0–2.0 (Low)	18.4	17.1	18.1	22.2	21.0	23.6	21.0	22.0	18.6	21.0	16.6	18.5	19.8	21.5	§	16.3	-5.2
2.5–3.0	17.1	17.4	15.7	18.1	20.4	21.2	21.3	21.4	19.9	18.4	15.8	18.1	19.6	24.1	§	12.0	-12.1 sss
3.5–4.0	14.9	15.1	15.4	16.0	17.7	18.9	17.5	19.7	18.5	17.3	15.9	17.7	18.2	20.5	§	11.1	-9.4 sss
4.5–5.0	12.0	11.2	11.2	14.4	13.7	14.8	15.4	15.5	13.4	12.1	12.0	13.5	14.4	14.5	§	9.6	-5.0 ss
5.5–6.0 (High)	11.8	11.3	10.1	13.1	12.9	13.2	14.0	12.8	13.7	9.8	12.1	13.3	13.0	13.8	§	6.9	-7.0 sss
Race (2-year average): ^b																	
White	15.2	14.7	14.2	14.6	15.7	16.4	16.7	16.5	16.2	14.6	14.1	15.1	15.3	15.9	§	12.9	-3.0 ss
African American	13.8	12.6	11.7	13.7	15.5	18.0	18.8	19.1	19.0	17.4	15.8	14.8	16.1	18.7	§	16.9	-1.8
Hispanic	14.0	13.0	15.3	17.4	18.2	18.2	18.5	20.0	19.4	18.1	15.5	14.9	16.9	18.2	§	15.2	-3.0

Source. The Monitoring the Future Study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 15
MARIJUANA: ^c Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	27.1	32.2	35.4	37.1	36.5	33.7	31.6	28.5	27.0	25.2	25.7	23.4	21.0	18.0	16.7	14.0
Gender:																
Male	32.3	37.7	40.7	42.6	41.4	37.8	35.3	31.4	31.0	28.2	28.7	26.8	23.1	20.7	19.5	16.1
Female	22.5	26.0	30.0	31.3	31.3	29.1	27.3	24.9	22.2	21.1	22.4	20.0	18.6	15.2	13.8	11.5
College Plans:																
None or under 4 years	—	34.5	38.7	39.2	39.6	37.7	36.1	32.9	30.7	29.2	29.0	28.3	25.1	20.4	21.0	17.6
Complete 4 years	—	28.4	31.0	33.2	32.2	29.4	27.4	23.9	22.9	20.9	22.7	20.0	18.5	16.4	14.6	11.9
Region:																
Northeast	32.2	38.6	40.4	46.7	44.7	39.3	38.2	33.3	32.0	33.3	32.0	28.5	25.3	20.2	18.6	17.6
Midwest	27.6	31.4	36.1	37.8	38.0	34.0	33.0	30.0	27.2	22.1	26.1	24.0	21.1	18.6	19.2	13.7
South	21.2	27.7	31.3	30.6	29.0	28.4	24.7	22.4	22.9	22.4	18.0	19.2	17.3	15.8	14.0	11.5
West	30.8	32.7	33.6	34.3	35.9	35.2	32.0	29.2	27.1	25.9	29.7	23.1	22.3	18.9	16.8	14.7
Population Density:																
Large MSA	36.2	37.9	40.4	44.0	42.2	39.6	36.3	34.3	31.7	28.5	28.5	26.7	23.1	19.4	15.2	14.1
Other MSA	26.4	32.5	36.2	37.1	37.5	34.5	31.4	28.3	28.1	25.1	25.9	22.9	21.3	19.3	17.5	14.6
Non-MSA	22.2	27.5	30.2	31.4	30.9	28.3	28.0	23.8	21.8	22.5	23.0	21.5	18.2	14.3	16.7	12.6
Parental Education: ^a																
1.0–2.0 (Low)	22.3	29.0	30.4	30.7	33.2	29.9	29.7	24.9	26.2	23.8	23.4	21.0	19.9	15.6	13.9	11.4
2.5–3.0	26.5	33.7	36.7	37.7	35.8	34.6	31.5	28.4	27.3	25.4	25.9	24.1	20.9	16.8	16.3	14.3
3.5–4.0	25.1	32.3	36.9	37.7	37.8	33.6	31.4	29.3	26.1	25.8	26.5	24.1	21.1	17.7	17.9	13.5
4.5–5.0	27.3	32.6	34.3	39.0	37.3	33.7	31.9	30.1	26.9	23.3	27.1	23.1	21.1	19.3	17.1	15.0
5.5–6.0 (High)	24.7	32.1	36.0	39.7	37.6	36.2	31.9	27.7	25.5	23.4	20.6	21.7	21.2	20.6	16.2	15.0
Race (2-year average): ^b																
White	—	—	34.0	36.9	37.6	35.8	33.3	30.8	27.9	26.0	25.9	25.5	23.4	21.1	19.2	17.1
African American	—	—	29.2	29.7	28.2	27.0	25.7	24.8	25.8	24.8	22.3	19.4	14.5	11.0	9.6	7.5
Hispanic	—	—	33.3	30.8	29.0	29.5	27.7	26.2	25.3	23.8	21.9	21.3	20.2	16.1	12.4	11.7

Table continued on next page.

TABLE 15 (cont.)
MARIJUANA: ^c Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700
Total	13.8	11.9	15.5	19.0	21.2	21.9	23.7	22.8	23.1	21.6	22.4	21.5	21.2	19.9	19.8
Gender:															
Male	16.1	13.4	18.2	23.0	24.6	25.1	26.4	26.5	26.3	24.7	25.6	25.3	24.7	23.0	23.6
Female	11.2	10.2	12.5	15.1	17.2	18.3	20.3	18.8	19.7	18.3	19.1	17.4	17.3	16.6	15.8
College Plans:															
None or under 4 years	15.8	15.0	17.9	21.6	23.7	26.4	29.3	28.3	26.6	26.0	27.5	27.0	26.9	24.1	24.9
Complete 4 years	12.5	10.4	14.2	17.7	19.6	20.0	21.5	20.5	21.8	19.6	20.4	19.6	18.9	18.3	18.1
Region:															
Northeast	17.6	14.4	19.5	22.7	23.8	25.9	27.0	27.2	24.2	26.0	28.4	25.6	26.5	25.1	25.0
Midwest	14.9	12.2	15.9	19.3	22.6	23.3	21.8	21.1	23.2	19.3	22.7	22.4	21.6	18.1	19.2
South	9.7	9.4	13.1	17.3	19.4	19.5	20.9	21.3	22.1	17.4	18.1	20.0	17.8	17.7	18.2
West	15.7	14.0	15.6	18.6	19.8	19.7	27.8	23.9	24.2	27.2	23.5	19.3	21.1	20.9	18.0
Population Density:															
Large MSA	14.3	12.9	18.0	20.7	23.6	22.9	21.7	23.4	23.2	22.6	23.4	22.9	19.6	19.1	20.4
Other MSA	16.3	11.5	15.3	19.8	21.5	23.7	25.6	23.9	24.4	22.6	22.5	22.7	23.7	22.3	20.8
Non-MSA	9.0	12.0	13.8	15.7	17.8	17.6	22.0	19.8	21.0	18.6	21.2	17.4	18.7	16.4	16.9
Parental Education: ^a															
1.0–2.0 (Low)	11.7	9.7	12.4	14.0	18.1	17.4	19.9	19.7	22.2	21.5	20.9	17.8	20.0	16.6	16.3
2.5–3.0	12.9	11.9	14.0	18.5	19.6	20.6	23.6	22.1	23.6	20.9	21.7	21.5	20.6	19.3	20.2
3.5–4.0	13.8	12.2	16.9	19.3	21.5	22.9	23.6	23.9	23.0	21.5	23.6	23.7	22.1	21.2	20.2
4.5–5.0	13.7	12.1	15.9	20.4	21.3	22.1	24.7	22.6	22.4	21.4	22.3	20.6	21.0	19.8	20.4
5.5–6.0 (High)	17.6	11.6	16.1	20.3	23.1	22.6	22.9	22.9	23.4	21.3	20.5	19.6	18.3	19.8	18.9
Race (2-year average): ^b															
White	15.3	14.1	14.9	18.4	20.8	22.0	23.6	24.4	23.8	22.7	22.9	23.3	22.8	22.1	21.6
African American	5.9	6.0	8.1	13.1	16.8	18.3	18.5	18.3	19.3	19.0	17.0	16.5	16.3	15.2	14.6
Hispanic	12.9	12.7	12.5	14.9	17.9	19.1	21.2	21.6	22.0	24.6	22.1	20.0	17.9	16.6	16.6

Table continued on next page.

TABLE 15 (cont.)
MARIJUANA: ^c Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days																2019-2021 change
	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	
Approximate weighted N =	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	18.3	18.8	19.4	20.6	21.4	22.6	22.9	22.7	21.2	21.3	22.5	22.9	22.2	22.3	21.1	19.5	-2.8* s
Gender:																	
Male	19.7	22.3	22.2	24.3	25.2	26.4	26.5	26.4	24.3	23.1	25.0	24.1	24.0	23.0	§	17.5	-5.5 sss
Female	16.4	15.0	16.2	16.8	16.9	18.4	18.8	18.7	17.9	19.2	19.7	21.5	19.2	21.1	§	20.6	-0.5
College Plans:																	
None or under 4 years	24.8	25.9	24.9	24.6	28.1	27.5	27.0	28.0	26.8	24.6	25.4	26.6	26.6	27.0	§	22.1	-5.0
Complete 4 years	16.4	16.8	17.7	19.2	19.3	21.1	21.4	21.3	19.5	20.0	21.2	21.6	20.2	20.3	§	18.0	-2.2
Region:																	
Northeast	24.8	23.7	23.1	24.6	26.1	26.4	24.0	25.8	25.6	26.2	25.8	26.7	27.4	22.7	§	22.0	-0.7
Midwest	17.4	18.6	17.5	20.2	20.0	21.7	19.5	20.0	21.0	19.6	20.4	22.7	23.2	20.7	§	20.1	-0.6
South	17.1	17.0	19.2	19.3	19.6	20.9	22.3	21.0	19.8	18.5	20.5	19.8	17.9	20.6	§	14.4	-6.2 sss
West	16.0	17.7	19.2	19.7	21.7	23.4	26.6	25.8	20.0	23.8	25.6	25.6	24.7	28.1	§	24.8	-3.3
Population Density:																	
Large MSA	19.6	20.9	21.8	24.0	22.8	23.3	25.9	25.7	22.0	24.5	24.4	25.2	24.0	22.3	§	21.2	-1.1
Other MSA	18.4	18.2	18.6	20.3	22.7	24.3	23.5	23.5	22.4	20.4	22.6	21.4	21.2	22.5	§	19.4	-3.1
Non-MSA	16.3	16.7	18.5	16.8	15.9	17.5	16.7	16.4	17.1	17.9	19.3	22.4	21.3	21.7	§	16.8	-4.9 s
Parental Education: ^a																	
1.0–2.0 (Low)	17.9	18.2	18.5	16.9	18.6	20.5	26.7	22.6	23.0	21.9	21.4	22.2	19.8	22.6	§	15.8	-6.8 s
2.5–3.0	19.6	20.1	18.7	22.0	21.2	23.0	23.2	25.2	23.0	22.9	24.4	24.5	23.8	24.4	§	22.6	-1.8
3.5–4.0	18.6	18.5	18.8	21.7	22.8	23.5	23.2	23.5	21.4	21.4	22.8	24.8	23.5	23.8	§	21.4	-2.4
4.5–5.0	16.3	18.3	19.3	19.1	20.8	21.0	22.3	20.8	19.7	20.3	21.6	21.1	20.8	18.5	§	16.9	-1.6
5.5–6.0 (High)	18.2	17.9	21.0	20.3	18.4	23.1	20.3	21.3	19.8	18.6	18.6	19.1	20.8	20.5	§	19.4	-1.2
Race (2-year average): ^b																	
White	20.5	19.5	20.1	20.8	21.4	22.2	22.6	21.8	21.3	21.0	21.1	22.1	22.3	21.3	§	18.7	-2.6
African American	15.9	16.0	16.3	18.6	20.1	21.0	22.3	24.0	23.0	20.8	22.2	23.9	24.0	23.8	§	23.7	-0.1
Hispanic	16.0	15.3	15.3	16.7	18.5	19.6	22.0	24.3	22.6	20.9	22.5	22.9	20.5	20.4	§	18.1	-2.3

Source. The Monitoring the Future Study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 16
MARIJUANA: ^c Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 8

	Percentage who used daily in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	
Total	0.2	0.2	0.4	0.7	0.8	1.5	1.1	1.1	1.4	1.3	1.3	1.2	1.0	0.8	1.0	
Gender																
Male	0.3	0.3	0.5	1.0	1.1	2.0	1.4	1.4	1.9	2.0	2.0	1.7	1.3	0.9	1.4	
Female	0.1	0.1	0.3	0.5	0.4	0.9	0.6	0.7	0.8	0.7	0.6	0.7	0.6	0.6	0.6	
College Plans:																
None or under 4 years	0.9	1.1	1.2	2.1	3.1	4.4	4.2	4.2	5.6	5.1	5.1	5.1	3.6	3.3	3.9	
Complete 4 years	0.1	0.1	0.3	0.5	0.5	1.0	0.6	0.7	0.8	0.9	0.8	0.8	0.7	0.4	0.7	
Region:																
Northeast	0.1	0.1	0.3	0.8	0.5	0.9	0.9	0.8	1.0	0.8	1.4	1.2	0.8	0.5	0.7	
Midwest	0.1	0.2	0.2	0.7	0.8	1.6	1.2	1.1	1.7	1.4	1.3	1.0	1.0	0.6	1.0	
South	0.3	0.2	0.4	0.6	0.8	1.4	0.9	1.4	1.5	1.4	1.5	1.6	1.1	0.9	1.2	
West	0.2	0.3	0.7	1.1	1.2	2.1	1.3	0.9	1.0	1.6	0.9	1.0	0.9	0.9	0.8	
Population Density:																
Large MSA	0.2	0.1	0.3	0.7	1.1	1.7	0.8	0.8	1.1	1.0	0.8	1.0	0.8	0.7	0.9	
Other MSA	0.2	0.2	0.5	0.9	0.9	1.6	1.2	1.1	1.5	1.3	1.6	1.3	1.0	0.8	1.2	
Non-MSA	0.2	0.3	0.2	0.5	0.5	1.0	1.1	1.4	1.3	1.8	1.4	1.4	1.2	0.8	0.7	
Parental Education: ^a																
1.0–2.0 (Low)	0.5	0.5	0.7	0.8	2.0	1.6	2.2	2.2	3.2	4.6	2.0	2.3	1.9	1.6	1.4	
2.5–3.0	0.3	0.3	0.4	0.8	0.9	2.0	0.9	1.5	1.3	1.6	1.9	1.7	1.4	0.8	1.2	
3.5–4.0	0.2	0.1	0.5	1.1	0.7	1.4	1.3	1.1	1.5	0.9	1.3	1.4	1.0	0.7	1.2	
4.5–5.0	0.1	0.1	0.1	0.5	0.6	0.8	0.8	0.5	0.8	0.5	0.8	0.7	0.7	0.5	0.5	
5.5–6.0 (High)	*	0.1	0.3	0.5	0.6	1.2	0.6	0.6	0.7	0.8	0.5	0.4	0.6	0.3	0.2	
Race (2-year average): ^b																
White	—	0.2	0.2	0.4	0.7	1.0	1.1	0.9	1.0	1.2	1.3	1.3	1.0	0.7	0.7	
African American	—	0.1	0.2	0.4	0.7	1.2	1.3	0.9	1.0	1.0	1.1	1.2	1.2	1.2	1.0	
Hispanic	—	0.3	0.6	1.1	1.2	1.6	1.8	1.6	2.0	1.9	1.6	1.5	1.2	1.1	1.3	

Table continued on next page.

TABLE 16 (cont.)
MARIJUANA: ^c Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 8

	Percentage who used daily in last 30 days																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	1.0	0.8	0.9	1.0	1.2	1.3	1.1	1.1	1.0	1.1	0.7	0.8	0.7	1.3	1.1	0.6	-0.6*
Gender																	
Male	1.3	1.2	1.2	1.4	1.8	1.8	1.6	1.3	1.2	1.3	0.9	0.9	0.8	1.4	§	0.6	-0.7
Female	0.7	0.5	0.6	0.5	0.7	0.7	0.5	0.9	0.7	0.9	0.5	0.6	0.5	1.2	§	0.7	-0.5
College Plans:																	
None or under 4 years	5.3	3.2	4.7	4.7	5.2	5.2	5.6	4.9	2.9	3.4	3.0	2.7	2.2	3.7	§	1.8	-1.9
Complete 4 years	0.6	0.6	0.6	0.6	0.9	1.0	0.7	0.8	0.8	0.8	0.5	0.5	0.4	0.9	§	0.4	-0.5 s
Region:																	
Northeast	0.7	0.5	0.3	0.9	0.9	0.5	0.8	0.6	0.9	0.8	0.4	0.8	0.5	0.7	§	1.0	+0.3
Midwest	0.8	0.9	1.3	1.0	1.3	1.2	0.5	0.9	1.3	1.1	0.7	0.5	0.5	1.1	§	0.9	-0.2
South	1.4	0.9	1.0	1.0	1.4	1.4	1.2	1.3	0.8	1.2	0.8	0.7	0.6	0.6	§	0.4	-0.2
West	0.9	0.7	0.8	0.9	1.1	1.9	1.6	1.5	1.1	1.1	0.9	1.0	1.3	2.8	§	0.6	-2.3
Population Density:																	
Large MSA	0.8	0.5	0.9	0.6	0.8	1.1	0.9	1.0	1.3	1.5	0.4	0.5	0.6	0.7	§	0.3	-0.4 s
Other MSA	1.3	0.8	0.8	1.2	1.3	1.5	1.2	1.2	0.8	0.9	0.8	0.9	0.6	1.3	§	0.9	-0.4
Non-MSA	0.9	1.4	1.0	1.0	1.7	1.0	1.2	1.3	0.9	0.9	1.1	0.7	1.0	2.2	§	0.7	-1.5
Parental Education: ^a																	
1.0–2.0 (Low)	2.6	1.3	2.6	1.9	2.5	2.2	1.9	2.3	2.0	2.5	1.3	1.2	1.4	1.4	§	2.3	+0.9
2.5–3.0	1.3	0.9	1.4	1.4	2.1	2.3	1.6	1.5	0.9	1.0	0.9	1.1	0.7	2.9	§	0.8	-2.0 s
3.5–4.0	0.9	1.0	0.7	0.8	1.0	1.3	1.0	1.0	1.1	0.8	0.9	0.8	0.6	1.1	§	0.9	-0.2
4.5–5.0	0.6	0.4	0.4	0.6	0.7	0.8	0.3	0.8	0.6	0.6	0.4	0.5	0.4	0.3	§	0.5	+0.2
5.5–6.0 (High)	0.3	0.5	0.5	0.6	0.3	0.5	0.6	0.5	0.7	0.7	0.4	0.4	0.3	1.0	§	0.1	-0.9 s
Race (2-year average): ^b																	
White	1.0	0.9	0.7	0.8	1.0	1.1	1.0	0.8	0.7	0.6	0.6	0.5	0.4	0.5	§	1.0	+0.5
African American	1.0	1.0	1.0	1.0	1.1	1.2	1.0	1.0	1.1	1.2	1.2	1.2	1.1	0.7	§	0.8	+0.1
Hispanic	1.1	0.7	1.0	1.3	1.4	1.9	2.1	1.7	1.4	1.3	0.9	0.8	0.9	1.4	§	0.3	-1.1 ss

Source. The Monitoring the Future Study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. '*' indicates less than .05 percent but greater than 0 percent. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 17
MARIJUANA: ^c Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 10

	Percentage who used daily in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Approximate weighted N =	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	0.8	0.8	1.0	2.2	2.8	3.5	3.7	3.6	3.8	3.8	4.5	3.9	3.6	3.2	3.1
Gender:															
Male	1.1	1.1	1.6	3.1	3.7	4.7	5.2	4.8	5.2	5.3	6.4	5.1	5.1	4.1	4.1
Female	0.5	0.4	0.4	1.2	1.7	2.3	2.1	2.4	2.4	2.3	2.6	2.6	2.2	2.2	2.0
College Plans:															
None or under 4 years	1.6	2.3	2.7	5.5	6.5	10.2	9.5	9.7	10.3	10.4	11.7	9.4	9.9	9.0	9.5
Complete 4 years	0.6	0.4	0.6	1.4	2.1	2.3	2.6	2.4	2.7	2.7	3.3	2.9	2.6	2.3	2.1
Region:															
Northeast	0.8	0.7	1.6	2.5	1.9	3.7	3.9	5.0	3.9	3.9	3.8	3.6	3.7	3.3	3.3
Midwest	0.9	0.6	1.0	2.0	2.9	3.8	3.8	3.4	4.2	4.2	4.1	4.2	2.8	3.0	3.3
South	0.6	0.6	0.7	2.2	3.2	3.0	3.9	3.4	3.6	3.4	4.5	3.2	4.2	3.4	2.9
West	1.1	1.3	1.1	2.1	2.7	3.8	2.7	2.9	3.7	3.7	6.2	5.2	3.8	3.0	2.9
Population Density:															
Large MSA	0.8	0.5	1.2	2.1	2.5	3.0	3.6	3.4	3.1	4.2	4.4	3.4	3.0	2.5	3.0
Other MSA	0.9	0.8	0.9	2.3	3.1	3.7	3.8	3.9	4.1	3.4	4.4	4.4	3.6	3.3	3.0
Non-MSA	0.7	0.9	1.1	2.0	2.4	3.7	3.6	3.3	3.9	4.0	4.8	3.7	4.8	3.8	3.4
Parental Education: ^a															
1.0–2.0 (Low)	1.1	1.1	1.4	2.0	4.1	3.6	3.2	3.8	4.6	4.9	6.0	4.6	5.5	5.3	3.7
2.5–3.0	0.9	0.8	1.0	2.6	3.6	4.6	4.3	4.9	5.5	4.3	5.3	4.9	4.8	3.8	4.8
3.5–4.0	0.8	0.9	1.1	2.4	2.8	4.1	4.2	3.4	3.5	4.2	4.7	3.9	3.4	3.4	3.3
4.5–5.0	0.8	0.6	0.6	1.8	1.9	2.4	3.0	3.1	2.6	3.4	3.8	2.7	2.9	2.0	2.0
5.5–6.0 (High)	0.4	0.8	1.0	1.0	2.0	2.1	2.2	2.3	2.5	2.1	3.3	3.2	2.4	1.7	1.8
Race (2-year average): ^b															
White	—	0.9	1.0	1.6	2.5	3.2	3.7	3.7	3.8	4.0	4.3	4.4	4.1	3.5	3.1
African American	—	0.3	0.3	0.8	1.7	2.2	2.8	3.4	3.0	3.2	3.6	3.0	2.9	2.8	2.8
Hispanic	—	0.8	1.1	1.9	2.5	2.8	3.5	3.6	3.2	3.2	4.2	3.9	3.0	2.9	3.1

Table continued on next page.

TABLE 17 (cont.)
MARIJUANA: ^c Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 10

	Percentage who used daily in last 30 days																	
	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	2019-2021 change	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000		
Total	2.8	2.8	2.7	2.8	3.3	3.6	3.5	4.0	3.4	3.0	2.5	2.9	3.4	4.8	4.4	3.2	-1.6* s	
Gender:																		
Male	3.5	3.9	3.8	4.1	5.1	5.4	4.6	6.2	4.3	3.6	3.1	3.3	4.2	5.2	§	3.2	-2.0 ss	
Female	2.0	1.8	1.7	1.7	1.6	1.7	2.4	2.0	2.4	2.4	1.9	2.4	2.5	4.2	§	2.7	-1.5	
College Plans:																		
None or under 4 years	8.9	8.3	9.2	7.9	9.5	9.5	10.5	10.0	8.8	8.8	7.4	8.1	8.9	10.3	§	8.0	-2.3	
Complete 4 years	1.9	2.1	1.9	2.2	2.6	2.9	2.8	3.3	2.8	2.4	1.9	2.2	2.5	3.5	§	2.0	-1.5 ss	
Region:																		
Northeast	2.8	3.2	2.8	1.9	2.9	4.3	3.2	2.8	3.4	3.0	2.4	3.2	4.1	4.3	§	4.0	-0.3	
Midwest	2.6	2.9	2.6	2.5	2.9	2.9	3.2	3.9	3.2	2.0	2.6	3.2	2.5	4.2	§	2.1	-2.1 ss	
South	3.0	2.7	3.2	3.5	3.3	3.6	4.1	5.0	3.7	3.7	2.6	2.7	3.6	5.0	§	3.1	-1.9	
West	2.5	2.7	2.4	3.1	4.3	3.8	3.5	3.7	3.3	3.0	2.4	2.9	3.5	5.3	§	3.6	-1.7	
Population Density:																		
Large MSA	2.8	2.3	2.5	2.2	2.8	3.6	2.9	3.4	3.3	2.6	2.1	2.7	3.2	3.8	§	2.3	-1.5 s	
Other MSA	2.6	2.9	3.0	3.0	3.7	3.5	3.6	4.4	3.6	3.2	2.5	2.7	3.8	5.7	§	3.5	-2.3	
Non-MSA	3.2	3.5	2.6	3.4	3.3	3.6	4.3	3.9	3.2	3.5	3.4	4.0	2.9	3.7	§	3.8	+0.1	
Parental Education: ^a																		
1.0–2.0 (Low)	4.2	4.9	4.2	4.6	4.5	5.8	5.5	5.8	4.3	5.1	3.0	4.3	4.9	5.6	§	5.6	+0.1	
2.5–3.0	3.9	3.4	3.7	3.4	4.6	4.3	5.3	5.7	4.4	4.8	3.3	3.9	5.0	7.5	§	5.3	-2.2	
3.5–4.0	2.9	3.2	3.2	2.9	3.8	3.7	3.2	4.4	4.6	3.3	3.2	3.2	3.3	5.8	§	3.1	-2.7 s	
4.5–5.0	1.8	2.0	1.8	2.1	2.4	2.5	3.0	2.8	2.0	1.7	1.8	2.1	2.3	2.7	§	2.8	0.0	
5.5–6.0 (High)	1.9	1.6	1.3	1.9	1.7	2.9	1.8	2.2	1.4	1.8	0.9	1.9	2.3	2.0	§	1.2	-0.8	
Race (2-year average): ^b																		
White	3.0	3.1	2.8	2.6	2.9	3.3	3.4	3.5	3.5	3.2	3.0	2.7	2.8	3.4	§	3.4	-0.1	
African American	2.8	2.5	2.8	2.9	3.2	3.8	4.0	4.7	4.4	3.0	2.3	2.7	3.8	4.6	§	6.3	+1.7	
Hispanic	2.6	1.9	2.5	2.9	3.2	3.3	3.2	3.7	3.7	3.3	2.7	2.6	2.9	4.5	§	2.9	-1.6	

Source. The Monitoring the Future Study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 18
MARIJUANA: ^cTrends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Approximate weighted N =	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	6.0	8.2	9.1	10.7	10.3	9.1	7.0	6.3	5.5	5.0	4.9	4.0	3.3	2.7	2.9	2.2
Gender:																
Male	8.1	10.8	12.4	14.2	12.7	11.9	9.6	8.2	7.3	7.0	6.9	5.7	4.3	3.9	4.1	3.2
Female	4.0	5.0	5.6	7.1	7.3	6.0	4.2	4.0	3.2	2.5	2.8	2.3	2.1	1.3	1.5	1.0
College Plans:																
None or under 4 years	—	9.9	11.1	12.8	13.0	11.9	9.4	8.6	7.3	6.9	6.7	6.2	5.2	4.0	4.6	3.5
Complete 4 years	—	5.5	6.3	7.4	6.8	5.9	4.8	3.9	3.4	2.9	3.3	2.3	2.0	1.8	2.0	1.5
Region:																
Northeast	6.7	10.2	9.9	14.5	13.6	11.1	9.1	8.0	6.9	7.5	6.8	5.0	4.1	2.5	3.3	2.7
Midwest	6.2	8.1	8.8	11.4	11.5	9.5	8.2	6.8	5.5	4.4	5.5	4.0	3.1	2.5	3.2	2.3
South	5.0	6.7	9.1	8.5	7.0	7.5	4.5	4.8	4.5	4.1	3.0	2.8	2.9	2.6	2.6	1.7
West	6.5	8.0	8.1	8.2	9.3	8.6	6.4	5.4	5.1	4.7	4.5	4.6	3.4	3.1	2.5	2.4
Population Density:																
Large MSA	8.4	10.7	9.5	12.7	10.6	10.3	8.3	7.9	7.3	5.7	5.8	4.6	3.8	2.6	2.3	2.4
Other MSA	5.9	8.2	10.0	10.9	11.3	9.5	7.1	6.0	5.7	5.0	5.0	3.9	3.5	3.4	3.0	2.0
Non-MSA	4.5	6.3	7.6	9.0	8.6	7.7	6.0	5.3	3.8	4.4	3.8	3.6	2.6	1.4	3.3	2.4
Parental Education: ^a																
1.0–2.0 (Low)	4.4	6.8	6.9	8.6	8.4	8.2	6.2	6.2	5.7	5.1	4.2	4.1	4.0	2.5	2.4	2.0
2.5–3.0	5.7	8.2	9.9	11.3	10.9	10.0	7.5	6.5	5.4	5.4	5.5	4.4	2.9	2.7	2.6	2.3
3.5–4.0	5.0	8.0	9.0	10.3	10.1	8.7	6.4	6.4	5.2	4.6	4.7	3.7	3.5	2.4	2.6	1.9
4.5–5.0	5.5	8.8	8.6	10.4	9.0	7.8	6.4	5.7	5.0	4.0	4.2	3.5	2.9	2.6	3.3	2.1
5.5–6.0 (High)	4.3	8.1	9.0	10.2	10.4	8.0	7.1	4.8	3.7	4.0	3.6	2.3	2.7	2.3	2.7	2.5
Race (2-year average): ^b																
White	—	—	8.6	10.2	10.8	9.7	8.0	6.8	5.9	5.1	4.9	4.5	3.7	3.2	3.1	2.8
African American	—	—	6.4	6.0	5.1	5.8	5.4	4.3	4.1	3.6	3.3	3.1	2.3	1.3	0.9	0.7
Hispanic	—	—	6.9	6.3	6.5	8.3	6.9	5.7	5.3	4.7	3.9	3.3	3.0	2.1	1.9	2.0

Table continued on next page.

TABLE 18 (cont.)
MARIJUANA: ^c Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700
Total	2.0	1.9	2.4	3.6	4.6	4.9	5.8	5.6	6.0	6.0	5.8	6.0	6.0	5.6	5.0
Gender:															
Male	3.0	2.8	3.3	5.1	6.5	6.8	8.1	7.7	7.9	8.2	7.9	8.7	8.3	7.7	7.2
Female	0.9	1.0	1.5	2.0	2.4	2.7	3.1	3.2	3.9	3.5	3.6	3.1	3.4	3.1	2.8
College Plans:															
None or under 4 years	3.3	3.5	3.6	4.9	6.8	7.1	8.0	9.8	9.2	9.8	8.8	10.3	9.4	8.1	8.4
Complete 4 years	1.4	1.2	2.0	2.9	3.5	3.9	4.7	4.0	4.7	4.4	4.4	4.6	4.5	4.5	4.0
Region:															
Northeast	2.4	2.4	3.0	4.1	5.1	6.2	6.9	6.8	5.7	7.3	7.7	7.4	7.8	6.8	5.5
Midwest	1.9	1.8	2.3	4.1	4.1	5.2	5.2	5.3	6.2	5.9	6.8	6.2	5.7	5.1	5.4
South	1.8	1.3	1.7	3.0	4.3	3.8	5.3	4.9	5.2	4.6	3.9	5.6	5.4	5.0	5.4
West	2.1	2.6	3.3	3.3	5.3	4.8	6.3	6.5	7.6	7.4	5.9	5.1	5.6	6.1	3.6
Population Density:															
Large MSA	1.9	1.9	2.5	3.7	5.3	5.5	4.6	5.4	5.4	5.7	5.9	6.0	4.4	5.3	5.3
Other MSA	2.5	1.7	2.4	3.9	4.5	5.0	6.6	5.9	5.8	6.3	5.7	6.1	7.6	6.3	5.0
Non-MSA	1.2	2.1	2.3	2.8	3.9	4.1	5.4	5.4	7.1	5.9	6.0	5.6	5.2	4.7	4.7
Parental Education: ^a															
1.0–2.0 (Low)	2.3	1.6	2.3	2.3	4.0	3.9	4.9	6.2	8.8	7.7	5.5	4.5	5.0	5.5	4.3
2.5–3.0	2.4	1.9	2.1	3.8	4.9	5.1	5.8	5.7	6.2	6.3	5.8	7.0	6.0	5.5	5.7
3.5–4.0	1.8	1.8	2.8	3.6	4.0	5.0	5.7	5.8	5.5	5.9	5.7	6.7	6.4	6.0	5.0
4.5–5.0	1.7	1.6	2.3	3.6	4.3	4.8	6.1	5.0	5.5	5.3	5.4	4.8	5.5	5.5	5.1
5.5–6.0 (High)	1.6	2.0	2.1	3.1	3.9	4.1	4.8	4.1	4.6	4.1	4.4	4.6	3.9	3.7	3.8
Race (2-year average): ^b															
White	2.2	2.0	2.4	3.2	4.1	4.7	5.5	5.9	5.8	6.0	6.2	6.4	6.5	6.3	5.8
African American	0.7	0.5	0.8	2.0	3.2	3.7	3.9	4.4	4.7	4.6	4.1	3.7	3.7	3.2	3.5
Hispanic	2.2	2.1	1.9	2.0	3.0	4.2	4.5	4.7	6.0	6.9	5.0	4.0	4.0	4.0	3.7

Table continued on next page.

TABLE 18 (cont.)
MARIJUANA: ^c Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	<u>change</u>
<i>Approximate weighted N =</i>	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	5.0	5.1	5.4	5.2	6.1	6.6	6.5	6.5	5.8	6.0	6.0	5.9	5.8	6.4	6.9	5.8	-0.6*
Gender:																	
Male	6.3	6.8	7.6	7.5	8.8	9.7	9.1	8.9	8.5	8.1	8.0	7.8	7.2	8.0	§	6.0	-2.0 ss
Female	3.2	3.2	3.0	2.6	3.2	3.3	3.6	3.8	3.3	3.8	3.6	3.6	3.4	4.6	§	5.3	+0.7
College Plans:																	
None or under 4 years	9.3	9.6	9.2	8.3	10.5	11.5	9.0	10.7	9.8	9.7	9.5	10.4	9.2	11.1	§	8.8	-2.2
Complete 4 years	3.7	3.8	4.2	4.1	4.7	5.4	5.5	5.1	4.6	5.0	4.9	4.6	4.1	4.7	§	4.4	-0.3
Region:																	
Northeast	7.9	7.0	6.8	5.3	7.5	7.1	7.1	8.0	7.0	7.0	7.6	7.3	6.5	5.7	§	5.5	-0.2
Midwest	4.9	5.7	4.8	6.0	5.8	6.8	6.0	5.5	6.2	5.6	5.6	6.0	6.4	5.7	§	6.2	+0.5
South	4.2	4.3	5.5	5.0	6.1	6.3	6.1	6.4	5.3	5.5	5.0	4.7	4.2	6.2	§	4.3	-1.9 ss
West	3.7	4.0	4.9	4.4	5.4	6.5	7.0	6.5	5.4	6.6	6.9	6.9	7.2	8.8	§	7.9	-0.9
Population Density:																	
Large MSA	4.9	4.7	5.6	5.4	6.3	6.2	7.4	6.6	5.2	6.4	6.7	6.1	6.4	5.6	§	5.9	+0.3
Other MSA	5.0	5.1	5.3	5.3	6.7	7.5	6.4	7.1	6.6	5.9	5.5	5.4	5.1	6.9	§	6.1	-0.7
Non-MSA	5.1	5.5	5.6	4.6	4.3	5.0	5.2	4.7	4.8	5.8	6.1	7.0	6.1	6.9	§	4.9	-2.0 s
Parental Education: ^a																	
1.0–2.0 (Low)	4.6	5.6	4.8	3.5	4.6	5.1	5.2	7.8	5.8	6.1	6.4	6.6	4.8	7.3	§	6.4	-0.9
2.5–3.0	6.0	5.8	5.4	5.5	6.1	7.4	7.6	8.7	6.5	7.1	6.8	7.6	6.7	7.1	§	7.7	+0.7
3.5–4.0	4.9	5.0	4.9	5.5	6.6	6.9	7.2	6.9	6.4	5.9	6.2	6.4	6.0	6.7	§	6.6	-0.1
4.5–5.0	4.3	4.3	5.8	4.7	5.4	5.8	6.1	4.3	4.6	5.0	5.0	4.0	4.1	5.2	§	4.1	-1.1
5.5–6.0 (High)	3.6	3.8	4.9	4.1	4.7	6.0	4.4	4.8	4.5	4.9	3.7	3.5	4.4	4.4	§	3.8	-0.7
Race (2-year average): ^b																	
White	5.4	5.4	5.7	5.8	5.9	6.6	6.9	6.5	6.2	5.9	5.5	5.9	5.8	5.6	§	5.7	+0.1
African American	4.4	4.5	3.8	4.3	6.1	6.5	6.0	6.2	6.1	6.3	7.1	6.6	5.9	6.6	§	7.1	+0.5
Hispanic	3.2	3.2	3.4	3.6	4.1	4.6	4.8	6.1	5.7	4.8	5.1	5.0	4.3	4.8	§	5.4	+0.6

Source. The Monitoring the Future Study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence-of-use estimates for the two most recent classes is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cFor 8th and 10th graders only: In 2021, the question on marijuana use was changed in half of the questionnaire forms to include smoking, vaping, and edibles in the list of examples. Data presented here for 2021 based on the forms that included the original question wording. N is one half of N indicated.

TABLE 19
SYNTHETIC MARIJUANA: Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months											2019– 2021 change
	1991– 2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	4.4	4.0	3.3	3.1	2.7	2.0	1.6	2.7	1.6	1.3	-1.4* sss
Gender:												
Male	—	4.1	3.8	2.9	2.6	2.6	2.3	1.8	2.3	§	1.1	-1.3 s
Female	—	4.6	4.1	3.6	3.5	2.8	1.6	1.4	3.1	§	1.5	-1.6 s
College Plans:												
None or under 4 years	—	10.2	13.8	11.6	9.9	8.5	7.4	2.2	5.5	§	4.0	-1.5
Complete 4 years	—	3.9	3.3	2.6	2.6	2.2	1.4	1.4	2.3	§	0.9	-1.5 sss
Region:												
Northeast	—	2.7	3.5	3.5	1.6	1.5	0.8	1.5	2.3	§	0.2	-2.1 sss
Midwest	—	2.4	2.8	2.6	2.9	2.7	1.6	1.2	2.0	§	1.2	-0.9
South	—	5.6	4.2	2.6	3.1	2.8	2.5	1.6	2.8	§	1.9	-0.8
West	—	5.5	5.1	5.1	4.6	3.3	2.3	2.0	3.4	§	1.2	-2.2 s
Population Density:												
Large MSA	—	4.0	3.8	3.5	2.8	2.4	1.7	1.3	2.1	§	0.7	-1.4 ss
Other MSA	—	4.9	3.9	2.6	3.4	2.5	2.4	1.8	3.1	§	1.7	-1.4 s
Non-MSA	—	3.8	4.2	4.3	3.0	3.6	1.4	1.5	2.5	§	1.4	-1.1
Parental Education: ^a												
1.0–2.0 (Low)	—	5.9	8.5	6.9	7.5	5.3	3.8	2.9	6.1	§	1.2	-5.0 s
2.5–3.0	—	6.4	4.0	4.3	3.5	3.3	2.0	1.5	5.1	§	1.6	-3.5 ss
3.5–4.0	—	5.5	4.8	2.9	3.1	4.3	2.2	1.5	2.6	§	2.8	+0.2
4.5–5.0	—	2.9	3.0	2.4	2.1	1.0	1.3	1.6	1.4	§	0.6	-0.9
5.5–6.0 (High)	—	1.8	2.7	2.7	1.3	1.4	1.2	1.0	1.7	§	1.2	-0.6
Race/Ethnicity (2-year data): ^b												
White	—	—	3.1	2.9	2.2	2.1	1.8	1.1	1.5	§	1.3	-0.2
African American	—	—	2.8	2.3	2.0	2.5	2.9	2.4	1.9	§	1.0	-0.9
Hispanic	—	—	7.6	5.3	4.3	3.7	2.7	2.5	3.5	§	0.9	-2.6 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 20
SYNTHETIC MARIJUANA: Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months											2019– 2021 change
	1991– 2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	8.8	7.4	5.4	4.3	3.3	2.7	2.9	2.6	2.5	1.6	-1.0* s
Gender:												
Male	—	10.0	7.9	5.1	4.3	2.9	2.9	3.4	2.8	§	1.1	-1.7 s
Female	—	7.7	6.9	5.6	4.4	3.6	2.4	2.4	2.3	§	2.1	-0.2
College Plans:												
None or under 4 years	—	18.7	16.5	12.9	11.4	7.8	5.5	4.7	4.3	§	1.8	-2.5 ss
Complete 4 years	—	7.8	6.5	4.5	3.6	2.8	2.4	2.6	2.4	§	1.6	-0.7
Region:												
Northeast	—	7.7	6.9	3.8	4.6	4.2	1.3	3.3	2.6	§	3.4	+0.8
Midwest	—	9.2	6.3	5.8	3.4	2.7	2.7	2.6	2.3	§	2.0	-0.3
South	—	10.0	7.0	5.6	4.0	3.4	1.6	2.0	1.7	§	1.3	-0.3
West	—	7.4	9.4	6.1	5.7	3.0	5.0	4.6	4.7	§	0.5	-4.2 sss
Population Density:												
Large MSA	—	7.7	6.4	5.6	4.6	2.8	1.9	3.7	1.8	§	0.6	-1.2 s
Other MSA	—	8.3	7.6	4.9	4.1	3.5	3.0	2.5	3.1	§	2.2	-0.9
Non-MSA	—	11.4	8.4	6.4	4.4	3.4	3.1	2.9	2.7	§	2.2	-0.5
Parental Education: ^a												
1.0–2.0 (Low)	—	10.5	8.6	6.6	8.5	4.4	5.0	5.5	3.0	§	5.0	+2.1
2.5–3.0	—	13.4	10.5	7.5	6.1	4.4	2.2	3.0	2.5	§	2.3	-0.2
3.5–4.0	—	9.0	8.4	6.9	5.0	3.8	3.2	2.8	2.4	§	0.8	-1.6 s
4.5–5.0	—	7.2	4.9	3.3	3.0	2.0	2.1	1.9	3.4	§	0.9	-2.4 s
5.5–6.0 (High)	—	6.4	4.8	3.4	2.0	2.4	1.4	3.2	1.6	§	2.1	+0.5
Race/Ethnicity (2-year data): ^b												
White	—	—	8.4	6.2	4.5	3.5	2.6	2.2	2.7	§	2.5	-0.2
African American	—	—	3.6	4.1	4.1	3.4	3.0	2.2	1.5	§	1.7	+0.2
Hispanic	—	—	8.8	7.4	5.7	5.1	4.0	4.3	3.4	§	1.0	-2.4 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 21
SYNTHETIC MARIJUANA: Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months												
	2009–											2019–	
	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
Approximate weighted N =	—	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	11.4	11.3	7.9	5.8	5.2	3.5	3.7	3.5	3.3	2.4	1.8	-1.5* ss
Gender:													
Male	—	14.7	14.1	8.7	6.2	5.2	3.1	3.0	3.0	2.6	§	1.7	-0.9
Female	—	7.9	7.9	6.8	5.3	5.2	3.5	4.2	3.4	3.8	§	1.5	-2.3 ss
College Plans:													
None or under 4 years	—	18.9	16.7	11.8	5.8	8.0	5.7	5.6	5.4	5.1	§	2.2	-2.9 s
Complete 4 years	—	10.0	9.9	7.0	5.8	4.4	2.8	3.3	2.9	2.8	§	1.7	-1.1 s
Region:													
Northeast	—	11.9	10.7	7.4	5.1	4.3	3.9	2.4	2.2	4.4	§	1.3	-3.0 s
Midwest	—	13.5	14.2	7.6	6.6	5.3	3.8	3.8	3.8	1.8	§	1.9	+0.1
South	—	11.0	9.3	7.9	6.2	4.3	3.4	3.4	4.0	3.1	§	2.4	-0.7
West	—	9.5	11.1	8.4	5.0	7.1	3.2	4.8	3.1	4.8	§	1.3	-3.5 ss
Population Density:													
Large MSA	—	9.7	13.3	6.8	5.9	5.4	3.3	3.3	3.3	4.1	§	1.9	-2.2 s
Other MSA	—	12.3	10.3	8.2	6.3	5.0	3.9	3.2	3.7	2.9	§	1.7	-1.2
Non-MSA	—	11.7	10.8	8.6	4.6	5.2	2.8	5.5	3.4	2.9	§	1.8	-1.1
Parental Education: ^a													
1.0–2.0 (Low)	—	13.2	10.7	8.8	7.5	6.1	4.1	5.1	7.1	3.9	§	2.6	-1.3
2.5–3.0	—	12.9	12.3	7.0	7.0	5.9	4.8	4.8	3.6	5.0	§	1.3	-3.7 ss
3.5–4.0	—	12.1	12.8	8.4	5.6	5.2	3.7	4.7	2.6	3.3	§	1.8	-1.6 s
4.5–5.0	—	9.2	11.0	8.5	5.2	4.2	2.5	1.8	3.0	2.8	§	1.0	-1.8 s
5.5–6.0 (High)	—	11.3	8.5	5.1	4.2	4.4	2.3	2.6	2.1	1.8	§	3.5	+1.7
Race/Ethnicity (2-year data): ^b													
White	—	—	12.9	10.4	6.9	5.3	4.3	3.6	3.0	2.5	§	1.3	-1.2 ss
African American	—	—	6.0	6.5	5.8	3.7	2.8	2.8	2.8	2.2	§	2.6	+0.3
Hispanic	—	—	9.9	9.6	7.6	6.9	6.1	4.4	5.0	5.0	§	3.7	-1.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 22
INHALANTS: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	9.0	9.5	11.0	11.7	12.8	12.2	11.8	11.1	10.3	9.4	9.1	7.7	8.7	9.6	9.5
Gender:															
Male	9.0	9.2	10.4	11.2	11.5	10.3	10.5	10.6	9.5	8.9	8.4	7.6	7.7	8.8	7.8
Female	9.0	9.8	11.9	12.2	14.0	14.1	12.9	11.6	11.1	9.9	9.9	7.8	9.6	10.5	11.1
College Plans:															
None or under 4 years	15.0	15.6	17.7	18.3	19.6	18.2	18.1	20.9	17.9	17.2	15.1	14.5	16.5	17.9	15.3
Complete 4 years	8.1	8.8	10.2	10.9	11.9	11.4	11.2	10.2	9.5	8.6	8.6	7.1	8.0	8.7	9.0
Region:															
Northeast	8.0	8.6	11.3	12.0	13.1	11.7	12.1	9.1	9.8	8.7	9.4	6.9	9.0	8.9	9.1
Midwest	9.8	10.5	9.9	10.3	13.8	13.3	11.3	11.3	10.6	10.6	8.8	8.0	9.6	9.6	8.5
South	8.9	9.1	10.0	11.3	12.1	11.3	11.6	11.3	9.9	8.4	9.5	8.4	7.9	9.9	9.8
West	8.8	9.8	14.2	14.0	12.4	12.9	12.6	12.4	10.9	10.5	8.6	6.4	8.9	9.6	10.4
Population Density:															
Large MSA	9.9	9.1	10.8	11.6	11.7	11.4	10.4	8.6	8.8	8.3	8.3	7.1	8.0	8.8	8.8
Other MSA	8.5	10.3	12.3	13.1	13.7	13.4	11.5	11.1	10.1	9.4	8.9	8.0	8.5	10.3	10.7
Non-MSA	9.1	8.6	8.5	9.3	12.3	11.0	13.9	14.0	12.3	10.9	10.4	7.8	10.1	9.2	8.3
Parental Education: ^a															
1.0–2.0 (Low)	12.0	11.4	11.5	12.4	13.0	11.3	12.1	14.4	12.9	13.1	10.7	10.3	10.6	12.6	11.6
2.5–3.0	9.5	9.9	10.9	12.1	13.9	12.6	12.6	12.0	11.8	11.3	9.7	8.3	10.1	10.2	11.6
3.5–4.0	8.9	10.0	11.5	12.3	14.7	13.4	13.5	12.8	10.8	9.9	9.4	8.9	10.3	11.0	10.0
4.5–5.0	8.0	8.4	10.6	11.0	12.3	13.2	11.4	9.7	9.2	7.1	9.0	7.3	7.4	9.3	8.4
5.5–6.0 (High)	8.4	10.3	12.6	12.2	11.6	11.7	10.8	10.6	9.1	9.2	7.7	6.2	6.5	6.9	8.0
Race/Ethnicity (2-year average): ^b															
White	—	10.1	11.3	12.4	13.8	14.6	14.1	13.3	12.1	10.9	10.1	9.1	8.8	9.6	9.8
African American	—	4.4	4.6	5.3	5.0	4.2	3.8	4.2	4.2	4.3	4.8	5.0	4.9	5.4	5.8
Hispanic	—	10.4	11.5	12.5	13.3	12.7	11.4	11.5	12.7	12.2	11.0	9.9	9.6	10.6	11.0

Table continued on next page.

TABLE 22 (cont.)
INHALANTS: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months															2019-2021 change		
	2006	2007	2008	2009	2010	2011	2012	2013	2014 ^c	2015 ^c	2016 ^c	2017 ^c	2018 ^c	2019 ^c	2020 ^c		2021 ^c	
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700		
Total	9.1	8.3	8.9	8.1	8.1	7.0	6.2	5.2	5.3	4.6	3.8	4.7	4.6	4.7	6.1	4.8	+0.2*	
Gender:																		
Male	7.7	7.3	7.0	7.0	6.4	5.5	4.5	4.0	4.0	3.8	2.9	4.0	3.6	3.1	§	3.5	+0.4	
Female	10.5	9.3	11.0	9.3	9.5	8.6	7.9	6.3	6.2	5.2	4.7	5.3	5.4	6.2	§	5.5	-0.7	
College Plans:																		
None or under 4 years	17.3	18.1	19.1	16.7	13.1	12.8	11.2	10.9	10.7	7.6	8.6	8.3	8.0	5.8	§	6.6	+0.7	
Complete 4 years	8.3	7.5	8.0	7.5	7.7	6.7	5.9	4.8	4.8	4.3	3.3	4.3	4.2	4.5	§	4.6	+0.1	
Region:																		
Northeast	7.2	5.9	7.9	7.4	7.4	6.7	5.6	5.3	4.1	4.6	2.6	4.3	3.7	4.0	§	3.8	-0.2	
Midwest	9.0	8.1	9.1	7.6	7.9	5.8	5.5	4.1	4.9	4.7	4.5	4.5	4.7	4.3	§	4.2	-0.1	
South	9.1	9.1	9.6	8.9	8.3	7.1	6.4	5.1	5.6	5.0	3.8	4.7	4.3	5.3	§	4.9	-0.4	
West	10.5	8.8	8.4	7.9	8.3	8.4	7.2	6.3	5.9	3.7	3.8	5.0	5.5	4.5	§	6.0	+1.6	
Population Density:																		
Large MSA	8.8	8.6	8.7	7.1	7.2	6.7	6.6	5.1	5.1	4.2	3.4	4.3	4.6	4.1	§	3.8	-0.3	
Other MSA	9.0	8.3	8.9	8.8	8.6	7.3	6.5	5.3	4.9	4.7	3.7	4.6	4.4	5.0	§	5.6	+0.6	
Non-MSA	9.7	7.9	9.2	8.1	8.0	7.1	5.0	5.1	6.3	4.9	4.6	5.2	5.0	4.9	§	4.9	0.0	
Parental Education: ^a																		
1.0–2.0 (Low)	10.5	11.9	13.3	11.8	12.2	10.7	8.9	8.6	7.2	5.3	6.1	5.8	5.8	6.2	§	4.6	-1.6	
2.5–3.0	10.7	9.3	10.8	9.5	9.8	8.1	7.8	6.0	6.7	5.5	4.3	5.7	4.8	5.6	§	6.1	+0.5	
3.5–4.0	10.3	9.2	9.5	9.3	8.2	8.1	6.8	6.2	5.9	4.9	3.5	5.5	5.4	4.9	§	5.9	+1.0	
4.5–5.0	7.8	7.6	7.3	6.9	6.7	5.5	5.0	4.0	4.2	3.9	3.6	4.1	3.9	4.4	§	5.3	+0.9	
5.5–6.0 (High)	7.9	6.3	7.8	6.1	6.6	6.2	5.2	3.8	4.1	3.7	3.2	4.2	4.1	4.5	§	4.3	-0.2	
Race/Ethnicity (2-year average): ^b																		
White	9.7	8.9	8.5	8.6	8.0	7.1	5.9	4.9	4.4	4.3	3.9	4.2	4.5	4.7	§	6.0	+1.3	
African American	5.4	5.5	5.6	5.9	5.5	5.1	5.6	5.1	4.7	4.4	3.6	3.3	3.6	4.0	§	3.3	-0.7	
Hispanic	9.5	9.2	10.2	10.3	10.7	10.5	9.0	7.6	6.5	4.8	4.1	4.4	4.7	4.6	§	4.1	-0.5	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cBeginning in 2014 data are based on three of four forms; *N* is two thirds of *N* indicated.

TABLE 23
INHALANTS: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>
Total	7.1	7.5	8.4	9.1	9.6	9.5	8.7	8.0	7.2	7.3	6.6	5.8	5.4	5.9	6.0
Gender:															
Male	7.4	7.6	9.1	9.7	10.3	10.1	9.1	8.4	7.6	7.7	6.7	5.4	5.2	5.8	5.0
Female	6.6	7.5	7.7	8.6	8.9	8.9	8.2	7.6	6.9	7.0	6.5	6.0	5.6	6.1	6.9
College Plans:															
None or under 4 years	12.0	12.4	14.0	15.1	14.6	14.3	14.4	13.5	11.6	11.2	11.0	9.8	9.6	11.2	9.9
Complete 4 years	5.9	6.4	7.3	7.8	8.7	8.7	7.7	7.0	6.5	6.7	5.9	5.2	4.8	5.2	5.5
Region:															
Northeast	7.2	7.8	10.6	9.8	10.4	11.5	8.9	9.3	8.3	7.2	6.5	6.0	5.9	6.1	6.2
Midwest	7.5	8.0	8.3	8.4	10.4	9.8	8.3	6.7	8.4	7.5	6.5	5.8	6.1	5.7	6.1
South	7.2	6.6	7.3	9.0	9.4	9.1	8.8	8.3	6.5	7.4	6.8	5.4	4.6	5.6	5.6
West	6.2	8.0	8.4	9.9	8.1	8.0	8.5	7.8	6.1	7.2	6.7	6.2	5.5	6.6	6.1
Population Density:															
Large MSA	7.7	7.8	8.5	7.9	8.7	8.1	8.1	6.7	6.9	7.1	6.0	5.1	4.8	5.0	6.2
Other MSA	7.1	7.4	8.4	9.8	9.7	9.6	8.4	7.7	7.0	6.8	6.9	5.4	5.2	6.3	5.6
Non-MSA	6.5	7.5	8.6	9.1	10.5	11.0	9.8	10.1	8.3	8.5	7.0	7.4	7.1	6.5	6.5
Parental Education: ^a															
1.0–2.0 (Low)	7.0	8.2	10.2	8.7	9.4	10.8	9.3	9.7	8.7	8.5	7.6	5.6	5.8	6.2	6.3
2.5–3.0	8.0	7.9	9.1	9.5	11.0	9.9	8.5	9.1	8.0	8.1	7.5	6.0	6.3	6.9	6.9
3.5–4.0	7.5	8.3	8.3	9.6	10.2	10.1	9.4	8.1	6.9	7.4	5.9	6.3	5.8	5.7	6.2
4.5–5.0	6.4	6.5	7.2	8.7	9.4	8.4	8.3	7.1	6.7	6.5	6.8	5.6	5.1	5.7	5.8
5.5–6.0 (High)	6.6	6.7	8.2	8.2	7.0	10.1	8.2	6.7	7.2	7.2	5.5	5.2	4.4	5.1	5.3
Race/Ethnicity (2-year average): ^b															
White	—	8.3	8.8	9.6	10.6	11.0	10.4	9.6	8.9	8.4	7.9	7.1	6.6	6.5	6.6
African American	—	3.6	3.7	3.3	2.8	2.3	2.3	2.4	2.0	2.0	2.1	2.4	2.0	2.1	2.8
Hispanic	—	6.4	8.3	9.0	8.5	8.2	7.9	7.6	7.3	6.3	5.9	4.8	4.8	5.7	6.2

Table continued on next page.

TABLE 23 (cont.)
INHALANTS: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014 ^c	2015 ^c	2016 ^c	2017 ^c	2018 ^c	2019 ^c	2020 ^c	2021 ^c	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	6.5	6.6	5.9	6.1	5.7	4.5	4.1	3.5	3.3	2.9	2.4	2.3	2.4	2.8	2.9	2.0	-0.8* s
Gender:																	
Male	5.9	6.3	5.4	5.4	5.0	4.1	3.4	3.4	2.8	2.7	2.2	2.1	2.0	2.6	§	1.5	-1.1 ss
Female	7.1	7.0	6.3	6.7	6.3	4.8	4.8	3.5	3.8	3.0	2.6	2.5	2.7	3.0	§	2.2	-0.8
College Plans:																	
None or under 4 years	10.6	12.0	11.1	11.7	12.1	9.7	8.8	6.0	6.5	5.7	4.3	4.6	4.1	4.3	§	3.6	-0.8
Complete 4 years	5.9	5.9	5.3	5.4	5.0	3.9	3.6	3.2	3.0	2.6	2.2	2.0	2.2	2.5	§	1.7	-0.9 ss
Region:																	
Northeast	6.5	6.4	4.9	6.1	4.9	4.8	3.8	2.6	2.5	2.7	2.1	2.0	2.0	2.2	§	2.9	+0.6
Midwest	6.7	6.8	5.7	5.7	5.8	3.9	3.8	3.3	3.6	2.4	2.1	1.6	2.0	3.0	§	2.3	-0.7
South	5.7	6.1	5.6	5.2	5.8	4.3	4.2	3.2	3.8	2.9	2.7	3.0	2.9	2.9	§	2.0	-1.0
West	7.2	7.3	7.2	7.6	6.3	5.1	4.4	4.5	3.2	3.5	2.6	2.1	2.2	2.8	§	1.2	-1.6
Population Density:																	
Large MSA	6.0	6.0	5.3	6.1	5.6	4.5	3.9	3.3	3.2	3.2	2.2	2.2	1.9	2.5	§	1.9	-0.7
Other MSA	6.4	6.8	6.4	6.2	5.4	4.2	3.6	3.4	3.1	2.7	2.4	2.1	2.6	2.6	§	1.9	-0.6
Non-MSA	7.4	7.1	5.5	5.9	6.4	4.9	5.2	3.7	4.2	2.9	2.9	2.9	2.5	4.0	§	2.4	-1.5
Parental Education: ^a																	
1.0–2.0 (Low)	6.9	9.3	8.7	7.1	9.0	6.0	5.4	4.1	4.6	5.7	2.9	2.0	1.0	3.4	§	1.7	-1.7
2.5–3.0	6.2	7.5	6.7	7.4	6.9	5.4	4.7	4.5	4.7	3.5	2.3	2.4	2.4	4.2	§	2.7	-1.4
3.5–4.0	7.1	7.0	6.3	6.5	5.7	4.5	3.9	3.8	3.3	3.2	2.6	2.7	2.9	2.0	§	2.4	+0.3
4.5–5.0	5.5	5.8	5.0	4.9	4.8	3.7	3.9	2.9	2.7	2.3	2.2	2.3	3.1	2.4	§	1.7	-0.7
5.5–6.0 (High)	7.5	5.4	4.5	4.8	4.4	3.8	3.6	2.3	2.3	1.9	2.9	1.5	1.7	3.2	§	1.7	-1.5
Race/Ethnicity (2-year average): ^b																	
White	6.8	7.1	6.5	5.9	5.7	4.8	4.1	3.6	3.2	2.9	2.5	2.3	2.3	2.7	§	2.5	-0.2
African American	2.9	2.9	2.9	3.2	3.1	2.9	2.4	2.6	3.1	2.7	1.9	2.0	2.8	3.0	§	1.9	-1.1
Hispanic	6.4	6.4	7.2	7.9	7.5	6.8	5.7	4.8	4.2	3.7	2.8	2.2	1.9	2.1	§	1.7	-0.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cBeginning in 2014 data are based on three of four forms; N is two thirds of N indicated.

TABLE 24
INHALANTS: ^aTrends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	3.0	3.7	4.1	5.4	4.6	4.1	4.5	4.3	5.1	5.7	6.1	6.9	6.5	5.9	6.9
Adjusted ^b	—	—	—	—	8.9	7.9	6.1	6.6	6.2	7.2	7.5	8.9	8.1	7.1	6.9	7.5
Gender:																
Male	—	3.8	5.1	5.6	6.7	5.9	5.1	5.8	5.8	6.5	6.9	7.8	8.3	8.2	7.8	8.8
Female	—	2.0	2.4	2.8	4.2	3.5	3.2	3.1	2.8	3.8	4.5	4.7	5.6	4.9	4.0	4.9
College Plans:																
None or under 4 years	—	3.6	4.7	5.0	6.3	5.0	4.3	4.9	4.7	5.8	5.8	7.7	8.0	8.1	7.1	7.8
Complete 4 years	—	2.2	2.9	3.4	4.5	4.3	4.0	4.1	3.9	4.7	5.7	5.2	6.4	6.0	5.4	6.4
Region:																
Northeast	—	3.2	4.1	4.4	6.4	6.0	5.2	6.2	5.0	6.1	8.0	5.6	6.7	6.0	6.3	7.4
Midwest	—	2.6	4.2	4.8	5.9	4.6	3.8	3.6	4.5	5.0	5.8	6.7	8.6	7.2	6.7	8.0
South	—	3.8	3.3	3.6	4.3	3.4	3.2	3.8	3.8	4.6	4.2	5.7	6.1	6.8	5.5	6.4
West	—	1.7	3.0	3.6	4.9	4.9	4.7	4.4	4.3	5.3	5.4	6.6	6.2	5.6	4.8	5.7
Population Density:																
Large MSA	—	2.9	3.4	3.4	5.1	5.7	4.7	5.5	4.8	5.3	5.9	5.2	6.0	6.5	5.1	6.7
Other MSA	—	2.6	3.6	3.7	4.8	4.2	4.0	3.9	4.4	5.0	5.9	6.3	6.9	6.0	5.8	6.8
Non-MSA	—	3.4	4.2	5.3	6.2	4.4	3.7	4.4	3.9	5.2	5.4	6.6	7.8	7.5	6.8	7.4
Parental Education: ^c																
1.0–2.0 (Low)	—	3.7	3.9	4.5	5.2	3.6	3.6	3.2	3.1	4.5	4.2	4.9	4.6	5.3	5.9	5.0
2.5–3.0	—	3.1	4.1	4.0	5.0	4.8	4.0	4.8	4.0	5.2	5.6	6.1	6.8	6.3	5.5	6.9
3.5–4.0	—	3.1	3.4	4.1	5.1	4.7	4.0	4.6	4.9	5.6	5.5	6.2	7.1	5.8	6.1	7.2
4.5–5.0	—	2.7	3.0	3.9	5.8	4.3	4.4	4.4	5.2	5.0	7.0	6.9	7.2	7.0	5.7	7.4
5.5–6.0 (High)	—	3.7	4.2	5.0	7.2	5.8	4.9	6.0	4.7	5.6	6.8	6.4	8.7	9.1	6.8	7.6
Race/Ethnicity (2-year average): ^d																
White	—	—	3.6	4.3	5.1	5.3	4.7	4.7	4.8	5.1	5.9	6.5	7.3	7.6	7.0	7.2
African American	—	—	1.5	1.3	2.1	2.2	2.1	1.9	1.8	2.2	2.0	2.1	3.0	3.1	2.2	2.1
Hispanic	—	—	2.7	3.0	2.9	2.9	3.5	4.1	3.4	4.6	6.5	5.5	4.6	4.1	4.7	4.8

Table continued on next page.

TABLE 24 (cont.)
INHALANTS: ^aTrends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	6.6	6.2	7.0	7.7	8.0	7.6	6.7	6.2	5.6	5.9	4.5	4.5	3.9	4.2	5.0	4.5
Adjusted ^b	6.9	6.4	7.4	8.2	8.4	8.5	7.3	7.1	6.0	6.2	4.9	4.9	4.5	4.6	5.4	4.7
Gender:																
Male	8.2	8.0	9.2	9.6	9.9	9.1	8.3	7.5	6.5	6.8	5.5	5.8	5.2	4.8	6.2	5.1
Female	5.0	4.5	4.8	6.0	6.2	6.1	5.2	5.1	4.9	5.1	3.5	3.3	2.9	3.4	4.1	3.9
College Plans:																
None or under 4 years	7.7	7.7	8.0	9.0	9.7	8.2	8.0	7.9	6.5	6.7	6.6	6.3	5.1	5.6	7.3	6.8
Complete 4 years	6.3	5.7	6.7	7.4	7.4	7.3	6.5	5.7	5.4	5.5	3.9	4.2	3.5	3.8	4.6	3.9
Region:																
Northeast	6.7	6.0	8.9	10.3	10.3	10.8	9.4	8.0	6.2	6.3	5.7	5.4	5.1	4.3	6.0	5.9
Midwest	8.6	7.4	6.3	9.5	8.6	7.6	6.9	7.6	6.3	5.5	5.0	5.6	3.7	3.8	5.6	4.8
South	5.0	4.8	6.5	6.2	7.0	6.5	5.6	5.1	5.2	5.5	3.4	4.1	3.5	4.1	4.6	3.6
West	6.8	7.5	7.0	5.7	6.7	6.0	5.4	4.7	4.9	6.7	4.8	3.2	3.6	4.6	4.3	4.7
Population Density:																
Large MSA	5.2	6.0	7.4	7.6	8.5	7.8	5.9	5.5	4.8	6.3	4.4	4.4	3.5	4.0	5.0	4.1
Other MSA	7.8	6.6	7.3	7.7	7.8	7.9	6.5	6.1	5.3	4.9	4.3	4.4	4.4	4.1	5.1	4.5
Non-MSA	5.8	5.6	6.0	7.6	7.8	7.0	8.1	7.4	6.9	7.2	5.2	4.8	3.7	4.3	4.9	5.2
Parental Education: ^c																
1.0–2.0 (Low)	6.1	4.2	4.3	5.3	7.5	5.8	5.4	6.3	4.8	4.0	3.2	5.7	3.4	4.6	5.2	5.0
2.5–3.0	6.6	6.7	6.0	7.8	8.0	7.9	6.3	6.0	5.4	5.2	5.7	4.3	3.8	4.9	5.7	4.8
3.5–4.0	6.1	6.3	7.7	7.1	6.7	7.8	7.1	7.3	6.1	6.1	4.2	4.5	3.9	5.0	4.8	4.7
4.5–5.0	7.4	6.3	7.6	8.9	8.9	7.4	7.9	5.3	5.6	5.9	3.8	4.7	4.4	3.4	5.8	4.3
5.5–6.0 (High)	7.1	6.7	9.4	9.7	9.7	8.5	6.0	6.2	6.1	7.5	5.8	4.6	4.0	3.2	3.0	3.9
Race/Ethnicity (2-year average): ^d																
White	7.6	7.2	7.6	8.6	9.1	9.0	8.6	7.9	7.0	6.4	5.9	5.2	4.9	4.7	5.1	5.3
African American	2.7	2.5	2.2	2.4	2.6	2.2	1.9	1.7	1.4	1.9	2.2	1.9	1.5	1.3	1.9	1.9
Hispanic	5.4	6.0	5.7	5.5	5.8	5.9	4.7	4.5	5.5	6.3	4.5	3.4	2.7	2.9	5.3	5.7

Table continued on next page.

TABLE 24 (cont.)
INHALANTS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	3.7	3.8	3.4	3.6	3.2	2.9	2.5	1.9	1.9	1.7	1.5	1.6	1.9	1.1	1.8	-0.1*
Adjusted ^b	4.1	4.0	4.1	—	—	—	—	—	—	—	—	—	—	—	—	—
Gender:																
Male	3.9	4.4	3.6	4.7	3.3	3.1	2.9	2.1	2.1	1.5	1.4	1.8	2.2	§	1.2	-1.0 s
Female	3.4	3.2	3.2	2.5	3.0	2.6	2.2	1.8	1.6	1.7	1.6	1.2	1.5	§	1.9	+0.3
College Plans:																
None or under 4 years	4.4	6.1	4.7	4.5	6.2	3.7	3.9	3.1	4.1	2.2	2.7	2.1	3.7	§	2.5	-1.2
Complete 4 years	3.5	3.3	3.1	3.4	2.6	2.6	2.2	1.7	1.4	1.5	1.3	1.4	1.3	§	1.4	+0.1
Region:																
Northeast	3.9	3.5	3.7	3.4	3.1	3.8	2.6	1.9	1.4	2.0	1.4	2.0	1.9	§	1.3	-0.6
Midwest	3.9	3.0	3.7	3.9	3.2	2.5	2.3	2.4	1.5	1.7	1.3	1.1	2.3	§	1.3	-1.0
South	3.4	3.7	3.1	3.6	2.7	2.2	2.0	1.8	2.4	1.6	1.8	1.7	1.7	§	1.8	+0.1
West	3.7	5.0	3.5	3.8	4.0	3.4	3.3	1.7	1.8	1.5	1.4	1.6	2.0	§	2.5	+0.5
Population Density:																
Large MSA	2.8	3.1	2.4	4.0	3.3	3.2	2.6	1.7	1.4	1.5	1.3	1.5	2.3	§	1.0	-1.3 s
Other MSA	3.8	3.7	3.8	3.5	3.0	2.7	2.4	2.0	1.9	1.6	1.6	1.5	1.7	§	2.0	+0.3
Non-MSA	4.7	4.7	3.9	3.4	3.5	2.8	2.6	2.1	2.7	2.1	1.7	1.8	1.8	§	2.5	+0.6
Parental Education: ^c																
1.0–2.0 (Low)	3.6	3.6	3.6	3.0	4.3	3.0	2.7	2.6	2.4	1.7	2.3	0.7	1.9	§	1.2	-0.6
2.5–3.0	3.8	3.6	4.1	3.5	3.4	3.4	3.0	1.5	2.2	0.7	1.1	2.0	2.2	§	1.9	-0.3
3.5–4.0	3.9	4.4	3.1	3.6	3.8	2.3	2.7	1.8	2.1	2.0	1.9	1.9	2.1	§	2.7	+0.5
4.5–5.0	3.7	3.6	3.9	3.9	2.3	3.1	1.9	2.1	1.4	2.0	0.8	1.6	1.3	§	1.2	-0.1
5.5–6.0 (High)	3.4	3.4	1.9	3.2	2.0	2.8	2.2	2.1	1.5	1.8	1.9	1.2	1.9	§	1.3	-0.6
Race/Ethnicity (2-year average): ^d																
White	4.7	4.3	4.1	3.8	3.5	3.0	2.6	2.3	1.9	1.6	1.4	1.5	1.8	§	1.5	-0.3
African American	1.6	1.5	1.6	2.0	2.2	1.9	2.0	2.1	1.9	1.8	1.9	1.6	1.9	§	0.6	-1.2
Hispanic	4.3	3.3	3.0	3.6	4.5	4.1	3.7	2.8	2.0	2.1	1.8	1.5	1.3	§	0.7	-0.5

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on four of five forms in 1976–1988; N is four fifths of N indicated in Table S3. Data based on five of six forms in 1989–1998; N is five sixths of N indicated in Table S3. Data based on three of six forms beginning in 1999; N is three sixths of N indicated in Table S3.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aAll data are unadjusted for underreporting of amyl and butyl nitrites, except where otherwise noted.

^bAdjusted for underreporting of amyl and butyl nitrites. See text for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection), and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE 25
HALLUCINOGENS: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^a	2002	2003	2004	2005
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	1.9	2.5	2.6	2.7	3.6	4.1	3.7	3.4	2.9	2.8†	3.4	2.6	2.6	2.2	2.4
Gender:															
Male	2.2	2.6	2.8	3.0	4.0	4.3	4.0	3.7	3.3	3.2†	3.8	2.9	2.9	2.3	2.5
Female	1.6	2.3	2.3	2.4	3.3	3.7	3.2	2.9	2.4	2.5†	2.9	2.2	2.3	2.1	2.2
College Plans:															
None or under 4 years	5.1	7.2	7.1	6.7	9.6	9.6	10.1	9.2	9.4	7.7†	9.5	7.8	8.7	7.8	7.6
Complete 4 years	1.4	1.8	1.9	2.2	2.9	3.2	2.9	2.7	2.1	2.3†	2.6	2.0	1.9	1.7	1.8
Region:															
Northeast	1.5	1.6	1.9	2.9	3.4	3.7	2.8	2.4	2.6	2.3†	2.9	1.5	2.5	2.0	1.9
Midwest	1.6	2.4	1.7	2.2	3.8	3.9	3.8	3.5	3.4	3.5†	2.7	2.7	2.6	1.9	2.2
South	1.9	2.7	2.8	2.4	3.3	3.9	3.4	3.7	2.9	2.7†	4.0	2.9	2.6	2.4	2.9
West	2.8	3.2	4.2	3.9	4.2	5.1	4.8	3.5	2.4	2.7†	3.3	2.9	2.7	2.6	2.3
Population Density:															
Large MSA	2.1	2.2	2.2	3.1	4.0	3.8	3.3	2.9	2.5	2.3†	2.7	2.0	2.2	1.9	2.2
Other MSA	2.0	3.0	3.1	3.1	3.8	4.8	4.0	3.4	3.1	3.0†	3.6	2.4	2.5	2.6	2.4
Non-MSA	1.5	2.0	1.8	1.6	3.0	3.2	3.5	3.8	2.8	3.2†	3.6	3.5	3.3	2.1	2.7
Parental Education: ^b															
1.0–2.0 (Low)	3.9	3.7	3.5	3.1	5.1	4.8	5.0	5.0	4.8	5.4†	5.9	4.4	4.3	4.8	3.4
2.5–3.0	2.2	2.3	2.7	2.8	3.8	4.7	3.9	3.4	3.5	3.2†	3.9	3.2	3.2	2.5	3.3
3.5–4.0	1.6	2.5	2.8	2.8	4.1	4.1	3.8	3.7	2.8	2.8†	3.7	2.6	2.6	2.1	3.0
4.5–5.0	1.6	2.0	2.3	2.8	3.2	4.0	3.4	3.0	2.1	2.1†	2.4	2.1	2.0	2.0	1.8
5.5–6.0 (High)	1.4	2.4	2.0	2.5	3.2	3.5	3.5	3.1	2.5	3.1†	2.3	1.7	1.9	1.7	1.1
Race/Ethnicity (2-year average): ^c															
White	—	2.2	2.6	2.8	3.6	4.5	4.5	3.9	3.2	3.1	2.9†	2.7 ^d	2.8	2.5	2.4
African American	—	0.7	0.7	0.6	0.6	0.7	0.7	0.7	0.6	0.7	0.7†	0.7 ^d	0.9	1.2	1.3
Hispanic	—	3.8	4.1	4.0	4.0	4.1	4.2	4.6	4.5	4.0	3.4†	3.6 ^d	2.9	2.9	2.7

Table continued on next page.

TABLE 25 (cont.)
HALLUCINOGENS: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	<u>change</u>
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	2.1	1.9	2.1	1.9	2.2	2.2	1.6	1.6	1.3	1.3	1.2	1.1	1.4	1.3	1.7	1.0	-0.3*
Gender:																	
Male	2.1	2.1	2.2	2.0	2.5	2.6	1.7	1.7	1.5	1.3	1.3	1.1	1.5	1.2	§	0.9	-0.3
Female	2.0	1.8	2.0	1.8	2.0	1.7	1.5	1.6	1.0	1.2	1.1	1.1	1.1	1.4	§	0.9	-0.5
College Plans:																	
None or under 4 years	6.7	6.9	8.4	7.4	6.9	7.7	5.2	5.3	4.1	3.8	3.5	2.9	3.9	3.3	§	2.1	-1.3 s
Complete 4 years	1.6	1.5	1.6	1.4	1.8	1.8	1.3	1.4	1.0	1.1	0.9	0.9	1.0	1.0	§	0.8	-0.3
Region:																	
Northeast	1.4	1.0	1.4	1.4	1.8	1.1	1.5	1.0	0.6	0.6	0.8	0.8	0.7	0.5	§	0.5	0.0
Midwest	1.9	2.0	2.0	1.9	2.5	1.8	0.8	1.5	1.0	1.3	0.9	0.6	0.9	1.3	§	0.8	-0.5 ss
South	2.3	1.7	2.3	1.8	1.9	2.0	2.0	1.8	1.3	1.4	1.5	1.2	1.3	1.1	§	1.0	-0.1
West	2.4	2.8	2.5	2.3	3.0	3.9	1.8	1.9	1.9	1.6	1.3	1.7	2.4	2.0	§	1.6	-0.5
Population Density:																	
Large MSA	1.4	1.5	2.1	1.5	1.9	2.0	1.7	1.5	1.1	1.1	1.3	1.0	1.8	0.9	§	0.7	-0.3
Other MSA	2.4	2.0	2.3	2.1	2.6	2.6	1.8	1.7	1.4	1.6	1.1	1.3	1.2	1.4	§	1.1	-0.3
Non-MSA	2.5	2.4	1.7	2.0	2.1	1.7	1.1	1.5	1.4	0.9	1.5	1.1	1.0	1.5	§	1.4	-0.1
Parental Education: ^b																	
1.0–2.0 (Low)	2.9	3.0	4.2	3.9	3.1	3.7	2.5	2.3	2.1	2.0	2.4	1.9	2.0	2.2	§	2.1	-0.1
2.5–3.0	2.6	2.0	2.4	1.8	2.8	3.2	1.9	1.8	1.4	1.1	1.6	1.4	1.8	2.2	§	1.3	-0.9
3.5–4.0	2.3	2.3	2.1	2.0	2.4	2.3	2.0	1.5	1.5	1.5	1.3	1.2	1.6	1.4	§	1.3	-0.1
4.5–5.0	1.9	1.6	1.7	1.6	2.0	1.6	1.0	1.4	0.7	1.0	0.6	0.8	1.0	0.9	§	0.8	-0.1
5.5–6.0 (High)	1.5	1.1	1.9	1.5	1.4	1.4	1.2	1.6	1.3	0.8	0.8	0.8	0.8	0.9	§	0.7	-0.2
Race/Ethnicity (2-year average): ^c																	
White	2.5	2.2	2.1	2.1	2.1	2.1	1.6	1.3	1.2	1.0	1.0	1.0	1.1	1.1	§	1.3	+0.2
African American	1.1	0.7	0.8	0.8	0.7	0.9	1.1	1.2	0.9	0.7	1.0	0.9	0.7	0.8	§	0.8	+0.1
Hispanic	1.8	2.1	2.5	2.4	2.7	3.1	2.8	2.1	1.8	1.5	1.7	1.5	1.5	1.8	§	1.8	-0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. The 2001 data are based on the changed forms only; N is one half of N indicated. In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 26
HALLUCINOGENS: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^a	2002	2003	2004	2005	
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
Total	4.0	4.3	4.7	5.8	7.2	7.8	7.6	6.9	6.9	6.1‡	6.2	4.7	4.1	4.1	4.0	
Gender:																
Male	4.4	4.7	5.7	6.6	8.1	8.5	8.7	7.4	8.1	7.2‡	7.9	5.5	4.9	4.6	4.8	
Female	3.6	3.8	3.6	4.8	6.1	7.0	6.4	6.3	5.7	4.9‡	4.6	3.9	3.4	3.5	3.1	
College Plans:																
None or under 4 years	7.5	7.5	9.1	10.4	12.5	14.5	13.6	14.2	14.3	12.3‡	15.0	10.3	8.9	9.3	9.4	
Complete 4 years	3.3	3.6	3.7	4.8	6.2	6.6	6.5	5.6	5.7	5.1‡	4.8	3.9	3.3	3.3	3.2	
Region:																
Northeast	4.0	2.7	4.7	5.8	5.6	7.7	6.2	8.1	8.6	5.8‡	6.0	4.2	4.0	4.7	5.0	
Midwest	3.4	4.3	4.6	5.7	7.8	9.0	7.0	5.6	6.7	6.1‡	6.0	5.1	3.5	3.7	3.9	
South	3.6	3.9	3.6	5.1	7.3	7.5	8.3	7.6	6.5	5.7‡	5.3	4.0	3.9	3.6	3.5	
West	5.2	6.5	6.7	7.1	7.6	6.6	8.5	6.1	6.1	6.9‡	8.7	5.9	5.0	4.7	3.7	
Population Density:																
Large MSA	4.1	4.6	4.9	6.0	7.1	8.6	7.8	6.3	5.8	7.5‡	5.0	4.7	3.2	2.9	3.4	
Other MSA	4.8	4.4	4.9	6.4	8.0	8.2	7.9	7.6	7.8	5.8‡	6.6	4.9	4.4	5.0	4.3	
Non-MSA	2.5	3.7	4.1	4.4	5.5	6.0	6.7	6.3	6.5	5.1‡	6.8	4.4	4.6	3.9	4.2	
Parental Education: ^b																
1.0–2.0 (Low)	3.7	4.9	6.0	6.1	7.7	8.0	6.5	8.3	7.0	5.2‡	6.4	5.3	6.5	3.2	3.7	
2.5–3.0	4.3	4.2	4.5	5.5	7.6	8.5	7.3	8.2	8.1	5.8‡	6.7	5.2	4.7	4.8	4.5	
3.5–4.0	3.7	4.6	4.8	5.9	7.6	8.6	8.2	6.6	6.6	6.7‡	6.1	4.8	4.0	3.9	4.0	
4.5–5.0	4.1	3.8	4.5	5.5	6.6	6.9	8.2	6.1	6.6	6.3‡	6.0	4.0	3.5	3.7	4.0	
5.5–6.0 (High)	4.6	4.2	4.6	6.2	6.5	7.2	6.8	6.0	6.5	6.2‡	5.8	5.2	3.4	4.1	3.7	
Race/Ethnicity (2-year average): ^c																
White	—	4.9	5.1	5.6	7.1	8.6	8.9	8.4	8.2	7.6	6.6‡	5.6 ^d	5.3	4.9	4.7	
African American	—	0.2	0.6	1.1	1.2	0.9	1.0	1.1	1.0	1.0	1.3‡	1.4 ^d	1.0	0.8	1.2	
Hispanic	—	3.6	4.5	5.7	6.3	6.6	7.3	7.3	6.4	5.2	4.4‡	4.5 ^d	3.9	3.8	4.0	

Table continued on next page.

TABLE 26 (cont.)
HALLUCINOGENS: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															2019-2021	
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	4.1	4.4	3.9	4.1	4.2	4.1	3.5	3.4	3.3	3.1	2.9	2.8	2.7	3.1	3.4	2.2	-1.0* ss
Gender:																	
Male	4.5	5.4	4.7	5.0	5.2	4.9	4.6	4.4	3.3	3.4	3.5	3.4	3.2	4.0	§	2.1	-2.0 sss
Female	3.5	3.4	3.1	3.1	3.0	3.2	2.5	2.4	3.4	2.5	2.3	2.2	2.1	2.2	§	2.1	-0.1
College Plans:																	
None or under 4 years	10.0	10.0	10.4	10.5	11.0	10.3	10.5	6.5	8.3	7.0	6.8	7.5	6.0	5.5	§	4.4	-1.1
Complete 4 years	3.3	3.6	3.1	3.4	3.4	3.4	2.9	3.1	2.8	2.6	2.4	2.2	2.2	2.5	§	1.6	-0.9 sss
Region:																	
Northeast	4.0	4.4	3.5	3.4	4.0	4.0	3.0	2.7	3.4	3.2	2.1	1.6	2.1	1.9	§	2.1	+0.2
Midwest	4.1	4.7	3.8	4.4	3.5	3.5	3.2	2.9	3.2	2.2	2.6	3.0	2.6	3.0	§	2.4	-0.7
South	3.5	3.6	4.1	3.8	4.2	4.0	3.4	3.4	3.3	3.5	2.7	2.2	2.9	2.9	§	2.1	-0.9 s
West	5.1	5.1	4.2	4.9	5.1	5.0	4.5	4.4	3.3	3.2	4.2	4.4	3.0	4.5	§	2.3	-2.3 s
Population Density:																	
Large MSA	4.1	3.9	3.9	4.0	3.9	4.0	3.4	2.6	2.8	2.7	2.9	2.8	2.6	3.0	§	2.0	-1.0 s
Other MSA	4.1	4.7	4.2	4.3	4.3	4.4	3.7	3.9	3.7	3.4	3.0	2.8	2.8	3.2	§	2.1	-1.0 s
Non-MSA	3.8	4.2	3.4	3.8	4.1	3.7	3.4	3.2	3.4	2.8	2.7	2.8	2.6	3.3	§	2.7	-0.7
Parental Education: ^b																	
1.0–2.0 (Low)	5.3	5.4	5.0	5.8	5.3	5.7	4.9	4.4	3.0	3.9	2.9	2.9	3.5	3.4	§	3.6	+0.2
2.5–3.0	4.5	4.6	4.1	3.9	5.6	4.6	4.5	4.0	3.5	4.1	3.8	3.8	3.0	3.7	§	3.0	-0.7
3.5–4.0	3.8	4.5	4.2	4.3	4.2	4.1	3.3	3.9	4.3	3.7	3.0	3.3	2.7	3.5	§	2.6	-1.0
4.5–5.0	3.9	4.2	3.3	3.6	3.5	3.5	3.2	2.9	2.7	2.5	2.5	2.4	2.3	2.5	§	1.8	-0.7
5.5–6.0 (High)	4.0	3.9	3.3	4.5	3.3	3.5	2.8	3.0	2.7	2.0	2.6	2.0	2.8	2.4	§	1.3	-1.1
Race/Ethnicity (2-year average): ^c																	
White	4.7	4.9	4.8	4.5	4.5	4.4	4.1	3.6	3.5	3.3	3.2	3.2	3.0	3.1	§	2.6	-0.5
African American	1.4	1.0	1.1	1.4	1.6	1.4	1.4	1.5	1.5	2.0	1.6	1.0	0.8	1.0	§	2.6	+1.6 ss
Hispanic	3.5	2.8	3.6	4.3	4.3	4.5	4.2	3.5	3.1	3.0	3.0	3.0	3.0	3.3	§	2.4	-1.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. '‡' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. The 2001 data are based on the changed forms only; N is one half of N indicated. In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 27
HALLUCINOGENS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Approximate weighted N =	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	11.2	9.4	8.8	9.6	9.9	9.3	9.0	8.1	7.3	6.5	6.3	6.0	6.4	5.5	5.6	5.9
Adjusted ^b	—	—	—	—	11.8	10.4	10.1	9.0	8.3	7.3	7.6	7.6	6.7	5.8	6.2	6.0
Gender:																
Male	13.7	11.6	10.8	11.6	11.8	11.7	10.9	9.6	8.6	7.9	8.1	7.2	7.5	7.2	7.4	7.7
Female	9.0	6.9	6.5	7.3	7.6	6.7	6.8	6.1	5.5	4.7	4.4	4.7	5.2	3.7	3.6	3.8
College Plans:																
None or under 4 years	—	11.2	10.6	11.0	11.3	11.2	10.7	9.5	8.9	8.3	7.7	7.4	7.9	6.4	7.1	6.6
Complete 4 years	—	6.9	6.4	7.3	7.5	7.1	7.4	6.2	5.4	4.7	5.0	4.7	5.4	4.7	4.8	5.3
Region:																
Northeast	13.2	10.9	10.6	13.0	12.9	12.2	12.9	11.4	8.7	11.3	9.9	7.9	7.5	5.8	5.6	6.6
Midwest	13.0	10.3	9.7	10.7	11.1	11.3	10.3	9.1	8.9	6.0	6.8	6.6	6.9	5.3	6.6	5.7
South	8.5	7.4	6.8	6.3	5.7	5.4	4.1	4.6	5.2	3.9	3.2	3.3	4.8	5.2	4.9	5.0
West	10.2	9.3	8.2	9.6	11.0	9.2	10.4	7.8	6.3	7.0	6.3	7.2	7.4	6.0	5.5	6.9
Population Density:																
Large MSA	13.9	11.1	9.9	11.9	12.3	11.6	12.0	10.9	9.2	8.8	8.3	7.6	7.9	6.5	5.4	5.7
Other MSA	12.1	9.8	9.1	9.3	10.5	9.8	9.0	7.6	7.6	6.3	6.1	5.9	6.3	6.0	5.9	6.6
Non-MSA	8.5	7.7	7.5	8.3	7.1	7.1	6.8	6.5	5.3	5.0	5.0	4.9	5.3	3.5	5.0	4.5
Parental Education: ^c																
1.0–2.0 (Low)	8.9	7.4	6.8	7.7	7.1	8.0	6.7	6.5	6.5	5.4	4.8	5.4	5.8	4.9	4.2	3.8
2.5–3.0	10.2	10.0	9.1	9.6	9.6	9.5	8.9	8.0	6.8	6.7	6.4	6.0	6.2	4.2	4.9	4.6
3.5–4.0	10.9	9.8	9.2	9.7	9.7	9.2	9.2	8.6	7.7	6.3	7.2	6.3	6.0	4.8	5.6	6.5
4.5–5.0	11.1	10.1	8.8	10.2	10.9	9.1	9.4	7.8	7.0	5.9	6.2	5.5	6.8	6.7	6.6	6.8
5.5–6.0 (High)	8.9	9.4	9.5	10.2	11.7	9.9	10.6	9.0	7.0	7.6	4.3	5.9	7.2	7.2	7.0	8.2
Race/Ethnicity (2-year average): ^d																
White	—	—	9.8	9.9	10.5	10.3	10.0	9.3	8.3	7.5	7.0	6.7	6.8	6.8	6.4	6.7
African American	—	—	2.4	2.3	2.0	1.9	1.9	1.8	2.2	1.7	1.2	1.6	1.5	1.0	0.9	0.8
Hispanic	—	—	7.9	7.2	7.0	7.1	7.0	7.7	6.6	5.2	5.7	5.7	5.0	4.0	3.2	3.3

Table continued on next page.

TABLE 27 (cont.)
HALLUCINOGENS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^o	2002	2003	2004	2005	2006
Approximate weighted N =	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	5.8	5.9	7.4	7.6	9.3	10.1	9.8	9.0	9.4	8.1‡	9.1	6.6	5.9	6.2	5.5	4.9
Adjusted ^b	6.1	6.2	7.8	7.8	9.7	10.7	10.0	9.2	9.8	8.7‡	9.7	7.2	6.5	6.4	5.9	5.3
Gender:																
Male	7.5	7.1	8.9	9.2	11.9	12.4	12.0	11.0	11.4	9.6‡	11.1	8.4	7.8	8.4	7.4	6.2
Female	3.9	4.7	5.6	5.8	6.3	7.3	7.4	6.8	7.4	6.3‡	6.8	4.7	3.8	3.8	3.4	3.6
College Plans:																
None or under 4 years	7.0	7.8	8.1	8.4	11.9	12.1	11.3	12.0	10.5	10.3‡	10.4	9.8	8.2	7.7	7.8	7.0
Complete 4 years	5.3	5.1	6.9	7.0	8.2	9.0	9.0	7.8	8.7	7.0‡	8.0	5.5	5.0	5.4	4.7	4.4
Region:																
Northeast	7.0	7.1	9.0	9.0	10.1	13.3	13.9	10.7	9.8	9.3‡	9.8	9.1	7.8	7.0	5.5	7.1
Midwest	6.5	5.9	6.8	8.1	9.2	8.8	7.6	8.4	9.8	7.0‡	11.4	6.4	5.4	5.3	6.7	4.9
South	3.7	4.7	5.9	6.7	8.8	8.9	9.2	8.5	8.6	6.9‡	5.8	5.6	4.9	5.7	5.2	4.1
West	7.3	7.3	9.2	7.1	9.6	10.5	9.5	9.1	10.0	10.5‡	10.8	6.2	6.3	7.4	4.4	4.3
Population Density:																
Large MSA	5.1	6.2	7.3	8.1	11.0	10.5	8.8	8.7	8.4	8.9‡	11.5	6.8	4.4	5.6	5.5	6.0
Other MSA	7.7	6.0	8.1	8.6	9.5	11.4	11.2	9.9	10.4	8.3‡	8.7	7.2	7.2	7.2	5.9	4.5
Non-MSA	3.3	5.5	6.3	5.1	7.0	7.4	8.3	7.4	8.8	7.0‡	7.3	5.2	5.5	4.8	4.5	4.1
Parental Education: ^c																
1.0–2.0 (Low)	4.9	3.6	4.9	5.0	7.2	7.4	7.3	7.9	9.0	7.0‡	6.3	5.1	5.3	4.9	3.0	4.9
2.5–3.0	4.9	5.6	5.9	7.0	8.7	8.8	8.5	8.8	8.6	7.4‡	9.1	6.6	4.9	5.6	5.7	5.1
3.5–4.0	6.2	6.0	7.5	8.0	9.5	10.3	9.9	9.5	10.6	8.2‡	9.4	7.1	6.4	6.6	5.4	5.1
4.5–5.0	6.1	6.2	8.9	7.7	9.6	10.5	10.4	8.6	9.3	7.7‡	8.6	6.7	6.5	6.4	5.7	4.6
5.5–6.0 (High)	7.3	7.4	8.9	9.0	9.5	11.4	11.6	9.4	8.4	9.6‡	8.8	5.9	5.3	6.1	5.7	4.6
Race/Ethnicity (2-year average): ^d																
White	6.8	6.9	7.9	8.6	9.5	10.8	11.6	11.3	10.7	9.9	9.0‡	8.4 ^f	7.2	7.0	6.7	6.0
African American	0.6	0.7	0.8	1.2	1.2	1.7	1.9	1.4	1.2	1.6	1.4‡	1.2 ^f	1.3	1.2	1.3	1.5
Hispanic	4.4	4.6	5.3	5.8	7.1	8.3	7.3	6.8	7.9	9.6	7.8‡	6.0 ^f	4.6	4.1	3.9	3.8

Table continued on next page.

TABLE 27 (cont.)
HALLUCINOGENS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014 ^g	2015 ^g	2016 ^g	2017 ^g	2018 ^g	2019 ^g	2020 ^g	2021 ^g	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	5.4	5.9	4.7	5.5	5.2	4.8	4.5	4.0	4.2	4.3	4.4	4.3	4.6	5.3	4.1	-0.5*
Adjusted ^b	5.8	6.1	5.2	6.0	5.8	5.0	4.9	—	—	—	—	—	—	—	—	—
Gender:																
Male	7.5	7.8	6.2	7.9	7.0	6.0	5.9	5.5	5.2	5.7	5.9	5.8	6.1	§	5.0	-1.2
Female	3.3	3.9	3.1	2.9	3.3	3.1	2.8	2.6	2.9	2.8	3.0	2.3	2.7	§	2.7	0.0
College Plans:																
None or under 4 years	8.3	8.4	5.6	8.4	7.7	5.8	6.0	5.8	6.1	4.7	6.1	6.7	6.6	§	5.9	-0.8
Complete 4 years	4.6	5.2	4.3	4.8	4.6	4.5	4.0	3.5	3.6	4.0	3.7	3.4	3.8	§	3.4	-0.4
Region:																
Northeast	6.7	8.0	4.7	6.2	5.7	5.2	4.5	4.4	3.6	5.0	4.2	4.8	4.0	§	4.1	+0.0
Midwest	5.3	4.5	5.9	5.4	5.2	4.5	4.9	4.0	3.8	4.9	3.6	4.5	4.1	§	3.4	-0.7
South	4.2	5.7	3.9	4.5	3.9	4.2	3.9	4.0	3.9	3.1	3.9	3.4	4.4	§	3.4	-1.0
West	6.5	6.2	4.6	6.6	6.7	5.5	4.8	3.6	5.3	5.5	6.2	5.1	6.3	§	5.6	-0.7
Population Density:																
Large MSA	6.2	6.9	5.1	5.6	4.9	5.4	4.5	4.2	4.4	4.3	5.2	3.8	4.2	§	4.5	+0.2
Other MSA	5.1	5.7	4.7	5.9	5.6	5.0	5.0	4.1	4.5	4.8	4.4	4.4	4.6	§	4.0	-0.6
Non-MSA	4.8	5.3	4.2	4.3	4.4	3.2	3.1	3.3	2.7	3.4	3.3	4.8	5.1	§	3.5	-1.7
Parental Education: ^c																
1.0–2.0 (Low)	3.9	4.4	3.6	3.7	4.9	3.3	4.4	3.0	4.8	4.0	4.1	4.0	4.1	§	3.1	-1.1
2.5–3.0	4.9	5.0	4.5	5.0	4.3	5.1	4.7	4.1	4.1	4.0	4.6	4.4	4.6	§	4.5	-0.1
3.5–4.0	5.5	5.5	4.6	6.2	5.4	4.8	4.6	4.4	4.3	4.5	4.7	4.3	5.1	§	3.7	-1.4
4.5–5.0	5.4	6.7	5.3	5.5	4.9	5.0	3.9	3.8	3.7	4.5	4.0	3.6	4.1	§	4.4	+0.2
5.5–6.0 (High)	6.7	7.2	4.6	5.4	5.7	4.5	4.7	4.1	4.1	4.0	4.5	4.4	4.0	§	3.4	-0.6
Race/Ethnicity (2-year average): ^d																
White	6.2	6.9	6.3	6.1	6.4	6.0	5.2	4.6	4.2	4.3	4.8	4.8	4.5	§	4.1	-0.5
African American	1.4	1.4	1.2	1.3	1.7	1.6	1.5	1.9	2.0	1.7	1.9	1.8	1.9	§	3.0	+1.1
Hispanic	3.9	3.5	3.3	3.3	3.0	2.9	3.5	3.5	4.0	4.7	3.9	3.4	4.0	§	5.2	+1.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aAll data are unadjusted for the underreporting of PCP, unless otherwise indicated.

^bAdjusted for the underreporting of PCP. See text for details.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^dEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^eIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. The 2001 data are based on the changed forms only; N is one half of N indicated. In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^fDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

^gIn 2014 questions on use of LSD and hallucinogens other than LSD, the two components of hallucinogen use, were each dropped from one of six forms. Beginning in 2014 data are based on five of six forms; N is five sixths of N indicated.

TABLE 28
LSD: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>
Total	1.7	2.1	2.3	2.4	3.2	3.5	3.2	2.8	2.4	2.4	2.2	1.5	1.3	1.1	1.2
Gender:															
Male	2.0	2.1	2.5	2.6	3.4	3.7	3.5	3.2	2.7	2.6	2.3	1.7	1.4	1.1	1.2
Female	1.3	2.0	2.1	2.1	2.9	3.2	2.8	2.4	2.0	2.2	2.1	1.3	1.1	1.1	1.1
College Plans:															
None or under 4 years	4.5	6.4	6.4	6.2	8.5	8.2	9.3	7.8	8.2	6.7	5.8	5.7	4.3	4.7	4.1
Complete 4 years	1.2	1.5	1.6	1.8	2.5	2.7	2.5	2.2	1.7	2.0	1.8	1.1	0.9	0.7	0.8
Region:															
Northeast	1.3	1.4	1.8	2.6	2.9	2.9	2.3	2.1	2.2	1.9	1.9	0.9	1.4	1.0	0.9
Midwest	1.4	1.8	1.4	1.7	3.5	3.4	3.3	2.5	2.7	3.0	1.7	1.8	1.2	0.9	0.9
South	1.8	2.4	2.4	2.1	2.8	3.4	3.0	3.2	2.5	2.4	2.7	1.8	1.3	1.2	1.5
West	2.2	2.9	3.7	3.3	3.8	4.3	4.3	3.2	1.9	2.3	2.0	1.2	1.2	1.2	1.1
Population Density:															
Large MSA	1.9	2.0	2.0	2.7	3.6	3.2	2.9	2.6	2.2	2.0	1.8	1.3	1.3	1.1	1.0
Other MSA	1.7	2.5	2.8	2.8	3.3	4.1	3.6	2.9	2.7	2.6	2.3	1.5	1.2	1.3	1.1
Non-MSA	1.3	1.6	1.4	1.3	2.4	2.6	2.8	2.9	1.9	2.8	2.4	1.9	1.4	0.7	1.4
Parental Education: ^a															
1.0–2.0 (Low)	3.5	3.1	3.1	2.8	4.6	4.4	4.7	4.4	3.7	4.9	4.1	3.4	2.3	2.8	1.7
2.5–3.0	1.8	2.1	2.3	2.6	3.1	4.0	3.2	2.8	2.7	2.9	2.6	2.0	1.4	1.2	1.7
3.5–4.0	1.4	2.0	2.4	2.4	3.6	3.5	3.4	3.1	2.4	2.2	2.4	1.3	1.3	1.0	1.5
4.5–5.0	1.4	1.5	2.1	2.1	2.6	3.4	2.9	2.5	1.9	1.8	1.5	1.2	1.1	1.0	0.6
5.5–6.0 (High)	1.3	2.0	2.0	2.1	2.9	3.0	2.9	2.4	1.9	2.3	1.3	0.8	0.8	0.6	0.4
Race/Ethnicity (2-year average): ^b															
White	—	1.9	2.3	2.5	3.1	3.9	3.9	3.2	2.6	2.6	2.6	2.0	1.5	1.1	1.0
African American	—	0.5	0.4	0.5	0.5	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.6	0.8	0.9
Hispanic	—	3.3	3.7	3.6	3.3	3.5	3.9	4.2	3.9	3.5	3.2	2.4	1.7	1.8	1.7

Table continued on next page.

TABLE 28 (cont.)
LSD: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																
	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	2019-2021 <u>change</u>
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	0.9	1.1	1.3	1.1	1.2	1.1	0.8	1.0	0.7	0.9	0.8	0.9	0.9	0.9	1.1	0.7	-0.2*
Gender:																	
Male	0.8	1.1	1.4	1.1	1.2	1.4	0.8	1.0	0.8	0.8	0.9	0.9	1.0	0.8	§	0.7	-0.1
Female	1.0	1.0	1.2	1.1	1.1	0.8	0.8	1.0	0.6	0.9	0.6	0.8	0.8	1.0	§	0.5	-0.5 ss
College Plans:																	
None or under 4 years	3.4	3.9	5.2	4.9	3.7	3.9	3.3	3.8	2.5	2.5	2.3	2.4	2.8	2.5	§	1.6	-0.9
Complete 4 years	0.7	0.8	1.0	0.8	0.9	0.9	0.6	0.8	0.6	0.7	0.6	0.7	0.6	0.7	§	0.5	-0.2
Region:																	
Northeast	0.5	0.6	0.8	0.8	1.0	0.6	0.7	0.6	0.3	0.3	0.6	0.5	0.5	0.5	§	0.2	-0.3
Midwest	0.7	1.2	1.3	1.1	1.4	0.9	0.3	0.9	0.6	1.0	0.6	0.6	0.4	1.1	§	0.6	-0.5 ss
South	1.1	1.0	1.4	1.0	1.0	1.0	1.1	1.2	0.8	0.9	1.1	0.9	0.8	0.9	§	0.7	-0.2
West	1.0	1.4	1.4	1.6	1.4	2.1	0.9	1.1	1.0	1.0	0.6	1.4	1.8	1.3	§	1.1	-0.1
Population Density:																	
Large MSA	0.6	0.7	1.4	1.0	1.0	1.0	0.9	1.0	0.7	0.8	1.1	0.8	1.3	0.7	§	0.5	-0.2
Other MSA	1.1	1.2	1.3	1.2	1.4	1.3	0.8	1.1	0.7	1.1	0.6	1.0	0.9	1.1	§	0.8	-0.3
Non-MSA	1.0	1.3	1.0	1.1	1.0	0.9	0.6	0.8	0.7	0.5	0.9	0.7	0.5	0.9	§	0.7	-0.2
Parental Education: ^a																	
1.0–2.0 (Low)	1.8	1.8	2.4	2.6	1.6	1.7	1.6	1.3	1.4	1.5	1.8	1.5	1.9	1.9	§	0.8	-1.1
2.5–3.0	1.0	1.1	1.6	1.1	1.1	1.7	1.1	1.0	0.8	0.7	0.9	1.2	1.3	1.5	§	0.9	-0.6
3.5–4.0	1.0	1.0	1.1	1.0	1.4	1.2	0.7	0.8	1.0	1.1	0.8	0.9	1.0	0.9	§	0.9	0.0
4.5–5.0	0.7	1.0	1.0	0.8	1.0	0.8	0.4	1.0	0.4	0.7	0.4	0.6	0.5	0.7	§	0.6	-0.1
5.5–6.0 (High)	0.7	0.6	1.3	0.9	0.8	0.8	0.7	1.1	0.6	0.6	0.5	0.5	0.5	0.5	§	0.6	+0.1
Race/Ethnicity (2-year average): ^b																	
White	1.0	1.0	1.1	1.1	1.1	1.0	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.8	§	0.8	0.0
African American	0.6	0.3	0.6	0.7	0.5	0.6	0.6	0.8	0.6	0.4	0.7	0.6	0.5	0.6	§	0.6	0.0
Hispanic	1.0	1.3	1.6	1.5	1.6	1.8	1.4	1.2	1.2	1.0	1.2	1.1	1.2	1.3	§	1.5	+0.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 29
LSD: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>
Total	3.7	4.0	4.2	5.2	6.5	6.9	6.7	5.9	6.0	5.1	4.1	2.6	1.7	1.6	1.5
Gender:															
Male	3.9	4.3	5.1	5.9	7.4	7.6	7.6	6.3	7.0	5.9	5.1	3.1	1.9	1.8	1.9
Female	3.4	3.6	3.2	4.3	5.5	6.2	5.8	5.4	5.1	4.3	3.1	2.0	1.6	1.4	1.0
College Plans:															
None or under 4 years	6.8	7.0	8.4	9.4	11.1	13.1	12.8	12.4	13.1	11.1	9.9	6.0	4.4	4.5	4.1
Complete 4 years	3.0	3.4	3.3	4.2	5.6	5.8	5.7	4.7	4.9	4.1	3.2	2.0	1.3	1.2	1.1
Region:															
Northeast	3.6	2.6	3.8	5.1	4.7	6.4	5.2	7.1	7.5	4.1	4.0	2.2	1.8	1.7	2.3
Midwest	3.2	4.1	4.4	5.2	7.3	8.3	6.0	4.5	6.0	5.4	4.3	2.8	1.7	1.5	1.2
South	3.3	3.7	3.2	4.6	6.8	6.8	7.9	6.5	5.8	5.0	3.5	2.3	1.7	1.4	1.4
West	4.8	5.9	6.1	6.3	6.5	5.7	7.4	5.2	5.1	5.9	5.3	3.2	1.7	1.9	1.0
Population Density:															
Large MSA	3.8	4.4	4.4	5.4	6.6	7.6	7.0	5.4	4.9	6.4	3.8	2.6	1.3	1.1	1.2
Other MSA	4.4	4.1	4.4	5.9	7.1	7.4	7.0	6.6	6.7	4.8	4.1	2.7	1.7	1.9	1.5
Non-MSA	2.3	3.5	3.7	3.7	5.0	5.2	6.0	5.0	5.9	4.4	4.6	2.3	2.3	1.5	1.8
Parental Education: ^a															
1.0–2.0 (Low)	3.1	4.4	5.5	5.5	6.9	7.6	5.9	7.9	6.3	4.9	3.9	2.7	3.7	1.3	1.8
2.5–3.0	4.0	4.2	4.2	5.1	6.9	7.6	6.6	7.0	7.3	5.1	4.8	2.9	2.0	1.7	1.7
3.5–4.0	3.4	4.1	4.2	5.3	6.9	7.9	7.4	5.6	5.8	5.6	4.3	2.8	1.7	1.4	1.6
4.5–5.0	3.8	3.6	3.9	4.8	6.0	6.0	7.0	5.0	5.7	5.0	3.8	2.0	1.2	1.4	1.2
5.5–6.0 (High)	4.2	3.9	3.9	5.4	5.9	5.8	6.0	4.6	5.3	5.0	3.5	2.4	1.2	1.6	1.4
Race/Ethnicity (2-year average): ^b															
White	—	4.6	4.6	5.0	6.4	7.7	7.9	7.3	7.0	6.5	5.2	3.8	2.4	1.9	1.7
African American	—	0.2	0.5	0.9	1.0	0.8	0.9	1.0	0.9	0.9	0.9	0.6	0.4	0.4	0.6
Hispanic	—	3.2	4.1	5.0	5.7	6.1	6.7	6.6	5.6	4.6	3.7	2.9	2.4	1.7	1.6

Table continued on next page.

TABLE 29 (cont.)
LSD: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															2019-2021 change	
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	1.7	1.9	1.8	1.9	1.9	1.8	1.7	1.7	1.9	2.0	2.1	2.1	2.0	2.3	2.5	1.5	-0.8* ss
Gender:																	
Male	2.1	2.3	2.2	2.3	2.4	2.2	2.3	2.3	2.1	2.2	2.6	2.5	2.4	2.7	§	1.4	-1.3 sss
Female	1.3	1.5	1.3	1.3	1.3	1.3	1.1	1.0	1.8	1.5	1.6	1.6	1.5	1.8	§	1.4	-0.4
College Plans:																	
None or under 4 years	5.1	4.3	5.1	5.9	5.1	4.2	6.3	3.5	5.2	4.2	4.7	5.3	4.4	4.3	§	3.0	-1.3
Complete 4 years	1.2	1.5	1.3	1.4	1.5	1.6	1.3	1.5	1.5	1.7	1.8	1.7	1.6	1.8	§	1.1	-0.8 sss
Region:																	
Northeast	1.7	1.8	1.5	1.6	2.0	1.8	1.4	1.7	1.7	2.1	1.6	1.1	1.7	1.4	§	1.3	-0.2
Midwest	1.6	2.2	2.0	2.3	1.4	1.7	1.6	1.5	2.0	1.4	2.0	2.6	1.9	2.2	§	1.6	-0.6
South	1.7	1.6	1.8	1.6	2.1	2.1	1.7	1.5	2.2	2.4	1.9	2.0	2.0	2.2	§	1.5	-0.7 s
West	1.8	1.9	1.7	2.0	2.0	1.8	2.1	1.9	1.5	1.6	3.0	2.5	2.1	3.0	§	1.6	-1.4
Population Density:																	
Large MSA	1.6	1.6	1.5	1.7	1.7	1.8	1.8	1.3	1.7	1.7	2.3	2.1	1.8	2.2	§	1.5	-0.7 s
Other MSA	1.7	2.1	1.9	2.2	1.9	1.8	1.7	2.2	2.1	2.4	2.1	2.1	2.2	2.3	§	1.4	-0.9
Non-MSA	1.8	1.7	1.7	1.4	2.1	1.9	1.8	0.9	1.6	1.4	1.9	1.9	1.7	2.3	§	1.7	-0.7
Parental Education: ^a																	
1.0–2.0 (Low)	2.5	2.9	2.2	2.5	2.4	3.0	2.0	2.2	1.7	2.5	2.4	2.0	2.4	2.8	§	2.4	-0.5
2.5–3.0	2.1	2.0	1.6	1.9	2.5	1.8	2.6	1.8	1.8	2.0	2.9	2.7	2.1	2.8	§	2.2	-0.6
3.5–4.0	1.6	1.8	1.9	2.0	1.8	1.8	1.4	1.7	2.3	2.4	2.1	2.6	2.0	2.5	§	1.8	-0.7
4.5–5.0	1.3	1.7	1.5	1.5	1.6	1.7	1.6	1.5	1.6	1.5	1.9	1.7	1.8	1.7	§	1.0	-0.7 s
5.5–6.0 (High)	1.3	1.7	1.9	2.2	1.9	1.6	1.3	1.7	1.6	1.6	1.8	1.6	2.0	1.5	§	0.9	-0.6
Race/Ethnicity (2-year average): ^b																	
White	1.8	2.0	2.0	1.9	2.0	2.0	1.9	1.9	2.0	2.1	2.2	2.3	2.2	2.2	§	1.8	-0.4
African American	0.7	0.6	0.7	0.9	1.1	1.0	0.8	0.8	1.1	1.2	0.8	0.9	0.7	0.7	§	2.1	+1.4 s
Hispanic	1.6	1.4	1.8	2.0	1.7	1.9	1.9	1.5	1.4	1.6	2.2	2.2	2.3	2.5	§	1.6	-0.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 30
LSD: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	7.2	6.4	5.5	6.3	6.6	6.5	6.5	6.1	5.4	4.7	4.4	4.5	5.2	4.8	4.9	5.4
Gender:																
Male	9.6	7.9	7.1	7.8	8.0	8.1	8.0	7.4	6.7	5.8	5.9	5.5	6.4	6.5	6.5	7.1
Female	5.6	4.6	3.9	4.5	4.8	4.8	4.7	4.3	3.8	3.1	2.8	3.4	3.9	3.0	3.2	3.6
College Plans:																
None or under 4 years	—	7.5	6.7	7.2	8.0	8.2	8.0	7.5	6.9	6.1	5.6	5.9	6.6	5.7	6.5	6.2
Complete 4 years	—	4.7	4.0	4.6	4.5	4.7	5.0	4.3	3.8	3.1	3.4	3.3	4.3	4.1	4.2	4.8
Region:																
Northeast	8.5	8.0	7.2	8.0	7.9	6.8	9.0	8.0	5.6	7.0	5.4	5.1	5.3	4.7	5.1	5.9
Midwest	8.7	7.0	6.5	7.9	7.9	8.5	7.8	7.3	7.0	4.4	5.3	5.3	5.7	4.7	6.0	5.3
South	5.4	4.7	3.7	3.7	3.4	4.3	3.4	3.9	4.4	3.5	2.8	2.6	4.2	4.7	4.2	4.7
West	7.6	5.9	5.0	5.8	8.3	6.5	6.3	4.8	4.2	4.5	4.6	5.9	6.2	5.2	4.4	6.4
Population Density:																
Large MSA	9.4	7.9	6.4	7.2	7.6	7.3	8.0	7.3	5.7	4.7	4.1	4.4	5.6	5.2	4.6	5.2
Other MSA	7.4	6.8	5.6	6.1	7.3	6.8	6.9	6.3	6.0	4.9	4.8	4.9	5.4	5.6	5.3	6.1
Non-MSA	5.7	4.8	4.8	5.8	4.9	5.6	4.9	4.8	4.4	4.2	4.1	4.0	4.4	3.1	4.3	4.2
Parental Education: ^a																
1.0–2.0 (Low)	6.1	4.8	4.5	5.0	4.5	5.2	4.8	5.0	4.9	4.1	3.0	3.9	4.4	4.1	3.6	3.4
2.5–3.0	6.5	6.8	5.8	6.1	6.3	6.8	6.5	6.1	5.1	4.8	4.5	4.6	4.9	3.8	4.3	4.4
3.5–4.0	6.4	6.7	5.6	6.1	6.7	6.7	6.7	6.4	5.7	4.3	4.7	4.6	4.9	4.2	5.1	6.0
4.5–5.0	7.0	6.4	5.3	6.7	7.5	5.7	6.4	5.7	5.2	4.3	4.8	4.1	5.8	6.2	5.9	6.2
5.5–6.0 (High)	6.5	6.4	6.1	7.0	7.4	7.2	7.7	6.0	4.8	5.0	3.8	4.7	6.1	6.2	5.5	7.4
Race/Ethnicity (2-year average): ^b																
White	—	—	6.3	6.3	6.8	7.0	7.2	6.9	6.2	5.5	5.0	4.9	5.4	5.8	5.7	6.1
African American	—	—	1.3	1.3	1.2	1.1	1.0	0.9	0.9	0.7	0.7	1.0	0.8	0.6	0.7	0.6
Hispanic	—	—	6.1	5.0	4.9	5.2	4.5	5.2	5.0	4.1	3.9	3.9	4.0	3.1	2.3	2.7

Table continued on next page.

TABLE 30 (cont.)
LSD: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	5.2	5.6	6.8	6.9	8.4	8.8	8.4	7.6	8.1	6.6	6.6	3.5	1.9	2.2	1.8	1.7
Gender:																
Male	6.8	6.7	8.4	8.4	10.7	10.9	10.3	9.3	10.0	7.6	7.9	4.4	2.5	3.1	2.7	2.1
Female	3.4	4.4	5.1	5.3	5.8	6.5	6.2	5.7	6.1	5.3	5.0	2.3	1.2	1.2	0.9	1.1
College Plans:																
None or under 4 years	6.4	7.6	7.5	7.7	11.2	11.4	10.3	10.9	9.4	8.7	8.2	5.7	3.0	3.2	3.4	2.6
Complete 4 years	4.7	4.8	6.4	6.3	7.3	7.7	7.4	6.3	7.3	5.6	5.7	2.7	1.4	1.8	1.3	1.4
Region:																
Northeast	6.1	6.6	8.6	8.2	8.8	11.9	11.8	8.2	7.8	7.1	6.7	5.4	2.4	2.3	1.7	3.0
Midwest	5.9	5.5	6.3	7.3	8.3	7.7	7.0	7.6	9.1	5.9	8.6	3.7	1.8	1.8	2.1	1.6
South	3.4	4.4	5.5	6.3	8.1	7.9	8.1	7.4	7.7	6.0	4.7	3.1	1.9	2.6	1.8	1.4
West	6.5	7.0	8.5	6.2	8.5	8.8	6.9	7.1	7.7	7.9	6.6	2.1	1.4	2.3	1.6	1.1
Population Density:																
Large MSA	4.3	5.7	6.7	7.3	9.7	9.0	7.7	7.2	6.8	6.7	7.7	3.3	1.5	1.9	1.8	2.2
Other MSA	7.0	5.8	7.6	7.9	8.7	10.0	9.3	8.4	9.2	6.9	6.4	4.0	2.2	2.6	1.9	1.5
Non-MSA	3.0	5.1	5.6	4.6	6.5	6.5	7.3	6.1	7.4	5.9	5.6	2.7	1.7	1.9	1.7	1.2
Parental Education: ^a																
1.0–2.0 (Low)	4.3	3.3	4.6	4.4	6.6	6.7	6.8	6.8	8.6	5.6	5.4	2.4	1.5	3.2	1.7	1.5
2.5–3.0	4.4	5.2	5.6	6.5	8.1	8.2	7.3	7.8	7.6	6.4	6.7	3.4	1.9	2.4	1.9	2.0
3.5–4.0	5.5	5.7	7.0	7.4	8.6	9.3	8.5	8.2	9.0	6.7	6.3	4.0	1.7	1.8	1.5	1.9
4.5–5.0	5.3	5.8	8.3	6.9	8.6	8.7	8.6	6.5	8.0	6.0	6.7	3.1	1.7	2.0	1.7	1.3
5.5–6.0 (High)	7.1	7.0	8.2	7.9	8.3	9.2	9.5	7.3	6.4	7.0	5.6	2.8	1.7	2.2	1.7	1.1
Race/Ethnicity (2-year average): ^b																
White	6.3	6.4	7.4	8.0	8.6	9.7	10.1	9.5	9.1	8.3	7.5	5.8	3.0	2.2	2.2	1.8
African American	0.6	0.6	0.6	0.9	1.0	1.3	1.6	1.1	0.8	1.3	1.3	0.8	0.8	0.8	0.7	0.9
Hispanic	3.6	4.1	5.1	5.4	6.4	7.4	6.3	5.9	7.0	7.6	5.8	3.8	1.8	1.7	1.7	1.4

Table continued on next page.

TABLE 30 (cont.)
LSD: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014 ^c	2015 ^c	2016 ^c	2017 ^c	2018 ^c	2019 ^c	2020 ^c	2021 ^c	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	2.1	2.7	1.9	2.6	2.7	2.4	2.2	2.5	2.9	3.0	3.3	3.2	3.6	3.9	2.5	-1.0* ss
Gender:																
Male	3.0	3.7	2.4	3.6	3.6	2.9	2.9	3.4	3.5	4.1	4.4	4.4	4.9	§	2.9	-2.0 ss
Female	1.2	1.6	1.3	1.4	1.7	1.6	1.3	1.5	2.1	1.9	2.1	1.6	2.1	§	1.9	-0.2
College Plans:																
None or under 4 years	4.1	4.8	2.5	4.2	3.7	3.1	3.4	3.7	4.4	3.6	4.9	5.0	4.7	§	3.6	-1.1
Complete 4 years	1.6	2.1	1.6	2.1	2.4	2.2	1.7	2.2	2.4	2.8	2.7	2.5	3.1	§	2.1	-1.0 s
Region:																
Northeast	2.1	3.0	2.6	3.7	3.0	2.9	2.7	2.6	2.3	3.2	2.9	3.7	3.3	§	2.5	-0.8
Midwest	2.7	2.5	2.4	2.8	3.0	1.9	2.3	2.3	2.4	3.5	2.8	3.5	3.1	§	2.3	-0.8
South	1.7	3.0	1.2	2.0	2.1	2.1	1.8	3.1	3.1	2.3	2.8	2.7	3.6	§	1.8	-1.7 sss
West	2.0	2.2	1.5	2.3	2.9	2.8	2.2	1.6	3.5	3.6	4.8	3.1	4.5	§	3.9	-0.5
Population Density:																
Large MSA	2.4	2.7	2.4	2.4	2.4	2.5	2.2	2.6	3.3	3.1	3.9	2.8	3.4	§	3.0	-0.4
Other MSA	2.0	2.8	1.7	2.9	3.0	2.7	2.4	2.6	3.2	3.4	3.4	3.6	3.7	§	2.4	-1.3
Non-MSA	1.8	2.6	1.4	2.0	2.3	1.4	1.5	2.1	1.5	1.9	2.1	2.9	3.5	§	2.2	-1.4
Parental Education: ^a																
1.0–2.0 (Low)	2.3	2.3	1.8	1.9	2.4	1.7	2.0	2.0	3.8	2.4	2.9	3.0	3.3	§	2.7	-0.7
2.5–3.0	1.9	2.4	1.6	2.3	2.3	2.6	2.8	2.5	2.7	2.8	3.4	3.1	3.8	§	3.3	-0.4
3.5–4.0	2.1	2.4	1.6	2.7	2.8	2.4	1.9	2.7	2.9	3.1	3.5	3.1	3.8	§	2.5	-1.3
4.5–5.0	2.0	2.8	1.9	2.3	2.7	2.4	1.9	2.1	2.5	3.0	3.1	2.5	3.3	§	1.8	-1.5 s
5.5–6.0 (High)	2.3	3.2	2.1	3.2	2.4	2.2	2.1	2.8	2.9	3.3	3.2	3.4	2.9	§	1.9	-1.0
Race/Ethnicity (2-year average): ^b																
White	2.1	2.8	2.6	2.7	3.3	3.1	2.5	2.4	2.6	2.9	3.5	3.6	3.4	§	2.3	-1.0 s
African American	0.8	0.8	0.7	0.8	1.0	0.9	0.8	1.1	1.3	1.1	1.5	1.5	1.6	§	2.5	+0.9
Hispanic	1.8	1.8	1.3	0.9	1.0	1.3	1.7	2.0	3.1	3.7	2.9	2.6	3.2	§	4.3	+1.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cIn 2014 questions on use of LSD were dropped from one of six forms. Beginning in 2014 data are based on five of six forms; N is five sixths of N indicated.

TABLE 31
HALLUCINOGENS OTHER THAN LSD: Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^a	2002	2003	2004	2005
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	0.7	1.1	1.0	1.3	1.7	2.0	1.8	1.6	1.5	1.4 [‡]	2.4	2.1	2.1	1.9	2.0
Gender:															
Male	0.8	1.1	1.2	1.6	1.9	2.1	2.1	1.8	1.8	1.5 [‡]	2.8	2.4	2.4	1.9	2.2
Female	0.6	1.0	0.9	0.9	1.4	1.7	1.4	1.4	1.1	1.3 [‡]	2.0	1.7	1.8	1.8	1.8
College Plans:															
None or under 4 years	1.7	3.6	2.9	3.6	4.8	5.2	4.7	5.1	5.3	3.9 [‡]	7.4	6.5	7.5	6.5	6.1
Complete 4 years	0.5	0.7	0.7	1.0	1.3	1.4	1.4	1.2	1.0	1.2 [‡]	1.8	1.6	1.5	1.4	1.6
Region:															
Northeast	0.4	0.6	0.7	1.4	1.8	2.1	1.7	1.2	1.4	1.3 [‡]	2.1	1.3	1.9	1.6	1.5
Midwest	0.6	1.3	0.6	1.1	1.6	1.8	1.8	1.9	1.7	1.8 [‡]	1.8	2.1	2.2	1.7	1.9
South	0.7	1.0	1.0	1.0	1.5	1.7	1.4	1.8	1.6	1.3 [‡]	2.9	2.2	2.2	1.9	2.4
West	1.4	1.3	1.9	1.9	1.9	2.6	2.7	1.3	1.1	1.4 [‡]	2.4	2.6	2.3	2.2	2.0
Population Density:															
Large MSA	0.7	0.8	0.7	1.6	1.6	1.8	1.4	1.3	1.1	1.2 [‡]	1.9	1.6	1.6	1.4	2.0
Other MSA	0.7	1.3	1.3	1.4	1.8	2.2	1.9	1.6	1.5	1.7 [‡]	2.6	1.9	2.1	2.1	2.0
Non-MSA	0.7	1.0	0.9	0.8	1.5	1.7	1.9	2.0	1.9	1.3 [‡]	2.6	3.0	3.0	2.0	2.2
Parental Education: ^b															
1.0–2.0 (Low)	1.5	1.8	1.4	1.6	2.4	2.2	2.6	2.7	2.6	2.5 [‡]	5.0	3.4	3.5	3.9	2.7
2.5–3.0	0.8	0.7	1.1	1.1	1.7	2.1	2.0	1.3	1.9	1.5 [‡]	2.3	2.6	2.8	2.0	2.9
3.5–4.0	0.4	1.2	1.2	1.2	1.8	2.0	1.6	1.8	1.3	1.3 [‡]	2.6	1.9	2.2	1.7	2.6
4.5–5.0	0.8	0.9	0.7	1.5	1.6	2.0	1.6	1.5	0.9	1.1 [‡]	1.6	1.7	1.5	1.6	1.6
5.5–6.0 (High)	0.8	1.3	0.9	1.5	1.7	1.4	1.9	1.9	1.5	2.0 [‡]	1.9	1.5	1.6	1.6	1.0
Race/Ethnicity (2-year average): ^c															
White	—	0.9	1.0	1.2	1.7	2.1	2.1	1.8	1.6	1.6	1.4 [‡]	1.5 ^d	2.3	2.2	2.1
African American	—	0.4	0.5	0.5	0.4	0.4	0.3	0.3	0.3	0.3 [‡]	0.4 ^d	0.6	0.6	0.8	0.9
Hispanic	—	1.5	1.5	1.5	1.8	2.1	2.0	2.2	2.2	1.8 [‡]	1.8 ^d	2.2	2.2	2.2	2.1

Table continued on next page.

TABLE 31 (cont.)
HALLUCINOGENS OTHER THAN LSD: Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
<i>Approximate weighted N =</i>																	
Total	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	-0.1*
Gender:																	
Male	1.8	1.7	1.6	1.7	2.1	2.1	1.3	1.2	1.2	1.0	0.8	0.6	1.1	0.8	§	0.7	-0.1
Female	1.7	1.4	1.5	1.4	1.6	1.4	1.2	1.1	0.7	0.7	0.7	0.7	0.7	0.9	§	0.6	-0.3
College Plans:																	
None or under 4 years	5.5	6.0	6.9	5.9	6.0	5.8	3.8	3.7	3.2	3.1	2.5	2.0	2.3	2.1	§	1.5	-0.6
Complete 4 years	1.4	1.2	1.1	1.2	1.5	1.5	1.1	1.0	0.8	0.6	0.6	0.5	0.7	0.7	§	0.6	-0.2
Region:																	
Northeast	1.2	0.8	1.0	1.3	1.4	0.9	1.2	0.5	0.5	0.6	0.4	0.4	0.5	0.2	§	0.4	+0.2
Midwest	1.7	1.7	1.4	1.4	2.0	1.4	0.7	1.1	0.8	0.8	0.6	0.3	0.8	0.9	§	0.6	-0.4
South	1.8	1.4	1.7	1.4	1.4	1.5	1.6	1.3	0.9	0.9	0.9	0.8	0.9	0.8	§	0.7	0.0
West	2.2	2.3	1.8	2.1	2.6	3.4	1.4	1.5	1.5	0.9	1.1	1.1	1.3	1.5	§	1.3	-0.2
Population Density:																	
Large MSA	1.2	1.3	1.4	1.2	1.6	1.7	1.4	1.0	0.8	0.7	0.7	0.5	1.1	0.5	§	0.5	0.0
Other MSA	2.0	1.6	1.7	1.6	2.0	2.2	1.4	1.2	1.0	1.0	0.8	0.8	0.8	1.0	§	0.8	-0.2
Non-MSA	2.1	2.0	1.3	1.8	1.7	1.1	0.9	1.2	1.1	0.8	1.0	0.7	0.8	1.2	§	1.1	-0.2
Parental Education: ^b																	
1.0–2.0 (Low)	2.3	2.5	3.1	3.3	2.7	3.1	1.6	1.8	1.6	1.1	1.3	1.1	1.3	1.4	§	1.8	+0.4
2.5–3.0	2.2	1.5	1.7	1.3	2.4	2.3	1.5	1.4	1.1	0.8	1.0	0.8	0.9	1.6	§	0.7	-0.9 s
3.5–4.0	1.8	1.9	1.7	1.6	2.0	1.9	1.7	1.2	1.0	0.8	1.1	0.8	1.2	0.8	§	0.9	+0.1
4.5–5.0	1.7	1.2	1.2	1.3	1.7	1.4	0.8	0.9	0.6	0.6	0.4	0.4	0.8	0.6	§	0.7	+0.1
5.5–6.0 (High)	1.2	0.9	1.3	1.3	1.0	1.2	1.1	1.0	1.0	0.7	0.6	0.6	0.5	0.8	§	0.5	-0.3
Race/Ethnicity (2-year average): ^c																	
White	2.2	1.8	1.7	1.6	1.8	1.8	1.3	1.1	1.0	0.8	0.7	0.7	0.7	0.8	§	1.0	+0.2
African American	0.9	0.6	0.6	0.6	0.6	0.5	0.7	0.7	0.4	0.5	0.7	0.6	0.3	0.3	§	0.3	-0.1
Hispanic	1.4	1.6	1.9	1.8	2.1	2.6	2.3	1.6	1.3	1.0	0.9	0.8	0.8	1.1	§	0.8	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. The 2001 data are based on the changed forms only; N is one half of N indicated. In 2002 have been the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and treated in a parallel manner.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 32
HALLUCINOGENS OTHER THAN LSD: Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^a	2002	2003	2004	2005	
Approximate weighted N =	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
Total	1.3	1.4	1.9	2.4	2.8	3.3	3.3	3.4	3.2	3.1‡	4.4	4.0	3.6	3.7	3.5	
Gender:																
Male	1.5	1.6	2.5	3.0	3.4	3.8	4.1	3.9	4.1	3.8‡	5.7	4.6	4.4	4.3	4.2	
Female	1.1	1.1	1.2	1.7	2.1	2.7	2.5	2.8	2.3	2.4‡	3.1	3.4	2.8	3.2	2.9	
College Plans:																
None or under 4 years	2.5	2.7	3.7	4.8	5.3	6.2	4.8	7.3	6.7	6.1‡	10.9	8.8	7.3	8.2	8.5	
Complete 4 years	1.1	1.1	1.5	1.9	2.3	2.7	3.0	2.6	2.6	2.6‡	3.3	3.3	2.9	3.1	2.8	
Region:																
Northeast	1.4	0.7	2.3	3.2	3.0	3.9	3.2	4.7	4.5	3.4‡	4.2	3.6	3.4	4.3	4.4	
Midwest	1.0	1.2	1.7	1.9	2.2	3.2	3.4	3.2	2.8	3.3‡	4.0	4.3	2.9	3.3	3.6	
South	1.3	1.2	1.5	2.1	2.7	3.1	3.1	3.2	2.7	2.9‡	3.6	3.4	3.5	3.3	3.0	
West	1.6	2.6	2.5	3.0	3.5	3.0	3.6	2.5	3.3	3.0‡	6.7	5.1	4.5	4.2	3.3	
Population Density:																
Large MSA	1.4	1.5	1.9	2.4	2.7	3.4	3.3	3.1	3.1	3.9‡	3.4	4.0	2.8	2.7	2.9	
Other MSA	1.4	1.4	1.9	2.5	3.0	3.5	3.2	3.4	3.5	2.9‡	4.8	4.0	3.9	4.4	3.9	
Non-MSA	1.0	1.2	2.0	2.3	2.5	2.7	3.5	3.5	2.7	2.8‡	4.5	3.9	3.8	3.6	3.6	
Parental Education: ^b																
1.0–2.0 (Low)	1.5	1.5	2.0	2.6	3.3	3.4	2.4	2.9	3.4	2.5‡	4.8	4.4	5.2	2.8	3.1	
2.5–3.0	1.2	1.0	1.4	1.8	2.7	3.4	2.7	4.2	3.0	2.6‡	4.4	4.3	4.2	4.5	4.1	
3.5–4.0	1.3	1.7	2.3	2.3	3.0	3.2	3.6	3.0	2.8	3.2‡	4.3	4.0	3.3	3.6	3.6	
4.5–5.0	1.2	1.5	1.9	2.6	2.5	3.3	3.7	3.0	3.7	3.7‡	4.4	3.5	3.2	3.5	3.5	
5.5–6.0 (High)	1.8	1.1	2.1	2.8	2.5	3.5	3.6	3.7	3.7	3.7‡	3.9	4.6	3.0	3.5	3.3	
Race/Ethnicity (2-year average): ^c																
White	—	1.5	1.8	2.3	2.8	3.4	3.9	4.0	4.0	3.8	3.4‡	4.0 ^d	4.7	4.4	4.2	
African American	—	0.1	0.4	0.7	0.7	0.4	0.3	0.4	0.5	0.6	1.1‡	1.3 ^d	0.9	0.7	1.1	
Hispanic	—	1.3	1.5	1.9	2.0	2.1	2.4	2.8	2.6	2.0	1.6‡	2.5 ^d	2.9	3.2	3.5	

Table continued on next page.

TABLE 32 (cont.)
HALLUCINOGENS OTHER THAN LSD: Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	3.7	3.8	3.3	3.5	3.5	3.5	3.0	2.7	2.6	1.9	2.0	1.8	1.7	2.1	2.2	1.5	-0.6* s
Gender:																	
Male	4.1	4.7	4.0	4.4	4.4	4.1	4.0	3.5	2.5	2.2	2.3	2.1	1.9	2.9	§	1.5	-1.4 sss
Female	3.1	3.0	2.6	2.6	2.5	2.9	2.1	2.0	2.7	1.6	1.5	1.4	1.4	1.3	§	1.4	+0.1
College Plans:																	
None or under 4 years	9.0	8.9	9.0	9.0	10.0	9.4	8.7	4.8	6.3	4.9	4.6	4.9	3.6	3.6	§	3.2	-0.4
Complete 4 years	3.0	3.2	2.6	2.9	2.8	2.8	2.4	2.5	2.1	1.6	1.7	1.4	1.4	1.7	§	1.1	-0.6 s
Region:																	
Northeast	3.7	4.1	2.9	2.7	3.3	3.3	2.8	1.8	2.9	2.1	1.4	1.0	1.1	1.2	§	1.6	+0.4
Midwest	3.7	4.0	3.3	3.8	2.9	2.9	2.4	2.3	2.4	1.3	1.7	1.6	1.5	1.8	§	1.6	-0.2
South	3.0	3.1	3.5	3.3	3.5	3.4	3.0	2.8	2.2	2.0	1.7	1.2	2.0	2.0	§	1.4	-0.5
West	4.7	4.4	3.6	4.3	4.4	4.6	3.9	3.7	2.8	2.4	3.0	3.4	2.1	3.3	§	1.6	-1.7 s
Population Density:																	
Large MSA	3.8	3.4	3.3	3.4	3.2	3.5	2.8	2.0	2.2	1.7	2.0	1.8	1.5	1.8	§	1.3	-0.5
Other MSA	3.7	4.1	3.6	3.6	3.7	3.8	3.2	3.1	2.8	2.0	2.1	1.6	1.9	2.2	§	1.6	-0.6
Non-MSA	3.3	3.8	2.9	3.4	3.6	3.0	2.9	2.7	2.5	2.1	1.8	2.1	1.6	2.4	§	1.9	-0.5
Parental Education: ^b																	
1.0–2.0 (Low)	4.7	4.5	4.7	5.0	4.7	4.9	4.1	3.6	2.5	2.8	2.0	1.9	2.2	1.6	§	2.8	+1.2
2.5–3.0	3.8	3.9	3.3	3.2	5.0	4.1	3.5	3.5	2.6	2.7	2.6	2.6	1.9	2.8	§	2.2	-0.6
3.5–4.0	3.4	4.1	3.5	3.7	3.4	3.3	2.8	3.0	3.1	2.3	2.2	2.0	1.7	2.7	§	1.7	-1.0
4.5–5.0	3.7	3.8	2.9	3.2	2.8	3.1	2.9	2.4	2.2	1.6	1.5	1.4	1.4	1.5	§	1.1	-0.4
5.5–6.0 (High)	3.8	3.3	2.6	3.8	2.6	3.2	2.4	2.0	2.1	1.2	1.6	1.3	1.9	1.7	§	1.0	-0.7
Race/Ethnicity (2-year average): ^c																	
White	4.2	4.4	4.2	3.9	3.8	3.8	3.5	3.0	2.7	2.4	2.2	2.2	1.9	2.1	§	1.7	-0.4
African American	1.2	0.8	0.7	1.1	1.2	1.0	1.1	1.1	0.9	1.4	1.2	0.6	0.4	0.6	§	1.8	+1.3 s
Hispanic	3.0	2.3	2.9	3.6	3.7	3.7	3.5	2.9	2.4	2.0	1.9	1.8	1.9	2.1	§	1.3	-0.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. The 2001 data are based on the changed forms only; N is one half of N indicated. In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 33
HALLUCINOGENS OTHER THAN LSD: Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
Approximate weighted N =	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	9.4	7.0	6.9	7.3	6.8	6.2	5.6	4.7	4.1	3.8	3.6	3.0	3.2	2.1	2.2	2.1
Gender:																
Male	12.1	8.8	8.9	8.8	8.0	8.0	6.9	5.7	4.9	4.8	4.6	3.4	3.7	2.7	3.1	3.0
Female	7.5	5.0	4.9	5.5	5.3	4.2	4.0	3.6	3.1	2.7	2.5	2.5	2.6	1.4	1.1	1.1
College Plans:																
None or under 4 years	—	8.3	8.6	8.1	7.6	6.7	6.0	5.1	4.8	4.5	4.1	3.3	3.5	2.5	2.7	2.1
Complete 4 years	—	5.2	4.9	5.7	5.3	5.1	5.0	4.1	3.1	3.0	2.9	2.5	2.8	1.8	1.9	1.9
Region:																
Northeast	12.0	7.8	8.2	10.3	10.2	10.0	9.0	8.1	6.1	8.4	7.1	5.1	4.8	2.6	2.5	2.9
Midwest	11.3	7.9	7.9	7.6	6.8	6.7	5.7	4.8	4.4	2.9	3.1	2.7	3.5	1.6	2.3	1.7
South	7.1	5.7	5.4	4.8	4.1	3.1	2.0	1.8	2.4	1.5	1.5	1.4	1.7	2.0	1.9	1.7
West	7.7	6.7	6.3	7.2	6.6	5.9	6.9	4.9	4.1	4.5	3.6	3.3	3.3	2.5	2.3	2.6
Population Density:																
Large MSA	11.1	7.8	7.5	9.3	8.8	8.5	8.1	7.3	6.6	6.9	6.1	5.1	4.8	3.3	2.6	2.3
Other MSA	10.7	7.3	7.2	6.9	7.2	6.5	5.0	3.7	4.0	3.1	3.0	2.4	2.5	1.7	2.0	2.4
Non-MSA	6.8	6.1	6.1	6.1	4.7	4.1	4.4	4.0	2.4	2.4	2.4	2.1	2.9	1.6	2.1	1.5
Parental Education: ^a																
1.0–2.0 (Low)	7.5	5.1	5.4	5.7	4.7	5.8	3.9	3.7	3.6	2.8	2.7	2.3	3.4	2.7	2.0	1.5
2.5–3.0	8.7	7.6	6.9	7.2	6.5	5.9	5.1	4.7	3.6	3.8	3.5	3.1	2.9	1.5	1.8	1.2
3.5–4.0	9.1	7.5	7.4	7.3	6.6	5.8	5.8	4.8	4.5	3.6	4.1	3.2	3.0	1.8	1.8	2.2
4.5–5.0	9.7	8.4	7.0	7.8	7.5	6.4	6.0	4.7	3.9	3.8	3.6	2.7	3.3	2.2	2.6	2.8
5.5–6.0 (High)	7.4	7.2	7.8	7.9	9.2	7.2	7.0	6.7	4.6	5.3	2.6	2.8	3.5	3.1	3.3	3.6
Race/Ethnicity (2-year average): ^b																
White	—	—	7.6	7.6	7.5	7.0	6.3	5.6	4.7	4.3	4.0	3.5	3.4	3.0	2.4	2.4
African American	—	—	1.6	1.6	1.3	1.4	1.3	1.2	1.6	1.3	0.8	0.9	0.9	0.6	0.5	0.4
Hispanic	—	—	5.1	5.2	4.8	4.4	4.5	4.8	4.2	3.4	3.7	3.3	2.1	1.7	1.6	1.5

Table continued on next page.

TABLE 33 (cont.)
HALLUCINOGENS OTHER THAN LSD: Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^c	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	2.0	1.7	2.2	3.1	3.8	4.4	4.6	4.6	4.3	4.4‡	5.9	5.4	5.4	5.6	5.0	4.6
Gender:																
Male	2.7	2.3	3.0	4.3	5.3	5.7	5.9	6.0	5.4	5.8‡	7.2	7.2	7.2	7.7	6.7	5.7
Female	1.3	1.2	1.3	1.9	2.1	2.8	3.2	3.0	3.1	2.9‡	4.2	3.5	3.4	3.4	3.2	3.3
College Plans:																
None or under 4 years	2.4	2.2	2.5	3.2	4.4	5.1	4.7	6.1	4.4	5.6‡	7.3	7.9	7.3	6.9	6.8	6.5
Complete 4 years	1.9	1.5	2.0	2.9	3.4	4.0	4.4	3.9	4.0	3.8‡	5.0	4.6	4.5	4.9	4.4	4.1
Region:																
Northeast	3.5	2.5	2.6	5.1	5.3	6.1	6.6	6.3	5.5	5.0‡	6.9	7.5	7.3	6.7	4.9	6.4
Midwest	1.9	1.8	2.0	3.1	3.2	3.5	2.8	4.1	3.7	3.6‡	6.9	5.0	4.7	4.6	6.3	4.7
South	1.0	1.3	1.8	1.9	3.1	3.6	4.0	3.5	3.3	3.3‡	3.1	4.5	4.3	4.9	4.7	3.8
West	2.6	1.8	2.9	3.6	4.5	5.4	6.1	5.5	5.7	6.9‡	8.0	5.6	6.0	6.9	4.0	4.2
Population Density:																
Large MSA	1.8	2.1	2.5	3.6	4.9	5.2	4.0	4.4	4.5	5.2‡	7.6	5.9	3.9	5.2	5.1	5.5
Other MSA	2.6	1.7	2.2	3.6	3.7	4.7	5.4	5.1	4.2	4.3‡	5.4	5.8	6.5	6.6	5.4	4.2
Non-MSA	1.2	1.4	2.0	1.8	2.7	3.1	3.6	3.7	4.2	3.8‡	4.8	4.3	5.1	4.1	4.0	3.9
Parental Education: ^a																
1.0–2.0 (Low)	1.9	1.3	1.7	2.2	2.7	3.5	2.7	3.7	2.2	3.8‡	3.8	4.3	4.7	3.3	2.9	4.9
2.5–3.0	1.6	1.6	1.5	2.4	3.4	3.0	3.5	3.9	3.3	3.6‡	5.4	5.4	4.3	4.9	5.1	4.5
3.5–4.0	2.0	1.8	2.4	2.9	3.6	4.0	4.8	4.6	5.1	4.2‡	6.1	5.6	5.9	6.0	4.9	4.9
4.5–5.0	2.4	1.7	2.7	3.7	4.2	5.2	5.3	5.0	4.4	4.6‡	5.5	5.8	6.0	6.3	5.3	4.3
5.5–6.0 (High)	2.4	2.1	3.0	4.4	4.1	5.9	5.9	5.4	4.7	5.8‡	6.6	5.1	4.8	5.6	5.3	4.5
Race/Ethnicity (2-year average): ^b																
White	2.4	2.2	2.2	3.0	3.8	4.4	5.2	5.6	5.2	4.8	4.7‡	5.4 ^d	6.3	6.4	6.1	5.5
African American	0.3	0.3	0.5	0.7	0.8	0.8	0.7	0.6	0.6	1.0	0.9‡	0.9 ^d	0.9	0.9	1.2	1.4
Hispanic	1.7	1.4	1.4	1.6	2.6	3.5	3.1	2.7	3.0	4.6	4.8‡	4.1 ^d	4.1	3.7	3.6	3.5

Table continued on next page.

TABLE 33 (cont.)
HALLUCINOGENS OTHER THAN LSD: Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014 ^e	2015 ^e	2016 ^e	2017 ^e	2018 ^e	2019 ^e	2020 ^e	2021 ^e	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	4.8	5.0	4.2	4.8	4.3	4.0	3.7	3.0	2.9	2.7	2.9	2.7	2.7	2.8	2.9	+0.1*
Gender:																
Male	6.8	6.7	5.7	6.9	5.9	5.3	5.0	4.1	3.6	3.5	4.1	3.6	3.7	§	3.8	+0.1
Female	2.9	3.4	2.6	2.5	2.7	2.6	2.2	2.1	1.9	1.8	1.8	1.4	1.5	§	1.6	0.0
College Plans:																
None or under 4 years	7.2	6.3	5.0	7.1	6.7	4.7	4.9	4.1	4.9	3.2	4.1	4.3	4.4	§	4.3	0.0
Complete 4 years	4.2	4.7	3.9	4.2	3.8	3.9	3.3	2.7	2.3	2.5	2.5	2.0	2.2	§	2.3	+0.2
Region:																
Northeast	6.0	7.2	3.8	5.0	4.7	4.0	3.3	3.4	2.8	3.1	2.9	3.0	2.2	§	2.9	+0.7
Midwest	4.6	3.9	5.3	4.5	4.2	4.1	4.1	3.2	2.7	2.9	2.1	2.9	2.7	§	2.5	-0.3
South	3.8	4.6	3.7	3.9	3.2	3.5	3.5	2.6	2.4	2.0	2.6	1.9	2.2	§	2.6	+0.4
West	5.7	5.4	4.0	6.2	5.8	4.7	3.7	3.4	4.1	3.6	4.3	3.6	4.4	§	3.5	-0.9
Population Density:																
Large MSA	5.5	6.0	4.5	5.0	4.1	4.8	3.7	3.1	2.8	2.5	3.4	2.3	2.4	§	3.3	+0.9
Other MSA	4.7	4.9	4.3	5.1	4.7	4.0	4.1	3.3	3.3	2.9	2.8	2.6	2.6	§	2.8	+0.2
Non-MSA	4.1	4.4	3.6	3.6	3.6	2.8	2.6	2.3	2.2	2.7	2.6	3.6	3.5	§	2.2	-1.3
Parental Education: ^a																
1.0–2.0 (Low)	3.5	3.4	3.0	3.1	4.5	2.8	3.5	2.2	3.0	2.9	3.0	2.7	2.2	§	1.8	-0.4
2.5–3.0	4.3	3.9	4.1	4.1	3.7	4.3	3.5	3.2	2.9	2.4	2.8	2.8	2.7	§	2.2	-0.4
3.5–4.0	4.9	4.8	4.4	5.5	4.3	4.1	4.2	3.3	3.3	2.5	3.2	2.5	2.8	§	2.7	-0.1
4.5–5.0	4.9	6.0	4.7	4.7	4.0	4.3	3.0	3.1	2.4	3.0	2.5	2.1	2.5	§	3.6	+1.1
5.5–6.0 (High)	5.8	6.1	4.1	4.8	5.1	3.4	4.3	2.8	3.0	2.7	3.3	2.7	3.0	§	2.7	-0.3
Race/Ethnicity (2-year average): ^b																
White	5.6	6.0	5.6	5.4	5.5	4.9	4.4	3.8	3.2	3.0	3.1	3.0	2.9	§	3.1	+0.2
African American	1.2	1.0	0.9	1.0	1.3	1.4	1.3	1.4	1.3	1.1	1.3	1.2	1.1	§	2.0	+0.9 s
Hispanic	3.4	3.0	2.8	3.0	2.8	2.6	3.0	2.7	2.7	2.8	2.3	2.1	2.2	§	1.9	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cIn 2001 the question text was changed on half of the questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. The 2001 data are based on the changed forms only; N is one half of N indicated. In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^dDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

^eIn 2014 questions on use of hallucinogens other than LSD were dropped from one of six forms. Beginning in 2014 data are based on five of six forms; N is five sixths of N indicated.

TABLE 34
MDMA (ECSTASY, MOLLY): Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months											
	1991- 1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Approximate weighted N =	—	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500
Total	—	2.3	2.3	1.8	1.7	3.1	3.5	2.9	2.1	1.7	1.7	1.4
Gender:												
Male	—	2.2	2.7	2.3	1.7	3.1	3.4	3.1	1.8	1.7	1.6	1.0
Female	—	2.3	2.0	1.3	1.7	3.0	3.7	2.6	2.2	1.8	1.7	1.7
College Plans:												
None or under 4 years	—	4.8	6.1	4.8	5.3	6.5	9.2	9.1	6.9	5.4	5.9	3.8
Complete 4 years	—	1.9	2.0	1.5	1.2	2.7	2.9	2.3	1.5	1.3	1.3	1.2
Region:												
Northeast	—	2.1	1.5	1.6	1.8	2.4	3.8	2.0	1.8	1.8	1.1	0.8
Midwest	—	1.7	1.7	1.7	1.4	3.5	2.9	2.3	2.1	1.4	1.7	1.6
South	—	2.8	2.3	2.7	1.8	3.2	3.7	3.7	2.5	1.9	2.1	2.0
West	—	2.3	3.8	0.8	1.7	2.9	3.6	3.0	1.6	1.7	1.5	0.7
Population Density:												
Large MSA	—	2.9	1.8	1.8	1.6	3.0	3.1	2.6	1.8	2.1	1.6	1.1
Other MSA	—	2.5	3.1	2.1	1.8	3.4	4.3	3.3	1.9	1.9	1.6	1.6
Non-MSA	—	1.2	1.5	1.5	1.6	2.5	2.5	2.5	2.7	1.0	1.9	1.6
Parental Education: ^a												
1.0–2.0 (Low)	—	2.9	2.2	2.1	2.5	4.2	5.8	6.0	3.6	3.6	2.5	2.8
2.5–3.0	—	2.2	2.4	1.4	2.0	3.9	4.3	3.6	3.1	2.2	3.0	1.5
3.5–4.0	—	2.2	2.9	2.6	1.5	2.8	2.4	2.9	2.1	1.7	1.7	1.2
4.5–5.0	—	2.6	2.0	1.4	1.2	3.1	3.3	2.4	1.4	1.2	1.1	1.3
5.5–6.0 (High)	—	2.4	2.7	2.5	2.3	2.0	3.0	1.2	2.0	1.7	0.9	1.2
Race/Ethnicity (2-year average): ^b												
White	—	—	2.7	2.4	1.9	2.5	3.2	2.9	2.4	1.9	1.5	1.5
African American	—	—	0.3	0.4	0.5	0.6	1.1	1.1	1.0	1.6	1.8	1.4
Hispanic	—	—	2.5	1.7	1.9	3.3	5.3	5.9	4.0	2.3	2.3	1.8

Table continued on next page.

TABLE 34 (cont.)
MDMA (ECSTASY, MOLLY): Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	1.5	1.7	1.3	2.4	1.7	1.1	1.1‡	1.5	1.4	1.0	0.9	1.1	1.1	0.8	0.6	-0.5* ss
Gender:																
Male	1.3	1.4	1.3	2.5	2.0	0.9	1.1‡	1.6	1.2	1.0	0.9	1.2	1.2	§	0.7	-0.5
Female	1.7	2.0	1.2	2.4	1.4	1.2	1.1‡	1.5	1.5	0.9	0.9	0.9	1.0	§	0.4	-0.7 sss
College Plans:																
None or under 4 years	5.4	6.9	4.9	9.9	4.7	3.0	3.1‡	4.1	3.3	2.9	3.1	3.3	3.4	§	1.3	-2.1 ss
Complete 4 years	1.1	1.3	1.0	1.9	1.5	0.9	1.0‡	1.3	1.2	0.8	0.7	0.7	0.8	§	0.5	-0.3
Region:																
Northeast	0.5	1.2	1.0	1.2	0.9	1.2	0.6‡	1.4	1.0	0.6	0.6	0.7	0.3	§	0.1	-0.2
Midwest	1.2	1.4	1.0	2.4	0.9	0.3	1.2‡	1.2	1.3	0.9	1.0	0.7	0.8	§	0.6	-0.2
South	1.8	1.8	1.1	2.5	1.5	1.4	1.3‡	1.7	1.7	1.1	0.9	1.2	1.3	§	0.5	-0.8 s
West	1.9	2.2	2.2	3.2	3.7	1.3	1.2‡	1.5	1.1	1.1	1.1	1.6	1.7	§	1.1	-0.6
Population Density:																
Large MSA	1.4	1.9	1.3	2.5	1.9	1.1	1.1‡	1.2	1.7	0.7	0.9	1.2	0.6	§	0.5	-0.2
Other MSA	1.5	1.7	1.4	2.6	1.9	1.3	1.3‡	1.8	1.4	1.1	0.9	1.1	1.4	§	0.8	-0.6 s
Non-MSA	1.8	1.3	1.0	2.0	1.2	0.6	0.8‡	1.2	0.9	1.2	0.8	0.9	1.3	§	0.5	-0.8
Parental Education: ^a																
1.0–2.0 (Low)	3.1	4.1	1.8	5.0	2.2	2.1	2.3‡	2.4	2.5	1.7	2.1	2.0	2.1	§	1.0	-1.1
2.5–3.0	1.9	1.7	1.6	3.8	2.2	1.7	1.2‡	2.1	1.5	1.1	1.5	1.4	2.3	§	0.6	-1.8 ss
3.5–4.0	1.1	2.1	1.4	2.2	2.1	1.0	1.0‡	1.3	1.2	1.1	0.8	1.1	0.8	§	0.8	0.0
4.5–5.0	1.4	1.0	1.1	1.9	1.5	0.6	1.1‡	1.0	1.0	1.1	0.5	1.0	0.8	§	0.4	-0.3
5.5–6.0 (High)	0.8	0.8	1.1	1.1	1.1	0.6	0.6‡	1.3	0.8	0.3	0.5	0.5	0.8	§	0.8	+0.1
Race/Ethnicity (2-year average): ^b																
White	1.4	1.6	1.5	1.9	1.8	1.0	0.9	1.0‡	1.3	0.9	0.7	0.7	0.9	§	0.8	-0.1
African American	1.2	0.7	0.4	0.7	1.1	1.0	0.9	1.0‡	1.3	1.3	1.2	0.5	0.2	§	0.2	0.0
Hispanic	1.8	2.1	2.1	2.6	3.1	2.4	1.8	1.5‡	1.3	1.3	1.0	1.3	1.5	§	0.7	-0.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1996; N is one half of N indicated in Table S2. Data based on one of four forms in 1997–2001; N is one third of N indicated in Table S2. Data based on two of four forms beginning in 2002; N is one half of N indicated in Table S2. In 2014 a revised question on use of ecstasy (MDMA) including "Molly" was added to one form. Beginning in 2014 data reported here are for the revised wording which includes "Molly"; N is two sixths of N indicated in 2014 and five sixths of N indicated beginning in 2015.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. Two year race

data for the 2014 are based on the 2013 and 2014 unchanged forms only. Two year race data for the 2015 are based on the 2014 revised wording data two and one half times weighted and the 2015 revised

wording data. Beginning in 2016 two year race data based only on revised wording data. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 35
MDMA (ECSTASY, MOLLY): Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months											
	1991- 1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Approximate weighted N =	—	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	16,200
Total	—	4.6	3.9	3.3	4.4	5.4	6.2	4.9	3.0	2.4	2.6	2.8
Gender:												
Male	—	4.9	4.7	3.5	4.7	5.7	7.6	4.3	2.8	2.4	2.6	3.3
Female	—	4.2	3.1	2.9	4.2	4.8	4.9	5.2	3.2	2.4	2.5	2.2
College Plans:												
None or under 4 years	—	7.7	7.5	5.4	8.5	10.7	13.6	10.1	5.7	5.4	5.6	6.9
Complete 4 years	—	4.0	3.3	2.9	3.7	4.5	5.1	4.1	2.6	2.0	2.2	2.2
Region:												
Northeast	—	4.4	3.0	3.8	7.0	6.4	8.2	4.4	2.7	2.6	2.2	2.2
Midwest	—	3.6	3.2	2.2	2.3	5.2	4.8	4.6	2.5	1.9	3.0	2.8
South	—	5.6	5.0	4.1	4.1	5.2	5.9	5.1	3.8	2.6	2.6	3.0
West	—	4.1	3.7	2.7	4.4	5.0	6.8	5.2	3.0	2.7	2.5	3.0
Population Density:												
Large MSA	—	5.6	4.0	2.5	5.2	7.0	7.3	5.1	2.4	1.8	3.0	2.9
Other MSA	—	4.6	3.6	4.1	4.7	5.3	5.5	5.2	3.1	3.1	2.5	2.7
Non-MSA	—	3.3	4.2	2.7	2.8	3.7	6.3	3.7	3.8	2.0	2.3	2.8
Parental Education: ^a												
1.0–2.0 (Low)	—	4.8	3.0	1.9	3.7	7.0	7.2	5.8	2.4	3.3	4.4	3.8
2.5–3.0	—	4.7	3.9	3.1	4.4	4.9	6.5	5.5	3.8	3.3	2.8	2.9
3.5–4.0	—	4.7	5.2	4.3	4.0	6.3	7.0	5.5	3.5	2.1	3.3	2.5
4.5–5.0	—	4.2	2.5	2.9	4.3	5.0	5.3	3.9	2.7	2.3	2.2	2.7
5.5–6.0 (High)	—	5.3	4.0	4.4	5.6	4.8	5.8	4.6	2.3	1.4	1.4	3.3
Race/Ethnicity (2-year average): ^b												
White	—	—	4.8	4.0	4.4	5.3	6.0	6.2	4.6	3.1	2.7	2.9
African American	—	—	0.8	1.2	1.3	1.5	2.2	1.8	1.5	0.9	0.6	1.0
Hispanic	—	—	3.6	2.3	2.4	4.6	5.4	4.3	3.4	3.0	4.3	3.9

Table continued on next page.

TABLE 35 (cont.)
MDMA (ECSTASY, MOLLY): Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	3.5	2.9	3.7	4.7	4.5	3.0	3.6‡	3.8	2.4	1.8	1.7	1.4	1.7	1.2	0.7	-0.9* sss
Gender:																
Male	3.7	3.4	3.7	5.4	4.9	3.3	4.4‡	3.9	2.6	2.2	2.1	1.5	2.1	§	0.8	-1.4 sss
Female	3.1	2.4	3.5	4.0	3.8	2.6	2.9‡	3.6	2.1	1.4	1.3	1.3	1.2	§	0.5	-0.7 ss
College Plans:																
None or under 4 years	7.4	7.5	8.6	9.2	10.6	8.3	7.0‡	8.7	5.9	4.8	4.5	3.4	3.7	§	1.5	-2.2 ss
Complete 4 years	3.0	2.3	3.1	4.2	3.8	2.4	3.2‡	3.2	2.0	1.5	1.4	1.1	1.2	§	0.5	-0.7 sss
Region:																
Northeast	2.9	2.0	2.0	3.7	3.2	1.9	3.3‡	3.4	1.9	1.2	1.6	0.8	0.9	§	0.4	-0.5
Midwest	3.5	2.0	2.8	3.1	2.1	2.4	3.1‡	3.1	1.9	1.6	2.1	1.6	1.1	§	0.7	-0.4
South	3.6	2.8	3.2	4.5	3.9	2.8	3.9‡	4.8	3.1	1.8	1.5	1.6	1.6	§	1.0	-0.6
West	3.8	4.5	6.8	7.8	9.2	4.8	3.8‡	3.3	2.3	2.4	1.8	1.4	2.8	§	0.5	-2.3 sss
Population Density:																
Large MSA	3.0	3.8	3.3	4.8	4.8	3.8	3.1‡	3.4	2.6	1.9	1.9	1.3	1.1	§	0.7	-0.4 s
Other MSA	3.7	2.7	4.0	5.3	5.1	2.9	4.6‡	4.4	2.1	1.7	1.6	1.5	2.0	§	0.7	-1.3 sss
Non-MSA	3.8	2.0	3.5	3.3	3.0	1.9	1.7‡	3.1	2.7	1.9	1.7	1.4	1.7	§	0.9	-0.8 s
Parental Education: ^a																
1.0–2.0 (Low)	4.3	4.8	6.0	8.5	9.5	5.6	3.5‡	2.5	3.7	2.3	2.0	1.7	1.3	§	2.1	+0.7
2.5–3.0	4.7	3.5	4.8	5.6	5.0	3.0	5.0‡	4.6	2.3	2.0	2.6	1.6	2.3	§	0.5	-1.8 sss
3.5–4.0	3.2	3.2	3.8	4.8	4.4	2.5	3.5‡	5.1	3.2	1.8	2.0	1.3	2.0	§	1.2	-0.8
4.5–5.0	3.2	2.0	2.5	3.6	2.5	2.9	3.1‡	3.8	1.9	1.8	1.1	1.3	1.2	§	0.5	-0.7
5.5–6.0 (High)	2.8	2.1	3.4	3.6	3.8	2.5	3.0‡	2.5	1.7	1.5	1.3	1.6	1.3	§	0.4	-0.9 s
Race/Ethnicity (2-year average): ^b																
White	3.5	3.4	2.9	3.7	4.0	3.2	2.9	3.2‡	3.5	2.2	1.7	1.4	1.4	§	0.9	-0.5 s
African American	1.3	1.3	1.5	1.9	2.2	1.8	1.3	1.3‡	1.9	1.5	1.3	1.0	0.8	§	0.8	0.0
Hispanic	2.7	3.8	5.6	7.3	8.7	7.3	5.1	3.9‡	3.2	2.1	2.1	2.2	2.0	§	0.7	-1.3 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1996; N is one half of N indicated in Table S2. Data based on one of four forms in 1997–2001; N is one third of N indicated in Table S2. Data based on two of four forms beginning in 2002; N is one half of N indicated in Table S2. In 2014 a revised question on use of ecstasy (MDMA) including "Molly" was added to one form. Beginning in 2014 data reported here are for the revised wording which includes "Molly"; N is two sixths of N indicated in 2014 and five sixths of N indicated beginning in 2015.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. Two year race

data for the 2014 are based on the 2013 and 2014 unchanged forms only. Two year race data for the 2015 are based on the 2014 revised wording data two and one half times weighted and the 2015 revised

wording data. Beginning in 2016 two year race data based only on revised wording data. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 36
MDMA (ECSTASY, MOLLY): Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months											
	1975– 1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Approximate weighted N =	—	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	—	4.6	4.0	3.6	5.6	8.2	9.2	7.4	4.5	4.0	3.0	4.1
Gender:												
Male	—	4.8	5.6	4.8	5.6	8.1	10.5	8.2	4.8	4.7	3.3	4.3
Female	—	4.2	2.5	2.7	5.6	8.2	8.0	6.4	4.0	3.2	2.7	4.0
College Plans:												
None or under 4 years	—	6.9	3.8	4.7	4.2	8.5	9.8	8.9	6.5	5.3	3.7	6.0
Complete 4 years	—	4.0	3.9	3.3	6.2	8.0	8.7	7.1	3.9	3.6	2.8	3.7
Region:												
Northeast	—	6.3	6.9	3.7	9.4	8.8	10.1	10.3	5.1	3.5	3.0	2.9
Midwest	—	3.7	1.3	2.7	3.3	5.7	11.8	5.0	4.7	3.0	3.0	3.4
South	—	4.6	4.3	4.0	5.7	5.9	5.7	7.9	4.2	5.1	3.4	4.9
West	—	3.9	4.1	4.0	5.0	14.4	10.3	6.8	4.2	3.8	2.5	4.6
Population Density:												
Large MSA	—	3.6	3.6	3.2	6.1	8.7	10.9	8.4	4.3	4.0	3.5	6.1
Other MSA	—	5.1	4.6	4.3	6.1	8.4	9.7	8.1	5.0	4.5	3.2	3.1
Non-MSA	—	4.5	3.4	2.7	4.2	7.4	6.4	4.6	4.0	2.8	2.1	3.1
Parental Education: ^a												
1.0–2.0 (Low)	—	5.5	3.5	4.2	6.8	7.3	3.8	4.8	4.1	4.6	3.0	3.8
2.5–3.0	—	5.0	3.1	3.2	5.1	7.7	10.3	8.0	3.8	3.7	3.9	3.6
3.5–4.0	—	4.9	3.8	3.2	5.7	6.2	8.4	7.5	5.7	4.6	2.8	3.9
4.5–5.0	—	4.0	2.9	4.3	6.2	8.3	11.2	7.3	5.0	3.8	3.2	5.1
5.5–6.0 (High)	—	4.1	8.7	3.5	4.7	10.6	8.1	7.6	3.3	3.1	2.0	3.6
Race/Ethnicity (2-year average): ^b												
White	—	—	5.2	4.7	5.1	7.6	9.6	8.5	6.4	4.7	3.9	4.0
African American	—	—	0.4	0.4	0.5	1.3	2.4	1.7	1.4	1.6	1.4	1.8
Hispanic	—	—	2.8	2.7	6.0	10.6	10.2	7.0	5.3	3.7	3.0	3.0

Table continued on next page.

TABLE 36 (cont.)
MDMA (ECSTASY, MOLLY): Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019– 2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	4.5	4.3	4.3	4.5	5.3	3.8	4.0‡	5.0	3.6	2.7	2.6	2.2	2.2	1.8	1.1	-1.1* sss
Gender:																
Male	5.0	5.5	5.5	5.3	6.3	4.1	5.0‡	6.6	4.0	3.3	3.5	2.9	2.9	§	1.3	-1.6 ss
Female	4.1	3.3	3.2	3.6	4.6	3.4	2.9‡	3.1	3.1	1.9	1.6	1.5	1.5	§	1.1	-0.4
College Plans:																
None or under 4 years	6.5	7.2	6.1	4.7	8.5	4.8	4.3‡	5.6	6.1	3.0	3.8	3.4	4.1	§	2.5	-1.6
Complete 4 years	3.9	3.7	3.9	4.4	4.7	3.4	4.0‡	4.6	3.1	2.4	2.1	1.8	1.7	§	0.7	-0.9 ss
Region:																
Northeast	4.3	2.9	3.8	4.5	5.9	3.7	4.6‡	6.0	3.9	2.1	1.7	1.6	1.3	§	0.5	-0.7
Midwest	4.4	2.8	4.5	3.5	3.7	2.4	3.9‡	2.8	2.3	2.9	1.9	2.2	1.5	§	1.1	-0.5
South	4.6	4.6	3.8	3.6	4.4	3.2	3.7‡	5.4	2.7	2.4	2.6	1.8	2.6	§	0.6	-2.0 sss
West	4.7	6.6	5.2	6.9	7.9	5.9	4.0‡	5.6	5.9	3.3	3.8	3.1	3.0	§	2.4	-0.6
Population Density:																
Large MSA	5.8	4.7	4.1	5.2	5.8	5.1	5.1‡	4.2	3.5	2.3	2.6	2.1	1.7	§	1.2	-0.4
Other MSA	4.0	4.8	4.7	4.9	5.7	3.7	3.9‡	6.8	3.6	2.9	2.9	2.2	2.4	§	1.3	-1.1
Non-MSA	3.7	2.8	3.5	2.2	3.7	1.8	2.6‡	1.8	3.9	2.6	1.9	2.2	2.5	§	0.5	-2.0 sss
Parental Education: ^a																
1.0–2.0 (Low)	4.7	4.0	6.2	3.8	6.3	3.7	5.3‡	6.4	3.4	2.9	3.3	1.7	1.8	§	0.2	-1.6 s
2.5–3.0	4.9	4.0	2.5	4.9	5.5	3.5	3.8‡	4.1	3.9	2.7	3.1	2.7	2.9	§	1.3	-1.6 s
3.5–4.0	4.4	5.3	4.6	4.7	5.0	4.1	4.1‡	4.2	4.3	2.4	2.5	2.0	1.8	§	2.3	+0.5
4.5–5.0	4.1	3.6	4.9	4.2	5.7	3.9	3.2‡	4.9	3.1	2.5	2.1	2.0	1.9	§	0.7	-1.3 s
5.5–6.0 (High)	4.6	4.1	4.0	4.3	4.4	3.6	5.9‡	6.7	3.7	2.9	2.4	2.5	2.5	§	0.9	-1.6
Race/Ethnicity (2-year average): ^b																
White	4.9	4.8	4.6	4.5	5.1	4.9	4.0	3.9‡	4.0	3.3	2.7	2.6	2.4	§	1.7	-0.7
African American	2.1	2.2	2.4	2.6	2.4	1.5	1.1	1.7‡	3.3	2.2	1.6	0.9	1.1	§	1.1	0.0
Hispanic	3.3	3.2	3.9	4.6	4.4	4.5	4.5	3.9‡	3.8	2.8	2.2	2.1	1.8	§	1.1	-0.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms in 1996–2001; N is one sixth of N indicated in Table S3. Data based on two of six forms beginning in 2002; N is two sixths of N indicated in Table S3. Beginning in 2014 a revised question on use of ecstasy (MDMA) including "Molly" was added to one form. Beginning in 2014 data reported here are for the revised wording which includes "Molly". N is one sixth of N indicated in 2014 and three sixths of N indicated beginning in 2015.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. Two year race

data for the 2014 are based on the 2013 and 2014 unchanged forms only. Two year race data for the 2015 are based on the 2014 revised wording data triple weighted and the 2015 revised

wording data. Beginning in 2016 two year race data based only on revised wording data. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size

of its usual sample because of pandemic-curtailed data collection).

TABLE 37
SALVIA: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														2019– 2021
	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change	
<i>Approximate weighted N =</i>	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700		
Total	—	1.7	1.6	1.4	1.2	0.6	0.7	0.9	0.4	0.6	0.8	0.5	0.5	-0.2*	
Gender:															
Male	—	2.6	2.0	1.4	1.6	0.6	0.7	1.2	0.5	1.0	1.0	§	0.4	-0.6	
Female	—	1.0	1.1	1.1	0.9	0.6	0.6	0.7	0.2	0.3	0.5	§	0.7	+0.2	
College Plans:															
None or under 4 years	—	4.4	5.4	5.2	4.2	2.7	1.7	5.5	1.4	2.0	2.1	§	1.9	-0.2	
Complete 4 years	—	1.4	1.3	1.1	1.0	0.4	0.6	0.5	0.2	0.4	0.6	§	0.3	-0.2	
Region:															
Northeast	—	1.9	1.8	0.5	0.8	0.8	0.5	0.7	0.1	0.5	0.2	§	0.0	-0.2	
Midwest	—	1.6	1.1	0.4	0.3	0.2	0.5	0.5	0.2	0.2	0.5	§	0.6	+0.1	
South	—	1.4	1.5	2.4	1.1	0.8	0.8	1.4	0.5	0.9	1.4	§	0.9	-0.4	
West	—	2.4	2.1	1.2	2.6	0.5	0.7	0.7	0.5	0.8	0.5	§	0.3	-0.2	
Population Density:															
Large MSA	—	2.0	1.3	1.9	1.4	0.5	0.7	0.9	0.3	1.0	1.0	§	0.6	-0.4	
Other MSA	—	1.6	2.1	1.3	1.1	0.8	0.7	1.0	0.5	0.5	0.9	§	0.6	-0.3	
Non-MSA	—	1.9	0.9	0.6	1.2	0.3	0.4	0.7	0.2	0.3	0.1	§	0.4	+0.3	
Parental Education: ^a															
1.0–2.0 (Low)	—	2.3	4.7	1.4	2.3	1.0	1.3	2.2	1.3	0.9	1.2	§	0.4	-0.8	
2.5–3.0	—	1.7	2.5	1.3	1.1	0.4	0.5	1.1	0.2	0.1	1.5	§	0.2	-1.3	
3.5–4.0	—	0.9	1.7	1.0	1.4	0.3	0.4	1.9	0.0	0.4	0.4	§	1.6	+1.1	
4.5–5.0	—	2.1	1.0	1.5	1.4	0.8	0.7	0.1	0.5	0.9	0.3	§	0.2	-0.1	
5.5–6.0 (High)	—	1.5	0.2	1.4	0.5	0.7	0.2	0.5	0.4	0.3	0.8	§	0.4	-0.4	
Race/Ethnicity (2-year average): ^b															
White	—	—	1.5	1.2	0.9	0.6	0.4	0.4	0.3	0.3	0.4	§	0.6	+0.2	
African American	—	—	1.4	1.5	1.4	0.8	0.6	1.1	1.2	1.3	1.6	§	0.7	-0.9	
Hispanic	—	—	2.1	2.4	2.2	1.6	0.9	1.3	1.1	0.6	0.8	§	0.1	-0.7 s	

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. See Table S1 for the number of subgroup cases. See appendix B for definition of variables

in table. Data for 8th graders based on one of four forms; N is one third of N indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 38
SALVIA: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months													2019– 2021 change
	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	3.7	3.9	2.5	2.3	1.8	1.2	0.9	0.9	0.7	0.9	1.2	0.4	-0.5* s
Gender:														
Male	—	5.4	5.0	3.7	3.4	2.2	1.8	1.3	1.0	0.8	1.4	§	0.5	-0.9 s
Female	—	1.9	2.7	1.4	1.3	1.4	0.5	0.5	0.8	0.6	0.5	§	0.3	-0.1
College Plans:														
None or under 4 years	—	8.1	12.1	8.2	5.4	5.7	3.8	1.9	1.4	1.4	1.7	§	1.1	-0.6
Complete 4 years	—	3.2	2.9	1.9	2.0	1.3	0.9	0.8	0.8	0.6	0.8	§	0.3	-0.4 s
Region:														
Northeast	—	2.8	5.0	3.6	2.6	1.4	1.5	0.9	0.3	1.0	0.2	§	0.4	+0.2
Midwest	—	3.1	3.4	1.3	0.9	1.3	0.7	1.0	0.9	0.6	0.6	§	0.1	-0.5 s
South	—	4.0	3.4	2.0	1.8	1.9	1.0	0.7	0.7	0.3	0.8	§	0.6	-0.2
West	—	4.7	4.2	3.8	4.1	2.6	1.7	1.3	1.7	1.4	1.9	§	0.5	-1.4 s
Population Density:														
Large MSA	—	4.3	3.8	2.3	2.5	2.1	1.1	1.0	1.1	1.0	0.6	§	0.3	-0.3
Other MSA	—	3.3	3.8	2.6	2.1	1.4	1.2	0.7	0.7	0.7	1.2	§	0.6	-0.7
Non-MSA	—	3.7	4.1	2.7	2.6	2.4	1.3	1.3	1.1	0.4	0.6	§	0.4	-0.2
Parental Education: ^a														
1.0–2.0 (Low)	—	5.8	2.6	3.6	3.3	1.6	1.6	0.8	1.9	2.3	1.1	§	0.3	-0.8
2.5–3.0	—	4.4	5.1	3.3	3.0	1.8	1.8	1.4	0.9	0.4	1.1	§	0.5	-0.6
3.5–4.0	—	3.6	4.6	2.2	2.1	2.3	1.1	0.8	0.9	0.5	1.3	§	0.8	-0.4
4.5–5.0	—	2.7	3.3	2.0	2.3	1.6	1.0	0.7	1.0	0.6	0.7	§	0.0	-0.7
5.5–6.0 (High)	—	4.0	2.8	2.5	1.2	1.3	0.6	0.6	0.3	0.3	0.5	§	0.3	-0.2
Race/Ethnicity (2-year average): ^b														
White	—	—	3.9	3.4	2.3	2.1	1.6	1.0	0.8	0.6	0.7	§	0.7	0.0
African American	—	—	1.6	0.7	1.3	1.3	1.7	1.2	0.7	0.7	0.9	§	1.7	+0.8
Hispanic	—	—	4.7	4.0	2.8	1.6	0.8	1.1	1.2	1.2	1.1	§	0.8	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data for 10th graders based on one of four forms; *N* is one third of *N* indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 39
SALVIA: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months													2019– 2021 change
	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	5.7	5.5	5.9	4.4	3.4	1.8	1.9	1.8	1.5	0.9	0.7	0.7	0.6	-0.1*
Gender:														
Male	9.3	8.5	8.6	5.6	5.0	2.3	2.4	2.2	1.0	1.0	0.7	§	0.7	0.0
Female	2.3	2.5	3.3	3.0	1.8	1.0	1.0	0.9	1.6	0.4	0.7	§	0.4	-0.3
College Plans:														
None or under 4 years	7.4	9.2	11.4	5.9	6.7	3.4	3.0	3.6	2.9	1.4	0.8	§	0.8	0.0
Complete 4 years	5.2	4.7	4.8	3.9	2.5	1.4	1.4	1.2	1.1	0.5	0.6	§	0.4	-0.2
Region:														
Northeast	7.1	6.2	6.0	5.3	3.9	2.9	1.3	1.8	2.8	2.3	0.0	§	0.6	+0.6
Midwest	5.0	5.0	3.1	2.6	1.3	1.5	1.3	1.0	1.6	0.4	0.3	§	1.4	+1.0
South	4.8	5.3	5.2	3.8	3.6	1.7	1.4	1.6	1.2	1.2	0.9	§	0.5	-0.4
West	6.9	6.1	9.9	6.5	5.2	1.5	3.5	2.8	1.3	0.2	1.2	§	0.3	-0.9
Population Density:														
Large MSA	7.8	6.7	7.0	5.0	2.8	2.3	2.0	1.4	1.7	1.3	1.0	§	0.4	-0.6
Other MSA	5.0	5.5	6.1	4.7	4.2	1.9	1.4	2.6	1.2	0.4	0.4	§	0.9	+0.5
Non-MSA	4.6	4.1	4.1	2.9	2.5	1.0	2.7	0.6	1.9	1.3	0.8	§	0.4	-0.4
Parental Education: ^a														
1.0–2.0 (Low)	5.8	5.4	6.3	8.3	7.1	1.9	2.5	4.5	1.0	1.1	1.8	§	0.0	-1.8
2.5–3.0	4.6	4.4	5.3	3.7	4.1	1.9	1.6	1.2	1.8	0.9	0.4	§	0.4	0.0
3.5–4.0	6.2	6.0	6.6	3.5	3.1	1.6	2.1	1.1	2.8	1.5	0.8	§	0.0	-0.8
4.5–5.0	5.5	5.6	5.4	4.6	2.6	1.7	1.0	1.6	0.5	0.6	0.0	§	0.8	+0.8
5.5–6.0 (High)	7.0	6.7	6.0	4.2	2.3	1.9	1.7	1.6	0.2	0.1	0.9	§	1.9	+1.0
Race/Ethnicity (2-year average): ^b														
White	—	6.3	6.2	5.3	3.8	2.5	1.7	1.5	1.6	1.4	0.5	§	0.5	0.0
African American	—	2.5	2.5	2.8	2.6	1.7	1.6	1.9	1.8	0.9	0.3	§	1.1	+0.8
Hispanic	—	4.4	4.6	5.6	4.8	2.0	1.0	1.8	1.6	1.4	1.8	§	0.9	-0.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table Data for 12th graders based on two of six forms; *N* is two sixths of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 40
COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>
Total	1.1	1.5	1.7	2.1	2.6	3.0	2.8	3.1	2.7	2.6	2.5	2.3	2.2	2.0	2.2
Gender:															
Male	1.4	1.5	1.9	2.1	2.5	2.7	3.1	2.9	2.8	2.6	2.4	2.2	1.9	1.7	2.0
Female	0.9	1.5	1.5	2.1	2.6	3.1	2.5	3.1	2.7	2.6	2.6	2.3	2.3	2.3	2.3
College Plans:															
None or under 4 years	3.2	4.8	5.4	6.6	7.0	7.9	7.5	9.4	8.1	7.5	7.0	7.2	7.1	6.8	6.1
Complete 4 years	0.8	1.0	1.1	1.5	2.0	2.2	2.2	2.3	2.0	2.0	2.0	1.8	1.6	1.5	1.8
Region:															
Northeast	1.3	0.8	1.0	2.2	2.2	2.6	2.4	1.9	2.2	1.6	1.9	1.6	1.6	1.9	1.8
Midwest	0.9	1.4	1.0	1.2	2.6	2.9	2.6	2.7	2.6	2.9	2.3	2.3	2.0	1.6	2.0
South	1.1	1.7	2.1	2.5	2.4	2.7	2.6	3.8	3.1	2.2	3.2	2.6	2.5	2.1	2.4
West	1.5	2.0	2.7	2.3	3.3	3.7	3.7	3.3	2.7	3.7	2.4	2.4	2.3	2.4	2.4
Population Density:															
Large MSA	1.1	1.4	1.3	1.9	2.4	2.8	2.4	2.3	2.1	2.3	2.1	2.0	1.7	2.2	2.2
Other MSA	1.1	1.7	2.2	2.5	2.8	3.2	2.9	3.3	2.8	2.5	2.8	2.3	2.2	2.1	2.2
Non-MSA	1.2	1.3	1.2	1.4	2.4	2.7	3.0	3.4	3.2	3.1	2.5	2.6	2.7	1.7	2.2
Parental Education: ^a															
1.0–2.0 (Low)	2.4	3.2	2.9	3.5	4.9	3.9	4.7	6.3	5.6	5.3	4.4	3.7	4.1	3.3	4.2
2.5–3.0	1.4	1.6	2.0	2.3	2.4	3.3	3.0	3.3	3.1	2.9	2.8	2.8	2.4	2.5	2.9
3.5–4.0	0.7	1.2	1.8	2.1	2.8	3.3	2.8	3.1	2.8	2.5	2.4	2.2	2.2	2.0	2.5
4.5–5.0	0.7	1.0	1.0	1.6	1.9	2.7	2.6	2.2	1.7	1.8	2.1	2.0	1.5	1.8	1.5
5.5–6.0 (High)	1.2	1.5	1.1	1.9	2.5	2.5	2.3	2.5	2.5	2.2	1.6	1.5	2.0	1.2	1.2
Race/Ethnicity (2-year average): ^b															
White	—	1.2	1.3	1.6	2.3	2.8	3.0	2.8	2.6	2.5	2.5	2.4	2.2	1.9	1.8
African American	—	0.7	0.7	0.7	0.6	0.6	0.5	0.7	0.8	0.8	0.8	0.8	0.9	1.0	1.0
Hispanic	—	3.1	4.0	4.5	4.7	4.8	4.3	5.2	5.9	4.7	4.2	4.1	3.8	3.6	3.6

Table continued on next page.

TABLE 40 (cont.)
COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021
	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	<u>change</u>
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	2.0	2.0	1.8	1.6	1.6	1.4	1.2	1.0	1.0	0.9	0.8	0.8	0.8	0.7	0.5	0.2	-0.5* sss
Gender:																	
Male	1.9	1.7	1.8	1.4	1.5	1.3	1.1	1.0	0.9	0.9	0.8	0.7	0.8	0.7	§	0.2	-0.5 sss
Female	2.1	2.2	1.8	1.7	1.7	1.3	1.2	1.0	1.1	0.8	0.7	0.8	0.6	0.8	§	0.2	-0.6 sss
College Plans:																	
None or under 4 years	7.1	6.5	6.9	6.4	5.6	5.5	4.6	3.4	3.0	2.5	2.3	2.6	2.4	1.3	§	0.6	-0.7
Complete 4 years	1.5	1.5	1.3	1.2	1.3	1.1	0.9	0.8	0.8	0.8	0.6	0.5	0.5	0.6	§	0.2	-0.5 sss
Region:																	
Northeast	1.3	1.2	1.2	1.4	1.0	0.9	1.1	0.8	0.5	0.6	0.5	0.5	0.5	0.3	§	0.0	-0.3
Midwest	1.3	1.6	1.9	1.8	2.0	1.3	0.6	0.8	0.8	0.8	0.6	0.6	0.5	0.4	§	0.2	-0.2
South	2.9	2.3	2.2	1.2	1.6	1.3	1.4	1.0	1.3	1.0	1.0	0.6	0.7	0.9	§	0.3	-0.5 s
West	1.9	2.2	1.6	2.1	1.7	2.1	1.4	1.3	1.3	1.1	0.9	1.3	1.4	1.0	§	0.3	-0.8 ss
Population Density:																	
Large MSA	1.9	1.9	2.0	1.1	1.4	1.5	1.3	1.0	1.2	1.0	0.8	0.7	0.8	0.8	§	0.1	-0.7 sss
Other MSA	2.0	2.1	1.7	1.9	1.8	1.5	1.2	1.0	1.0	1.1	0.7	0.9	0.7	0.7	§	0.4	-0.3 s
Non-MSA	2.2	1.8	1.7	1.4	1.5	1.2	0.9	0.8	1.0	0.4	1.0	0.5	0.7	0.5	§	0.2	-0.3
Parental Education: ^a																	
1.0–2.0 (Low)	4.4	3.8	3.6	3.9	3.3	2.6	2.6	2.1	1.8	1.2	1.8	1.3	1.7	1.5	§	0.6	-0.9
2.5–3.0	2.2	2.2	2.3	1.8	2.0	2.1	1.6	1.1	1.1	0.9	0.8	1.1	0.8	0.8	§	0.2	-0.6 s
3.5–4.0	2.0	1.8	1.8	1.5	1.6	1.6	1.2	0.9	1.1	0.8	0.8	0.8	0.7	0.8	§	0.4	-0.4
4.5–5.0	1.4	1.4	1.2	0.8	1.2	0.9	0.5	0.7	0.9	0.6	0.6	0.5	0.7	0.6	§	0.2	-0.3
5.5–6.0 (High)	1.5	1.3	1.4	1.5	1.2	0.8	0.7	0.8	0.6	1.0	0.5	0.3	0.4	0.6	§	0.1	-0.4
Race/Ethnicity (2-year average): ^b																	
White	1.9	1.7	1.6	1.5	1.4	1.3	1.0	0.9	0.7	0.7	0.6	0.5	0.5	0.5	§	0.0	-0.5 sss
African American	0.9	0.9	0.8	0.8	0.7	0.5	0.6	0.8	0.8	0.7	0.6	1.1	1.2	1.3	§	0.2	-1.1 sss
Hispanic	3.5	3.8	3.5	2.8	2.8	2.9	2.6	1.8	1.4	1.3	1.2	1.7	1.8	1.5	§	0.6	-0.9 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 41
COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	
<i>Approximate weighted N =</i>	<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>	
Total	2.2	1.9	2.1	2.8	3.5	4.2	4.7	4.7	4.9	4.4	3.6	4.0	3.3	3.7	3.5	
Gender:																
Male	2.2	2.0	2.5	3.1	3.5	4.5	4.7	4.9	5.2	4.7	3.8	4.2	3.3	3.9	3.6	
Female	2.2	1.7	1.6	2.5	3.3	4.0	4.6	4.4	4.6	4.1	3.2	3.9	3.2	3.4	3.3	
College Plans:																
None or under 4 years	4.7	4.0	5.1	6.6	7.2	10.0	10.4	10.7	11.6	9.7	9.1	10.1	8.3	9.4	8.8	
Complete 4 years	1.7	1.4	1.4	2.0	2.8	3.2	3.7	3.6	3.8	3.5	2.7	3.1	2.4	2.9	2.8	
Region:																
Northeast	1.5	1.0	2.0	2.4	2.5	3.0	3.0	4.9	4.6	3.1	2.3	2.4	3.1	3.4	3.6	
Midwest	1.7	1.7	1.4	2.2	2.9	4.1	4.0	3.7	4.4	4.6	3.4	3.6	2.7	3.2	3.4	
South	2.0	1.8	1.9	2.6	3.5	4.2	5.4	4.3	5.2	4.2	3.8	3.8	3.5	3.5	2.9	
West	3.6	3.2	3.7	4.7	5.3	5.9	6.4	6.4	5.3	5.7	4.9	6.5	3.7	4.8	4.4	
Population Density:																
Large MSA	1.9	1.6	1.6	2.3	3.4	3.8	4.5	4.3	4.1	4.3	3.2	3.6	2.5	3.3	3.3	
Other MSA	2.7	2.1	2.3	3.1	3.5	4.7	4.3	4.7	5.1	4.2	3.7	4.4	3.2	3.8	3.4	
Non-MSA	1.6	1.7	2.1	2.7	3.6	3.7	5.7	5.2	5.4	4.7	3.8	3.8	4.6	3.9	3.8	
Parental Education: ^a																
1.0–2.0 (Low)	3.3	3.5	3.2	3.8	5.3	7.4	6.3	8.1	8.2	7.7	5.6	7.9	7.0	6.5	5.3	
2.5–3.0	2.4	1.7	2.2	2.9	4.3	4.5	5.0	5.5	5.9	5.4	5.2	4.8	3.9	4.2	4.6	
3.5–4.0	2.4	2.1	2.5	3.2	3.7	4.3	5.4	4.4	4.7	4.4	2.9	3.8	3.4	4.0	3.4	
4.5–5.0	1.6	1.4	1.6	2.1	2.6	3.4	3.7	3.5	3.9	3.2	2.7	2.4	2.2	3.0	2.8	
5.5–6.0 (High)	1.9	1.5	1.1	1.9	1.9	3.4	3.3	3.2	3.9	2.9	2.0	3.2	1.6	1.8	2.4	
Race/Ethnicity (2-year average): ^b																
White	—	2.1	2.0	2.2	3.0	3.8	4.4	4.7	4.9	4.7	3.9	3.9	3.9	3.7	3.7	
African American	—	0.6	0.6	1.0	0.9	0.7	0.8	1.0	0.9	0.6	0.8	1.0	0.9	0.8	1.0	
Hispanic	—	3.7	3.7	4.9	5.5	7.0	8.5	8.3	8.2	8.0	6.6	6.0	6.1	5.8	6.4	

Table continued on next page.

TABLE 41 (cont.)
COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	3.2	3.4	3.0	2.7	2.2	1.9	2.0	1.9	1.5	1.8	1.3	1.4	1.5	1.5	1.1	0.6	-0.8* ss
Gender:																	
Male	3.2	3.6	3.0	2.8	2.7	2.3	2.3	2.5	1.8	1.9	1.5	1.7	1.8	1.5	§	0.6	-0.9 s
Female	3.1	3.3	2.8	2.4	1.6	1.4	1.6	1.3	1.0	1.7	1.0	1.0	1.3	1.3	§	0.4	-0.9 ss
College Plans:																	
None or under 4 years	9.0	8.6	8.7	8.5	7.2	5.5	6.8	4.1	4.8	4.3	4.0	3.8	3.8	2.3	§	1.1	-1.2 s
Complete 4 years	2.4	2.8	2.2	2.0	1.6	1.5	1.5	1.6	1.1	1.6	0.9	1.1	1.2	1.2	§	0.3	-0.9 ss
Region:																	
Northeast	2.5	2.9	2.0	2.4	2.0	2.0	1.3	1.2	1.2	1.7	0.8	0.9	1.0	0.6	§	0.8	+0.2
Midwest	3.2	3.6	2.5	2.4	1.8	1.3	1.4	1.3	1.3	1.2	1.0	1.4	1.1	1.0	§	0.5	-0.6 s
South	3.2	3.5	3.5	2.4	1.9	1.9	2.2	2.0	1.5	1.8	1.4	1.3	2.1	1.8	§	0.7	-1.1
West	3.8	3.6	3.3	3.6	3.3	2.4	2.7	2.7	1.7	2.5	1.6	1.8	1.4	1.8	§	0.5	-1.4
Population Density:																	
Large MSA	2.8	3.3	2.8	2.4	2.1	1.9	2.1	1.8	1.2	1.7	1.3	1.8	1.4	0.9	§	0.5	-0.4
Other MSA	3.2	3.4	3.3	3.1	2.3	2.0	1.7	2.1	1.3	1.7	1.2	1.3	1.9	1.9	§	0.6	-1.3 s
Non-MSA	3.8	3.7	2.4	2.2	2.1	1.6	2.2	1.3	2.2	2.1	1.4	0.9	1.0	1.3	§	0.9	-0.4
Parental Education: ^a																	
1.0–2.0 (Low)	6.5	6.4	6.8	6.0	4.4	3.4	5.3	3.1	2.7	3.9	1.4	2.0	2.5	2.2	§	1.3	-0.9
2.5–3.0	4.1	3.9	3.5	3.3	2.5	2.0	2.3	2.4	1.5	2.3	1.9	1.9	1.8	1.9	§	0.9	-1.0
3.5–4.0	2.9	3.5	2.7	2.7	1.9	1.9	1.6	1.6	1.8	1.9	1.0	1.1	1.7	1.7	§	0.6	-1.1
4.5–5.0	2.0	2.5	2.2	1.6	1.8	1.4	1.5	1.4	0.8	1.1	1.1	1.0	1.0	0.9	§	0.1	-0.7 ss
5.5–6.0 (High)	3.0	2.5	1.6	2.0	1.4	1.3	1.2	1.6	1.1	1.2	0.8	1.2	1.5	1.0	§	0.4	-0.6
Race/Ethnicity (2-year average): ^b																	
White	3.5	3.5	3.2	2.6	2.1	1.7	1.6	1.6	1.5	1.5	1.5	1.3	1.3	1.2	§	0.5	-0.7 ss
African American	1.0	0.8	0.8	0.9	0.9	0.7	1.3	1.6	0.9	1.0	0.8	0.7	0.6	0.3	§	0.6	+0.3
Hispanic	5.9	4.9	5.2	5.2	4.6	4.1	3.8	3.7	2.8	2.2	1.9	1.5	2.3	2.6	§	1.4	-1.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 42
COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Approximate weighted N =	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	5.6	6.0	7.2	9.0	12.0	12.3	12.4	11.5	11.4	11.6	13.1	12.7	10.3	7.9	6.5	5.3
Gender:																
Male	7.5	7.5	9.3	11.4	14.6	14.8	13.8	13.1	13.2	13.8	14.8	14.3	11.3	9.1	8.1	6.6
Female	3.9	4.4	4.9	6.5	9.3	9.8	10.4	9.6	9.3	9.1	11.2	10.9	9.2	6.5	4.9	3.8
College Plans:																
None or under 4 years	—	6.6	8.1	9.5	13.7	13.2	12.4	12.5	12.2	13.2	14.7	15.7	12.4	9.7	9.3	7.8
Complete 4 years	—	5.0	5.5	7.7	9.5	10.8	11.5	9.9	9.9	9.7	11.4	10.4	9.0	6.7	5.3	4.1
Region:																
Northeast	5.3	6.6	7.9	11.8	13.8	14.2	16.8	16.9	15.2	19.5	20.8	17.9	13.3	9.1	7.3	6.5
Midwest	5.1	5.5	6.3	8.5	10.5	10.9	9.4	9.0	8.0	5.8	8.2	10.1	7.5	6.1	5.3	4.1
South	5.4	5.1	6.0	6.8	8.5	7.8	6.8	6.3	7.7	7.7	7.5	7.1	7.0	6.2	6.0	4.8
West	7.8	7.9	10.2	10.7	18.6	20.6	22.1	17.9	19.2	19.3	19.7	20.0	16.4	12.1	8.5	6.6
Population Density:																
Large MSA	7.3	8.6	8.6	12.3	16.6	18.7	17.5	17.2	16.9	16.8	18.8	18.8	12.9	9.3	6.4	5.6
Other MSA	5.9	5.8	7.3	8.9	11.7	11.3	11.5	10.1	11.2	11.0	12.4	12.0	10.1	8.5	7.1	5.4
Non-MSA	4.3	4.3	5.8	6.4	8.9	8.9	9.4	8.5	7.3	8.3	9.2	9.0	8.1	5.3	5.4	4.8
Parental Education: ^a																
1.0–2.0 (Low)	4.5	5.3	5.5	6.3	8.4	9.0	8.3	7.6	9.0	9.4	12.0	10.5	8.7	7.6	6.7	4.7
2.5–3.0	4.6	6.1	6.8	8.7	11.1	11.2	10.5	11.0	9.8	10.9	12.7	12.9	9.9	7.4	6.4	5.6
3.5–4.0	4.5	5.9	7.2	9.0	13.2	13.3	13.3	12.5	11.7	12.2	14.0	13.6	11.2	7.2	6.4	5.6
4.5–5.0	6.3	7.6	8.1	10.4	14.0	13.6	14.9	13.6	13.1	12.2	13.7	12.2	10.0	8.7	7.1	4.4
5.5–6.0 (High)	5.2	7.1	9.5	11.6	15.2	16.3	16.2	13.8	15.1	13.4	11.9	12.5	10.8	8.1	5.8	5.5
Race/Ethnicity (2-year average): ^b																
White	—	—	6.5	8.3	10.9	12.8	13.0	12.6	11.8	11.9	13.0	13.5	12.0	9.6	7.6	6.3
African American	—	—	4.8	4.6	4.6	5.2	4.8	5.2	7.2	6.3	5.3	5.8	4.8	3.8	2.9	1.7
Hispanic	—	—	7.2	7.5	8.9	11.2	12.4	12.1	11.4	13.3	16.3	16.7	14.0	9.9	7.8	7.4

Table continued on next page.

TABLE 42 (cont.)
COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	3.5	3.1	3.3	3.6	4.0	4.9	5.5	5.7	6.2	5.0	4.8	5.0	4.8	5.3	5.1	5.7
Gender:																
Male	4.1	3.7	4.0	4.5	4.8	6.0	6.6	6.8	7.3	5.8	5.4	5.9	5.9	6.5	5.8	6.0
Female	2.6	2.4	2.3	2.8	3.1	3.5	4.2	4.5	5.0	3.9	4.1	4.0	3.7	4.1	4.2	5.2
College Plans:																
None or under 4 years	4.9	5.1	4.5	5.3	5.6	7.5	8.1	9.7	9.1	7.1	8.2	8.6	6.0	8.3	7.5	8.1
Complete 4 years	2.8	2.4	2.8	3.0	3.4	4.0	4.4	4.5	5.4	4.2	3.7	3.9	4.2	4.3	4.4	5.1
Region:																
Northeast	3.8	2.8	3.1	3.1	3.8	5.5	6.6	5.9	4.3	4.1	4.8	5.0	5.2	5.0	4.7	6.7
Midwest	3.2	2.5	2.4	3.7	3.4	3.8	4.7	5.8	6.2	4.8	5.7	5.2	3.9	4.2	5.0	4.2
South	3.0	3.2	3.1	3.4	3.6	4.6	4.8	5.8	6.9	4.7	3.9	5.0	4.7	5.4	5.3	6.4
West	4.4	4.3	4.9	4.5	5.8	6.1	6.8	5.4	6.9	6.3	5.0	4.6	5.8	7.0	5.0	5.5
Population Density:																
Large MSA	4.1	3.6	2.7	3.3	4.4	4.8	4.7	5.4	5.0	4.1	4.3	4.1	3.8	4.7	5.2	7.4
Other MSA	3.7	3.3	3.9	4.1	3.9	4.9	5.6	5.8	6.6	4.9	5.0	5.4	5.7	5.8	5.2	4.8
Non-MSA	2.5	2.4	2.7	3.2	3.9	4.9	6.0	6.0	6.9	6.1	5.2	5.3	4.6	5.0	4.7	4.9
Parental Education: ^a																
1.0–2.0 (Low)	3.5	3.9	3.5	4.1	4.8	5.3	6.5	6.9	9.0	6.2	5.7	6.6	4.8	7.3	4.3	6.1
2.5–3.0	3.8	3.3	3.0	4.0	3.9	5.0	5.5	6.3	6.0	4.6	6.0	5.3	4.1	5.2	6.6	6.3
3.5–4.0	3.7	3.0	3.8	3.8	4.2	5.0	5.6	6.0	6.8	5.0	4.8	5.3	5.2	5.3	5.2	5.7
4.5–5.0	3.1	2.9	3.0	3.1	3.7	4.8	5.2	5.0	5.4	5.3	4.4	4.2	4.7	5.2	4.3	5.1
5.5–6.0 (High)	2.4	2.6	2.4	3.3	3.4	4.3	4.4	4.4	5.2	3.9	2.6	4.3	4.4	4.3	4.1	5.7
Race/Ethnicity (2-year average): ^b																
White	4.6	3.3	3.1	3.5	4.0	4.5	5.5	6.3	6.7	6.2	5.5	5.7	5.6	5.6	5.6	5.8
African American	1.5	1.2	0.8	0.9	1.0	0.8	0.9	0.9	0.9	1.0	1.0	0.9	1.1	1.2	1.2	1.6
Hispanic	6.1	5.2	5.8	5.4	5.5	7.3	7.6	6.7	7.5	7.6	6.1	5.5	4.9	5.5	6.2	6.5

Table continued on next page.

TABLE 42 (cont.)
COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	5.2	4.4	3.4	2.9	2.9	2.7	2.6	2.6	2.5	2.3	2.7	2.3	2.2	2.9	1.2	-1.0* ss
Gender:																
Male	5.9	5.2	4.1	4.0	3.6	3.3	3.4	3.5	3.3	3.0	3.7	2.8	2.6	§	0.9	-1.7 sss
Female	4.4	3.5	2.7	1.9	2.1	2.0	1.7	1.6	1.7	1.5	1.9	1.4	1.7	§	1.3	-0.4
College Plans:																
None or under 4 years	8.8	7.4	5.4	5.2	5.4	4.2	5.1	3.9	4.4	3.2	4.5	4.0	4.1	§	2.1	-1.9 ss
Complete 4 years	4.4	3.6	2.8	2.4	2.3	2.3	1.9	2.2	2.1	1.9	2.2	1.8	1.6	§	0.8	-0.8 ss
Region:																
Northeast	6.0	4.5	2.8	3.3	2.8	2.1	2.7	2.5	2.3	1.6	2.6	2.3	1.3	§	0.8	-0.5
Midwest	5.0	3.2	3.6	2.4	2.9	2.2	1.9	2.3	1.8	1.9	2.0	2.0	1.5	§	0.8	-0.7
South	5.2	4.8	3.2	2.5	2.6	2.5	2.2	2.7	2.0	1.7	2.5	1.7	2.1	§	1.0	-1.1 ss
West	4.8	4.9	3.9	4.0	3.3	4.1	4.0	2.7	4.4	4.0	3.9	3.5	4.4	§	2.1	-2.2
Population Density:																
Large MSA	5.0	4.4	3.4	3.3	2.9	2.9	2.6	2.3	3.0	2.4	3.0	2.3	2.0	§	0.8	-1.1 ss
Other MSA	5.4	4.3	3.3	3.1	2.9	2.9	3.0	3.1	2.3	2.4	2.9	2.2	2.2	§	1.4	-0.8
Non-MSA	5.3	4.5	3.6	1.9	2.7	2.1	1.9	1.7	2.4	1.7	2.0	2.5	2.7	§	1.3	-1.4 s
Parental Education: ^a																
1.0–2.0 (Low)	6.2	4.7	4.1	3.2	4.2	3.0	4.2	3.0	4.0	2.7	3.2	3.3	2.3	§	1.0	-1.3 s
2.5–3.0	5.9	4.0	3.6	3.2	2.6	3.1	2.9	3.1	2.6	2.1	3.1	2.8	2.2	§	1.8	-0.4
3.5–4.0	5.5	4.9	3.4	3.0	3.0	2.6	3.0	2.4	2.3	2.0	2.8	1.7	2.1	§	1.3	-0.8
4.5–5.0	4.9	4.0	3.2	2.5	2.3	2.6	1.6	2.2	2.2	2.0	2.0	2.0	2.3	§	0.9	-1.4 sss
5.5–6.0 (High)	4.0	4.1	2.5	2.4	3.0	2.4	2.1	2.6	2.2	2.4	2.7	1.9	1.6	§	0.7	-0.9
Race/Ethnicity (2-year average): ^b																
White	6.1	5.6	4.4	3.4	3.1	3.1	2.6	2.3	2.3	2.2	2.6	2.7	2.2	§	1.6	-0.6
African American	1.3	0.8	0.7	0.9	1.1	0.9	0.9	1.6	1.5	0.9	0.8	0.8	0.9	§	1.0	0.0
Hispanic	6.0	4.7	4.0	3.5	3.3	3.4	3.8	3.6	3.4	3.6	3.2	2.9	2.7	§	1.5	-1.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 43
CRACK: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>
Total	0.7	0.9	1.0	1.3	1.6	1.8	1.7	2.1	1.8	1.8	1.7	1.6	1.6	1.3	1.4
Gender:															
Male	0.8	0.9	1.1	1.3	1.5	1.7	1.8	2.1	1.8	1.7	1.6	1.6	1.3	1.1	1.3
Female	0.5	0.9	0.9	1.2	1.6	1.9	1.5	2.1	1.8	1.8	1.8	1.6	1.7	1.5	1.5
College Plans:															
None or under 4 years	2.0	2.9	3.4	4.6	4.2	4.9	4.7	7.5	5.3	5.2	4.4	5.7	5.8	4.9	3.3
Complete 4 years	0.4	0.6	0.6	0.8	1.3	1.3	1.3	1.5	1.4	1.4	1.4	1.2	1.1	1.0	1.2
Region:															
Northeast	0.5	0.4	0.4	1.4	1.4	1.7	1.6	1.2	1.5	1.2	1.2	1.2	1.3	1.2	0.9
Midwest	0.6	1.0	0.8	0.9	1.4	1.9	1.6	1.9	1.9	1.9	1.7	1.5	1.3	1.2	1.4
South	0.7	1.0	1.2	1.6	1.4	1.7	1.4	2.5	1.9	1.4	2.0	1.7	1.8	1.3	1.5
West	0.8	1.3	1.4	1.3	2.3	2.1	2.3	2.6	1.8	2.9	1.7	2.0	1.8	1.8	1.6
Population Density:															
Large MSA	0.5	0.8	0.7	1.3	1.5	1.8	1.5	1.6	1.2	1.6	1.4	1.5	1.2	1.5	1.5
Other MSA	0.7	1.1	1.2	1.5	1.7	2.0	1.8	2.2	2.0	1.8	1.8	1.6	1.7	1.3	1.3
Non-MSA	0.8	0.8	0.9	1.0	1.4	1.7	1.7	2.6	2.1	2.0	1.8	1.8	1.7	1.2	1.4
Parental Education: ^a															
1.0–2.0 (Low)	1.7	2.2	1.8	2.8	3.0	2.7	3.0	5.0	3.6	3.7	3.2	3.1	3.3	2.2	3.1
2.5–3.0	0.7	0.8	1.0	1.4	1.2	2.1	2.0	2.2	2.0	2.1	1.8	2.1	1.7	1.7	1.8
3.5–4.0	0.4	0.7	1.2	0.9	1.7	2.0	1.4	2.1	2.0	1.8	1.3	1.4	1.6	1.3	1.6
4.5–5.0	0.4	0.6	0.5	1.1	1.3	1.5	1.5	1.6	1.0	1.0	1.5	1.2	1.0	1.1	0.9
5.5–6.0 (High)	0.8	1.0	0.6	1.4	1.6	1.5	1.5	1.6	1.9	1.8	1.3	1.3	1.3	1.0	0.6
Race/Ethnicity (2-year average): ^b															
White	—	0.7	0.8	1.0	1.4	1.7	1.7	1.7	1.8	1.7	1.7	1.5	1.5	1.3	1.2
African American	—	0.4	0.4	0.5	0.5	0.4	0.4	0.5	0.5	0.4	0.5	0.6	0.7	0.8	0.8
Hispanic	—	1.9	2.0	2.1	2.7	3.0	2.8	3.6	3.9	2.9	2.6	2.7	2.8	2.6	2.5

Table continued on next page.

TABLE 43 (cont.)
CRACK: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	1.3	1.3	1.1	1.1	1.0	0.9	0.6	0.6	0.7	0.5	0.5	0.5	0.4	0.4	0.2	0.2	-0.2* ss
Gender:																	
Male	1.2	1.2	1.0	0.9	0.9	0.8	0.6	0.5	0.6	0.5	0.5	0.5	0.4	0.5	§	0.1	-0.4 sss
Female	1.3	1.5	1.2	1.2	1.1	1.0	0.7	0.7	0.8	0.6	0.5	0.6	0.4	0.3	§	0.2	-0.2
College Plans:																	
None or under 4 years	4.4	4.1	4.3	4.8	3.2	3.5	2.8	2.2	2.2	1.6	1.6	1.9	1.3	0.8	§	0.4	-0.4
Complete 4 years	1.0	1.1	0.8	0.8	0.8	0.7	0.5	0.5	0.6	0.4	0.4	0.4	0.3	0.4	§	0.1	-0.2 ss
Region:																	
Northeast	0.9	0.8	0.8	1.0	0.8	0.5	0.8	0.6	0.3	0.4	0.2	0.4	0.2	0.3	§	0.0	-0.3
Midwest	0.7	1.1	1.2	1.2	1.1	0.9	0.4	0.6	0.6	0.3	0.5	0.5	0.3	0.3	§	0.2	-0.1
South	1.8	1.5	1.2	0.7	0.9	0.9	0.7	0.7	0.9	0.6	0.7	0.4	0.4	0.5	§	0.3	-0.3
West	1.3	1.8	1.2	1.7	1.2	1.3	0.6	0.6	0.6	0.7	0.5	0.9	0.8	0.5	§	0.2	-0.3 s
Population Density:																	
Large MSA	1.3	1.1	1.4	0.8	1.0	1.0	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5	§	0.1	-0.4 ss
Other MSA	1.1	1.6	1.0	1.3	1.1	0.9	0.6	0.6	0.7	0.6	0.5	0.6	0.4	0.4	§	0.3	-0.1
Non-MSA	1.6	1.2	1.0	0.9	0.8	0.9	0.4	0.6	0.7	0.3	0.6	0.3	0.4	0.4	§	0.2	-0.2
Parental Education: ^a																	
1.0–2.0 (Low)	3.0	2.9	2.8	3.3	1.8	2.0	1.6	1.2	1.2	0.9	1.4	1.0	1.1	0.6	§	0.4	-0.2
2.5–3.0	1.5	1.4	1.6	0.9	1.1	1.4	0.8	0.7	0.9	0.7	0.6	0.7	0.4	0.3	§	0.2	-0.1
3.5–4.0	1.2	1.3	0.8	1.0	0.9	0.9	0.7	0.5	0.6	0.4	0.6	0.5	0.5	0.4	§	0.2	-0.2
4.5–5.0	0.9	0.9	0.7	0.6	0.8	0.6	0.2	0.5	0.5	0.4	0.3	0.4	0.2	0.4	§	0.2	-0.2
5.5–6.0 (High)	1.0	1.0	0.9	1.1	0.9	0.5	0.3	0.6	0.5	0.3	0.2	0.3	0.2	0.6	§	0.1	-0.4
Race/Ethnicity (2-year average): ^b																	
White	1.1	1.0	1.0	0.8	0.8	0.7	0.6	0.4	0.4	0.4	0.3	0.3	0.3	0.3	§	0.1	-0.1
African American	0.6	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.6	0.6	0.6	0.5	0.5	0.4	§	0.0	-0.3 sss
Hispanic	2.6	2.7	2.5	2.2	2.1	2.1	1.7	1.1	0.9	0.8	0.8	0.8	0.8	0.8	§	0.1	-0.6 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 44
CRACK: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>
Total	0.9	0.9	1.1	1.4	1.8	2.1	2.2	2.5	2.4	2.2	1.8	2.3	1.6	1.7	1.7
Gender:															
Male	0.9	0.9	1.3	1.6	1.9	2.1	2.3	2.7	2.5	2.3	1.9	2.5	1.6	1.8	1.6
Female	0.8	0.9	0.7	1.0	1.6	2.1	2.2	2.2	2.3	2.1	1.7	2.2	1.6	1.5	1.7
College Plans:															
None or under 4 years	2.4	2.1	2.7	3.4	3.7	5.0	4.9	6.0	5.5	5.2	4.7	5.6	4.5	4.6	4.8
Complete 4 years	0.6	0.6	0.7	0.9	1.5	1.5	1.8	1.9	1.8	1.7	1.4	1.8	1.1	1.2	1.3
Region:															
Northeast	0.5	0.4	1.1	1.4	1.1	1.4	1.5	2.6	2.5	1.7	1.0	1.4	1.3	1.4	1.8
Midwest	0.9	0.9	0.8	1.0	1.5	2.2	2.1	2.1	2.1	1.8	1.8	2.2	1.4	1.5	1.4
South	1.0	0.8	0.9	1.3	1.9	2.0	2.0	1.9	2.0	1.9	1.7	1.7	1.7	1.4	1.3
West	1.1	1.4	1.7	1.9	2.8	2.8	3.8	3.9	3.2	3.8	3.3	4.4	2.0	2.6	2.5
Population Density:															
Large MSA	0.9	0.8	0.7	0.9	1.9	1.7	2.3	2.2	2.2	2.4	1.9	2.2	1.2	1.5	1.8
Other MSA	0.9	0.9	1.1	1.5	1.6	2.4	1.7	2.4	2.3	2.0	1.8	2.5	1.7	1.8	1.5
Non-MSA	0.9	0.9	1.2	1.6	2.3	1.9	3.3	3.1	2.8	2.5	1.8	2.2	2.1	1.7	1.9
Parental Education: ^a															
1.0–2.0 (Low)	1.3	1.7	1.8	1.9	3.0	3.9	3.4	4.5	3.4	4.8	3.0	3.9	2.9	3.5	2.6
2.5–3.0	1.0	0.8	1.0	1.1	2.4	2.5	2.4	3.1	2.9	2.9	2.5	2.6	1.8	1.9	2.5
3.5–4.0	0.9	1.0	1.4	1.5	1.7	1.9	2.6	2.0	2.5	2.2	1.4	2.2	1.6	1.6	1.4
4.5–5.0	0.7	0.6	0.7	1.0	1.3	1.4	1.8	2.0	1.9	1.1	1.6	1.4	1.3	1.4	1.3
5.5–6.0 (High)	0.7	0.9	0.5	1.1	1.1	1.8	1.2	1.8	1.8	1.4	1.2	2.2	0.8	0.8	1.0
Race/Ethnicity (2-year average): ^b															
White	—	0.9	0.9	1.1	1.5	1.9	2.2	2.3	2.4	2.2	1.8	2.0	2.0	1.6	1.5
African American	—	0.3	0.4	0.8	0.6	0.4	0.4	0.5	0.5	0.5	0.7	0.8	0.6	0.4	0.6
Hispanic	—	1.5	1.7	1.9	2.5	3.7	3.7	4.1	4.4	4.0	3.7	3.6	3.5	3.1	3.6

Table continued on next page.

TABLE 44 (cont.)
CRACK: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															2019-2021 change	
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	1.3	1.3	1.3	1.2	1.0	0.9	0.8	0.8	0.5	0.7	0.4	0.6	0.6	0.6	0.5	0.3	-0.2*
Gender:																	
Male	1.3	1.5	1.3	1.1	1.0	1.0	0.9	0.9	0.6	0.7	0.6	0.6	0.5	0.6	§	0.3	-0.3
Female	1.2	1.1	1.2	1.2	0.9	0.8	0.6	0.7	0.5	0.6	0.3	0.5	0.6	0.5	§	0.3	-0.2
College Plans:																	
None or under 4 years	3.7	3.8	3.9	4.1	3.4	2.5	3.5	1.3	1.8	2.2	1.7	2.4	1.3	0.9	§	0.6	-0.3
Complete 4 years	1.0	1.0	0.9	0.9	0.7	0.7	0.5	0.8	0.4	0.5	0.3	0.4	0.5	0.4	§	0.2	-0.2 s
Region:																	
Northeast	0.9	1.1	1.0	1.2	1.0	1.1	0.5	0.6	0.5	0.7	0.3	0.6	0.3	0.3	§	0.3	0.0
Midwest	1.3	1.1	1.1	1.4	0.9	0.6	0.7	0.7	0.6	0.5	0.3	0.4	0.4	0.5	§	0.2	-0.3
South	1.2	1.4	1.1	0.8	0.9	0.8	0.6	0.8	0.5	0.7	0.6	0.5	0.9	0.8	§	0.5	-0.3
West	1.7	1.6	1.8	1.5	1.4	1.3	1.3	1.2	0.5	1.1	0.6	0.8	0.4	0.5	§	0.3	-0.2
Population Density:																	
Large MSA	1.3	1.3	1.1	0.8	1.0	0.9	1.0	0.7	0.5	0.7	0.5	0.6	0.6	0.3	§	0.2	0.0
Other MSA	1.1	1.3	1.4	1.5	1.0	1.0	0.5	1.0	0.4	0.7	0.4	0.6	0.6	0.7	§	0.3	-0.4
Non-MSA	1.6	1.3	1.2	1.0	1.2	0.8	0.9	0.6	0.8	1.0	0.6	0.5	0.4	0.7	§	0.6	0.0
Parental Education: ^a																	
1.0–2.0 (Low)	3.9	3.8	3.3	2.5	2.5	1.6	1.9	1.5	0.4	2.1	0.5	1.0	1.0	0.8	§	0.8	0.0
2.5–3.0	1.6	1.6	1.3	1.6	0.9	0.9	1.0	1.0	0.6	0.8	0.7	0.9	0.5	0.9	§	0.3	-0.6
3.5–4.0	1.0	0.9	1.1	1.3	0.8	0.9	0.5	0.6	0.8	0.6	0.3	0.5	0.4	0.6	§	0.4	-0.2
4.5–5.0	0.8	0.9	1.0	0.7	0.8	0.6	0.6	0.7	0.3	0.4	0.3	0.3	0.4	0.2	§	0.0	-0.2 s
5.5–6.0 (High)	1.0	0.7	0.7	0.6	0.9	0.8	0.4	0.8	0.4	0.6	0.4	0.3	0.8	0.5	§	0.3	-0.2
Race/Ethnicity (2-year average): ^b																	
White	1.4	1.2	1.1	1.0	0.9	0.7	0.6	0.5	0.6	0.5	0.4	0.4	0.5	0.5	§	0.2	-0.3 s
African American	0.6	0.5	0.5	0.5	0.6	0.7	0.6	0.8	0.7	0.8	0.7	0.4	0.3	0.2	§	0.6	+0.4
Hispanic	3.1	2.3	2.4	2.5	2.0	1.9	2.0	1.8	1.0	0.8	0.9	0.6	1.0	1.0	§	0.5	-0.5

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 45
CRACK: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	—	—	—	—	—	—	—	—	—	—	—	15,200	16,300	16,300	16,700	15,200
Total	—	—	—	—	—	—	—	—	—	—	—	4.1	3.9	3.1	3.1	1.9
Gender:																
Male	—	—	—	—	—	—	—	—	—	—	—	4.2	4.6	4.0	4.3	2.3
Female	—	—	—	—	—	—	—	—	—	—	—	3.6	3.0	2.0	1.8	1.4
College Plans:																
None or under 4 years	—	—	—	—	—	—	—	—	—	—	—	5.2	5.1	4.1	3.8	3.5
Complete 4 years	—	—	—	—	—	—	—	—	—	—	—	2.8	2.7	2.3	2.7	1.2
Region:																
Northeast	—	—	—	—	—	—	—	—	—	—	—	6.0	4.0	2.3	3.3	2.0
Midwest	—	—	—	—	—	—	—	—	—	—	—	3.1	3.5	2.4	2.2	1.6
South	—	—	—	—	—	—	—	—	—	—	—	1.6	2.8	2.6	3.3	1.8
West	—	—	—	—	—	—	—	—	—	—	—	7.5	6.1	5.6	3.8	2.7
Population Density:																
Large MSA	—	—	—	—	—	—	—	—	—	—	—	5.9	4.7	3.9	3.4	1.6
Other MSA	—	—	—	—	—	—	—	—	—	—	—	3.5	3.5	3.2	3.3	2.0
Non-MSA	—	—	—	—	—	—	—	—	—	—	—	3.5	3.7	2.0	2.2	2.0
Parental Education: ^a																
1.0–2.0 (Low)	—	—	—	—	—	—	—	—	—	—	—	1.2	3.6	3.3	3.1	2.2
2.5–3.0	—	—	—	—	—	—	—	—	—	—	—	5.3	4.2	2.6	3.1	2.2
3.5–4.0	—	—	—	—	—	—	—	—	—	—	—	4.0	4.0	3.4	2.8	1.8
4.5–5.0	—	—	—	—	—	—	—	—	—	—	—	2.9	3.4	3.1	2.6	1.1
5.5–6.0 (High)	—	—	—	—	—	—	—	—	—	—	—	3.7	2.4	2.1	3.7	1.8
Race/Ethnicity (2-year average): ^b																
White	—	—	—	—	—	—	—	—	—	—	—	—	3.8	3.4	3.1	2.1
African American	—	—	—	—	—	—	—	—	—	—	—	—	1.9	2.5	2.0	1.3
Hispanic	—	—	—	—	—	—	—	—	—	—	—	—	5.5	3.7	3.2	4.2

Table continued on next page.

TABLE 45 (cont.)
CRACK: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700
Total	1.5	1.5	1.5	1.9	2.1	2.1	2.4	2.5	2.7	2.2	2.1	2.3	2.2	2.3	1.9
Gender:															
Male	1.8	1.7	1.9	2.4	2.5	2.6	3.0	3.1	2.9	2.5	2.4	2.6	2.3	2.5	2.2
Female	1.0	1.0	1.1	1.3	1.5	1.6	1.8	2.0	2.2	1.7	1.8	1.8	1.9	1.9	1.6
College Plans:															
None or under 4 years	2.3	2.6	2.7	3.3	3.0	4.0	4.3	4.6	5.0	3.5	4.4	4.5	2.7	3.9	3.4
Complete 4 years	1.1	1.0	1.2	1.4	1.7	1.6	1.7	1.9	1.9	1.7	1.5	1.7	1.8	1.7	1.6
Region:															
Northeast	1.3	1.3	1.2	1.5	1.6	2.1	2.6	3.0	2.4	1.8	2.0	2.0	2.0	1.7	1.5
Midwest	1.5	1.4	1.3	2.2	2.0	2.3	2.2	2.6	2.8	2.0	2.6	2.6	1.9	2.1	2.3
South	1.2	1.2	1.5	1.6	1.7	1.7	1.8	2.0	2.3	1.8	1.2	1.9	2.0	2.2	1.8
West	1.8	2.1	2.1	2.3	3.5	2.6	3.4	3.2	3.4	3.2	3.0	2.8	3.0	3.3	2.3
Population Density:															
Large MSA	1.2	1.3	1.3	1.5	2.0	2.1	2.2	2.5	2.0	1.8	1.5	1.6	1.8	2.0	1.7
Other MSA	1.7	1.6	1.8	2.1	2.1	1.9	2.3	2.4	2.5	2.1	2.2	2.4	2.3	2.2	1.8
Non-MSA	1.2	1.3	1.4	1.9	2.1	2.5	2.8	2.9	3.6	2.8	2.8	2.8	2.4	2.8	2.5
Parental Education: ^a															
1.0–2.0 (Low)	1.6	1.9	2.6	2.7	3.4	3.4	3.6	3.9	3.2	2.8	3.8	4.8	2.5	3.3	2.3
2.5–3.0	1.5	1.9	1.6	2.2	2.3	2.4	2.5	2.8	2.8	2.1	2.6	2.2	1.6	2.9	2.4
3.5–4.0	1.7	1.3	1.5	1.8	1.7	2.2	2.0	2.4	3.0	2.2	1.8	2.3	2.3	1.9	2.0
4.5–5.0	0.9	1.0	1.4	1.1	1.9	1.6	2.5	2.1	2.2	1.8	1.9	1.7	2.2	1.8	1.5
5.5–6.0 (High)	1.1	0.8	1.0	1.8	1.5	1.6	1.8	2.1	1.6	2.2	1.2	1.5	1.9	1.7	1.5
Race/Ethnicity (2-year average): ^b															
White	1.6	1.3	1.3	1.6	1.9	2.0	2.2	2.6	2.8	2.5	2.2	2.3	2.2	2.2	2.1
African American	1.0	0.6	0.6	0.9	1.0	0.7	0.5	0.3	0.4	0.5	0.5	0.7	1.2	1.3	0.9
Hispanic	3.4	2.7	2.5	2.4	3.1	4.1	4.2	3.9	3.5	3.4	3.1	3.1	2.9	3.0	3.1

Table continued on next page.

TABLE 45 (cont.)
CRACK: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																2019-2021 change	
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Approximate weighted N =	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	2.1	1.9	1.6	1.3	1.4	1.0	1.2	1.1	1.1	1.1	0.8	1.0	0.9	1.0	1.2	0.7	-0.3*	
Gender:																		
Male	2.1	2.1	1.7	1.6	1.9	1.2	1.4	1.1	1.4	1.3	1.0	1.2	1.4	1.4	§	0.6	-0.8 ss	
Female	1.9	1.7	1.3	1.0	0.9	0.8	0.9	0.9	0.6	0.6	0.5	0.8	0.4	0.7	§	0.8	+0.1	
College Plans:																		
None or under 4 years	3.5	3.7	3.3	2.8	2.5	2.4	1.9	2.1	1.8	2.2	1.2	2.1	1.9	2.2	§	1.7	-0.5	
Complete 4 years	1.7	1.5	1.1	0.9	1.2	0.7	1.0	0.7	0.9	0.7	0.7	0.7	0.6	0.7	§	0.4	-0.3	
Region:																		
Northeast	2.5	2.1	1.8	0.6	1.4	1.2	0.8	1.1	0.8	0.7	0.7	0.9	0.7	0.5	§	0.2	-0.3	
Midwest	1.5	1.7	1.3	1.7	1.1	1.2	1.0	1.0	0.9	1.3	0.6	0.7	1.2	0.9	§	0.5	-0.4	
South	2.1	2.2	1.6	1.2	1.4	0.8	1.1	0.8	1.0	0.9	0.5	1.1	0.9	1.1	§	1.0	-0.1	
West	2.2	1.6	1.7	1.6	1.8	1.2	1.9	1.6	1.6	1.4	1.6	1.2	0.9	1.6	§	0.9	-0.7	
Population Density:																		
Large MSA	2.6	1.9	1.3	1.1	1.4	1.0	1.2	0.9	0.7	1.0	1.1	0.7	0.9	1.0	§	0.4	-0.5	
Other MSA	1.7	1.9	1.6	1.3	1.6	1.1	1.3	1.2	1.5	1.0	0.7	1.2	1.1	1.0	§	0.8	-0.2	
Non-MSA	2.0	2.0	1.6	1.6	1.0	0.9	1.0	0.9	0.6	1.2	0.6	1.0	0.7	1.2	§	1.1	-0.1	
Parental Education: ^a																		
1.0–2.0 (Low)	3.6	3.6	2.1	2.2	2.5	2.3	1.9	1.5	1.8	1.8	1.7	1.6	1.7	1.6	§	0.7	-0.9	
2.5–3.0	2.4	2.2	1.4	0.9	1.7	0.8	1.3	1.0	1.2	1.2	1.0	1.1	1.0	1.1	§	0.7	-0.3	
3.5–4.0	2.0	1.9	1.6	1.4	1.4	1.2	1.0	1.2	0.7	1.0	0.3	1.0	0.8	1.0	§	1.2	+0.2	
4.5–5.0	1.8	1.7	1.4	1.4	1.0	0.8	1.1	0.8	0.8	0.7	0.8	0.5	0.7	0.6	§	0.6	0.0	
5.5–6.0 (High)	1.2	1.0	1.3	0.6	0.8	0.8	1.1	0.8	1.4	0.8	0.6	0.6	0.5	1.0	§	0.1	-0.8 s	
Race/Ethnicity (2-year average): ^b																		
White	1.9	2.0	1.8	1.4	1.2	1.1	1.1	1.0	0.8	0.8	0.7	0.8	0.9	0.8	§	0.5	-0.2	
African American	1.0	1.0	0.9	0.9	0.7	0.7	0.7	0.9	1.3	1.3	0.9	0.8	0.7	0.9	§	0.8	0.0	
Hispanic	3.1	2.7	2.0	1.6	1.8	1.9	1.7	1.7	1.7	1.5	1.3	1.1	1.2	1.3	§	1.2	-0.1	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1986; N is one fifth of N indicated in Table S3. Data based on two forms in 1987–1989; N is two fifths of N indicated in 1987–1988 and two sixths of N indicated in 1989 in Table S3. Data based on six forms beginning in 1990.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 46
OTHER COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	
Total	1.0	1.2	1.3	1.7	2.1	2.5	2.2	2.4	2.3	1.9	1.9	1.8	1.6	1.6	1.7	
Gender:																
Male	1.1	1.2	1.5	1.7	2.0	2.2	2.5	2.3	2.3	1.9	1.8	1.7	1.5	1.3	1.5	
Female	0.8	1.2	1.2	1.8	2.2	2.6	1.9	2.4	2.2	1.8	2.0	1.9	1.6	1.8	1.8	
College Plans:																
None or under 4 years	2.7	4.2	4.1	5.6	5.9	6.6	6.0	7.7	7.1	6.1	5.3	6.4	5.0	5.4	4.5	
Complete 4 years	0.6	0.7	0.9	1.2	1.6	1.8	1.7	1.8	1.7	1.4	1.5	1.3	1.2	1.1	1.4	
Region:																
Northeast	1.2	0.7	0.9	1.9	1.8	2.1	1.6	1.4	1.7	1.1	1.6	1.2	1.0	1.6	1.5	
Midwest	0.6	1.0	0.7	0.9	2.0	2.4	2.0	1.9	2.2	2.3	1.6	1.8	1.7	1.1	1.3	
South	1.0	1.5	1.6	2.0	2.0	2.3	2.1	3.1	2.6	1.7	2.4	2.1	1.9	1.7	1.9	
West	1.3	1.5	2.1	2.0	2.7	3.1	2.9	2.5	2.2	2.5	1.7	1.9	1.6	1.8	1.8	
Population Density:																
Large MSA	0.9	1.1	1.0	1.6	2.0	2.5	1.7	1.8	1.8	1.6	1.6	1.4	1.2	1.7	1.6	
Other MSA	0.9	1.4	1.8	2.1	2.1	2.6	2.2	2.5	2.3	1.8	2.2	1.8	1.6	1.6	1.7	
Non-MSA	1.1	0.9	0.7	1.2	2.2	2.2	2.5	2.8	2.8	2.5	1.6	2.3	2.2	1.3	1.7	
Parental Education: ^a																
1.0–2.0 (Low)	2.1	2.7	2.2	3.1	4.3	3.2	3.5	4.7	5.1	4.2	3.7	2.8	2.9	3.1	2.9	
2.5–3.0	1.2	1.1	1.5	2.0	2.0	2.6	2.4	2.5	2.8	2.1	1.9	2.3	1.8	1.9	2.2	
3.5–4.0	0.6	1.0	1.5	1.9	2.2	2.8	2.1	2.4	2.3	1.9	1.8	1.9	1.6	1.5	2.0	
4.5–5.0	0.6	0.8	0.8	1.1	1.6	2.4	1.9	1.8	1.3	1.4	1.6	1.6	1.1	1.3	1.2	
5.5–6.0 (High)	1.0	1.2	0.8	1.2	2.0	1.9	1.7	1.8	1.8	1.3	1.1	0.8	1.5	0.9	0.9	
Race/Ethnicity (2-year average): ^b																
White	—	0.9	1.0	1.2	1.8	2.4	2.5	2.2	2.1	1.9	1.8	1.8	1.8	1.5	1.4	
African American	—	0.6	0.5	0.6	0.5	0.4	0.3	0.5	0.7	0.6	0.5	0.5	0.6	0.8	0.9	
Hispanic	—	2.6	3.3	4.0	4.3	4.1	3.3	4.0	4.9	3.9	3.2	3.1	2.7	2.8	2.9	

Table continued on next page.

TABLE 46 (cont.)
OTHER COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021
Approximate weighted N =	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	1.6	1.5	1.4	1.3	1.3	1.1	1.0	0.8	0.8	0.8	0.6	0.6	0.7	0.6	0.5	0.2	-0.4* sss
Gender:																	
Male	1.4	1.3	1.4	1.3	1.3	1.1	0.9	0.8	0.8	0.9	0.6	0.5	0.7	0.4	§	0.1	-0.3 ss
Female	1.7	1.6	1.4	1.3	1.4	0.9	1.0	0.8	0.7	0.6	0.6	0.7	0.5	0.7	§	0.1	-0.7 sss
College Plans:																	
None or under 4 years	5.7	5.7	5.5	5.6	4.7	4.6	3.8	3.0	2.6	2.0	2.0	2.1	2.1	1.0	§	0.4	-0.7 s
Complete 4 years	1.2	1.1	1.1	1.0	1.1	0.8	0.7	0.6	0.6	0.7	0.4	0.4	0.4	0.5	§	0.1	-0.4 sss
Region:																	
Northeast	1.0	1.0	0.9	1.2	0.9	0.7	0.8	0.5	0.4	0.4	0.5	0.5	0.4	0.1	§	0.0	-0.1
Midwest	1.1	1.3	1.5	1.5	1.6	1.0	0.5	0.6	0.6	0.6	0.3	0.5	0.4	0.3	§	0.1	-0.2
South	2.3	1.8	1.8	1.0	1.3	1.1	1.2	0.8	0.9	0.9	0.8	0.5	0.5	0.7	§	0.2	-0.4 s
West	1.5	1.5	1.3	1.8	1.4	1.6	1.2	1.1	1.1	0.9	0.8	1.0	1.3	0.9	§	0.2	-0.7 ss
Population Density:																	
Large MSA	1.5	1.5	1.5	0.9	1.1	1.2	1.1	0.8	1.1	0.8	0.5	0.6	0.7	0.5	§	0.1	-0.4 ss
Other MSA	1.6	1.5	1.4	1.6	1.5	1.2	1.0	0.9	0.7	0.9	0.6	0.7	0.7	0.7	§	0.3	-0.4 s
Non-MSA	1.7	1.5	1.4	1.4	1.3	0.8	0.8	0.7	0.7	0.3	0.8	0.4	0.6	0.4	§	0.1	-0.3 s
Parental Education: ^a																	
1.0–2.0 (Low)	3.6	3.1	3.0	3.4	2.8	2.1	2.1	1.8	1.4	1.0	1.4	1.2	1.2	1.4	§	0.4	-1.0
2.5–3.0	1.7	1.6	1.8	1.4	1.9	1.6	1.2	1.0	0.7	0.8	0.5	0.9	0.7	0.8	§	0.2	-0.6 s
3.5–4.0	1.7	1.4	1.7	1.2	1.3	1.3	0.9	0.8	0.8	0.8	0.6	0.7	0.6	0.5	§	0.3	-0.3
4.5–5.0	1.2	1.0	0.8	0.6	1.0	0.7	0.4	0.5	0.8	0.4	0.4	0.3	0.6	0.3	§	0.1	-0.2
5.5–6.0 (High)	0.9	0.9	1.1	1.3	0.9	0.6	0.6	0.7	0.4	0.8	0.5	0.4	0.4	0.4	§	0.1	-0.3
Race/Ethnicity (2-year average): ^b																	
White	1.5	1.3	1.2	1.2	1.2	1.0	0.8	0.7	0.5	0.5	0.5	0.5	0.4	0.4	§	0.3	-0.1
African American	0.7	0.6	0.6	0.8	0.7	0.4	0.4	0.6	0.6	0.5	0.4	0.3	0.4	0.4	§	0.0	-0.4
Hispanic	2.8	3.1	2.8	2.2	2.2	2.4	2.2	1.5	1.1	1.0	0.9	0.8	0.9	1.0	0.1		-0.8 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 47
OTHER COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
Total	2.1	1.7	1.8	2.4	3.0	3.5	4.1	4.0	4.4	3.8	3.0	3.4	2.8	3.3	3.0	
Gender:																
Male	2.0	1.9	2.2	2.7	3.1	3.7	4.1	4.1	4.6	4.2	3.3	3.6	2.9	3.6	3.3	
Female	2.1	1.5	1.4	2.1	2.9	3.3	4.0	3.8	4.1	3.4	2.6	3.3	2.7	3.0	2.7	
College Plans:																
None or under 4 years	4.4	3.3	4.5	5.9	6.3	8.4	9.0	9.3	10.5	8.3	8.2	8.7	7.3	8.3	8.1	
Complete 4 years	1.6	1.3	1.3	1.7	2.5	2.7	3.2	3.0	3.4	3.0	2.1	2.6	2.1	2.6	2.3	
Region:																
Northeast	1.3	1.0	1.8	2.0	2.2	2.2	2.5	4.3	4.1	2.5	2.0	1.8	2.7	3.2	3.0	
Midwest	1.6	1.3	1.3	1.8	2.5	3.4	3.4	3.1	3.9	4.0	3.0	3.2	2.3	3.0	3.0	
South	1.9	1.6	1.7	2.2	2.9	3.5	4.8	3.7	4.7	3.9	3.2	3.3	3.1	3.2	2.6	
West	3.4	3.1	3.2	4.3	4.8	5.2	5.3	5.2	4.6	4.6	3.9	5.5	3.2	4.1	3.6	
Population Density:																
Large MSA	1.6	1.5	1.4	1.9	2.8	3.3	3.9	3.8	3.7	3.7	2.5	3.1	2.2	2.9	2.7	
Other MSA	2.6	2.0	2.0	2.7	3.1	3.9	3.8	4.0	4.5	3.7	3.2	3.7	2.8	3.4	3.1	
Non-MSA	1.4	1.4	1.9	2.5	3.1	3.2	4.9	4.2	4.8	4.0	3.2	3.3	4.1	3.6	3.3	
Parental Education: ^a																
1.0–2.0 (Low)	3.1	2.7	2.7	3.1	5.0	6.1	5.5	6.7	7.6	6.3	4.8	7.1	6.4	5.7	4.5	
2.5–3.0	2.2	1.6	2.0	2.6	3.6	3.6	4.3	4.5	5.1	4.4	4.5	3.9	3.5	3.7	4.0	
3.5–4.0	2.2	2.0	2.2	2.7	3.3	3.8	4.6	4.0	4.2	4.0	2.5	3.2	2.9	3.7	3.0	
4.5–5.0	1.6	1.3	1.4	1.8	2.2	3.0	3.3	3.0	3.5	3.0	2.2	2.0	1.8	2.8	2.4	
5.5–6.0 (High)	1.8	1.3	0.9	1.6	1.7	2.8	2.9	2.5	3.5	2.7	1.4	2.7	1.2	1.6	2.1	
Race/Ethnicity (2-year average): ^b																
White	—	1.9	1.8	1.9	2.6	3.2	3.7	4.1	4.3	4.1	3.4	3.3	3.4	3.4	3.3	
African American	—	0.5	0.5	0.9	0.8	0.6	0.6	0.9	0.8	0.5	0.5	0.7	0.7	0.7	0.8	
Hispanic	—	3.4	3.4	4.6	5.2	6.1	7.5	7.0	6.8	7.1	5.6	5.0	5.2	5.1	5.4	

Table continued on next page.

TABLE 47 (cont.)
OTHER COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	2.9	3.1	2.6	2.3	1.9	1.7	1.8	1.6	1.3	1.5	1.1	1.2	1.4	1.4	1.0	0.5	-0.8* ss
Gender:																	
Male	3.0	3.2	2.7	2.5	2.3	2.1	2.1	2.2	1.7	1.5	1.4	1.5	1.7	1.4	§	0.5	-0.9 s
Female	2.8	2.9	2.4	2.1	1.3	1.2	1.5	1.0	0.9	1.4	0.9	0.9	1.2	1.3	§	0.3	-0.9 ss
College Plans:																	
None or under 4 years	8.3	7.5	7.8	7.2	6.8	5.1	6.2	3.9	4.5	3.1	3.6	2.9	3.6	2.2	§	1.0	-1.3 s
Complete 4 years	2.2	2.5	1.8	1.8	1.3	1.3	1.4	1.3	0.9	1.3	0.9	1.0	1.1	1.2	§	0.3	-0.8 ss
Region:																	
Northeast	2.5	2.5	1.7	2.1	1.7	1.7	1.1	1.1	1.0	1.4	0.7	0.6	0.9	0.5	§	0.8	+0.3
Midwest	2.8	3.3	2.2	2.1	1.5	1.2	1.2	0.8	1.1	1.2	1.0	1.4	1.0	1.0	§	0.4	-0.7 s
South	2.9	3.3	3.2	2.1	1.5	1.8	2.2	1.8	1.5	1.5	1.2	1.1	2.0	1.8	§	0.6	-1.1
West	3.5	3.0	2.7	3.1	3.0	2.1	2.5	2.4	1.6	2.0	1.5	1.5	1.3	1.8	§	0.4	-1.4 s
Population Density:																	
Large MSA	2.6	3.1	2.4	2.2	1.9	1.8	2.0	1.5	1.0	1.4	1.1	1.7	1.2	0.9	§	0.5	-0.4
Other MSA	3.0	2.9	2.9	2.6	2.0	1.7	1.6	1.9	1.3	1.5	1.1	1.0	1.8	1.8	§	0.6	-1.2 s
Non-MSA	3.4	3.3	2.0	1.9	1.7	1.5	2.0	0.9	1.9	1.8	1.3	0.8	0.9	1.3	§	0.6	-0.7 s
Parental Education: ^a																	
1.0–2.0 (Low)	5.6	5.1	5.9	5.1	3.6	3.1	5.3	2.5	2.6	2.5	1.2	1.7	2.3	2.2	§	1.3	-0.9
2.5–3.0	3.7	3.3	3.1	2.7	2.3	1.9	2.2	2.3	1.3	1.9	1.8	1.5	1.8	1.8	§	0.9	-0.9
3.5–4.0	2.8	3.2	2.4	2.5	1.7	1.7	1.5	1.4	1.5	1.7	0.9	1.0	1.6	1.6	§	0.5	-1.1
4.5–5.0	1.8	2.4	1.9	1.4	1.6	1.3	1.3	1.1	0.8	1.0	1.0	0.9	0.9	0.8	§	0.1	-0.7 ss
5.5–6.0 (High)	2.8	2.4	1.3	2.0	1.0	1.0	1.0	1.2	0.9	1.1	0.7	1.1	1.4	0.9	§	0.4	-0.5
Race/Ethnicity (2-year average): ^b																	
White	3.2	3.2	2.8	2.3	1.9	1.6	1.5	1.4	1.3	1.3	1.3	1.2	1.2	1.2	§	0.5	-0.7 ss
African American	0.9	0.6	0.7	0.7	0.6	0.6	1.2	1.4	0.7	0.6	0.5	0.6	0.6	0.3	§	0.5	+0.2
Hispanic	5.0	4.5	4.6	4.5	4.0	3.7	3.4	3.4	2.6	1.9	1.5	1.2	2.1	2.5	§	1.4	-1.2

Source: The Monitoring the Future study, the University of Michigan.

Notes: Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 48
OTHER COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

		Percentage who used in last 12 months															
		1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Approximate weighted N =		—	—	—	—	—	—	—	—	—	—	—	—	16,300	16,300	16,700	15,200
Total		—	—	—	—	—	—	—	—	—	—	—	—	9.8	7.4	5.2	4.6
Gender:																	
Male		—	—	—	—	—	—	—	—	—	—	—	—	10.1	8.0	6.5	5.8
Female		—	—	—	—	—	—	—	—	—	—	—	—	9.1	6.2	4.0	3.2
College Plans:																	
None or under 4 years		—	—	—	—	—	—	—	—	—	—	—	—	9.8	6.0	7.3	6.3
Complete 4 years		—	—	—	—	—	—	—	—	—	—	—	—	8.3	6.7	4.2	3.7
Region:																	
Northeast		—	—	—	—	—	—	—	—	—	—	—	—	12.9	7.0	4.9	5.6
Midwest		—	—	—	—	—	—	—	—	—	—	—	—	8.2	5.6	4.8	3.7
South		—	—	—	—	—	—	—	—	—	—	—	—	5.8	5.8	4.6	4.1
West		—	—	—	—	—	—	—	—	—	—	—	—	15.3	13.4	7.5	6.1
Population Density:																	
Large MSA		—	—	—	—	—	—	—	—	—	—	—	—	13.3	9.8	5.6	5.0
Other MSA		—	—	—	—	—	—	—	—	—	—	—	—	8.9	7.8	5.4	4.7
Non-MSA		—	—	—	—	—	—	—	—	—	—	—	—	8.0	4.5	4.4	4.1
Parental Education: ^a																	
1.0–2.0 (Low)		—	—	—	—	—	—	—	—	—	—	—	—	5.3	4.9	3.3	3.4
2.5–3.0		—	—	—	—	—	—	—	—	—	—	—	—	10.5	6.5	4.6	5.0
3.5–4.0		—	—	—	—	—	—	—	—	—	—	—	—	10.5	7.2	5.1	4.7
4.5–5.0		—	—	—	—	—	—	—	—	—	—	—	—	9.0	7.7	6.1	4.1
5.5–6.0 (High)		—	—	—	—	—	—	—	—	—	—	—	—	9.7	9.0	6.5	5.4
Race/Ethnicity (2-year average): ^b																	
White		—	—	—	—	—	—	—	—	—	—	—	—	—	9.3	7.0	5.3
African American		—	—	—	—	—	—	—	—	—	—	—	—	—	2.8	1.4	0.7
Hispanic		—	—	—	—	—	—	—	—	—	—	—	—	—	6.3	5.1	5.1

Table continued on next page.

TABLE 48 (cont.)
OTHER COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	3.2	2.6	2.9	3.0	3.4	4.2	5.0	4.9	5.8	4.5	4.4	4.4	4.2	4.7	4.5	5.2
Gender:																
Male	3.7	3.1	3.7	3.7	4.0	4.9	5.7	5.6	7.1	5.5	4.9	5.0	5.4	5.8	5.4	5.7
Female	2.4	2.0	2.0	2.3	2.5	3.2	4.0	3.9	4.2	3.4	3.7	3.7	2.9	3.5	3.6	4.4
College Plans:																
None or under 4 years	4.0	4.0	3.9	4.3	4.5	5.7	7.0	8.9	7.6	6.7	7.4	7.1	4.8	6.8	6.7	6.9
Complete 4 years	2.8	2.0	2.5	2.5	2.9	3.5	4.1	3.5	5.2	3.7	3.4	3.4	3.8	3.9	3.9	4.7
Region:																
Northeast	3.4	2.8	2.3	2.8	4.2	5.2	5.9	4.7	4.1	3.7	4.4	4.3	5.3	4.4	4.3	6.2
Midwest	2.9	2.2	2.3	3.5	2.7	3.2	4.1	5.3	5.7	4.5	5.6	4.8	3.2	3.9	4.2	3.6
South	2.8	2.5	2.6	2.6	3.1	4.2	4.6	4.9	6.6	4.0	3.9	4.4	4.1	4.9	4.7	5.8
West	3.9	3.1	4.6	3.5	4.0	4.5	6.2	4.4	6.1	5.9	3.5	3.9	4.7	5.8	4.7	5.1
Population Density:																
Large MSA	3.7	3.1	2.6	2.6	3.7	3.9	4.2	4.8	4.8	3.7	4.1	3.5	3.0	3.9	4.6	7.2
Other MSA	3.3	2.5	3.6	3.5	3.3	4.4	5.2	4.9	6.0	4.5	4.3	4.7	5.3	5.4	4.7	4.4
Non-MSA	2.5	2.3	2.0	2.6	3.1	4.2	5.6	4.9	6.7	5.4	5.0	4.9	3.9	4.2	4.0	3.8
Parental Education: ^a																
1.0–2.0 (Low)	3.5	3.7	3.9	2.7	2.9	5.2	6.4	5.5	7.4	5.4	3.7	5.1	4.1	6.3	4.1	4.6
2.5–3.0	3.5	2.3	2.3	3.2	3.4	3.8	4.9	5.3	5.0	3.8	5.6	4.7	3.4	3.6	5.5	5.5
3.5–4.0	3.2	2.6	3.3	3.4	3.6	4.6	4.9	5.3	6.9	4.7	4.7	4.9	4.6	5.1	4.7	5.2
4.5–5.0	2.7	2.3	2.9	2.6	3.2	3.9	4.4	4.2	5.5	4.5	3.9	3.4	4.3	5.1	4.2	4.6
5.5–6.0 (High)	2.4	2.0	1.7	3.1	2.7	3.8	4.8	3.5	3.6	4.2	2.2	3.9	3.8	3.7	3.7	5.9
Race/Ethnicity (2-year average): ^b																
White	4.2	2.9	2.6	2.9	3.3	3.9	5.0	5.6	6.0	5.7	5.0	5.1	4.9	4.8	4.9	5.2
African American	1.0	1.0	0.7	0.8	0.8	0.7	0.7	0.6	0.8	1.0	0.9	0.8	1.0	1.1	1.2	1.4
Hispanic	5.0	4.3	5.1	5.1	4.0	5.6	6.9	6.0	5.8	6.6	5.8	4.7	3.9	4.8	5.8	5.7

Table continued on next page.

TABLE 48 (cont.)
OTHER COCAINE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	4.5	4.0	3.0	2.6	2.6	2.4	2.4	2.4	2.1	2.0	2.3	2.0	1.9	2.9	0.9	-0.9* s
Gender:																
Male	4.8	4.8	3.6	3.6	3.4	2.8	3.1	3.6	2.7	2.5	3.4	2.2	2.2	§	0.6	-1.6 sss
Female	3.9	3.1	2.3	1.7	1.9	1.8	1.6	1.3	1.6	1.4	1.4	1.2	1.3	§	1.3	0.0
College Plans:																
None or under 4 years	7.7	6.8	4.7	5.3	4.3	3.2	4.4	3.3	3.8	3.3	4.3	3.3	3.5	§	1.9	-1.5
Complete 4 years	3.6	3.1	2.5	2.1	2.2	2.0	1.8	2.1	1.7	1.7	1.8	1.4	1.2	§	0.6	-0.7 s
Region:																
Northeast	5.3	4.4	2.7	3.3	2.9	1.8	2.4	2.2	1.9	1.0	2.2	2.1	1.2	§	0.8	-0.5
Midwest	4.4	3.4	3.5	2.2	2.6	1.7	1.7	2.0	1.9	1.6	1.7	1.8	1.5	§	0.4	-1.0 s
South	4.5	4.2	2.7	1.9	2.4	2.5	2.1	2.5	1.6	1.7	2.0	1.4	1.8	§	0.2	-1.6 sss
West	3.8	4.1	2.9	3.6	2.9	3.3	3.5	2.7	3.5	3.7	3.5	2.9	3.4	§	2.6	-0.8
Population Density:																
Large MSA	4.3	4.1	2.8	2.8	2.6	2.6	2.1	2.0	2.4	2.2	2.6	2.0	1.6	§	0.4	-1.2 sss
Other MSA	4.6	3.9	3.0	3.0	2.7	2.6	2.7	2.9	2.0	2.1	2.3	1.8	1.8	§	1.4	-0.4
Non-MSA	4.6	4.2	3.2	1.5	2.5	1.5	1.9	1.9	1.9	1.8	1.9	2.3	2.4	§	0.6	-1.8 sss
Parental Education: ^a																
1.0–2.0 (Low)	5.2	3.0	3.8	3.0	3.2	3.5	3.3	1.7	3.1	2.7	2.7	2.4	1.9	§	0.9	-1.0
2.5–3.0	5.0	3.8	3.2	2.6	2.6	2.4	3.1	3.1	2.3	1.9	2.7	2.5	2.1	§	1.5	-0.5
3.5–4.0	4.5	4.6	2.5	2.9	2.9	2.4	2.7	2.3	2.1	1.9	2.2	1.3	1.6	§	1.4	-0.3
4.5–5.0	4.3	3.5	2.9	2.1	2.2	1.9	1.4	2.2	1.8	2.0	1.7	1.6	1.8	§	0.5	-1.3 ss
5.5–6.0 (High)	3.7	4.2	2.6	2.6	2.5	2.2	1.9	2.6	1.9	2.0	2.4	1.6	1.2	§	0.3	-0.9 s
Race/Ethnicity (2-year average): ^b																
White	5.5	5.2	4.3	3.2	3.0	2.8	2.3	2.0	2.0	2.1	2.3	2.3	1.9	§	1.6	-0.3
African American	1.2	0.7	0.6	0.7	0.7	0.8	1.0	1.5	1.4	0.8	0.6	0.4	0.7	§	0.9	+0.2
Hispanic	4.8	3.7	3.1	2.5	2.5	3.1	3.6	3.4	3.0	2.9	2.7	2.4	2.1	§	1.6	-0.5

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one form in 1987–1989; *N* is one fifth of *N* indicated in 1987–1988 and one sixth of *N* indicated in 1989 in Table S3. Data based on four of six forms beginning in 1990; *N* is four sixths of *N* indicated in Table S3.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 49
HEROIN: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995 ^a	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	
Total	0.7	0.7	0.7	1.2	1.4	1.6	1.3	1.3	1.4	1.1	1.0	0.9	0.9	1.0	0.8	
Gender:																
Male	0.9	0.8	0.8	1.3	1.6	1.5	1.4	1.5	1.4	1.0	1.0	0.8	0.8	0.9	0.8	
Female	0.5	0.7	0.5	0.9	1.2	1.5	1.1	1.1	1.3	1.2	1.0	1.0	0.9	1.0	0.8	
College Plans:																
None or under 4 years	2.1	2.7	2.0	3.9	4.4	4.1	3.4	5.0	3.7	3.5	3.1	2.9	3.1	4.2	2.3	
Complete 4 years	0.4	0.4	0.5	0.7	1.0	1.1	1.1	0.9	1.1	0.8	0.8	0.7	0.6	0.6	0.6	
Region:																
Northeast	0.5	0.6	0.7	1.3	1.4	1.8	1.1	1.1	1.3	1.1	1.1	0.8	0.7	1.1	0.7	
Midwest	0.4	0.8	0.5	1.1	1.4	1.6	1.4	1.3	1.6	1.4	1.0	1.0	1.0	0.8	0.5	
South	0.8	0.7	0.7	1.1	1.5	1.4	1.2	1.4	1.4	0.7	1.2	1.0	1.0	0.9	1.0	
West	1.0	0.7	1.1	1.1	1.2	1.6	1.4	1.3	1.2	1.4	0.7	1.0	0.9	1.2	0.8	
Population Density:																
Large MSA	0.5	0.7	0.7	1.2	1.2	1.4	1.0	1.0	1.3	1.0	0.9	0.7	0.8	1.2	0.6	
Other MSA	0.7	0.8	0.9	1.2	1.5	1.7	1.3	1.3	1.4	1.1	1.0	1.0	0.9	1.0	0.8	
Non-MSA	0.8	0.7	0.4	1.0	1.5	1.5	1.5	1.6	1.5	1.0	1.1	1.1	1.2	0.6	1.0	
Parental Education: ^b																
1.0–2.0 (Low)	1.5	1.4	0.8	2.0	2.6	2.0	2.3	3.5	2.8	1.7	2.6	1.2	2.1	1.9	0.5	
2.5–3.0	0.9	0.7	0.6	1.1	1.0	1.7	1.4	1.2	1.4	1.5	1.1	1.3	1.0	1.1	1.3	
3.5–4.0	0.6	0.6	0.7	1.3	1.6	1.7	1.0	1.1	1.2	0.7	1.0	0.9	0.8	1.0	1.0	
4.5–5.0	0.4	0.5	0.8	0.8	1.2	1.4	1.1	1.1	1.1	0.8	0.6	0.8	0.6	0.7	0.5	
5.5–6.0 (High)	0.5	0.8	0.6	1.3	1.6	1.0	1.5	1.4	1.8	1.0	0.8	0.6	0.7	0.6	0.4	
Race/Ethnicity (2-year average): ^c																
White	—	0.6	0.6	0.8	1.2	1.6	1.6	1.3	1.2	1.2	1.0	1.0	0.9	0.9	0.8	
African American	—	0.4	0.3	0.6	0.7	0.5	0.4	0.5	0.7	0.5	0.6	0.6	0.5	0.7	0.5	
Hispanic	—	1.4	1.4	1.5	1.8	2.1	1.7	1.7	2.2	2.0	1.4	1.2	1.4	1.7	1.5	

Table continued on next page.

TABLE 49
HEROIN: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015 ^a	2016 ^a	2017 ^a	2018 ^a	2019 ^a	2020 ^a	2021 ^a	
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	0.8	0.8	0.9	0.7	0.8	0.7	0.5	0.5	0.5	0.3	0.3	0.3	0.3	0.3	0.2	0.2	-0.2*
Gender:																	
Male	0.6	0.8	0.9	0.8	0.9	0.7	0.5	0.4	0.6	0.2	0.2	0.2	0.4	0.3	§	0.2	-0.2
Female	0.9	0.9	0.7	0.7	0.7	0.7	0.5	0.5	0.4	0.3	0.2	0.4	0.2	0.4	§	0.1	-0.2
College Plans:																	
None or under 4 years	3.0	3.7	3.5	3.5	2.3	3.1	1.8	2.1	2.2	0.6	0.9	1.7	0.9	0.7	§	0.6	-0.1
Complete 4 years	0.6	0.5	0.6	0.5	0.7	0.5	0.4	0.4	0.3	0.2	0.1	0.2	0.2	0.3	§	0.1	-0.2 s
Region:																	
Northeast	0.4	0.5	0.7	0.4	0.6	0.6	0.4	0.3	0.3	0.1	0.2	0.2	0.2	0.2	§	0.0	-0.2
Midwest	0.7	0.8	1.0	0.7	0.8	0.8	0.4	0.4	0.5	0.3	0.1	0.3	0.1	0.3	§	0.1	-0.2
South	0.9	0.8	0.8	0.6	0.8	0.6	0.6	0.6	0.4	0.4	0.4	0.3	0.4	0.4	§	0.2	-0.2
West	0.9	1.0	0.9	1.2	0.9	0.8	0.5	0.6	0.8	0.1	0.2	0.7	0.5	0.4	§	0.3	0.0
Population Density:																	
Large MSA	0.7	0.5	1.2	0.6	0.7	0.6	0.6	0.5	0.6	0.2	0.2	0.4	0.6	0.3	§	0.0	-0.3 ss
Other MSA	0.9	0.9	0.8	0.8	0.9	0.8	0.5	0.6	0.4	0.2	0.3	0.4	0.2	0.4	§	0.2	-0.2
Non-MSA	0.8	1.0	0.6	0.7	0.7	0.7	0.3	0.4	0.5	0.5	0.2	0.1	0.2	0.2	§	0.3	+0.1
Parental Education: ^b																	
1.0–2.0 (Low)	1.8	1.9	2.3	2.0	1.4	1.4	0.9	1.2	1.1	0.3	0.3	1.0	0.6	0.5	§	0.3	-0.2
2.5–3.0	0.8	0.8	1.0	0.8	1.0	1.3	0.8	0.6	0.4	0.2	0.3	0.3	0.6	0.3	§	0.0	-0.2
3.5–4.0	0.8	0.6	0.7	0.6	0.8	0.6	0.8	0.5	0.4	0.2	0.3	0.3	0.4	0.6	§	0.1	-0.5
4.5–5.0	0.6	0.9	0.4	0.5	0.5	0.3	0.2	0.3	0.3	0.3	0.2	0.0	0.2	0.1	§	0.1	0.0
5.5–6.0 (High)	0.4	0.4	1.2	0.6	0.8	0.5	0.3	0.5	0.5	0.1	0.1	0.3	0.1	0.4	§	0.1	-0.3
Race/Ethnicity (2-year average): ^c																	
White	0.7	0.7	0.8	0.7	0.6	0.6	0.5	0.4	0.4	0.3	0.2	0.2	0.2	0.2	§	0.1	-0.1
African American	0.4	0.4	0.4	0.6	0.7	0.4	0.5	0.5	0.4	0.2	0.1	0.2	0.2	0.4	§	0.7	+0.3
Hispanic	1.1	1.2	1.3	1.2	1.3	1.4	1.0	0.7	0.6	0.3	0.1	0.4	0.6	0.5	§	0.1	-0.5 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 1995, the heroin question was changed in half of the forms. Separate questions were asked for use with and without injection. In 1996, the remaining forms were changed. Data presented here represent the combined data from all forms. Beginning in 2015 Heroin use with and without a needle were dropped in one form; data based on three of four forms; N is two thirds of N indicated.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 50
HEROIN: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995^a</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	
<i>Approximate weighted N =</i>	<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>	
Total	0.5	0.6	0.7	0.9	1.1	1.2	1.4	1.4	1.4	1.4	0.9	1.1	0.7	0.9	0.9	
Gender:																
Male	0.7	0.8	0.9	1.0	1.3	1.5	1.6	1.7	1.7	1.5	1.0	1.3	0.8	1.0	1.1	
Female	0.4	0.4	0.4	0.8	0.8	0.9	1.3	1.1	1.2	1.2	0.8	0.8	0.7	0.7	0.8	
College Plans:																
None or under 4 years	1.4	1.4	1.9	2.0	2.2	2.4	2.9	2.7	3.4	3.8	2.5	2.7	2.0	2.8	3.2	
Complete 4 years	0.3	0.4	0.4	0.7	0.9	1.0	1.2	1.2	1.1	1.0	0.7	0.8	0.5	0.6	0.6	
Region:																
Northeast	0.4	0.6	0.6	0.6	0.9	0.9	1.3	1.8	1.9	1.5	1.0	0.9	0.6	1.1	1.3	
Midwest	0.6	0.6	0.8	0.9	1.0	1.5	1.5	1.4	1.3	1.6	1.1	1.2	0.7	0.7	1.0	
South	0.6	0.5	0.6	1.0	1.3	1.4	1.5	1.3	1.4	1.5	0.9	0.8	0.9	1.0	0.8	
West	0.4	0.8	0.5	1.2	1.0	1.0	1.3	1.1	1.1	0.7	0.7	1.4	0.7	0.8	0.8	
Population Density:																
Large MSA	0.6	0.6	0.7	0.8	1.0	1.1	1.6	1.2	1.2	1.8	0.9	0.9	0.6	0.9	0.8	
Other MSA	0.5	0.6	0.6	0.9	1.0	1.3	1.3	1.5	1.4	1.2	1.0	1.0	0.8	0.9	1.0	
Non-MSA	0.4	0.6	0.7	1.0	1.3	1.2	1.6	1.5	1.6	1.2	0.9	1.4	0.9	0.7	0.9	
Parental Education: ^b																
1.0–2.0 (Low)	0.4	0.5	1.2	1.3	1.8	1.0	1.3	1.7	2.4	1.3	1.1	2.0	1.1	1.5	1.7	
2.5–3.0	0.8	0.7	0.7	0.8	1.0	1.2	1.3	1.2	1.5	1.7	1.3	0.8	0.8	1.0	1.2	
3.5–4.0	0.5	0.6	0.8	0.9	1.2	1.3	1.6	1.6	1.2	1.5	0.6	1.1	0.7	0.8	0.8	
4.5–5.0	0.4	0.5	0.3	0.9	0.9	1.1	1.5	1.3	1.4	1.3	0.8	0.7	0.6	0.7	0.8	
5.5–6.0 (High)	0.4	0.5	0.8	0.9	0.9	1.5	1.1	1.3	1.2	1.1	0.9	1.6	0.3	0.5	0.6	
Race/Ethnicity (2-year average): ^c																
White	—	0.6	0.7	0.8	1.0	1.2	1.4	1.4	1.5	1.5	1.2	1.0	0.9	0.8	0.9	
African American	—	0.3	0.4	0.6	0.6	0.2	0.2	0.4	0.5	0.5	0.5	0.3	0.3	0.4	0.6	
Hispanic	—	0.7	0.7	0.7	1.0	1.0	1.3	1.6	1.7	1.4	1.4	1.1	1.1	1.2	1.2	

Table continued on next page.

TABLE 50
HEROIN: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015 ^a	2016 ^a	2017 ^a	2018 ^a	2019 ^a	2020 ^a	2021 ^a	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	0.9	0.8	0.8	0.9	0.8	0.8	0.6	0.6	0.5	0.5	0.3	0.2	0.2	0.3	0.2	0.1	-0.1*
Gender:																	
Male	0.9	0.9	1.0	1.1	0.9	0.9	0.6	0.9	0.5	0.4	0.5	0.2	0.2	0.4	§	0.1	-0.3 s
Female	0.8	0.8	0.6	0.6	0.7	0.6	0.6	0.4	0.5	0.4	0.1	0.2	0.1	0.1	§	0.2	0.0
College Plans:																	
None or under 4 years	2.7	2.0	2.5	2.9	2.8	2.6	2.5	2.0	1.9	1.3	1.0	0.7	0.5	0.5	§	0.4	-0.1
Complete 4 years	0.6	0.7	0.6	0.7	0.6	0.6	0.4	0.5	0.4	0.3	0.2	0.1	0.1	0.2	§	0.1	-0.1
Region:																	
Northeast	0.9	1.1	0.7	0.9	0.8	0.9	0.6	0.5	0.2	0.6	0.2	0.2	0.2	0.2	§	0.0	-0.1
Midwest	0.9	0.7	1.0	1.1	0.8	0.5	0.6	0.5	0.4	0.4	0.4	0.4	0.1	0.3	§	0.1	-0.1
South	0.9	0.8	0.9	0.8	0.8	0.9	0.6	0.5	0.8	0.5	0.3	0.1	0.2	0.3	§	0.1	-0.1
West	0.7	0.9	0.7	1.0	0.8	0.7	0.6	0.9	0.5	0.4	0.4	0.3	0.2	0.4	§	0.2	-0.2
Population Density:																	
Large MSA	0.7	0.9	0.6	0.8	0.8	0.7	0.7	0.5	0.4	0.4	0.3	0.3	0.2	0.2	§	0.2	0.0
Other MSA	0.9	0.8	0.8	1.2	0.8	0.8	0.4	0.8	0.4	0.3	0.3	0.2	0.2	0.4	§	0.1	-0.3
Non-MSA	1.0	0.7	1.1	0.5	0.9	0.8	0.9	0.5	1.1	0.8	0.4	0.3	0.2	0.2	§	0.1	-0.1
Parental Education: ^b																	
1.0–2.0 (Low)	1.4	1.8	1.9	1.5	1.3	0.9	1.7	0.8	0.5	1.1	0.6	0.6	0.2	0.3	§	0.0	-0.3
2.5–3.0	0.9	0.8	0.5	1.1	0.6	1.0	0.8	0.9	0.7	0.5	0.3	0.3	0.1	0.5	§	0.0	-0.5 s
3.5–4.0	0.6	0.8	1.1	1.2	0.7	0.8	0.5	0.4	0.6	0.4	0.2	0.2	0.2	0.2	§	0.1	-0.1
4.5–5.0	0.7	0.7	0.6	0.6	0.9	0.5	0.5	0.5	0.4	0.3	0.3	0.1	0.1	0.1	§	0.1	0.0
5.5–6.0 (High)	1.2	0.5	0.6	0.6	0.8	0.5	0.3	0.9	0.5	0.4	0.4	0.1	0.3	0.5	§	0.3	-0.2
Race/Ethnicity (2-year average): ^c																	
White	1.0	0.8	0.7	0.8	0.8	0.7	0.6	0.5	0.6	0.5	0.3	0.2	0.1	0.2	§	0.1	-0.1
African American	0.6	0.4	0.5	0.4	0.4	0.8	0.8	0.7	0.6	0.5	0.4	0.4	0.2	0.1	§	0.1	+0.1
Hispanic	1.2	1.0	1.0	1.1	1.0	0.9	1.0	1.0	0.7	0.5	0.4	0.4	0.3	0.2	§	0.2	+0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 1995, the heroin question was changed in half of the forms. Separate questions were asked for use with and without injection. In 1996, the remaining forms were changed. Data presented here represent the combined data from all forms. Beginning in 2015 Heroin use with and without a needle were dropped in one form; data based on three of four forms; N is two thirds of N indicated.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 51
HEROIN: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	1.0	0.8	0.8	0.8	0.5	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.5	0.5	0.6	0.5
Gender:																
Male	1.2	1.0	1.2	1.1	0.6	0.6	0.6	0.8	0.7	0.7	0.8	0.7	0.7	0.7	0.9	0.6
Female	0.8	0.5	0.4	0.6	0.3	0.4	0.3	0.4	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.3
College Plans:																
None or under 4 years	—	0.9	1.1	1.0	0.7	0.6	0.5	0.7	0.9	0.6	0.7	0.8	0.5	0.8	0.9	0.6
Complete 4 years	—	0.6	0.5	0.6	0.3	0.3	0.5	0.4	0.3	0.4	0.5	0.4	0.4	0.3	0.5	0.4
Region:																
Northeast	1.1	0.7	0.7	0.6	0.6	0.5	0.5	0.9	0.6	0.6	0.8	0.7	0.6	0.5	0.9	0.8
Midwest	1.3	1.0	1.0	0.8	0.5	0.7	0.6	0.5	0.4	0.6	0.6	0.4	0.6	0.3	0.6	0.3
South	0.9	0.7	0.9	1.1	0.6	0.3	0.5	0.5	0.7	0.5	0.6	0.5	0.4	0.5	0.6	0.5
West	0.7	0.6	0.5	0.8	0.2	0.4	0.5	0.3	0.5	0.4	0.3	0.5	0.5	0.7	0.7	0.3
Population Density:																
Large MSA	1.3	1.0	0.5	0.7	0.4	0.3	0.3	0.7	0.6	0.6	0.7	0.7	0.3	0.4	0.5	0.4
Other MSA	0.9	1.0	0.8	0.8	0.6	0.5	0.5	0.4	0.4	0.4	0.7	0.4	0.6	0.5	0.7	0.5
Non-MSA	1.0	0.4	1.1	1.0	0.5	0.6	0.7	0.6	0.7	0.7	0.4	0.5	0.5	0.5	0.8	0.5
Parental Education: ^a																
1.0–2.0 (Low)	1.2	0.8	0.8	0.8	0.6	0.6	0.4	0.4	0.5	0.6	0.8	0.9	0.5	0.5	0.9	0.8
2.5–3.0	0.8	0.9	0.8	0.9	0.5	0.6	0.6	0.7	0.6	0.5	0.5	0.4	0.4	0.7	0.7	0.4
3.5–4.0	0.6	0.8	0.9	0.7	0.4	0.4	0.5	0.3	0.5	0.5	0.5	0.5	0.6	0.4	0.6	0.4
4.5–5.0	1.2	1.4	0.6	0.9	0.6	0.4	0.3	0.6	0.4	0.4	0.7	0.3	0.3	0.3	0.6	0.4
5.5–6.0 (High)	1.2	0.6	1.1	1.0	0.8	0.4	0.7	1.1	0.8	0.5	0.6	0.5	0.7	0.4	0.4	0.5
Race/Ethnicity (2-year average): ^b																
White	—	—	0.8	0.8	0.6	0.5	0.4	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.5	0.6
African American	—	—	0.6	0.6	0.5	0.5	0.6	0.7	0.6	0.4	0.5	0.5	0.5	0.7	0.6	0.3
Hispanic	—	—	1.2	2.0	1.7	0.4	0.3	0.4	0.6	1.1	1.0	0.9	0.9	0.5	0.5	0.6

Table continued on next page.

TABLE 51 (cont.)
HEROIN: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995 ^c	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	0.4	0.6	0.5	0.6	1.1	1.0	1.2	1.0	1.1	1.5	0.9	1.0	0.8	0.9	0.8	0.8
Gender:																
Male	0.6	0.8	0.7	0.8	1.4	1.3	1.5	1.4	1.4	1.7	1.3	1.1	0.8	1.2	1.2	1.0
Female	0.3	0.3	0.3	0.4	0.8	0.7	0.9	0.7	0.8	1.2	0.6	0.8	0.5	0.4	0.5	0.4
College Plans:																
None or under 4 years	0.5	0.9	1.0	1.1	1.5	1.8	1.8	1.7	1.3	2.2	1.3	2.3	0.9	1.6	1.8	1.3
Complete 4 years	0.4	0.5	0.4	0.5	0.9	0.8	1.0	0.8	1.0	1.1	0.7	0.7	0.6	0.6	0.6	0.6
Region:																
Northeast	0.2	0.5	0.9	0.7	1.0	1.6	1.7	1.3	1.2	1.0	1.2	1.2	0.8	0.9	1.0	1.1
Midwest	0.8	0.6	0.5	0.9	0.7	0.7	0.9	1.0	0.9	1.4	1.3	1.0	0.6	0.6	0.8	0.4
South	0.4	0.6	0.4	0.6	1.4	1.0	1.1	1.1	1.3	1.6	0.6	1.2	1.0	1.0	0.9	0.8
West	0.3	0.8	0.5	0.4	1.0	0.9	1.2	0.6	0.9	1.8	0.8	0.5	0.5	0.9	0.7	0.7
Population Density:																
Large MSA	0.4	0.4	0.6	0.4	1.4	1.1	1.1	0.9	0.8	1.9	1.1	1.0	0.7	0.8	0.8	1.0
Other MSA	0.4	0.7	0.5	0.8	0.9	1.1	1.3	1.3	1.3	1.3	0.8	1.0	0.8	1.0	0.9	0.6
Non-MSA	0.6	0.7	0.5	0.5	1.0	0.9	1.0	0.6	1.0	1.3	0.9	1.1	0.9	0.6	0.8	0.6
Parental Education: ^a																
1.0–2.0 (Low)	0.5	0.7	0.3	0.9	1.8	1.1	2.1	1.2	2.3	2.3	1.3	2.1	1.1	1.4	1.0	1.2
2.5–3.0	0.4	0.6	0.4	0.8	1.1	0.9	1.0	1.0	0.9	1.6	0.5	1.0	0.7	0.9	1.2	0.9
3.5–4.0	0.4	0.6	0.6	0.4	0.9	1.1	1.2	1.2	1.0	1.3	1.0	0.9	0.8	0.7	0.8	0.7
4.5–5.0	0.6	0.7	0.7	0.3	1.1	1.0	1.1	1.0	1.1	1.5	1.0	0.7	0.6	0.9	0.5	0.5
5.5–6.0 (High)	0.5	0.3	0.4	0.9	1.0	0.8	1.1	0.7	1.1	1.1	0.8	1.0	0.6	0.6	0.8	0.7
Race/Ethnicity (2-year average): ^b																
White	0.5	0.5	0.5	0.5	0.8	1.0	1.2	1.2	1.1	1.3	1.3	1.0	0.8	0.7	0.7	0.7
African American	0.2	0.5	0.4	0.3	0.4	0.5	0.5	0.4	0.3	0.5	0.4	0.5	0.7	0.7	0.8	0.7
Hispanic	0.6	0.9	0.7	0.5	1.2	1.5	1.1	0.8	1.0	2.0	1.6	0.8	0.8	1.0	1.3	1.2

Table continued on next page.

TABLE 51 (cont.)
HEROIN: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	0.9	0.7	0.7	0.9	0.8	0.6	0.6	0.6	0.5	0.3	0.4	0.4	0.4	0.3	0.1	-0.3* sss
Gender:																
Male	1.2	1.0	1.0	1.1	1.0	0.8	0.8	0.9	0.7	0.4	0.4	0.3	0.5	§	0.1	-0.4 sss
Female	0.6	0.4	0.5	0.5	0.4	0.4	0.3	0.2	0.2	0.2	0.3	0.2	0.2	§	0.0	-0.2 ss
College Plans:																
None or under 4 years	2.2	1.4	1.1	1.7	1.2	1.2	1.0	1.1	1.3	0.8	0.7	0.8	0.8	§	0.2	-0.6 ss
Complete 4 years	0.7	0.5	0.6	0.6	0.7	0.5	0.4	0.5	0.3	0.1	0.2	0.2	0.3	§	0.0	-0.2 sss
Region:																
Northeast	1.1	0.7	0.8	1.3	0.7	0.8	0.6	0.8	0.3	0.2	0.1	0.2	0.4	§	0.1	-0.3 s
Midwest	0.8	0.6	0.8	0.9	0.7	0.7	0.6	0.6	0.6	0.4	0.2	0.6	0.1	§	0.0	-0.1
South	0.9	0.7	0.6	0.7	0.7	0.4	0.5	0.5	0.4	0.2	0.6	0.4	0.5	§	0.2	-0.4 ss
West	0.9	1.0	0.9	0.7	1.0	0.8	0.7	0.5	0.6	0.4	0.4	0.1	0.4	§	0.2	-0.2
Population Density:																
Large MSA	0.9	0.9	0.9	0.8	0.9	0.7	0.5	0.4	0.4	0.4	0.3	0.3	0.5	§	0.1	-0.3 s
Other MSA	0.9	0.8	0.8	0.9	0.8	0.7	0.6	0.7	0.4	0.2	0.4	0.4	0.3	§	0.1	-0.2 s
Non-MSA	1.0	0.5	0.6	0.8	0.5	0.5	0.6	0.5	0.6	0.4	0.5	0.3	0.4	§	0.1	-0.3
Parental Education: ^a																
1.0–2.0 (Low)	1.7	1.3	1.5	1.1	1.0	1.3	1.0	1.3	0.8	0.5	0.7	1.1	0.9	§	0.0	-0.9
2.5–3.0	1.0	0.5	0.8	1.0	0.5	0.8	0.6	0.6	0.4	0.2	0.6	0.3	0.4	§	0.0	-0.4
3.5–4.0	0.8	0.9	0.6	0.9	0.8	0.4	0.4	0.3	0.6	0.3	0.3	0.3	0.4	§	0.1	-0.3 s
4.5–5.0	0.9	0.5	0.6	0.7	0.8	0.5	0.5	0.7	0.4	0.2	0.2	0.1	0.1	§	0.1	+0.1
5.5–6.0 (High)	0.5	0.7	0.5	0.6	0.7	0.4	0.6	0.5	0.2	0.3	0.2	0.2	0.3	§	0.2	-0.1
Race/Ethnicity (2-year average): ^b																
White	0.9	0.9	0.7	0.8	0.9	0.7	0.5	0.4	0.4	0.3	0.2	0.2	0.2	§	0.0	-0.2 sss
African American	0.5	0.4	0.4	0.6	0.7	0.8	0.8	0.9	1.0	0.6	0.5	0.6	0.7	§	0.6	0.0
Hispanic	1.1	0.9	0.7	0.6	0.5	0.4	0.5	0.6	0.5	0.4	0.4	0.4	0.5	§	0.2	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cIn 1995, the heroin question was changed in half of the forms. Separate questions were asked for use with and without injection. In 1996, the remaining forms were changed. Data presented here represent the combined data from all forms.

TABLE 52
HEROIN WITH A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months											
	1991– 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	—	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	—	0.9	1.0	0.8	0.8	0.9	0.6	0.7	0.6	0.6	0.7	0.6
Gender:												
Male	—	1.2	1.0	0.8	1.0	1.0	0.7	0.7	0.6	0.7	0.7	0.6
Female	—	0.5	1.0	0.7	0.7	0.8	0.6	0.7	0.6	0.5	0.7	0.6
College Plans:												
None or under 4 years	—	3.2	3.1	2.0	3.1	2.2	2.1	2.2	1.9	2.0	3.1	1.5
Complete 4 years	—	0.6	0.7	0.6	0.6	0.7	0.5	0.6	0.5	0.4	0.4	0.4
Region:												
Northeast	—	0.8	1.3	0.5	0.6	1.0	0.8	0.9	0.6	0.4	0.7	0.4
Midwest	—	0.9	1.1	0.7	0.9	1.2	0.9	0.7	0.8	0.6	0.6	0.3
South	—	0.8	0.9	0.8	1.0	0.9	0.3	0.9	0.6	0.7	0.6	0.8
West	—	1.0	1.0	1.0	0.8	0.7	0.8	0.3	0.6	0.7	0.9	0.6
Population Density:												
Large MSA	—	0.9	0.7	0.6	0.8	0.8	0.7	0.7	0.4	0.5	1.0	0.3
Other MSA	—	0.9	1.2	0.8	0.8	0.9	0.6	0.6	0.7	0.6	0.7	0.7
Non-MSA	—	0.9	1.1	0.9	1.1	1.1	0.7	1.0	0.8	0.9	0.5	0.7
Parental Education: ^a												
1.0–2.0 (Low)	—	1.3	0.9	1.5	2.3	2.0	1.2	2.0	1.1	1.6	1.6	0.3
2.5–3.0	—	0.5	1.2	0.8	0.8	0.9	0.5	0.7	0.9	0.6	0.8	1.0
3.5–4.0	—	0.8	1.2	0.6	0.6	0.9	0.4	0.6	0.6	0.4	0.7	0.7
4.5–5.0	—	1.0	0.8	0.7	0.7	0.5	0.4	0.5	0.5	0.5	0.6	0.3
5.5–6.0 (High)	—	1.0	0.6	0.9	0.9	1.2	0.9	0.7	0.4	0.5	0.5	0.3
Race/Ethnicity (2-year average): ^b												
White	—	—	1.1	1.0	0.7	0.8	0.7	0.6	0.7	0.6	0.6	0.5
African American	—	—	0.3	0.2	0.3	0.4	0.3	0.5	0.5	0.4	0.5	0.4
Hispanic	—	—	1.1	1.0	1.1	1.5	1.2	0.8	0.7	0.9	1.3	1.1

Table continued on next page.

TABLE 52 (cont.)
HEROIN WITH A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021 change	
	<i>Approximate weighted N =</i>	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
Total	16,500	0.5	0.6	0.5	0.5	0.6	0.5	0.4	0.3	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.1	-0.1*
Gender:																		
Male		0.4	0.5	0.6	0.6	0.6	0.5	0.4	0.2	0.5	0.2	0.2	0.2	0.2	0.2	§	0.1	-0.1
Female		0.6	0.5	0.4	0.5	0.5	0.4	0.3	0.4	0.3	0.2	0.1	0.3	0.1	0.2	§	0.0	-0.2 s
College Plans:																		
None or under 4 years		2.1	2.7	2.1	2.6	1.3	2.1	1.3	1.8	2.2	0.6	0.7	1.1	0.5	0.3	§	0.4	+0.1
Complete 4 years		0.4	0.3	0.4	0.4	0.5	0.3	0.3	0.2	0.2	0.1	0.1	0.1	0.1	0.2	§	0.0	-0.1 s
Region:																		
Northeast		0.3	0.4	0.5	0.3	0.5	0.5	0.3	0.2	0.2	0.1	0.2	0.1	0.2	0.2	§	0.0	-0.2
Midwest		0.4	0.6	0.5	0.5	0.6	0.3	0.3	0.3	0.4	0.2	0.1	0.3	0.1	0.0	§	0.0	0.0
South		0.6	0.6	0.5	0.5	0.5	0.4	0.5	0.4	0.3	0.3	0.3	0.2	0.2	0.3	§	0.2	-0.1
West		0.7	0.6	0.6	0.9	0.7	0.6	0.3	0.3	0.7	0.1	0.1	0.3	0.4	0.3	§	0.2	-0.1
Population Density:																		
Large MSA		0.5	0.3	0.7	0.5	0.6	0.4	0.4	0.2	0.5	0.2	0.2	0.3	0.3	0.2	§	0.0	-0.2 s
Other MSA		0.6	0.6	0.5	0.6	0.7	0.5	0.4	0.4	0.4	0.2	0.2	0.2	0.2	0.3	§	0.2	-0.1
Non-MSA		0.4	0.7	0.3	0.5	0.3	0.4	0.2	0.2	0.4	0.3	0.2	0.1	0.1	0.0	§	0.2	+0.2
Parental Education: ^a																		
1.0–2.0 (Low)		1.4	1.3	1.6	1.6	0.6	1.0	0.6	0.8	0.9	0.3	0.2	0.7	0.5	0.3	§	0.0	-0.3
2.5–3.0		0.5	0.6	0.5	0.5	0.7	0.9	0.5	0.4	0.3	0.1	0.2	0.2	0.4	0.3	§	0.0	-0.3
3.5–4.0		0.4	0.3	0.4	0.3	0.6	0.4	0.7	0.3	0.3	0.1	0.2	0.2	0.2	0.3	§	0.1	-0.2
4.5–5.0		0.4	0.6	0.3	0.4	0.4	0.2	0.1	0.2	0.3	0.2	0.2	0.0	0.1	0.1	§	0.0	-0.1
5.5–6.0 (High)		0.2	0.3	0.7	0.5	0.6	0.4	0.2	0.3	0.4	0.1	0.1	0.1	0.1	0.1	§	0.1	0.0
Race/Ethnicity (2-year average): ^b																		
White		0.4	0.5	0.5	0.4	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.1	0.1	0.2	§	0.1	-0.1
African American		0.3	0.2	0.2	0.5	0.6	0.4	0.4	0.5	0.4	0.2	0.1	0.2	0.2	0.1	§	0.7	+0.5
Hispanic		0.9	0.9	1.0	0.9	1.0	1.1	0.7	0.4	0.4	0.2	0.1	0.2	0.3	0.3	§	0.1	-0.3 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1995; *N* is one half of *N* indicated in Table S1. Data based on all forms beginning in 1996. Beginning in 2015 Heroin use with and without a needle were dropped in one form; data based on three of four forms; *N* is two thirds of *N* indicated.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 53
HEROIN WITH A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months											
	1991– 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Approximate weighted N =	—	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	—	0.6	0.7	0.7	0.8	0.6	0.5	0.4	0.6	0.5	0.5	0.5
Gender:												
Male	—	0.8	0.8	0.9	1.0	0.8	0.6	0.5	0.6	0.5	0.6	0.6
Female	—	0.4	0.4	0.5	0.5	0.5	0.4	0.3	0.4	0.4	0.3	0.4
College Plans:												
None or under 4 years	—	1.6	1.4	1.5	1.6	1.7	1.7	1.4	1.5	1.4	1.6	1.6
Complete 4 years	—	0.4	0.5	0.6	0.6	0.5	0.3	0.3	0.4	0.3	0.3	0.3
Region:												
Northeast	—	0.5	0.3	0.6	0.8	0.7	0.5	0.5	0.4	0.4	0.5	0.6
Midwest	—	0.6	1.0	0.8	0.9	0.7	0.5	0.5	0.6	0.5	0.4	0.5
South	—	0.6	0.7	0.8	0.8	0.6	0.6	0.3	0.5	0.5	0.5	0.4
West	—	0.5	0.5	0.6	0.6	0.6	0.4	0.5	0.8	0.5	0.5	0.6
Population Density:												
Large MSA	—	0.5	0.5	0.6	0.7	0.6	0.6	0.4	0.5	0.4	0.5	0.4
Other MSA	—	0.5	0.7	0.6	0.7	0.5	0.4	0.5	0.4	0.4	0.5	0.6
Non-MSA	—	0.8	0.8	1.1	1.0	1.0	0.6	0.4	1.0	0.6	0.3	0.5
Parental Education: ^a												
1.0–2.0 (Low)	—	1.6	0.6	0.7	1.3	1.2	1.0	0.5	1.2	0.8	1.0	1.4
2.5–3.0	—	0.5	0.8	0.6	0.6	0.7	0.8	0.7	0.3	0.4	0.5	0.6
3.5–4.0	—	0.7	0.7	0.9	0.9	0.6	0.5	0.2	0.6	0.5	0.4	0.4
4.5–5.0	—	0.4	0.5	0.7	0.6	0.6	0.3	0.3	0.3	0.5	0.3	0.4
5.5–6.0 (High)	—	0.3	0.6	0.7	0.8	0.6	0.3	0.5	1.1	0.2	0.3	0.3
Race/Ethnicity (2-year average): ^b												
White	—	—	0.6	0.7	0.7	0.6	0.5	0.4	0.5	0.5	0.4	0.4
African American	—	—	0.1	0.1	0.3	0.3	0.3	0.3	0.2	0.1	0.2	0.5
Hispanic	—	—	0.6	0.6	0.8	1.0	0.9	0.9	0.7	0.7	0.8	0.8

Table continued on next page.

TABLE 53 (cont.)
HEROIN WITH A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	0.5	0.5	0.5	0.6	0.5	0.5	0.4	0.5	0.4	0.2	0.3	0.2	0.1	0.2	0.2	0.1	-0.1*
Gender:																	
Male	0.6	0.5	0.6	0.8	0.5	0.6	0.5	0.6	0.4	0.2	0.5	0.2	0.2	0.3	§	0.1	-0.3
Female	0.4	0.5	0.4	0.3	0.4	0.4	0.3	0.3	0.4	0.2	0.1	0.2	0.1	0.1	§	0.1	0.0
College Plans:																	
None or under 4 years	1.3	1.3	1.7	2.1	1.7	1.6	1.6	1.1	1.7	0.5	1.0	0.6	0.3	0.4	§	0.3	-0.1
Complete 4 years	0.4	0.4	0.3	0.4	0.3	0.4	0.3	0.4	0.3	0.2	0.2	0.1	0.1	0.2	§	0.1	-0.1
Region:																	
Northeast	0.6	0.6	0.4	0.8	0.5	0.4	0.3	0.4	0.2	0.3	0.1	0.1	0.1	0.2	§	0.0	-0.1
Midwest	0.5	0.4	0.5	0.7	0.5	0.4	0.5	0.5	0.4	0.3	0.4	0.3	0.1	0.1	§	0.1	0.0
South	0.6	0.4	0.6	0.4	0.5	0.7	0.3	0.4	0.5	0.2	0.2	0.1	0.1	0.3	§	0.1	-0.1
West	0.3	0.6	0.5	0.5	0.4	0.4	0.4	0.6	0.5	0.1	0.4	0.2	0.1	0.4	§	0.2	-0.1
Population Density:																	
Large MSA	0.4	0.5	0.4	0.5	0.5	0.4	0.4	0.3	0.3	0.3	0.3	0.2	0.1	0.1	§	0.2	+0.1
Other MSA	0.5	0.5	0.5	0.8	0.4	0.5	0.3	0.6	0.3	0.1	0.3	0.2	0.1	0.3	§	0.1	-0.2
Non-MSA	0.7	0.4	0.7	0.4	0.6	0.6	0.5	0.4	1.0	0.4	0.3	0.1	0.1	0.2	§	0.1	-0.1
Parental Education: ^a																	
1.0–2.0 (Low)	1.0	1.3	1.7	1.0	0.9	0.6	1.2	0.4	0.5	0.4	0.5	0.6	0.1	0.2	§	0.0	-0.2
2.5–3.0	0.5	0.5	0.2	0.7	0.2	0.5	0.5	0.5	0.5	0.3	0.3	0.2	0.1	0.4	§	0.0	-0.4
3.5–4.0	0.4	0.4	0.5	0.8	0.4	0.5	0.3	0.3	0.5	0.2	0.2	0.1	0.1	0.1	§	0.0	-0.1
4.5–5.0	0.3	0.5	0.3	0.3	0.4	0.4	0.3	0.4	0.3	0.3	0.2	0.1	0.1	0.0	§	0.1	+0.1
5.5–6.0 (High)	1.0	0.2	0.3	0.4	0.7	0.2	0.3	0.7	0.3	0.1	0.3	0.1	0.2	0.5	§	0.3	-0.2
Race/Ethnicity (2-year average): ^b																	
White	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.4	0.3	0.2	0.2	0.1	0.2	§	0.1	-0.1
African American	0.4	0.2	0.4	0.4	0.4	0.6	0.5	0.4	0.5	0.4	0.4	0.3	0.2	0.1	§	0.1	+0.1
Hispanic	0.8	0.6	0.7	0.8	0.6	0.4	0.6	0.6	0.5	0.3	0.3	0.4	0.2	0.1	§	0.2	+0.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: *s* = .05, *ss* = .01, *sss* = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1995; *N* is one half of *N* indicated in Table S2. Data based on all forms beginning in 1996. Beginning in 2015 Heroin use with and without a needle were dropped in one form; data based on three of four forms; *N* is two thirds of *N* indicated.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 54
HEROIN WITH A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months											
	1975– 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	—	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700
Total	—	0.5	0.5	0.5	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.5
Gender:												
Male	—	0.7	0.9	0.9	0.6	0.6	0.6	0.5	0.5	0.6	0.6	0.6
Female	—	0.3	0.1	0.2	0.3	0.2	0.2	0.1	0.2	0.3	0.1	0.3
College Plans:												
None or under 4 years	—	0.9	1.2	0.9	0.8	0.5	1.0	0.7	0.9	0.4	1.0	0.9
Complete 4 years	—	0.4	0.4	0.4	0.3	0.4	0.2	0.2	0.3	0.4	0.2	0.4
Region:												
Northeast	—	0.4	0.6	0.6	0.7	0.3	0.2	0.7	0.6	0.6	0.6	0.6
Midwest	—	0.5	0.4	0.8	0.4	0.5	0.4	0.2	0.6	0.3	0.2	0.7
South	—	0.4	0.6	0.3	0.5	0.4	0.8	0.1	0.3	0.6	0.4	0.4
West	—	0.8	0.5	0.5	0.1	0.3	0.1	0.4	0.2	0.2	0.5	0.3
Population Density:												
Large MSA	—	0.6	0.7	0.3	0.3	0.4	0.8	0.4	0.5	0.5	0.6	0.4
Other MSA	—	0.4	0.5	0.6	0.5	0.3	0.3	0.3	0.3	0.4	0.4	0.6
Non-MSA	—	0.6	0.3	0.7	0.5	0.6	0.3	0.2	0.5	0.4	0.2	0.5
Parental Education: ^a												
1.0–2.0 (Low)	—	0.7	0.9	1.8	0.8	1.5	1.2	0.7	0.8	0.5	0.2	0.8
2.5–3.0	—	0.6	0.5	0.4	0.5	0.1	0.4	0.3	0.4	0.7	0.6	0.5
3.5–4.0	—	0.2	0.3	0.3	0.4	0.5	0.2	0.1	0.5	0.3	0.3	0.4
4.5–5.0	—	0.5	0.7	0.6	0.4	0.3	0.4	0.4	0.2	0.4	0.4	0.3
5.5–6.0 (High)	—	0.6	0.5	0.5	0.2	0.4	0.2	0.3	0.2	0.3	0.4	0.4
Race/Ethnicity (2-year average): ^b												
White	—	—	0.4	0.5	0.5	0.4	0.3	0.2	0.2	0.3	0.3	0.3
African American	—	—	0.2	0.5	0.4	0.2	0.5	0.4	0.2	0.4	0.5	0.6
Hispanic	—	—	1.0	0.7	0.5	0.6	1.0	0.6	0.4	0.6	0.7	0.8

Table continued on next page.

TABLE 54 (cont.)
HEROIN WITH A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																2019-2021 change
	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	
<i>Approximate weighted N =</i>	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	0.5	0.4	0.4	0.3	0.7	0.6	0.4	0.4	0.5	0.3	0.3	0.2	0.3	0.3	0.1	0.1	-0.2* s
Gender:																	
Male	0.7	0.5	0.6	0.4	1.0	0.8	0.5	0.5	0.8	0.4	0.3	0.2	0.3	0.4	§	0.1	-0.3 s
Female	0.3	0.1	0.1	0.2	0.2	0.3	0.3	0.3	0.1	0.1	0.2	0.2	0.1	0.2	§	0.0	-0.2
College Plans:																	
None or under 4 years	1.0	1.0	1.0	0.6	1.6	1.3	1.1	0.7	0.9	0.8	1.0	0.4	1.0	0.7	§	0.2	-0.5
Complete 4 years	0.3	0.2	0.2	0.2	0.4	0.5	0.3	0.2	0.4	0.2	0.1	0.2	0.1	0.2	§	0.0	-0.2 s
Region:																	
Northeast	0.8	0.7	0.3	0.3	0.6	0.8	0.6	0.3	0.7	0.0	0.1	0.1	0.2	0.6	§	0.1	-0.5 s
Midwest	0.1	0.1	0.3	0.3	0.8	0.4	0.6	0.3	0.7	0.6	0.3	0.1	0.6	0.1	§	0.0	-0.1
South	0.6	0.4	0.3	0.2	0.8	0.5	0.2	0.4	0.3	0.3	0.3	0.4	0.4	0.3	§	0.1	-0.2
West	0.3	0.2	0.5	0.3	0.4	1.0	0.4	0.5	0.5	0.4	0.3	0.1	0.1	0.1	§	0.2	0.0
Population Density:																	
Large MSA	0.7	0.6	0.2	0.5	0.7	0.8	0.5	0.3	0.4	0.5	0.2	0.2	0.4	0.4	§	0.2	-0.2
Other MSA	0.4	0.2	0.5	0.2	0.6	0.7	0.4	0.4	0.6	0.2	0.3	0.3	0.3	0.2	§	0.1	-0.1
Non-MSA	0.4	0.4	0.2	0.1	0.7	0.3	0.2	0.3	0.4	0.4	0.5	0.2	0.2	0.4	§	0.1	-0.3
Parental Education: ^a																	
1.0–2.0 (Low)	1.0	1.0	0.9	0.8	1.4	1.4	0.4	0.8	1.5	0.8	0.7	0.6	1.4	0.5	§	0.0	-0.5
2.5–3.0	0.5	0.4	0.3	0.5	0.9	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	§	0.0	-0.3
3.5–4.0	0.5	0.2	0.4	0.2	0.5	0.8	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.4	§	0.2	-0.2
4.5–5.0	0.2	0.2	0.3	0.1	0.5	0.4	0.5	0.3	0.6	0.4	0.1	0.1	0.1	0.1	§	0.0	-0.1
5.5–6.0 (High)	0.6	0.3	0.2	0.1	0.3	0.6	0.3	0.0	0.1	0.1	0.2	0.3	0.1	0.3	§	0.3	0.0
Race/Ethnicity (2-year average): ^b																	
White	0.3	0.3	0.3	0.3	0.4	0.6	0.5	0.3	0.2	0.3	0.3	0.2	0.1	0.1	§	0.0	-0.1 s
African American	0.7	0.4	0.3	0.3	0.7	0.9	0.5	0.5	0.8	0.6	0.2	0.4	0.6	0.9	§	0.1	-0.7 s
Hispanic	0.9	0.7	0.4	0.3	0.5	0.6	0.2	0.3	0.6	0.7	0.4	0.3	0.4	0.4	§	0.1	-0.4 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on three of six forms; N is three sixths of N indicated in Table S3.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 55
HEROIN WITHOUT A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months											
	1991– 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	—	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	—	0.8	1.0	0.8	0.8	0.9	0.7	0.6	0.6	0.6	0.6	0.5
Gender:												
Male	—	1.0	0.9	0.9	0.9	0.8	0.5	0.6	0.6	0.6	0.4	0.4
Female	—	0.6	0.9	0.7	0.8	0.8	0.8	0.6	0.7	0.6	0.7	0.5
College Plans:												
None or under 4 years	—	3.2	2.1	2.2	3.4	2.7	2.4	1.9	1.7	2.3	2.5	1.6
Complete 4 years	—	0.5	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4
Region:												
Northeast	—	0.9	1.1	0.8	0.7	0.7	0.6	0.7	0.3	0.5	0.6	0.5
Midwest	—	1.0	1.0	0.9	0.9	1.0	0.8	0.6	0.7	0.8	0.4	0.3
South	—	0.8	0.8	0.7	0.9	0.9	0.5	0.6	0.7	0.7	0.6	0.6
West	—	0.7	1.1	1.1	0.9	0.9	0.8	0.5	0.7	0.5	0.7	0.4
Population Density:												
Large MSA	—	0.9	1.0	0.6	0.5	0.8	0.7	0.5	0.6	0.6	0.7	0.4
Other MSA	—	0.8	1.0	0.9	1.0	0.8	0.8	0.8	0.5	0.6	0.6	0.5
Non-MSA	—	0.9	0.9	1.0	1.0	1.1	0.5	0.4	0.8	0.7	0.3	0.6
Parental Education: ^a												
1.0–2.0 (Low)	—	1.9	1.5	1.3	2.4	1.7	1.1	1.3	0.5	1.4	1.5	0.2
2.5–3.0	—	0.7	0.9	1.1	0.7	1.0	1.3	0.6	0.8	0.6	0.6	0.9
3.5–4.0	—	0.5	1.2	0.5	0.8	0.6	0.5	0.6	0.6	0.7	0.6	0.5
4.5–5.0	—	0.8	0.8	0.8	0.6	0.8	0.5	0.5	0.6	0.4	0.4	0.3
5.5–6.0 (High)	—	0.7	0.8	1.0	1.1	1.1	0.3	0.4	0.4	0.5	0.3	0.1
Race/Ethnicity (2-year average): ^b												
White	—	—	1.0	1.0	0.8	0.8	0.8	0.7	0.6	0.6	0.5	0.4
African American	—	—	0.2	0.2	0.4	0.5	0.3	0.3	0.3	0.3	0.5	0.4
Hispanic	—	—	1.5	1.2	1.1	1.4	1.1	0.8	1.0	1.0	1.0	0.9

Table continued on next page.

TABLE 55 (cont.)
HEROIN WITHOUT A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	0.5	0.5	0.6	0.4	0.5	0.4	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.1	-0.1*
Gender:																	
Male	0.5	0.4	0.6	0.4	0.5	0.4	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.2	§	0.0	-0.1
Female	0.6	0.5	0.6	0.4	0.5	0.5	0.3	0.3	0.2	0.2	0.1	0.3	0.1	0.3	§	0.1	-0.1
College Plans:																	
None or under 4 years	2.3	1.9	2.6	1.9	1.8	1.8	1.1	1.3	0.5	0.4	0.9	1.3	0.8	0.4	§	0.2	-0.2
Complete 4 years	0.4	0.3	0.4	0.2	0.4	0.3	0.2	0.3	0.2	0.1	0.1	0.1	0.1	0.2	§	0.0	-0.1 s
Region:																	
Northeast	0.2	0.2	0.4	0.3	0.4	0.3	0.1	0.3	0.1	0.1	0.1	0.2	0.1	0.0	§	.	.
Midwest	0.7	0.5	0.8	0.3	0.4	0.5	0.2	0.3	0.1	0.2	0.1	0.2	0.1	0.3	§	0.1	-0.2
South	0.6	0.4	0.7	0.3	0.5	0.4	0.3	0.4	0.2	0.3	0.3	0.2	0.3	0.3	§	0.1	-0.2
West	0.5	0.7	0.5	0.7	0.6	0.5	0.4	0.3	0.4	0.0	0.1	0.5	0.4	0.2	§	0.1	-0.1
Population Density:																	
Large MSA	0.4	0.3	0.8	0.3	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.4	0.1	§	0.0	-0.1
Other MSA	0.5	0.5	0.5	0.5	0.6	0.5	0.2	0.3	0.2	0.2	0.1	0.3	0.2	0.3	§	0.1	-0.1
Non-MSA	0.7	0.7	0.4	0.3	0.5	0.4	0.2	0.3	0.2	0.3	0.2	0.1	0.1	0.2	§	0.1	-0.1
Parental Education: ^a																	
1.0–2.0 (Low)	1.4	1.0	1.6	1.1	1.4	0.7	0.7	0.9	0.5	0.3	0.3	0.5	0.6	0.2	§	0.3	+0.1
2.5–3.0	0.6	0.4	0.8	0.4	0.6	1.0	0.4	0.4	0.2	0.2	0.3	0.3	0.4	0.2	§	0.0	-0.1
3.5–4.0	0.6	0.3	0.4	0.4	0.4	0.4	0.3	0.2	0.2	0.2	0.3	0.3	0.2	0.4	§	0.0	-0.4
4.5–5.0	0.4	0.7	0.3	0.2	0.2	0.2	0.1	0.1	0.1	0.2	0.1	0.0	0.1	0.0	§	0.1	+0.1
5.5–6.0 (High)	0.3	0.3	0.8	0.2	0.5	0.2	0.1	0.4	0.2	0.0	0.0	0.2	0.1	0.4	§	0.0	-0.4
Race/Ethnicity (2-year average): ^b																	
White	0.5	0.5	0.5	0.4	0.3	0.4	0.3	0.2	0.2	0.1	0.2	0.1	0.1	0.1	§	0.1	-0.1
African American	0.3	0.3	0.3	0.3	0.2	0.1	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.3	§	0.6	+0.2
Hispanic	0.6	0.7	0.8	0.8	0.8	0.8	0.6	0.5	0.5	0.2	0.1	0.3	0.5	0.4	§	0.0	-0.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1995; N is one half of N indicated in Table S1. Data based on all forms beginning in 1996. Beginning in 2015 Heroin use with and without a needle were dropped in one form; data based on three of four forms; N is two thirds of N indicated.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 56
HEROIN WITHOUT A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months											
	1991– 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Approximate weighted N =	—	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	—	0.8	0.9	1.1	1.0	1.1	1.1	0.7	0.8	0.5	0.7	0.7
Gender:												
Male	—	0.9	1.1	1.2	1.2	1.3	1.1	0.8	0.9	0.6	0.8	0.7
Female	—	0.7	0.7	1.0	0.8	1.0	1.0	0.6	0.6	0.4	0.5	0.6
College Plans:												
None or under 4 years	—	1.7	1.8	2.1	1.8	2.5	2.7	1.7	2.2	1.4	2.2	2.3
Complete 4 years	—	0.6	0.7	1.0	0.9	0.8	0.8	0.5	0.6	0.4	0.5	0.4
Region:												
Northeast	—	0.5	0.7	1.0	1.4	1.6	1.2	0.8	0.7	0.4	0.9	0.9
Midwest	—	0.5	0.8	1.2	0.7	1.0	1.2	0.8	0.9	0.5	0.5	0.8
South	—	1.1	1.1	1.2	1.0	1.0	1.3	0.7	0.5	0.7	0.7	0.6
West	—	0.9	0.9	1.0	0.8	0.8	0.5	0.5	1.2	0.4	0.5	0.5
Population Density:												
Large MSA	—	0.9	0.9	1.3	0.8	0.9	1.5	0.8	0.6	0.4	0.6	0.7
Other MSA	—	0.7	1.0	1.0	1.1	1.2	1.0	0.7	0.8	0.6	0.8	0.7
Non-MSA	—	0.9	0.7	1.1	1.0	1.2	0.9	0.7	1.0	0.6	0.5	0.6
Parental Education: ^a												
1.0–2.0 (Low)	—	1.7	0.8	0.9	1.2	1.8	1.1	0.9	1.5	0.9	1.0	1.1
2.5–3.0	—	0.6	0.7	1.0	1.0	1.1	1.1	0.9	0.7	0.7	0.7	0.9
3.5–4.0	—	0.9	1.0	1.3	1.2	0.9	1.3	0.5	0.8	0.4	0.7	0.6
4.5–5.0	—	0.5	0.8	1.1	1.0	1.0	1.0	0.6	0.5	0.5	0.5	0.6
5.5–6.0 (High)	—	0.9	1.1	1.0	0.7	1.1	1.1	0.6	1.1	0.1	0.4	0.5
Race/Ethnicity (2-year average): ^b												
White	—	—	0.9	1.0	1.1	1.1	1.2	1.0	0.8	0.7	0.6	0.7
African American	—	—	0.1	0.1	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.3
Hispanic	—	—	0.8	1.1	1.4	1.3	0.9	0.8	0.8	0.8	0.9	1.0

Table continued on next page.

TABLE 56 (cont.)
HEROIN WITHOUT A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	0.6	0.6	0.6	0.6	0.6	0.5	0.4	0.4	0.3	0.3	0.2	0.1	0.1	0.2	0.1	0.1	-0.1*
Gender:																	
Male	0.6	0.6	0.8	0.8	0.6	0.5	0.4	0.5	0.3	0.3	0.2	0.2	0.1	0.3	§	0.1	-0.2
Female	0.6	0.5	0.4	0.4	0.5	0.4	0.4	0.3	0.4	0.3	0.1	0.1	0.1	0.1	§	0.1	0.0
College Plans:																	
None or under 4 years	2.0	1.1	1.6	2.0	2.2	1.6	1.8	1.2	1.3	1.2	0.7	0.4	0.4	0.3	§	0.3	0.0
Complete 4 years	0.4	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.2	0.2	0.1	0.1	0.1	0.1	§	0.0	-0.1
Region:																	
Northeast	0.8	0.7	0.6	0.5	0.5	0.6	0.3	0.4	0.2	0.4	0.1	0.2	0.1	0.1	§	0.0	-0.1
Midwest	0.6	0.5	0.8	0.7	0.6	0.2	0.4	0.2	0.2	0.3	0.3	0.2	0.0	0.2	§	0.0	-0.1
South	0.6	0.6	0.6	0.4	0.5	0.6	0.5	0.3	0.5	0.3	0.1	0.1	0.2	0.2	§	0.1	-0.1
West	0.5	0.5	0.4	0.8	0.6	0.5	0.5	0.6	0.3	0.3	0.1	0.1	0.1	0.3	§	0.1	-0.2
Population Density:																	
Large MSA	0.5	0.7	0.5	0.4	0.6	0.5	0.5	0.3	0.3	0.2	0.2	0.1	0.1	0.2	§	0.1	-0.1
Other MSA	0.6	0.5	0.6	0.8	0.5	0.4	0.3	0.5	0.2	0.2	0.1	0.1	0.1	0.3	§	0.0	-0.2 s
Non-MSA	0.8	0.6	0.8	0.3	0.6	0.5	0.6	0.3	0.6	0.6	0.2	0.2	0.2	0.1	§	0.1	0.0
Parental Education: ^a																	
1.0–2.0 (Low)	1.2	0.9	1.1	1.1	0.9	0.7	1.3	0.5	0.3	0.8	0.4	0.2	0.1	0.1	§	0.0	-0.1
2.5–3.0	0.6	0.6	0.4	0.6	0.5	0.7	0.6	0.7	0.4	0.2	0.1	0.1	0.0	0.3	§	0.0	-0.3
3.5–4.0	0.5	0.6	0.8	0.7	0.5	0.4	0.4	0.1	0.3	0.4	0.1	0.2	0.2	0.1	§	0.0	0.0
4.5–5.0	0.5	0.5	0.5	0.3	0.6	0.3	0.3	0.2	0.3	0.1	0.1	0.0	0.1	0.1	§	0.1	0.0
5.5–6.0 (High)	0.4	0.3	0.4	0.5	0.5	0.4	0.1	0.7	0.4	0.4	0.2	0.1	0.2	0.5	§	0.1	-0.4
Race/Ethnicity (2-year average): ^b																	
White	0.7	0.5	0.5	0.6	0.6	0.5	0.4	0.3	0.3	0.3	0.2	0.1	0.1	0.2	§	0.1	-0.1
African American	0.4	0.3	0.4	0.3	0.2	0.5	0.6	0.5	0.4	0.4	0.1	0.0	0.1	0.0	§	0.1	+0.1
Hispanic	0.9	0.8	0.7	0.7	0.7	0.7	0.6	0.5	0.3	0.3	0.2	0.2	0.2	0.1	§	0.0	-0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1995; *N* is one half of *N* indicated in Table S2. Data based on all forms beginning in 1996. Beginning in 2015 Heroin use with and without a needle were dropped in one form; data based on three of four forms; *N* is two thirds of *N* indicated.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 57
HEROIN WITHOUT A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months											
	1975– 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Approximate weighted N =	—	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700
Total	—	1.0	1.0	1.2	0.8	1.0	1.6	0.8	0.8	0.8	0.7	0.8
Gender:												
Male	—	1.4	1.3	1.4	1.0	1.3	1.8	1.3	0.9	0.9	0.9	1.1
Female	—	0.7	0.7	0.9	0.7	0.8	1.5	0.4	0.7	0.7	0.4	0.5
College Plans:												
None or under 4 years	—	1.4	1.7	1.6	1.3	1.5	2.7	1.3	2.2	0.8	1.5	1.7
Complete 4 years	—	0.9	0.8	1.0	0.6	1.0	1.3	0.7	0.4	0.7	0.4	0.6
Region:												
Northeast	—	0.9	1.9	1.9	1.1	1.3	0.9	1.3	1.1	0.6	1.1	1.1
Midwest	—	0.7	0.5	1.0	0.6	0.7	1.6	1.1	0.9	0.7	0.4	0.7
South	—	1.4	0.9	1.1	1.1	1.2	1.8	0.3	0.9	1.2	0.5	0.7
West	—	0.8	0.7	0.6	0.3	1.0	2.0	0.9	0.4	0.3	1.0	0.7
Population Density:												
Large MSA	—	1.3	1.1	1.2	0.9	0.9	2.5	1.0	0.6	0.8	0.8	0.7
Other MSA	—	0.8	1.2	1.1	0.9	1.2	1.2	0.8	1.1	0.7	0.9	0.7
Non-MSA	—	1.0	0.4	1.2	0.4	0.8	1.4	0.7	0.6	1.0	0.2	1.0
Parental Education: ^a												
1.0–2.0 (Low)	—	1.0	0.8	1.7	0.5	2.0	2.7	1.5	1.8	1.0	1.1	1.1
2.5–3.0	—	1.1	0.6	0.8	1.1	1.1	1.8	0.5	0.9	0.9	0.8	1.0
3.5–4.0	—	0.9	1.1	1.3	0.9	0.8	1.7	0.8	1.0	0.7	0.6	0.9
4.5–5.0	—	1.2	1.2	1.2	0.8	1.1	1.6	1.1	0.2	0.8	0.7	0.5
5.5–6.0 (High)	—	0.6	1.0	1.0	0.5	1.0	0.6	0.8	0.8	0.7	0.3	0.5
Race/Ethnicity (2-year average): ^b												
White	—	—	1.1	1.3	1.2	1.1	1.4	1.3	1.0	0.8	0.7	0.6
African American	—	—	0.2	0.3	0.3	0.1	0.4	0.4	0.4	0.6	0.7	0.5
Hispanic	—	—	0.9	0.6	0.4	0.6	1.8	1.3	0.5	0.7	0.7	0.9

Table continued on next page.

TABLE 57 (cont.)
HEROIN WITHOUT A NEEDLE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	0.6	1.0	0.5	0.6	0.8	0.7	0.4	0.4	0.5	0.4	0.3	0.2	0.2	0.2	0.1	0.1	-0.1*
Gender:																	
Male	0.8	1.2	0.7	0.8	1.1	0.9	0.5	0.5	0.8	0.5	0.2	0.1	0.2	0.2	§	0.1	-0.1
Female	0.3	0.6	0.3	0.4	0.4	0.5	0.2	0.3	0.2	0.1	0.3	0.2	0.1	0.1	§	0.0	-0.1
College Plans:																	
None or under 4 years	1.2	2.6	1.0	0.7	1.8	1.1	0.8	0.8	0.7	0.8	0.5	0.4	0.6	0.6	§	0.3	-0.3
Complete 4 years	0.4	0.7	0.3	0.5	0.4	0.6	0.3	0.4	0.3	0.2	0.2	0.2	0.1	0.1	§	0.0	-0.1
Region:																	
Northeast	1.0	1.1	0.7	0.7	1.3	0.8	0.2	0.4	0.9	0.3	0.1	0.0	0.2	0.4	§	0.0	-0.4
Midwest	0.2	0.9	0.3	0.7	0.7	0.9	0.3	0.3	0.5	0.6	0.2	0.0	0.4	0.1	§	0.0	-0.1
South	0.7	1.0	0.3	0.6	0.7	0.4	0.1	0.4	0.3	0.3	0.3	0.4	0.2	0.3	§	0.1	-0.2
West	0.6	0.9	0.8	0.7	0.6	0.9	0.8	0.8	0.4	0.3	0.3	0.2	0.1	0.1	§	0.2	+0.2
Population Density:																	
Large MSA	0.8	0.9	0.5	0.7	1.1	0.8	0.2	0.2	0.4	0.3	0.3	0.1	0.3	0.4	§	0.2	-0.1
Other MSA	0.4	0.9	0.6	0.6	0.8	0.7	0.4	0.5	0.6	0.4	0.2	0.3	0.1	0.2	§	0.1	-0.1
Non-MSA	0.5	1.2	0.3	0.5	0.5	0.4	0.5	0.6	0.2	0.4	0.3	0.2	0.4	0.3	§	0.0	-0.3
Parental Education: ^a																	
1.0–2.0 (Low)	1.2	2.2	0.7	1.1	0.8	1.0	0.1	1.0	0.9	1.0	0.6	0.4	0.3	0.7	§	0.0	-0.7
2.5–3.0	0.5	0.9	0.5	0.7	0.9	0.4	0.5	0.7	0.3	0.4	0.2	0.1	0.4	0.2	§	0.0	-0.2
3.5–4.0	0.5	0.8	0.3	0.9	1.1	1.0	0.2	0.3	0.2	0.2	0.2	0.3	0.1	0.4	§	0.2	-0.2
4.5–5.0	0.4	0.8	0.6	0.3	0.5	0.7	0.5	0.3	0.7	0.3	0.1	0.1	0.1	0.1	§	0.1	+0.1
5.5–6.0 (High)	0.7	0.7	0.6	0.2	0.5	0.4	0.2	0.3	0.5	0.3	0.3	0.2	0.2	0.1	§	0.2	+0.1
Race/Ethnicity (2-year average): ^b																	
White	0.6	0.8	0.8	0.6	0.7	0.8	0.6	0.3	0.3	0.3	0.2	0.2	0.1	0.1	§	0.0	-0.1
African American	0.4	0.4	0.4	0.3	0.7	0.8	0.4	0.3	0.7	0.8	0.3	0.3	0.4	0.8	§	0.1	-0.7 s
Hispanic	1.0	1.1	0.7	0.4	0.6	0.7	0.3	0.3	0.6	0.6	0.4	0.2	0.2	0.2	§	0.1	-0.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on three of six forms; N is three sixths of N indicated in Table S3.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 58
NARCOTICS OTHER THAN HEROIN: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
<i>Approximate weighted N =</i>	<i>9,400</i>	<i>15,400</i>	<i>17,100</i>	<i>17,800</i>	<i>15,500</i>	<i>15,900</i>	<i>17,500</i>	<i>17,700</i>	<i>16,300</i>	<i>15,900</i>	<i>16,000</i>	<i>15,200</i>	<i>16,300</i>	<i>16,300</i>	<i>16,700</i>	<i>15,200</i>
Total	5.7	5.7	6.4	6.0	6.2	6.3	5.9	5.3	5.1	5.2	5.9	5.2	5.3	4.6	4.4	4.5
Gender:																
Male	6.6	6.8	7.3	6.9	7.3	7.1	6.5	6.0	6.0	6.2	6.8	5.9	5.6	5.1	4.9	5.0
Female	4.8	4.7	5.4	5.1	5.1	5.4	5.3	4.6	4.2	4.2	5.1	4.6	4.9	4.1	3.8	3.9
College Plans:																
None or under 4 years	—	6.8	8.0	6.8	7.3	7.4	7.2	6.1	6.1	6.1	6.6	6.7	6.1	4.8	5.3	5.7
Complete 4 years	—	4.6	4.7	4.9	5.0	5.1	4.8	4.6	4.3	4.3	5.4	4.3	4.8	4.6	3.9	4.0
Region:																
Northeast	6.1	6.5	6.6	6.8	7.0	5.7	7.2	5.6	5.6	6.7	7.3	5.7	6.0	3.7	4.7	4.1
Midwest	6.2	6.2	7.5	6.7	6.1	7.6	6.2	5.5	5.3	4.8	6.3	5.8	5.2	4.4	5.7	4.6
South	4.9	5.0	5.2	4.5	5.2	5.0	4.1	4.5	4.4	4.5	3.8	4.2	4.3	4.7	3.2	4.1
West	5.4	5.0	6.0	6.7	7.1	6.8	7.2	6.2	5.2	5.3	7.1	5.4	6.1	5.7	4.9	5.3
Population Density:																
Large MSA	7.3	6.7	6.7	6.9	7.3	6.9	6.9	5.2	6.0	5.2	6.0	4.8	5.2	4.0	4.1	3.8
Other MSA	5.5	6.1	6.3	5.9	6.3	7.0	6.3	5.7	5.3	5.1	6.4	5.6	5.3	5.2	4.9	4.6
Non-MSA	4.8	4.6	6.2	5.4	5.3	4.8	4.8	4.9	4.1	5.2	5.2	5.0	5.2	4.4	3.8	4.8
Parental Education: ^b																
1.0–2.0 (Low)	5.4	5.0	5.1	5.0	5.2	5.2	4.8	4.8	4.8	4.7	4.5	4.7	4.1	3.9	3.6	3.8
2.5–3.0	5.1	5.9	6.4	6.2	5.9	5.8	5.6	4.9	5.0	5.2	5.5	5.0	4.4	4.3	4.0	4.1
3.5–4.0	4.2	6.3	6.7	6.0	6.3	6.9	6.6	5.2	4.5	5.1	6.5	6.0	5.6	4.3	4.6	4.6
4.5–5.0	6.4	6.3	6.6	6.4	6.7	7.0	6.3	6.4	6.0	5.6	6.4	4.8	5.4	5.4	4.2	4.7
5.5–6.0 (High)	6.5	6.5	7.9	6.1	7.8	6.8	6.8	7.1	5.3	4.9	6.8	5.4	7.8	5.6	6.4	5.7
Race/Ethnicity (2-year average): ^c																
White	—	—	6.6	6.7	6.6	6.8	6.7	6.2	5.8	5.7	6.3	6.3	6.0	5.8	5.3	5.2
African American	—	—	2.2	2.0	1.8	1.7	1.9	1.8	1.7	1.6	1.6	1.7	1.6	1.5	1.5	1.4
Hispanic	—	—	3.8	3.5	3.5	3.7	4.3	4.1	4.0	4.2	3.6	3.0	2.4	2.2	2.5	2.4

Table continued on next page.

TABLE 58 (cont.)
NARCOTICS OTHER THAN HEROIN: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002 ^d	2003 ^d	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	3.5	3.3	3.6	3.8	4.7	5.4	6.2	6.3	6.7	7.0	6.7†	9.4	9.3	9.5	9.0	9.0
Gender:																
Male	3.9	3.3	3.6	4.3	5.6	6.4	7.1	7.4	8.2	8.0	8.0†	11.6	10.7	10.9	10.7	10.3
Female	3.1	3.3	3.3	3.4	3.8	4.4	5.4	5.1	5.2	5.9	5.6†	7.4	7.8	8.1	7.4	7.8
College Plans:																
None or under 4 years	3.8	4.3	4.2	4.9	5.6	7.0	8.2	8.4	7.0	7.8	7.5†	12.4	12.3	12.4	12.2	12.4
Complete 4 years	3.5	3.0	3.3	3.5	4.4	4.9	5.7	5.5	6.6	6.6	6.4†	8.8	8.4	8.7	8.3	8.1
Region:																
Northeast	3.2	3.7	4.6	3.5	4.3	6.1	7.8	6.5	6.2	6.7	7.2†	10.6	9.3	9.9	8.3	10.1
Midwest	4.2	3.6	3.2	4.7	5.2	6.0	6.1	6.5	7.5	7.2	7.8†	8.8	9.1	8.7	10.7	9.9
South	2.7	2.7	3.2	3.8	4.5	5.1	6.1	6.5	7.5	6.6	5.7†	9.3	8.0	9.1	8.6	7.9
West	4.4	3.5	4.0	3.1	4.7	4.1	4.7	5.2	4.7	7.5	6.5†	9.5	11.4	11.1	8.3	8.9
Population Density:																
Large MSA	3.3	3.5	3.1	4.1	4.8	4.6	4.6	5.2	5.4	7.0	8.4†	8.9	7.4	8.1	8.4	8.1
Other MSA	3.9	3.1	3.7	3.7	4.7	5.4	7.2	6.8	7.4	7.2	5.6†	10.3	10.3	10.7	9.3	9.0
Non-MSA	3.1	3.6	3.7	3.6	4.7	6.0	6.0	6.5	6.8	6.6	6.9†	8.5	9.7	8.9	9.1	10.4
Parental Education: ^b																
1.0–2.0 (Low)	3.8	3.5	3.8	3.0	4.0	4.5	4.7	4.7	4.6	4.6	5.1†	6.2	7.9	6.8	5.2	6.1
2.5–3.0	3.2	3.5	2.9	3.8	4.2	5.6	5.3	5.9	5.9	6.4	6.4†	8.9	9.3	8.8	9.5	9.5
3.5–4.0	3.7	3.2	3.7	3.4	4.4	5.5	6.5	6.8	7.2	7.3	7.1†	10.4	9.8	10.8	10.5	10.3
4.5–5.0	3.6	3.4	3.7	4.3	5.5	5.4	6.8	6.2	7.4	7.9	7.0†	10.3	9.5	10.5	9.3	9.1
5.5–6.0 (High)	4.1	3.2	4.5	4.8	5.5	5.6	7.6	6.6	7.4	7.5	7.0†	8.7	8.8	8.8	7.6	8.0
Race/Ethnicity (2-year average): ^c																
White	4.7	4.1	4.1	4.3	5.0	5.9	7.1	7.6	7.7	8.3	8.5	10.3†	10.2 ^f	11.2	11.2	11.2
African American	1.1	0.9	1.0	1.5	1.4	1.2	1.8	2.4	2.0	1.7	1.8	1.9†	2.1 ^f	2.7	2.5	2.4
Hispanic	2.3	2.1	2.3	2.2	2.5	3.7	3.1	2.8	3.6	4.6	4.5	4.3†	5.2 ^f	5.5	5.3	5.0

Table continued on next page.

TABLE 58 (cont.)
NARCOTICS OTHER THAN HEROIN: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007 ^a	2008	2009	2010	2011	2012	2013 ^a	2014	2015	2016	2017	2018	2019	2020	2021	
	<i>Approximate weighted N =</i>															
	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	9.2	9.1	9.2	8.7	8.7	7.9	7.1	6.1	5.4	4.8	4.2	3.4	2.7	2.1	1.0	-1.7* sss
Gender:																
Male	10.9	10.6	10.7	9.9	9.6	9.3	8.4	7.2	6.1	5.6	5.3	4.4	3.4	§	0.9	-2.6 sss
Female	7.6	7.9	7.9	7.4	7.8	6.5	5.6	5.0	4.6	3.8	3.2	2.2	1.8	§	1.0	-0.8 ss
College Plans:																
None or under 4 years	12.9	13.1	11.0	12.3	13.2	9.7	10.1	7.3	6.8	4.9	6.0	4.6	3.7	§	1.0	-2.7 sss
Complete 4 years	8.3	8.3	8.8	7.8	7.8	7.4	6.3	5.8	5.0	4.6	3.8	3.0	2.3	§	1.0	-1.4 sss
Region:																
Northeast	9.3	9.7	9.3	9.2	8.3	6.3	7.1	4.0	3.9	3.1	3.0	2.8	1.6	§	1.1	-0.5
Midwest	10.5	10.0	11.6	9.0	9.7	9.1	6.8	6.3	5.0	5.6	3.7	3.5	2.5	§	0.9	-1.5 sss
South	8.3	8.3	8.0	8.4	7.3	6.1	7.1	6.4	5.5	4.8	4.9	3.2	2.8	§	1.1	-1.7 sss
West	9.2	9.0	8.2	8.4	10.2	10.0	7.4	7.1	6.6	5.0	4.5	4.0	3.6	§	0.7	-2.9 sss
Population Density:																
Large MSA	7.9	7.9	7.9	8.8	7.6	8.4	6.4	4.6	4.2	4.4	3.4	2.8	2.5	§	1.1	-1.4 sss
Other MSA	9.3	9.0	9.4	9.0	9.0	7.4	7.9	7.4	5.9	5.0	4.6	3.6	2.8	§	1.0	-1.8 sss
Non-MSA	10.7	10.7	10.6	7.7	9.6	8.3	6.1	4.8	6.1	4.7	4.9	4.0	2.6	§	0.8	-1.9 sss
Parental Education: ^b																
1.0–2.0 (Low)	7.1	7.1	6.1	5.2	6.6	7.2	5.5	5.8	5.7	4.0	3.3	3.2	2.8	§	0.6	-2.2 sss
2.5–3.0	8.8	9.2	9.9	9.4	8.5	9.0	7.6	6.8	4.7	4.0	5.1	3.9	2.9	§	0.8	-2.1 sss
3.5–4.0	9.5	9.9	9.9	10.0	10.1	8.4	8.3	6.1	6.1	5.1	4.6	3.7	2.9	§	0.8	-2.1 sss
4.5–5.0	9.9	9.0	9.9	8.1	8.1	8.0	6.6	5.8	5.3	5.2	3.7	3.1	2.2	§	1.0	-1.2 s
5.5–6.0 (High)	9.5	9.7	8.2	8.3	9.2	6.0	6.2	6.3	5.7	5.7	4.6	3.1	2.7	§	1.5	-1.3
Race/Ethnicity (2-year average): ^c																
White	11.5	11.7	11.7	11.1	10.6	10.0	8.5	7.0	6.0	5.6	5.0	4.2	3.3	§	1.3	-1.9 sss
African American	2.6	2.7	2.8	4.0	4.1	3.4	4.0	4.7	4.6	3.7	3.2	2.8	1.7	§	1.7	0.0
Hispanic	4.5	4.2	4.7	5.1	5.4	5.5	5.7	5.4	4.1	3.9	3.8	2.9	2.5	§	1.3	-1.2 ss

Table continued on next page.

TABLE 58 (cont.)
NARCOTICS OTHER THAN HEROIN: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dIn 2002 the question text was changed in half of the questionnaire forms. In the list of examples of narcotics other than heroin, Talwin, laudanum, and paregoric were replaced with Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; N is one half of N indicated. In 2003 the remaining forms were changed to the new wording. Beginning in 2003, the data are based on all forms.

^eIn 2007, in the list of narcotics other than heroin, Dilaudid was replaced with Ultram.

^fThe 2003 data comprise half of the 2002 sample data double-weighted and all of the 2003 sample data.

^gIn 2013 the list of examples was changed on one form: MS Contin, Roxycodone, Hydrocodone (Lortab, Lorcet, Norco), Suboxone, Tylox, and Tramadol were added to the list. An examination of the data did not show any effect from the wording change.

TABLE 59
OXYCONTIN: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months																				2019– 2021 change	
	1991– 2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
Approximate weighted N =	—	15,100	16,500	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	1.3	1.7	1.7	1.8	2.6	1.8	2.1	2.0	2.1	1.8	1.6	2.0	1.0	0.8	0.9	0.8	0.8	1.2	0.9	0.8	-0.4*
Gender:																						
Male	—	1.9	1.6	1.8	1.9	2.8	1.7	2.3	2.7	2.1	1.7	1.8	1.8	0.9	1.2	1.2	1.0	0.8	1.4	§	0.4	-1.0 ss
Female	—	0.9	1.7	1.6	1.7	2.2	2.0	1.8	1.4	2.2	1.8	1.2	2.2	1.2	0.6	0.7	0.6	0.7	1.0	§	1.4	+0.4
College Plans:																						
None or under 4 years	—	4.3	7.5	6.9	6.8	8.8	5.1	6.1	8.4	6.4	7.7	8.2	7.6	3.8	3.4	3.3	3.2	2.5	3.8	§	1.8	-2.0
Complete 4 years	—	1.0	1.1	1.2	1.2	1.9	1.5	1.8	1.6	1.8	1.3	1.1	1.6	0.8	0.6	0.7	0.6	0.5	0.9	§	0.7	-0.2
Region:																						
Northeast	—	1.0	1.2	1.1	1.7	1.6	1.1	0.8	1.2	1.6	1.3	0.8	3.0	0.8	0.3	0.5	0.2	0.6	0.5	§	0.0	-0.5
Midwest	—	1.7	1.7	1.3	2.4	3.2	2.0	1.4	2.2	2.2	1.9	0.9	1.0	0.6	0.4	0.8	0.7	0.2	0.8	§	1.6	+0.8
South	—	1.5	2.1	2.2	2.0	2.4	2.0	2.9	2.1	2.5	1.8	2.6	1.9	1.1	1.4	1.2	0.9	0.9	1.8	§	1.1	-0.6
West	—	0.9	1.3	1.8	1.0	3.1	1.6	2.3	2.2	1.8	2.2	1.0	2.4	1.5	0.7	0.7	1.2	1.2	1.3	§	0.3	-1.0 s
Population Density:																						
Large MSA	—	1.2	1.1	0.9	1.1	1.3	0.9	1.7	0.7	1.3	0.8	1.8	1.4	0.8	0.6	0.9	0.5	0.8	1.4	§	0.8	-0.6
Other MSA	—	1.4	1.7	2.6	2.3	3.1	2.2	2.2	2.3	2.3	2.7	1.3	2.0	1.2	1.3	0.8	1.0	0.9	1.2	§	1.1	-0.1
Non-MSA	—	1.5	2.5	1.1	1.8	3.2	2.3	2.6	2.8	2.9	1.5	1.7	2.7	0.9	0.3	1.0	0.7	0.4	0.9	§	0.3	-0.6
Parental Education: ^b																						
1.0–2.0 (Low)	—	2.9	4.0	3.5	3.7	3.6	2.0	4.0	6.1	3.0	3.8	1.4	1.9	1.6	1.2	1.9	2.5	0.8	4.3	§	0.4	-3.9 s
2.5–3.0	—	2.0	2.3	2.1	2.9	3.4	2.2	2.3	2.1	2.7	2.6	2.7	1.7	1.3	1.1	1.2	0.3	0.7	2.1	§	0.4	-1.6 s
3.5–4.0	—	1.0	1.9	2.2	2.5	2.4	2.2	2.3	1.7	2.5	2.4	1.2	3.0	1.0	0.9	1.0	1.2	0.6	0.9	§	2.5	+1.6
4.5–5.0	—	0.9	0.7	1.3	0.7	2.0	1.6	1.7	1.7	1.3	1.1	1.2	1.9	1.0	0.7	0.5	0.4	1.1	0.5	§	0.3	-0.2
5.5–6.0 (High)	—	1.3	1.6	0.7	0.8	2.2	0.8	1.7	1.0	1.1	0.5	1.4	1.4	0.5	0.3	0.5	0.6	0.3	0.4	§	0.5	+0.1
Race/Ethnicity (2-year average): ^c																						
White	—	—	1.6	1.7	1.7	2.3	2.5	2.3	2.2	2.1	1.9	1.5	1.3	1.2	1.1	0.7	0.5	0.6	0.7	§	0.9	+0.3
African American	—	—	0.9	0.6	1.0	1.4	1.2	1.4	1.6	1.5	1.3	1.5	1.8	1.0	0.6	1.5	1.8	1.4	1.6	§	1.3	-0.3
Hispanic	—	—	1.3	2.8	2.6	2.4	2.3	1.5	1.8	2.2	2.6	2.1	1.9	1.9	1.0	1.0	0.9	0.7	1.3	§	0.2	-1.1 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 60
OXYCONTIN: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months																				2019– 2021 change	
	1991– 2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
Approximate weighted N =	—	14,300	15,800	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	3.0	3.6	3.5	3.2	3.8	3.9	3.6	5.1	4.6	3.9	3.0	3.4	3.0	2.6	2.1	2.2	2.2	2.0	1.0	0.9	-1.1* ss
Gender:																						
Male	—	3.6	4.3	3.8	3.4	3.7	4.2	3.8	5.6	4.9	3.8	3.2	4.7	2.8	2.9	2.1	1.9	2.5	2.8	§	0.9	-1.9 s
Female	—	2.4	2.9	3.1	3.0	4.0	3.5	3.5	4.5	4.1	4.0	2.8	2.2	2.8	2.3	2.1	2.4	1.7	1.3	§	0.8	-0.5
College Plans:																						
None or under 4 years	—	6.5	10.8	7.2	8.0	11.0	8.8	10.2	12.6	13.9	11.8	9.1	6.1	6.9	7.5	6.5	4.6	3.7	3.2	§	1.0	-2.2 s
Complete 4 years	—	2.5	2.5	3.0	2.6	2.9	3.2	2.9	4.2	3.5	2.9	2.4	3.1	2.5	2.1	1.7	1.9	1.9	1.8	§	0.8	-1.0 s
Region:																						
Northeast	—	3.6	3.7	4.6	3.8	3.1	2.8	3.2	3.6	4.4	4.5	3.0	3.1	1.0	2.4	1.3	1.2	1.6	1.1	§	1.7	+0.6
Midwest	—	2.1	2.9	2.5	3.5	4.1	4.3	3.9	5.8	4.9	4.2	2.0	4.2	2.3	2.7	1.9	1.8	1.9	1.3	§	0.9	-0.4
South	—	3.7	4.2	4.8	3.6	4.4	4.7	4.1	5.4	4.8	4.0	3.5	2.7	3.6	2.5	2.6	3.0	2.1	2.1	§	0.8	-1.3
West	—	2.6	3.5	1.9	1.6	3.3	3.3	3.0	5.2	4.2	2.9	3.4	3.7	4.2	3.0	2.4	2.1	3.1	3.2	§	0.3	-2.9 ss
Population Density:																						
Large MSA	—	2.0	2.3	2.1	2.8	3.4	3.5	2.5	4.5	4.2	2.9	2.3	3.2	2.6	2.0	1.5	1.9	2.3	0.8	§	1.0	+0.3
Other MSA	—	3.3	3.6	3.7	2.7	3.3	3.5	4.1	5.8	4.3	3.6	2.6	3.4	2.5	2.8	2.0	2.1	2.1	2.6	§	0.7	-1.9 s
Non-MSA	—	3.7	5.7	5.1	4.9	5.8	5.4	4.1	4.4	5.7	5.7	4.9	3.6	4.7	3.2	3.6	2.9	2.2	2.6	§	1.0	-1.7 s
Parental Education: ^b																						
1.0–2.0 (Low)	—	6.5	5.9	3.7	3.3	6.9	3.9	3.1	6.5	7.9	5.5	4.4	5.2	2.5	4.3	1.4	2.3	5.5	1.4	§	0.3	-1.1
2.5–3.0	—	3.8	3.9	4.0	4.8	5.3	5.9	5.1	6.7	6.9	5.4	4.1	4.4	3.1	2.8	2.3	2.7	3.7	2.0	§	1.2	-0.8
3.5–4.0	—	2.0	4.1	4.1	3.0	4.3	3.5	3.6	5.5	4.0	4.3	3.9	3.2	3.8	4.0	2.8	2.3	1.4	2.9	§	1.3	-1.7
4.5–5.0	—	1.7	2.9	2.6	2.6	2.3	3.4	3.2	3.9	2.9	2.4	1.7	2.7	2.4	1.6	1.9	2.6	1.2	1.8	§	0.7	-1.1
5.5–6.0 (High)	—	4.4	2.8	2.8	2.6	2.4	2.8	3.5	4.6	3.6	3.2	2.2	2.8	2.1	1.7	1.9	0.9	0.8	2.1	§	0.8	-1.3
Race/Ethnicity (2-year average): ^c																						
White	—	—	3.6	4.1	4.1	4.1	4.6	4.7	5.0	5.3	4.4	3.5	3.5	3.6	3.0	2.7	2.3	1.7	1.7	§	1.1	-0.6
African American	—	—	2.3	2.0	1.1	1.5	1.8	1.5	1.7	2.4	2.9	1.8	1.5	2.1	2.4	1.7	1.6	1.6	1.8	§	2.0	+0.1
Hispanic	—	—	2.5	2.9	2.3	1.8	1.9	1.8	2.9	4.0	4.4	3.6	1.9	1.9	2.0	1.7	2.2	4.0	3.1	§	0.1	-3.1 sss

Source: The Monitoring the Future study, the University of Michigan.

Notes: Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 61
OXYCONTIN: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months																				2019– 2021 change	
	1975– 2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
<i>Approximate weighted N =</i>	—	12,900	14,600	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	4.0	4.5	5.0	5.5	4.3	5.2	4.7	4.9	5.1	4.9	4.3	3.6	3.3	3.7	3.4	2.7	2.3	1.7	2.4	0.9	-0.8* s
Gender:																						
Male	—	5.6	6.2	6.1	7.4	5.3	6.3	5.3	6.5	5.9	5.9	5.1	4.7	4.3	4.2	3.6	3.4	2.4	2.3	§	1.0	-1.4 ss
Female	—	2.6	2.8	3.4	3.5	2.9	4.0	4.0	3.4	3.8	3.8	3.1	2.5	2.2	3.0	2.6	1.8	1.7	1.0	§	0.6	-0.5
College Plans:																						
None or under 4 years	—	7.0	8.8	8.2	9.9	7.4	7.8	7.6	6.7	10.2	9.7	5.2	6.0	4.5	4.3	5.3	4.5	3.8	3.5	§	0.8	-2.7 sss
Complete 4 years	—	3.2	3.1	3.8	4.3	3.3	4.5	4.1	4.5	3.9	3.9	4.0	2.8	2.9	3.4	2.9	2.1	1.7	1.2	§	0.9	-0.2
Region:																						
Northeast	—	5.0	5.5	4.6	6.0	5.5	5.5	4.6	5.0	6.3	4.4	3.6	3.6	3.1	2.8	2.1	2.0	1.3	0.7	§	0.5	-0.2
Midwest	—	5.0	4.5	4.2	6.0	4.3	5.1	4.4	5.4	5.0	4.3	4.5	3.0	3.4	3.8	3.6	2.1	2.0	1.4	§	0.8	-0.6
South	—	3.6	4.0	5.9	6.3	4.0	5.2	4.9	4.3	4.8	5.0	4.3	4.1	4.4	3.6	3.6	3.1	2.8	2.1	§	1.4	-0.7
West	—	2.6	4.4	4.7	3.0	3.6	5.1	4.9	5.0	4.7	5.9	4.5	3.3	1.6	4.2	3.7	3.1	2.5	2.2	§	0.7	-1.5 s
Population Density:																						
Large MSA	—	3.7	2.8	4.6	4.8	4.4	4.2	3.9	4.2	5.4	3.9	4.6	2.8	3.6	2.6	3.7	2.0	2.5	1.3	§	0.7	-0.6
Other MSA	—	3.8	5.3	5.2	5.1	3.6	5.4	5.0	5.2	4.8	5.2	4.0	4.2	3.4	4.3	3.6	3.4	2.0	1.8	§	1.3	-0.5
Non-MSA	—	4.7	5.2	4.9	7.1	5.4	6.2	5.1	5.2	5.3	6.0	4.6	3.1	2.7	3.9	2.7	2.5	2.6	2.3	§	0.6	-1.7 s
Parental Education: ^b																						
1.0–2.0 (Low)	—	6.3	6.9	3.7	5.5	4.2	4.4	3.8	4.9	4.9	3.4	4.3	6.0	3.6	3.2	4.2	3.7	2.6	3.3	§	1.1	-2.2
2.5–3.0	—	5.3	6.8	4.8	6.3	4.4	5.6	4.5	4.6	5.5	4.8	4.2	3.4	3.0	3.3	2.9	3.4	3.0	1.9	§	0.5	-1.4 ss
3.5–4.0	—	3.9	3.4	6.5	6.3	5.1	5.0	6.0	5.8	5.4	5.7	4.6	3.4	3.4	3.4	2.4	2.8	2.2	1.7	§	0.8	-0.9 s
4.5–5.0	—	2.2	3.6	3.1	4.2	3.4	5.3	4.3	4.8	4.2	4.4	4.4	3.5	2.8	3.9	3.6	1.6	1.6	1.3	§	1.0	-0.3
5.5–6.0 (High)	—	4.0	2.5	4.3	4.8	3.5	4.3	3.8	4.1	4.7	5.0	3.5	2.2	4.1	4.3	5.0	2.8	1.3	0.8	§	1.6	+0.9
Race/Ethnicity (2-year average): ^c																						
White	—	—	4.7	5.1	5.8	5.4	5.5	6.0	5.8	5.7	5.8	5.2	4.0	3.3	3.7	3.7	3.0	2.5	1.9	§	0.9	-1.0 sss
African American	—	—	2.5	2.4	2.0	2.2	2.4	2.1	2.0	3.1	3.2	2.8	3.4	4.0	3.9	3.3	2.5	1.7	1.3	§	2.5	+1.2
Hispanic	—	—	2.5	2.2	3.0	3.7	3.2	2.1	2.9	3.5	3.0	3.2	3.4	2.7	2.1	2.9	3.3	2.1	1.6	§	2.1	+0.5

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms in 2002–2005; *N* is two sixths of *N* indicated in Table S3. Data based on three of six forms beginning in 2006; *N* is three sixths of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 62
VICODIN:^a Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months																						
	1991–	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019–
	2001–																						2021
	Approximate weighted N =	—	15,100	16,500	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	change
Total	—	2.5	2.8	2.5	2.6	3.0	2.7	2.9	2.5	2.7	2.1	1.3	1.4	1.0	0.9	0.8	0.7	0.6	0.9	0.5	0.6	0.6	-0.3*
Gender:																							
Male	—	2.7	2.8	2.4	2.9	2.6	2.1	2.9	3.0	2.8	2.1	0.8	1.1	0.9	0.8	1.2	0.9	0.8	1.2	§	0.4	0.4	-0.8 s
Female	—	2.4	2.8	2.6	2.2	3.4	3.2	2.7	2.2	2.6	1.9	1.7	1.6	1.1	1.0	0.5	0.4	0.3	0.7	§	0.8	0.8	+0.1
College Plans:																							
None or under 4 years	—	5.9	6.3	7.1	7.3	9.5	8.1	7.0	6.5	7.2	6.4	3.0	4.0	1.9	2.2	2.9	2.7	1.9	2.3	§	1.4	1.4	-0.9
Complete 4 years	—	2.1	2.4	2.0	1.9	2.4	2.1	2.5	2.3	2.2	1.8	1.2	1.2	0.9	0.7	0.6	0.4	0.4	0.7	§	0.4	0.4	-0.3
Region:																							
Northeast	—	1.5	1.1	1.0	1.5	1.9	1.4	1.0	1.8	2.8	1.7	0.7	1.6	0.3	0.3	0.6	0.1	0.5	0.5	§	0.0	0.0	-0.5
Midwest	—	3.3	3.9	3.2	3.3	3.9	3.6	3.1	3.5	3.4	2.8	1.8	1.0	1.4	1.0	0.8	1.0	0.0	0.9	§	0.5	0.5	-0.4
South	—	2.5	2.1	2.1	2.3	2.2	2.1	2.9	1.7	1.8	1.5	1.5	1.1	0.6	1.0	0.9	0.5	0.8	1.2	§	1.0	1.0	-0.2
West	—	2.6	4.2	3.6	3.1	4.4	3.5	3.7	3.3	3.2	3.0	1.1	2.0	1.8	1.0	1.0	0.9	0.8	0.8	§	0.4	0.4	-0.3
Population Density:																							
Large MSA	—	2.6	2.4	1.8	2.5	2.4	1.6	2.7	1.9	1.8	1.2	1.8	1.3	0.8	0.7	0.9	0.4	0.6	1.4	§	0.5	0.5	-0.9
Other MSA	—	3.0	3.1	3.0	2.6	3.3	3.4	3.0	2.6	3.2	2.7	1.0	1.4	1.2	1.2	0.8	1.0	0.6	0.8	§	0.8	0.8	-0.1
Non-MSA	—	1.6	2.5	2.4	2.6	3.3	2.8	2.8	3.2	2.8	2.4	1.6	1.5	1.0	0.4	0.9	0.2	0.5	0.4	§	0.4	0.4	-0.1
Parental Education: ^b																							
1.0–2.0 (Low)	—	3.3	3.3	4.9	3.9	4.8	3.1	3.4	5.4	3.9	4.0	1.7	1.7	1.5	2.8	1.7	1.4	0.6	2.2	§	0.8	0.8	-1.4
2.5–3.0	—	3.9	3.4	4.1	3.8	4.0	3.0	3.3	3.8	3.3	3.0	1.2	1.1	0.9	1.1	0.7	0.5	0.1	1.3	§	0.0	0.0	-1.3
3.5–4.0	—	3.4	2.9	3.2	2.9	2.7	3.1	2.8	2.5	1.6	2.8	1.3	2.3	1.5	0.6	1.1	0.6	0.6	0.6	§	1.7	1.7	+1.1
4.5–5.0	—	1.4	2.1	1.1	1.9	2.6	2.4	3.2	1.5	2.7	1.4	1.3	1.4	0.8	0.3	0.7	0.4	0.5	0.7	§	0.3	0.3	-0.4
5.5–6.0 (High)	—	1.3	2.1	0.9	1.6	2.2	1.8	2.3	2.1	2.7	0.9	1.0	0.8	0.8	0.3	0.3	0.4	0.5	0.6	§	0.6	0.6	0.0
Race/Ethnicity (2-year average): ^c																							
White	—	—	3.0	2.7	2.4	2.9	3.2	3.2	2.8	2.6	2.5	1.8	1.2	0.9	0.7	0.7	0.6	0.4	0.3	§	0.6	0.6	+0.3
African American	—	—	1.4	1.4	1.6	1.7	1.6	1.6	1.9	2.0	1.3	1.2	1.3	0.9	0.5	1.0	1.1	1.5	1.8	§	0.8	0.8	-1.0
Hispanic	—	—	2.1	3.3	3.6	2.8	2.6	2.1	2.5	3.0	2.8	2.1	1.9	1.8	1.2	0.8	0.6	0.5	0.7	§	0.3	0.3	-0.5

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 63
VICODIN:^a Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months																				2019– 2021 change	
	1991– 2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	14,300	15,800	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	6.9	7.2	6.2	5.9	7.0	7.2	6.7	8.1	7.7	5.9	4.4	4.6	3.4	2.5	1.7	1.5	1.1	1.1	0.9	0.5	-0.6* s
Gender:																						
Male	—	7.4	8.4	6.2	5.5	7.0	7.5	7.1	9.2	8.1	6.0	4.5	5.1	2.9	2.7	1.8	1.4	1.4	1.3	§	0.3	-1.0 ss
Female	—	6.1	6.1	6.0	6.2	6.9	6.8	6.4	7.0	7.0	5.8	4.4	4.1	3.9	2.2	1.8	1.5	0.7	0.8	§	0.5	-0.3
College Plans:																						
None or under 4 years	—	13.4	15.8	11.6	13.2	17.2	15.2	16.5	17.6	16.9	14.1	8.8	9.3	7.1	7.8	4.9	3.7	3.0	1.4	§	0.6	-0.8
Complete 4 years	—	5.7	5.8	5.4	4.8	5.6	6.1	5.6	7.1	6.5	4.9	4.0	4.1	3.0	2.0	1.4	1.2	0.8	1.0	§	0.4	-0.6 s
Region:																						
Northeast	—	6.2	5.6	6.4	5.7	4.9	6.5	5.3	5.1	5.6	4.6	3.3	3.5	2.4	1.9	1.8	1.0	0.8	0.5	§	0.7	+0.1
Midwest	—	7.2	9.4	6.4	7.1	8.7	8.8	8.4	10.5	9.3	8.1	4.6	5.8	4.4	2.9	2.6	1.3	1.6	0.8	§	0.6	-0.2
South	—	5.3	5.0	5.0	5.0	4.2	4.5	5.2	5.6	6.0	4.3	3.4	3.1	2.7	1.8	0.7	1.5	0.7	1.1	§	0.3	-0.8 s
West	—	10.0	9.1	7.1	5.8	11.1	9.4	7.9	12.1	10.1	6.6	6.6	6.0	4.4	3.9	2.4	1.9	1.6	1.5	§	0.3	-1.2
Population Density:																						
Large MSA	—	5.7	7.1	4.8	6.5	6.1	6.8	6.2	7.7	7.1	5.7	4.3	4.8	3.9	1.5	1.4	1.5	0.9	0.3	§	0.6	+0.3
Other MSA	—	7.9	6.4	6.9	5.3	7.6	7.2	6.7	8.9	7.5	5.8	4.5	4.7	3.0	2.8	1.5	1.3	1.1	1.5	§	0.4	-1.1 ss
Non-MSA	—	6.2	9.2	6.4	6.3	6.9	7.8	7.6	7.1	8.9	6.3	4.5	4.0	3.9	3.5	3.0	1.8	1.5	1.1	§	0.2	-0.9 ss
Parental Education: ^b																						
1.0–2.0 (Low)	—	6.1	6.4	6.4	4.9	7.6	6.5	6.3	10.4	12.1	5.8	6.6	8.2	3.0	4.8	1.1	2.4	3.0	0.2	§	0.3	+0.1
2.5–3.0	—	8.0	8.3	8.2	7.7	8.0	8.6	7.5	9.1	9.6	7.5	5.8	4.4	3.5	3.0	2.1	1.1	1.6	0.5	§	0.7	+0.1
3.5–4.0	—	7.5	8.5	6.2	6.8	8.1	8.2	7.9	9.0	8.9	7.3	4.9	5.3	5.1	3.5	2.3	1.1	0.8	1.7	§	1.1	-0.5
4.5–5.0	—	5.7	6.3	5.1	5.0	5.6	7.0	6.0	7.3	4.5	4.3	3.6	3.7	2.4	1.7	1.7	1.6	0.4	0.9	§	0.2	-0.8 s
5.5–6.0 (High)	—	6.6	5.6	5.0	4.6	6.3	4.7	5.7	6.0	5.8	4.1	3.1	3.7	2.7	1.2	1.2	1.3	0.9	1.8	§	0.1	-1.7 s
Race/Ethnicity (2-year average): ^c																						
White	—	—	8.0	7.6	7.1	7.4	8.3	8.5	8.3	8.3	7.2	5.3	4.7	4.8	3.6	2.6	1.8	1.2	1.0	§	0.7	-0.3
African American	—	—	3.1	3.0	1.6	2.2	2.4	2.2	2.7	3.4	3.0	1.7	1.9	2.4	2.2	1.2	1.2	0.9	1.1	§	1.9	+0.8
Hispanic	—	—	6.5	6.0	4.8	4.4	4.2	4.1	6.2	8.2	7.3	5.6	4.6	3.3	2.2	1.5	1.5	1.8	1.0	§	0.1	-0.9 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 64
VICODIN:^a Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months																				2019– 2021 change	
	1975– 2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
<i>Approximate weighted N =</i>	—	12,900	14,600	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	9.6	10.5	9.3	9.5	9.7	9.6	9.7	9.7	8.0	8.1	7.5	5.3	4.8	4.4	2.9	2.0	1.7	1.1	1.2	0.9	-0.2*
Gender:																						
Male	—	12.0	13.0	10.7	11.9	10.9	10.9	10.6	11.5	9.5	9.0	8.6	6.5	5.5	5.1	3.2	2.2	1.6	1.4	§	1.1	-0.2
Female	—	7.5	8.1	7.6	7.2	8.7	8.1	8.9	7.9	6.4	7.1	5.9	4.1	3.8	3.8	2.3	1.5	1.3	0.7	§	0.4	-0.4
College Plans:																						
None or under 4 years	—	13.0	14.7	11.6	14.3	11.8	13.2	15.0	13.1	12.0	13.2	9.2	8.3	5.7	5.8	4.1	2.7	3.2	2.4	§	1.1	-1.3 s
Complete 4 years	—	8.4	9.1	8.3	8.4	9.1	8.5	8.5	8.9	7.1	7.1	6.8	4.5	4.4	4.0	2.6	1.7	1.2	0.7	§	0.7	0.0
Region:																						
Northeast	—	8.1	9.4	9.1	8.8	10.8	9.3	9.9	9.0	9.0	5.8	5.4	5.4	3.2	3.4	1.3	1.5	1.1	0.5	§	0.2	-0.3
Midwest	—	11.6	13.2	8.4	12.4	12.0	14.1	11.7	14.3	9.6	9.6	9.7	6.7	7.4	5.6	4.4	2.5	2.2	1.3	§	0.5	-0.8 s
South	—	8.1	6.1	7.0	7.8	6.9	6.3	7.3	6.2	5.0	5.6	4.9	3.1	3.6	3.7	2.3	1.9	1.8	0.9	§	1.3	+0.5
West	—	10.9	16.3	15.4	9.3	11.2	10.9	11.1	10.4	10.6	11.9	9.7	7.1	5.7	5.2	3.9	1.8	1.6	1.9	§	0.9	-1.0
Population Density:																						
Large MSA	—	9.7	7.3	8.3	8.9	10.4	9.1	8.4	8.6	8.4	8.7	8.7	5.5	4.2	3.4	3.0	2.0	2.1	0.9	§	0.4	-0.5
Other MSA	—	10.6	12.2	10.9	9.3	9.4	9.3	10.0	9.7	8.5	8.2	6.8	5.5	5.3	5.4	3.3	2.0	1.3	1.3	§	1.2	-0.2
Non-MSA	—	7.8	11.4	7.2	10.6	9.5	10.7	10.5	11.3	6.2	7.2	7.3	4.6	4.3	3.6	1.9	1.7	2.0	0.8	§	0.9	+0.1
Parental Education: ^b																						
1.0–2.0 (Low)	—	12.1	9.4	7.8	7.8	7.9	6.3	7.8	9.6	6.0	8.6	6.8	6.0	4.9	4.5	2.6	2.0	2.0	0.9	§	1.2	+0.3
2.5–3.0	—	9.6	10.8	8.8	10.5	9.3	10.2	10.5	9.2	8.5	7.6	9.0	5.0	5.0	3.7	3.0	2.5	2.1	1.1	§	0.5	-0.7
3.5–4.0	—	9.5	12.0	10.1	10.0	11.6	9.9	11.2	10.9	8.3	9.0	7.8	6.5	4.1	4.9	2.6	2.0	1.8	1.5	§	1.0	-0.5
4.5–5.0	—	9.9	9.7	8.4	9.0	8.6	10.5	8.8	10.3	7.9	7.3	7.1	5.3	5.1	4.8	2.4	1.2	0.8	0.8	§	0.9	+0.2
5.5–6.0 (High)	—	8.6	8.8	10.1	8.9	10.4	7.9	8.8	7.7	8.4	7.7	5.5	3.4	4.8	4.2	4.2	2.0	1.6	0.4	§	0.6	+0.2
Race/Ethnicity (2-year average): ^c																						
White	—	—	11.7	11.2	10.5	11.3	11.5	11.6	11.8	10.4	9.4	8.7	6.6	5.0	4.7	3.8	2.5	1.9	1.2	§	0.7	-0.5 s
African American	—	—	3.3	3.1	3.3	3.7	3.3	3.6	4.3	4.3	4.1	3.8	3.2	2.7	2.9	2.9	1.8	1.1	1.2	§	1.5	+0.3
Hispanic	—	—	7.1	7.2	7.7	6.5	5.7	4.9	5.7	6.2	6.0	7.1	7.1	4.7	3.5	3.4	2.3	1.7	1.4	§	1.5	+0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms in 2002–2005; N is two sixths of N indicated in Table S3. Data based on three of six forms beginning in 2006; N is three sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 65
AMPHETAMINES: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	6.2	6.5	7.2	7.9	8.7	9.1	8.1	7.2	6.9	6.5	6.7	5.5	5.5	4.9	4.9
Gender:															
Male	5.5	5.2	5.6	6.5	7.0	6.7	6.6	5.6	5.6	5.1	5.7	4.8	4.4	3.7	3.5
Female	6.9	7.9	8.8	9.3	10.3	11.3	9.6	8.7	8.2	7.7	7.5	6.2	6.5	6.1	6.2
College Plans:															
None or under 4 years	11.6	12.9	14.6	14.5	17.1	15.5	14.1	15.4	14.0	14.2	12.0	12.2	12.4	8.9	11.2
Complete 4 years	5.4	5.7	6.3	7.0	7.6	8.3	7.5	6.3	6.2	5.7	6.1	4.8	4.8	4.5	4.3
Region:															
Northeast	5.1	4.3	5.9	6.9	7.3	7.6	6.5	5.5	5.6	5.0	5.5	3.4	4.1	2.8	3.5
Midwest	7.1	8.0	7.3	7.8	10.6	10.8	9.3	7.2	8.3	7.8	7.1	5.9	5.9	5.1	5.0
South	6.1	6.6	7.3	8.3	8.6	8.7	8.1	8.4	7.5	7.0	8.1	6.7	6.4	5.7	5.8
West	6.0	6.6	8.6	8.4	7.9	9.1	8.3	6.7	5.4	5.4	4.6	4.7	4.6	4.9	4.6
Population Density:															
Large MSA	5.8	4.8	5.6	6.6	7.2	7.9	6.4	5.4	5.2	5.2	5.0	4.2	3.9	4.0	3.6
Other MSA	6.2	7.5	8.2	8.8	8.9	10.0	8.1	7.4	6.8	6.4	7.6	5.9	5.5	5.4	5.8
Non-MSA	6.7	7.0	7.5	7.5	10.1	8.9	9.9	8.8	9.3	8.5	6.9	6.3	7.5	5.0	5.2
Parental Education: ^b															
1.0–2.0 (Low)	8.3	8.4	10.2	11.2	11.8	10.1	9.6	11.2	8.9	9.7	7.4	6.3	8.4	9.7	8.3
2.5–3.0	6.6	7.3	8.2	9.0	10.6	9.9	9.2	8.1	7.8	8.1	8.2	7.3	7.4	5.2	6.6
3.5–4.0	6.7	7.4	7.8	8.5	10.1	10.3	8.9	7.7	8.2	6.8	7.8	5.7	5.9	5.6	5.6
4.5–5.0	5.3	5.5	6.4	6.6	6.8	8.6	7.5	6.2	5.6	5.2	5.9	4.9	4.3	4.2	3.7
5.5–6.0 (High)	5.7	5.4	5.3	5.7	6.4	8.7	7.3	6.4	5.8	5.2	4.5	4.3	3.9	3.5	3.1
Race/Ethnicity (2-year average): ^c															
White	—	6.8	7.4	8.1	9.3	10.2	9.9	9.0	8.4	8.0	7.8	7.2	6.4	5.9	5.4
African American	—	3.3	3.4	3.9	3.9	3.4	3.0	2.8	2.7	2.4	2.3	2.7	2.5	2.1	2.2
Hispanic	—	7.2	7.7	8.6	8.7	8.6	8.1	7.2	7.0	6.8	6.6	5.9	5.0	5.4	5.3

Table continued on next page.

TABLE 65 (cont.)
AMPHETAMINES: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021
	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	<u>change</u>
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	4.7	4.2	4.5	4.1	3.9	3.5	2.9†	4.2	4.3	4.1	3.5	3.5	3.7	4.1	5.3	3.0	-1.1* sss
Gender:																	
Male	3.5	3.3	3.5	3.7	3.4	3.1	2.4†	3.5	3.4	3.1	2.9	2.4	3.6	2.8	§	2.1	-0.7 s
Female	5.7	4.9	5.5	4.4	4.3	3.8	3.3†	4.8	4.8	5.1	4.1	4.4	3.9	5.3	§	3.5	-1.8 ss
College Plans:																	
None or under 4 years	10.7	10.0	12.4	10.8	8.8	8.1	6.8†	8.7	10.4	6.5	7.4	6.0	7.0	7.5	§	4.8	-2.8 s
Complete 4 years	4.1	3.6	3.8	3.5	3.5	3.1	2.6†	3.9	3.7	3.9	3.1	3.2	3.3	3.5	§	2.7	-0.8 s
Region:																	
Northeast	2.2	2.6	2.6	2.9	2.0	2.3	2.1†	2.6	3.3	3.4	1.6	2.3	2.1	3.5	§	1.6	-1.9 ss
Midwest	5.3	4.8	5.1	4.1	4.0	3.5	2.2†	4.0	3.8	3.8	3.5	3.0	3.2	4.0	§	3.5	-0.5
South	5.6	4.6	5.5	5.1	4.8	3.8	4.0†	5.3	4.4	4.6	4.2	3.9	4.3	4.2	§	3.1	-1.1 s
West	4.3	3.7	3.8	3.0	3.6	3.9	2.2†	3.8	5.2	4.0	3.7	4.4	4.5	4.4	§	3.3	-1.2
Population Density:																	
Large MSA	3.9	3.4	4.1	2.7	2.6	2.6	2.6†	3.8	3.3	3.1	2.8	3.5	4.0	3.4	§	2.6	-0.9 s
Other MSA	4.8	4.6	4.8	4.8	4.6	4.1	3.1†	4.3	4.7	5.0	3.6	3.6	3.7	4.4	§	3.2	-1.2 s
Non-MSA	5.4	4.4	4.4	4.3	3.9	3.6	3.0†	4.8	4.8	3.5	4.3	3.6	3.3	4.4	§	3.2	-1.1
Parental Education: ^b																	
1.0–2.0 (Low)	6.5	7.3	7.9	6.5	5.9	5.1	3.8†	6.0	5.3	6.0	5.0	5.8	4.8	5.4	§	3.8	-1.6
2.5–3.0	6.0	4.9	6.0	4.5	4.6	5.4	3.7†	5.0	5.3	5.3	4.0	4.1	4.2	5.8	§	3.4	-2.4 ss
3.5–4.0	5.0	4.5	5.4	5.3	4.1	3.8	3.4†	6.1	4.8	4.4	3.5	3.7	4.5	4.1	§	4.1	0.0
4.5–5.0	4.2	2.9	3.2	3.3	3.5	2.5	1.9†	3.5	3.2	3.3	3.2	2.7	3.2	3.8	§	2.7	-1.1 s
5.5–6.0 (High)	3.6	3.5	3.4	2.8	2.5	2.4	2.6†	2.3	3.6	3.3	3.3	2.7	3.0	3.1	§	2.6	-0.5
Race/Ethnicity (2-year average): ^c																	
White	5.3	4.9	4.9	4.9	4.5	3.9	3.2	2.7†	3.9	4.1	3.6	3.2	3.3	3.8	§	4.2	+0.4
African American	2.0	1.7	1.8	2.4	2.3	1.9	1.9	1.6†	3.2	3.2	3.3	3.3	3.3	2.9	§	2.0	-1.0
Hispanic	4.5	3.9	4.0	3.9	3.5	3.9	3.6	2.6†	4.5	4.1	3.8	3.7	4.0	4.4	§	3.2	-1.2

Table continued
on next page.

TABLE 65 (cont.)
AMPHETAMINES:^a Trends in Annual Prevalence of Use
by Subgroups in Grade 8

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. In 2009, the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording changes. In 2010 the remaining forms were changed in a like manner. In 2013 the question wording was changed two of the 8th and 10th grade questionnaires. The new wording in 2013 asked "On how many occasions (if any) have you taken amphetamines or other prescription stimulant drugs..." In contrast, the old wording did not include "or other prescription stimulant drugs." Results in 2013 indicated higher prevalence in questionnaires with the new as compared to the old wording; it was proportionally 61% higher in 8th grade and 34% higher in 10th grade. 2013 data are based on the changed forms only; *N* is one half of *N* indicated. Beginning in 2014 all questionnaires included the new, updated wording.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined. 2013 race/ethnicity data are based on the 2012 data combined with the 2013 data from the unchanged forms which were double weighted. The 2014 race/ethnicity data are based on the 2013 changed data which were double weighted and the 2014 data. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 66
AMPHETAMINES: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	8.2	8.2	9.6	10.2	11.9	12.4	12.1	10.7	10.4	11.1	11.7	10.7	9.0	8.5	7.8
Gender:															
Male	7.0	7.0	8.2	8.6	9.6	10.5	10.3	9.0	9.2	10.3	10.6	9.6	7.8	7.3	6.6
Female	9.3	9.3	10.9	11.7	14.1	14.2	13.9	12.3	11.5	11.8	12.7	11.8	10.1	9.6	9.0
College Plans:															
None or under 4 years	13.4	14.4	15.5	16.6	19.9	20.3	19.3	17.9	16.3	18.2	20.4	17.5	15.7	14.8	13.7
Complete 4 years	7.1	6.9	8.4	8.9	10.6	11.1	10.9	9.5	9.5	10.0	10.3	9.7	7.9	7.6	7.0
Region:															
Northeast	6.1	5.4	7.8	8.7	9.8	11.5	10.7	11.0	12.1	9.8	10.5	9.3	7.4	8.5	7.4
Midwest	10.3	9.4	9.5	10.5	13.3	14.0	11.0	9.8	10.3	11.1	10.7	10.4	9.7	8.2	8.5
South	8.1	8.7	10.9	11.2	12.8	12.6	14.2	12.6	10.8	12.0	14.0	12.2	10.2	9.6	8.6
West	7.7	8.4	9.5	9.4	10.6	10.6	11.1	8.5	8.2	10.9	9.9	9.6	8.1	7.3	5.8
Population Density:															
Large MSA	7.5	6.7	7.6	8.0	9.2	10.5	9.9	8.9	9.3	10.7	9.8	9.8	6.2	6.2	6.5
Other MSA	7.9	8.0	9.5	10.8	12.8	12.8	11.5	10.3	10.6	9.9	11.1	11.0	9.5	9.6	8.0
Non-MSA	9.3	10.0	11.6	11.2	13.3	13.7	15.5	13.8	11.5	13.6	15.1	11.2	12.2	9.2	9.0
Parental Education: ^b															
1.0–2.0 (Low)	10.0	11.9	12.3	10.8	14.3	15.1	12.2	12.6	10.7	13.3	12.3	13.1	10.9	9.2	8.9
2.5–3.0	9.7	8.9	10.5	11.6	14.2	13.0	14.1	12.8	11.3	12.9	13.6	12.8	11.2	9.9	8.9
3.5–4.0	7.9	8.4	10.5	11.1	12.4	14.1	13.5	11.1	11.2	12.2	12.6	11.1	9.9	10.1	8.1
4.5–5.0	7.4	6.6	7.5	8.9	10.7	10.7	10.6	9.0	9.8	9.7	10.5	9.5	7.6	7.2	7.4
5.5–6.0 (High)	6.9	6.9	8.3	7.3	8.8	10.1	9.2	9.4	9.8	8.8	9.2	8.1	6.4	6.3	6.6
Race/Ethnicity (2-year average): ^c															
White	—	9.4	10.1	11.0	12.4	13.9	14.2	13.6	12.6	12.7	13.4	13.4	12.2	10.6	9.5
African American	—	2.8	3.0	4.0	4.0	3.4	3.1	2.9	2.8	2.5	2.9	3.5	2.8	2.7	3.0
Hispanic	—	6.2	7.0	7.7	8.9	10.3	9.8	8.9	8.8	9.1	8.3	7.9	7.7	7.3	7.3

Table continued on next page.

TABLE 66 (cont.)
AMPHETAMINES: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	7.9	8.0	6.4	7.1	7.6	6.6	6.5†	7.9	7.6	6.8	6.1	5.6	5.7	5.2	4.3	2.7	-2.6* sss
Gender:																	
Male	6.7	7.4	5.8	6.8	7.4	7.2	6.9†	8.3	6.9	6.4	6.1	5.3	5.4	4.9	§	1.9	-2.9 sss
Female	8.9	8.7	6.9	7.4	7.7	6.0	6.2†	7.5	8.3	7.3	6.2	5.7	5.9	5.5	§	2.8	-2.7 sss
College Plans:																	
None or under 4 years	14.8	15.4	12.5	15.3	15.7	14.6	13.8†	11.5	14.7	12.8	12.1	9.3	10.2	7.3	§	4.2	-3.1 sss
Complete 4 years	7.0	7.1	5.6	6.3	6.7	5.7	5.8†	7.6	6.9	6.3	5.5	5.2	5.0	4.8	§	2.2	-2.6 sss
Region:																	
Northeast	6.7	7.3	4.2	5.2	5.9	5.7	6.2†	6.2	6.6	6.4	4.0	4.6	4.6	3.7	§	3.0	-0.7
Midwest	8.7	8.7	7.3	7.0	8.6	7.4	6.4†	8.5	8.7	6.0	6.9	6.9	6.6	4.6	§	2.4	-2.2 sss
South	8.5	8.1	6.8	8.3	7.6	7.2	7.5†	9.0	7.9	7.5	6.8	5.4	5.6	5.5	§	2.7	-2.8 sss
West	7.1	7.7	6.6	7.4	8.1	5.6	5.2†	7.3	7.0	7.1	6.2	5.5	5.8	6.5	§	2.6	-3.9 sss
Population Density:																	
Large MSA	6.8	7.5	5.4	6.3	6.9	6.0	5.9†	7.5	7.3	6.6	6.0	6.0	5.8	4.4	§	2.6	-1.8 ss
Other MSA	8.3	8.0	6.6	7.7	8.0	6.4	6.5†	8.4	7.3	6.7	6.2	5.6	5.7	5.7	§	2.4	-3.3 sss
Non-MSA	8.7	9.0	7.2	7.3	8.1	8.0	7.3†	7.3	8.9	7.6	6.3	5.1	5.4	5.3	§	3.5	-1.8 ss
Parental Education: ^b																	
1.0–2.0 (Low)	9.3	7.9	7.8	9.8	9.6	6.6	9.8†	10.9	9.5	9.5	6.7	6.0	6.5	6.1	§	3.9	-2.2
2.5–3.0	9.0	10.2	7.6	8.0	8.5	7.9	6.8†	9.9	8.7	8.0	6.7	6.0	6.7	7.2	§	3.3	-3.9 sss
3.5–4.0	9.1	8.9	7.7	7.8	8.3	7.3	6.7†	8.2	8.7	8.5	7.3	5.6	5.4	5.4	§	3.2	-2.1 s
4.5–5.0	6.6	6.9	5.4	6.0	6.9	5.8	6.1†	7.3	6.5	6.1	5.6	5.5	5.0	3.8	§	2.2	-1.7 ss
5.5–6.0 (High)	6.8	6.0	4.3	6.5	6.3	5.3	5.7†	5.6	6.2	5.0	5.3	5.8	5.9	4.8	§	2.1	-2.8 sss
Race/Ethnicity (2-year average): ^c																	
White	9.2	9.4	8.7	7.8	8.5	8.3	7.4	7.1†	8.5	7.7	7.0	6.5	6.2	5.5	§	3.4	-2.1 sss
African American	2.8	2.3	2.3	2.3	2.8	3.0	2.9	2.6†	5.3	4.8	4.6	3.9	3.3	3.5	§	2.5	-0.9
Hispanic	6.3	5.8	6.0	6.1	6.3	6.1	6.1	4.7†	7.4	6.7	6.6	5.6	5.4	5.8	§	2.9	-2.9 sss

Table continued
on next page.

TABLE 66 (cont.)
AMPHETAMINES:^a Trends in Annual Prevalence of Use by Subgroups in Grade 10

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data ' not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. In 2009, the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording changes. In 2010 the remaining forms were changed in a like manner. In 2013 the question wording was changed two of the 8th and 10th grade questionnaires. The new wording in 2013 asked "On how many occasions (if any) have you taken amphetamines or other prescription stimulant drugs..." In contrast, the old wording did not include "or other prescription stimulant drugs." Results in 2013 indicated higher prevalence in questionnaires with the new as compared to the old wording; it was proportionally 61% higher in 8th grade and 34% higher in 10th grade. 2013 data are based on the changed forms only; *N* is one half of *N* indicated. Beginning in 2014 all questionnaires included the new, updated wording.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined. 2013 race/ethnicity data are based on the 2012 data combined with the 2013 data from the unchanged forms which were double weighted. The 2014 race/ethnicity data are based on the 2013 changed data which were double weighted and the 2014 data. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 67
AMPHETAMINES: ^{a,b} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
Approximate weighted N =	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	16.2	15.8	16.3	17.1	18.3	20.8	26.0	20.3	17.9	17.7	15.8	13.4	12.2	10.9	10.8	9.1
Gender:																
Male	15.6	15.8	16.0	16.9	18.4	19.7	24.8	19.6	17.2	16.8	14.9	12.7	11.8	10.8	11.1	9.4
Female	16.5	15.4	16.4	17.1	17.8	21.8	26.9	20.3	17.9	18.2	16.4	13.8	12.4	10.9	10.5	8.6
College Plans:																
None or under 4 years	—	19.3	20.5	20.0	21.8	25.8	30.9	23.7	20.9	22.2	19.7	17.7	16.0	13.9	15.1	12.6
Complete 4 years	—	11.9	11.5	13.7	14.5	16.5	22.3	16.8	14.5	14.2	13.3	10.9	10.2	9.5	9.1	7.4
Region:																
Northeast	16.5	14.7	16.8	19.6	22.0	22.0	28.8	21.5	17.9	19.0	16.8	12.6	10.4	8.4	9.0	6.3
Midwest	18.7	17.8	19.0	18.2	18.3	22.2	30.1	24.1	20.4	20.3	17.3	15.2	13.5	12.2	13.3	10.7
South	12.6	13.7	13.2	14.0	14.0	17.7	19.6	16.4	15.4	15.1	12.8	11.5	11.5	10.8	9.9	8.9
West	18.5	17.2	16.0	17.8	20.7	22.1	26.6	18.7	18.2	16.9	17.3	15.0	13.4	11.8	11.1	10.2
Population Density:																
Large MSA	19.6	15.4	15.3	17.7	19.5	21.9	28.0	21.6	18.1	17.7	15.0	11.2	10.9	8.8	7.1	6.5
Other MSA	15.5	16.3	17.1	17.5	18.9	20.8	25.5	20.7	19.6	17.1	15.7	14.2	11.9	11.9	11.4	9.6
Non-MSA	14.8	15.4	15.9	16.0	16.6	19.9	25.1	18.8	15.6	18.5	16.6	14.1	14.0	11.3	13.3	10.6
Parental Education: ^c																
1.0–2.0 (Low)	15.7	13.4	14.5	14.9	16.0	19.1	22.3	18.7	15.7	17.1	14.5	11.9	11.9	9.8	10.4	7.6
2.5–3.0	16.7	16.9	17.4	17.3	18.4	22.2	26.7	21.9	19.6	19.2	17.0	15.2	13.3	11.1	11.7	9.7
3.5–4.0	14.9	16.6	16.1	18.2	19.6	21.5	26.9	21.7	19.4	18.5	17.2	14.3	12.6	11.8	12.3	10.6
4.5–5.0	14.5	16.8	15.9	16.9	17.1	20.0	26.2	19.1	18.9	15.9	15.1	12.0	11.7	10.3	9.4	8.1
5.5–6.0 (High)	12.0	14.6	16.0	17.2	20.4	17.9	26.8	20.5	16.1	14.0	10.9	10.1	10.4	10.0	9.1	7.3
Race/Ethnicity (2-year average): ^d																
White	—	—	17.3	18.2	19.2	21.3	26.4	23.6	22.3	20.5	18.9	16.4	14.3	13.0	12.4	11.4
African American	—	—	5.3	4.7	4.2	5.3	5.8	6.0	5.7	4.7	4.3	4.0	3.8	3.9	3.6	3.1
Hispanic	—	—	12.3	12.2	12.8	14.5	17.5	12.3	11.5	13.2	14.6	10.8	8.7	9.6	9.0	7.0

Table continued on next page.

TABLE 67 (cont.)
AMPHETAMINES: ^{a,b} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Approximate weighted N =	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	8.2	7.1	8.4	9.4	9.3	9.5	10.2	10.1	10.2	10.5	10.9	11.1	9.9	10.0	8.6	8.1
Gender:																
Male	8.3	7.2	8.2	9.2	9.5	9.6	10.1	10.3	10.6	10.4	10.9	11.3	9.8	9.9	9.1	7.4
Female	7.9	6.9	8.5	9.4	8.9	8.8	10.2	9.8	9.6	10.5	10.6	10.7	9.5	9.9	7.9	8.5
College Plans:																
None or under 4 years	11.0	9.7	11.0	13.4	12.3	12.8	14.1	13.6	12.7	13.9	14.4	14.8	14.4	13.3	11.5	10.6
Complete 4 years	7.0	6.1	7.6	8.0	8.3	8.4	8.9	9.0	9.5	9.6	9.8	10.1	8.4	8.9	7.8	7.5
Region:																
Northeast	6.5	6.2	8.1	7.4	9.6	10.4	11.1	9.0	9.9	10.6	12.0	11.5	9.7	9.5	7.4	8.5
Midwest	10.1	8.4	8.9	12.0	9.5	10.0	10.8	11.0	10.5	10.4	12.7	11.3	10.3	10.2	10.3	7.6
South	7.9	6.7	8.3	9.0	9.2	9.1	9.8	10.4	10.8	10.2	9.5	11.7	9.7	10.3	9.4	8.8
West	7.8	6.9	8.3	8.4	8.9	8.3	9.1	9.6	8.8	10.9	9.7	9.6	9.7	9.5	6.1	7.0
Population Density:																
Large MSA	6.2	6.0	6.5	7.8	9.1	7.9	8.9	9.0	7.0	8.3	10.6	10.7	7.5	8.0	7.8	8.7
Other MSA	8.4	6.7	8.5	9.4	8.5	8.9	9.5	9.9	10.8	10.9	10.8	11.0	10.5	10.6	8.9	7.6
Non-MSA	9.5	9.0	9.8	10.9	10.8	11.9	13.0	12.2	12.4	12.4	11.4	12.1	11.7	11.1	8.9	8.2
Parental Education: ^c																
1.0–2.0 (Low)	9.5	7.0	9.0	10.4	9.9	8.1	9.8	9.7	9.9	11.0	8.9	8.1	10.6	7.5	6.1	8.2
2.5–3.0	9.1	7.7	8.6	10.3	9.9	10.5	10.3	10.6	11.3	11.0	12.2	11.4	9.9	11.0	9.2	8.2
3.5–4.0	8.9	7.7	9.1	9.4	9.1	9.3	10.8	11.4	10.8	10.6	11.6	12.3	10.7	12.3	9.5	8.9
4.5–5.0	6.5	6.3	8.0	9.5	9.2	8.9	9.4	9.4	9.7	10.3	10.7	11.1	9.4	8.2	8.5	7.8
5.5–6.0 (High)	5.7	5.8	7.6	7.1	8.1	9.1	10.2	8.7	8.4	10.2	8.7	10.2	7.7	8.4	7.4	7.0
Race/Ethnicity (2-year average): ^d																
White	9.8	8.8	9.0	10.4	10.7	10.5	11.4	12.1	11.9	11.9	12.6	13.2	12.4	11.6	11.0	9.9
African American	2.7	2.2	2.3	3.4	3.4	2.9	2.8	2.8	2.5	2.6	3.0	2.9	2.8	2.4	2.4	2.6
Hispanic	6.1	6.0	6.2	6.4	7.1	7.8	7.3	7.0	7.6	9.2	9.2	7.9	6.8	7.0	6.7	6.4

Table continued on next page.

TABLE 67 (cont.)
AMPHETAMINES: ^{a,b} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	7.5	6.8	6.6	7.4	8.2	7.9‡	9.2	8.1	7.7	6.7	5.9	5.5	4.5	4.3	2.3	-2.1* sss
Gender:																
Male	7.4	6.5	7.5	8.3	8.8	8.2‡	10.0	8.8	8.1	7.5	6.9	6.3	4.9	§	2.2	-2.7 sss
Female	7.3	6.8	5.7	6.4	7.4	7.5‡	7.9	7.3	7.1	5.8	5.0	4.2	3.8	§	2.1	-1.8 sss
College Plans:																
None or under 4 years	10.3	10.7	8.4	10.5	11.6	8.9‡	13.1	9.5	9.3	6.7	6.8	6.7	6.1	§	2.9	-3.2 sss
Complete 4 years	6.7	5.7	6.1	6.7	7.4	7.6‡	8.4	7.6	7.3	6.6	5.7	4.9	3.9	§	2.0	-1.9 sss
Region:																
Northeast	7.5	6.5	6.7	8.6	7.1	7.0‡	8.1	7.1	6.6	5.8	5.0	5.4	3.1	§	2.3	-0.8
Midwest	8.2	6.9	7.8	8.1	8.6	9.0‡	10.7	8.9	9.0	7.7	5.9	6.0	4.9	§	2.6	-2.4 ss
South	7.9	7.1	6.5	6.9	8.1	7.0‡	9.3	8.9	7.5	6.6	6.3	5.5	4.6	§	2.3	-2.3 sss
West	5.8	6.3	5.2	6.5	8.7	8.5‡	8.5	7.0	7.7	6.5	5.8	5.0	4.7	§	2.2	-2.5 ss
Population Density:																
Large MSA	6.7	6.2	6.3	6.8	7.4	9.2‡	8.5	7.5	7.7	6.2	5.1	4.5	4.4	§	2.3	-2.2 sss
Other MSA	7.4	6.7	7.0	8.0	8.7	7.8‡	10.2	9.0	7.8	7.0	6.6	5.8	4.7	§	2.8	-2.0 ss
Non-MSA	8.6	7.5	6.2	6.8	8.1	6.1‡	7.9	7.0	7.5	6.7	5.8	6.6	3.7	§	1.4	-2.4 sss
Parental Education: ^c																
1.0–2.0 (Low)	7.5	7.3	5.5	5.0	6.8	6.0‡	7.9	6.8	6.4	5.8	4.2	4.1	3.8	§	0.8	-3.0 sss
2.5–3.0	7.6	7.3	7.3	7.2	8.1	7.8‡	10.1	8.9	7.3	6.0	6.0	5.1	3.8	§	1.6	-2.2 sss
3.5–4.0	7.6	6.5	7.3	7.9	8.3	8.0‡	9.0	7.8	7.7	7.4	5.9	6.0	4.6	§	3.7	-0.8
4.5–5.0	7.4	6.2	6.3	8.1	7.6	8.7‡	10.0	8.5	8.5	6.9	6.1	5.5	5.0	§	2.3	-2.7 sss
5.5–6.0 (High)	7.1	7.1	5.6	7.5	9.7	8.4‡	9.3	8.6	9.5	7.8	7.4	6.2	5.1	§	2.0	-3.1 sss
Race/Ethnicity (2-year average): ^d																
White	9.3	8.7	8.1	8.6	9.6	10.0	10.1‡	9.8	9.1	8.6	7.6	6.8	5.7	§	3.7	-2.0 ss
African American	2.5	2.2	2.3	2.8	2.9	2.9	3.3‡	5.9	4.3	3.7	3.4	3.1	2.7	§	2.5	-0.2
Hispanic	5.8	4.7	4.4	4.4	4.8	4.7	6.3‡	6.9	6.5	5.8	4.7	3.9	3.6	§	1.5	-2.1 sss

TABLE 67 (cont.)
AMPHETAMINES: ^{a,b} Trends in Annual Prevalence of Use by Subgroups in Grade 12

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. In 2013 the question wording was changed slightly in four of the six questionnaire forms. Vyvanse was added to the list of examples in one questionnaire form. 2013 data presented here are based on the unchanged forms only; N is two sixths of N indicated.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bBeginning in 1982, the question about amphetamine use was revised to get respondents to exclude the inappropriate reporting of nonprescription amphetamines. The prevalence rate dropped slightly as a result of this methodological change. (In 1982 and 1983, the data are based on three of the five questionnaire forms.) Only drug use not under a doctor's orders is included here. In 2009, the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording changes. In 2010 the remaining forms were changed in a like manner. In 2011 bennies, Benzedrine, and Methadrine were deleted from the list of examples in one questionnaire form. An examination of the data did not show any effect from the wording changes. In 2013 the question wording was changed in three of the 12th grade questionnaires. The new wording in 2013 asked "On how many occasions (if any) have you taken amphetamines or other prescription stimulant drugs..." In contrast, the old wording did not include "or other prescription stimulant drugs." Results in 2013 indicated higher prevalence in questionnaires with the new as compared to the old wording; it was proportionally 21% higher in 12th grade. 2013 data are based on the changed forms only; N is one half of N indicated. Beginning in 2014 all questionnaires included the new, updated wording.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^eTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined. 2013 race/ethnicity data are based on the 2012 data combined with the 2013 data from the unchanged forms which were tripleweighted. The 2014 race/ethnicity data are based on the 2013 changed data which were double weighted and the 2014 data. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 68
RITALIN: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																					2019– 2021 change	
	1991– 2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
Approximate weighted N =	—	16,200	15,100	16,500	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	2.9	2.8	2.6	2.5	2.4	2.6	2.1	1.6	1.8	1.5	1.3	0.7	1.1	0.9	0.6	0.8	0.4	0.5	1.0	0.5	0.6	-0.4*
Gender:																							
Male	—	3.2	3.2	2.9	2.6	2.4	2.7	1.9	1.5	2.4	1.5	1.1	0.5	1.3	0.8	0.8	1.2	0.5	0.7	1.2	§	0.5	-0.7 s
Female	—	2.6	2.5	2.3	2.4	2.4	2.5	2.3	1.5	1.4	1.5	1.3	0.8	0.8	1.0	0.5	0.4	0.3	0.3	0.7	§	0.7	0.0
College Plans:																							
None or under 4 years	—	8.1	7.7	6.8	8.8	7.5	9.0	4.2	5.7	5.3	5.8	3.7	3.0	3.2	4.6	2.6	2.9	1.1	2.1	3.4	§	1.9	-1.5
Complete 4 years	—	2.4	2.2	2.1	1.8	1.8	1.9	1.8	1.2	1.6	1.1	1.1	0.5	0.9	0.6	0.4	0.6	0.3	0.4	0.6	§	0.3	-0.3
Region:																							
Northeast	—	1.5	2.8	1.7	1.6	2.0	2.0	0.9	0.4	1.2	1.8	1.5	0.1	1.3	1.3	0.2	0.5	0.1	0.5	0.6	§	0.0	-0.6
Midwest	—	3.0	3.0	2.9	2.4	1.9	3.2	2.4	2.0	2.3	1.8	1.2	0.1	0.7	0.2	0.5	0.4	0.2	0.4	1.0	§	0.2	-0.7
South	—	3.5	2.8	2.9	3.0	3.0	2.5	2.5	2.2	1.5	1.4	1.4	1.4	1.2	0.9	0.8	1.2	0.7	0.5	1.4	§	1.1	-0.3
West	—	2.9	2.5	2.2	2.5	2.1	2.7	1.6	1.2	2.2	1.2	1.0	0.5	1.3	1.4	0.7	0.7	0.5	0.7	0.7	§	0.5	-0.1
Population Density:																							
Large MSA	—	2.2	2.3	2.3	1.7	1.5	2.3	1.2	1.1	1.6	1.1	0.8	0.8	0.9	1.0	0.6	0.9	0.3	0.5	1.2	§	0.5	-0.8
Other MSA	—	2.9	3.1	2.8	3.0	3.4	3.0	2.7	1.8	1.9	1.9	1.8	0.8	1.2	1.0	0.7	1.0	0.6	0.7	0.9	§	0.8	-0.1
Non-MSA	—	3.5	2.8	2.4	2.4	1.7	2.3	2.0	1.9	2.0	1.3	0.9	0.2	1.0	0.8	0.3	0.3	0.3	0.2	0.9	§	0.4	-0.5
Parental Education: ^b																							
1.0–2.0 (Low)	—	4.8	5.3	2.5	4.2	5.2	6.5	1.9	4.4	4.6	2.3	2.7	0.7	2.3	2.0	0.8	1.8	0.9	0.6	2.0	§	0.4	-1.6
2.5–3.0	—	3.3	4.0	2.8	2.9	2.7	3.1	2.3	2.2	2.2	1.4	1.6	0.8	1.2	0.7	1.1	0.7	0.2	0.6	1.4	§	0.0	-1.4
3.5–4.0	—	3.5	3.3	3.3	2.7	2.7	2.6	2.4	1.7	2.1	0.5	1.5	1.1	1.2	1.0	0.4	1.0	0.4	0.5	1.2	§	1.5	+0.3
4.5–5.0	—	2.4	1.5	1.8	2.1	1.5	2.0	1.8	1.0	1.3	1.4	0.9	0.3	1.2	1.0	0.4	0.4	0.2	0.5	0.7	§	0.2	-0.5
5.5–6.0 (High)	—	2.1	1.7	2.1	1.9	1.6	1.2	1.8	0.7	1.4	1.6	0.8	0.6	0.4	0.4	0.3	0.6	0.6	0.3	0.5	§	0.9	+0.3
Race/Ethnicity (2-year average): ^c																							
White	—	—	3.0	2.8	2.7	2.5	2.6	2.4	1.9	1.7	1.7	1.4	1.0	0.7	0.8	0.6	0.4	0.4	0.3	0.4	§	0.7	+0.3
African American	—	—	1.0	1.3	1.5	1.6	1.8	1.8	1.8	1.3	1.0	1.1	1.0	1.4	1.2	0.7	1.5	1.6	0.9	1.2	§	0.9	-0.3
Hispanic	—	—	3.8	2.7	2.5	3.3	3.1	2.8	2.1	1.5	1.5	1.3	1.2	1.3	1.3	0.7	0.7	0.6	0.6	0.9	§	0.2	-0.7 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 69
RITALIN: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																					2019– 2021 change	
	1991– 2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
Approximate weighted N =	—	14,000	14,300	15,800	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	4.8	4.8	4.1	3.4	3.4	3.6	2.8	2.9	3.6	2.7	2.6	1.9	1.8	1.8	1.6	1.2	0.8	0.9	0.7	1.0	0.3	-0.3*
Gender:																							
Male	—	5.2	4.9	4.2	3.6	3.2	3.7	2.6	3.0	3.8	3.1	2.9	2.8	2.5	2.2	2.1	1.5	0.7	1.5	1.1	§	0.3	-0.8 s
Female	—	4.3	4.6	3.8	3.0	3.5	3.7	2.9	2.9	3.4	2.1	2.5	1.0	1.2	1.5	1.2	0.9	0.7	0.4	0.2	§	0.2	0.0
College Plans:																							
None or under 4 years	—	7.2	10.2	11.0	5.5	6.7	9.1	6.0	8.6	9.3	6.2	7.3	4.5	3.7	6.0	4.7	3.2	2.4	1.9	0.8	§	0.6	-0.1
Complete 4 years	—	4.3	3.8	2.9	3.0	2.9	2.8	2.3	2.3	3.0	2.2	2.1	1.6	1.6	1.4	1.3	0.9	0.6	0.7	0.7	§	0.3	-0.4
Region:																							
Northeast	—	5.5	3.6	3.8	4.5	4.1	3.0	2.8	3.3	4.0	2.2	3.5	2.1	1.8	1.6	1.9	1.2	0.4	0.8	0.5	§	0.1	-0.4
Midwest	—	4.6	4.8	4.2	2.7	4.8	4.0	3.6	3.1	4.6	3.2	2.9	2.1	1.9	2.2	1.0	1.6	1.0	1.3	0.5	§	0.8	+0.2
South	—	5.3	5.2	4.2	3.8	3.0	3.4	3.0	3.0	3.5	2.7	2.4	1.9	1.6	1.7	1.4	1.0	0.8	0.8	0.6	§	0.4	-0.2
West	—	3.1	5.1	3.9	2.4	1.4	4.2	1.6	2.4	2.3	2.4	2.1	1.2	1.9	1.9	2.4	1.1	0.8	0.7	1.0	§	0.1	-1.0 s
Population Density:																							
Large MSA	—	4.3	3.6	4.0	2.6	4.0	2.8	2.9	2.0	3.3	3.0	2.8	2.0	1.7	1.7	0.9	0.8	0.8	1.0	0.1	§	0.2	+0.1
Other MSA	—	4.6	5.8	3.7	3.6	3.3	4.1	2.5	3.3	3.6	2.6	2.8	1.7	1.7	1.7	1.8	1.1	0.6	0.8	1.0	§	0.4	-0.6
Non-MSA	—	5.8	4.3	5.1	3.7	2.8	3.7	3.0	3.5	4.1	2.3	2.1	2.1	2.1	2.3	2.5	2.1	1.2	1.0	0.6	§	0.4	-0.3
Parental Education: ^b																							
1.0–2.0 (Low)	—	5.0	7.7	4.2	4.8	3.3	3.9	3.9	2.8	4.1	4.0	3.0	1.2	2.4	1.2	2.2	0.4	1.0	1.8	0.0	§	0.3	+0.3
2.5–3.0	—	5.4	5.9	5.0	3.4	4.4	3.8	3.2	2.9	3.9	2.7	3.3	3.2	3.3	1.3	1.6	2.1	0.9	1.4	0.6	§	0.7	+0.1
3.5–4.0	—	5.0	4.2	4.1	3.0	2.6	3.6	2.7	2.9	3.8	3.1	2.9	1.5	1.6	2.5	2.2	1.2	0.8	0.5	0.8	§	0.4	-0.4
4.5–5.0	—	4.2	3.6	3.7	2.9	4.0	3.4	2.6	3.4	3.3	1.4	1.9	1.7	1.5	1.5	1.1	0.4	0.4	0.7	1.0	§	0.3	-0.7
5.5–6.0 (High)	—	4.6	4.7	3.3	3.7	2.8	4.0	2.2	2.8	3.8	3.3	2.2	1.8	0.7	1.7	1.3	1.6	0.6	0.8	0.6	§	0.2	-0.4
Race/Ethnicity (2-year average): ^c																							
White	—	—	5.5	4.8	4.1	3.9	3.9	3.7	3.5	3.8	3.4	2.9	2.6	2.0	2.1	1.9	1.6	1.0	0.7	0.8	§	0.5	-0.3
African American	—	—	1.8	2.5	1.9	1.0	1.5	1.7	1.4	1.2	2.0	2.2	1.1	1.2	1.3	1.3	0.9	1.1	0.8	1.1	§	2.2	+1.1
Hispanic	—	—	3.1	4.4	3.6	2.2	2.1	1.9	1.6	2.3	2.8	2.6	2.2	1.7	1.5	1.8	1.7	1.2	1.0	0.5	§	0.1	-0.4 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 70
RITALIN: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																					2019– 2021 change	
	1975– 2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
<i>Approximate weighted N =</i>	—	12,800	12,900	14,600	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	5.1	4.0	4.0	5.1	4.4	4.4	3.8	3.4	2.1	2.7	2.6	2.6	2.3	1.8	2.0	1.2	1.3	0.9	1.1	1.7	0.5	-0.6* s
Gender:																							
Male	—	6.0	5.1	5.5	6.0	5.4	4.5	3.6	3.7	3.2	3.5	2.7	2.7	2.9	1.9	2.4	1.3	1.6	1.4	1.5	§	0.5	-1.1 s
Female	—	4.1	2.8	2.6	4.0	3.3	3.8	3.5	2.9	1.1	1.7	2.3	1.9	1.6	1.5	1.4	0.6	0.9	0.4	0.7	§	0.2	-0.5 s
College Plans:																							
None or under 4 years	—	6.4	7.7	4.4	6.8	5.3	5.3	5.6	4.2	3.2	5.4	3.1	2.9	4.1	1.4	2.4	2.4	1.5	1.3	2.3	§	1.0	-1.2
Complete 4 years	—	4.7	3.0	3.7	4.4	4.1	4.0	3.3	3.2	1.8	2.1	2.5	2.3	1.9	1.6	1.8	0.8	1.1	0.7	0.8	§	0.3	-0.4
Region:																							
Northeast	—	4.7	4.5	4.6	4.1	4.7	6.1	4.1	4.2	1.3	3.5	2.1	2.5	2.0	1.6	1.6	0.9	1.5	1.8	0.7	§	0.3	-0.4
Midwest	—	7.4	5.4	3.8	6.0	5.3	3.8	5.2	3.4	3.0	2.9	3.0	2.4	2.0	2.1	2.0	1.3	1.1	0.9	1.4	§	0.0	-1.4
South	—	4.1	3.4	3.5	4.7	4.6	4.3	3.0	3.4	2.1	2.1	2.7	2.8	2.3	2.5	1.8	1.2	1.3	0.7	0.8	§	1.0	+0.2
West	—	3.8	2.6	4.4	5.5	2.7	3.8	3.5	2.8	1.7	2.8	2.2	2.5	2.8	0.4	2.5	1.4	1.2	0.7	1.9	§	0.6	-1.4
Population Density:																							
Large MSA	—	7.2	3.9	3.2	4.2	4.1	5.4	4.0	4.1	2.4	2.7	2.2	2.4	1.9	2.1	2.0	1.7	1.5	0.9	0.9	§	0.4	-0.4
Other MSA	—	3.7	4.0	5.1	5.6	5.1	3.9	3.4	3.5	2.2	2.8	2.6	2.4	2.5	1.8	1.6	1.0	1.5	0.7	1.2	§	0.7	-0.5
Non-MSA	—	5.4	4.0	2.9	4.9	3.4	3.8	4.2	2.4	1.5	2.2	3.1	3.1	2.4	1.2	2.7	1.0	0.4	1.3	1.5	§	0.4	-1.1
Parental Education: ^b																							
1.0–2.0 (Low)	—	5.6	5.3	3.1	5.2	5.5	5.2	2.1	2.0	2.6	2.9	1.9	4.1	2.3	1.9	2.0	1.5	0.8	0.4	2.6	§	1.5	-1.1
2.5–3.0	—	4.5	4.2	3.7	4.2	4.2	3.6	4.1	3.4	3.3	3.0	2.5	2.5	2.1	2.0	2.0	1.0	1.8	1.1	0.7	§	0.0	-0.7
3.5–4.0	—	5.0	3.9	4.8	6.3	4.1	4.6	3.5	3.6	1.3	2.6	2.7	2.7	2.6	1.1	1.6	0.9	1.3	0.8	1.4	§	0.3	-1.1 s
4.5–5.0	—	4.7	3.5	3.5	4.6	3.9	4.7	4.7	3.1	1.9	2.2	2.0	2.1	2.3	1.2	2.0	1.0	0.8	1.0	1.0	§	0.6	-0.4
5.5–6.0 (High)	—	6.6	3.9	4.0	4.5	6.0	4.6	2.7	4.5	1.7	2.4	3.3	2.0	1.3	3.3	1.9	1.4	1.2	0.9	0.5	§	0.9	+0.4
Race/Ethnicity (2-year average): ^c																							
White	—	—	5.4	4.6	5.2	5.3	4.9	4.9	4.3	3.1	2.6	2.7	2.6	2.3	1.9	1.9	1.5	1.2	1.1	0.8	§	0.7	-0.1
African American	—	—	0.8	1.2	1.2	2.0	2.3	1.8	1.5	1.5	2.1	1.9	1.6	2.3	2.2	2.3	2.2	1.1	0.9	0.7	§	1.5	+0.8
Hispanic	—	—	3.1	3.3	3.6	3.6	3.9	2.7	1.5	1.9	2.1	2.3	2.7	2.3	1.6	1.4	1.2	1.1	1.0	1.3	§	2.2	+0.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 71
ADDERALL: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months													2019– 2021
	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	<u>change</u>
Approximate weighted N =	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	2.0	2.3	1.7	1.7	1.8	1.3	1.0	1.5	1.3	1.8	2.5	2.7	1.8	-0.7*
Gender:														
Male	2.8	2.5	1.5	1.3	1.6	1.0	1.2	2.0	1.3	1.5	2.7	§	1.3	-1.4 s
Female	1.3	2.1	1.7	1.9	1.9	1.4	1.0	1.2	1.2	1.9	2.4	§	2.0	-0.4
College Plans:														
None or under 4 years	6.1	5.4	5.1	4.2	7.4	4.9	3.3	4.5	3.4	2.9	6.0	§	3.5	-2.5
Complete 4 years	1.7	2.0	1.4	1.5	1.4	1.0	0.9	1.2	1.0	1.6	2.0	§	1.5	-0.5
Region:														
Northeast	1.0	1.5	1.4	1.0	2.2	1.0	1.1	0.6	0.3	1.1	1.0	§	0.3	-0.7
Midwest	2.3	3.1	2.3	1.0	0.8	0.9	0.6	1.8	1.7	1.5	2.1	§	2.4	+0.2
South	2.3	2.3	1.7	2.6	2.1	1.6	1.5	1.7	1.5	2.2	3.6	§	2.3	-1.3
West	1.8	1.9	1.2	1.3	2.0	1.3	0.6	1.6	1.1	2.0	2.3	§	1.5	-0.9
Population Density:														
Large MSA	1.9	1.7	0.7	2.1	1.5	1.3	0.9	1.2	0.9	1.8	3.2	§	0.9	-2.4 sss
Other MSA	2.0	2.8	2.5	1.4	1.7	1.3	1.4	1.5	1.4	2.1	2.4	§	2.7	+0.4
Non-MSA	2.0	2.0	1.5	1.5	2.3	1.2	0.5	2.1	1.3	1.2	2.0	§	1.3	-0.6
Parental Education: ^b														
1.0–2.0 (Low)	5.3	3.5	3.7	1.1	2.2	1.9	1.9	1.8	0.9	1.3	4.6	§	0.6	-4.0 sss
2.5–3.0	1.8	2.5	2.6	1.9	2.1	1.3	1.0	1.7	1.5	1.9	5.6	§	0.6	-5.1 sss
3.5–4.0	2.9	2.0	1.6	1.5	2.3	1.3	1.3	1.7	1.6	1.9	1.5	§	3.4	+1.9
4.5–5.0	1.4	2.4	1.3	1.4	1.8	1.0	1.1	1.0	1.4	1.9	2.3	§	1.8	-0.5
5.5–6.0 (High)	1.1	1.2	0.7	1.7	0.6	1.0	0.3	1.4	0.2	1.0	1.1	§	1.9	+0.8
Race/Ethnicity (2-year average): ^c														
White	—	2.4	2.1	1.7	1.4	1.3	1.2	1.3	1.5	1.3	1.9	§	2.7	+0.8
African American	—	1.2	1.5	1.4	1.6	1.3	1.1	2.0	2.0	1.5	2.4	§	3.7	+1.4
Hispanic	—	1.9	1.9	1.7	1.8	1.8	0.9	0.8	1.0	1.2	2.1	§	0.5	-1.6 sss

Source: The Monitoring the Future study, the University of Michigan.

Notes: '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table D-S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Tables D-S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 72
ADDERALL: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months													2019– 2021
	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	change
	<i>Approximate weighted N =</i>													
Total	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
	5.7	5.3	4.6	4.5	4.4	4.6	5.2	4.2	4.0	4.1	3.1	2.9	1.6	-1.5* ss
Gender:														
Male	6.1	5.5	5.4	5.0	5.2	4.7	5.3	4.1	4.3	4.8	3.7	§	1.6	-2.2 ss
Female	5.3	4.9	3.8	4.1	3.7	4.4	5.2	4.1	3.6	3.2	2.3	§	1.5	-0.8
College Plans:														
None or under 4 years	15.1	10.6	11.5	9.3	8.1	10.4	9.2	7.9	9.9	7.6	3.9	§	3.2	-0.7
Complete 4 years	4.8	4.6	3.7	4.0	4.0	4.0	4.8	3.7	3.3	3.5	2.9	§	1.2	-1.6 sss
Region:														
Northeast	4.2	4.1	4.9	4.0	5.4	3.9	6.1	3.1	2.7	1.9	1.9	§	1.4	-0.4
Midwest	8.4	7.6	6.3	5.3	5.8	6.1	5.2	4.2	5.5	5.7	2.1	§	2.5	+0.3
South	6.5	5.2	4.6	5.5	4.1	4.9	5.3	4.8	3.7	4.2	3.2	§	1.5	-1.7 s
West	3.4	3.7	2.3	2.5	2.9	3.4	4.2	4.1	3.7	3.6	4.6	§	1.1	-3.5 sss
Population Density:														
Large MSA	6.2	6.0	4.8	3.7	4.3	5.4	3.8	3.6	5.2	6.0	1.9	§	1.5	-0.4
Other MSA	5.9	5.0	4.1	4.4	4.7	4.3	6.0	4.5	3.0	2.6	3.8	§	1.8	-2.1 ss
Non-MSA	4.8	4.8	5.1	5.7	4.0	4.2	5.6	4.3	4.3	4.4	3.2	§	1.4	-1.7 s
Parental Education: ^b														
1.0–2.0 (Low)	7.9	6.7	4.7	5.4	4.2	4.4	5.8	3.1	7.3	5.9	2.6	§	2.2	-0.4
2.5–3.0	6.6	6.0	5.4	6.3	5.2	4.8	5.2	3.7	2.7	4.5	3.1	§	2.0	-1.1
3.5–4.0	5.4	6.2	4.9	4.8	4.5	5.8	6.8	5.5	4.0	4.2	3.6	§	2.2	-1.4
4.5–5.0	4.9	3.8	4.5	4.1	4.7	3.9	5.1	3.4	3.1	2.9	3.2	§	0.9	-2.3 ss
5.5–6.0 (High)	6.7	4.6	3.2	2.9	3.2	4.2	4.0	4.9	4.5	4.8	3.2	§	1.6	-1.6
Race/Ethnicity (2-year average): ^c														
White	—	6.6	5.7	5.3	5.6	5.9	6.0	5.6	4.5	4.1	3.6	§	2.7	-0.9
African American	—	2.2	2.7	1.7	2.0	3.0	3.5	2.9	2.9	2.9	2.5	§	2.4	-0.1
Hispanic	—	4.2	3.9	3.8	2.5	2.0	3.3	4.0	4.6	4.5	3.4	§	0.7	-2.7 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table D-S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table D-S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 73
ADDERALL: ^a Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months													2019– 2021
	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
Approximate weighted N =	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	5.4	6.5	6.5	7.6	7.4	6.8	7.5	6.2	5.5	4.6	3.9	4.4	1.8	-2.1* sss
Gender:														
Male	7.4	7.5	7.4	8.0	8.7	7.7	8.0	7.7	7.1	4.9	4.8	§	1.8	-3.0 sss
Female	3.8	5.5	5.5	6.9	6.0	6.0	6.9	4.4	4.2	3.9	3.0	§	1.3	-1.7 ss
College Plans:														
None or under 4 years	7.0	10.9	7.5	7.8	9.2	5.4	8.6	7.3	7.0	7.0	4.9	§	2.4	-2.5 s
Complete 4 years	5.0	5.7	6.4	7.4	6.8	7.0	7.0	5.8	5.1	4.0	3.5	§	1.5	-2.0 sss
Region:														
Northeast	5.1	7.7	7.2	7.6	5.2	7.2	6.9	5.1	5.9	5.6	3.7	§	2.5	-1.2
Midwest	7.9	7.3	7.7	10.1	8.7	9.2	10.3	7.6	5.7	4.8	3.7	§	1.1	-2.6 ss
South	5.1	6.2	6.3	6.4	8.3	7.0	6.8	6.0	5.7	4.6	3.6	§	1.9	-1.7 s
West	3.2	5.3	5.3	6.6	6.3	3.9	6.4	5.7	4.9	3.7	4.9	§	1.7	-3.2 s
Population Density:														
Large MSA	5.9	7.0	4.7	9.1	7.2	7.8	7.5	6.8	5.0	4.2	3.6	§	2.2	-1.4
Other MSA	5.5	6.5	7.9	7.4	8.1	7.2	7.3	6.1	5.9	4.8	4.4	§	1.7	-2.7 sss
Non-MSA	4.5	6.0	6.1	6.2	5.7	4.3	8.0	5.2	5.6	4.8	2.9	§	1.0	-1.9 s
Parental Education: ^b														
1.0–2.0 (Low)	4.0	5.2	5.6	5.8	7.0	2.7	5.7	4.9	3.7	3.0	4.1	§	1.7	-2.4
2.5–3.0	5.7	6.0	7.2	6.3	5.7	7.3	7.0	4.5	6.9	4.6	3.1	§	1.4	-1.7 s
3.5–4.0	6.1	6.7	5.9	8.4	8.4	6.6	7.9	7.5	5.7	4.8	3.4	§	1.6	-1.8 s
4.5–5.0	5.5	7.0	7.1	9.4	7.7	7.7	7.8	6.9	3.6	4.4	3.9	§	1.6	-2.3 s
5.5–6.0 (High)	4.7	7.0	6.2	6.6	8.6	8.8	8.5	5.8	8.5	6.0	6.3	§	3.0	-3.3 s
Race/Ethnicity (2-year average): ^c														
White	—	7.5	8.1	8.8	9.1	8.1	8.4	8.6	7.7	6.6	5.1	§	3.0	-2.2 ss
African American	—	2.9	2.2	2.1	3.6	4.6	5.9	5.1	2.2	1.6	1.7	§	2.2	+0.5
Hispanic	—	3.1	2.9	3.7	4.8	4.9	4.4	4.7	4.3	3.6	3.1	§	3.6	+0.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table D-S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 74
PROVIGIL: ^a Trends in Annual Prevalence of Use
 by Subgroups in Grade 12

	Percentage who used in last 12 months													2019– 2021 change
	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	13,700	14,400	14,100	—	—	—	—	—	—	—	—	—	—	—
Total	1.8	1.3	1.5	—	—	—	—	—	—	—	—	—	—	—
Gender:														
Male	2.3	1.8	1.4	—	—	—	—	—	—	—	—	—	—	—
Female	1.3	0.6	1.3	—	—	—	—	—	—	—	—	—	—	—
College Plans:														
None or under 4 years	4.0	4.0	2.5	—	—	—	—	—	—	—	—	—	—	—
Complete 4 years	1.3	0.8	1.1	—	—	—	—	—	—	—	—	—	—	—
Region:														
Northeast	1.9	1.7	2.0	—	—	—	—	—	—	—	—	—	—	—
Midwest	2.0	1.7	1.0	—	—	—	—	—	—	—	—	—	—	—
South	1.9	1.2	1.7	—	—	—	—	—	—	—	—	—	—	—
West	1.4	0.9	1.2	—	—	—	—	—	—	—	—	—	—	—
Population Density:														
Large MSA	1.9	1.0	1.4	—	—	—	—	—	—	—	—	—	—	—
Other MSA	1.7	1.5	1.3	—	—	—	—	—	—	—	—	—	—	—
Non-MSA	2.0	1.6	1.8	—	—	—	—	—	—	—	—	—	—	—
Parental Education: ^b														
1.0–2.0 (Low)	3.8	1.7	2.4	—	—	—	—	—	—	—	—	—	—	—
2.5–3.0	2.0	1.6	2.0	—	—	—	—	—	—	—	—	—	—	—
3.5–4.0	1.7	1.2	1.6	—	—	—	—	—	—	—	—	—	—	—
4.5–5.0	1.2	0.8	0.5	—	—	—	—	—	—	—	—	—	—	—
5.5–6.0 (High)	1.1	1.8	0.9	—	—	—	—	—	—	—	—	—	—	—
Race/Ethnicity (2-year data): ^c														
White	—	1.5	1.3	—	—	—	—	—	—	—	—	—	—	—
African American	—	1.5	1.5	—	—	—	—	—	—	—	—	—	—	—
Hispanic	—	1.5	1.2	—	—	—	—	—	—	—	—	—	—	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on two of six forms; *N* is two sixths of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE 75
METHAMPHETAMINE: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																								2019– 2021 change
	1991– 1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	3.2	2.5	2.8	2.2	2.5	1.5	1.8	1.8	1.1	1.2	1.0	1.2	0.8	1.0	1.0	0.6	0.5	0.4	0.5	0.4	0.5	0.5	0.2	-0.3* s
Gender:																									
Male	—	3.1	2.2	2.5	2.0	2.0	1.1	1.7	1.4	0.7	0.9	1.0	1.3	0.5	0.8	0.7	0.5	0.5	0.4	0.4	0.3	0.3	§	0.1	-0.2
Female	—	3.2	2.8	3.0	2.4	3.0	1.7	1.9	2.3	1.5	1.5	1.0	1.1	1.0	1.1	1.0	0.6	0.5	0.4	0.6	0.5	0.7	§	0.2	-0.5 s
College Plans:																									
None or under 4 years	—	7.4	6.8	6.6	6.6	7.9	6.6	6.8	6.3	3.2	3.8	3.1	2.7	3.6	5.0	4.4	3.0	1.2	1.9	1.8	1.8	2.1	§	0.1	-2.0 s
Complete 4 years	—	2.7	2.0	2.3	1.7	2.0	1.0	1.2	1.3	0.9	1.0	0.9	1.1	0.6	0.7	0.7	0.4	0.4	0.2	0.3	0.2	0.3	§	0.2	-0.1
Region:																									
Northeast	—	1.6	2.0	1.2	0.8	1.7	1.0	0.6	0.7	0.5	0.8	0.3	0.6	0.6	0.8	0.4	0.5	0.2	0.2	0.0	0.4	0.4	§	0.1	-0.3
Midwest	—	4.4	3.0	3.2	2.5	3.5	2.1	2.0	1.7	1.5	1.6	0.9	1.6	0.5	0.3	0.7	0.8	0.8	0.4	0.4	0.1	0.1	§	0.3	+0.2
South	—	3.4	2.4	3.4	2.8	2.4	1.4	2.0	1.6	1.1	1.4	1.1	1.0	0.9	1.2	1.0	0.4	0.4	0.5	0.6	0.6	0.5	§	0.2	-0.4
West	—	2.8	2.5	2.4	2.0	2.4	1.6	2.1	3.1	1.2	1.0	1.4	1.6	1.1	1.4	1.5	0.9	0.6	0.4	0.8	0.5	0.7	§	0.0	-0.7
Population Density:																									
Large MSA	—	2.2	2.1	1.6	1.3	2.0	1.5	1.3	1.5	0.4	0.7	1.1	0.9	0.6	0.9	1.5	0.7	0.3	0.3	0.4	0.5	0.3	§	0.0	-0.3
Other MSA	—	3.6	2.6	3.4	2.5	2.4	1.7	2.0	2.1	1.4	1.5	1.0	1.3	1.2	1.2	0.6	0.6	0.6	0.3	0.6	0.5	0.7	§	0.3	-0.4
Non-MSA	—	3.5	3.0	2.8	2.7	3.4	1.2	2.1	1.8	1.6	1.4	0.9	1.3	0.3	0.7	0.9	0.5	0.6	0.7	0.5	0.1	0.2	§	0.1	-0.1
Parental Education: ^a																									
1.0–2.0 (Low)	—	6.3	5.0	4.4	4.3	4.2	4.4	4.1	3.9	1.9	2.1	1.7	2.2	3.1	1.5	2.1	0.2	0.4	0.8	1.2	1.4	1.3	§	1.0	-0.2
2.5–3.0	—	4.3	3.1	4.0	3.0	3.4	1.3	1.5	1.8	1.4	1.7	0.9	1.0	1.5	1.3	0.7	0.8	0.2	0.2	0.5	0.7	0.9	§	0.3	-0.7
3.5–4.0	—	3.3	2.4	3.0	2.0	2.5	1.8	2.9	2.2	1.2	1.3	1.1	1.3	0.5	1.4	0.7	0.6	0.0	0.5	0.6	0.6	0.4	§	0.3	-0.1
4.5–5.0	—	1.9	1.3	1.6	1.9	1.6	1.0	0.9	1.2	0.9	0.9	0.6	1.1	0.4	0.6	0.7	0.6	0.5	0.2	0.3	0.1	0.2	§	0.0	-0.2
5.5–6.0 (High)	—	3.2	2.2	1.2	1.1	2.3	0.9	0.7	1.7	0.8	1.1	1.6	1.1	0.4	0.7	0.8	0.2	0.6	0.2	0.2	0.0	0.5	§	0.0	-0.5
Race/Ethnicity (2-year average): ^b																									
White	—	—	3.2	3.1	2.8	2.7	2.2	1.6	1.6	1.4	1.1	1.0	1.0	0.8	0.7	0.7	0.6	0.6	0.5	0.3	0.2	0.2	§	0.2	+0.1
African American	—	—	1.1	0.8	0.6	0.8	0.6	0.5	0.8	0.6	0.5	0.7	0.5	0.6	0.6	0.4	0.3	0.3	0.2	0.2	0.3	0.3	§	1.2	+0.9
Hispanic	—	—	4.0	3.3	3.2	3.2	3.0	2.8	2.8	2.3	1.7	1.5	1.8	1.9	1.5	1.5	1.1	0.4	0.6	0.7	1.0	1.3	§	0.1	-1.2 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 76
METHAMPHETAMINE: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																							2019– 2021 change	
	1991– 1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
Approximate weighted N =	—	14,000	13,600	14,000	14,300	15,800	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	4.6	4.0	3.7	3.9	3.3	3.0	2.9	1.8	1.6	1.5	1.6	1.6	1.4	1.0	1.0	0.8	0.8	0.4	0.4	0.4	0.5	0.3	0.2	-0.3*
Gender:																									
Male	—	4.5	4.5	3.8	3.9	3.0	2.7	2.6	1.7	1.4	1.4	1.4	1.5	1.3	1.1	1.0	0.7	0.6	0.4	0.4	0.4	0.5	§	0.1	-0.4
Female	—	4.7	3.6	3.5	3.6	3.7	3.3	3.0	2.0	1.8	1.6	1.7	1.8	1.4	0.9	0.8	0.8	0.9	0.5	0.3	0.4	0.3	§	0.1	-0.2
College Plans:																									
None or under 4 years	—	9.1	8.8	7.0	9.1	9.4	6.0	7.3	5.8	3.9	3.6	4.6	4.8	4.6	4.1	2.4	3.6	3.4	2.2	1.2	1.5	0.2	§	0.5	+0.3
Complete 4 years	—	3.9	3.3	3.1	3.0	2.3	2.6	2.2	1.3	1.2	1.2	1.3	1.2	1.0	0.7	0.9	0.5	0.5	0.2	0.2	0.2	0.4	§	0.0	-0.4
Region:																									
Northeast	—	5.1	4.1	2.3	1.5	2.1	2.5	3.0	1.5	1.4	0.5	1.3	1.4	2.0	1.0	0.4	0.1	0.3	0.2	0.1	0.3	0.1	§	0.0	-0.1
Midwest	—	4.6	4.0	3.8	3.0	3.4	2.2	3.4	1.7	0.9	1.4	1.8	1.1	0.9	0.5	0.7	0.9	0.6	0.9	0.8	0.3	0.6	§	0.3	-0.4
South	—	4.0	3.8	4.0	4.6	3.8	2.9	2.2	2.3	1.9	1.9	1.4	1.5	1.4	1.0	0.9	1.2	1.2	0.3	0.2	0.6	0.1	§	0.1	-0.1
West	—	5.1	4.4	4.6	5.8	3.8	4.7	3.0	1.7	2.3	1.8	2.1	2.6	1.4	1.4	1.9	0.6	0.8	0.3	0.3	0.4	1.0	§	0.3	-0.8
Population Density:																									
Large MSA	—	3.8	4.4	4.0	3.1	2.3	2.7	2.9	2.2	1.7	1.2	1.5	1.5	1.0	1.1	1.0	0.7	0.7	0.1	0.3	0.4	0.2	§	0.3	0.0
Other MSA	—	4.8	3.4	3.6	4.4	3.8	3.4	3.0	1.5	1.5	1.8	1.6	1.9	1.7	0.6	1.2	0.5	0.8	0.4	0.4	0.3	0.6	§	0.1	-0.5
Non-MSA	—	5.2	4.8	3.6	3.7	3.7	2.5	2.5	2.1	1.8	1.3	1.6	1.3	1.4	1.7	0.6	1.6	0.9	1.3	0.4	0.7	0.5	§	0.1	-0.4
Parental Education: ^a																									
1.0–2.0 (Low)	—	7.2	7.0	5.6	6.2	7.3	4.5	4.2	3.8	3.8	2.6	3.7	3.4	2.9	2.5	1.9	1.2	1.0	0.7	0.3	0.7	0.4	§	0.0	-0.4
2.5–3.0	—	4.2	4.3	4.6	4.9	4.1	3.7	3.8	2.7	1.6	2.3	2.7	2.3	1.2	1.7	1.4	0.4	1.0	0.3	0.9	0.3	0.1	§	0.3	+0.2
3.5–4.0	—	4.8	4.3	3.2	3.7	3.2	3.3	2.0	1.4	1.7	1.0	1.4	1.9	1.3	0.7	1.0	1.2	1.1	0.5	0.1	0.7	0.3	§	0.1	-0.2
4.5–5.0	—	4.8	3.5	3.6	2.4	2.2	2.3	3.2	1.4	1.0	1.6	0.7	0.8	1.3	0.8	0.3	0.4	0.6	0.2	0.2	0.1	0.3	§	0.1	-0.2
5.5–6.0 (High)	—	3.3	3.5	2.5	4.0	2.1	1.7	1.3	1.3	1.0	0.6	0.7	0.8	1.0	0.3	1.2	0.6	0.3	0.6	0.1	0.7	1.2	§	0.0	-1.2
Race/Ethnicity (2-year average): ^b																									
White	—	—	4.7	4.1	4.1	4.2	3.4	2.9	2.4	1.8	1.6	1.4	1.5	1.5	1.2	0.9	0.9	0.8	0.6	0.4	0.3	0.4	§	0.1	-0.2
African American	—	—	0.4	1.0	1.1	0.6	0.8	1.1	1.0	0.8	0.5	0.4	0.6	0.7	0.3	0.3	0.6	0.9	0.4	0.5	0.4	0.0	§	0.4	+0.3
Hispanic	—	—	3.9	3.0	4.4	4.6	4.6	5.0	3.3	1.9	2.3	2.7	2.5	1.9	1.8	2.1	1.2	1.0	0.8	0.3	0.4	0.3	§	0.0	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 77
METHAMPHETAMINE: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																							2019– 2021 change		
	1975– 1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021	
Approximate weighted N =	—	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	4.7	4.3	3.9	3.6	3.2	3.4	2.5	2.5	1.7	1.2	1.2	1.0	1.4	1.1	0.9	1.0	0.6	0.6	0.6	0.5	0.5	1.4	0.2	-0.3* s	
Gender:																										
Male	—	5.0	4.4	4.3	4.3	3.6	3.7	2.7	2.0	1.8	1.8	1.3	1.5	1.6	1.2	1.3	1.3	0.7	0.9	0.6	0.7	0.7	§	0.1	-0.5 ss	
Female	—	4.5	4.2	3.4	3.0	2.9	3.1	2.3	3.0	1.6	0.7	1.0	0.5	1.1	0.9	0.4	0.7	0.4	0.4	0.5	0.2	0.2	§	0.1	0.0	
College Plans:																										
None or under 4 years	—	6.8	6.2	6.9	6.5	5.8	6.7	4.9	3.7	3.9	1.9	3.1	1.8	1.9	2.3	1.6	1.3	2.0	1.3	1.6	1.0	0.8	§	0.2	-0.7 s	
Complete 4 years	—	4.0	3.8	3.2	2.7	2.4	2.4	2.0	2.1	1.2	1.0	0.8	0.8	1.3	0.8	0.6	0.9	0.3	0.5	0.4	0.4	0.3	§	0.1	-0.2	
Region:																										
Northeast	—	3.1	4.5	3.4	1.6	1.8	1.3	0.6	1.4	1.4	1.1	1.0	1.2	2.1	0.5	1.3	0.9	0.0	0.6	0.6	0.1	0.1	§	0.1	0.0	
Midwest	—	5.1	4.1	4.7	4.5	4.2	3.3	4.4	1.2	1.2	0.8	1.2	1.1	1.2	0.9	0.8	0.7	1.2	0.4	0.6	1.3	0.5	§	0.0	-0.5	
South	—	3.9	3.7	3.4	3.0	2.8	4.0	2.2	3.1	2.1	1.2	1.4	0.9	1.1	0.9	0.5	1.2	0.4	0.4	0.6	0.5	0.7	§	0.3	-0.3	
West	—	7.1	5.4	4.4	5.4	4.0	4.9	2.7	3.8	1.5	1.9	1.0	1.2	1.5	1.8	1.4	0.7	0.7	1.4	0.7	0.2	0.2	§	0.2	-0.1	
Population Density:																										
Large MSA	—	4.2	3.6	2.7	2.1	1.8	2.8	1.5	3.4	1.3	1.3	1.1	1.0	0.9	1.2	0.8	0.9	0.5	0.7	0.6	0.4	0.6	§	0.1	-0.4	
Other MSA	—	4.0	4.9	4.0	4.4	3.0	2.6	2.4	2.0	1.7	1.2	1.1	1.1	2.0	1.2	0.9	1.0	0.5	0.8	0.7	0.6	0.3	§	0.3	0.0	
Non-MSA	—	6.4	4.3	5.3	4.1	5.3	5.5	4.1	2.0	2.2	1.3	1.4	1.0	0.6	0.5	1.2	0.8	1.0	0.2	0.4	0.6	0.7	§	0.1	-0.7 s	
Parental Education: ^a																										
1.0–2.0 (Low)	—	6.0	7.7	5.0	3.5	4.3	5.5	4.0	2.7	3.1	1.6	3.5	2.0	3.1	1.4	1.4	1.6	1.1	2.2	1.0	0.9	1.0	§	0.0	-1.0	
2.5–3.0	—	4.8	3.5	4.8	4.9	3.6	4.1	3.0	2.4	2.2	0.6	1.4	1.1	1.0	0.6	1.2	1.1	0.8	0.3	0.2	0.4	0.3	§	0.1	-0.3	
3.5–4.0	—	5.2	3.9	3.8	3.8	3.6	3.1	3.0	2.9	1.4	1.3	0.8	0.9	1.2	1.3	0.6	0.7	0.5	0.3	0.8	0.4	0.4	§	0.5	+0.1	
4.5–5.0	—	3.7	4.2	3.1	2.9	2.9	2.8	2.1	2.2	1.7	0.8	1.0	1.0	1.2	1.2	0.7	0.9	0.6	0.4	0.6	0.6	0.1	§	0.0	-0.1	
5.5–6.0 (High)	—	4.2	5.5	3.2	3.1	1.0	3.0	1.2	1.2	0.8	1.6	0.4	0.4	1.3	0.7	0.9	0.9	0.2	0.7	0.1	0.3	0.3	§	0.4	+0.1	
Race/Ethnicity (2-year average): ^b																										
White	—	—	4.9	4.6	4.2	3.5	3.5	3.3	2.6	2.1	1.5	1.1	1.1	1.3	1.2	0.9	0.7	0.6	0.4	0.5	0.7	0.4	§	0.3	-0.1	
African American	—	—	1.1	1.0	0.5	1.4	1.8	0.7	0.4	0.7	1.1	1.1	0.5	0.4	0.5	0.4	1.1	1.0	0.3	0.2	0.2	0.6	§	0.9	+0.3	
Hispanic	—	—	4.9	4.6	3.9	3.4	3.4	3.4	3.7	2.7	1.1	1.2	1.1	1.2	1.2	1.2	1.5	1.0	0.8	0.8	0.5	0.5	§	2.0	+1.5	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 78
CRYSTAL METHAMPHETAMINE (ICE): Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																
	1975- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Approximate weighted N =	—	15,200	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700
Total	—	1.3	1.4	1.3	1.7	1.8	2.4	2.8	2.3	3.0	1.9	2.2	2.5	3.1	2.0	2.1	2.3
Gender:																	
Male	—	1.5	1.9	1.5	1.9	2.2	2.6	3.9	2.6	3.9	2.2	2.5	2.7	3.5	2.5	2.6	2.5
Female	—	1.0	0.9	1.0	1.2	1.3	2.1	1.7	2.1	2.1	1.6	1.9	2.1	2.5	1.5	1.6	2.0
College Plans:																	
None or under 4 years	—	1.4	1.9	2.2	2.6	2.4	4.3	5.1	3.8	5.0	2.3	3.8	4.2	4.8	2.6	4.4	4.2
Complete 4 years	—	1.2	1.2	1.0	1.4	1.5	1.8	2.1	1.9	2.4	1.8	1.7	2.0	2.6	1.9	1.5	1.8
Region:																	
Northeast	—	1.6	1.1	0.9	1.1	0.9	1.4	3.2	2.8	2.6	1.0	1.0	1.9	2.6	1.3	0.9	1.2
Midwest	—	1.1	1.4	1.1	1.5	2.3	2.2	2.1	1.9	2.7	1.7	2.8	2.1	2.0	1.0	1.8	2.2
South	—	0.5	1.0	1.0	1.2	1.2	1.8	2.0	1.6	3.1	2.3	2.0	1.7	3.5	2.3	2.3	2.7
West	—	2.5	2.2	2.6	3.2	2.8	4.7	4.9	3.7	3.4	2.5	2.9	4.7	4.0	3.4	3.5	3.1
Population Density:																	
Large MSA	—	1.2	1.3	1.5	2.2	2.0	3.0	4.6	2.9	2.5	1.6	2.3	2.7	2.2	1.9	2.3	2.1
Other MSA	—	1.3	1.7	1.3	1.7	1.8	2.0	2.0	2.0	3.3	2.1	1.7	2.3	3.6	2.0	1.4	2.3
Non-MSA	—	1.2	0.8	1.2	1.2	1.6	2.3	2.6	2.1	2.8	2.0	3.0	2.5	3.2	2.1	3.2	2.7
Parental Education: ^a																	
1.0-2.0 (Low)	—	1.1	1.0	0.9	1.6	1.0	3.2	3.4	2.2	2.8	3.6	3.2	5.9	3.7	3.7	4.0	2.6
2.5-3.0	—	1.4	1.1	1.1	1.2	3.0	2.6	3.0	2.5	3.2	1.3	1.7	2.4	3.6	1.7	2.7	2.6
3.5-4.0	—	1.4	1.2	1.5	2.3	1.6	1.9	2.6	2.2	3.6	2.5	1.9	2.8	2.4	2.2	2.4	3.1
4.5-5.0	—	1.2	1.8	1.4	1.8	1.3	2.5	2.9	2.9	2.6	1.3	2.8	1.5	2.8	1.5	1.0	1.4
5.5-6.0 (High)	—	1.1	1.0	1.4	1.3	1.0	2.2	2.5	1.4	1.9	2.1	1.3	2.0	2.8	1.8	0.9	1.1
Race/Ethnicity (2-year average): ^b																	
White	—	—	1.3	1.3	1.5	1.7	2.0	2.5	2.6	2.9	2.8	2.3	2.4	2.6	2.3	2.0	2.0
African American	—	—	0.8	1.0	0.8	0.5	0.5	0.3	0.5	1.0	0.7	0.7	1.1	1.6	1.4	0.7	1.0
Hispanic	—	—	1.3	1.8	2.3	2.1	2.7	4.0	2.8	1.7	1.7	2.4	3.4	3.6	2.5	2.6	3.4

Table continued on next page.

TABLE 78 (cont.)
CRYSTAL METHAMPHETAMINE (ICE): Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	1.9	1.6	1.1	0.9	0.9	1.2	0.8	1.1	0.8	0.5	0.8	0.8	0.6	0.6	0.0	0.4	-0.3*
Gender:																	
Male	1.5	1.8	1.4	1.1	1.1	1.0	1.0	1.3	0.8	0.5	1.0	0.4	0.6	0.7	§	0.4	-0.2
Female	2.2	1.4	0.7	0.8	0.6	1.1	0.7	0.8	0.7	0.5	0.6	1.2	0.2	0.4	§	0.2	-0.2
College Plans:																	
None or under 4 years	3.5	3.7	1.7	1.8	2.1	2.9	1.9	2.4	1.7	1.0	1.6	1.3	1.4	1.6	§	0.8	-0.8
Complete 4 years	1.3	1.2	0.8	0.7	0.6	0.9	0.5	0.8	0.5	0.4	0.6	0.7	0.2	0.4	§	0.2	-0.2
Region:																	
Northeast	1.0	1.1	1.4	0.5	0.6	0.4	0.9	1.0	0.3	0.4	0.4	0.5	0.5	0.4	§	0.1	-0.3
Midwest	1.0	1.5	0.5	1.0	0.9	0.8	1.0	0.6	0.9	1.2	0.8	0.6	0.6	0.6	§	0.1	-0.5
South	2.9	1.7	0.9	0.8	0.9	1.4	0.6	1.1	0.8	0.4	1.0	1.1	0.7	0.7	§	0.7	0.0
West	1.9	2.1	1.7	1.5	1.2	1.8	0.9	1.6	1.1	0.4	1.0	0.6	0.4	0.7	§	0.2	-0.5
Population Density:																	
Large MSA	2.2	1.3	0.6	0.9	0.9	1.1	1.2	1.3	1.2	0.8	0.8	0.7	0.6	0.9	§	0.3	-0.6 s
Other MSA	1.5	1.7	1.2	0.8	0.9	1.3	0.7	1.2	0.7	0.3	0.7	0.7	0.2	0.4	§	0.5	+0.1
Non-MSA	2.1	1.9	1.3	1.3	1.1	1.0	0.7	0.7	0.6	0.7	1.3	1.1	1.2	0.8	§	0.2	-0.6
Parental Education: ^a																	
1.0–2.0 (Low)	4.2	3.2	1.4	2.8	1.9	3.6	1.1	3.8	3.0	1.4	1.2	1.6	0.9	0.7	§	0.4	-0.3
2.5–3.0	2.6	2.1	0.6	0.8	1.6	0.3	1.6	1.3	0.8	1.0	0.7	0.7	0.4	1.2	§	0.1	-1.1 s
3.5–4.0	1.4	1.4	1.0	1.2	0.4	1.5	0.6	0.9	0.6	0.4	1.0	1.2	0.7	0.6	§	0.7	+0.1
4.5–5.0	1.0	1.5	0.7	0.3	0.5	0.5	0.4	0.4	0.3	0.3	0.6	0.2	0.4	0.3	§	0.3	+0.1
5.5–6.0 (High)	1.4	0.8	1.2	0.6	0.7	1.3	0.2	0.4	0.2	0.0	1.0	0.5	0.3	0.3	§	0.1	-0.1
Race/Ethnicity (2-year average): ^b																	
White	1.9	1.7	1.3	0.7	0.5	0.6	0.7	0.6	0.6	0.3	0.3	0.5	0.4	0.3	§	0.1	-0.2
African American	1.0	0.6	0.6	0.7	0.9	1.1	0.8	1.0	1.3	0.6	0.6	0.8	0.8	1.5	§	0.5	-0.9
Hispanic	3.3	2.9	1.8	1.4	1.7	2.4	2.1	1.4	2.0	2.0	1.6	1.2	0.9	0.6	§	0.0	-0.6 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 79
BATH SALTS (SYNTHETIC STIMULANTS): Trends in
Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months											
	1991–											2019–
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2021
<i>Approximate weighted N =</i>	—	15,100	14,600	14,600	14,400	16,900	15,300	14,000	—	—	—	
Total	—	0.8	1.0	0.5	0.4	0.9	0.5	0.9	—	—	—	—
Gender:												
Male	—	0.6	1.1	0.5	0.5	1.2	0.9	1.4	—	—	—	—
Female	—	0.9	0.9	0.5	0.3	0.7	0.3	0.4	—	—	—	—
College Plans:												
None or under 4 years	—	3.6	3.1	2.3	1.0	3.2	2.5	3.3	—	—	—	—
Complete 4 years	—	0.6	0.8	0.4	0.3	0.6	0.3	0.6	—	—	—	—
Region:												
Northeast	—	0.3	0.7	0.9	0.1	0.5	0.4	0.6	—	—	—	—
Midwest	—	0.5	0.6	0.3	0.2	0.7	0.3	0.2	—	—	—	—
South	—	1.3	1.0	0.5	0.6	1.0	0.5	1.6	—	—	—	—
West	—	0.7	1.5	0.4	0.3	1.1	0.9	0.7	—	—	—	—
Population Density:												
Large MSA	—	1.0	1.4	0.8	0.5	1.2	0.5	1.1	—	—	—	—
Other MSA	—	0.7	0.8	0.4	0.3	0.7	0.6	0.8	—	—	—	—
Non-MSA	—	0.8	0.8	0.3	0.3	1.0	0.4	0.8	—	—	—	—
Parental Education: ^a												
1.0–2.0 (Low)	—	0.9	2.7	1.0	0.5	2.5	1.6	1.1	—	—	—	—
2.5–3.0	—	0.6	0.6	0.4	0.4	0.6	0.4	0.3	—	—	—	—
3.5–4.0	—	0.8	0.8	0.2	0.5	1.3	0.4	1.1	—	—	—	—
4.5–5.0	—	0.6	1.0	0.7	0.3	0.3	0.4	0.8	—	—	—	—
5.5–6.0 (High)	—	0.8	0.4	0.6	0.2	0.5	0.5	0.9	—	—	—	—
Race/Ethnicity (2-year data): ^b												
White	—	—	0.5	0.4	0.3	0.4	0.4	0.4	—	—	—	—
African American	—	—	1.3	1.0	0.7	1.4	1.6	1.9	—	—	—	—
Hispanic	—	—	1.7	1.3	0.5	0.8	0.8	0.9	—	—	—	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE 80
BATH SALTS (SYNTHETIC STIMULANTS): Trends in
Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months											
	1991–											2019–
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2021
												change
<i>Approximate weighted N =</i>	—	15,000	12,900	13,000	15,600	14,700	13,500	14,300	—	—	—	
Total	—	0.6	0.9	0.9	0.7	0.8	0.4	0.5	—	—	—	—
Gender:												
Male	—	0.9	1.3	1.0	0.8	1.1	0.4	0.7	—	—	—	—
Female	—	0.4	0.6	0.7	0.5	0.4	0.4	0.3	—	—	—	—
College Plans:												
None or under 4 years	—	1.9	3.7	3.6	3.6	2.4	0.8	1.6	—	—	—	—
Complete 4 years	—	0.5	0.6	0.5	0.4	0.6	0.4	0.4	—	—	—	—
Region:												
Northeast	—	0.7	1.4	0.6	0.4	1.3	0.2	0.7	—	—	—	—
Midwest	—	0.8	0.8	0.7	0.4	0.7	0.4	0.4	—	—	—	—
South	—	0.5	0.8	1.0	0.9	0.6	0.3	0.6	—	—	—	—
West	—	0.6	0.9	1.1	0.9	0.7	0.9	0.6	—	—	—	—
Population Density:												
Large MSA	—	1.0	0.7	1.0	0.5	0.4	0.4	0.6	—	—	—	—
Other MSA	—	0.3	0.8	0.6	0.7	0.8	0.3	0.5	—	—	—	—
Non-MSA	—	0.8	1.7	1.4	0.9	1.5	0.7	0.5	—	—	—	—
Parental Education: ^a												
1.0–2.0 (Low)	—	0.7	0.4	0.9	1.8	1.0	1.0	1.2	—	—	—	—
2.5–3.0	—	0.9	2.0	0.6	1.2	0.7	0.4	0.5	—	—	—	—
3.5–4.0	—	0.4	0.6	1.5	0.4	1.0	0.6	0.5	—	—	—	—
4.5–5.0	—	0.7	0.4	0.4	0.3	0.3	0.2	0.4	—	—	—	—
5.5–6.0 (High)	—	0.2	1.1	0.9	0.5	1.0	0.1	0.2	—	—	—	—
Race/Ethnicity (2-year data): ^b												
White	—	—	0.7	0.8	0.6	0.6	0.6	0.3	—	—	—	—
African American	—	—	0.8	1.4	1.7	1.3	0.9	0.5	—	—	—	—
Hispanic	—	—	0.7	0.6	0.5	0.8	0.9	0.9	—	—	—	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE 81
BATH SALTS (SYNTHETIC STIMULANTS): Trends in
Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months											
	1991–											2019–
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	—	13,700	12,600	12,400	12,900	11,800	12,600	13,300	—	—	—	—
Total	—	1.3	0.9	0.9	1.0	0.8	0.6	0.6	—	—	—	—
Gender:												
Male	—	1.2	1.0	1.1	1.2	0.8	0.6	0.4	—	—	—	—
Female	—	1.1	0.6	0.5	0.6	0.3	0.7	0.5	—	—	—	—
College Plans:												
None or under 4 years	—	1.8	1.8	1.5	1.6	2.1	1.8	1.4	—	—	—	—
Complete 4 years	—	1.1	0.6	0.6	0.8	0.3	0.3	0.3	—	—	—	—
Region:												
Northeast	—	1.2	0.7	1.4	0.1	0.6	0.6	0.3	—	—	—	—
Midwest	—	1.3	1.0	0.7	1.5	1.0	0.4	0.5	—	—	—	—
South	—	1.7	1.0	1.2	1.3	0.9	0.7	0.8	—	—	—	—
West	—	1.0	0.8	0.2	0.7	0.4	0.6	0.5	—	—	—	—
Population Density:												
Large MSA	—	1.3	0.9	1.0	1.1	0.5	0.6	0.7	—	—	—	—
Other MSA	—	1.4	0.7	1.0	0.8	0.9	0.5	0.2	—	—	—	—
Non-MSA	—	1.2	1.3	0.5	1.4	0.7	0.9	1.2	—	—	—	—
Parental Education: ^a												
1.0–2.0 (Low)	—	2.1	1.7	2.0	1.6	1.9	1.5	1.2	—	—	—	—
2.5–3.0	—	1.7	0.9	1.5	0.9	1.0	0.9	1.0	—	—	—	—
3.5–4.0	—	0.8	1.0	0.3	0.6	0.6	0.4	0.1	—	—	—	—
4.5–5.0	—	1.2	0.6	0.5	1.0	0.3	0.4	0.4	—	—	—	—
5.5–6.0 (High)	—	1.3	0.2	0.2	0.4	0.1	0.2	0.3	—	—	—	—
Race/Ethnicity (2-year data): ^b												
White	—	—	1.0	0.5	0.6	0.7	0.5	0.4	—	—	—	—
African American	—	—	1.6	2.2	2.3	1.7	1.2	1.0	—	—	—	—
Hispanic	—	—	1.2	0.8	0.6	0.8	0.8	0.7	—	—	—	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms; *N* is two sixths of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE 82
SEDATIVES (BARBITURATES): ^{a,b} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	10.7	9.6	9.3	8.1	7.5	6.8	6.6	5.5	5.2	4.9	4.6	4.2	3.6	3.2	3.3	3.4
Gender:																
Male	12.3	9.9	10.2	8.4	7.6	7.3	7.2	5.9	5.9	5.5	5.2	4.7	4.0	3.4	3.5	3.8
Female	9.9	9.2	8.4	7.7	7.0	6.0	5.8	5.0	4.2	4.0	3.9	3.8	3.2	3.0	3.0	3.0
College Plans:																
None or under 4 years	—	11.6	11.4	9.1	9.3	9.0	8.1	7.4	6.7	6.2	6.2	6.1	4.7	4.1	4.8	4.7
Complete 4 years	—	7.3	6.8	6.8	5.2	4.8	5.1	3.8	3.8	3.7	3.6	3.0	3.0	2.7	2.5	2.8
Region:																
Northeast	11.5	10.4	9.2	9.6	9.6	6.9	6.8	5.6	4.7	5.1	5.3	5.2	4.2	2.5	3.2	2.9
Midwest	12.8	10.4	10.7	7.9	6.9	7.3	7.5	5.4	6.1	4.9	4.9	4.2	3.3	2.5	3.2	3.5
South	9.9	9.7	9.3	7.8	7.3	7.0	5.5	6.3	5.2	5.2	4.2	4.1	3.7	4.1	3.7	4.0
West	10.0	6.7	6.6	6.6	5.7	5.2	6.5	3.9	4.0	4.2	4.1	3.3	3.2	3.2	2.7	2.9
Population Density:																
Large MSA	11.1	10.2	8.1	8.1	8.3	6.6	6.9	5.3	5.2	4.4	4.4	3.7	3.3	2.8	2.6	2.6
Other MSA	11.3	9.8	9.9	8.2	7.3	6.5	6.4	5.7	5.3	4.9	4.2	4.4	3.6	3.4	3.1	3.6
Non-MSA	9.8	9.0	9.5	8.1	7.0	7.2	6.6	5.5	5.0	5.5	5.4	4.5	3.9	3.2	4.4	3.9
Parental Education: ^c																
1.0–2.0 (Low)	10.3	9.1	8.0	7.5	7.8	8.0	6.5	5.8	6.1	4.7	5.0	4.8	3.8	4.3	4.1	3.1
2.5–3.0	10.3	10.2	10.3	8.2	7.3	7.2	6.5	5.7	5.7	5.2	5.3	4.6	3.6	3.1	3.4	3.7
3.5–4.0	9.5	9.6	9.0	8.3	7.4	6.3	6.5	5.1	4.6	5.0	4.4	4.4	3.2	2.9	3.2	3.9
4.5–5.0	10.7	10.1	9.1	7.8	6.6	5.9	6.4	5.0	4.4	4.3	4.1	3.3	3.9	3.3	2.8	3.1
5.5–6.0 (High)	9.0	10.3	8.3	8.0	7.2	5.4	6.8	5.8	3.7	4.0	3.1	3.4	3.6	3.1	3.4	2.9
Race/Ethnicity (2-year average): ^d																
White	—	—	10.2	9.3	8.2	7.5	7.2	6.5	5.8	5.5	5.1	4.7	4.2	3.7	3.5	3.7
African American	—	—	3.3	3.2	2.6	2.5	2.4	2.0	1.7	1.6	1.6	1.6	1.7	1.5	1.1	1.1
Hispanic	—	—	7.4	5.8	5.8	5.8	5.7	5.1	4.1	4.4	4.6	3.6	2.8	2.8	3.2	2.8

Table continued on next page.

TABLE 82 (cont.)
SEDATIVES (BARBITURATES): ^{a,b} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	3.4	2.8	3.4	4.1	4.7	4.9	5.1	5.5	5.8	6.2	5.7	6.7	6.0	6.5	7.2	6.6
Gender:																
Male	3.4	2.9	3.4	4.3	5.1	5.2	5.3	6.3	6.3	6.9	6.3	7.4	6.7	7.0	7.7	6.4
Female	3.2	2.6	3.3	3.8	4.2	4.4	4.8	4.8	5.3	5.4	5.1	5.9	5.4	6.0	6.6	6.7
College Plans:																
None or under 4 years	4.3	3.9	3.8	5.4	5.9	6.7	6.6	6.9	7.0	7.0	7.1	9.8	9.6	9.3	10.7	10.0
Complete 4 years	2.9	2.3	3.2	3.7	4.4	4.3	4.6	5.1	5.5	6.0	5.4	6.0	5.1	5.6	6.3	5.7
Region:																
Northeast	2.8	2.7	3.5	4.0	4.1	5.6	5.7	5.5	4.7	5.6	5.3	5.9	4.8	5.4	5.7	5.7
Midwest	3.5	2.7	3.5	4.1	4.5	4.9	4.4	4.8	6.0	5.2	5.6	6.4	5.1	5.9	7.5	6.2
South	3.6	3.0	3.6	4.8	5.3	5.4	5.8	6.8	6.8	7.5	6.1	8.4	7.1	7.8	8.5	7.5
West	3.3	2.5	2.7	2.8	4.3	3.3	4.2	4.2	4.9	5.7	5.6	5.1	6.5	6.3	5.9	6.2
Population Density:																
Large MSA	2.4	2.4	2.6	3.6	4.1	3.6	4.2	4.6	3.7	5.1	5.9	6.5	4.2	5.5	6.1	6.6
Other MSA	3.9	2.6	3.1	4.3	4.9	5.4	5.0	5.6	6.2	6.6	6.1	6.7	6.6	7.1	7.5	6.6
Non-MSA	3.3	3.4	4.3	4.1	5.0	5.4	6.4	6.8	7.4	6.8	4.9	7.0	7.4	6.6	7.8	6.6
Parental Education: ^c																
1.0–2.0 (Low)	3.6	3.9	3.8	4.5	4.9	4.8	4.3	6.4	7.5	5.2	6.3	5.1	6.7	6.6	6.2	7.3
2.5–3.0	3.7	2.4	3.6	4.5	5.2	4.9	4.9	6.3	6.2	7.1	5.3	7.1	6.5	7.0	8.1	6.5
3.5–4.0	3.0	2.8	2.8	4.0	4.6	4.9	6.0	5.6	6.4	5.8	6.1	7.7	7.2	8.0	7.7	7.3
4.5–5.0	3.3	2.9	3.4	4.0	4.4	5.0	5.1	5.0	5.0	6.7	5.9	6.3	4.7	5.1	7.0	6.2
5.5–6.0 (High)	3.6	2.4	3.8	3.6	4.1	4.6	4.6	5.0	5.1	5.5	5.8	6.6	4.3	5.5	6.3	6.0
Race/Ethnicity (2-year average): ^d																
White	3.8	3.5	3.6	4.3	4.9	5.4	5.9	6.5	6.7	6.9	7.0	7.5	7.6	7.4	7.9	7.9
African American	1.2	1.1	1.0	1.5	1.6	1.1	1.0	1.4	1.4	1.3	1.5	1.4	1.7	2.4	2.6	2.5
Hispanic	2.4	2.2	1.9	2.6	3.5	4.0	3.7	3.3	4.0	5.1	4.7	4.5	4.1	4.7	5.8	5.8

Table continued on next page.

TABLE 82 (cont.)
SEDATIVES (BARBITURATES): ^{a,b} Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																2019-2021 change
	2007	2008	2009	2010	2011	2012	2013 ^e	2014	2015	2016	2017	2018	2019	2020	2021		
<i>Approximate weighted N =</i>																	
	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	6.2	5.8	5.2	4.8	4.3	4.5	4.8	4.3	3.6	3.0	2.9	2.7	2.5	2.4	1.8	-0.7* s	
Gender:																	
Male	6.3	5.9	5.6	4.8	4.2	4.2	4.7	4.4	3.1	2.7	2.9	2.4	2.5	§	2.0	-0.6	
Female	6.2	5.7	4.9	4.6	4.4	4.7	4.9	4.2	4.1	3.2	2.9	2.5	2.3	§	1.7	-0.6	
College Plans:																	
None or under 4 years	8.2	8.9	7.6	6.7	7.5	5.1	6.1	5.9	4.1	3.6	3.9	3.5	3.3	§	2.3	-1.0	
Complete 4 years	5.7	5.2	4.7	4.3	3.7	4.3	4.4	3.9	3.5	2.9	2.8	2.4	2.2	§	1.7	-0.4	
Region:																	
Northeast	5.5	5.1	4.3	4.2	3.4	4.1	3.9	3.0	3.3	2.2	1.6	2.5	1.7	§	1.7	0.0	
Midwest	5.9	5.7	6.5	5.3	4.7	4.4	4.7	4.1	2.9	3.7	2.7	2.9	2.1	§	0.8	-1.3 sss	
South	7.2	6.8	5.6	5.3	4.2	4.4	5.5	5.1	3.8	2.9	3.2	2.5	2.8	§	1.6	-1.2 ss	
West	5.4	5.0	4.0	4.2	4.7	5.0	4.5	4.1	4.3	3.2	3.7	2.7	3.1	§	3.1	0.0	
Population Density:																	
Large MSA	5.7	4.3	4.5	4.1	3.4	4.3	4.7	3.6	3.3	2.5	2.3	2.3	2.3	§	1.7	-0.6	
Other MSA	6.3	6.2	5.4	5.0	4.6	4.8	5.2	5.0	4.0	3.4	3.4	2.6	2.5	§	2.1	-0.4	
Non-MSA	6.9	6.7	5.9	5.5	4.9	4.0	3.9	3.5	3.3	3.0	3.1	3.2	2.7	§	1.2	-1.5 ss	
Parental Education: ^c																	
1.0–2.0 (Low)	6.3	6.0	4.8	4.3	4.8	5.1	4.9	4.3	3.3	3.4	3.4	2.8	2.5	§	1.2	-1.4	
2.5–3.0	5.7	6.3	5.7	5.5	4.6	5.0	5.0	4.8	3.2	3.0	3.2	2.5	2.7	§	1.6	-1.0	
3.5–4.0	6.5	6.2	5.5	5.2	4.6	4.8	5.1	4.0	4.6	3.4	3.3	2.9	2.5	§	1.8	-0.7	
4.5–5.0	7.0	5.0	5.6	4.5	4.2	4.3	4.6	4.6	3.1	2.7	2.2	2.8	2.2	§	1.6	-0.6	
5.5–6.0 (High)	5.3	5.4	3.5	3.6	3.0	3.7	4.6	3.7	4.1	3.3	2.8	2.1	2.3	§	3.4	+1.2	
Race/Ethnicity (2-year average): ^d																	
White	7.5	7.3	6.6	5.8	5.2	4.9	5.1	5.0	4.5	3.8	3.2	2.8	2.6	§	2.0	-0.6	
African American	2.5	2.2	2.0	2.7	2.5	2.1	2.5	3.1	2.6	2.0	1.7	1.5	2.0	§	2.7	+0.7	
Hispanic	5.2	4.3	4.0	3.8	3.5	3.7	3.8	3.8	3.3	2.7	3.1	2.9	2.4	§	1.4	-0.9	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bIn 2004 the barbiturate question text was changed on half of the questionnaire forms. Barbiturates was changed to Sedatives, including barbiturates, and "have you taken barbiturates . . ." was changed to "have you taken sedatives . . ." In the list of examples downs, downers, goofballs, yellow, reds, blues, rainbows were changed to downs, or downers, and include Phenobarbital, Tuinal, Nembutal, and Seconal. An examination of the data did not show any effect from the wording change. In 2005 the remaining forms were changed in a like manner.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^dEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^eIn 2013 the question text was changed in all forms: Tuinal, Nembutal, and Seconal were replaced with Ambien, Lunesta, and Sonata. In one form the list of examples was also changed: Tuinal was dropped from the list and Dalmane, Restoril, Halcion, Intermezzo, and Zolpimist were added. An examination of the data did not show any effect from the wording changes.

TABLE 83
TRANQUILIZERS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001^b</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>	
Total	1.8	2.0	2.1	2.4	2.7	3.3	2.9	2.6	2.5	2.6‡	2.8	2.6	2.7	2.5	2.8	
Gender:																
Male	1.5	1.6	1.8	1.9	2.0	2.3	2.6	2.3	2.1	2.1‡	2.5	2.2	2.2	1.9	2.1	
Female	2.1	2.3	2.4	2.8	3.3	4.0	3.2	3.0	2.9	3.1‡	2.9	2.8	3.0	3.2	3.4	
College Plans:																
None or under 4 years	3.9	4.9	3.6	5.1	5.9	6.4	5.8	6.4	5.8	6.8‡	5.8	6.5	6.3	6.2	6.9	
Complete 4 years	1.5	1.5	1.9	2.0	2.3	2.8	2.6	2.2	2.0	2.2‡	2.5	2.1	2.3	2.1	2.4	
Region:																
Northeast	1.0	1.6	1.7	2.5	2.3	2.7	2.5	2.0	2.3	2.2‡	2.0	1.8	1.9	1.7	1.6	
Midwest	1.4	1.9	1.3	1.7	2.6	3.5	2.4	2.6	2.6	2.2‡	2.2	2.4	2.2	2.1	2.8	
South	2.6	2.5	2.4	2.6	3.0	3.7	3.3	3.2	2.8	3.2‡	3.4	3.4	3.5	3.4	3.7	
West	1.8	1.6	3.0	2.7	2.4	2.9	3.0	2.3	1.9	2.5‡	3.2	1.9	2.6	2.2	2.2	
Population Density:																
Large MSA	1.8	2.1	1.7	2.5	1.8	3.2	2.1	1.9	1.8	2.5‡	2.8	2.0	1.8	1.8	2.0	
Other MSA	1.7	1.8	2.5	2.6	3.2	3.4	3.2	2.6	2.4	2.7‡	3.1	2.7	2.9	3.1	3.1	
Non-MSA	2.2	2.2	1.6	1.9	2.6	3.1	3.2	3.6	3.4	2.7‡	2.3	2.9	3.4	2.4	3.4	
Parental Education: ^c																
1.0–2.0 (Low)	3.6	3.8	2.5	3.2	3.9	4.9	4.3	5.5	5.0	5.6‡	3.7	5.0	3.9	4.7	5.0	
2.5–3.0	1.6	2.1	2.5	2.6	2.7	3.5	3.1	2.8	3.3	3.6‡	3.7	3.6	3.5	2.8	4.3	
3.5–4.0	2.0	2.2	2.1	2.6	3.2	3.9	3.2	2.7	2.1	2.6‡	2.7	2.2	2.8	3.0	3.3	
4.5–5.0	1.4	0.9	1.8	2.0	2.2	2.8	2.4	2.4	1.7	1.4‡	2.7	2.0	2.1	2.0	1.3	
5.5–6.0 (High)	1.8	1.9	1.7	2.1	1.6	2.7	2.7	2.1	1.9	2.2‡	2.1	1.6	1.6	1.7	2.0	
Race/Ethnicity (2-year average): ^d																
White	—	2.0	2.0	2.2	2.7	3.2	3.4	3.1	2.9	3.0	3.0‡	3.0 ^e	2.8	2.8	2.9	
African American	—	0.9	1.1	1.2	1.2	1.3	1.2	0.9	0.7	0.5	0.5‡	0.6 ^e	1.0	1.2	1.0	
Hispanic	—	2.7	3.1	3.4	3.3	3.5	3.5	3.4	3.5	3.5	3.6‡	3.8 ^e	3.4	3.4	3.5	

Table continued on next page.

TABLE 83 (cont.)
TRANQUILIZERS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	2.6	2.4	2.4	2.6	2.8	2.0	1.8	1.8	1.7	1.7	1.7	2.0	2.0	2.4	2.2	1.1	-1.3* sss
Gender:																	
Male	2.2	1.6	1.7	2.3	2.2	1.6	1.3	1.3	1.3	1.2	1.4	1.5	2.0	2.1	§	0.6	-1.5 sss
Female	2.9	3.1	3.2	3.0	3.3	2.4	2.3	2.4	2.1	2.2	1.9	2.4	2.1	2.7	§	1.5	-1.2 sss
College Plans:																	
None or under 4 years	6.0	6.8	5.9	7.0	5.5	4.8	4.1	5.3	4.8	3.1	3.3	4.2	4.0	5.1	§	2.1	-3.1 sss
Complete 4 years	2.3	2.0	2.2	2.3	2.6	1.8	1.6	1.6	1.5	1.6	1.5	1.8	1.8	2.0	§	0.9	-1.0 sss
Region:																	
Northeast	1.3	1.4	1.6	1.2	1.6	1.1	1.3	1.2	1.5	1.6	1.1	0.9	0.9	1.1	§	0.6	-0.4
Midwest	2.4	2.8	2.5	2.4	2.6	1.8	1.0	1.5	1.5	1.8	1.2	1.6	1.3	2.1	§	1.1	-1.0 ss
South	3.6	2.9	3.3	3.6	3.6	2.5	2.7	2.3	2.0	1.9	2.0	2.3	2.5	2.8	§	1.2	-1.5 sss
West	2.1	1.9	1.8	2.2	2.4	2.2	1.3	1.9	1.8	1.4	2.1	2.7	2.9	3.2	§	1.4	-1.8 sss
Population Density:																	
Large MSA	1.9	1.9	1.8	2.0	2.2	1.7	1.5	1.6	1.8	2.2	1.5	2.0	2.2	2.5	§	0.9	-1.6 sss
Other MSA	2.6	2.8	2.8	3.0	3.1	2.4	2.2	1.9	1.8	1.7	1.7	2.1	2.2	2.6	§	1.4	-1.1 sss
Non-MSA	3.5	2.4	2.6	2.5	2.8	1.6	1.5	1.9	1.6	1.0	1.8	1.7	1.5	1.9	§	0.9	-1.1 s
Parental Education: ^c																	
1.0–2.0 (Low)	5.1	5.2	4.7	5.9	4.8	3.4	3.2	3.6	3.0	3.1	2.5	4.4	3.2	3.4	§	1.1	-2.3 ss
2.5–3.0	3.2	3.3	3.0	3.4	3.4	2.8	2.4	2.4	2.3	1.7	1.7	2.4	2.6	4.1	§	1.0	-3.1 sss
3.5–4.0	3.1	2.9	2.5	2.8	3.2	2.5	1.9	2.2	1.9	1.8	2.0	1.7	2.4	2.3	§	1.4	-0.9 s
4.5–5.0	1.7	1.3	2.1	1.7	2.0	1.3	1.1	1.5	1.5	1.5	1.7	1.5	1.9	2.0	§	1.5	-0.6
5.5–6.0 (High)	1.8	1.0	1.7	1.4	1.7	1.3	1.5	0.9	1.1	1.3	1.2	1.5	1.1	1.7	§	0.8	-1.0 s
Race/Ethnicity (2-year average): ^d																	
White	3.0	2.7	2.6	2.8	2.9	2.6	2.0	1.7	1.7	1.7	1.4	1.5	1.6	1.8	§	2.1	+0.2
African American	0.8	0.9	0.9	1.0	1.2	1.3	1.2	1.1	1.1	1.2	1.5	1.4	1.0	1.0	§	0.1	-0.9 sss
Hispanic	3.0	2.9	2.9	2.9	3.3	2.9	2.4	2.4	2.2	2.0	2.1	2.3	2.6	3.0	§	1.2	-1.9 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. * ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bIn 2001, for the tranquilizer list of examples, Miltown was replaced with Xanax. The 2001 data are based on the changed forms only; N is one half of N indicated.

In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^dEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^eDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 84
TRANQUILIZERS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^b	2002	2003	2004	2005	
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
Total	3.2	3.5	3.3	3.3	4.0	4.6	4.9	5.1	5.4	5.6‡	7.3	6.3	5.3	5.1	4.8	
Gender:																
Male	2.5	2.7	3.2	3.0	4.0	4.3	4.7	4.7	5.2	5.8‡	7.9	5.7	4.7	4.7	4.3	
Female	3.8	4.3	3.2	3.6	4.0	4.9	5.2	5.4	5.4	5.5‡	6.8	6.9	5.8	5.5	5.3	
College Plans:																
None or under 4 years	5.0	6.0	5.8	6.0	7.4	9.4	8.6	8.3	8.6	10.0‡	13.5	11.6	9.2	9.6	9.6	
Complete 4 years	2.8	3.1	2.7	2.8	3.4	3.8	4.3	4.5	4.8	4.9‡	6.4	5.5	4.7	4.5	4.2	
Region:																
Northeast	2.7	2.8	3.4	2.8	2.6	3.9	3.6	5.7	5.5	3.9‡	6.2	4.6	4.3	5.0	5.0	
Midwest	2.4	3.0	2.5	2.6	3.2	4.4	3.7	3.4	4.6	5.4‡	5.6	5.5	4.5	3.5	3.3	
South	4.2	4.5	3.9	4.2	5.1	5.7	7.3	6.6	6.0	6.9‡	9.6	8.1	7.5	7.3	6.6	
West	2.9	3.2	3.2	3.6	4.3	3.6	3.7	4.1	4.9	5.4‡	6.7	5.8	4.1	4.1	3.8	
Population Density:																
Large MSA	3.2	3.3	2.7	2.6	3.2	4.2	3.9	4.1	5.0	5.5‡	5.8	5.1	3.9	3.2	3.5	
Other MSA	3.0	3.8	3.3	3.9	4.1	4.6	4.4	5.2	5.5	5.4‡	7.0	6.8	5.8	5.7	5.4	
Non-MSA	3.5	3.3	3.6	3.0	4.7	5.2	7.0	6.0	5.6	6.1‡	9.9	6.7	6.4	6.4	5.3	
Parental Education: ^c																
1.0–2.0 (Low)	3.3	5.3	4.8	4.2	5.0	5.9	4.7	6.5	5.0	8.2‡	7.6	7.1	7.5	5.8	5.2	
2.5–3.0	3.6	3.5	3.1	3.3	4.5	5.2	5.3	6.3	5.9	6.2‡	7.9	7.5	7.3	5.6	5.7	
3.5–4.0	3.2	3.4	3.5	3.4	4.3	5.0	5.5	4.9	5.2	5.6‡	8.1	6.7	5.7	5.7	5.4	
4.5–5.0	2.5	3.9	2.9	2.9	3.5	4.1	4.2	4.5	5.7	5.2‡	6.7	5.8	4.1	4.6	4.0	
5.5–6.0 (High)	3.5	2.3	3.1	3.4	3.2	3.6	4.4	4.0	5.3	4.7‡	6.9	4.6	3.4	4.1	4.0	
Race/Ethnicity (2-year average): ^d																
White	—	4.0	3.8	3.6	4.1	4.9	5.6	6.0	6.1	6.4	7.0‡	7.6 ^e	7.4	6.5	6.0	
African American	—	0.9	0.9	0.9	0.8	0.7	0.9	1.0	1.2	1.1	1.5‡	1.5 ^e	1.3	1.3	1.3	
Hispanic	—	2.9	3.3	3.1	3.1	3.4	3.5	3.5	3.7	4.0	4.0‡	4.3 ^e	4.4	4.5	4.3	

Table continued on next page.

TABLE 84 (cont.)
TRANQUILIZERS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																	2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000		
Total	5.2	5.3	4.6	5.0	5.1	4.5	4.3	3.7	3.9	3.9	4.1	4.1	3.9	3.4	2.6	1.3	-2.2* sss	
Gender:																		
Male	5.0	4.6	4.1	4.4	4.5	4.7	4.0	3.8	3.2	3.2	3.5	3.8	3.9	3.7	§	0.8	-2.8 sss	
Female	5.3	6.0	5.0	5.7	5.5	4.3	4.5	3.6	4.6	4.5	4.5	4.2	3.7	3.1	§	1.5	-1.7 sss	
College Plans:																		
None or under 4 years	10.2	9.9	9.6	11.4	10.6	9.9	10.7	6.4	7.9	8.5	8.7	7.2	7.4	6.0	§	1.5	-4.5 sss	
Complete 4 years	4.5	4.7	3.9	4.4	4.4	3.9	3.6	3.4	3.5	3.4	3.6	3.7	3.4	2.9	§	1.2	-1.8 sss	
Region:																		
Northeast	3.8	4.1	2.8	3.0	4.0	3.5	3.0	2.9	3.0	3.4	2.3	2.4	2.1	1.9	§	0.8	-1.1 ss	
Midwest	4.2	5.3	4.7	5.1	5.1	4.4	4.3	3.9	4.7	2.9	3.9	4.3	3.9	3.3	§	1.2	-2.1 sss	
South	6.9	6.5	6.2	6.2	6.1	5.4	5.3	4.3	4.3	4.8	4.6	3.8	4.6	4.0	§	1.7	-2.3 sss	
West	5.3	4.7	3.7	5.2	4.3	4.2	3.9	3.3	3.2	3.8	4.8	5.5	4.0	3.7	§	1.0	-2.6 sss	
Population Density:																		
Large MSA	4.4	4.1	3.3	3.9	4.6	4.0	3.7	3.0	3.6	3.8	4.5	4.5	3.8	2.8	§	1.3	-1.4 sss	
Other MSA	5.3	5.8	5.2	5.8	5.2	4.7	4.2	4.1	3.9	3.9	3.7	4.1	3.9	3.8	§	1.2	-2.6 sss	
Non-MSA	6.3	6.0	5.1	5.1	5.5	5.1	5.2	3.7	4.5	3.8	4.5	3.4	3.8	3.6	§	1.3	-2.2 sss	
Parental Education: ^c																		
1.0–2.0 (Low)	6.6	5.5	5.4	7.4	6.5	4.8	5.5	5.4	5.4	4.8	4.9	4.6	5.3	4.0	§	2.0	-1.9 s	
2.5–3.0	6.2	6.4	5.9	5.9	6.7	5.0	5.4	4.6	4.0	4.8	5.4	4.8	3.7	4.0	§	1.7	-2.3 ss	
3.5–4.0	5.6	5.7	5.2	5.9	5.8	4.8	5.0	4.4	4.9	5.0	4.2	4.5	4.7	4.4	§	1.4	-2.9 sss	
4.5–5.0	4.3	4.9	3.5	4.1	4.0	4.4	3.6	2.4	3.3	3.4	3.7	3.7	3.2	2.5	§	0.8	-1.7 sss	
5.5–6.0 (High)	4.3	3.9	3.6	3.6	3.3	4.1	3.2	3.2	2.8	2.5	3.1	3.8	3.4	2.8	§	1.2	-1.7 sss	
Race/Ethnicity (2-year average): ^d																		
White	6.0	6.2	6.1	5.7	5.8	5.6	5.1	4.4	4.0	4.1	4.0	4.1	4.0	3.6	§	1.8	-1.8 sss	
African American	1.6	1.3	1.0	1.3	1.8	2.1	1.8	1.7	2.1	2.5	2.7	2.2	1.5	1.8	§	1.1	-0.7	
Hispanic	3.8	3.3	3.5	4.5	4.9	4.1	4.0	4.2	3.8	3.8	4.4	5.2	5.3	4.2	§	2.2	-2.1 ss	

Source. The Monitoring the Future study, the University of Michigan.

Notes. † ‡ * indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bIn 2001, for the tranquilizer list of examples, Miltown was replaced with Xanax. The 2001 data are based on the changed forms only; N is one half of N indicated.

In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^cParental education is an average score of mother's education and father's education. See appendix B for details.

^dEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^eDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 85
TRANQUILIZERS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
Approximate weighted N =	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	10.6	10.3	10.8	9.9	9.6	8.7	8.0	7.0	6.9	6.1	6.1	5.8	5.5	4.8	3.8	3.5
Gender:																
Male	10.0	9.4	10.2	9.7	9.9	9.0	8.0	6.9	7.0	6.3	6.4	5.9	5.2	4.7	4.0	3.5
Female	11.1	11.0	11.4	10.1	9.3	8.5	7.7	7.1	6.7	5.8	5.7	5.8	5.8	4.8	3.5	3.5
College Plans:																
None or under 4 years	—	11.5	12.3	11.1	11.0	10.7	9.4	8.0	8.0	7.4	6.8	7.2	6.7	5.1	4.8	4.3
Complete 4 years	—	8.9	9.0	8.6	8.1	7.2	6.9	6.3	5.8	5.2	5.5	5.1	4.9	4.6	3.3	3.2
Region:																
Northeast	9.2	9.7	10.4	10.9	11.5	8.6	8.3	7.8	6.8	6.8	7.1	6.4	6.9	4.5	3.7	2.9
Midwest	10.6	10.1	11.0	8.8	7.5	8.2	7.8	6.2	6.8	5.6	6.0	5.5	4.5	3.7	3.1	2.9
South	11.3	11.7	11.4	10.5	10.4	9.5	7.8	7.4	7.4	6.9	5.9	6.3	5.7	6.0	4.4	4.3
West	11.7	8.5	9.6	8.9	9.4	8.6	8.0	6.4	6.2	4.9	5.3	4.8	5.2	4.4	3.4	3.9
Population Density:																
Large MSA	11.2	9.6	9.6	10.3	9.9	8.7	8.3	7.0	7.0	5.4	5.8	5.3	5.8	4.7	3.1	3.6
Other MSA	11.0	11.3	11.4	10.1	10.2	9.3	8.1	7.2	7.2	6.1	6.0	5.7	5.6	5.0	3.5	3.7
Non-MSA	9.9	9.5	11.0	9.2	8.7	8.0	7.5	6.8	6.5	6.8	6.5	6.4	5.2	4.5	4.9	3.3
Parental Education: ^b																
1.0–2.0 (Low)	—	11.2	10.1	9.4	9.4	9.1	7.8	7.1	6.1	6.0	6.5	5.3	6.7	5.7	3.9	3.6
2.5–3.0	—	9.8	10.3	11.5	10.1	8.8	9.1	8.0	7.3	7.2	6.5	6.2	5.8	5.4	4.6	3.9
3.5–4.0	—	9.8	11.2	11.1	9.5	10.4	8.9	8.3	6.7	6.9	5.8	6.4	6.5	5.3	4.5	3.4
4.5–5.0	—	11.3	11.7	11.4	10.5	10.0	8.1	7.4	7.6	6.6	5.8	6.3	4.7	5.9	5.5	3.8
5.5–6.0 (High)	—	9.3	12.0	10.1	11.0	11.4	10.3	9.1	7.6	7.1	6.3	5.5	5.4	5.4	5.6	4.9
Race/Ethnicity (2-year average): ^c																
White	—	—	11.4	11.1	10.5	9.9	9.1	8.3	7.8	7.3	6.8	6.6	6.3	5.9	5.0	4.2
African American	—	—	4.3	4.2	3.6	3.1	3.0	2.5	2.3	2.1	1.7	1.7	2.0	2.0	1.2	0.7
Hispanic	—	—	8.4	8.2	7.4	6.4	5.7	5.8	5.1	5.3	5.0	4.4	3.7	2.5	1.6	1.9

Table continued on next page.

TABLE 85 (cont.)
TRANQUILIZERS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
Approximate weighted N =	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001 ^d	2002	2003	2004	2005	2006
	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	3.6	2.8	3.5	3.7	4.4	4.6	4.7	5.5	5.8	5.7 [‡]	6.9	7.7	6.7	7.3	6.8	6.6
Gender:																
Male	3.5	2.7	3.5	4.0	4.7	5.0	5.4	6.3	6.9	6.4 [‡]	7.9	8.4	6.9	8.0	7.2	7.1
Female	3.6	3.0	3.3	3.5	4.1	4.0	3.9	4.7	4.8	4.9 [‡]	5.8	6.9	6.3	6.5	6.2	6.1
College Plans:																
None or under 4 years	4.2	3.9	3.9	4.5	5.6	5.6	6.2	6.8	6.4	6.7 [‡]	7.6	9.9	9.6	9.1	10.3	8.5
Complete 4 years	3.4	2.5	3.3	3.5	4.1	4.2	4.0	5.1	5.6	5.2 [‡]	6.7	7.1	6.0	6.8	5.9	6.0
Region:																
Northeast	3.0	3.0	3.7	3.5	3.9	4.8	5.3	4.9	5.6	5.2 [‡]	5.7	6.8	5.4	6.5	5.7	7.2
Midwest	3.0	2.3	2.8	3.1	4.0	4.4	3.5	3.7	5.1	4.7 [‡]	8.1	6.5	5.5	6.2	7.0	5.8
South	4.0	3.5	4.2	4.8	5.0	5.3	5.4	7.5	7.6	6.7 [‡]	7.4	10.4	8.5	9.0	8.3	7.6
West	4.4	2.3	3.0	2.8	4.3	3.0	4.3	4.4	3.9	5.6 [‡]	5.5	5.8	6.6	7.0	4.9	5.3
Population Density:																
Large MSA	2.5	2.9	2.9	3.9	4.0	3.7	4.2	4.8	4.0	4.7 [‡]	7.9	7.8	4.7	5.5	5.5	6.5
Other MSA	4.1	2.7	3.6	3.7	4.5	4.9	4.8	5.7	6.6	6.2 [‡]	6.4	8.2	7.8	8.5	7.4	6.6
Non-MSA	3.7	3.1	3.7	3.5	4.8	4.7	5.1	5.9	6.5	6.0 [‡]	6.7	6.8	7.4	7.3	7.1	6.9
Parental Education: ^b																
1.0–2.0 (Low)	4.0	3.9	3.3	4.2	3.9	4.2	3.9	6.4	5.2	6.4 [‡]	5.4	5.2	7.1	5.0	4.3	5.7
2.5–3.0	3.6	2.8	3.3	3.5	4.7	4.0	4.3	5.2	6.2	5.3 [‡]	6.7	7.9	6.5	7.9	7.7	6.9
3.5–4.0	3.1	2.7	3.5	3.6	4.3	4.7	4.6	6.0	6.4	5.6 [‡]	7.7	8.4	7.6	9.0	7.3	6.9
4.5–5.0	3.9	3.0	3.4	3.7	4.5	4.6	5.4	4.9	5.4	5.9 [‡]	7.4	7.9	5.9	7.1	6.8	6.6
5.5–6.0 (High)	4.0	2.2	4.2	4.2	4.1	5.3	5.1	5.5	5.6	5.8 [‡]	6.0	7.9	6.2	5.4	5.9	6.4
Race/Ethnicity (2-year average): ^c																
White	4.1	3.7	3.7	4.2	4.6	5.1	5.5	6.2	6.8	6.9	7.7 [‡]	9.2 ^f	8.7	8.4	8.6	8.2
African American	0.9	1.3	1.0	1.1	1.2	0.9	0.8	1.0	1.0	0.7	0.6 [‡]	1.1 ^f	1.3	1.7	1.9	1.6
Hispanic	2.7	2.4	2.0	2.4	3.5	4.3	3.8	3.3	3.5	4.3	3.9 [‡]	4.1 ^f	4.5	4.5	4.3	4.3

Table continued on next page.

TABLE 85 (cont.)
TRANQUILIZERS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change	
	<u>2007^e</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>		
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	6.2	6.2	6.3	5.6	5.6	5.3	4.6	4.7	4.7	4.9	4.7	3.9	3.4	3.2	1.2	-2.1* sss	
Gender:																	
Male	6.9	6.8	6.8	5.9	5.4	5.1	4.6	4.9	4.6	4.8	5.2	4.2	3.2	§	1.0	-2.2 sss	
Female	5.4	5.7	5.7	5.2	5.6	5.5	4.4	4.4	4.6	4.9	4.0	3.1	3.4	§	1.4	-2.0 sss	
College Plans:																	
None or under 4 years	9.1	9.3	8.7	8.6	9.1	6.7	6.0	5.5	5.5	5.2	6.2	5.4	4.7	§	2.2	-2.4 ss	
Complete 4 years	5.5	5.5	5.7	4.9	4.9	4.9	4.2	4.4	4.5	4.6	4.2	3.2	2.9	§	1.0	-2.0 sss	
Region:																	
Northeast	5.9	5.3	6.7	5.2	4.8	4.6	4.2	3.3	4.1	3.7	3.6	3.4	2.2	§	1.1	-1.2 s	
Midwest	6.4	5.2	7.0	5.7	6.1	5.3	4.7	4.4	4.5	4.9	4.4	4.3	3.8	§	0.5	-3.3 sss	
South	7.0	7.8	6.9	6.4	5.5	5.3	5.0	5.0	4.6	5.1	5.4	4.0	3.5	§	1.5	-2.0 sss	
West	4.7	5.5	4.1	4.3	5.7	6.0	4.4	5.8	5.6	5.3	4.6	3.5	3.6	§	1.6	-2.0 s	
Population Density:																	
Large MSA	5.9	5.2	5.5	4.9	4.7	5.3	4.5	3.7	4.5	4.8	3.9	3.1	2.9	§	1.4	-1.5 sss	
Other MSA	6.4	6.6	6.7	5.8	5.8	5.6	5.2	5.6	5.3	5.1	5.5	4.2	3.7	§	1.3	-2.4 sss	
Non-MSA	6.2	6.5	6.3	6.0	6.5	4.9	3.6	4.0	3.6	4.6	4.2	4.5	3.2	§	0.7	-2.5 sss	
Parental Education: ^b																	
1.0–2.0 (Low)	6.0	5.9	5.1	5.0	5.4	5.3	4.3	4.4	4.6	4.9	4.3	3.2	3.1	§	1.8	-1.3	
2.5–3.0	6.5	5.5	7.0	6.4	5.5	5.9	4.5	5.1	4.9	4.2	4.9	4.4	4.1	§	0.7	-3.4 sss	
3.5–4.0	6.4	7.4	6.2	6.1	6.7	5.7	5.8	4.7	5.5	5.3	5.5	4.2	3.8	§	1.7	-2.1 s	
4.5–5.0	6.3	5.7	6.5	5.1	5.2	4.9	4.0	4.8	4.5	5.0	4.0	3.3	2.9	§	1.2	-1.7 ss	
5.5–6.0 (High)	5.5	6.0	5.9	4.2	4.6	4.6	4.0	4.4	4.3	5.0	4.9	3.6	2.5	§	1.0	-1.5 ss	
Race/Ethnicity (2-year average): ^c																	
White	7.9	7.9	8.0	7.3	6.8	6.7	5.9	5.1	5.0	5.1	5.1	4.7	4.0	§	1.8	-2.2 sss	
African American	1.3	1.0	1.2	2.2	2.3	1.6	1.7	2.5	2.8	2.9	2.9	2.7	2.3	§	2.5	+0.2	
Hispanic	4.5	4.2	4.0	3.9	3.9	3.9	3.8	4.2	4.6	4.7	4.7	3.7	3.0	§	2.1	-0.8	

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dIn 2001, for the tranquilizer list of examples, Miltown was replaced with Xanax. The 2001 data are based on the changed forms only; N is one half of N indicated.

In 2002 the remaining forms were changed. Beginning in 2002, the data are based on all forms. Data for any illicit drug other than marijuana and hallucinogens are also affected by these changes and have been treated in a parallel manner.

^eIn 2007, for the list of tranquilizers, Equanil and meprobamate were replaced with Soma while Atarax, Tranxene, and Vistaril were replaced with Ativan and Klonopin.

^fDue to changes in the question wording in half of the questionnaire forms in 2001, the 2002 entry comprises half of the 2001 sample data double-weighted (the half with the new version of the question) and all of the 2002 sample data.

TABLE 86
ANY PRESCRIPTION DRUG^a WITHOUT MEDICAL SUPERVISION:
Trends in Annual Prevalence of Use by Subgroups in Grade 12

	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
Approx. weighted N =	14,700	14,200	14,500	14,000	13,700	13,700	13,700	13,600	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	17.1	16.8	15.8	15.4	14.4	15.0	15.2	14.8‡	15.9	13.9	12.9	12.0	10.9	9.9	8.6	7.6	4.4	-4.2* sss
Gender																		
Male	18.2	17.0	16.4	16.3	16.1	16.1	15.9	15.5‡	16.5	14.2	12.9	12.3	12.0	10.7	9.0	§	4.4	-4.6 sss
Female	15.7	16.3	14.8	14.4	13.5	13.7	14.0	13.7‡	14.6	13.0	12.6	11.0	9.6	8.1	7.8	§	4.1	-3.7 sss
College Plans																		
None or under 4 years	22.2	22.3	21.7	20.9	18.6	20.0	22.3	16.7‡	21.1	16.4	14.8	12.3	12.9	12.0	10.7	§	5.2	-5.5 sss
Complete 4 years	15.8	15.2	14.3	14.1	13.9	13.8	13.7	14.0‡	14.7	13.0	12.3	11.5	10.3	8.8	7.7	§	4.0	-3.6 sss
Region																		
Northeast	15.5	17.4	15.8	15.2	14.7	15.1	13.3	12.4‡	14.6	11.7	11.2	9.7	9.2	9.1	5.7	§	4.3	-1.4
Midwest	18.9	16.1	16.5	15.0	17.3	15.0	15.9	15.6‡	17.0	14.3	13.1	13.5	10.1	9.9	9.6	§	3.6	-5.9 sss
South	17.9	17.4	16.3	16.1	14.3	15.6	14.3	13.5‡	16.1	15.1	13.0	12.0	11.7	10.0	8.6	§	4.2	-4.4 sss
West	15.2	16.2	14.1	15.0	12.8	14.0	17.2	17.0‡	15.5	13.3	13.9	11.9	11.7	10.2	10.3	§	5.5	-4.8 ss
Population Density																		
Large MSA	15.5	16.9	15.0	13.3	13.6	14.9	14.1	15.8‡	15.9	12.3	12.2	12.0	9.6	8.5	8.0	§	4.6	-3.4 sss
Other MSA	18.0	16.5	15.7	15.7	15.3	15.5	15.9	14.7‡	16.8	15.9	13.5	11.9	11.8	10.5	9.3	§	4.9	-4.4 sss
Non-MSA	17.4	17.5	17.1	17.3	15.4	14.1	15.0	13.2‡	13.5	11.2	12.7	12.0	11.3	10.9	8.0	§	3.0	-5.0 sss
Parental Education ^b																		
1.0–2.0 (Low)	11.8	15.7	13.7	13.2	11.5	11.9	13.2	13.3‡	13.6	12.1	12.5	10.3	8.6	8.6	7.5	§	3.6	-3.9 ss
2.5–3.0	18.1	17.2	16.4	15.3	15.6	15.6	15.1	16.1‡	16.9	14.7	12.3	10.6	11.7	9.8	8.5	§	3.3	-5.3 sss
3.5–4.0	18.9	18.2	16.2	16.3	15.5	16.3	16.1	15.5‡	16.3	13.7	14.0	13.1	11.3	10.0	9.7	§	5.1	-4.6 sss
4.5–5.0	17.4	16.6	15.9	15.2	15.7	14.6	14.6	14.6‡	16.6	13.9	12.7	12.4	10.7	10.1	8.2	§	4.1	-4.1 sss
5.5–6.0 (High)	15.0	15.5	14.8	15.1	13.5	14.1	15.5	13.0‡	15.2	14.8	14.0	12.7	12.0	10.1	8.1	§	5.9	-2.1
Race/Ethnicity (2-year average) ^c																		
White	—	19.8	19.3	18.9	18.2	17.8	17.8	17.5	17.2‡	15.5	14.4	13.5	12.5	11.4	10.1	§	6.1	-4.0 sss
African American	—	6.1	5.8	5.7	5.7	7.3	7.5	6.5	10.1‡	10.2	9.3	8.7	7.8	6.4	5.6	§	6.1	+0.5
Hispanic	—	12.8	11.9	10.3	10.1	9.9	10.3	10.9	11.9‡	12.4	11.1	10.5	10.1	8.5	7.3	§	4.2	-3.1 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. " — " indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table D-S3 for the number of subgroup cases. See appendix B for definition of variables in table. ‡ indicates that the question changed the following year.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aThe use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers ... without a doctor telling you to use them. In 2013 the question wording for amphetamines was changed in three of the 12th grade questionnaires. 2013 data are based on the three changed forms only. N is three sixths of N indicated. Data for any prescription drug use are affected by these changes and have been handled in a parallel manner.

^bParental education is an average score of mother's education and father's education reported on the following scale: (1) Completed grade school or less, (2) Some high school, (3) Completed high school, (4) Some college, (5) Completed college, (6) Graduate or professional school after college. Missing data were allowed on one of the two variables.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 87
OVER-THE-COUNTER COUGH/COLD MEDICINES: ^c
Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																	2019– 2021 change
	1991– 2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	4.2	4.0	3.6	3.8	3.2	2.7	3.0	2.9	2.0	1.6	2.6	2.1	2.8	3.2	4.6	3.5	+0.3*
Gender:																		
Male	—	3.4	3.1	2.7	3.7	2.7	2.2	1.7	2.7	1.4	1.1	2.4	2.1	2.6	3.4	§	3.0	-0.4
Female	—	4.7	5.0	4.3	3.8	3.7	3.1	3.9	3.0	2.5	2.1	2.9	2.1	2.9	3.1	§	4.2	+1.1
College Plans:																		
None or under 4 years	—	11.9	8.0	7.1	12.9	5.9	5.9	6.4	8.2	5.2	4.5	7.0	6.0	4.2	5.2	§	4.2	-1.0
Complete 4 years	—	3.4	3.6	3.3	3.1	2.9	2.5	2.8	2.6	1.8	1.4	2.3	1.7	2.6	3.0	§	3.3	+0.3
Region:																		
Northeast	—	3.4	2.4	2.5	2.7	2.2	2.3	1.6	2.6	1.8	1.2	1.8	0.9	1.4	2.1	§	0.7	-1.4
Midwest	—	4.5	4.0	3.8	3.8	3.0	2.4	2.6	1.9	1.6	1.6	3.2	2.4	2.4	2.6	§	4.3	+1.7
South	—	4.2	4.8	4.1	4.5	3.2	2.7	3.9	3.1	2.2	2.0	2.5	2.1	3.2	3.6	§	4.4	+0.9
West	—	4.4	3.6	3.3	3.3	4.1	3.5	3.0	3.8	2.3	1.2	2.9	2.6	3.7	4.1	§	3.3	-0.8
Population Density:																		
Large MSA	—	3.5	2.7	3.1	3.4	2.6	1.7	3.1	3.6	1.6	1.3	2.3	1.3	2.3	3.3	§	2.7	-0.6
Other MSA	—	4.4	4.8	3.5	3.8	3.7	3.5	3.0	2.7	2.2	1.7	2.8	2.7	2.9	3.3	§	4.0	+0.7
Non-MSA	—	4.6	4.2	4.3	4.3	3.0	2.7	2.8	2.3	2.2	1.8	2.8	1.8	3.5	2.9	§	3.8	+0.9
Parental Education: ^a																		
1.0–2.0 (Low)	—	6.2	5.7	4.8	9.2	4.1	5.0	4.6	3.1	2.8	1.5	4.1	3.4	4.3	5.8	§	0.8	-5.0 ss
2.5–3.0	—	5.1	4.5	4.4	4.6	4.0	3.7	3.3	3.9	1.0	2.7	2.9	1.9	2.8	5.0	§	4.0	-1.0
3.5–4.0	—	4.4	5.4	3.8	4.2	3.1	3.7	4.0	3.2	2.7	1.3	3.5	2.6	3.0	3.4	§	5.1	+1.7
4.5–5.0	—	3.8	3.4	3.4	3.3	2.1	1.9	1.9	3.2	1.7	1.6	1.8	1.9	3.5	2.4	§	3.5	+1.1
5.5–6.0 (High)	—	3.0	2.0	2.5	1.7	3.3	1.0	2.0	1.5	2.5	1.1	2.4	1.6	1.4	2.7	§	3.1	+0.5
Race/Ethnicity (2-year average): ^b																		
White	—	—	4.5	4.1	3.7	3.3	2.9	2.6	2.6	2.6	1.9	2.1	2.4	2.4	2.9	§	5.0	+2.1 s
African American	—	—	2.2	2.3	2.7	2.8	2.3	3.0	3.2	2.1	1.9	2.4	2.3	2.4	3.3	§	3.5	+0.2
Hispanic	—	—	4.4	3.8	3.6	3.9	3.2	3.8	3.9	2.8	1.5	1.9	2.3	2.4	2.8	§	3.4	+0.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cTaken to get high.

TABLE 88
OVER-THE-COUNTER COUGH/COLD MEDICINES: ^c
Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																	2019– 2021 change
	1991– 2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	5.3	5.4	5.3	6.0	5.1	5.5	4.7	4.3	3.7	3.3	3.0	3.6	3.3	2.6	3.3	2.7	+0.1*
Gender:																		
Male	—	5.1	4.6	4.9	5.9	5.5	5.4	5.5	4.6	3.8	3.0	2.7	3.5	3.7	3.1	§	2.8	-0.3
Female	—	5.5	6.3	5.7	6.0	4.9	5.8	4.1	3.5	3.3	3.5	3.3	3.5	3.0	2.1	§	2.4	+0.3
College Plans:																		
None or under 4 years	—	9.9	9.7	11.3	13.5	10.8	13.6	10.2	8.0	9.2	9.3	4.8	7.6	4.8	4.6	§	4.3	-0.3
Complete 4 years	—	4.6	4.8	4.6	5.2	4.5	4.6	4.2	3.9	3.0	2.6	2.8	3.2	3.1	2.3	§	2.3	+0.1
Region:																		
Northeast	—	4.1	5.8	3.5	4.1	4.4	5.3	4.5	4.4	3.1	2.2	3.4	2.1	2.6	3.2	§	3.2	-0.1
Midwest	—	4.3	5.4	5.4	6.2	4.9	6.0	4.3	3.5	3.3	3.2	2.0	3.7	3.3	2.1	§	2.1	0.0
South	—	6.3	5.9	6.6	6.9	5.0	6.2	5.7	4.2	4.4	3.4	3.7	3.5	3.4	2.1	§	3.3	+1.3
West	—	6.0	4.7	4.8	6.3	6.3	4.2	4.0	5.1	3.5	4.2	2.6	4.8	3.9	3.7	§	1.9	-1.8
Population Density:																		
Large MSA	—	4.3	4.7	4.3	5.3	5.1	4.1	4.2	4.3	3.8	2.6	2.9	3.6	3.7	1.6	§	1.8	+0.3
Other MSA	—	5.5	5.6	6.2	6.8	4.7	6.4	4.7	4.0	3.3	3.6	3.1	3.2	3.2	2.7	§	2.8	+0.1
Non-MSA	—	6.3	6.3	4.7	5.3	6.1	5.9	5.5	5.0	4.3	3.7	2.7	4.7	3.1	4.0	§	3.9	-0.1
Parental Education: ^a																		
1.0–2.0 (Low)	—	5.1	6.4	6.8	9.7	9.3	8.0	6.2	5.4	3.5	5.1	4.4	5.4	6.7	2.8	§	3.1	+0.3
2.5–3.0	—	6.2	7.4	5.9	7.3	6.0	6.2	5.9	5.3	2.9	3.9	3.6	4.2	4.2	3.3	§	2.8	-0.5
3.5–4.0	—	6.2	6.0	5.0	6.5	5.3	5.5	5.8	5.2	4.7	3.4	2.4	4.8	2.4	3.3	§	4.0	+0.7
4.5–5.0	—	4.1	4.5	5.6	4.5	3.4	5.1	3.5	3.0	2.9	2.9	2.6	3.1	2.4	2.2	§	2.0	-0.1
5.5–6.0 (High)	—	5.1	3.0	3.9	5.2	5.1	4.6	3.6	2.9	3.6	2.6	3.3	2.0	3.0	2.0	§	2.2	+0.2
Race/Ethnicity (2-year average): ^b																		
White	—	—	6.0	6.2	6.4	5.8	5.4	5.1	4.7	4.2	3.7	3.5	3.5	3.4	3.2	§	3.5	+0.3
African American	—	—	3.9	3.1	2.1	2.6	3.8	2.7	2.3	3.6	3.4	2.4	2.8	2.6	2.5	§	1.7	-0.8
Hispanic	—	—	3.2	4.0	5.5	5.6	5.7	6.5	5.0	3.5	2.4	2.1	3.4	4.2	3.0	§	3.7	+0.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cTaken to get high.

TABLE 89
OVER-THE-COUNTER COUGH/COLD MEDICINES: ^c
Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Approximate weighted N =	Percentage who used in last 12 months																2019– 2021 change																				
		1991– 2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021																			
Total	—	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	—	6.9	5.8	5.5	5.9	6.6	5.3	5.6	5.0	4.1	4.6	4.0	3.2	3.4	2.5	3.2	1.7	-0.8*			
Gender:																																						
Male	—	7.6	5.9	6.3	8.1	7.7	5.6	6.1	5.9	4.8	5.8	4.7	4.0	4.0	3.4	§	1.7	-1.7 s																				
Female	—	5.9	5.1	4.7	4.1	5.4	4.7	4.5	4.2	3.3	3.2	3.3	2.5	2.6	1.5	§	1.3	-0.3																				
College Plans:																																						
None or under 4 years	—	8.9	8.3	7.6	7.1	8.6	7.3	8.4	7.3	4.3	5.6	5.7	4.7	5.9	4.3	§	3.3	-1.0																				
Complete 4 years	—	6.2	5.2	4.9	5.5	6.1	4.9	4.8	4.4	3.9	4.2	3.6	2.9	2.8	2.0	§	1.2	-0.7																				
Region:																																						
Northeast	—	5.9	7.8	4.4	5.1	7.4	5.6	4.9	5.1	3.8	3.8	3.9	2.7	3.4	1.4	§	1.0	-0.4																				
Midwest	—	6.6	6.8	4.4	6.7	5.9	6.1	5.8	4.6	4.3	4.2	3.7	2.9	3.7	2.1	§	0.4	-1.7 s																				
South	—	7.6	5.2	6.7	6.2	6.6	5.5	5.6	5.7	4.6	4.6	4.7	3.3	3.1	2.3	§	2.8	+0.5																				
West	—	6.7	4.2	5.8	4.8	6.6	4.1	5.8	4.5	3.5	5.4	3.2	3.7	3.6	4.3	§	1.8	-2.5																				
Population Density:																																						
Large MSA	—	7.6	5.4	4.9	5.7	6.3	5.2	5.9	5.0	2.8	4.7	3.4	2.7	2.6	1.6	§	1.0	-0.6																				
Other MSA	—	6.0	5.9	5.5	6.6	6.8	5.4	5.2	4.9	4.8	4.4	4.6	3.9	3.4	2.7	§	2.7	0.0																				
Non-MSA	—	7.5	6.1	6.4	4.4	6.2	5.4	6.0	5.4	4.3	4.7	3.6	2.8	4.7	3.4	§	0.6	-2.8 sss																				
Parental Education: ^a																																						
1.0–2.0 (Low)	—	7.7	7.2	6.8	7.8	7.4	7.3	5.7	5.9	3.6	3.3	4.6	3.0	3.6	5.6	§	1.9	-3.7																				
2.5–3.0	—	6.2	6.7	5.0	6.1	6.2	5.7	5.5	4.3	5.6	4.8	4.1	5.1	4.0	2.8	§	1.1	-1.7 s																				
3.5–4.0	—	7.7	4.6	6.3	5.7	7.9	4.9	5.6	6.3	2.8	5.9	3.9	3.7	3.3	2.0	§	2.2	+0.2																				
4.5–5.0	—	6.7	6.8	4.8	5.6	5.3	4.9	6.0	4.1	4.5	4.0	3.4	2.0	2.5	2.2	§	1.1	-1.1																				
5.5–6.0 (High)	—	4.9	4.0	4.4	5.1	5.9	5.6	5.3	3.9	4.1	3.2	4.5	0.9	3.8	1.6	§	3.0	+1.4																				
Race/Ethnicity (2-year average): ^b																																						
White	—	—	6.3	5.8	5.9	6.4	6.2	5.6	5.3	4.5	4.2	4.2	3.4	3.2	2.8	§	2.1	-0.7																				
African American	—	—	5.0	4.2	4.5	4.6	4.6	3.8	3.7	4.5	4.8	4.9	3.3	2.2	2.1	§	2.9	+0.8																				
Hispanic	—	—	7.9	6.0	6.0	7.1	5.4	5.4	5.8	3.8	4.2	5.0	4.2	3.3	3.1	§	3.6	+0.5																				

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cTaken to get high.

TABLE 90
ROHYPNOL: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months											
	1991– 1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	—	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500
Total	—	1.0	0.8	0.8	0.5	0.5	0.7	0.3	0.5	0.6	0.7	0.5
Gender:												
Male	—	1.1	0.7	0.8	0.6	0.5	0.6	0.2	0.4	0.5	0.7	0.7
Female	—	1.0	0.9	0.9	0.2	0.6	0.7	0.2	0.4	0.7	0.7	0.4
College Plans:												
None or under 4 years	—	2.5	1.5	3.0	0.9	2.2	2.0	1.0	1.3	2.4	1.8	1.1
Complete 4 years	—	0.8	0.7	0.6	0.4	0.4	0.5	0.2	0.4	0.4	0.6	0.5
Region:												
Northeast	—	0.9	0.3	0.9	0.3	0.6	0.9	0.2	0.2	0.3	0.6	1.4
Midwest	—	0.9	0.8	0.8	0.3	0.4	1.0	0.4	0.2	0.8	0.6	0.0
South	—	1.3	1.1	0.9	0.7	0.9	0.6	0.4	0.6	0.5	0.5	0.6
West	—	1.0	0.7	0.8	0.5	*	0.4	0.1	0.7	0.7	1.1	0.2
Population Density:												
Large MSA	—	1.2	0.8	0.7	0.4	0.9	1.2	0.1	0.1	0.5	0.4	0.6
Other MSA	—	1.1	0.9	1.0	0.6	0.5	0.5	0.3	0.4	0.8	0.7	0.5
Non-MSA	—	0.8	0.7	0.8	0.3	0.2	0.4	0.5	1.0	0.3	1.0	0.5
Parental Education: ^a												
1.0–2.0 (Low)	—	2.0	2.1	2.2	0.2	0.9	1.3	0.9	1.8	2.4	0.6	0.7
2.5–3.0	—	1.1	1.1	0.8	0.3	0.8	0.3	0.0	0.9	0.1	1.3	0.1
3.5–4.0	—	1.2	0.7	0.9	0.6	0.2	1.2	0.2	0.1	0.2	0.4	0.7
4.5–5.0	—	0.8	0.6	0.6	0.5	0.4	0.7	0.6	0.4	0.6	0.9	0.1
5.5–6.0 (High)	—	0.8	0.5	0.9	0.7	0.9	0.2	0.2	0.0	0.1	0.5	1.5
Race/Ethnicity (2-year average): ^b												
White	—	—	0.9	0.8	0.7	0.5	0.6	0.5	0.4	0.4	0.5	0.5
African American	—	—	0.4	0.4	0.4	0.3	0.4	0.4	0.3	0.9	0.7	0.2
Hispanic	—	—	1.4	1.5	1.5	0.6	0.9	0.8	0.4	1.0	0.8	0.6

Table continued on next page.

TABLE 90 (cont.)
ROHYPNOL: Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	0.7	0.5	0.4	0.5	0.8	0.4	0.4	0.3	0.3	0.5	0.4	0.3	0.4	§	0.2	-0.2*
Gender:																
Male	0.6	0.6	0.4	0.3	0.7	0.4	0.5	0.3	0.5	0.4	0.2	0.4	0.2	§	0.0	-0.2
Female	0.7	0.3	0.4	0.7	0.8	0.3	0.3	0.4	0.2	0.3	0.6	0.3	0.7	§	0.0	-0.7
College Plans:																
None or under 4 years	3.3	2.6	2.4	0.5	1.8	3.1	2.2	0.9	1.5	3.3	1.1	1.3	0.2	§	0.4	+0.2
Complete 4 years	0.4	0.3	0.2	0.5	0.7	0.2	0.3	0.3	0.2	0.2	0.3	0.1	0.4	§	0.1	-0.4
Region:																
Northeast	0.1	0.0	0.0	0.5	0.9	0.4	0.5	0.2	0.0	1.2	0.3	0.0	0.0	§	.	.
Midwest	0.8	0.7	0.6	0.4	0.6	0.1	0.0	0.5	0.0	0.2	0.5	0.4	0.5	§	0.2	-0.3
South	0.7	0.6	0.4	0.7	0.4	0.6	0.5	0.2	0.4	0.5	0.3	0.4	0.7	§	0.3	-0.4
West	0.8	0.5	0.5	0.5	1.7	0.4	0.6	0.4	0.6	0.1	0.4	0.3	0.2	§	0.0	-0.2
Population Density:																
Large MSA	0.4	0.6	0.1	0.4	1.1	0.4	0.3	0.2	0.3	1.0	0.6	0.2	0.2	§	0.1	-0.1
Other MSA	0.8	0.5	0.6	0.7	0.7	0.3	0.6	0.3	0.5	0.1	0.4	0.3	0.3	§	0.1	-0.2
Non-MSA	0.8	0.4	0.4	0.3	0.5	0.6	0.2	0.4	0.0	0.3	0.1	0.6	0.9	§	0.3	-0.6
Parental Education: ^a																
1.0–2.0 (Low)	0.5	0.7	1.4	1.4	0.3	1.6	0.3	0.7	0.0	1.9	0.9	0.3	0.0	§	.	.
2.5–3.0	0.5	0.5	0.0	0.0	1.0	0.5	0.7	0.4	0.2	0.5	0.7	0.2	1.9	§	0.0	-1.9
3.5–4.0	0.7	0.7	0.3	0.6	1.0	0.4	0.4	0.2	0.8	0.4	0.2	0.6	0.3	§	0.0	-0.3
4.5–5.0	0.9	0.2	0.6	0.6	0.6	0.3	0.7	0.1	0.3	0.3	0.2	0.4	0.2	§	0.0	-0.2
5.5–6.0 (High)	0.4	0.3	0.1	0.4	1.0	0.2	0.0	0.4	0.0	0.2	0.1	0.2	0.0	§	0.2	+0.2
Race/Ethnicity (2-year average): ^b																
White	0.4	0.5	0.6	0.4	0.3	0.4	0.4	0.3	0.2	0.2	0.2	0.3	0.4	§	0.5	+0.1
African American	0.2	0.2	0.2	0.1	0.3	0.7	0.6	0.6	0.6	0.3	0.2	0.4	0.3	§	0.0	-0.3
Hispanic	1.1	1.0	0.5	0.8	1.6	1.2	0.3	0.4	0.3	0.6	0.7	0.4	0.5	§	0.2	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. ' * ' indicates less than 0.05% but greater than 0%. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1996; N is one half of N indicated in Table S1. Data based on three of four forms in 1997–1998; N is two thirds of N indicated in Table S1. Data based on two of four forms in 1999–2001; N is one third of N indicated in Table S1. Data are based on one of four forms beginning in 2002; N is one sixth of N indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 91
ROHYPNOL: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months											
	1991– 1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	—	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	16,200
Total	—	1.1	1.3	1.2	1.0	0.8	1.0	0.7	0.6	0.7	0.5	0.5
Gender:												
Male	—	1.1	1.4	1.4	1.2	1.0	1.1	0.4	0.7	0.8	0.3	0.4
Female	—	1.1	1.1	1.1	0.9	0.6	0.9	0.9	0.6	0.6	0.6	0.7
College Plans:												
None or under 4 years	—	2.9	2.7	2.9	2.8	1.4	2.8	0.5	0.3	1.4	1.6	1.5
Complete 4 years	—	0.8	1.1	0.9	0.8	0.6	0.7	0.7	0.6	0.6	0.3	0.4
Region:												
Northeast	—	0.6	0.7	0.9	0.4	0.6	0.9	0.4	1.0	0.8	0.7	0.3
Midwest	—	0.5	0.8	1.0	0.9	0.4	0.9	0.3	0.2	0.3	0.2	1.0
South	—	1.9	2.4	2.0	1.8	1.2	1.0	0.4	0.8	0.9	0.7	0.7
West	—	1.1	0.6	0.5	0.6	0.7	1.3	2.0	0.4	0.9	0.2	0.0
Population Density:												
Large MSA	—	1.1	1.3	1.3	0.9	0.9	1.2	0.7	0.3	0.6	0.3	0.2
Other MSA	—	1.4	1.3	1.3	1.3	0.9	0.8	0.9	0.4	0.7	0.5	0.7
Non-MSA	—	0.6	1.3	1.0	0.7	0.5	1.2	0.3	1.5	0.8	0.5	0.8
Parental Education: ^a												
1.0–2.0 (Low)	—	1.7	1.3	1.7	1.2	1.3	2.3	0.4	1.3	0.4	1.1	1.4
2.5–3.0	—	1.0	1.1	1.6	1.6	0.5	1.1	0.9	0.7	0.4	0.5	0.3
3.5–4.0	—	1.5	1.6	1.0	0.8	0.8	0.7	1.0	0.9	1.0	0.4	0.7
4.5–5.0	—	0.7	1.5	0.9	0.6	0.7	1.1	0.7	0.2	0.6	0.6	0.3
5.5–6.0 (High)	—	1.1	0.8	1.3	1.1	0.8	0.3	0.2	0.4	1.0	0.0	0.0
Race/Ethnicity (2-year average): ^b												
White	—	—	1.4	1.5	1.5	1.0	1.0	1.0	0.6	0.9	0.8	0.5
African American	—	—	0.1	0.1	0.1	0.1	0.2	0.3	0.1	0.2	0.4	0.3
Hispanic	—	—	1.0	1.2	1.1	0.7	0.9	1.0	0.7	0.3	0.2	0.6

Table continued on next page.

TABLE 91 (cont.)
ROHYPNOL: Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	0.7	0.4	0.4	0.6	0.6	0.5	0.6	0.5	0.2	0.5	0.3	0.3	0.6	§	0.2	-0.3*
Gender:																
Male	0.7	0.3	0.3	0.6	0.8	0.4	0.7	0.7	0.1	0.6	0.0	0.1	0.3	§	0.2	-0.1
Female	0.7	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.3	0.2	0.5	0.5	0.9	§	0.3	-0.5
College Plans:																
None or under 4 years	1.4	0.5	0.9	1.2	0.9	2.3	0.6	1.8	0.7	2.4	0.4	0.9	0.0	§	1.2	+1.2
Complete 4 years	0.6	0.4	0.4	0.5	0.6	0.3	0.6	0.4	0.2	0.3	0.2	0.2	0.7	§	0.0	-0.6 s
Region:																
Northeast	0.5	0.7	0.2	0.9	0.6	0.8	0.3	0.1	0.1	0.3	0.5	0.1	0.1	§	0.0	-0.1
Midwest	1.4	0.2	0.9	0.3	0.5	0.3	1.1	0.5	0.1	0.9	0.3	0.3	0.1	§	0.2	+0.1
South	0.5	0.4	0.3	0.8	0.8	0.5	0.4	0.8	0.4	0.1	0.3	0.3	1.1	§	0.4	-0.7
West	0.4	0.4	0.3	0.2	0.6	0.5	0.4	0.7	0.1	1.0	0.1	0.5	0.6	§	0.2	-0.4
Population Density:																
Large MSA	0.7	0.1	0.3	0.5	0.7	0.5	0.8	0.3	0.4	0.5	0.4	0.2	1.2	§	0.4	-0.8
Other MSA	0.6	0.7	0.5	0.7	0.5	0.4	0.4	0.4	0.1	0.4	0.3	0.3	0.3	§	0.2	-0.2
Non-MSA	0.9	0.3	0.4	0.4	0.9	0.5	0.7	1.3	0.0	0.8	0.0	0.6	0.1	§	0.1	0.0
Parental Education: ^a																
1.0–2.0 (Low)	0.4	1.2	0.2	0.7	0.0	0.8	0.6	0.6	0.9	1.8	0.9	0.4	0.7	§	0.7	-0.1
2.5–3.0	0.6	0.3	0.4	0.3	0.6	0.8	0.5	1.3	0.0	0.7	0.1	0.0	0.8	§	0.0	-0.8
3.5–4.0	0.5	0.5	0.9	0.4	0.5	0.0	0.8	0.8	0.0	0.3	0.2	0.4	0.3	§	0.4	+0.1
4.5–5.0	0.6	0.2	0.1	0.6	1.2	0.2	0.6	0.1	0.2	0.2	0.3	0.4	0.0	§	.	.
5.5–6.0 (High)	1.1	0.6	0.6	1.0	0.0	1.1	0.4	0.0	0.5	0.5	0.1	0.2	1.8	§	0.2	-1.5
Race/Ethnicity (2-year average): ^b																
White	0.6	0.5	0.5	0.6	0.8	0.6	0.5	0.7	0.5	0.4	0.3	0.4	0.6	§	0.3	-0.3
African American	0.0	0.0	0.1	0.3	0.5	0.3	0.1	0.5	0.3	0.3	0.3	0.1	0.1	§	0.0	-0.1
Hispanic	0.8	0.6	0.3	0.4	0.3	0.3	0.5	0.7	0.3	0.4	0.6	0.4	0.3	§	0.3	0.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1996; N is one half of N indicated in Table S2. Data based on three of four forms in 1997–1998; N is two thirds of N indicated in Table S2. Data based on two of four forms in 1999–2001; N is one third of N indicated in Table S2. Data based on one of four forms beginning in 2002; N is one sixth of N indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 92
ROHYPNOL: Trends in Annual Prevalence of Use by Subgroups in Grade 12

		Percentage who used in last 12 months											
		1975–											
		1995	1996	1997	1998	1999	2000	2001 ^a	2002 ^a	2003	2004	2005	2006
Approximate weighted N =		—	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total		—	1.1	1.2	1.4	1.0	0.8	0.9‡	1.6	1.3	1.6	1.2	1.1
Gender:													
Male		—	1.8	1.2	1.7	1.0	0.8	1.1‡	2.3	2.0	2.3	1.5	1.3
Female		—	0.3	1.1	1.1	1.0	0.7	0.6‡	1.0	0.5	0.7	0.8	0.5
College Plans:													
None or under 4 years		—	1.4	0.5	1.8	1.0	0.1	1.2‡	2.5	2.1	3.0	2.4	1.6
Complete 4 years		—	0.8	1.3	1.3	1.0	0.9	0.8‡	1.4	0.9	1.0	0.9	0.8
Region:													
Northeast		—	0.9	0.6	1.2	0.8	0.3	1.7‡	1.2	1.4	0.8	1.1	1.3
Midwest		—	1.0	0.5	1.0	0.8	0.9	0.5‡	1.8	1.2	1.3	1.7	0.8
South		—	1.8	2.2	2.6	1.6	1.3	1.0‡	1.5	1.4	2.1	1.3	1.3
West		—	0.1	0.8	0.0	0.2	0.1	0.8‡	2.0	1.1	1.8	0.6	1.3
Population Density:													
Large MSA		—	1.3	0.9	0.8	0.3	0.3	1.0‡	1.6	1.1	1.8	1.4	1.2
Other MSA		—	1.3	1.5	2.2	1.4	1.1	1.2‡	2.0	1.7	1.6	1.2	1.0
Non-MSA		—	0.7	0.7	0.6	0.9	0.7	0.3‡	1.1	0.9	1.2	1.1	1.4
Parental Education: ^b													
1.0–2.0 (Low)		—	1.6	0.4	0.6	4.7	0.0	4.2‡	3.7	2.8	3.4	3.9	1.2
2.5–3.0		—	0.5	0.6	1.0	0.3	0.4	0.4‡	2.9	1.8	1.8	1.2	1.2
3.5–4.0		—	0.8	1.4	1.1	1.1	1.5	0.8‡	0.8	1.2	1.5	1.4	1.0
4.5–5.0		—	0.9	1.7	2.7	0.3	0.6	0.8‡	1.6	0.9	0.6	0.3	1.2
5.5–6.0 (High)		—	1.8	1.7	1.4	0.7	0.0	0.2‡	0.8	0.2	1.5	0.9	0.7
Race/Ethnicity (2-year average): ^c													
White		—	—	1.2	1.6	1.5	0.9	0.8‡	—	1.5	1.3	1.1	0.9
African American		—	—	0.1	0.0	0.2	0.4	0.2‡	—	0.8	1.2	1.7	1.3
Hispanic		—	—	2.0	1.3	1.0	0.6	1.3‡	—	1.6	1.7	1.9	2.1

Table continued on next page.

TABLE 92 (cont.)
ROHYPNOL: Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	1.0	1.3	1.0	1.5	1.3	1.5	0.9	0.7	1.0	1.1	0.8	0.7	0.5	§	0.4	-0.1*
Gender:																
Male	0.9	1.5	1.1	2.2	2.0	1.2	0.9	0.9	0.9	1.0	0.7	0.5	0.7	§	0.4	-0.3
Female	0.7	0.8	0.8	0.7	0.5	0.7	0.7	0.4	0.7	1.0	0.8	0.4	0.3	§	0.2	-0.2
College Plans:																
None or under 4 years	1.8	2.3	1.8	3.8	1.7	2.7	1.7	0.6	1.0	3.9	2.5	1.5	1.5	§	0.6	-0.9
Complete 4 years	0.7	1.0	0.8	0.9	1.2	0.8	0.6	0.7	0.8	0.4	0.3	0.3	0.2	§	0.3	+0.1
Region:																
Northeast	1.7	0.8	0.7	2.2	1.9	0.9	0.9	1.3	0.2	0.7	2.0	0.3	0.0	§	0.9	+0.9
Midwest	0.9	1.0	1.3	1.5	0.6	1.1	1.0	0.5	1.6	1.6	0.6	0.6	0.8	§	0.3	-0.5
South	1.0	2.0	1.1	1.0	1.7	2.3	1.0	0.7	1.0	1.3	0.6	1.3	0.3	§	0.4	+0.1
West	0.5	0.9	0.5	1.8	1.0	1.1	0.7	0.3	0.8	0.6	0.4	0.1	1.0	§	0.0	-1.0
Population Density:																
Large MSA	1.0	0.6	1.0	1.5	0.7	1.4	0.5	0.7	1.4	0.6	0.6	0.7	0.6	§	0.3	-0.3
Other MSA	1.1	1.8	1.2	1.5	1.4	1.1	0.9	0.8	0.9	1.7	1.0	0.6	0.4	§	0.6	+0.2
Non-MSA	0.8	1.0	0.4	1.3	1.8	2.6	1.5	0.4	0.6	0.5	0.6	1.0	0.6	§	0.0	-0.6
Parental Education: ^b																
1.0–2.0 (Low)	1.3	1.6	2.1	1.0	3.6	1.4	2.1	3.2	0.7	3.8	1.0	1.0	1.8	§	0.0	-1.8
2.5–3.0	0.8	1.7	1.2	1.2	1.0	1.1	1.2	0.5	0.3	1.6	1.6	1.2	0.6	§	0.5	-0.1
3.5–4.0	0.5	1.3	0.8	1.3	1.1	1.4	0.3	0.3	0.8	0.2	0.4	0.6	0.7	§	0.3	-0.4
4.5–5.0	1.4	0.7	0.4	1.4	1.0	1.4	0.4	0.5	0.6	0.8	0.3	0.3	0.0	§	0.1	+0.1
5.5–6.0 (High)	0.9	0.6	0.9	2.0	1.0	1.1	0.0	0.0	0.3	0.0	0.2	0.3	0.0	§	1.2	+1.2
Race/Ethnicity (2-year average): ^c																
White	0.7	0.9	1.0	1.1	1.3	0.9	0.7	0.4	0.4	0.6	0.8	0.7	0.4	§	0.5	+0.1
African American	1.6	2.1	1.7	1.2	1.3	1.8	2.0	1.9	2.1	2.2	1.4	0.3	0.4	§	1.3	+0.9
Hispanic	1.5	0.8	1.3	1.5	1.0	1.5	1.4	0.8	0.5	0.8	0.7	0.8	1.1	§	3.8	+2.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. '±' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms in 1996–2001; N is one sixth of N indicated in Table S3. Data based on two of six forms beginning in 2002; N is two sixths of N indicated in Table S3. Data based on one of six forms beginning in 2010; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aThe 2001 and 2002 data are not comparable due to changes in the questionnaire forms.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 93
ALCOHOL: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days															
	<u>1991</u>	<u>1992</u>	<u>1993^a</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004^a</u>	<u>2005</u>	
Approximate weighted N =	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	
Total	25.1	26.1‡	24.3	25.5	24.6	26.2	24.5	23.0	24.0	22.4	21.5	19.6	19.7	18.6	17.1	
Gender:																
Male	26.3	26.3‡	25.3	26.5	25.0	26.6	25.2	24.0	24.8	22.5	22.3	19.1	19.4	17.9	16.2	
Female	23.8	25.9‡	23.7	24.7	24.0	25.8	23.9	21.9	23.3	22.0	20.6	20.0	19.8	19.0	17.9	
College Plans:																
None or under 4 years	37.2	39.6‡	41.1	41.4	40.0	41.7	40.2	41.2	41.6	38.3	37.0	35.3	35.3	32.6	32.6	
Complete 4 years	23.1	24.2‡	22.2	23.6	22.6	24.0	22.8	21.0	22.0	20.4	19.7	18.2	18.1	17.2	15.5	
Region:																
Northeast	24.3	23.8‡	21.0	25.4	24.1	26.9	24.8	21.2	25.7	25.6	23.3	19.3	18.4	18.0	14.5	
Midwest	26.6	28.3‡	24.7	24.2	24.7	26.9	22.8	23.9	25.7	24.1	21.2	19.1	21.9	19.3	16.9	
South	25.1	26.8‡	25.4	25.6	25.5	26.3	26.4	23.8	24.4	20.9	22.8	21.6	20.8	18.8	19.0	
West	23.1	23.5‡	25.6	27.2	23.1	24.8	22.7	22.2	19.8	20.2	18.1	17.0	16.0	17.9	16.4	
Population Density:																
Large MSA	25.4	27.4‡	21.2	23.8	22.3	24.9	23.1	21.4	21.7	21.2	19.4	17.4	18.2	18.1	17.2	
Other MSA	24.3	26.1‡	26.0	27.4	25.3	27.4	24.9	22.4	23.4	21.3	21.5	20.1	19.4	19.1	17.9	
Non-MSA	26.2	24.2‡	24.9	23.8	26.0	25.7	25.4	26.0	28.1	26.1	24.1	21.4	22.3	18.1	15.6	
Parental Education: ^b																
1.0–2.0 (Low)	30.7	32.8‡	28.0	33.5	30.8	28.1	29.7	28.9	30.7	30.2	28.5	27.6	25.5	24.2	23.8	
2.5–3.0	27.0	27.2‡	28.0	27.4	27.8	30.1	26.2	26.5	27.9	26.3	27.0	23.2	24.0	22.1	20.4	
3.5–4.0	25.1	26.3‡	25.9	26.7	26.8	27.6	27.8	24.5	25.2	23.0	23.3	21.2	22.0	20.0	18.8	
4.5–5.0	22.8	24.6‡	20.6	22.6	21.0	25.0	22.6	20.2	20.4	18.4	17.2	17.0	16.0	16.3	14.6	
5.5–6.0 (High)	24.0	25.2‡	22.3	23.6	20.5	21.5	20.5	21.3	22.1	19.0	16.4	15.1	15.7	15.8	13.1	
Race/Ethnicity (2-year average): ^c																
White	—	26.6	27.1‡	25.3	25.4	26.6	26.7	24.8	24.7	24.7	23.2	21.5	20.1	19.2	17.9	
African American	—	18.6	19.7‡	19.4	18.7	18.1	17.9	16.1	16.1	16.0	15.0	14.8	15.5	16.2	14.9	
Hispanic	—	31.0	32.3‡	33.5	32.4	29.7	29.8	29.5	29.0	26.7	25.7	26.5	25.3	23.5	20.6	

Table continued on next page.

TABLE 93 (cont.)
ALCOHOL: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days																	2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change	
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700		
Total	17.2	15.9	15.9	14.9	13.8	12.7	11.0	10.2	9.0	9.7	7.3	8.0	8.2	7.9	9.9	7.3	-0.6*	
Gender:																		
Male	16.3	15.6	15.4	14.7	13.2	12.1	10.3	9.3	8.2	9.1	6.7	6.8	7.4	7.3	§	5.9	-1.5	
Female	17.6	16.0	16.4	14.9	14.3	12.8	11.6	11.2	9.5	9.9	7.8	9.1	9.1	8.3	§	8.5	+0.2	
College Plans:																		
None or under 4 years	30.8	33.5	30.4	31.7	25.7	24.6	23.3	23.6	17.0	15.5	14.6	14.5	15.7	13.5	§	9.6	-3.8 s	
Complete 4 years	15.8	14.3	14.7	13.5	12.8	11.7	10.0	9.4	8.3	9.1	6.7	7.3	7.3	7.1	§	6.7	-0.3	
Region:																		
Northeast	16.9	12.9	15.9	15.3	13.3	8.8	9.3	9.1	7.6	8.2	5.9	5.8	6.8	5.3	§	4.8	-0.5	
Midwest	17.0	15.5	16.1	13.0	12.2	11.7	7.8	8.5	8.9	7.9	6.3	6.9	7.6	8.9	§	8.3	-0.7	
South	18.5	17.1	18.2	17.4	15.7	14.2	13.3	12.5	9.8	10.8	8.4	8.1	8.0	9.1	§	7.7	-1.4	
West	15.3	15.9	12.6	12.5	12.5	14.2	11.5	9.1	8.6	10.2	7.3	10.3	9.9	7.3	§	7.7	+0.4	
Population Density:																		
Large MSA	16.6	14.8	15.2	12.9	13.5	11.5	10.3	9.5	8.8	9.2	6.4	7.5	8.7	6.7	§	5.0	-1.7	
Other MSA	17.4	16.1	16.3	15.9	14.0	13.1	11.8	10.4	9.0	9.6	7.0	8.0	7.9	7.8	§	8.2	+0.5	
Non-MSA	17.5	16.8	16.2	15.5	13.5	13.6	10.6	11.1	9.2	10.4	9.3	8.7	7.9	10.5	§	9.2	-1.3	
Parental Education: ^b																		
1.0–2.0 (Low)	24.1	26.1	25.5	22.8	19.2	19.7	17.1	16.2	11.3	13.4	10.3	11.8	10.8	9.6	§	4.7	-4.9 ss	
2.5–3.0	20.0	20.3	19.3	18.3	17.4	16.6	15.3	13.2	11.5	11.6	9.5	10.5	10.9	10.9	§	9.2	-1.8	
3.5–4.0	19.0	16.9	16.8	16.8	15.3	14.1	11.7	12.7	10.7	10.4	6.9	8.5	9.5	9.4	§	10.9	+1.6	
4.5–5.0	13.9	11.6	13.4	11.5	11.4	9.5	8.9	7.5	7.6	7.7	6.3	6.9	6.3	6.9	§	7.0	+0.1	
5.5–6.0 (High)	13.8	11.2	11.5	10.4	9.2	9.3	7.3	7.4	6.5	8.0	6.2	5.5	6.4	5.6	§	5.8	+0.2	
Race/Ethnicity (2-year average): ^c																		
White	16.9	15.6	15.2	15.4	13.9	12.3	10.7	9.5	9.0	8.9	8.2	7.2	7.3	8.0	§	9.2	+1.2	
African American	13.1	12.3	12.9	12.3	11.8	11.6	10.0	9.7	8.8	8.2	6.9	6.0	6.0	4.9	§	6.9	+2.1	
Hispanic	21.2	23.0	21.5	19.2	18.1	18.0	17.5	14.3	11.2	10.4	9.5	9.0	9.8	9.6	§	6.2	-3.4	

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any 'apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 1993, the question text was changed slightly in one form to indicate that a drink meant more than a few sips. The 1993 data are based on the changed form only;

N is one half of N indicated. In 1994 the question text was changed in the remaining form. Beginning in 1994, the data are based on all forms. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 94
ALCOHOL: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days															
	1991	1992	1993 ^a	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004 ^a	2005	
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
Total	42.8	39.9‡	38.2	39.2	38.8	40.4	40.1	38.8	40.0	41.0	39.0	35.4	35.4	35.2	33.2	
Gender:																
Male	45.5	41.6‡	40.6	43.5	39.7	42.6	42.5	40.0	42.3	43.3	41.1	35.3	35.3	36.3	32.8	
Female	40.2	38.3‡	35.6	34.8	37.8	38.3	37.9	37.7	38.1	38.6	36.8	35.7	35.3	34.0	33.6	
College Plans:																
None or under 4 years	53.6	49.5‡	48.6	52.0	52.2	53.3	51.6	52.4	53.7	53.9	52.2	47.1	46.6	47.5	44.9	
Complete 4 years	40.6	37.9‡	36.1	36.4	36.4	38.3	38.1	36.5	37.9	39.1	36.8	33.5	33.6	33.6	31.6	
Region:																
Northeast	48.0	42.3‡	42.4	37.4	38.3	41.4	41.1	41.6	44.8	42.0	37.8	36.3	38.7	40.4	37.0	
Midwest	43.5	40.3‡	37.4	39.6	38.9	39.1	38.6	37.6	40.9	42.5	41.0	35.7	34.4	30.9	31.9	
South	41.7	38.2‡	38.0	40.5	39.4	41.7	40.8	39.9	38.8	39.1	38.3	33.7	34.8	36.4	33.4	
West	39.6	39.8‡	35.6	38.2	38.0	38.9	39.9	35.5	36.1	41.1	38.2	37.2	34.1	33.7	30.7	
Population Density:																
Large MSA	43.6	40.4‡	39.0	36.3	34.6	37.9	37.8	34.2	39.7	42.4	37.6	32.0	33.1	32.0	31.1	
Other MSA	41.4	38.6‡	36.2	40.1	39.9	41.0	40.2	39.0	39.7	39.3	38.3	35.2	35.6	37.0	34.4	
Non-MSA	44.8	41.9‡	41.3	40.6	41.3	42.1	42.6	43.7	41.0	42.4	41.8	40.4	38.3	35.8	33.5	
Parental Education: ^b																
1.0–2.0 (Low)	42.1	40.4‡	37.5	38.6	43.5	43.2	39.2	39.9	40.6	41.1	38.6	38.2	38.4	38.6	36.5	
2.5–3.0	43.9	40.9‡	40.6	41.5	42.3	42.6	41.1	41.2	42.3	42.8	41.4	38.0	37.4	38.5	36.0	
3.5–4.0	44.2	40.0‡	38.0	40.6	38.8	42.2	41.6	40.1	40.2	42.8	41.0	36.4	36.6	37.5	33.5	
4.5–5.0	40.7	39.4‡	36.2	37.7	37.9	37.8	39.3	36.9	38.7	40.2	37.0	33.7	32.7	32.6	31.8	
5.5–6.0 (High)	44.9	41.7‡	39.3	35.4	34.3	39.6	38.9	37.0	40.9	39.0	37.1	32.0	34.1	32.4	31.8	
Race/Ethnicity (2-year average): ^c																
White	—	44.1	43.1‡	40.4	41.0	42.2	43.0	42.7	43.0	43.9	42.7	40.0	38.7	37.8	37.0	
African American	—	30.2	29.3‡	29.7	28.0	23.9	24.6	25.1	24.4	24.7	25.3	24.3	23.7	24.6	23.0	
Hispanic	—	41.0	39.9‡	37.7	40.5	44.0	42.8	39.4	39.6	40.5	40.2	37.9	37.1	39.5	38.2	

Table continued on next page.

TABLE 94 (cont.)
ALCOHOL: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	33.8	33.4	28.8	30.4	28.9	27.2	27.6	25.7	23.5	21.5	19.9	19.7	18.6	18.4	20.3	13.1	-5.3* sss
Gender:																	
Male	33.8	33.4	28.6	31.0	30.1	28.2	28.0	26.0	23.0	20.6	18.6	17.1	18.2	18.1	§	11.2	-6.9 sss
Female	33.8	33.3	29.0	29.8	27.7	26.0	27.1	25.3	23.9	22.5	20.9	22.1	19.1	18.6	§	14.4	-4.2 ss
College Plans:																	
None or under 4 years	44.8	47.4	44.6	45.5	45.6	41.4	40.9	37.1	37.0	32.0	28.9	26.6	25.3	24.1	§	16.0	-8.1 sss
Complete 4 years	32.5	31.7	26.9	28.8	26.9	25.6	26.3	24.4	22.1	20.5	19.0	18.8	17.7	17.3	§	12.2	-5.1 sss
Region:																	
Northeast	35.7	36.2	31.4	30.5	29.5	31.5	33.3	28.4	27.0	24.0	19.8	21.1	20.4	19.5	§	16.4	-3.0
Midwest	33.7	34.4	29.5	28.9	27.4	23.2	24.5	22.4	20.4	19.1	18.1	17.2	16.6	17.7	§	14.0	-3.8
South	32.8	31.2	29.0	32.2	29.9	27.7	28.3	27.4	24.3	22.9	21.7	19.2	19.5	18.6	§	13.5	-5.1 s
West	33.9	32.9	25.9	29.0	28.7	27.5	24.9	24.6	22.6	19.4	19.2	21.8	17.8	18.0	§	9.1	-8.9 sss
Population Density:																	
Large MSA	32.5	31.8	27.4	30.3	29.3	26.5	27.4	24.8	23.3	21.2	21.0	22.4	19.6	16.7	§	9.7	-7.0 ss
Other MSA	34.5	33.3	28.7	29.7	28.4	27.7	26.9	26.5	24.0	21.4	18.1	17.9	18.8	19.1	§	14.6	-4.5 s
Non-MSA	34.4	36.2	31.1	31.9	29.4	27.1	29.1	24.9	22.6	22.4	23.0	19.8	16.8	19.5	§	15.4	-4.1 s
Parental Education: ^b																	
1.0–2.0 (Low)	33.1	36.5	35.5	37.4	34.2	30.3	32.2	29.4	26.3	23.2	20.0	21.2	20.3	19.4	§	14.8	-4.6
2.5–3.0	38.0	37.0	30.9	32.9	33.1	30.2	28.6	30.3	25.0	22.8	22.0	18.4	19.0	21.4	§	13.4	-8.0 sss
3.5–4.0	35.6	35.6	30.4	31.3	30.3	29.4	28.6	26.8	26.2	23.6	21.2	22.0	18.8	19.6	§	12.3	-7.2 sss
4.5–5.0	32.0	31.5	25.9	28.2	26.3	24.3	27.6	22.6	21.3	19.9	19.2	18.6	17.3	16.7	§	13.2	-3.5 s
5.5–6.0 (High)	31.7	28.8	26.2	27.4	24.5	24.0	26.2	23.4	22.1	21.6	20.3	21.0	22.0	18.2	§	14.7	-3.4
Race/Ethnicity (2-year average): ^c																	
White	36.3	35.9	33.3	31.5	30.8	29.1	29.1	28.2	26.4	24.6	23.0	22.4	21.9	20.8	§	19.8	-0.9
African American	21.5	21.7	20.6	20.2	20.7	20.8	20.2	19.0	16.6	15.1	13.9	11.4	10.2	11.4	§	9.1	-2.3
Hispanic	36.1	34.8	33.7	34.1	34.9	31.8	28.4	28.8	27.1	24.0	21.4	20.8	20.0	19.2	§	15.1	-4.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 1993, the question text was changed slightly in one form to indicate that a drink meant more than a few sips. The 1993 data are based on the changed form only;

N is one half of N indicated. In 1994 the question text was changed in the remaining form. Beginning in 1994, the data are based on all forms. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 95
ALCOHOL: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	68.2	68.3	71.2	72.1	71.8	72.0	70.7	69.7	69.4	67.2	65.9	65.3	66.4	63.9	60.0	57.1
Gender:																
Male	75.0	74.5	77.8	77.5	76.7	77.4	75.7	74.1	74.4	71.4	69.8	69.0	69.9	68.0	65.1	61.3
Female	62.2	61.8	65.0	67.1	67.0	66.8	65.7	65.4	64.3	62.8	62.1	61.9	63.1	59.9	54.9	52.3
College Plans:																
None or under 4 years	—	69.9	72.8	72.7	72.2	73.5	72.1	71.6	70.5	69.0	67.9	66.6	68.6	65.0	61.6	58.7
Complete 4 years	—	66.5	69.4	71.6	71.4	70.8	70.0	68.6	68.1	65.7	64.6	64.8	65.7	63.6	59.1	56.4
Region:																
Northeast	76.9	75.7	76.6	78.0	81.1	79.4	80.4	76.7	74.4	73.6	72.3	67.6	69.1	66.7	61.7	65.3
Midwest	71.1	73.2	76.4	77.2	73.9	75.1	73.6	75.0	74.4	70.6	66.8	71.3	70.7	67.9	65.9	61.5
South	62.8	60.2	64.7	67.0	65.7	65.5	62.9	61.3	64.3	62.1	60.0	58.2	60.7	58.6	55.1	51.0
West	60.0	62.2	64.4	63.1	65.5	67.6	65.3	63.8	62.9	63.6	66.2	64.5	66.7	65.0	59.3	51.6
Population Density:																
Large MSA	75.3	72.6	74.0	75.5	77.3	78.0	75.5	72.9	69.2	66.6	67.4	66.2	66.3	63.8	56.9	59.2
Other MSA	68.5	67.0	72.0	72.7	72.0	70.8	69.1	69.3	69.8	66.2	65.1	64.8	66.9	64.1	60.7	57.4
Non-MSA	63.2	66.5	67.8	68.4	67.3	69.0	68.9	67.6	69.0	69.0	65.9	65.2	65.5	63.8	61.7	54.4
Parental Education: ^b																
1.0–2.0 (Low)	58.7	62.5	62.0	62.7	64.6	65.9	62.1	61.3	61.2	58.1	58.7	56.1	56.3	54.5	47.8	47.2
2.5–3.0	70.0	71.4	72.5	71.9	71.1	72.0	70.7	69.4	69.2	67.4	65.9	65.3	67.0	64.6	59.7	57.2
3.5–4.0	69.2	67.9	73.5	75.0	74.6	73.3	71.5	72.7	70.4	69.6	66.9	66.7	67.2	64.3	62.9	57.7
4.5–5.0	69.6	71.3	74.5	77.0	76.0	74.4	73.1	74.5	73.1	69.3	68.9	68.0	68.8	66.0	62.1	60.8
5.5–6.0 (High)	67.3	72.5	77.1	79.2	75.9	77.2	77.4	74.1	75.0	70.3	67.9	69.9	70.5	67.3	62.2	60.8
Race/Ethnicity (2-year average): ^c																
White	—	—	72.8	75.0	75.3	75.4	75.4	74.6	73.9	72.8	71.2	70.2	71.0	70.6	67.3	63.8
African American	—	—	49.5	48.7	47.2	47.6	46.7	46.0	47.7	45.5	42.8	42.1	39.4	39.8	39.5	35.8
Hispanic	—	—	63.0	64.5	63.8	63.6	62.0	60.3	59.1	59.7	58.1	56.3	57.2	57.8	52.9	49.1

Table continued on next page.

TABLE 95 (cont.)
ALCOHOL: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1991	1992	1993 ^a	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004 ^c	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	54.0	51.3‡	48.6	50.1	51.3	50.8	52.7	52.0	51.0	50.0	49.8	48.6	47.5	48.0	47.0	45.3
Gender:																
Male	58.4	55.8‡	54.2	55.5	55.7	54.8	56.2	57.3	55.3	54.0	54.7	52.3	51.7	51.1	50.7	47.3
Female	49.0	46.8‡	43.4	45.2	47.0	46.9	48.9	46.9	46.8	46.1	45.1	45.1	43.8	45.1	43.3	43.0
College Plans:																
None or under 4 years	57.1	54.9‡	52.4	53.6	55.9	54.8	56.1	56.0	55.2	54.3	55.5	53.0	55.4	52.1	52.8	54.0
Complete 4 years	52.7	50.0‡	47.4	48.9	49.6	49.3	51.4	50.9	49.8	48.3	47.9	47.4	45.2	47.0	45.5	43.2
Region:																
Northeast	59.6	51.5‡	56.1	53.1	55.0	56.5	56.7	56.2	57.2	58.0	54.3	50.9	51.6	53.9	54.5	52.4
Midwest	59.7	58.0‡	51.6	53.8	55.3	51.5	51.5	51.9	51.1	52.3	54.5	52.1	50.8	49.5	48.0	46.3
South	49.1	48.1‡	47.7	49.2	50.6	51.1	51.1	51.4	49.5	44.9	45.9	46.8	43.0	43.5	43.9	43.2
West	49.7	46.7‡	39.8	44.2	43.2	42.1	52.7	49.2	47.8	48.3	44.9	45.0	47.0	47.3	43.6	41.3
Population Density:																
Large MSA	52.9	49.0‡	50.6	49.5	50.6	51.6	51.1	49.1	48.9	51.2	49.7	50.3	43.0	44.1	46.6	47.6
Other MSA	55.7	50.8‡	47.1	49.2	50.6	50.1	53.4	53.9	52.8	48.8	49.6	48.8	49.6	51.7	47.6	43.9
Non-MSA	52.0	54.1‡	49.8	52.5	53.4	51.4	52.9	51.6	50.1	50.8	50.0	45.9	49.6	45.5	46.1	44.7
Parental Education: ^b																
1.0–2.0 (Low)	49.9	45.6‡	36.6	43.5	45.9	41.2	43.8	43.8	46.8	43.4	42.9	42.2	43.6	42.1	38.2	44.4
2.5–3.0	53.3	52.3‡	49.0	49.9	52.0	48.2	51.0	50.1	50.5	50.0	49.0	47.9	47.8	46.8	47.8	46.7
3.5–4.0	54.3	51.2‡	51.2	50.1	50.6	51.4	52.1	55.6	51.1	51.3	51.4	50.9	47.9	52.4	47.8	46.3
4.5–5.0	54.8	51.0‡	49.8	52.6	51.8	53.6	55.3	52.4	50.2	48.1	51.5	48.9	47.5	47.7	50.2	42.9
5.5–6.0 (High)	58.0	55.7‡	53.2	52.2	55.1	54.2	57.4	54.7	56.0	54.0	49.5	51.1	49.3	47.4	46.0	48.2
Race/Ethnicity (2-year average): ^c																
White	60.0	56.8	55.6‡	54.0	54.5	54.8	56.4	57.7	56.3	55.1	55.3	54.0	52.3	52.2	52.3	50.7
African American	33.7	31.7	32.4‡	33.8	35.2	36.5	34.3	33.3	32.2	30.0	29.4	30.1	29.9	29.2	29.0	29.2
Hispanic	51.5	53.8	50.5‡	45.9	48.7	47.5	48.2	49.8	50.2	51.2	48.9	47.5	46.4	45.4	43.3	43.4

Table continued on next page.

TABLE 95 (cont.)
ALCOHOL: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	44.4	43.1	43.5	41.2	40.0	41.5	39.2	37.4	35.3	33.2	33.2	30.2	29.3	33.6	25.8	-3.5*
Gender:																
Male	47.1	45.8	47.8	44.2	42.1	43.8	41.8	37.4	36.0	34.5	34.1	30.4	29.8	§	23.1	-6.6 ss
Female	41.4	40.9	38.9	37.9	37.5	38.8	36.3	37.1	35.0	32.0	32.3	30.1	28.5	§	28.6	+0.1
College Plans:																
None or under 4 years	48.6	49.7	51.0	47.9	45.2	44.2	45.3	40.2	37.9	35.6	35.7	30.9	31.1	§	21.8	-9.3 sss
Complete 4 years	43.3	41.7	41.6	39.3	38.8	40.7	37.9	36.6	35.0	32.5	32.8	30.0	28.7	§	27.0	-1.7
Region:																
Northeast	51.8	49.1	48.4	49.2	46.3	48.1	45.7	41.6	42.7	36.9	37.3	32.6	30.5	§	35.8	+5.3
Midwest	47.9	42.8	45.0	40.8	41.2	41.8	39.0	40.2	33.6	33.7	34.0	30.4	29.4	§	26.5	-2.8
South	43.1	42.9	42.0	39.6	37.5	36.6	35.4	34.6	33.1	31.3	31.7	30.0	28.3	§	21.6	-6.7 ss
West	36.6	39.2	39.2	37.4	38.0	43.1	39.8	36.1	35.4	33.8	32.1	28.5	30.4	§	24.3	-6.1
Population Density:																
Large MSA	47.1	46.1	46.7	42.4	41.1	44.1	39.2	37.7	38.5	33.0	34.3	30.0	27.9	§	31.7	+3.7
Other MSA	42.1	40.4	41.5	41.6	41.1	41.8	40.8	38.7	33.4	33.6	30.5	28.3	29.1	§	21.2	-7.9 ss
Non-MSA	45.2	45.8	43.7	38.5	35.8	36.9	35.3	33.7	35.0	32.7	37.5	34.9	32.1	§	26.7	-5.4 s
Parental Education: ^b																
1.0–2.0 (Low)	38.3	41.4	39.8	36.4	36.7	41.4	36.1	32.5	32.5	31.7	28.9	23.7	23.4	§	18.3	-5.1
2.5–3.0	46.0	42.4	43.4	42.4	38.8	41.4	40.3	38.2	35.3	31.4	31.1	27.1	28.5	§	21.0	-7.4 ss
3.5–4.0	43.6	43.2	46.9	41.6	41.2	42.7	41.1	37.9	36.5	32.8	33.2	32.4	29.3	§	25.1	-4.2
4.5–5.0	45.0	43.9	42.0	42.1	40.1	41.7	39.5	38.7	36.1	35.7	35.8	32.0	30.0	§	31.3	+1.3
5.5–6.0 (High)	47.6	45.9	43.3	38.4	41.9	41.1	38.7	40.6	36.3	34.4	38.3	35.5	35.6	§	32.0	-3.7
Race/Ethnicity (2-year average): ^c																
White	49.3	48.6	47.2	45.4	43.8	43.8	43.6	42.5	40.9	39.0	39.0	38.0	34.9	§	35.3	+0.4
African American	28.7	28.6	30.5	31.4	30.1	29.6	28.4	25.9	24.0	21.8	20.3	21.2	19.4	§	18.2	-1.2
Hispanic	41.4	38.9	40.1	40.1	39.7	39.8	39.0	37.0	36.3	34.9	30.9	26.4	24.4	§	21.8	-2.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 1993, the question text was changed slightly in one form to indicate that a drink meant more than a few sips. The 1993 data are based on the changed form only;

N is one half of N indicated. In 1994 the question text was changed in the remaining form. Beginning in 1994, the data are based on all forms. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 96
ALCOHOL: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 8

	Percentage who used daily in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993^a</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004^b</u>	<u>2005</u>
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	0.6	0.6‡	1.0	1.0	0.7	1.0	0.8	0.9	1.0	0.8	0.9	0.7	0.8	0.6	0.5
Gender:															
Male	0.7	0.8‡	1.5	1.2	1.0	1.4	0.9	1.2	1.3	1.0	1.2	0.8	0.8	0.6	0.6
Female	0.4	0.4‡	0.5	0.7	0.5	0.6	0.6	0.5	0.6	0.5	0.6	0.4	0.7	0.6	0.3
College Plans:															
None or under 4 years	1.6	2.2‡	2.7	2.8	1.9	3.0	2.0	3.1	3.4	1.8	2.8	2.8	2.3	2.2	1.9
Complete 4 years	0.4	0.4‡	0.7	0.7	0.6	0.7	0.6	0.6	0.7	0.6	0.6	0.5	0.6	0.5	0.3
Region:															
Northeast	0.4	0.5‡	0.7	1.0	0.6	1.0	0.8	1.0	1.0	0.6	0.8	0.7	0.6	0.3	0.3
Midwest	0.6	0.6‡	0.9	0.7	0.7	1.1	0.8	1.0	1.3	1.2	0.7	0.7	0.9	0.6	0.6
South	0.6	0.8‡	1.3	1.4	0.7	1.1	0.8	1.0	1.0	0.7	1.2	0.8	0.8	0.8	0.5
West	0.5	0.5‡	0.9	0.7	0.8	0.8	0.7	0.5	0.9	0.6	0.6	0.4	0.6	0.6	0.4
Population Density:															
Large MSA	0.4	0.6‡	1.1	0.6	0.6	0.9	0.4	0.7	1.0	0.4	0.5	0.4	0.7	0.5	0.5
Other MSA	0.5	0.6‡	0.6	1.0	0.6	1.0	0.9	0.8	0.8	0.7	0.9	0.8	0.7	0.5	0.4
Non-MSA	0.8	0.7‡	1.6	1.4	1.1	1.3	0.9	1.1	1.5	1.4	1.3	0.8	1.0	0.9	0.6
Parental Education: ^b															
1.0–2.0 (Low)	1.6	1.4‡	1.5	1.5	1.4	1.3	1.6	1.8	1.9	0.8	1.7	1.1	2.5	1.5	0.9
2.5–3.0	0.8	0.6‡	0.6	0.9	0.8	1.3	0.5	0.8	0.9	1.1	1.2	0.8	1.1	0.6	0.5
3.5–4.0	0.3	0.5‡	0.8	1.0	1.0	1.2	0.8	0.6	1.2	0.6	0.5	0.7	0.6	0.6	0.5
4.5–5.0	0.3	0.5‡	0.5	0.9	0.4	0.5	0.7	0.7	0.4	0.6	0.6	0.5	0.3	0.2	0.3
5.5–6.0 (High)	0.4	0.6‡	1.8	0.9	0.6	1.0	0.5	1.0	1.1	0.6	0.7	0.4	0.5	0.4	0.1
Race/Ethnicity (2-year average): ^c															
White	—	0.5	0.6‡	0.7	0.8	0.8	0.9	0.7	0.8	0.8	0.8	0.7	0.6	0.6	0.4
African American	—	0.5	1.0‡	1.2	0.9	0.8	0.6	0.6	0.9	0.5	0.4	0.5	0.5	0.6	0.6
Hispanic	—	1.1	1.2‡	1.6	1.5	1.4	1.4	1.3	1.3	1.1	1.2	1.3	1.4	1.4	0.8

Table continued on next page.

TABLE 96 (cont.)
ALCOHOL: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 8

	Percentage who used daily in last 30 days																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	0.5	0.6	0.7	0.5	0.5	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.1	0.2	0.4	0.3	0.0*
Gender:																	
Male	0.5	0.8	1.0	0.6	0.5	0.4	0.4	0.3	0.3	0.2	0.2	0.2	0.1	0.3	§	0.3	0.0
Female	0.4	0.3	0.4	0.4	0.5	0.3	0.2	0.2	0.3	0.1	0.1	0.1	0.1	0.1	§	0.1	0.0
College Plans:																	
None or under 4 years	1.5	2.4	2.7	1.8	2.2	1.6	1.9	1.5	1.3	1.0	0.5	0.6	0.3	0.7	§	0.4	-0.3
Complete 4 years	0.3	0.4	0.6	0.4	0.4	0.3	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	§	0.2	+0.1
Region:																	
Northeast	0.3	0.4	0.6	0.1	0.4	0.1	0.3	0.0	0.2	0.1	0.1	0.2	0.1	0.1	§	0.0	-0.1
Midwest	0.5	0.6	0.8	0.3	0.4	0.3	0.2	0.1	0.2	0.2	0.0	0.1	0.0	0.2	§	0.3	+0.1
South	0.6	0.7	0.9	0.7	0.7	0.4	0.4	0.5	0.4	0.3	0.3	0.1	0.2	0.3	§	0.2	-0.1
West	0.4	0.5	0.4	0.6	0.5	0.7	0.4	0.2	0.4	0.1	0.1	0.2	0.2	0.3	§	0.6	+0.3
Population Density:																	
Large MSA	0.4	0.4	0.7	0.3	0.5	0.3	0.4	0.2	0.2	0.2	0.1	0.2	0.1	0.1	§	0.2	+0.1
Other MSA	0.5	0.6	0.8	0.4	0.5	0.4	0.4	0.2	0.3	0.2	0.1	0.1	0.2	0.3	§	0.2	0.0
Non-MSA	0.6	0.8	0.7	0.8	0.6	0.5	0.2	0.5	0.5	0.2	0.5	0.2	0.1	0.4	§	0.5	+0.1
Parental Education: ^b																	
1.0–2.0 (Low)	0.7	1.3	1.8	2.1	1.3	0.5	0.3	0.2	0.8	0.2	0.2	0.3	0.1	0.3	§	0.7	+0.3
2.5–3.0	0.6	0.5	0.9	0.5	0.5	0.7	0.5	0.2	0.2	0.3	0.3	0.3	0.2	0.2	§	0.4	+0.2
3.5–4.0	0.6	0.7	0.7	0.4	0.6	0.4	0.4	0.6	0.3	0.2	0.3	0.2	0.0	0.3	§	0.1	-0.2
4.5–5.0	0.3	0.2	0.2	0.1	0.3	0.1	0.1	0.1	0.4	0.1	0.1	0.1	0.1	0.0	§	0.1	+0.1
5.5–6.0 (High)	0.3	0.4	0.6	0.4	0.3	0.4	0.2	0.2	0.2	0.1	0.2	0.0	0.2	0.2	§	0.4	+0.2
Race/Ethnicity (2-year average): ^c																	
White	0.4	0.4	0.6	0.5	0.4	0.4	0.3	0.2	0.2	0.2	0.2	0.2	0.1	0.2	§	0.4	+0.3 s
African American	0.5	0.5	0.6	0.4	0.3	0.4	0.3	0.2	0.2	0.1	0.1	0.2	0.2	0.1	§	0.1	0.0
Hispanic	0.6	0.8	1.0	1.0	0.9	0.8	0.7	0.4	0.3	0.3	0.2	0.2	0.2	0.1	§	0.2	+0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. 'Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 1993, the question text was changed slightly in one form to indicate that a drink meant more than a few sips. The 1993 data are based on the changed form only;

N is one half of N indicated. In 1994 the question text was changed in the remaining form. Beginning in 1994, the data are based on all forms. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 97
ALCOHOL: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 10

	Percentage who used daily in last 30 days														
	1991	1992	1993 ^a	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004 ^a	2005
Approximate weighted N =	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	1.4	1.3‡	1.8	1.7	1.7	1.6	1.7	1.9	1.9	1.8	1.9	1.8	1.5	1.3	1.3
Gender:															
Male	2.3	1.6‡	2.6	2.5	2.6	2.4	2.5	2.8	2.9	2.7	2.9	2.6	2.0	1.9	1.9
Female	0.4	0.8‡	1.0	0.8	0.8	0.8	0.8	1.2	0.9	0.8	0.9	1.0	1.0	0.7	0.7
College Plans:															
None or under 4 years	2.0	3.2‡	3.6	3.6	4.0	3.2	4.0	4.3	4.5	4.0	4.6	3.7	3.0	3.7	3.9
Complete 4 years	1.2	0.8‡	1.5	1.2	1.3	1.3	1.3	1.5	1.5	1.4	1.4	1.4	1.2	1.0	1.0
Region:															
Northeast	1.3	0.9‡	1.8	0.9	1.5	1.0	1.5	2.3	1.9	1.3	1.6	1.4	1.5	0.9	1.4
Midwest	1.4	1.5‡	1.8	1.7	1.8	1.6	1.6	2.1	2.3	1.9	1.6	1.7	1.2	1.0	1.1
South	1.6	1.2‡	1.8	2.3	1.8	1.8	1.9	1.9	1.6	2.0	1.9	1.8	1.5	1.6	1.4
West	1.0	1.4‡	1.7	1.6	1.8	1.9	1.8	1.5	1.9	1.7	2.4	2.0	1.8	1.6	1.4
Population Density:															
Large MSA	1.3	1.2‡	1.1	1.3	1.3	1.5	1.2	1.5	2.0	1.6	1.8	1.4	1.4	0.7	0.9
Other MSA	1.2	1.0‡	1.4	1.7	1.8	1.5	1.6	1.7	1.7	1.6	1.6	1.8	1.4	1.5	1.4
Non-MSA	1.6	1.9‡	3.2	2.1	2.2	1.9	2.5	2.9	2.2	2.3	2.5	2.1	2.0	1.5	1.7
Parental Education: ^b															
1.0–2.0 (Low)	2.6	2.0‡	3.1	1.9	2.9	1.6	2.1	2.7	2.4	2.2	2.9	2.0	2.3	1.6	1.7
2.5–3.0	1.5	1.1‡	2.1	1.9	2.1	1.6	2.1	2.4	2.1	2.3	1.9	2.1	1.6	1.6	2.0
3.5–4.0	1.1	1.3‡	1.7	1.7	1.7	1.6	1.9	2.0	1.9	1.9	1.5	1.5	1.4	1.1	1.2
4.5–5.0	1.1	0.8‡	1.0	1.3	1.3	1.5	1.2	1.2	1.5	1.4	1.7	1.6	1.6	1.1	0.7
5.5–6.0 (High)	1.2	1.5‡	2.0	1.2	0.9	1.4	0.8	1.9	2.0	1.2	1.7	1.8	0.9	1.2	0.9
Race/Ethnicity (2-year average): ^c															
White	—	1.3	1.4‡	1.6	1.7	1.7	1.6	1.8	1.9	1.8	1.7	1.8	1.7	1.5	1.4
African American	—	0.9	1.2‡	1.2	1.1	0.6	0.7	1.0	1.0	1.1	1.4	1.2	0.9	0.5	0.5
Hispanic	—	1.6	1.9‡	1.8	2.3	2.3	2.0	2.4	2.4	2.0	2.6	2.4	1.8	1.7	1.6

Table continued on next page.

TABLE 97 (cont.)
ALCOHOL: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 10

	Percentage who used daily in last 30 days																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>																	
Total	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	-0.2*
Gender:																	
Male	1.4	1.4	1.0	1.1	1.1	0.8	1.0	0.9	0.8	0.5	0.5	0.6	0.5	0.6	1.0	0.4	-0.2*
Female	1.8	1.9	1.4	1.5	1.7	1.1	1.4	1.3	1.1	0.7	0.8	0.7	0.6	0.9	§	0.5	-0.4 s
College Plans:																	
None or under 4 years	1.0	0.9	0.6	0.7	0.6	0.4	0.5	0.5	0.6	0.3	0.3	0.4	0.3	0.4	§	0.3	0.0
Complete 4 years	3.2	3.3	2.1	3.5	3.3	2.2	3.0	1.9	2.7	1.3	1.1	1.5	0.9	0.9	§	0.9	-0.3 s
Region:																	
Northeast	1.2	1.2	0.9	0.9	0.9	0.6	0.7	0.8	0.6	0.4	0.5	0.4	0.4	0.6	§	0.3	-0.3 s
Midwest	1.6	1.5	1.1	0.8	0.7	0.8	1.2	0.9	0.5	0.3	0.6	0.5	0.8	0.2	§	0.6	+0.4
South	1.3	1.6	0.9	1.1	1.2	0.5	0.8	0.6	0.6	0.3	0.5	0.6	0.3	0.6	§	0.3	-0.4
West	1.6	1.2	1.1	1.3	1.1	0.9	1.2	1.3	1.2	0.7	0.6	0.5	0.6	0.9	§	0.5	-0.4
Population Density:																	
Large MSA	1.2	1.6	0.9	1.2	1.5	0.8	0.8	0.8	0.7	0.5	0.5	0.6	0.4	0.5	§	0.4	-0.1
Other MSA	1.3	1.3	0.9	1.0	1.2	0.8	0.8	1.0	0.6	0.4	0.5	0.6	0.4	0.2	§	0.2	0.0
Non-MSA	1.4	1.5	1.0	1.1	1.0	0.7	0.8	0.8	0.7	0.5	0.4	0.3	0.5	0.8	§	0.5	-0.3
Parental Education: ^b																	
1.0–2.0 (Low)	1.8	1.6	1.2	1.3	1.3	0.9	1.5	0.9	1.4	0.6	0.9	1.1	0.7	0.9	§	0.8	-0.1
2.5–3.0	2.9	3.6	2.1	1.6	2.3	0.8	1.5	1.1	1.8	0.7	0.7	0.5	0.3	1.3	§	1.3	0.0
3.5–4.0	1.9	1.6	1.0	1.4	1.6	0.9	1.2	1.5	0.9	0.9	0.5	0.8	0.6	1.2	§	0.4	-0.8 s
4.5–5.0	1.5	1.2	1.0	1.3	0.6	0.8	0.7	0.9	0.9	0.4	0.7	0.5	0.2	0.6	§	0.3	-0.3
5.5–6.0 (High)	0.9	1.1	0.4	0.6	1.0	0.8	0.8	0.7	0.4	0.4	0.4	0.2	0.6	0.3	§	0.3	0.0
Race/Ethnicity (2-year average): ^c																	
White	0.7	1.0	0.9	0.6	0.7	0.5	1.1	0.4	0.7	0.3	0.3	0.9	0.9	0.4	§	0.3	-0.1
African American	1.4	1.4	1.2	1.0	1.0	0.9	0.9	1.0	0.9	0.6	0.5	0.6	0.6	0.6	§	0.5	0.0
Hispanic	0.8	0.8	0.7	0.9	0.8	0.5	0.5	0.4	0.3	0.4	0.2	0.2	0.2	0.2	§	0.3	0.0
	2.0	2.1	2.0	1.9	1.9	1.6	1.1	1.0	1.0	0.9	0.6	0.5	0.4	0.6	§	1.3	+0.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' ‡ ' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 1993, the question text was changed slightly in one form to indicate that a drink meant more than a few sips. The 1993 data are based on the changed form only;

N is one half of N indicated. In 1994 the question text was changed in the remaining form. Beginning in 1994, the data are based on all forms. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 98
ALCOHOL: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total		5.6	6.1	5.7	6.9	6.0	6.0	5.8	5.5	4.8	5.0	4.7	4.8	4.2	4.2	3.7
Gender:																
Male	—	8.1	8.6	8.3	9.7	8.6	8.4	7.7	7.7	6.6	7.0	6.7	7.2	6.2	6.0	5.2
Female	—	2.7	3.6	3.2	4.0	3.6	3.4	3.4	2.9	2.7	3.0	2.8	2.5	2.3	2.2	1.9
College Plans:																
None or under 4 years	—	7.3	8.0	7.3	9.1	8.1	7.7	7.5	6.7	6.0	6.5	6.8	7.0	6.0	6.1	4.9
Complete 4 years	—	3.6	4.0	4.1	5.0	4.4	4.6	4.1	4.0	3.6	4.0	3.5	3.6	3.4	3.2	3.0
Region:																
Northeast	—	6.3	6.5	6.2	8.7	7.4	7.5	6.3	5.9	6.5	6.6	4.9	4.2	4.2	4.3	4.6
Midwest	—	6.9	6.7	7.0	6.8	6.7	6.6	7.6	6.2	4.3	4.4	4.9	5.3	4.8	5.1	3.9
South	—	4.6	5.9	5.0	7.2	5.2	5.2	5.0	5.4	5.3	5.0	4.9	5.1	4.0	4.0	3.4
West	—	3.8	4.3	3.8	3.8	4.5	4.3	2.8	3.6	2.8	4.0	4.2	4.5	3.9	3.5	2.9
Population Density:																
Large MSA	—	5.4	5.9	6.2	7.0	7.1	6.5	6.4	5.6	5.1	5.3	4.1	3.7	3.5	4.0	4.7
Other MSA	—	5.4	5.8	5.5	6.1	5.4	5.3	5.1	5.7	4.5	5.1	4.5	5.4	4.5	3.9	3.2
Non-MSA	—	6.1	6.5	5.7	7.9	6.1	6.6	6.1	5.9	5.1	4.8	5.8	4.9	4.5	5.2	3.8
Parental Education: ^b																
1.0–2.0 (Low)	—	5.7	5.0	4.7	6.2	6.3	5.8	5.2	6.1	5.2	5.5	5.5	5.8	5.0	3.9	3.1
2.5–3.0	—	6.2	7.1	6.4	7.2	6.5	6.3	6.1	5.2	5.1	5.2	4.7	4.5	4.3	5.0	3.9
3.5–4.0	—	5.0	5.3	5.3	6.4	5.5	6.1	5.5	5.3	4.6	4.7	4.6	4.9	3.8	4.0	3.5
4.5–5.0	—	5.6	5.7	5.6	6.7	5.8	4.8	5.6	5.1	4.1	4.5	4.4	4.8	3.9	3.8	3.4
5.5–6.0 (High)	—	5.0	6.3	5.6	8.8	5.5	6.3	5.2	5.6	4.1	4.8	4.0	4.5	4.8	3.0	4.2
Race/Ethnicity (2-year average): ^c																
White	—	—	6.0	6.2	6.6	6.8	6.5	6.3	5.9	5.3	5.1	5.2	5.1	4.8	4.5	4.2
African American	—	—	2.2	2.0	2.1	2.5	2.3	2.2	2.3	2.2	2.2	2.1	2.1	2.2	2.4	2.1
Hispanic	—	—	5.3	4.6	4.0	4.3	4.5	3.7	3.2	4.0	5.0	4.7	4.6	4.6	4.3	3.8

Table continued on next page.

TABLE 98 (cont.)
ALCOHOL: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															
	1991	1992	1993 ^a	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004 ^c	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	3.6	3.4‡	3.4	2.9	3.6	3.7	3.9	4.0	3.4	2.9	3.6	3.5	3.2	2.8	3.1	3.0
Gender:																
Male	5.3	5.2‡	5.3	4.8	5.5	5.2	5.7	6.4	5.2	4.7	5.5	5.3	4.7	4.1	4.4	4.3
Female	1.6	1.6‡	1.6	1.2	1.6	2.0	1.9	1.4	1.4	1.1	1.7	1.7	1.6	1.4	1.6	1.5
College Plans:																
None or under 4 years	5.4	5.3‡	4.3	4.4	5.4	5.0	6.0	6.8	5.9	4.8	5.3	4.8	5.0	4.3	5.6	4.9
Complete 4 years	2.9	2.7‡	2.9	2.4	2.9	3.1	3.2	3.0	2.6	2.2	2.9	3.1	2.4	2.2	2.3	2.4
Region:																
Northeast	3.3	2.9‡	5.3	3.0	2.8	4.1	3.7	4.1	2.8	3.0	3.1	2.8	3.7	3.3	3.1	3.3
Midwest	3.9	3.7‡	3.3	3.1	4.3	3.8	4.5	4.1	3.2	3.1	4.5	4.6	3.4	2.6	3.6	3.3
South	4.1	3.7‡	3.5	3.0	3.8	3.6	4.1	4.2	3.9	2.8	3.5	3.3	3.0	2.8	3.2	3.1
West	2.7	2.9‡	1.8	2.5	2.8	3.1	3.2	3.2	3.2	2.9	3.0	3.0	2.6	2.5	2.1	2.4
Population Density:																
Large MSA	3.3	3.4‡	2.7	2.5	3.5	4.4	3.9	3.4	2.3	3.0	3.6	3.4	2.1	2.5	2.8	3.4
Other MSA	3.5	3.1‡	3.6	2.8	3.5	3.3	4.0	4.1	3.4	2.7	3.6	3.2	3.5	3.3	3.1	2.8
Non-MSA	4.1	4.1‡	3.6	3.6	3.6	3.7	3.9	4.3	4.4	3.4	3.7	4.2	3.9	2.3	3.4	3.0
Parental Education: ^b																
1.0–2.0 (Low)	4.2	3.0‡	3.6	2.3	3.9	2.6	3.8	5.3	4.0	3.5	3.5	2.5	2.7	3.4	2.6	3.1
2.5–3.0	4.1	4.0‡	2.6	3.4	3.9	3.5	3.7	4.1	3.6	2.9	3.4	3.4	3.6	3.0	3.7	3.4
3.5–4.0	3.0	2.9‡	3.7	2.5	2.7	3.4	3.2	4.0	3.3	3.0	3.4	3.7	2.9	2.9	2.5	2.8
4.5–5.0	3.1	2.8‡	3.6	2.8	3.9	4.0	3.9	3.6	2.9	2.4	3.3	3.1	3.0	2.0	3.1	2.7
5.5–6.0 (High)	4.1	4.7‡	2.9	3.1	3.5	3.6	4.7	3.1	3.2	2.9	3.9	4.0	2.3	2.6	2.7	2.8
Race/Ethnicity (2-year average): ^c																
White	3.8	3.6	3.1‡	3.1	3.3	3.7	4.1	4.3	3.9	3.3	3.5	3.8	3.6	3.2	3.3	3.4
African American	1.6	1.9	2.3‡	2.6	2.5	2.2	1.8	1.8	1.6	1.4	1.6	1.6	1.5	1.2	1.1	1.5
Hispanic	3.6	3.6	2.8‡	3.1	3.5	4.0	4.0	4.3	4.8	5.0	4.1	3.3	2.8	2.7	2.9	2.9

Table continued on next page.

TABLE 98 (cont.)
ALCOHOL: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	3.1	2.8	2.5	2.7	2.1	2.5	2.2	1.9	1.9	1.3	1.6	1.2	1.7	2.7	0.9	-0.8* ss
Gender:																
Male	4.3	4.0	3.6	3.6	2.9	3.6	3.0	2.8	2.3	1.7	2.2	1.6	2.4	§	1.1	-1.3 sss
Female	1.9	1.7	1.3	1.6	1.2	1.2	1.2	1.0	1.4	0.8	0.7	0.5	0.9	§	0.6	-0.3
College Plans:																
None or under 4 years	4.8	5.1	5.0	4.4	4.1	3.8	4.8	3.7	3.9	2.6	2.9	2.4	3.1	§	1.2	-1.9 sss
Complete 4 years	2.6	2.2	1.8	2.2	1.6	2.1	1.5	1.4	1.4	1.0	1.2	0.7	1.1	§	0.7	-0.5 s
Region:																
Northeast	3.5	2.2	2.2	2.9	2.8	3.3	2.2	1.3	2.0	1.6	1.2	1.2	0.9	§	0.9	0.0
Midwest	3.3	2.9	3.2	3.2	2.6	2.6	2.3	2.1	2.1	1.4	1.2	0.8	1.6	§	1.1	-0.5
South	3.4	3.8	2.3	2.9	1.7	2.1	2.2	1.9	1.8	1.3	2.1	1.5	2.2	§	1.1	-1.1 ss
West	2.0	1.7	2.1	1.8	1.8	2.5	1.9	1.9	1.8	1.3	1.4	0.9	1.4	§	0.6	-0.8
Population Density:																
Large MSA	3.1	2.1	2.5	2.7	2.3	2.2	1.8	1.6	2.1	1.3	1.5	1.0	1.1	§	1.3	+0.2
Other MSA	3.3	2.6	2.6	2.7	2.1	2.8	2.2	2.0	1.5	1.2	1.6	1.2	1.5	§	0.6	-0.9 ss
Non-MSA	2.6	4.2	2.3	2.8	2.0	2.4	2.7	1.9	2.7	1.7	1.7	1.6	3.1	§	1.1	-2.0 ss
Parental Education: ^b																
1.0–2.0 (Low)	3.6	2.6	3.3	2.9	2.4	2.8	2.3	2.6	3.1	1.6	1.3	1.0	2.0	§	0.9	-1.1
2.5–3.0	3.6	2.6	2.4	2.9	2.9	2.2	3.0	1.5	1.8	1.4	1.6	1.0	1.8	§	0.6	-1.2 ss
3.5–4.0	2.9	2.7	2.5	2.3	2.1	2.5	1.7	1.5	1.8	0.9	1.4	1.2	1.3	§	0.9	-0.4
4.5–5.0	3.0	2.9	2.1	2.5	1.3	2.5	2.1	2.2	1.6	1.5	1.5	1.1	1.5	§	1.1	-0.4
5.5–6.0 (High)	2.4	2.6	1.5	2.0	2.0	2.5	1.7	1.8	1.5	0.9	1.1	1.4	1.6	§	0.9	-0.7
Race/Ethnicity (2-year average): ^c																
White	3.4	3.4	2.9	2.6	2.5	2.5	2.4	2.0	1.9	1.7	1.7	1.7	1.6	§	1.3	-0.3
African American	1.5	1.3	1.4	2.1	2.0	1.5	1.5	1.4	1.1	0.8	0.9	0.9	0.8	§	0.3	-0.5 s
Hispanic	3.1	2.8	2.5	2.5	2.4	2.3	2.7	2.3	2.0	1.9	1.2	0.8	1.0	§	2.7	+1.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. '‡' indicates some change in the question. See relevant footnote. See relevant figure to assess the impact of the wording changes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn 1993, the question text was changed slightly in one form to indicate that a drink meant more than a few sips. The 1993 data are based on the changed form only;

N is one half of N indicated. In 1994 the question text was changed in the remaining form. Beginning in 1994, the data are based on all forms. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 99
BEEN DRUNK: Trends in 30-Day Prevalence by Subgroups in Grade 8

	Percentage who had been drunk in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
	<i>Approximate weighted N =</i>														
	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	7.6	7.5	7.8	8.7	8.3	9.6	8.2	8.4	9.4	8.3	7.7	6.7	6.7	6.2	6.0
Gender:															
Male	8.4	7.4	7.8	9.0	8.2	9.7	8.4	8.5	10.2	8.2	7.8	7.1	6.6	5.8	5.9
Female	7.0	7.6	7.8	8.3	8.2	9.5	7.9	8.2	8.6	8.1	7.4	6.3	6.8	6.6	6.2
College Plans:															
None or under 4 years	15.8	17.2	18.4	20.0	17.2	19.3	18.7	21.4	22.4	18.9	18.9	15.5	17.0	16.1	15.7
Complete 4 years	6.4	6.1	6.4	7.3	7.3	8.2	7.1	6.9	8.0	7.0	6.5	5.9	5.7	5.2	5.1
Region:															
Northeast	5.7	6.4	6.2	8.2	8.2	9.7	7.9	6.9	9.4	8.3	8.0	5.3	4.9	4.8	4.2
Midwest	7.7	7.6	7.3	8.3	8.3	10.2	8.2	10.4	11.6	9.7	8.2	7.0	8.1	6.8	6.2
South	8.8	8.2	8.3	8.8	8.4	9.1	8.3	7.8	9.5	7.4	8.1	7.6	7.2	6.4	7.1
West	7.3	6.9	9.4	9.6	8.2	9.8	8.3	8.3	6.6	8.1	6.0	5.9	6.0	6.4	5.8
Population Density:															
Large MSA	7.4	7.0	6.0	7.6	7.2	8.9	6.7	6.9	7.8	7.1	6.3	5.2	5.6	5.4	5.4
Other MSA	7.3	7.4	8.4	9.7	8.9	9.9	8.6	7.5	8.4	7.2	7.9	7.3	6.8	6.5	6.1
Non-MSA	8.4	8.2	8.8	7.9	8.6	10.0	9.2	11.7	13.3	12.0	8.9	7.3	8.1	6.5	6.9
Parental Education: ^a															
1.0–2.0 (Low)	13.4	11.0	10.4	12.5	13.1	11.1	11.5	13.1	14.5	14.0	12.1	10.5	13.4	10.0	9.3
2.5–3.0	9.2	8.8	9.2	9.3	9.6	11.9	9.3	9.5	11.7	10.5	9.7	8.5	9.4	7.4	8.4
3.5–4.0	6.9	7.6	8.5	9.3	9.4	10.4	10.2	9.1	9.9	8.0	8.7	6.7	7.1	6.8	6.7
4.5–5.0	6.1	6.5	5.9	7.5	6.4	8.7	6.7	7.0	6.9	5.6	5.7	5.5	4.5	5.0	4.6
5.5–6.0 (High)	6.8	4.9	6.7	7.6	6.0	7.1	5.8	6.9	8.7	7.1	4.6	4.8	4.2	4.4	3.5
Race/Ethnicity (2-year average): ^b															
White	—	7.7	7.8	8.4	8.9	9.7	9.7	9.1	9.8	10.0	9.0	8.0	7.2	6.8	6.6
African American	—	5.4	5.1	5.6	5.6	5.5	4.6	3.9	4.9	4.7	4.0	4.0	4.2	4.1	3.8
Hispanic	—	9.9	9.9	10.8	10.8	10.8	10.4	9.8	9.9	8.5	7.8	8.4	8.5	8.1	7.0

Table continued on next page.

TABLE 99 (cont.)
BEEN DRUNK: Trends in 30-Day Prevalence by Subgroups in Grade 8

	Percentage who had been drunk in last 30 days																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	6.2	5.5	5.4	5.4	5.0	4.4	3.6	3.5	2.7	3.1	1.8	2.2	2.1	2.6	3.4	2.0	-0.6*
Gender:																	
Male	6.1	5.3	5.3	5.3	4.8	4.4	3.3	3.0	2.7	2.9	1.8	1.6	1.8	2.3	§	1.3	-1.1 ss
Female	6.2	5.6	5.4	5.3	5.2	4.2	3.8	4.0	2.7	3.0	1.7	2.8	2.3	2.9	§	2.4	-0.5
College Plans:																	
None or under 4 years	15.4	16.3	14.9	14.9	12.9	10.9	9.5	10.9	6.7	6.7	5.3	5.2	4.5	4.9	§	3.0	-1.8
Complete 4 years	5.4	4.6	4.6	4.6	4.4	3.9	3.1	3.0	2.4	2.7	1.5	1.9	1.8	2.3	§	1.8	-0.5
Region:																	
Northeast	5.8	3.5	4.5	4.7	4.2	2.9	3.5	2.7	2.0	2.6	0.9	1.5	1.5	1.8	§	1.1	-0.7
Midwest	6.3	5.7	5.3	5.0	4.7	4.1	2.2	2.6	2.3	2.4	1.4	2.0	2.0	2.2	§	2.6	+0.4
South	6.9	6.1	6.8	6.5	5.7	4.4	4.4	4.5	2.8	3.3	2.2	2.4	2.0	2.9	§	1.9	-1.0
West	5.2	5.7	4.2	4.4	4.8	5.7	3.5	3.3	3.4	3.6	1.9	2.7	2.8	3.1	§	2.3	-0.8
Population Density:																	
Large MSA	5.1	4.8	5.4	4.0	4.7	3.5	3.1	3.1	2.5	2.6	1.2	2.1	2.0	2.1	§	1.1	-1.0 ss
Other MSA	6.4	5.6	5.2	5.8	5.2	4.8	4.1	3.3	2.8	3.3	1.8	2.1	2.1	2.7	§	2.2	-0.5
Non-MSA	7.5	6.4	5.9	6.3	5.0	4.7	3.2	4.6	3.0	3.2	2.6	2.9	2.3	3.3	§	3.1	-0.1
Parental Education: ^a																	
1.0–2.0 (Low)	9.4	10.6	10.1	10.1	7.8	7.9	6.0	6.4	3.8	4.9	3.5	2.8	2.2	2.9	§	2.2	-0.7
2.5–3.0	7.9	7.9	7.4	6.3	7.1	6.6	5.6	4.6	3.4	4.1	2.2	3.0	2.5	3.5	§	2.8	-0.7
3.5–4.0	7.1	6.0	5.8	5.4	5.3	4.8	4.1	4.3	2.9	2.6	1.5	2.6	2.3	3.7	§	2.4	-1.3
4.5–5.0	4.4	3.1	3.7	4.4	4.2	3.2	2.6	2.4	2.4	2.3	1.4	1.9	1.7	2.3	§	2.0	-0.3
5.5–6.0 (High)	4.6	3.1	3.3	3.4	2.6	2.7	1.6	2.2	2.2	2.5	1.4	1.8	1.9	1.6	§	1.7	+0.1
Race/Ethnicity (2-year average): ^b																	
White	6.4	5.9	5.5	5.5	5.3	4.7	3.9	3.3	2.9	2.6	2.3	2.0	2.1	2.5	§	3.0	+0.5
African American	3.6	3.7	3.8	3.6	3.4	2.9	2.7	3.0	2.4	2.0	1.7	1.4	1.6	1.2	§	1.7	+0.5
Hispanic	7.2	7.4	6.7	6.2	5.8	5.6	5.5	4.5	3.6	3.5	2.8	2.2	2.2	2.4	§	2.3	0.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 100
BEEN DRUNK: Trends in 30-Day Prevalence by Subgroups in Grade 10

		Percentage who had been drunk in last 30 days														
		<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>		<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>
Total		20.5	18.1	19.8	20.3	20.8	21.3	22.4	21.1	22.5	23.5	21.9	18.3	18.2	18.5	17.6
Gender:																
Male		22.3	18.6	21.4	23.2	21.9	23.0	24.6	22.3	25.4	26.2	24.2	19.3	18.8	20.2	18.2
Female		18.7	17.5	18.1	17.2	19.6	19.8	20.2	19.9	19.8	20.9	19.7	17.4	17.7	17.0	16.8
College Plans:																
None or under 4 years		29.5	26.3	29.0	31.1	31.4	32.0	35.5	33.5	34.6	35.1	34.8	27.4	27.4	27.5	27.8
Complete 4 years		18.6	16.4	17.9	18.0	19.0	19.7	20.3	19.1	20.7	21.8	19.8	16.8	16.9	17.4	16.3
Region:																
Northeast		23.9	18.8	20.0	19.0	19.5	22.4	21.9	23.1	25.8	22.8	19.9	18.1	20.3	22.4	20.5
Midwest		21.8	18.9	20.1	21.0	22.6	22.0	23.3	21.8	26.0	27.6	24.0	18.7	18.8	16.6	17.4
South		19.2	16.8	19.8	20.9	20.9	21.4	22.0	21.9	20.3	21.4	21.5	17.5	17.6	18.7	17.2
West		18.2	18.3	19.0	19.5	19.5	19.3	22.6	17.0	19.0	22.4	21.2	19.5	16.5	16.7	15.2
Population Density:																
Large MSA		20.6	17.6	17.6	16.1	18.2	19.6	20.7	17.2	21.6	23.4	19.5	16.5	15.7	15.1	16.2
Other MSA		20.1	17.3	18.2	21.7	21.8	21.9	21.8	21.2	22.7	22.5	21.6	18.0	18.6	20.5	18.3
Non-MSA		21.1	19.9	24.7	21.8	21.8	22.4	25.5	25.4	23.4	25.4	25.3	21.4	21.4	19.0	17.8
Parental Education: ^a																
1.0–2.0 (Low)		20.9	18.2	22.2	20.0	23.4	22.1	19.7	20.1	21.8	21.3	22.0	17.3	19.0	17.7	15.8
2.5–3.0		22.5	18.5	21.4	21.2	22.9	23.4	22.5	23.3	23.4	24.5	22.6	19.9	18.9	21.0	19.4
3.5–4.0		20.4	19.4	19.4	22.1	21.4	22.1	24.1	21.3	23.3	24.5	22.8	19.5	19.3	18.9	17.5
4.5–5.0		19.7	17.1	18.2	18.7	19.7	19.5	22.3	20.2	21.7	24.1	21.4	17.4	17.5	17.3	17.5
5.5–6.0 (High)		20.6	18.5	18.6	17.9	17.9	22.3	22.4	20.4	24.0	23.1	21.5	16.8	17.7	19.3	17.0
Race/Ethnicity (2-year average): ^b																
White		—	21.6	20.8	22.0	22.7	23.7	25.0	25.5	25.7	26.7	25.8	23.2	21.9	21.7	21.0
African American		—	9.4	10.3	10.1	9.8	8.5	8.6	8.8	7.6	8.6	9.5	8.6	8.0	8.3	8.0
Hispanic		—	16.2	15.9	17.0	18.6	20.1	19.5	18.0	17.8	18.0	18.7	17.4	15.7	17.0	17.3

Table continued on next page.

TABLE 100 (cont.)
BEEN DRUNK: Trends in 30-Day Prevalence by Subgroups in Grade 10

	Percentage who had been drunk in last 30 days																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	18.8	18.1	14.4	15.5	14.7	13.7	14.5	12.8	11.2	10.3	9.0	8.9	8.4	8.8	9.3	5.4	-3.4* sss
Gender:																	
Male	19.1	18.9	14.6	16.4	15.5	14.9	15.4	13.2	11.1	10.3	8.4	7.6	8.2	8.6	§	4.5	-4.2 sss
Female	18.4	17.4	14.3	14.8	13.7	12.4	13.5	12.4	11.5	10.3	9.6	10.1	8.6	8.9	§	6.0	-2.8 ss
College Plans:																	
None or under 4 years	29.0	28.9	25.2	26.8	27.0	23.8	23.1	18.5	20.9	16.3	13.9	13.4	11.8	11.2	§	6.9	-4.3 ss
Complete 4 years	17.5	16.8	13.2	14.4	13.3	12.6	13.7	12.3	10.3	9.7	8.5	8.3	7.9	8.3	§	5.0	-3.4 sss
Region:																	
Northeast	20.4	20.5	15.3	14.6	14.7	16.6	18.6	13.6	13.3	11.6	10.0	9.9	9.1	9.2	§	6.2	-3.0
Midwest	19.8	19.6	16.0	15.0	15.3	11.9	12.8	11.4	9.7	9.5	8.4	8.1	8.2	8.1	§	6.4	-1.7
South	17.6	16.3	14.9	16.6	14.5	14.0	15.0	13.9	11.6	11.4	9.2	8.5	8.7	8.8	§	5.7	-3.1 s
West	17.9	16.9	11.8	15.5	14.3	13.0	12.1	12.4	10.7	7.9	8.4	9.6	7.6	9.0	§	3.5	-5.6 sss
Population Density:																	
Large MSA	17.7	16.3	12.7	15.3	14.2	12.9	14.9	12.8	11.3	10.0	9.3	11.1	9.4	7.8	§	3.8	-4.0 ss
Other MSA	19.0	18.5	14.8	15.3	14.8	14.1	13.7	13.0	10.8	9.9	7.9	7.4	8.1	9.4	§	5.9	-3.5 ss
Non-MSA	20.0	20.0	16.4	16.4	15.1	14.0	15.7	12.6	12.3	11.6	11.5	9.2	7.7	8.6	§	7.1	-1.5
Parental Education: ^a																	
1.0–2.0 (Low)	18.4	17.7	16.9	19.7	18.2	13.6	15.7	15.1	12.0	8.5	8.5	8.4	8.5	8.1	§	7.2	-1.0
2.5–3.0	21.6	19.9	15.0	17.6	16.4	15.8	14.9	13.6	11.5	11.4	8.9	8.0	8.1	9.5	§	5.4	-4.1 ss
3.5–4.0	19.9	19.1	15.7	16.4	14.3	14.8	14.7	13.4	12.9	11.3	9.4	9.3	8.4	10.1	§	4.7	-5.4 sss
4.5–5.0	17.4	16.8	13.3	13.8	13.9	12.1	15.0	12.1	10.4	10.0	8.7	8.9	7.8	7.8	§	5.7	-2.0
5.5–6.0 (High)	17.6	17.7	14.2	14.6	13.2	13.1	14.6	12.3	11.3	9.8	10.8	10.7	10.9	9.6	§	6.2	-3.4 s
Race/Ethnicity (2-year average): ^b																	
White	20.9	21.3	19.2	17.0	16.4	15.6	16.0	15.4	14.1	12.8	11.7	11.3	11.0	10.5	§	9.1	-1.4
African American	7.8	8.3	8.0	8.4	9.3	8.3	7.6	7.2	5.5	5.3	4.8	3.7	3.1	3.7	§	3.9	+0.2
Hispanic	16.6	15.0	14.0	14.9	15.7	13.8	12.5	13.1	11.6	10.0	8.5	7.8	7.7	8.0	§	5.4	-2.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 101
BEEN DRUNK: Trends in 30-Day Prevalence by Subgroups in Grade 12

Percentage who had been drunk in last 30 days																	
	1975–	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	—	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	
Total	—	31.6	29.9	28.9	30.8	33.2	31.3	34.2	32.9	32.9	32.3	32.7	30.3	30.9	32.5	30.2	
Gender:																	
Male	—	37.1	35.2	34.5	34.5	37.8	35.4	39.2	39.0	37.9	38.4	37.0	34.3	34.9	36.0	33.6	
Female	—	25.4	24.5	23.5	26.8	28.8	27.3	29.1	26.6	27.7	26.7	28.4	26.9	26.9	29.0	26.4	
College Plans:																	
None or under 4 years	—	32.2	31.4	32.6	32.2	37.6	31.4	38.1	33.7	36.1	35.0	36.3	31.7	37.6	34.7	34.3	
Complete 4 years	—	30.9	29.2	27.4	29.4	31.4	31.0	32.3	32.0	31.7	30.6	31.3	29.3	28.7	31.2	28.5	
Region:																	
Northeast	—	36.4	30.0	35.0	35.2	35.5	37.2	35.9	35.6	37.5	39.3	33.9	33.6	35.4	38.1	38.7	
Midwest	—	37.2	38.2	32.5	34.1	38.2	31.5	33.7	34.8	33.4	34.8	39.2	35.0	34.9	34.2	31.3	
South	—	26.5	25.2	26.4	29.1	31.2	31.0	34.5	30.1	30.8	26.5	28.8	28.4	25.9	29.3	26.9	
West	—	28.5	26.6	23.2	25.4	27.1	24.7	32.7	33.5	32.2	32.8	28.8	25.0	30.1	29.5	26.5	
Population Density:																	
Large MSA	—	30.4	26.1	29.4	28.7	32.0	31.5	31.5	32.2	29.2	30.5	30.6	32.9	25.4	27.4	28.2	
Other MSA	—	33.5	29.8	26.9	29.9	31.7	33.0	33.7	34.0	35.4	34.5	31.7	29.1	33.5	37.3	31.5	
Non-MSA	—	29.4	33.7	32.0	34.4	36.9	28.2	38.2	31.4	32.5	30.5	36.7	29.2	33.2	29.1	29.9	
Parental Education: ^a																	
1.0–2.0 (Low)	—	20.4	20.5	23.6	25.7	25.4	18.8	23.7	24.6	20.8	28.4	22.2	20.0	25.6	25.5	22.0	
2.5–3.0	—	30.2	30.0	26.4	30.3	30.0	27.4	31.5	28.0	30.5	33.1	32.2	27.0	28.7	32.3	29.9	
3.5–4.0	—	31.0	31.3	29.2	29.9	34.4	31.1	32.7	34.1	34.0	31.2	32.0	32.4	30.6	35.1	31.3	
4.5–5.0	—	34.4	29.4	32.8	33.5	36.5	35.8	37.7	36.0	32.8	31.9	36.1	31.0	33.3	33.7	31.3	
5.5–6.0 (High)	—	40.5	34.3	30.4	30.7	34.9	34.6	39.8	39.9	40.6	35.5	33.6	34.5	33.4	32.6	32.2	
Race/Ethnicity (2-year average): ^b																	
White	—	—	34.7	33.6	34.0	36.4	36.6	37.7	39.3	37.8	37.0	37.7	36.6	35.6	36.4	36.5	
African American	—	—	11.0	12.5	14.1	13.2	13.0	13.8	13.8	14.9	14.9	12.0	12.1	11.7	14.3	15.4	
Hispanic	—	—	27.2	24.8	23.0	24.2	26.2	26.9	25.9	27.5	29.8	25.5	23.5	23.9	24.1	22.2	

Table continued on next page.

TABLE 101 (cont.)
BEEN DRUNK: Trends in 30-Day Prevalence by Subgroups in Grade 12

	Percentage who had been drunk in last 30 days																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>																	
	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	30.0	28.7	27.6	27.4	26.8	25.0	28.1	26.0	23.5	20.6	20.4	19.1	17.5	17.5	19.8	15.5	-2.0*
Gender:																	
Male	32.9	31.7	29.2	32.7	31.2	27.5	30.6	29.1	24.9	21.1	22.1	20.4	19.0	17.2	§	15.6	-1.6
Female	26.3	25.7	26.2	22.5	21.8	22.0	24.8	22.7	22.4	19.6	19.1	18.1	16.2	17.4	§	15.8	-1.6
College Plans:																	
None or under 4 years	37.2	30.6	32.0	31.0	32.9	27.9	29.6	28.9	25.3	21.2	23.8	21.7	17.1	19.6	§	13.9	-5.7
Complete 4 years	27.6	28.2	26.4	25.9	24.8	24.1	27.3	25.0	22.6	20.3	19.4	18.6	17.6	16.4	§	15.5	-0.9
Region:																	
Northeast	37.9	32.6	31.1	36.3	32.4	31.5	34.7	30.1	28.6	25.1	21.8	25.3	20.7	17.9	§	26.1	+8.2
Midwest	32.2	32.0	31.2	29.4	28.9	27.5	29.6	26.8	25.9	20.3	22.8	17.7	18.0	18.9	§	17.3	-1.6
South	27.4	26.5	25.5	23.6	24.9	21.5	22.2	21.7	21.0	18.4	17.6	17.8	15.4	16.2	§	9.7	-6.4 ss
West	25.2	26.0	24.1	23.5	23.4	23.3	29.3	28.3	21.4	21.3	22.1	18.4	18.3	18.4	§	14.3	-4.1
Population Density:																	
Large MSA	30.4	29.8	31.4	29.8	26.6	23.8	30.7	25.6	22.5	22.9	21.1	19.7	18.4	16.7	§	22.3	+5.6
Other MSA	29.0	27.6	25.0	25.9	26.8	26.9	28.0	28.1	25.2	18.1	20.4	17.3	16.4	17.3	§	10.8	-6.5 s
Non-MSA	31.4	29.4	28.8	27.8	27.2	22.3	24.1	21.1	20.7	22.6	19.4	21.9	18.6	19.1	§	14.9	-4.2
Parental Education: ^a																	
1.0–2.0 (Low)	27.7	19.7	26.7	19.8	22.4	19.1	19.7	25.7	18.7	14.0	13.9	15.8	9.5	12.7	§	7.5	-5.2
2.5–3.0	29.5	31.6	25.3	29.0	25.4	22.9	26.6	24.8	25.2	21.4	19.3	15.7	16.0	15.6	§	11.6	-4.0
3.5–4.0	31.0	26.8	28.2	28.2	28.6	26.3	29.5	25.6	21.7	22.4	21.8	20.0	16.8	17.1	§	11.2	-5.9 s
4.5–5.0	29.7	28.4	30.0	27.0	26.5	25.6	29.7	27.9	25.9	21.8	21.5	19.7	19.1	19.7	§	18.8	-0.9
5.5–6.0 (High)	30.1	33.0	27.8	29.0	26.5	27.5	30.5	27.1	25.8	21.1	23.4	24.1	27.5	21.2	§	27.9	+6.7
Race/Ethnicity (2-year average): ^b																	
White	35.2	33.7	32.7	31.8	31.6	29.9	29.7	30.6	29.0	26.4	25.0	24.2	23.1	21.9	§	23.4	+1.5
African American	13.5	14.6	14.8	15.1	14.7	14.2	14.3	13.4	15.1	13.8	9.9	10.1	10.9	10.5	§	7.0	-3.5
Hispanic	23.1	24.0	22.2	21.7	20.5	20.0	21.2	22.7	19.5	16.0	17.2	16.8	13.1	11.6	§	9.1	-2.5

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms; *N* is two sixths of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 102
BINGE DRINKING: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 8

	Percentage reporting 5+ drinks in a row on one or more occasions														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>
Total	10.9	11.3	11.3	12.1	12.3	13.3	12.3	11.5	13.1	11.7	11.0	10.3	9.8	9.4	8.4
Gender:															
Male	12.1	11.6	12.4	13.2	12.5	14.1	13.0	12.1	14.0	11.7	11.5	10.4	10.2	8.8	8.2
Female	9.6	10.8	10.3	11.1	12.1	12.4	11.5	10.8	12.0	11.3	10.2	10.0	9.4	9.9	8.6
College Plans:															
None or under 4 years	21.1	23.0	25.7	25.3	25.0	26.1	26.5	26.7	30.2	25.0	24.7	24.7	23.2	21.2	21.0
Complete 4 years	9.3	9.7	9.4	10.5	10.8	11.3	10.7	9.8	11.2	10.1	9.4	9.0	8.4	8.2	7.2
Region:															
Northeast	8.6	8.8	8.1	10.9	11.1	12.8	10.9	9.7	12.4	11.5	10.5	8.4	7.8	8.1	6.0
Midwest	11.5	12.5	10.9	11.5	12.2	13.5	12.0	12.4	15.5	13.2	10.8	10.5	10.8	10.1	8.7
South	11.9	12.3	12.9	12.4	13.0	13.7	13.2	11.9	13.3	11.0	12.5	11.4	10.6	9.6	9.8
West	10.2	10.4	12.8	13.8	12.3	12.9	12.3	11.6	10.3	10.9	9.0	9.9	9.0	9.5	8.0
Population Density:															
Large MSA	10.6	10.4	8.6	10.4	10.3	12.2	11.3	10.1	11.0	10.6	9.2	8.3	8.8	8.7	8.0
Other MSA	10.5	11.7	12.3	13.1	12.2	13.7	12.1	10.9	12.1	10.5	11.0	10.8	8.9	9.3	8.3
Non-MSA	12.0	11.7	12.9	12.1	15.0	14.2	13.9	14.3	17.4	15.3	13.1	11.8	12.9	10.5	9.2
Parental Education: ^a															
1.0–2.0 (Low)	18.3	17.8	16.3	16.8	20.1	16.9	17.6	16.5	20.8	18.9	16.5	16.6	16.3	14.9	14.4
2.5–3.0	12.6	13.4	13.3	14.2	14.8	15.4	13.5	13.8	17.0	15.3	14.3	12.8	13.5	12.1	10.7
3.5–4.0	10.7	11.3	11.5	12.5	13.3	14.3	14.2	12.4	14.0	11.0	11.7	11.1	10.5	10.0	9.4
4.5–5.0	8.8	8.7	8.5	10.3	9.3	11.1	10.3	9.3	9.5	8.2	8.1	7.9	6.7	8.0	6.4
5.5–6.0 (High)	8.8	8.4	8.9	9.1	8.7	10.7	9.0	9.4	9.2	9.5	6.6	7.3	6.8	6.2	5.0
Race/Ethnicity (2-year average): ^b															
White	—	11.2	10.9	11.1	12.1	13.2	13.3	12.3	12.6	13.0	11.9	10.9	10.0	9.6	9.0
African American	—	6.5	7.6	8.6	8.3	8.0	7.5	6.9	7.7	7.3	6.4	7.1	7.5	6.9	6.1
Hispanic	—	16.5	17.7	18.9	18.4	17.5	17.4	16.9	17.4	16.0	14.4	14.4	13.8	13.2	12.1

Table continued on next page.

TABLE 102 (cont.)
BINGE DRINKING: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 8

	Percentage reporting 5+ drinks in a row on one or more occasions																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	8.7	8.3	8.1	7.8	7.2	6.4	5.1	5.1	4.1	4.6	3.4	3.7	3.7	3.8	4.5	2.8	-1.1* ss
Gender:																	
Male	8.6	8.2	8.1	7.8	6.5	6.1	4.6	4.5	3.5	4.6	3.2	3.1	3.3	3.5	§	2.0	-1.5 sss
Female	8.5	8.2	8.0	7.7	7.8	6.5	5.5	5.7	4.6	4.6	3.6	4.2	3.9	4.0	§	3.2	-0.8
College Plans:																	
None or under 4 years	19.3	21.5	21.3	19.1	17.6	15.6	15.3	14.2	9.9	10.5	7.7	8.3	8.9	8.2	§	4.7	-3.6 ss
Complete 4 years	7.6	7.1	7.0	6.8	6.3	5.7	4.4	4.5	3.7	4.1	3.0	3.2	2.9	3.1	§	2.3	-0.9 s
Region:																	
Northeast	7.8	5.6	7.1	8.2	6.2	3.7	4.3	4.2	3.3	3.6	2.5	2.8	2.9	2.0	§	1.2	-0.8
Midwest	8.8	7.5	8.2	7.0	6.5	6.1	2.6	4.1	3.4	3.8	2.7	3.1	2.9	3.6	§	3.6	+0.1
South	9.7	9.4	9.7	8.8	8.4	7.1	6.3	6.4	4.7	5.4	4.3	3.9	3.6	4.8	§	2.7	-2.1 ss
West	7.6	8.9	6.3	6.6	6.6	8.0	6.1	4.8	4.5	4.8	3.2	4.7	5.1	3.9	§	3.1	-0.9
Population Density:																	
Large MSA	8.1	7.4	8.1	6.4	6.5	5.8	4.5	4.9	4.0	4.3	2.7	3.5	3.7	3.1	§	1.6	-1.5 ss
Other MSA	8.3	8.5	7.9	8.2	7.6	6.6	5.6	5.0	4.0	4.6	3.4	3.5	3.7	3.9	§	2.9	-0.9
Non-MSA	10.2	9.2	8.4	8.9	7.0	7.0	5.1	5.8	4.5	5.3	4.6	4.4	3.4	5.1	§	4.4	-0.7
Parental Education: ^a																	
1.0–2.0 (Low)	13.3	15.5	17.0	15.2	12.0	11.0	9.5	9.5	5.7	7.5	6.2	6.2	5.9	5.6	§	2.8	-2.9 s
2.5–3.0	10.6	10.5	10.5	9.5	9.3	8.8	7.6	6.7	5.4	5.8	4.6	5.1	4.7	6.5	§	3.2	-3.3 ss
3.5–4.0	9.9	9.2	7.9	8.7	8.0	7.2	5.3	6.4	4.9	4.9	2.8	3.7	3.7	4.2	§	4.2	-0.1
4.5–5.0	6.7	5.5	5.7	5.3	5.7	4.6	3.8	3.0	3.4	3.1	3.1	2.9	2.6	3.0	§	2.3	-0.7
5.5–6.0 (High)	5.9	5.0	4.7	4.7	4.3	4.1	2.0	3.9	3.0	3.6	2.3	2.5	2.5	2.1	§	2.3	+0.3
Race/Ethnicity (2-year average): ^b																	
White	8.4	8.0	7.8	7.7	7.1	6.2	4.9	4.2	4.2	4.0	3.6	3.0	2.9	3.4	§	3.6	+0.2
African American	5.7	5.6	5.7	5.2	5.3	5.1	4.3	4.5	4.4	4.1	3.4	2.9	2.6	1.9	§	3.4	+1.5
Hispanic	11.6	12.5	12.3	11.5	10.8	10.4	9.9	7.8	5.7	5.4	5.3	4.9	5.2	5.3	§	2.4	-2.9 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 103
BINGE DRINKING: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 10

	Percentage reporting 5+ drinks in a row on one or more occasions															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	
<i>Approximate weighted N =</i>	<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>	
Total	21.0	19.1	21.0	21.9	22.0	22.8	23.1	22.4	23.5	24.1	22.8	20.3	20.0	19.9	19.0	
Gender:																
Male	24.1	21.5	24.1	26.4	24.1	25.1	26.3	24.7	27.4	27.6	26.1	21.6	21.1	21.6	19.9	
Female	18.1	17.0	17.7	17.3	19.7	20.4	20.1	20.4	20.0	20.6	19.7	19.0	19.0	18.1	17.9	
College Plans:																
None or under 4 years	30.5	28.6	31.8	34.1	34.5	35.7	36.9	35.7	36.5	36.7	37.2	31.0	31.5	29.6	31.3	
Complete 4 years	19.1	17.2	18.8	19.3	19.8	20.7	20.8	20.2	21.5	22.2	20.5	18.5	18.2	18.6	17.3	
Region:																
Northeast	23.7	18.0	21.1	19.9	20.3	22.3	21.8	24.1	26.0	23.8	20.7	19.5	21.5	23.8	21.0	
Midwest	21.7	19.8	22.0	23.0	23.3	23.4	22.4	22.2	25.0	26.2	24.8	20.3	19.9	16.7	18.8	
South	20.8	18.9	20.4	22.9	22.4	23.5	23.4	23.1	22.1	22.5	22.1	19.7	19.1	20.1	18.6	
West	18.6	19.8	20.0	20.4	21.1	21.4	25.3	19.9	21.7	24.6	23.1	22.2	20.1	19.6	17.6	
Population Density:																
Large MSA	20.2	17.5	19.1	17.6	18.4	20.5	20.4	18.3	22.4	24.0	20.8	17.6	18.0	16.9	17.4	
Other MSA	20.4	18.1	19.2	22.7	22.1	23.0	22.9	22.0	22.7	23.4	21.7	20.4	19.9	21.6	19.4	
Non-MSA	22.9	22.8	25.8	24.6	25.9	25.3	26.6	28.0	26.7	25.7	27.2	23.8	23.4	20.1	20.3	
Parental Education: ^a																
1.0–2.0 (Low)	23.2	23.0	24.3	22.6	26.7	24.7	22.7	24.1	25.2	24.7	23.0	23.9	24.4	23.0	21.0	
2.5–3.0	23.8	20.0	23.4	23.4	24.7	25.5	23.5	25.5	26.3	27.0	24.6	22.5	21.8	21.9	21.3	
3.5–4.0	19.9	19.2	20.6	23.3	22.8	23.7	24.6	22.8	23.6	23.8	24.2	20.7	20.5	20.4	18.9	
4.5–5.0	19.3	18.1	18.6	20.2	20.2	20.3	21.6	20.1	21.3	23.3	21.5	17.7	17.8	18.2	17.9	
5.5–6.0 (High)	20.9	18.4	18.9	18.5	17.3	22.4	22.4	20.1	23.6	23.1	21.2	18.7	18.4	19.0	17.9	
Race/Ethnicity (2-year average): ^b																
White	—	21.6	21.4	22.9	23.8	24.5	25.2	25.3	25.4	26.2	25.6	23.6	22.5	22.2	21.8	
African American	—	12.3	11.9	11.8	11.1	9.4	10.0	10.6	10.7	10.8	10.3	10.0	9.8	9.7	9.1	
Hispanic	—	20.3	20.5	20.9	23.3	25.7	24.4	23.6	24.4	25.1	24.8	23.3	22.9	23.3	22.4	

Table continued on next page.

TABLE 103 (cont.)
BINGE DRINKING: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 10

	Percentage reporting 5+ drinks in a row on one or more occasions																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	19.9	19.6	16.0	17.5	16.3	14.7	15.6	13.7	12.6	10.9	9.7	9.8	8.7	8.5	9.6	5.9	-2.6* sss
Gender:																	
Male	21.0	20.9	16.6	18.8	17.9	16.5	16.4	14.7	13.1	11.3	9.6	9.0	9.1	9.0	§	5.4	-3.7 sss
Female	18.9	18.3	15.4	16.1	14.6	12.7	14.8	12.5	12.2	10.6	9.8	10.5	8.3	7.9	§	5.9	-2.0 ss
College Plans:																	
None or under 4 years	30.8	30.7	29.1	29.9	30.5	27.1	25.9	21.7	24.3	18.3	16.4	15.0	14.0	12.1	§	9.2	-2.8
Complete 4 years	18.6	18.2	14.4	16.1	14.6	13.4	14.6	12.8	11.4	10.1	9.0	9.1	7.9	7.7	§	4.9	-2.8 sss
Region:																	
Northeast	21.2	20.4	16.8	16.4	15.2	16.8	19.5	14.5	13.8	11.9	9.4	9.8	9.0	9.0	§	6.5	-2.5
Midwest	20.0	20.7	16.5	17.2	16.2	13.0	13.8	12.3	11.4	9.7	9.2	8.5	7.6	8.2	§	6.7	-1.6
South	19.1	18.4	16.3	18.4	16.7	15.4	16.2	15.0	13.3	11.9	10.4	9.7	9.6	8.7	§	6.5	-2.3
West	20.1	19.5	14.4	17.4	16.9	14.1	13.2	12.8	11.9	9.6	9.6	11.3	8.1	8.1	§	3.8	-4.3 ss
Population Density:																	
Large MSA	19.2	17.9	14.5	16.7	16.0	14.3	15.9	13.7	12.3	11.0	10.2	12.2	9.4	7.2	§	4.3	-3.0 ss
Other MSA	19.7	19.6	16.0	17.0	16.5	14.7	14.8	13.7	12.4	10.3	8.7	8.2	8.1	8.9	§	6.2	-2.7 s
Non-MSA	21.9	22.5	18.1	19.8	16.7	15.4	16.6	13.7	13.8	12.4	11.8	10.0	8.7	9.6	§	7.9	-1.7
Parental Education: ^a																	
1.0–2.0 (Low)	19.8	22.0	20.6	25.7	22.8	17.6	20.5	17.8	15.4	13.9	10.5	11.7	11.2	8.9	§	9.8	+0.9
2.5–3.0	22.8	22.6	17.8	19.6	19.5	17.1	17.3	15.1	14.0	11.9	10.8	9.4	9.5	9.4	§	6.4	-3.0 s
3.5–4.0	21.2	20.7	16.6	18.4	16.3	15.8	15.0	13.6	13.7	12.1	10.0	10.9	8.5	9.5	§	5.2	-4.2 sss
4.5–5.0	18.1	17.7	14.7	14.7	14.1	12.7	15.4	12.5	10.7	10.1	9.2	9.2	7.2	7.0	§	5.5	-1.5
5.5–6.0 (High)	19.0	17.2	12.8	14.8	13.1	12.3	14.7	12.1	11.6	9.6	10.0	9.9	9.4	8.4	§	5.6	-2.8 ss
Race/Ethnicity (2-year average): ^b																	
White	21.7	21.8	19.7	17.9	17.2	16.1	16.3	15.7	14.4	13.0	11.6	11.0	10.5	9.7	§	9.0	-0.7
African American	9.1	10.0	9.8	9.8	10.7	9.4	8.2	8.6	7.5	6.9	5.8	4.7	3.9	4.2	§	4.8	+0.6
Hispanic	21.2	20.1	19.6	20.6	22.2	19.7	17.1	16.9	15.0	13.2	11.4	11.3	10.8	9.3	§	7.6	-1.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 104
BINGE DRINKING: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	36.8	37.1	39.4	40.3	41.2	41.2	41.4	40.5	40.8	38.7	36.7	36.8	37.5	34.7	33.0	32.2
Gender:																
Male	49.0	47.9	50.0	51.4	51.9	52.1	51.6	49.8	50.4	47.5	45.3	46.1	46.1	43.0	41.2	39.1
Female	26.4	25.9	29.3	29.6	30.9	30.5	30.8	31.1	31.0	29.6	28.2	28.1	29.2	26.5	24.9	24.4
College Plans:																
None or under 4 years	—	41.8	44.7	44.3	44.5	46.3	46.7	45.7	44.9	43.5	41.6	41.3	42.7	38.5	38.2	35.8
Complete 4 years	—	31.5	33.9	35.9	37.7	36.9	37.4	36.5	37.2	34.6	33.0	34.1	35.0	32.8	30.5	30.3
Region:																
Northeast	43.0	40.8	40.0	43.5	47.4	48.0	49.3	43.3	42.2	42.9	42.4	37.1	37.2	34.3	33.3	37.2
Midwest	40.6	42.8	44.5	45.3	44.8	45.4	44.9	47.9	47.2	44.3	39.7	42.6	43.5	39.9	40.4	37.9
South	32.1	30.8	36.3	36.4	36.7	34.4	34.7	34.6	37.6	33.5	29.7	31.7	33.4	30.4	28.5	27.2
West	29.0	32.8	34.2	33.3	34.0	36.0	35.6	32.5	33.3	34.5	36.1	35.9	36.6	35.4	30.8	26.3
Population Density:																
Large MSA	37.9	37.0	38.1	39.5	42.2	44.8	43.4	40.9	38.8	37.9	37.6	36.4	34.8	32.5	28.8	34.5
Other MSA	36.1	36.8	39.5	40.1	40.8	38.9	39.5	39.7	41.0	37.3	35.4	35.5	38.6	35.3	33.7	31.8
Non-MSA	36.9	38.0	40.5	41.3	40.9	41.4	42.2	41.3	42.0	41.2	37.6	39.1	38.3	35.9	35.8	30.6
Parental Education: ^a																
1.0–2.0 (Low)	31.6	34.1	35.6	36.3	36.0	37.0	37.0	35.3	37.2	34.8	31.8	31.7	33.9	30.7	25.4	25.3
2.5–3.0	37.5	41.1	41.8	40.9	42.3	43.3	43.2	41.4	41.2	39.8	38.2	37.9	38.9	35.7	34.0	32.7
3.5–4.0	35.1	36.4	39.5	41.3	41.4	42.1	42.4	42.4	40.9	39.3	36.9	37.9	38.3	34.7	34.3	32.0
4.5–5.0	34.4	36.9	37.2	42.4	43.8	40.8	40.8	41.9	41.9	38.6	37.1	37.1	37.2	35.1	34.2	34.5
5.5–6.0 (High)	29.9	34.5	41.1	37.2	41.9	38.5	39.3	40.9	42.1	38.2	34.9	36.7	37.2	34.7	31.8	34.1
Race/Ethnicity (2-year average): ^b																
White	—	—	40.5	42.4	43.5	44.3	44.9	44.9	44.5	43.6	41.5	40.3	40.9	40.0	37.9	36.6
African American	—	—	19.0	19.3	18.9	17.7	17.1	17.1	18.3	17.2	15.7	16.4	15.8	15.2	15.7	14.4
Hispanic	—	—	36.4	37.2	33.6	33.1	34.8	32.9	32.5	33.0	31.7	30.8	33.0	33.7	28.8	25.6

Table continued on next page.

TABLE 104 (cont.)
BINGE DRINKING: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	29.8	27.9	27.5	28.2	29.8	30.2	31.3	31.5	30.8	30.0	29.7	28.6	27.9	29.2	27.1	25.4
Gender:																
Male	37.8	35.6	34.6	37.0	36.9	37.0	37.9	39.2	38.1	36.7	36.0	34.2	34.2	34.3	32.6	28.9
Female	21.2	20.3	20.7	20.2	23.0	23.5	24.4	24.0	23.6	23.5	23.7	23.0	22.1	24.2	21.6	21.5
College Plans:																
None or under 4 years	34.4	32.8	32.7	34.0	35.2	33.9	36.2	36.3	35.4	35.7	35.9	34.0	34.5	34.6	34.3	32.3
Complete 4 years	27.9	26.0	25.8	26.3	27.8	28.8	29.5	30.0	29.5	27.6	27.8	27.2	25.9	27.6	25.1	23.7
Region:																
Northeast	33.4	25.8	30.3	29.2	31.2	33.7	33.5	33.5	33.7	35.1	31.8	30.7	30.9	33.2	32.2	30.9
Midwest	34.6	34.6	30.1	31.9	34.3	31.5	31.6	32.6	31.6	33.9	34.5	33.4	32.5	30.2	29.1	26.5
South	26.3	24.7	27.1	26.9	28.6	30.2	30.6	30.7	30.0	24.9	26.6	26.8	23.7	26.1	24.2	23.9
West	26.3	26.0	22.0	24.5	24.2	24.0	29.6	29.5	28.6	29.3	26.0	23.9	26.5	28.6	24.5	22.0
Population Density:																
Large MSA	28.6	25.5	27.6	26.8	28.3	29.8	29.9	27.9	27.1	29.7	29.0	29.9	23.3	25.5	26.5	26.6
Other MSA	30.1	27.0	26.5	27.1	28.4	30.3	31.1	33.1	32.2	29.3	29.0	28.0	29.8	31.9	27.4	25.0
Non-MSA	30.4	31.9	29.2	31.5	34.0	30.5	33.2	32.4	32.4	31.7	31.9	28.2	30.5	28.2	27.0	24.5
Parental Education: ^a																
1.0–2.0 (Low)	26.8	23.4	21.9	24.0	26.6	24.8	22.6	26.3	27.4	25.1	24.8	22.2	23.5	22.9	19.4	25.4
2.5–3.0	29.9	28.1	27.6	28.5	31.2	28.6	30.4	30.3	30.8	29.2	30.2	28.5	27.6	28.8	28.3	25.8
3.5–4.0	30.4	27.9	28.4	28.4	29.5	29.8	31.0	33.2	30.5	30.5	30.3	28.6	28.4	30.9	27.6	26.5
4.5–5.0	29.9	28.1	28.4	29.3	29.9	32.4	32.4	32.3	30.6	29.9	30.9	30.9	28.6	30.5	29.4	24.0
5.5–6.0 (High)	30.6	30.4	29.0	29.0	30.7	33.1	34.9	32.4	34.2	32.7	28.6	30.6	28.7	27.8	25.2	26.3
Race/Ethnicity (2-year average): ^b																
White	34.6	32.1	31.3	31.5	32.3	33.4	35.1	36.4	35.7	34.6	34.5	33.7	32.4	32.5	32.5	30.4
African American	11.7	11.3	12.6	14.4	14.9	15.3	13.4	12.3	12.3	11.5	11.8	11.5	10.8	11.4	11.3	11.4
Hispanic	27.9	31.1	27.2	24.3	26.6	27.1	27.6	28.1	29.3	31.0	28.4	26.4	25.9	26.0	23.9	23.3

Table continued on next page.

TABLE 104 (cont.)
BINGE DRINKING: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions														2019-2021	
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	25.9	24.6	25.2	23.2	21.6	23.7	22.1	19.4	17.2	15.5	16.6	13.8	14.4	16.8	11.8	-2.6*
Gender:																
Male	30.7	28.4	30.5	28.0	25.5	27.2	26.1	22.3	19.3	17.2	18.5	15.9	16.1	§	11.1	-5.0 ss
Female	21.5	21.3	20.2	18.4	17.6	19.7	18.1	16.6	14.9	13.5	14.6	11.9	12.4	§	12.6	+0.2
College Plans:																
None or under 4 years	30.8	31.5	31.3	32.1	27.4	27.5	29.2	24.0	21.5	18.2	20.7	16.5	16.7	§	11.3	-5.3 ss
Complete 4 years	24.7	23.2	23.8	21.2	20.4	22.6	20.6	18.3	16.4	14.7	15.8	12.9	13.5	§	11.8	-1.7
Region:																
Northeast	29.7	27.3	28.9	28.7	26.1	29.0	26.6	22.3	21.3	18.0	18.7	15.2	13.6	§	16.4	+2.8
Midwest	28.9	25.2	28.8	24.6	24.3	24.8	22.7	20.4	16.7	16.0	15.7	13.5	13.5	§	12.7	-0.8
South	25.4	24.7	23.1	21.0	18.8	19.6	18.6	17.9	16.4	14.6	16.9	14.0	15.0	§	9.6	-5.3 ss
West	20.5	21.9	20.9	20.7	19.9	24.1	23.1	18.8	16.2	14.9	15.6	12.7	15.0	§	10.9	-4.1
Population Density:																
Large MSA	28.0	25.8	25.7	23.0	21.7	25.1	21.3	19.4	18.9	15.6	16.5	13.7	12.8	§	14.3	+1.5
Other MSA	24.2	22.5	24.1	23.6	22.5	23.3	23.3	20.1	15.3	15.2	14.7	12.4	14.2	§	9.7	-4.5 s
Non-MSA	26.4	28.1	27.2	22.4	19.6	22.4	20.5	17.9	19.3	16.2	21.2	17.1	17.6	§	12.4	-5.2 s
Parental Education: ^a																
1.0–2.0 (Low)	20.4	21.3	23.2	20.1	17.9	23.6	21.5	19.0	17.2	14.6	13.0	10.8	11.7	§	8.4	-3.3
2.5–3.0	27.3	23.0	24.6	24.6	21.1	23.9	23.1	20.1	17.1	13.9	15.5	12.7	14.8	§	10.2	-4.7 s
3.5–4.0	24.8	25.1	27.6	23.3	23.2	24.3	23.9	19.2	17.4	15.9	16.4	13.8	13.8	§	12.0	-1.8
4.5–5.0	27.5	26.2	24.4	24.2	21.4	23.6	21.4	20.7	17.9	16.5	18.2	15.0	14.5	§	13.9	-0.6
5.5–6.0 (High)	27.1	26.4	25.6	21.2	22.1	22.8	20.3	19.0	17.6	15.9	20.1	17.2	17.5	§	14.5	-3.0
Race/Ethnicity (2-year average): ^b																
White	29.7	29.9	29.0	27.6	25.9	25.7	25.6	23.8	21.2	19.1	19.6	19.4	17.6	§	17.2	-0.4
African American	11.5	10.9	12.0	13.1	11.3	11.3	12.5	11.3	9.8	8.3	7.7	7.4	6.7	§	5.8	-0.9
Hispanic	22.5	21.5	22.6	22.1	20.8	21.8	22.4	20.4	18.5	16.7	14.4	11.7	10.8	§	9.8	-1.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 105
EXTREME BINGE DRINKING:
Trends in 2-Week Prevalence of Having 10 or More Drinks in a Row
by Subgroups in Grades 8 and 10

	8th Graders								10th Graders							
	1991– 2015	2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
<i>Approximate weighted N =</i>	—	16,900	15,300	14,000	13,600	3,100	10,700		—	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	1.2	1.1	1.1	1.7	0.9	1.0	-0.7*	—	3.0	3.6	3.3	3.3	2.5	2.1	-1.2* s
Gender:																
Male	—	1.2	1.0	1.1	1.0	§	0.9	0.0	—	3.5	4.1	3.5	3.8	§	1.5	-2.3 sss
Female	—	0.9	1.0	1.0	2.4	§	1.1	-1.3	—	2.5	2.8	2.9	2.8	§	2.4	-0.4
College Plans:																
None or under 4 years	—	3.4	2.7	3.8	3.2	§	1.8	-1.4	—	6.6	7.2	8.2	6.9	§	3.9	-3.0
Complete 4 years	—	0.9	0.8	0.7	1.5	§	0.9	-0.6	—	2.6	2.8	2.7	2.7	§	1.6	-1.0 s
Region:																
Northeast	—	0.7	1.2	0.5	0.7	§	0.5	-0.3	—	3.2	2.7	3.0	2.3	§	1.7	-0.7
Midwest	—	1.0	0.7	1.0	1.2	§	0.8	-0.4	—	3.5	3.5	2.9	3.3	§	2.7	-0.6
South	—	1.4	0.9	0.6	2.2	§	0.9	-1.2	—	2.8	3.5	4.0	3.6	§	2.4	-1.2
West	—	1.2	1.7	2.7	2.0	§	1.8	-0.3	—	2.7	4.6	2.7	3.5	§	1.3	-2.1
Population Density:																
Large MSA	—	0.7	1.2	0.7	0.8	§	0.6	-0.2	—	3.3	4.8	2.9	2.5	§	0.9	-1.5 s
Other MSA	—	1.1	1.1	1.2	1.8	§	1.1	-0.7	—	2.3	2.2	3.4	3.3	§	2.5	-0.8
Non-MSA	—	2.0	1.0	1.8	2.8	§	1.7	-1.2	—	4.5	5.2	3.5	4.6	§	2.9	-1.7
Parental Education: ^a																
1.0–2.0 (Low)	—	2.8	2.1	1.8	2.6	§	0.6	-2.0	—	4.9	6.1	4.4	4.9	§	3.8	-1.1
2.5–3.0	—	1.2	1.5	1.8	3.3	§	2.8	-0.5	—	2.7	3.2	2.9	4.7	§	1.9	-2.8 s
3.5–4.0	—	1.0	0.7	0.6	2.2	§	1.4	-0.9	—	2.5	3.1	3.6	4.8	§	2.1	-2.6 s
4.5–5.0	—	0.8	0.9	0.5	0.7	§	0.7	0.0	—	2.9	2.9	2.7	1.3	§	1.6	+0.3
5.5–6.0 (High)	—	1.0	0.6	0.8	0.9	§	0.6	-0.3	—	3.3	4.2	3.8	2.3	§	2.3	0.0
Race/Ethnicity (2-year data): ^b																
White	—	—	0.8	0.6	1.1	§	0.6	-0.5	—	—	3.7	3.9	3.8	§	2.4	-1.4 s
African American	—	—	1.0	0.8	0.6	§	1.2	+0.6	—	—	0.9	1.0	0.8	§	0.9	+0.1
Hispanic	—	—	1.6	1.6	1.8	§	1.2	-0.7	—	—	3.5	3.4	4.1	§	2.3	-1.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in S1 and S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 105 (cont.)
EXTREME BINGE DRINKING:
Trends in 2-Week Prevalence of Having 10 or More Drinks in a Row
by Subgroups in Grade 12

	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
<i>Approx. weighted N =</i>	14,700	14,200	14,500	14,000	13,700	13,700	13,700	13,600	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	10.6	12.9	11.1	10.4	10.6	9.9	9.8	10.4	8.1	7.1	6.1	4.4	6.0	4.6	5.3	§	3.2	-2.1* s
Gender																		
Male	16.0	17.8	15.6	14.3	15.2	14.5	13.6	14.1	11.7	10.9	8.0	5.4	7.6	6.4	6.2	§	4.1	-2.1
Female	4.1	6.8	6.2	5.8	5.5	4.9	4.6	5.3	4.1	3.0	3.3	2.8	2.6	2.4	3.6	§	1.8	-1.8 s
College Plans																		
None or under 4 years	14.0	19.4	14.3	16.4	13.4	15.4	14.9	10.9	11.0	10.3	10.4	7.1	7.8	8.5	6.8	§	3.7	-3.1
Complete 4 years	8.2	10.0	9.7	8.2	9.4	8.0	8.3	9.5	6.2	5.7	4.6	3.0	4.7	3.4	3.4	§	2.4	-1.0
Region																		
Northeast	11.0	13.3	13.9	11.0	14.3	9.3	11.2	12.7	8.7	7.5	7.3	4.9	7.4	3.9	2.1	§	3.5	+1.4
Midwest	12.6	15.3	12.1	11.9	13.1	13.8	11.9	10.5	9.4	7.6	8.6	4.3	5.7	3.5	5.6	§	4.7	-0.9
South	10.3	14.2	11.1	10.3	9.6	9.5	9.3	9.0	6.3	7.6	5.1	4.1	6.0	5.5	6.8	§	3.2	-3.6 s
West	8.3	7.1	7.5	8.1	5.7	7.1	7.5	10.8	9.1	5.4	4.8	4.6	5.4	4.5	4.4	§	1.9	-2.6
Population Density																		
Large MSA	9.2	12.6	9.5	10.4	9.5	10.0	10.7	12.5	8.2	5.9	6.9	4.8	7.1	3.8	4.1	§	5.0	+0.9
Other MSA	9.1	12.7	12.2	8.6	10.2	9.0	9.4	8.8	8.8	7.0	4.7	2.5	4.4	4.3	4.8	§	1.4	-3.4 ss
Non-MSA	15.1	13.8	10.9	14.3	13.0	11.9	9.7	11.2	6.4	9.0	8.5	8.4	7.8	6.6	8.6	§	5.2	-3.5
Parental Education ^a																		
1.0–2.0 (Low)	6.8	15.8	6.0	10.6	8.8	7.8	9.4	14.9	9.1	6.9	3.0	3.3	6.3	5.6	5.0	§	2.7	-2.2
2.5–3.0	11.8	13.5	10.6	10.8	11.8	9.2	7.6	9.3	7.8	11.4	5.2	4.7	6.0	4.3	3.8	§	4.4	+0.6
3.5–4.0	10.7	11.4	11.1	8.0	10.8	10.4	10.0	9.9	8.5	4.9	7.2	4.4	3.7	4.5	6.7	§	2.2	-4.5 s
4.5–5.0	10.2	11.3	11.8	11.8	8.4	8.8	9.2	9.9	8.0	6.2	6.1	3.7	6.3	3.7	3.8	§	2.2	-1.6
5.5–6.0 (High)	6.4	10.7	11.2	8.0	9.0	7.1	10.2	9.6	2.7	6.6	5.2	2.4	4.9	4.9	3.8	§	3.6	-0.1
Race/Ethnicity (2-year average) ^b																		
White	—	13.2	13.8	13.3	12.2	11.9	11.9	11.4	9.8	7.9	7.3	6.1	5.9	6.2	5.8	§	5.1	-0.7
African American	—	4.4	4.1	1.4	2.8	4.4	3.5	5.0	5.1	5.0	4.5	1.9	3.2	3.8	2.5	§	0.8	-1.7
Hispanic	—	9.6	8.8	6.7	9.0	8.0	6.8	7.1	8.1	8.0	5.4	3.5	3.0	3.6	3.9	§	3.3	-0.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. " — " indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table D-S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on 1 of 6 forms; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 106
EXTREME BINGE DRINKING:
Trends in 2-Week Prevalence of Having 15 or More Drinks in a Row
by Subgroups in Grade 12

	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change	
<i>Approx. weighted N =</i>	14,700	14,200	14,500	14,000	13,700	13,700	13,700	13,600	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	5.7	7.2	5.6	5.6	6.0	6.3	4.6	5.5	4.4	4.1	3.5	2.3	3.1	2.5	3.2	§	1.3	-1.9* sss	
Gender																			
Male	8.5	10.7	8.3	8.7	8.8	9.4	6.4	7.1	6.4	6.4	4.9	3.6	4.4	3.0	3.7	§	1.0	-2.7 ss	
Female	1.8	2.9	2.7	2.1	3.0	2.3	1.9	2.1	1.7	1.2	1.5	1.0	0.7	0.9	1.8	§	0.7	-1.2	
College Plans																			
None or under 4 years	7.0	12.2	8.3	10.5	7.9	11.0	9.1	6.0	5.9	5.3	8.4	4.2	6.3	4.8	5.2	§	0.8	-4.4 sss	
Complete 4 years	4.1	5.1	4.5	4.1	5.3	4.6	3.3	4.6	2.9	3.1	2.0	1.7	2.0	1.6	1.7	§	0.8	-0.9	
Region																			
Northeast	5.6	5.5	6.0	4.7	6.6	5.7	4.3	5.1	5.5	4.0	3.8	1.7	2.2	1.7	1.7	§	1.7	-0.0	
Midwest	7.1	8.9	6.7	6.5	8.2	9.6	6.8	5.9	4.9	4.2	4.2	2.8	2.3	1.0	2.7	§	1.9	-0.9	
South	5.9	8.4	6.1	6.2	6.1	6.1	4.6	5.6	3.4	4.9	3.0	2.2	3.7	3.9	4.3	§	0.7	-3.6 sss	
West	4.1	4.3	3.5	4.1	2.4	3.9	2.7	5.2	4.4	2.6	3.6	2.5	3.6	2.0	2.7	§	1.4	-1.3	
Population Density																			
Large MSA	4.9	7.2	4.3	4.0	5.5	6.5	4.0	7.1	4.9	3.3	3.9	2.6	3.3	1.3	2.4	§	2.4	+0.0	
Other MSA	5.5	6.6	6.4	4.2	5.4	5.8	4.3	4.2	4.5	4.5	3.0	1.6	2.6	2.2	3.2	§	0.5	-2.7 sss	
Non-MSA	7.4	8.1	6.0	10.5	7.8	7.2	6.3	6.3	3.3	4.2	4.3	3.5	4.2	5.2	4.7	§	1.5	-3.2 s	
Parental Education ^a																			
1.0–2.0 (Low)	5.5	8.0	5.7	7.9	5.6	6.6	6.3	9.7	4.3	4.9	1.9	1.6	2.6	4.1	3.8	§	1.8	-2.0	
2.5–3.0	6.3	7.6	6.4	5.3	7.4	5.2	4.7	4.1	3.2	4.7	2.9	2.8	3.6	1.8	2.5	§	1.2	-1.3	
3.5–4.0	4.0	6.1	5.6	4.6	6.1	6.6	5.5	4.2	5.2	3.4	4.0	2.3	2.7	2.1	3.7	§	0.7	-3.0 ss	
4.5–5.0	4.9	6.9	5.7	6.1	4.0	4.0	2.6	5.0	3.7	3.4	3.3	2.1	2.7	1.1	2.4	§	0.6	-1.8 s	
5.5–6.0 (High)	4.1	4.7	2.7	2.6	4.2	4.9	3.8	5.2	0.6	3.6	2.4	0.7	0.9	3.2	1.0	§	0.6	-0.4	
Race/Ethnicity (2-year average) ^b																			
White	—	6.9	7.0	6.5	6.7	6.9	6.1	5.2	4.9	4.4	4.1	3.1	2.7	2.9	2.7	§	1.5	-1.2	
African American	—	1.1	2.2	1.5	1.8	3.1	2.5	2.8	2.8	2.7	1.6	0.4	2.3	3.2	2.5	§	0.6	-1.9	
Hispanic	—	5.6	5.8	5.1	5.0	4.3	4.4	4.5	4.6	3.8	2.9	2.8	1.8	1.3	1.9	§	2.0	+0.0	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. " — " indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table D-S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on 1 of 6 forms; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 107
BEER: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days															
<i>Approximate weighted N =</i>	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	
Total	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	
	16.2	16.9	17.4	18.3	18.8	18.4	16.7	16.2	16.6	15.2	15.0	12.3	12.0	14.4	12.8	
Gender:																
Male	19.1	18.6	19.1	21.2	20.4	19.9	18.6	17.7	19.4	16.8	16.5	12.2	12.4	14.0	12.9	
Female	13.5	15.3	16.1	16.0	17.0	17.1	14.6	14.7	14.1	13.5	13.6	12.2	11.4	14.6	12.9	
College Plans:																
None or under 4 years	26.4	27.7	32.2	32.8	36.0	33.0	33.9	32.6	35.5	29.6	32.9	27.4	28.7	26.4	26.3	
Complete 4 years	14.6	15.4	15.6	16.6	16.8	16.3	14.6	14.4	14.3	13.8	12.9	11.1	10.2	13.1	11.5	
Region:																
Northeast	14.0	14.0	14.9	18.6	19.3	21.7	18.4	14.7	18.5	14.2	18.7	10.8	12.3	15.1	11.4	
Midwest	16.4	18.0	16.7	18.0	17.4	19.3	16.8	18.4	17.5	17.0	13.2	10.2	14.0	15.9	11.4	
South	17.8	18.5	19.1	18.2	20.1	16.6	15.5	16.1	17.2	14.2	14.3	14.0	11.4	14.4	15.0	
West	14.9	15.1	18.0	18.8	17.6	17.8	17.2	15.1	12.6	15.7	15.3	13.9	10.3	12.2	11.4	
Population Density:																
Large MSA	15.0	18.7	15.1	15.2	12.0	16.2	15.0	13.8	14.3	14.0	12.0	11.5	11.0	14.2	11.4	
Other MSA	16.1	16.8	17.6	19.4	19.1	19.6	15.9	14.3	16.5	14.1	15.5	12.4	11.9	13.9	13.6	
Non-MSA	17.8	15.4	19.8	18.3	21.8	18.8	19.6	22.7	19.4	18.8	17.2	13.2	13.5	15.6	13.1	
Parental Education: ^a																
1.0–2.0 (Low)	22.0	22.5	21.5	24.4	23.9	19.7	21.6	21.9	25.4	23.2	21.3	24.1	17.1	23.7	22.6	
2.5–3.0	18.2	18.4	19.5	19.7	22.8	21.3	21.2	17.4	19.7	20.7	18.4	14.8	15.1	17.8	15.8	
3.5–4.0	15.7	18.0	19.4	19.7	19.6	20.2	18.5	17.9	16.9	13.5	18.1	13.5	14.3	15.0	13.9	
4.5–5.0	14.4	13.4	14.9	15.3	15.6	17.8	12.7	12.6	12.3	12.0	10.0	8.4	7.4	11.9	11.2	
5.5–6.0 (High)	13.5	14.6	14.4	17.2	14.2	13.2	14.1	17.0	16.0	11.4	9.7	10.3	10.2	13.3	6.7	
Race/Ethnicity (2-year average): ^b																
White	—	16.7	17.6	17.9	19.0	19.7	18.8	17.4	17.4	16.8	15.5	13.5	12.0	13.5	14.0	
African American	—	12.6	13.0	15.8	14.5	11.1	9.7	8.1	9.7	9.7	7.9	7.2	7.5	8.1	7.6	
Hispanic	—	23.9	24.2	22.2	23.3	23.7	21.8	19.9	19.5	19.8	20.7	21.8	19.6	18.4	18.3	

Table continued on next page.

TABLE 107 (cont.)
BEER: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days																2019-2021
Approximate weighted N =	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
Total	12.5	12.2	11.8	10.0	10.6	9.8	7.9	6.8	6.3	7.3	4.7	5.7	6.8	5.8	8.4	5.2	-0.6*
Gender:																	
Male	13.0	11.3	12.3	10.6	11.3	10.2	8.5	6.9	6.9	8.1	4.9	5.3	6.3	5.2	§	4.7	-0.5
Female	11.9	12.2	11.2	9.2	10.1	9.2	7.2	6.6	5.8	6.5	4.7	6.3	7.5	6.3	§	5.3	-1.0
College Plans:																	
None or under 4 years	24.7	27.3	30.5	20.9	26.1	16.6	19.5	14.5	16.0	14.5	11.8	11.3	13.0	12.2	§	7.5	-4.6
Complete 4 years	11.4	10.8	10.3	9.0	9.4	9.4	7.0	6.3	5.6	6.8	4.3	5.1	6.1	5.0	§	4.6	-0.4
Region:																	
Northeast	11.4	9.5	10.4	9.2	10.4	6.8	6.8	5.0	4.6	5.8	4.0	3.1	4.8	3.4	§	3.6	+0.2
Midwest	13.2	10.6	12.5	8.4	10.2	9.1	5.0	7.0	6.1	7.6	3.1	4.7	7.0	5.0	§	6.4	+1.4
South	13.3	14.0	14.1	12.2	12.2	10.5	8.8	8.2	6.4	7.8	6.3	6.4	7.1	6.4	§	4.4	-2.0
West	11.4	12.0	8.9	8.8	8.6	11.7	10.1	5.6	7.6	7.1	4.2	7.7	7.5	6.9	§	6.4	-0.5
Population Density:																	
Large MSA	11.9	10.3	13.0	8.9	9.0	8.2	7.4	6.6	4.7	6.8	4.6	4.3	5.6	5.0	§	3.1	-1.9
Other MSA	12.1	12.3	11.3	9.5	10.9	9.9	8.6	7.5	6.5	7.5	4.3	5.8	8.0	5.3	§	5.4	+0.1
Non-MSA	14.3	14.6	11.3	12.5	12.3	12.0	6.9	5.9	8.1	7.5	6.2	7.7	5.7	8.3	§	8.3	-0.1
Parental Education: ^a																	
1.0–2.0 (Low)	17.3	21.0	21.3	15.0	13.6	15.4	12.1	11.8	8.6	13.7	12.4	10.1	9.5	9.0	§	3.8	-5.2
2.5–3.0	16.8	19.5	16.7	13.2	14.6	12.5	9.6	8.3	8.8	8.5	6.2	6.3	9.8	8.4	§	7.2	-1.2
3.5–4.0	13.6	10.6	13.1	10.6	11.5	11.4	8.4	8.1	9.0	8.7	3.2	7.1	7.9	6.7	§	9.5	+2.7
4.5–5.0	9.4	9.3	8.8	8.6	8.8	7.5	6.3	6.1	4.1	4.6	4.4	5.2	5.9	4.8	§	4.4	-0.4
5.5–6.0 (High)	9.3	6.7	7.0	6.5	6.5	7.4	4.7	4.9	4.5	6.1	3.0	3.3	4.8	3.8	§	3.9	0.0
Race/Ethnicity (2-year average): ^b																	
White	12.7	11.5	11.5	11.2	10.7	10.2	8.2	6.8	6.4	6.5	5.4	4.6	5.8	6.5	§	7.4	+0.9
African American	8.5	9.0	7.4	6.3	5.6	5.9	5.8	4.2	3.3	3.7	4.1	3.9	3.1	2.1	§	3.1	+0.9
Hispanic	17.1	17.8	17.8	14.9	13.3	14.0	13.8	11.3	9.0	8.8	8.9	7.8	8.4	8.4	§	4.3	-4.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996; N is one half of N indicated in Table S1. Data based on one of four forms beginning in 1997; N is one third of N indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 108
BEER: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>14,800</i>	<i>14,800</i>	<i>15,300</i>	<i>15,800</i>	<i>17,000</i>	<i>15,600</i>	<i>15,500</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>
Total	31.1	28.9	28.7	30.2	29.9	30.5	30.4	28.3	29.5	30.6	28.0	24.6	23.2	26.5	24.8
Gender:															
Male	37.6	33.1	33.3	37.3	33.5	34.7	34.1	32.5	35.4	35.4	33.7	28.7	27.9	30.7	26.5
Female	25.2	25.1	24.2	23.3	26.2	26.7	27.1	24.3	23.8	25.6	22.9	20.9	19.3	22.3	23.1
College Plans:															
None or under 4 years	39.4	37.5	37.4	41.4	41.9	44.8	43.8	40.5	43.7	42.7	42.3	37.2	30.3	36.4	36.4
Complete 4 years	29.4	27.2	26.9	28.0	27.8	28.1	28.2	26.1	27.0	28.6	25.9	22.7	22.2	25.1	23.2
Region:															
Northeast	37.1	29.4	32.0	31.9	31.7	33.1	33.2	31.0	34.7	31.7	27.0	23.8	25.6	30.4	25.2
Midwest	31.7	29.4	29.1	31.6	31.1	30.4	30.1	27.8	29.8	33.1	29.0	25.6	23.3	22.1	26.2
South	29.4	28.4	26.6	29.5	29.4	30.6	28.5	27.2	27.8	27.1	26.6	22.6	21.9	27.2	23.6
West	28.4	28.8	28.7	27.5	27.3	28.1	31.1	28.1	26.6	32.6	30.6	27.6	22.6	27.0	24.4
Population Density:															
Large MSA	31.0	30.1	28.8	28.9	23.9	28.9	28.8	24.3	29.3	30.1	25.5	22.5	21.6	23.4	21.8
Other MSA	28.9	27.4	26.4	30.8	30.3	30.1	31.2	28.9	29.4	29.8	26.2	24.7	22.9	27.0	25.2
Non-MSA	35.2	30.8	32.8	30.0	32.2	33.0	30.5	31.4	30.0	32.7	34.4	27.1	26.1	29.3	27.9
Parental Education: ^a															
1.0–2.0 (Low)	33.5	27.7	33.3	30.0	34.0	28.9	28.6	29.3	26.5	26.7	26.7	22.4	26.3	28.7	28.9
2.5–3.0	32.0	30.3	28.4	31.1	33.0	31.1	30.3	30.9	32.3	30.8	27.2	28.6	24.1	30.3	29.1
3.5–4.0	31.4	29.9	29.0	31.3	30.1	30.9	31.3	30.0	29.5	32.3	29.0	25.1	24.3	26.2	23.8
4.5–5.0	29.9	27.7	28.0	30.3	28.5	29.0	31.4	25.0	27.4	32.4	27.7	22.7	21.9	26.4	24.4
5.5–6.0 (High)	30.7	27.4	28.1	27.1	26.5	33.7	30.7	26.5	31.1	28.7	30.4	21.9	21.0	24.2	21.7
Race/Ethnicity (2-year average): ^b															
White	—	31.7	30.2	31.1	32.1	32.5	32.4	31.9	32.2	33.1	31.7	28.7	27.0	27.5	28.2
African American	—	20.9	20.3	19.0	18.8	15.9	15.6	13.1	10.0	13.5	14.8	11.6	10.2	11.3	12.5
Hispanic	—	32.2	29.6	29.3	31.2	31.9	31.3	29.8	28.3	28.9	30.7	28.6	25.1	27.8	31.3

Table continued on next page.

TABLE 108 (cont.)
BEER: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days																2019-2021
Approximate weighted N =	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
Total	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	-3.5* ss
Gender:																	
Male	26.8	24.4	22.4	22.6	22.4	19.6	19.6	18.4	17.3	15.7	13.0	14.0	14.1	12.2	13.5	8.7	
Female	29.2	26.8	24.5	25.2	26.3	23.0	22.1	21.0	19.2	16.3	13.4	13.6	16.1	§	9.2	9.2	-4.3 s
College Plans:	24.0	21.8	20.6	19.8	18.4	16.1	17.3	15.7	15.8	15.2	12.3	14.4	12.3	11.1	§	8.3	-2.8
None or under 4 years	39.2	39.2	38.1	38.0	36.9	31.0	27.6	26.9	28.6	24.9	23.6	20.4	16.5	16.2	§	13.4	-2.8
Complete 4 years	25.2	22.6	20.4	20.9	20.6	18.4	18.7	17.4	16.1	14.8	11.8	13.1	13.9	11.5	§	7.7	-3.9 ss
Region:																	
Northeast	31.4	27.3	24.9	23.3	22.3	22.9	23.6	19.7	20.2	17.2	15.2	15.6	15.2	15.1	§	10.3	-4.8 s
Midwest	25.5	24.7	23.0	21.3	21.3	16.6	16.6	14.1	13.8	13.7	10.7	11.3	11.6	13.6	§	9.5	-4.1
South	25.0	21.2	22.7	23.5	22.2	20.8	20.9	21.5	17.6	17.1	13.4	14.7	16.2	10.9	§	8.8	-2.1
West	27.2	25.8	19.7	21.8	23.8	19.0	17.6	17.7	18.0	14.2	12.9	14.5	12.5	11.1	§	6.8	-4.3
Population Density:																	
Large MSA	25.5	24.6	21.7	22.1	22.0	18.3	21.6	16.3	16.7	15.3	14.9	15.8	14.9	11.6	§	5.4	-6.2 ss
Other MSA	27.5	22.4	21.8	22.6	22.1	20.7	18.9	19.0	18.0	15.4	10.6	13.0	14.1	11.2	§	10.3	-0.9
Non-MSA	27.3	28.7	24.8	23.1	23.5	19.6	18.2	19.8	16.5	17.3	16.3	13.7	13.1	15.7	§	11.1	-4.7 s
Parental Education: ^a																	
1.0–2.0 (Low)	27.8	28.9	29.2	26.8	30.4	25.9	23.1	20.6	23.7	19.1	12.5	15.4	17.9	13.9	§	12.4	-1.5
2.5–3.0	30.7	29.7	25.7	25.7	26.8	22.7	21.1	23.3	18.3	20.5	16.6	15.9	14.2	14.5	§	10.2	-4.3
3.5–4.0	28.4	25.4	23.5	22.8	23.3	20.6	18.8	18.0	18.4	16.7	13.0	14.9	13.0	13.6	§	7.7	-5.9 ss
4.5–5.0	23.4	21.5	20.6	20.3	18.5	16.3	19.7	16.8	14.2	13.5	13.4	11.7	13.0	8.9	§	8.6	-0.3
5.5–6.0 (High)	26.0	20.5	17.3	20.0	18.8	17.0	19.9	16.8	18.7	14.6	10.4	13.8	16.3	13.5	§	8.8	-4.6 s
Race/Ethnicity (2-year average): ^b																	
White	28.4	27.8	25.3	23.9	23.3	21.6	21.0	20.3	19.6	18.7	16.3	15.0	15.8	15.3	§	12.0	-3.3 s
African American	11.9	9.7	10.0	11.7	12.5	12.2	8.2	6.9	6.9	7.2	6.1	4.2	4.1	4.4	§	5.1	+0.7
Hispanic	30.7	29.3	28.2	28.0	29.9	28.3	24.4	23.3	20.5	18.2	16.0	16.0	17.5	16.2	§	13.0	-3.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996; N is one half of N indicated in Table S2. Data based on one of four forms beginning in 1997; N is one third of N indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 109
BEER: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	—	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	59.8	62.1	62.3	63.7	62.9	62.7	60.3	61.7	59.5	56.7	55.5	56.2	53.3	51.4	47.2
Gender:																
Male	—	68.6	73.1	73.0	74.0	74.3	71.4	68.3	69.7	67.4	64.5	64.3	64.9	61.8	59.7	55.7
Female	—	51.3	52.8	52.7	54.5	52.4	53.9	52.7	54.0	51.8	49.4	47.9	48.1	46.3	43.4	38.0
College Plans:																
None or under 4 years	—	62.1	64.6	65.2	65.7	68.3	65.0	64.6	62.4	62.7	59.8	54.2	58.2	57.3	55.5	46.5
Complete 4 years	—	57.6	60.3	59.5	61.6	58.4	61.0	57.4	61.1	57.4	55.1	55.7	55.1	51.8	49.4	47.5
Region:																
Northeast	—	64.9	65.9	64.7	71.9	67.6	67.1	65.7	68.3	62.6	64.4	51.5	56.8	54.8	54.2	52.7
Midwest	—	65.4	66.9	68.3	66.1	64.9	67.5	66.1	66.6	66.8	57.4	66.3	61.4	59.8	58.5	50.3
South	—	53.2	55.3	58.4	59.2	60.3	57.5	53.0	56.5	53.7	51.0	48.6	52.1	47.6	46.1	42.0
West	—	52.8	59.4	54.2	56.1	57.7	56.9	52.8	53.8	55.3	56.4	54.4	54.8	52.7	48.5	45.4
Population Density:																
Large MSA	—	62.5	64.2	62.8	65.3	67.7	63.8	63.3	63.1	58.8	56.0	55.1	56.9	52.6	50.1	52.3
Other MSA	—	58.6	62.0	62.4	62.3	62.1	61.9	58.6	59.4	56.6	56.8	55.1	56.4	53.0	51.6	47.3
Non-MSA	—	59.3	60.7	61.7	64.4	60.5	63.0	60.0	63.6	63.7	57.3	56.3	55.4	54.3	52.1	42.4
Parental Education: ^a																
1.0–2.0 (Low)	—	58.5	48.3	54.6	52.9	55.9	54.7	56.8	58.8	51.1	52.2	46.8	55.8	47.1	45.3	42.2
2.5–3.0	—	62.7	64.8	63.6	64.5	66.1	65.4	58.5	59.8	62.4	56.8	54.6	53.6	51.9	52.4	47.5
3.5–4.0	—	58.4	66.0	62.7	67.1	64.4	64.0	63.9	62.3	61.2	57.9	56.0	56.0	51.5	52.2	45.8
4.5–5.0	—	62.6	64.4	64.7	68.4	58.4	60.1	60.0	66.2	61.9	60.5	57.6	59.9	58.2	51.4	49.3
5.5–6.0 (High)	—	64.1	69.3	68.3	63.6	69.3	68.9	68.3	65.9	54.2	55.4	62.5	57.7	57.4	53.2	53.4
Race/Ethnicity (2-year average): ^b																
White	—	—	63.8	65.2	65.9	66.3	66.3	65.4	64.6	64.6	62.2	59.9	59.6	59.1	57.0	54.1
African American	—	—	38.7	36.4	37.4	38.7	38.8	37.2	39.2	36.9	33.1	32.7	28.6	30.3	32.6	28.4
Hispanic	—	—	52.7	54.3	52.3	52.5	47.7	47.5	43.4	40.7	46.5	50.9	55.4	49.2	40.4	36.6

Table continued on next page.

TABLE 109 (cont.)
BEER: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	47.2	42.0	43.4	42.6	44.9	46.9	44.4	45.6	42.7	42.7	41.5	39.7	37.8	38.3	38.0	35.5
Gender:																
Male	55.3	50.1	50.0	51.2	53.0	53.5	49.7	55.0	48.2	51.7	50.9	44.3	44.5	44.1	47.3	39.7
Female	39.3	34.6	37.2	35.6	37.4	40.2	39.9	37.0	36.9	33.9	32.1	34.4	31.7	32.1	30.4	31.1
College Plans:																
None or under 4 years	52.4	48.8	48.2	47.1	50.1	46.2	44.6	49.6	47.5	42.4	54.1	45.7	39.4	44.0	45.1	46.2
Complete 4 years	45.2	39.9	41.5	41.6	42.8	47.0	44.4	44.4	41.7	42.4	38.4	37.9	37.2	36.6	36.6	33.4
Region:																
Northeast	49.5	40.8	47.0	43.3	47.0	56.9	47.5	52.6	54.3	48.9	47.2	43.4	41.9	42.6	41.3	42.9
Midwest	53.5	49.1	46.3	48.0	49.7	48.1	43.3	44.6	45.4	47.6	48.1	42.9	43.1	41.4	40.5	35.8
South	41.1	39.5	43.0	40.4	43.8	44.4	43.3	43.2	36.6	35.1	33.6	35.1	32.0	34.2	33.8	34.2
West	46.0	36.7	37.0	38.2	37.8	38.8	44.5	45.5	40.2	43.5	39.9	39.8	36.9	36.1	38.8	31.3
Population Density:																
Large MSA	44.4	42.3	44.0	42.8	45.9	49.1	45.5	42.1	39.9	45.2	38.0	40.5	36.3	34.2	37.9	35.1
Other MSA	47.6	40.3	42.0	39.4	43.7	46.9	45.0	47.6	42.8	40.7	42.6	38.8	38.9	38.9	38.4	37.2
Non-MSA	48.8	45.1	45.3	47.6	46.0	45.3	41.9	45.6	45.4	43.2	43.3	40.5	37.5	41.4	37.1	33.0
Parental Education: ^a																
1.0–2.0 (Low)	48.1	36.6	41.1	37.0	43.1	31.2	30.4	37.0	35.4	37.2	39.1	41.6	30.9	34.1	32.9	27.5
2.5–3.0	46.7	41.4	43.7	44.4	44.0	42.4	41.5	42.8	43.0	37.4	48.0	36.6	36.2	38.0	35.8	38.1
3.5–4.0	48.4	40.7	44.4	42.2	42.8	48.2	46.3	48.9	42.5	45.1	35.8	40.9	39.9	43.0	38.4	35.7
4.5–5.0	46.4	43.8	41.9	41.1	47.1	47.1	47.7	48.9	40.0	43.9	43.6	39.9	38.4	36.1	42.4	35.3
5.5–6.0 (High)	46.1	47.0	47.7	46.6	49.6	58.0	49.9	44.5	51.8	52.0	42.3	45.2	41.4	38.7	37.6	38.4
Race/Ethnicity (2-year average): ^b																
White	51.6	48.3	46.0	46.5	47.8	50.1	50.7	49.9	47.7	45.8	46.7	45.4	43.1	42.1	42.3	41.8
African American	25.3	24.2	28.0	29.9	25.6	28.4	25.5	22.8	22.3	21.7	20.6	16.9	14.0	15.7	18.0	14.8
Hispanic	40.3	44.7	39.8	40.5	41.7	41.3	39.8	39.0	43.6	45.3	42.6	40.8	37.2	36.7	39.2	35.4

Table continued on next page.

TABLE 109 (cont.)
BEER: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	36.6	33.7	34.9	31.7	29.0	32.2	30.8	28.6	25.4	23.6	25.5	21.8	21.9	§	16.5	-5.4* s
Gender:																
Male	43.7	40.7	42.2	39.7	35.2	39.1	36.9	33.1	28.9	28.2	32.6	25.0	27.3	§	17.0	-10.2 ss
Female	29.3	27.7	28.7	23.0	22.4	25.6	24.6	24.4	22.4	19.0	19.7	18.5	16.7	§	16.6	-0.1
College Plans:																
None or under 4 years	39.0	43.7	38.5	40.6	35.8	33.9	41.5	31.4	27.1	23.5	30.8	23.5	27.8	§	16.2	-11.6 ss
Complete 4 years	35.7	31.7	34.1	30.2	27.5	31.9	28.4	27.6	25.2	23.4	24.7	21.5	20.9	§	16.4	-4.5
Region:																
Northeast	44.1	39.7	42.6	41.2	30.5	40.7	35.7	32.4	38.1	26.5	30.6	23.7	22.1	§	23.8	+1.7
Midwest	39.2	32.7	37.8	31.8	31.2	31.1	33.6	28.8	22.2	26.3	29.6	22.5	19.8	§	15.1	-4.7
South	34.2	33.4	31.0	29.0	25.5	26.4	28.2	26.5	20.8	20.2	23.5	20.4	22.3	§	13.1	-9.1 ss
West	31.7	31.8	30.5	28.9	31.0	35.1	27.9	28.7	26.7	25.6	21.6	22.0	24.0	§	17.6	-6.4
Population Density:																
Large MSA	40.7	36.9	39.5	31.6	30.9	36.2	30.7	30.0	29.4	20.7	25.0	22.7	21.6	§	19.1	-2.5
Other MSA	32.8	31.5	31.7	33.1	27.9	32.4	32.5	29.0	23.4	24.7	22.5	18.4	20.1	§	13.5	-6.6
Non-MSA	38.2	35.0	36.6	28.3	29.0	26.0	26.9	25.8	23.7	25.4	32.4	27.9	27.3	§	17.4	-9.9 s
Parental Education: ^a																
1.0–2.0 (Low)	32.8	28.5	32.2	24.1	28.7	32.1	24.5	26.3	28.2	23.5	11.0	18.0	21.7	§	13.3	-8.4
2.5–3.0	35.2	34.7	30.4	34.8	28.9	30.9	33.7	26.6	26.2	20.6	24.5	21.8	23.3	§	10.4	-13.0 sss
3.5–4.0	36.8	30.9	36.9	31.5	26.2	31.4	33.2	30.5	22.7	23.8	26.9	19.5	19.7	§	18.1	-1.6
4.5–5.0	39.4	33.8	38.9	34.3	32.3	33.6	32.7	25.9	28.3	24.5	28.7	23.6	20.3	§	18.1	-2.2
5.5–6.0 (High)	38.8	43.1	33.7	28.2	29.5	34.7	26.1	39.5	23.2	26.4	33.8	29.1	26.6	§	23.1	-3.5
Race/Ethnicity (2-year average): ^b																
White	41.0	39.7	37.8	36.4	34.0	33.8	34.5	34.0	31.5	29.1	31.0	30.8	25.8	§	22.6	-3.2
African American	13.6	14.6	19.3	18.9	12.0	12.2	15.2	13.4	11.1	9.7	7.3	6.6	6.2	§	5.9	-0.3
Hispanic	32.7	31.2	31.6	32.1	32.7	35.5	32.9	30.7	29.6	23.8	21.6	20.0	21.5	§	11.8	-9.7 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1976–1988; N is one fifth of N indicated in Table S3. Data based on one of six forms beginning in 1989; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 110
BEER: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 8

	Percentage reporting 5+ drinks in a row on one or more occasions														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	7.2	7.3	8.2	8.5	8.5	9.2	7.6	8.0	8.3	7.5	8.1	6.1	5.5	6.8	5.8
Gender:															
Male	9.2	8.9	9.4	10.6	9.8	10.1	9.2	10.0	10.5	8.8	9.4	6.5	5.8	6.4	5.5
Female	5.2	5.9	7.1	6.7	7.4	8.2	6.0	6.3	6.2	6.3	7.0	5.5	5.1	6.9	6.2
College Plans:															
None or under 4 years	15.4	18.2	18.4	20.7	17.5	19.7	22.0	21.6	24.8	21.9	25.4	19.0	17.5	19.0	16.7
Complete 4 years	5.9	5.8	6.9	7.1	7.5	7.6	6.0	6.6	6.2	6.1	6.3	5.0	4.2	5.5	4.9
Region:															
Northeast	4.8	5.5	5.8	8.1	9.3	10.7	7.6	6.6	9.1	5.7	9.4	3.5	5.6	6.5	3.7
Midwest	6.9	7.0	7.6	8.8	8.0	9.2	8.4	10.0	9.3	8.9	7.8	5.6	7.0	8.3	4.7
South	8.9	9.0	9.4	8.1	8.8	8.4	6.9	8.0	8.9	7.1	7.9	6.7	5.1	6.4	7.4
West	7.0	6.7	9.2	9.1	8.1	9.2	7.9	7.1	5.3	8.0	7.9	8.4	4.2	6.0	6.0
Population Density:															
Large MSA	6.6	7.4	6.1	6.9	5.3	8.5	5.6	6.7	6.2	6.0	5.4	5.0	4.2	5.6	4.9
Other MSA	7.0	7.6	8.2	9.1	8.8	9.3	7.4	6.6	8.6	6.4	8.4	6.0	5.3	6.4	6.0
Non-MSA	8.4	6.8	10.7	8.4	9.8	9.8	9.6	12.3	10.2	11.5	10.5	7.7	7.3	8.9	6.6
Parental Education: ^a															
1.0–2.0 (Low)	12.6	12.5	11.0	11.6	12.5	12.8	9.8	12.6	17.9	14.8	11.5	11.2	9.4	11.9	14.0
2.5–3.0	7.7	8.9	9.9	9.3	10.2	10.7	10.1	8.0	9.8	10.9	10.0	8.1	7.2	8.8	8.1
3.5–4.0	7.8	7.0	7.9	9.0	9.8	10.1	8.3	9.0	10.4	6.6	9.1	6.8	6.0	7.9	5.3
4.5–5.0	5.0	4.5	7.3	6.8	6.1	7.5	5.3	5.9	3.9	5.0	5.1	3.7	3.5	5.3	4.3
5.5–6.0 (High)	6.2	6.0	6.2	8.4	5.9	6.7	5.7	8.7	6.1	4.3	5.3	4.8	4.1	4.7	2.4
Race/Ethnicity (2-year average): ^b															
White	—	7.0	7.6	8.0	8.7	9.2	8.6	7.9	8.3	8.3	7.7	6.7	5.7	6.1	5.9
African American	—	5.2	5.1	7.2	6.4	4.9	4.4	3.6	4.7	4.9	4.0	3.2	3.0	3.6	3.1
Hispanic	—	12.7	13.2	12.1	11.6	12.7	12.3	10.5	10.2	9.8	12.3	13.2	10.3	9.1	10.0

Table continued on next page.

TABLE 110 (cont.)
BEER: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 8

	Percentage reporting 5+ drinks in a row on one or more occasions																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
Approximate weighted N =	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	5.5	5.4	5.1	4.9	4.8	4.3	3.5	3.0	2.5	2.5	2.0	2.6	2.9	2.5	3.2	1.9	-0.6*
Gender:																	
Male	5.9	5.5	6.0	5.3	5.2	4.4	3.5	2.8	2.6	2.3	1.9	2.6	3.1	2.2	§	2.1	-0.1
Female	4.9	4.8	4.3	4.1	4.7	4.0	3.6	3.1	2.4	2.6	2.1	2.5	2.7	2.9	§	1.8	-1.1
College Plans:																	
None or under 4 years	13.9	15.3	19.2	15.4	16.8	8.4	12.2	7.8	6.7	8.6	4.9	6.4	7.7	7.0	§	3.0	-4.0 s
Complete 4 years	4.7	4.4	4.1	4.1	3.9	4.0	2.9	2.7	2.2	2.0	1.8	2.1	2.2	1.9	§	1.6	-0.3
Region:																	
Northeast	4.1	4.8	5.3	4.6	4.8	2.1	2.1	2.0	1.8	1.9	1.3	0.8	1.9	0.8	§	0.7	0.0
Midwest	5.9	3.6	4.6	4.4	4.3	4.9	1.5	2.8	2.9	1.9	1.6	2.2	2.9	2.1	§	2.2	+0.1
South	6.3	6.6	6.2	5.8	5.8	4.3	4.1	3.7	2.6	3.1	2.7	3.1	2.8	3.0	§	2.2	-0.7
West	4.9	5.4	3.9	4.2	3.9	5.4	5.8	2.6	2.6	2.3	1.5	3.5	3.8	3.3	§	2.0	-1.3
Population Density:																	
Large MSA	4.1	4.5	5.6	4.4	4.4	3.0	3.6	2.3	1.7	2.1	2.2	1.9	2.8	1.5	§	1.0	-0.5
Other MSA	5.4	5.2	4.7	4.4	5.0	4.6	3.7	3.4	2.7	2.5	1.6	2.6	3.2	2.6	§	2.0	-0.6
Non-MSA	7.8	6.9	5.2	6.8	5.2	5.5	2.9	2.8	3.3	2.9	2.4	3.6	2.2	3.7	§	3.0	-0.7
Parental Education: ^a																	
1.0–2.0 (Low)	8.9	11.6	11.3	9.5	9.2	7.6	7.7	6.1	4.7	6.2	7.3	3.9	4.8	4.0	§	1.9	-2.1
2.5–3.0	7.1	8.8	8.8	6.8	7.6	5.5	5.1	3.1	3.7	4.3	2.5	2.8	4.7	4.1	§	2.1	-1.9
3.5–4.0	5.4	4.8	5.7	5.1	4.8	5.7	2.7	4.1	2.9	1.8	1.7	2.6	2.8	2.8	§	3.2	+0.5
4.5–5.0	4.3	3.4	2.5	4.3	3.4	3.0	2.4	2.2	1.4	1.4	1.4	2.6	1.8	1.8	§	1.5	-0.3
5.5–6.0 (High)	4.7	2.1	2.3	2.0	2.5	2.7	1.4	2.1	1.6	0.9	0.9	1.6	1.9	1.8	§	1.9	+0.1
Race/Ethnicity (2-year average): ^b																	
White	5.2	4.9	4.8	5.0	4.8	4.3	3.3	2.9	2.5	1.9	1.8	2.0	2.3	2.5	§	2.6	+0.1
African American	2.7	2.6	2.4	2.0	1.5	2.9	3.0	1.3	1.7	1.6	0.9	1.4	1.9	1.1	§	0.2	-0.9
Hispanic	9.0	9.0	9.0	7.3	7.1	7.0	7.3	5.9	3.6	3.9	4.3	3.4	3.7	4.3	§	1.8	-2.5

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996; *N* is one half of *N* indicated in Table S1. Data based on one of four forms beginning in 1997; *N* is one third of *N* indicated in Table S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 111
BEER: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 10

Percentage reporting 5+ drinks in a row on one or more occasions

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	16.4	15.1	16.1	17.0	17.1	17.9	17.6	16.0	18.1	17.5	16.3	14.1	13.8	13.6	13.1
Gender:															
Male	21.7	18.9	20.4	23.0	21.1	22.5	21.2	20.7	24.0	22.5	22.2	16.7	18.3	17.1	15.2
Female	11.7	11.7	11.8	11.1	13.1	13.7	14.4	11.8	12.5	12.7	11.2	11.7	9.9	10.4	10.8
College Plans:															
None or under 4 years	25.4	22.8	25.4	26.6	28.2	30.7	31.6	26.1	29.9	26.8	31.2	26.2	22.7	20.5	23.0
Complete 4 years	14.5	13.5	14.2	15.0	15.3	15.8	15.4	14.2	16.2	15.9	14.1	12.2	12.6	12.7	11.8
Region:															
Northeast	19.0	14.6	17.1	18.4	17.4	19.9	17.5	18.8	21.5	16.4	15.2	12.6	16.4	16.9	15.2
Midwest	16.5	16.7	17.3	18.2	18.3	18.8	18.4	16.3	18.3	20.9	16.4	14.9	13.5	10.9	14.6
South	15.4	13.9	14.9	16.9	16.9	17.7	16.6	15.4	17.5	15.7	15.9	13.2	12.4	12.8	10.8
West	15.9	15.5	15.2	13.5	15.5	15.1	18.3	14.0	15.5	16.9	18.3	16.0	14.0	15.0	12.4
Population Density:															
Large MSA	16.0	14.8	15.2	14.5	12.7	16.4	15.9	12.8	18.0	16.9	13.0	12.0	13.0	11.5	10.6
Other MSA	15.8	13.9	13.9	17.0	17.2	17.6	17.6	15.9	17.3	16.3	15.7	13.5	13.1	14.4	13.8
Non-MSA	17.9	17.7	20.9	18.7	19.6	20.1	19.1	19.6	20.1	20.1	21.2	18.1	17.0	14.7	14.6
Parental Education: ^a															
1.0–2.0 (Low)	21.3	15.6	18.9	19.2	20.3	20.5	18.1	12.8	14.2	16.7	17.9	13.8	19.0	14.1	15.1
2.5–3.0	18.5	16.6	15.3	18.5	19.7	18.8	18.0	19.0	20.5	16.5	15.9	15.8	14.8	17.1	16.3
3.5–4.0	15.0	15.6	16.5	16.7	17.4	18.5	18.9	15.8	17.6	17.9	17.8	14.4	13.7	13.9	13.6
4.5–5.0	15.1	14.0	16.3	16.8	16.0	15.9	16.5	14.9	17.3	19.0	14.3	12.5	13.1	12.0	11.6
5.5–6.0 (High)	15.4	13.1	15.3	14.5	13.8	18.3	16.1	16.2	19.3	16.8	17.5	13.1	11.8	13.7	11.3
Race/Ethnicity (2-year average): ^b															
White	—	16.7	16.4	17.4	18.1	19.0	19.3	18.6	19.7	19.9	18.1	16.5	16.1	15.5	15.1
African American	—	9.8	10.1	9.3	8.7	6.5	6.6	5.5	4.0	6.2	6.8	6.1	4.7	4.1	4.2
Hispanic	—	17.2	16.1	16.8	18.6	19.7	19.5	16.9	14.8	17.5	20.3	17.5	16.2	17.5	16.5

Table continued on next page.

TABLE 111 (cont.)
BEER: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 10

	Percentage reporting 5+ drinks in a row on one or more occasions																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	14.5	13.7	10.8	11.3	10.9	10.3	10.4	9.1	8.2	7.4	5.6	6.4	6.8	5.4	6.1	3.9	-1.5* s
Gender:																	
Male	16.3	16.5	12.8	13.6	14.0	12.7	12.9	10.8	9.9	8.6	6.2	7.1	9.4	6.8	§	4.8	-2.0
Female	12.3	10.7	9.2	9.1	8.0	7.7	8.1	7.1	6.6	6.2	4.9	5.9	4.4	4.2	§	3.0	-1.2
College Plans:																	
None or under 4 years	26.1	26.4	21.2	24.3	23.9	20.0	18.3	15.2	18.3	16.4	11.3	13.7	11.2	6.5	§	8.0	+1.6
Complete 4 years	12.8	12.3	9.6	10.0	9.4	9.3	9.7	8.4	7.0	6.5	5.0	5.6	6.2	5.3	§	3.0	-2.3 ss
Region:																	
Northeast	17.7	14.7	12.0	12.1	10.3	12.2	12.7	8.8	9.0	8.7	6.1	7.0	8.2	5.5	§	4.3	-1.2
Midwest	14.8	14.2	11.2	11.0	10.9	8.0	9.9	6.4	6.2	6.4	5.2	4.5	5.3	5.9	§	5.3	-0.6
South	12.7	12.5	11.3	11.6	10.7	11.2	11.1	12.0	9.0	8.0	4.8	7.2	8.2	5.4	§	3.5	-1.8
West	13.7	13.9	9.0	10.4	11.8	10.1	8.0	8.4	8.3	6.5	6.7	6.9	5.1	4.9	§	2.8	-2.1
Population Density:																	
Large MSA	13.3	12.7	10.2	10.4	10.6	9.5	11.4	7.5	7.2	6.8	6.6	6.4	7.0	4.9	§	2.3	-2.6 s
Other MSA	15.3	12.5	10.9	11.5	10.5	10.6	9.4	9.3	8.1	7.1	4.5	6.6	6.1	5.3	§	4.8	-0.5
Non-MSA	14.1	17.8	11.7	12.2	12.3	10.6	11.1	10.8	10.1	9.6	6.8	6.1	8.2	6.5	§	4.7	-1.8
Parental Education: ^a																	
1.0–2.0 (Low)	16.4	19.2	15.1	17.0	19.0	16.0	17.5	13.5	10.9	10.3	5.1	10.2	10.2	6.8	§	9.4	+2.6
2.5–3.0	17.8	18.1	13.2	13.7	11.7	12.1	11.2	10.8	11.1	8.4	7.1	8.1	5.5	6.7	§	5.9	-0.9
3.5–4.0	14.5	13.9	10.9	12.6	11.4	10.5	9.3	9.3	8.8	8.7	5.6	6.4	6.1	5.9	§	3.6	-2.3
4.5–5.0	12.3	11.0	9.5	8.4	8.9	9.2	10.2	7.7	6.0	6.6	5.7	4.8	6.4	2.6	§	3.2	+0.5
5.5–6.0 (High)	14.3	10.6	7.6	9.6	7.9	7.6	10.2	7.6	7.2	6.0	4.6	5.5	7.1	7.4	§	3.3	-4.1 ss
Race/Ethnicity (2-year average): ^b																	
White	15.8	15.7	13.7	12.2	11.4	10.7	11.0	10.1	9.4	8.8	7.2	6.6	7.6	7.0	§	4.8	-2.2 s
African American	3.8	3.7	3.7	4.0	4.4	4.2	3.1	3.1	3.0	3.2	2.8	1.9	1.1	0.9	§	3.9	+3.0
Hispanic	16.0	16.6	15.7	14.7	17.2	17.1	14.3	12.6	8.9	7.3	7.2	8.0	8.8	7.9	§	7.0	-1.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996; N is one half of N indicated in Table S2. Data based on one of four forms beginning in 1997; N is one third of N indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 112
BEER: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	—	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	33.0	35.1	35.4	36.4	36.6	38.0	37.6	38.1	36.2	32.0	33.1	33.8	31.0	31.3	27.0
Gender:																
Male	—	45.6	49.0	49.8	50.6	51.1	49.4	48.5	49.8	47.7	42.6	45.4	45.4	41.5	39.5	37.1
Female	—	21.4	23.5	22.6	24.2	23.9	26.5	27.3	27.4	25.0	22.4	22.4	23.1	22.2	23.8	16.5
College Plans:																
None or under 4 years	—	37.7	39.2	41.6	39.5	43.5	42.8	42.1	43.1	41.2	35.5	36.1	37.8	36.9	37.7	29.0
Complete 4 years	—	27.8	31.0	29.4	33.3	30.9	34.2	34.2	34.3	33.6	29.8	30.8	31.7	28.4	28.5	26.0
Region:																
Northeast	—	39.0	35.3	36.5	41.9	41.4	42.1	40.0	37.9	39.8	35.5	32.6	32.1	32.7	32.1	27.4
Midwest	—	35.8	37.6	39.9	39.0	42.3	43.6	45.3	45.1	43.7	34.2	40.8	41.3	38.5	38.2	31.4
South	—	26.6	33.9	31.8	34.0	30.5	33.9	31.7	35.1	31.5	26.6	29.4	30.7	25.7	27.5	23.8
West	—	31.0	31.6	32.0	28.3	31.1	28.5	28.1	31.3	29.4	33.8	27.5	29.8	28.5	27.4	25.1
Population Density:																
Large MSA	—	32.8	34.4	34.4	35.0	40.9	38.9	36.2	37.4	34.5	31.2	33.9	32.3	29.6	27.3	32.0
Other MSA	—	33.9	35.4	35.7	36.3	33.9	37.7	36.6	36.6	33.2	30.2	31.5	34.5	30.9	32.3	26.9
Non-MSA	—	32.1	35.2	35.7	37.6	37.0	37.6	39.8	40.6	41.4	35.1	35.0	33.9	32.6	32.9	22.6
Parental Education: ^a																
1.0–2.0 (Low)	—	30.7	31.9	33.6	32.8	31.1	32.3	35.0	38.4	26.8	27.4	28.5	32.9	31.3	26.9	28.1
2.5–3.0	—	37.5	36.4	37.8	38.0	38.8	42.9	37.5	37.4	39.2	33.6	34.2	34.3	31.0	32.2	26.7
3.5–4.0	—	32.9	34.8	34.4	36.5	37.6	39.2	41.6	39.1	37.1	32.1	32.9	34.9	29.4	32.8	24.9
4.5–5.0	—	33.9	34.1	36.2	38.2	36.0	31.7	35.3	39.1	39.9	35.9	34.4	34.7	32.8	31.2	27.7
5.5–6.0 (High)	—	27.2	38.0	28.7	35.5	37.6	38.5	38.9	36.5	30.2	27.4	33.8	29.7	31.4	26.5	32.3
Race/Ethnicity (2-year average): ^b																
White	—	—	35.5	36.8	37.6	38.5	40.1	40.9	40.8	40.0	37.1	35.3	35.8	35.2	34.4	32.5
African American	—	—	18.8	19.5	19.4	17.7	16.4	16.9	18.1	18.9	15.5	13.2	13.6	16.1	18.5	13.5
Hispanic	—	—	34.4	33.5	31.5	32.7	30.8	27.6	27.9	27.4	24.6	29.4	32.8	27.5	22.0	20.6

Table continued on next page.

TABLE 112 (cont.)

BEER: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	28.5	25.4	25.5	24.6	27.6	29.2	26.7	28.5	25.6	27.5	26.7	26.1	21.4	23.7	22.4	21.4
Gender:																
Male	37.0	34.0	34.2	35.7	36.5	38.7	32.9	39.3	34.7	38.2	35.9	33.7	31.1	32.1	30.9	26.9
Female	20.1	17.6	18.1	15.5	19.5	19.2	21.2	19.5	17.0	17.6	18.0	18.2	12.9	15.6	15.3	16.2
College Plans:																
None or under 4 years	35.1	32.8	30.2	29.6	32.7	29.4	29.0	39.7	29.5	31.0	40.5	37.0	24.9	32.0	28.9	31.5
Complete 4 years	26.0	23.3	23.8	23.4	25.4	28.8	26.2	25.1	24.7	26.0	23.3	22.9	20.3	21.5	21.1	19.0
Region:																
Northeast	27.8	25.7	28.5	23.4	30.5	35.4	32.0	34.2	33.0	32.7	26.5	28.6	27.1	26.1	25.8	25.9
Midwest	34.7	32.5	27.6	28.3	30.2	29.9	25.7	31.4	28.9	32.4	33.9	28.9	24.8	26.2	26.2	22.1
South	24.2	22.9	25.2	24.8	27.2	27.5	24.1	25.5	21.3	20.0	19.7	22.4	16.8	21.8	20.2	20.6
West	27.3	18.4	20.8	19.7	21.8	24.6	27.6	25.6	22.8	29.4	27.9	26.3	19.2	20.9	18.0	18.1
Population Density:																
Large MSA	24.7	26.0	23.8	23.6	26.9	32.4	26.2	24.4	21.3	29.8	23.4	25.7	18.9	20.9	21.4	21.2
Other MSA	28.8	24.2	24.8	22.6	26.7	29.3	27.9	30.3	26.8	26.1	26.7	25.2	22.9	22.9	23.7	21.2
Non-MSA	31.0	27.2	28.1	28.7	30.0	26.6	25.0	29.5	28.1	27.4	30.2	28.1	21.7	28.2	21.0	22.1
Parental Education: ^a																
1.0–2.0 (Low)	34.9	23.4	27.3	23.7	28.9	23.8	18.7	23.5	22.1	16.8	28.2	32.0	20.1	18.5	17.0	19.3
2.5–3.0	29.2	26.6	26.1	25.3	27.6	25.0	26.0	26.5	27.7	24.3	31.9	24.5	22.9	27.0	21.3	23.2
3.5–4.0	29.4	25.2	26.6	24.2	25.6	29.3	28.3	31.9	26.5	29.1	21.4	27.4	20.5	26.3	23.0	23.9
4.5–5.0	26.2	23.9	24.1	24.1	27.6	28.1	29.1	29.3	24.2	28.7	28.7	24.4	23.3	20.9	25.5	19.6
5.5–6.0 (High)	25.7	26.9	26.1	26.0	32.2	41.0	28.3	27.6	24.3	35.1	28.0	27.3	19.0	20.2	21.8	19.3
Race/Ethnicity (2-year average): ^b																
White	30.6	29.6	27.6	27.3	29.0	31.3	31.4	31.3	30.0	28.8	31.0	30.2	26.4	25.3	26.1	25.8
African American	10.2	13.1	15.5	15.4	12.6	14.6	13.3	11.7	10.6	9.9	9.1	7.6	6.1	9.3	11.2	7.1
Hispanic	23.8	27.2	24.8	24.9	23.8	26.1	26.0	24.4	27.5	29.0	27.1	24.7	21.6	20.0	19.7	19.8

Table continued on next page.

TABLE 112 (cont.)
BEER: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions														2019-2021 change	
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	22.0	20.9	20.8	19.5	16.8	18.8	17.3	16.8	13.6	11.5	14.7	11.2	11.5	§	8.8	-2.7*
Gender:																
Male	27.7	27.7	28.0	26.3	24.2	25.1	23.3	22.3	17.5	14.7	20.5	14.4	15.8	§	10.7	-5.1 s
Female	16.1	15.1	14.3	12.4	10.2	12.3	11.8	11.7	10.0	7.6	9.8	8.1	7.6	§	7.3	-0.3
College Plans:																
None or under 4 years	24.2	30.9	26.7	25.9	21.1	21.3	25.6	18.7	16.6	13.3	21.7	16.5	19.9	§	12.3	-7.6
Complete 4 years	21.2	18.7	19.5	18.3	16.0	18.0	15.3	16.0	12.9	10.7	13.1	9.6	9.9	§	7.9	-2.0
Region:																
Northeast	29.5	23.7	26.0	25.2	17.7	24.9	20.9	18.9	20.0	15.0	19.8	12.1	12.2	§	10.8	-1.4
Midwest	24.3	21.0	23.6	23.3	21.5	18.6	19.8	14.4	13.7	12.7	19.0	10.8	9.9	§	9.7	-0.3
South	18.8	21.1	16.7	16.6	14.2	16.1	15.6	17.1	10.4	9.0	13.3	11.1	12.4	§	7.2	-5.1 s
West	18.6	18.7	18.8	16.0	15.1	18.5	14.2	16.7	14.0	12.7	9.9	11.0	11.4	§	8.8	-2.5
Population Density:																
Large MSA	25.1	22.7	21.0	17.8	17.7	22.4	17.3	17.4	14.8	11.5	14.0	12.4	11.5	§	9.9	-1.6
Other MSA	19.0	18.8	19.4	21.4	15.6	18.3	16.9	15.8	13.2	10.6	12.9	9.0	8.9	§	6.7	-2.2
Non-MSA	23.3	23.4	23.6	17.3	18.3	14.6	18.0	18.2	12.6	13.7	19.6	13.9	18.7	§	10.6	-8.1 s
Parental Education: ^a																
1.0–2.0 (Low)	21.5	14.0	21.8	12.3	15.8	22.5	15.4	19.6	12.5	15.1	7.7	11.7	11.3	§	9.9	-1.4
2.5–3.0	23.1	20.4	17.6	22.3	16.9	19.7	18.4	14.2	15.3	9.6	13.7	10.6	10.9	§	5.6	-5.3
3.5–4.0	21.1	19.8	21.2	19.9	16.1	18.9	19.2	19.3	10.6	10.2	15.9	9.3	11.6	§	9.6	-2.0
4.5–5.0	24.2	21.1	23.6	20.9	18.7	18.1	18.5	15.1	16.2	11.7	16.7	13.3	11.1	§	9.6	-1.5
5.5–6.0 (High)	21.0	27.6	19.6	15.8	15.6	15.9	12.1	17.0	12.6	13.6	17.2	11.3	12.2	§	9.4	-2.7
Race/Ethnicity (2-year average): ^b																
White	25.8	25.3	23.6	23.0	21.4	19.7	19.6	20.2	18.1	14.8	16.7	17.2	13.6	§	10.9	-2.7
African American	5.0	5.7	8.2	6.1	4.8	5.3	5.9	7.7	6.8	4.3	3.1	3.7	3.4	§	1.0	-2.4
Hispanic	19.8	17.5	19.4	20.6	18.6	22.9	22.1	16.6	17.7	15.0	11.8	11.2	11.7	§	5.7	-6.0 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1976–1988; *N* is one fifth of *N* indicated in Table S3. Data based on one of six forms beginning in 1989; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 113
LIQUOR: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Approximate weighted N =	—	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	44.1	45.0	48.2	47.3	47.9	44.6	45.2	46.4	42.3	40.0	41.0	39.0	35.6	35.7	30.8
Gender:																
Male	—	44.4	44.7	48.9	48.5	48.8	45.3	45.2	48.8	43.3	40.3	43.7	42.8	38.5	39.6	36.9
Female	—	43.8	45.3	47.6	46.6	47.4	44.3	45.1	43.7	40.8	39.5	38.6	35.4	33.6	32.1	24.6
College Plans:																
None or under 4 years	—	44.6	47.8	49.3	46.1	51.4	46.9	47.5	47.4	44.6	41.0	38.9	39.4	36.8	38.1	30.6
Complete 4 years	—	43.8	42.5	47.7	48.5	45.1	43.0	43.6	45.3	40.6	39.6	41.8	39.0	35.0	35.0	30.4
Region:																
Northeast	—	52.5	47.4	47.8	56.7	53.2	49.5	50.8	50.8	46.4	42.1	37.2	43.3	36.1	40.3	34.5
Midwest	—	47.0	49.7	53.7	45.4	46.5	47.2	49.1	50.5	45.6	40.1	49.8	42.2	42.2	40.5	31.0
South	—	37.6	39.8	46.0	43.6	48.5	38.8	38.7	43.3	38.5	37.1	35.9	34.6	30.4	33.0	29.6
West	—	38.3	41.3	42.0	45.0	42.9	43.2	40.0	38.4	39.6	42.1	39.6	37.4	35.0	30.0	28.5
Population Density:																
Large MSA	—	47.3	45.4	49.3	52.0	51.7	44.4	48.5	48.1	40.8	37.7	41.2	39.9	36.0	36.4	33.8
Other MSA	—	44.7	46.0	48.3	47.1	48.4	45.9	45.6	45.5	40.9	39.3	39.9	40.5	36.3	33.5	31.1
Non-MSA	—	41.4	43.3	47.2	44.1	44.8	43.2	42.1	46.1	45.1	42.6	42.4	35.5	34.0	40.3	27.7
Parental Education: ^a																
1.0–2.0 (Low)	—	40.8	38.2	37.3	38.2	44.2	43.1	40.3	41.6	33.1	36.2	36.1	32.2	33.5	32.1	25.7
2.5–3.0	—	46.7	45.9	49.8	46.6	47.8	43.5	44.8	44.7	46.3	37.9	38.1	36.0	34.6	32.1	29.5
3.5–4.0	—	44.7	47.3	51.9	51.7	47.5	46.6	48.0	46.5	42.0	39.2	42.3	38.2	33.1	38.3	31.6
4.5–5.0	—	47.9	45.6	49.1	51.7	48.7	40.7	43.6	50.0	44.9	48.0	43.9	43.4	38.7	36.9	34.2
5.5–6.0 (High)	—	46.6	53.4	54.2	49.3	53.4	53.8	54.2	55.2	40.4	40.9	47.4	45.7	39.4	37.7	33.8
Race/Ethnicity (2-year average): ^b																
White	—	—	47.5	49.5	50.3	50.1	48.9	48.2	49.0	47.3	44.1	43.5	43.4	41.1	39.8	37.7
African American	—	—	21.2	19.9	23.3	27.5	25.1	21.5	25.3	23.7	18.5	17.4	15.0	16.0	16.8	14.4
Hispanic	—	—	47.7	48.8	43.3	47.2	45.6	41.2	35.6	35.4	42.1	40.4	36.9	31.6	24.6	21.0

Table continued on next page.

TABLE 113 (cont.)
LIQUOR: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	31.3	28.6	31.4	28.0	34.3	34.7	34.6	37.3	34.3	36.0	35.1	36.0	34.3	35.6	36.4	34.2
Gender:																
Male	34.9	32.0	36.4	32.8	38.2	36.4	36.3	41.2	38.0	41.7	35.6	37.9	36.1	39.3	38.4	33.7
Female	28.0	25.9	27.3	23.2	30.9	32.6	33.4	33.3	30.0	30.7	33.7	33.8	32.9	32.2	34.9	33.9
College Plans:																
None or under 4 years	37.0	31.9	37.3	28.3	36.8	35.8	34.2	38.3	35.0	38.1	40.4	42.5	36.9	40.3	42.6	39.7
Complete 4 years	29.0	27.4	29.9	28.1	33.4	34.1	34.9	36.6	33.9	35.0	33.6	34.2	33.6	34.3	35.5	33.0
Region:																
Northeast	36.6	25.8	32.4	25.9	37.7	40.9	36.6	46.3	39.8	40.9	40.8	41.4	35.8	41.1	41.9	40.0
Midwest	33.5	33.5	32.1	29.0	35.4	36.2	35.2	36.6	33.7	40.5	37.4	39.6	37.4	35.4	39.3	35.2
South	28.1	27.1	34.7	28.1	34.9	33.6	32.9	33.9	32.1	28.3	31.7	32.9	30.7	33.3	33.7	32.5
West	29.0	26.3	24.0	28.2	28.7	27.8	35.1	37.2	34.5	38.5	33.3	31.8	35.4	33.8	32.3	30.9
Population Density:																
Large MSA	30.4	29.8	32.2	30.3	33.3	30.6	36.0	37.2	34.5	41.1	36.0	37.5	32.4	30.1	36.2	34.4
Other MSA	33.1	27.4	30.4	26.5	34.9	38.0	34.6	38.9	34.9	33.8	35.4	37.0	35.6	37.7	37.8	36.3
Non-MSA	29.1	30.0	32.7	28.1	34.3	32.4	33.4	33.9	33.3	34.0	33.9	32.3	34.3	37.4	34.2	30.2
Parental Education: ^a																
1.0–2.0 (Low)	29.7	25.4	29.1	23.6	26.8	24.7	27.7	33.5	35.3	30.8	28.2	29.4	27.6	29.5	30.9	30.2
2.5–3.0	32.6	29.9	29.4	28.2	34.8	37.5	31.8	33.5	33.7	28.7	36.3	33.5	35.2	34.0	33.0	36.7
3.5–4.0	33.6	29.0	34.1	25.8	32.4	34.7	36.4	40.6	32.7	40.0	35.4	37.3	35.7	38.2	33.8	34.7
4.5–5.0	27.4	25.9	32.2	29.3	35.7	30.1	35.8	38.2	33.4	35.2	38.3	38.2	36.5	36.4	43.3	33.8
5.5–6.0 (High)	33.5	33.1	29.9	32.0	41.1	42.1	40.4	38.3	40.8	47.8	34.5	38.8	33.0	36.6	38.4	35.9
Race/Ethnicity (2-year average): ^b																
White	35.0	32.9	32.5	32.6	34.1	37.2	38.5	39.5	37.6	37.2	38.8	38.9	37.8	37.8	39.8	39.5
African American	14.1	15.1	17.5	17.1	17.9	24.9	20.4	17.6	21.5	22.2	22.1	22.0	23.0	23.7	25.3	23.0
Hispanic	23.2	29.2	28.9	26.3	27.4	29.4	28.1	31.7	36.6	36.3	36.0	36.3	33.3	29.6	30.2	28.4

Table continued on next page.

TABLE 113 (cont.)
LIQUOR: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days														2019-2021 change	
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		2021
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	34.1	32.4	33.2	29.8	29.8	31.2	31.0	28.1	26.1	25.1	26.8	21.8	22.4	§	18.2	-4.3*
Gender:																
Male	36.2	33.8	35.2	31.9	31.1	32.3	31.8	29.2	26.0	25.6	26.7	20.0	22.2	§	14.1	-8.1 ss
Female	31.4	30.7	30.9	27.3	28.6	30.4	29.8	27.6	26.3	24.1	27.2	23.3	22.3	§	22.7	+0.4
College Plans:																
None or under 4 years	37.5	38.0	40.6	34.9	29.5	30.0	33.6	25.3	27.8	20.5	29.3	21.4	27.1	§	14.5	-12.6 ss
Complete 4 years	33.5	31.4	31.7	28.8	29.7	31.3	30.4	28.5	26.0	26.1	26.8	22.0	21.7	§	19.4	-2.3
Region:																
Northeast	41.2	37.1	37.7	38.5	38.0	41.8	33.4	29.7	35.9	29.6	28.7	24.4	21.4	§	29.5	+8.1
Midwest	38.1	33.4	34.7	31.1	30.4	27.9	32.5	30.9	23.6	26.3	30.1	22.3	23.5	§	18.0	-5.5
South	30.6	32.2	32.1	26.4	26.8	28.2	27.7	25.4	22.6	21.8	25.1	20.6	22.0	§	11.0	-11.0 ss
West	29.8	28.7	29.3	27.6	28.2	31.7	32.1	28.8	26.9	27.0	25.2	21.6	23.0	§	21.0	-2.0
Population Density:																
Large MSA	37.4	37.1	38.7	30.1	34.3	36.5	35.7	29.9	26.8	25.7	28.3	21.1	24.0	§	23.5	-0.5
Other MSA	31.7	29.2	30.9	31.9	29.2	32.4	30.8	29.0	25.9	24.8	23.7	22.0	20.3	§	14.3	-6.0
Non-MSA	34.1	34.1	31.7	24.2	24.7	20.7	24.9	23.7	25.8	24.9	30.7	22.8	25.4	§	16.5	-8.8
Parental Education: ^a																
1.0–2.0 (Low)	27.1	33.3	32.5	22.5	28.7	28.8	24.7	21.7	27.3	24.3	10.8	16.8	16.6	§	14.0	-2.6
2.5–3.0	33.7	30.8	28.1	33.3	28.9	31.7	29.9	28.9	24.4	22.2	27.9	19.1	22.7	§	13.1	-9.6 s
3.5–4.0	33.2	33.1	33.9	29.3	29.8	32.4	34.6	29.1	23.4	24.4	28.1	22.9	23.3	§	15.8	-7.6
4.5–5.0	37.3	30.3	36.8	30.5	29.5	31.9	34.2	28.1	28.9	29.9	30.0	25.0	22.0	§	22.7	+0.7
5.5–6.0 (High)	39.4	37.5	35.9	28.7	35.3	30.3	26.6	32.5	28.6	24.5	34.5	27.5	27.2	§	25.4	-1.8
Race/Ethnicity (2-year average): ^b																
White	37.5	35.2	33.8	33.0	31.9	32.5	32.5	32.2	30.4	28.2	30.3	29.6	25.8	§	20.8	-5.0
African American	21.6	23.7	29.4	27.4	22.8	26.7	27.1	21.6	19.2	17.6	16.3	15.3	15.0	§	10.0	-5.0
Hispanic	27.5	29.3	30.5	30.5	30.9	30.9	30.6	29.1	26.4	25.4	22.6	18.7	17.9	§	12.1	-5.7 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1976–1988; N is one fifth of N indicated in Table S3. Data based on one of six forms beginning in 1989; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 114
LIQUOR: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	—	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	19.8	19.9	22.0	22.8	23.0	21.3	23.8	22.0	21.3	19.6	21.4	20.5	18.2	20.0	16.8
Gender:																
Male	—	23.3	21.1	24.4	26.3	26.1	23.0	25.8	26.4	24.6	20.9	24.7	25.0	20.3	24.5	20.7
Female	—	16.1	18.5	19.6	19.8	20.2	19.7	21.9	17.8	18.2	18.2	18.6	16.2	16.6	15.7	12.8
College Plans:																
None or under 4 years	—	23.8	23.4	26.7	24.2	26.5	26.4	26.9	25.7	24.7	22.0	24.0	24.5	20.7	23.7	17.3
Complete 4 years	—	15.8	16.8	17.9	21.5	19.7	18.0	21.8	19.2	19.2	18.4	19.9	18.6	17.2	18.6	16.4
Region:																
Northeast	—	24.9	20.2	21.8	28.8	27.2	25.0	26.1	24.7	24.9	23.8	20.1	22.3	17.4	23.7	18.0
Midwest	—	22.1	23.3	24.0	23.0	23.7	22.9	27.5	24.0	24.1	19.3	25.9	21.9	23.1	21.4	17.7
South	—	16.0	16.4	22.5	20.6	22.1	18.0	19.6	20.7	20.2	17.0	19.1	18.7	15.2	18.6	15.4
West	—	14.8	19.2	16.2	18.5	18.2	19.2	20.1	17.0	15.3	19.8	19.6	19.6	17.2	17.5	16.5
Population Density:																
Large MSA	—	20.3	18.3	20.5	22.6	22.8	22.6	26.1	24.3	21.1	18.7	21.7	20.3	16.8	18.4	17.9
Other MSA	—	19.9	21.8	21.4	23.4	23.9	20.7	22.2	21.8	20.1	19.6	19.7	20.8	19.8	19.1	16.4
Non-MSA	—	19.4	18.8	23.8	22.3	22.1	21.2	24.1	20.5	23.0	20.2	23.9	20.2	16.6	23.7	16.8
Parental Education: ^a																
1.0–2.0 (Low)	—	22.0	18.1	20.9	19.1	22.0	24.3	21.0	21.5	21.0	19.6	19.0	20.6	19.9	17.6	18.0
2.5–3.0	—	22.8	20.4	23.7	23.6	23.9	21.3	24.7	23.6	25.5	20.4	24.2	21.2	21.6	18.1	16.4
3.5–4.0	—	17.4	21.5	22.0	22.9	20.5	24.9	25.2	19.9	19.7	18.1	20.1	18.1	15.7	23.2	15.0
4.5–5.0	—	18.0	16.7	18.6	24.7	24.8	14.7	21.3	22.6	20.0	20.2	20.4	21.8	16.7	20.9	18.5
5.5–6.0 (High)	—	14.0	24.9	22.5	23.2	23.6	19.5	28.1	20.4	17.5	20.6	22.0	22.2	16.4	16.3	19.4
Race/Ethnicity (2-year average): ^b																
White	—	—	20.5	22.1	23.6	24.1	23.3	24.2	24.5	23.0	22.0	22.0	22.6	21.3	21.0	20.7
African American	—	—	10.5	7.7	8.4	11.1	10.6	8.6	10.9	11.2	8.0	7.5	6.2	7.2	10.4	9.6
Hispanic	—	—	25.5	26.8	25.9	22.9	25.3	26.7	24.5	21.4	20.9	20.8	21.9	19.0	14.6	11.9

Table continued on next page.

TABLE 114 (cont.)
LIQUOR: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	17.4	16.0	18.2	17.1	20.2	21.6	21.1	23.6	22.3	23.8	22.7	25.6	21.1	23.8	25.0	23.3
Gender:																
Male	20.5	20.0	24.6	22.4	24.7	24.6	23.9	27.3	27.6	30.8	26.4	27.2	24.1	29.2	28.5	24.7
Female	14.4	12.6	13.2	12.6	16.2	17.7	18.7	20.0	16.1	17.4	18.8	23.6	18.9	18.2	22.0	22.0
College Plans:																
None or under 4 years	21.7	21.4	24.6	19.1	22.5	21.1	23.4	31.2	23.2	26.4	29.2	31.9	23.0	29.2	32.6	27.1
Complete 4 years	15.8	14.1	16.5	16.6	19.2	21.4	20.6	21.7	21.9	22.9	20.7	23.7	20.5	22.1	23.7	22.6
Region:																
Northeast	21.3	14.8	15.1	14.2	19.2	25.4	24.6	29.8	25.5	24.7	27.9	27.8	23.5	28.2	28.7	29.0
Midwest	19.0	19.0	18.9	17.7	21.5	22.0	20.6	23.6	19.9	29.7	21.5	28.1	24.8	23.4	28.3	26.2
South	15.3	15.5	21.7	17.6	21.8	21.9	20.0	21.7	21.7	17.4	20.4	24.3	17.4	21.6	22.8	20.5
West	15.2	13.7	13.6	18.0	16.4	16.3	20.5	22.3	23.6	26.1	24.3	22.7	20.7	23.4	21.2	19.8
Population Density:																
Large MSA	16.4	15.6	16.8	16.1	19.3	19.5	23.0	23.9	21.4	26.6	23.6	25.8	20.0	20.7	25.3	22.6
Other MSA	17.5	15.2	17.3	16.3	20.3	24.9	21.3	25.1	22.8	21.7	22.2	26.3	21.9	24.6	26.2	25.0
Non-MSA	18.1	17.9	20.7	19.5	21.1	17.7	18.7	20.3	22.2	24.1	22.6	24.0	21.2	25.5	22.5	21.0
Parental Education: ^a																
1.0–2.0 (Low)	19.9	15.9	20.7	14.4	15.2	14.7	16.7	23.9	25.5	24.4	15.2	18.4	17.9	17.8	18.5	19.1
2.5–3.0	18.1	17.9	18.3	18.5	19.7	22.1	18.3	21.3	20.7	17.8	25.4	24.2	24.2	25.9	22.1	22.7
3.5–4.0	18.6	15.9	19.7	16.2	20.7	20.4	23.3	25.4	23.6	25.8	21.8	24.4	21.1	26.2	23.5	26.1
4.5–5.0	16.4	13.5	17.4	17.7	21.6	18.0	22.2	24.3	20.6	24.5	24.9	29.0	21.0	22.2	29.8	24.5
5.5–6.0 (High)	13.3	16.5	16.0	18.8	22.0	31.3	23.5	23.0	22.8	30.4	22.4	31.0	21.0	20.3	28.7	20.3
Race/Ethnicity (2-year average): ^b																
White	19.0	18.0	18.7	19.4	20.4	22.3	23.4	24.7	24.4	24.3	25.3	26.3	25.4	24.6	26.9	27.5
African American	6.8	7.1	8.9	9.5	11.3	15.4	12.5	9.9	10.4	10.2	12.8	14.6	14.9	15.8	17.2	14.9
Hispanic	13.4	19.4	18.1	16.8	18.1	18.9	20.0	22.2	24.3	25.2	24.0	24.0	21.1	18.9	20.6	18.4

Table continued on next page.

TABLE 114 (cont.)
LIQUOR: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															2019-2021 change		
<i>Approximate weighted N =</i>	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021			
Total	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	§	11.4	-3.3*
Gender:																		
Male																§	9.6	-5.1 s
Female																§	13.9	-0.5
College Plans:																		
None or under 4 years																§	11.2	-7.3
Complete 4 years																§	11.5	-2.6
Region:																		
Northeast																§	13.8	+1.2
Midwest																§	13.4	-1.8
South																§	6.6	-8.4 ss
West																§	15.0	+0.4
Population Density:																		
Large MSA																§	15.0	+1.7
Other MSA																§	8.7	-5.0
Non-MSA																§	10.3	-9.4 s
Parental Education: ^a																		
1.0–2.0 (Low)																§	11.1	+1.9
2.5–3.0																§	7.4	-8.6 s
3.5–4.0																§	10.3	-5.7
4.5–5.0																§	13.6	0.0
5.5–6.0 (High)																§	14.8	-2.2
Race/Ethnicity (2-year average): ^b																		
White																§	15.2	-1.4
African American																§	8.4	-0.9
Hispanic																§	5.8	-4.8 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1976–1988; *N* is one fifth of *N* indicated in Table S3. Data based on one of six forms beginning in 1989; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 115
WINE: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988 ^a	1989	1990
<i>Approximate weighted N =</i>	—	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	34.5	37.4	37.7	36.6	39.5	36.6	37.8	36.0	34.1	36.0	34.5	33.5	22.7	19.7	14.7
Gender:																
Male	—	34.1	35.5	34.8	33.9	38.1	35.3	35.6	33.8	30.3	34.2	30.8	29.3	20.7	18.9	16.1
Female	—	34.5	39.1	40.2	39.0	41.1	38.2	39.9	37.7	38.1	37.4	38.1	37.1	24.3	20.4	13.5
College Plans:																
None or under 4 years	—	31.5	33.1	33.8	33.8	37.1	32.8	36.3	34.5	33.7	34.6	31.5	28.2	19.7	18.6	11.9
Complete 4 years	—	38.0	41.4	41.5	38.5	41.0	39.1	39.5	37.2	34.0	37.3	36.2	35.7	24.0	20.5	16.0
Region:																
Northeast	—	45.2	42.6	47.5	45.3	48.0	51.6	51.4	46.0	41.5	43.9	32.0	35.6	27.7	22.1	15.0
Midwest	—	33.8	36.3	40.8	33.8	36.6	38.5	38.9	36.0	34.3	34.1	41.0	34.0	23.0	23.7	13.4
South	—	27.5	34.2	30.2	30.8	34.8	26.3	27.7	30.9	29.5	30.7	28.1	31.7	18.9	17.4	14.5
West	—	34.0	38.1	32.7	40.8	42.4	31.6	32.5	32.4	34.7	38.4	38.1	33.8	24.5	16.0	16.9
Population Density:																
Large MSA	—	46.6	47.0	45.6	44.2	46.9	46.0	46.3	45.0	36.7	40.7	36.6	35.3	27.6	20.6	15.3
Other MSA	—	33.0	36.5	36.8	36.0	42.1	34.6	36.6	32.4	34.0	36.4	36.0	34.9	22.0	20.5	16.8
Non-MSA	—	28.0	31.4	32.8	31.9	31.0	32.4	32.8	33.8	32.3	31.7	30.7	29.2	19.3	16.7	9.9
Parental Education: ^b																
1.0–2.0 (Low)	—	33.4	30.1	32.3	31.4	28.0	28.0	30.2	26.6	28.3	28.6	28.0	23.9	16.4	19.0	9.5
2.5–3.0	—	33.8	32.8	35.3	33.0	38.0	33.2	33.8	35.3	34.0	32.4	31.3	30.7	17.8	15.6	12.4
3.5–4.0	—	36.9	42.0	40.2	38.2	39.6	37.2	42.0	32.9	35.9	36.9	37.1	31.5	23.4	18.8	14.2
4.5–5.0	—	39.6	42.5	40.8	43.6	45.2	41.5	43.4	46.1	37.1	43.1	36.3	39.7	26.5	24.1	18.9
5.5–6.0 (High)	—	48.4	57.3	49.8	46.7	51.7	52.9	47.5	43.2	33.9	42.6	40.1	40.9	31.4	23.5	19.3
Race/Ethnicity (2-year average): ^c																
White	—	—	35.9	38.1	38.1	39.2	40.2	39.6	38.4	36.5	36.7	37.1	35.7	29.9	23.0	18.6
African American	—	—	36.2	30.8	27.4	27.9	24.9	26.0	28.9	24.4	21.3	21.0	21.8	18.4	15.0	11.2
Hispanic	—	—	42.9	38.5	30.0	31.0	27.3	26.8	28.3	30.9	37.0	37.4	33.0	22.2	15.0	14.2

Table continued on next page.

TABLE 115 (cont.)
WINE: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>
<i>Approximate weighted N =</i>	<i>15,000</i>	<i>15,800</i>	<i>16,300</i>	<i>15,400</i>	<i>15,400</i>	<i>14,300</i>	<i>15,400</i>	<i>15,200</i>	<i>13,600</i>	<i>12,800</i>	<i>12,800</i>	<i>12,900</i>	<i>14,600</i>	<i>14,600</i>	<i>14,700</i>	<i>14,200</i>
Total	16.1	16.9	14.9	14.2	14.3	18.3	17.0	16.0	15.8	16.2	14.1	13.4	13.4	13.9	14.4	12.6
Gender:																
Male	16.9	17.2	15.9	13.8	13.2	18.1	16.7	17.4	17.6	17.7	12.8	13.2	13.9	13.8	13.8	11.6
Female	15.5	16.6	13.9	14.3	15.3	18.2	17.3	15.0	14.4	15.0	15.5	13.2	13.5	14.1	15.1	13.6
College Plans:																
None or under 4 years	16.1	12.6	16.5	11.7	10.6	14.5	14.2	19.0	11.9	12.0	15.9	11.7	13.8	12.7	14.4	13.9
Complete 4 years	16.4	18.4	14.4	14.8	15.3	19.3	18.0	15.3	16.8	17.1	13.3	13.7	13.5	14.5	14.3	11.9
Region:																
Northeast	22.1	18.5	15.2	15.9	14.2	23.4	19.4	18.7	23.5	20.8	19.2	17.3	12.5	18.2	15.2	13.9
Midwest	17.3	21.9	14.0	14.3	14.3	13.1	14.2	17.7	16.2	11.9	13.5	14.9	13.5	13.0	15.3	13.9
South	11.8	11.0	16.7	13.0	15.9	20.0	17.7	13.8	14.6	17.2	13.9	12.7	14.0	12.6	13.4	12.1
West	16.2	18.3	13.0	15.0	11.6	16.8	17.1	15.9	11.2	15.9	11.3	9.8	13.5	12.8	14.1	10.7
Population Density:																
Large MSA	13.9	17.0	20.0	16.8	14.8	19.5	18.2	19.5	17.4	20.4	16.7	17.2	11.1	12.5	16.9	15.0
Other MSA	17.8	19.0	15.0	14.4	15.1	18.6	18.2	15.9	16.5	14.6	11.5	11.8	15.0	15.4	14.0	12.8
Non-MSA	15.1	12.6	11.3	11.3	12.4	16.6	13.6	12.3	13.3	14.2	15.6	11.9	13.5	12.8	12.2	8.9
Parental Education: ^b																
1.0–2.0 (Low)	10.8	11.7	14.3	9.8	13.5	9.7	13.1	14.6	16.7	19.7	12.3	14.8	9.4	12.8	10.9	4.0
2.5–3.0	16.0	13.9	12.1	13.6	10.2	15.2	13.8	10.2	10.5	10.4	13.6	8.6	13.1	10.9	13.4	10.6
3.5–4.0	16.4	15.4	14.3	11.7	10.5	17.9	15.7	15.1	11.5	14.6	10.2	13.1	13.1	15.6	12.2	11.4
4.5–5.0	17.5	19.2	17.7	14.6	18.1	18.0	18.0	20.8	19.6	16.2	19.1	15.4	15.5	14.1	16.8	15.5
5.5–6.0 (High)	19.4	27.1	16.5	23.8	24.9	27.2	29.1	20.0	29.8	29.1	16.7	19.0	15.8	18.3	18.2	18.4
Race/Ethnicity (2-year average): ^c																
White	16.6	17.4	16.2	14.4	14.4	16.8	19.3	18.1	16.6	16.4	15.9	15.2	14.5	14.7	15.5	15.6
African American	10.2	11.7	14.5	17.6	13.6	12.8	11.1	9.9	9.3	9.5	10.9	8.5	8.0	7.9	7.8	6.1
Hispanic	10.9	14.9	14.3	14.2	15.0	13.8	13.5	13.4	16.4	17.9	12.9	9.4	12.2	13.5	13.1	10.5

Table continued on next page.

TABLE 115 (cont.)
WINE: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	14.1	14.0	11.5	9.3	10.2	10.6	11.5	11.5	8.4	9.5	10.8	9.7	10.3	§	8.5	-1.8*
Gender:																
Male	12.1	13.4	12.0	8.7	9.2	8.8	10.2	10.7	5.5	6.8	8.7	5.9	8.0	§	5.1	-2.9
Female	15.3	14.4	10.9	10.2	11.0	12.5	12.7	12.6	10.2	11.6	12.7	13.6	12.0	§	11.4	-0.6
College Plans:																
None or under 4 years	13.1	17.3	10.3	10.6	8.3	5.6	8.9	12.9	8.0	7.5	9.7	4.9	7.8	§	5.9	-2.0
Complete 4 years	14.4	13.4	11.6	9.0	10.6	11.9	12.1	11.1	8.7	9.6	11.2	11.2	11.0	§	9.4	-1.6
Region:																
Northeast	17.6	17.0	17.6	13.4	12.2	13.2	11.2	13.5	13.9	11.8	15.5	15.6	11.4	§	14.4	+3.0
Midwest	14.1	13.9	14.6	12.3	8.8	10.4	12.4	12.4	10.4	8.6	12.0	9.3	8.7	§	8.1	-0.6
South	13.9	13.3	7.6	6.8	10.0	9.3	10.5	10.9	6.3	8.6	9.2	10.2	11.7	§	4.7	-7.0 sss
West	11.5	13.3	8.2	7.3	10.8	10.7	12.0	10.0	5.9	10.2	9.3	5.5	8.1	§	10.2	+2.1
Population Density:																
Large MSA	16.7	18.8	12.6	9.6	11.1	16.0	12.9	13.6	9.7	9.5	11.3	9.8	9.6	§	11.9	+2.4
Other MSA	13.5	13.0	12.3	10.3	11.0	9.4	10.3	11.5	7.6	8.2	11.2	8.7	10.2	§	8.2	-2.1
Non-MSA	11.3	11.0	8.3	6.6	7.4	5.6	12.2	8.7	7.9	12.4	9.4	11.7	11.5	§	3.6	-7.9 ss
Parental Education: ^b																
1.0–2.0 (Low)	9.9	11.3	9.1	5.7	9.0	8.9	5.6	6.6	6.3	4.9	3.1	5.9	3.2	§	9.8	+6.6
2.5–3.0	9.0	11.9	8.8	9.5	5.8	9.0	10.5	7.9	7.2	7.0	9.3	6.7	5.4	§	3.3	-2.1
3.5–4.0	15.3	13.3	10.2	7.5	11.2	11.1	12.0	11.5	7.8	10.4	10.4	9.0	11.1	§	5.6	-5.5
4.5–5.0	16.4	13.7	13.6	12.3	8.8	11.6	14.9	10.8	9.4	10.6	11.7	12.1	12.0	§	13.0	+1.0
5.5–6.0 (High)	19.9	17.2	17.9	10.0	17.3	12.4	11.4	25.5	12.4	10.9	19.8	16.7	16.8	§	12.2	-4.6
Race/Ethnicity (2-year average): ^c																
White	15.4	14.3	12.8	11.5	10.8	11.5	12.2	13.0	12.1	10.6	11.8	13.3	12.7	§	8.9	-3.8 s
African American	6.5	11.2	11.6	7.6	8.8	10.4	10.5	7.7	5.4	6.7	6.6	4.5	7.3	§	5.9	-1.4
Hispanic	11.0	15.7	14.2	9.4	8.2	9.4	7.8	6.6	6.5	6.6	7.4	6.9	6.4	§	6.4	0.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1976–1988; N is one fifth of N indicated in Table S3. Data based on one of six forms beginning in 1989; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 116
WINE: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988 ^a	1989	1990
Approximate weighted N =	—	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	12.5	13.1	13.5	12.9	13.2	13.0	14.9	14.3	13.0	12.8	13.8	12.7	7.8	6.8	4.9
Gender:																
Male	—	14.5	13.9	14.3	13.9	13.3	12.2	12.6	14.4	12.6	12.9	12.4	12.3	7.8	6.1	5.5
Female	—	10.4	12.1	12.8	12.0	12.8	13.5	16.8	13.9	13.1	12.5	14.6	12.8	7.7	7.3	4.3
College Plans:																
None or under 4 years	—	14.1	14.6	14.6	14.8	15.3	13.2	16.1	15.7	14.4	13.9	14.6	13.7	9.5	9.0	5.1
Complete 4 years	—	10.4	11.6	12.7	11.4	11.3	12.6	13.9	13.2	11.9	11.9	13.3	12.0	7.1	6.0	4.9
Region:																
Northeast	—	14.8	14.8	15.0	16.4	18.2	17.9	18.3	19.1	16.9	15.7	11.5	10.7	8.1	7.4	3.6
Midwest	—	13.8	12.9	15.4	12.6	12.2	13.9	16.0	13.6	14.1	12.3	17.2	13.5	7.8	6.9	3.9
South	—	10.1	11.8	11.7	11.8	11.6	10.0	12.4	12.5	11.0	10.9	12.9	13.1	6.8	6.7	5.4
West	—	10.8	13.4	10.9	10.4	11.6	9.8	12.0	13.1	10.9	13.3	12.2	13.2	9.5	6.3	7.0
Population Density:																
Large MSA	—	17.0	14.9	14.7	14.3	14.9	16.8	17.9	17.1	14.1	14.0	13.5	11.0	9.7	6.1	4.4
Other MSA	—	11.4	13.0	12.5	11.8	13.2	11.6	13.3	12.6	12.4	13.7	13.8	13.6	7.8	8.0	4.9
Non-MSA	—	10.7	11.8	14.0	13.1	12.0	11.8	14.7	14.4	13.0	10.6	13.9	12.9	6.0	4.8	5.2
Parental Education: ^b																
1.0–2.0 (Low)	—	14.0	16.4	14.9	12.7	10.2	13.7	15.7	12.1	12.8	11.9	14.7	12.2	8.2	8.3	4.6
2.5–3.0	—	13.4	12.0	13.7	12.4	15.4	13.1	12.1	16.8	14.6	12.1	13.6	13.0	7.2	6.2	5.4
3.5–4.0	—	11.7	14.0	12.2	10.3	13.0	13.4	16.7	12.4	15.0	13.1	14.9	12.1	7.5	6.3	5.4
4.5–5.0	—	12.4	9.4	12.9	16.5	12.7	11.2	14.9	14.4	10.2	13.4	12.2	13.3	7.7	8.0	4.3
5.5–6.0 (High)	—	14.4	15.4	15.0	16.5	12.0	14.2	21.2	12.5	7.7	14.5	12.9	12.3	9.5	5.0	4.4
Race/Ethnicity (2-year average): ^c																
White	—	—	12.0	13.1	13.4	13.1	13.4	14.4	14.8	13.8	13.3	13.6	13.4	10.4	7.5	6.0
African American	—	—	16.7	12.6	10.8	10.4	9.0	10.0	11.4	10.1	8.3	8.3	7.6	8.3	7.8	4.7
Hispanic	—	—	19.3	14.7	9.7	9.5	15.3	15.1	15.8	14.3	13.4	16.1	14.7	8.4	5.1	6.4

Table continued on next page.

TABLE 116 (cont.)
WINE: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	5.8	5.7	5.8	4.9	5.8	7.0	6.4	6.4	5.0	5.4	5.9	4.7	5.0	4.9	5.4	4.4
Gender:																
Male	7.1	6.5	7.2	4.9	6.1	8.3	6.9	6.9	6.5	7.3	5.8	4.6	5.7	6.2	5.1	4.9
Female	4.6	4.5	4.4	4.9	5.6	5.7	6.1	5.8	3.8	3.6	5.3	4.1	4.3	3.5	5.6	4.1
College Plans:																
None or under 4 years	8.0	6.7	8.3	4.0	6.1	5.7	8.7	9.6	5.0	4.2	7.8	6.4	7.0	5.2	7.1	6.0
Complete 4 years	5.3	5.4	5.1	5.3	5.4	7.2	5.7	5.5	5.1	5.6	5.2	4.1	4.5	4.8	4.8	4.0
Region:																
Northeast	7.1	4.5	3.4	4.2	5.9	10.2	6.6	8.0	5.4	5.9	8.3	5.3	4.1	4.6	5.2	5.6
Midwest	5.6	7.2	4.6	4.6	6.3	5.7	5.5	6.4	6.4	5.3	4.5	4.6	5.5	5.1	6.1	5.0
South	5.1	4.1	8.2	4.7	6.9	6.7	7.2	5.7	5.0	5.2	5.8	3.9	5.4	4.9	5.1	3.6
West	6.0	7.1	5.3	6.5	3.0	6.1	6.3	6.5	3.2	5.5	6.1	5.8	4.5	4.9	5.3	4.1
Population Density:																
Large MSA	5.3	6.1	5.4	4.7	5.6	9.6	7.3	6.6	5.3	6.7	5.5	5.4	4.1	5.3	5.8	5.3
Other MSA	6.5	5.8	6.8	6.2	5.4	7.0	6.4	6.6	4.2	5.4	5.4	3.2	5.0	4.6	5.3	3.9
Non-MSA	4.9	5.1	4.5	3.2	6.6	4.9	5.6	5.8	6.2	3.9	7.0	6.7	5.9	5.0	5.2	4.1
Parental Education: ^b																
1.0–2.0 (Low)	5.7	8.0	12.4	4.9	6.1	4.2	5.2	10.7	8.2	11.3	8.0	8.2	6.8	6.4	5.6	4.3
2.5–3.0	6.8	4.8	5.3	3.8	5.6	6.5	6.8	6.0	3.5	3.7	5.6	4.6	6.6	4.4	5.3	2.8
3.5–4.0	6.5	4.1	4.3	5.5	4.2	7.1	4.8	5.7	4.1	5.3	4.7	4.5	4.6	6.5	4.6	3.6
4.5–5.0	4.9	6.1	6.0	3.7	5.6	4.4	8.1	7.6	4.7	4.8	7.2	4.4	4.1	4.1	4.7	5.2
5.5–6.0 (High)	3.5	8.3	4.4	8.4	11.1	13.0	8.2	4.5	8.8	6.5	5.5	4.4	4.2	4.2	8.1	7.3
Race/Ethnicity (2-year average): ^c																
White	5.3	5.5	5.0	4.8	5.2	6.2	6.9	6.5	5.4	4.7	4.8	4.9	5.0	4.9	5.1	5.2
African American	5.1	5.1	7.6	7.5	4.0	4.9	5.1	5.3	4.5	5.2	6.5	5.0	4.3	3.5	2.3	1.7
Hispanic	4.3	7.5	7.8	7.2	7.3	7.6	7.6	7.1	7.8	7.8	7.8	4.7	3.5	5.2	5.0	3.9

Table continued on next page.

TABLE 116 (cont.)
WINE: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	5.1	5.0	4.0	3.7	3.5	3.9	4.0	4.9	4.5	3.6	4.4	3.6	3.6	§	3.2	-0.4*
Gender:																
Male	4.7	4.8	3.6	3.5	4.0	3.3	4.4	4.5	2.9	2.8	3.5	2.1	2.6	§	1.7	-0.9
Female	5.6	5.1	4.3	3.7	2.9	4.1	4.2	5.4	5.7	3.8	4.9	5.0	4.5	§	4.6	+0.2
College Plans:																
None or under 4 years	5.4	7.7	3.4	4.6	3.8	1.6	3.7	7.5	4.1	3.1	4.2	2.3	2.9	§	2.9	0.0
Complete 4 years	5.1	4.3	3.9	3.3	3.5	4.4	3.7	4.1	4.7	3.3	4.4	4.0	3.7	§	3.4	-0.4
Region:																
Northeast	8.3	4.2	5.8	5.7	4.8	5.7	4.7	5.8	5.6	5.4	8.2	4.6	4.2	§	5.5	+1.3
Midwest	4.9	6.2	5.1	4.3	2.2	4.0	3.9	6.2	5.5	4.2	4.5	3.5	3.0	§	3.8	+0.8
South	4.4	4.7	2.7	2.7	4.1	3.4	3.9	4.5	4.6	2.8	3.5	4.6	4.0	§	1.7	-2.3 s
West	4.1	4.8	3.0	3.0	3.3	3.3	3.8	3.7	2.6	3.2	3.3	1.6	2.8	§	3.2	+0.4
Population Density:																
Large MSA	6.4	4.2	3.9	3.6	3.8	5.2	4.4	5.2	6.3	3.8	4.3	4.6	2.7	§	6.4	+3.7 s
Other MSA	5.3	5.4	4.2	4.2	4.0	3.5	3.2	4.8	3.4	2.6	5.1	2.4	4.0	§	0.8	-3.2 ss
Non-MSA	2.9	5.2	3.8	2.4	2.1	3.0	5.4	4.8	4.1	5.7	3.2	4.4	3.9	§	2.4	-1.5
Parental Education: ^b																
1.0–2.0 (Low)	2.3	5.1	4.3	2.4	2.5	2.1	2.2	3.6	4.2	4.3	1.5	4.0	0.7	§	0.0	-0.7
2.5–3.0	3.1	5.2	3.5	3.4	3.0	4.1	5.7	3.1	5.4	2.8	5.2	2.0	1.4	§	1.4	0.0
3.5–4.0	6.5	5.2	3.6	3.2	3.3	3.3	5.4	6.6	4.2	4.6	4.1	2.3	4.7	§	2.7	-2.0
4.5–5.0	5.3	3.9	5.3	4.5	3.6	5.2	3.6	3.7	3.4	2.9	4.4	4.8	3.6	§	4.4	+0.8
5.5–6.0 (High)	7.5	4.5	3.8	3.7	5.4	3.6	1.8	7.8	7.1	3.5	6.8	5.6	5.2	§	5.8	+0.6
Race/Ethnicity (2-year average): ^c																
White	5.2	4.9	4.1	4.1	4.0	3.6	3.7	5.1	5.8	4.4	4.6	4.8	4.0	§	3.7	-0.3
African American	2.6	3.8	4.5	4.3	3.6	4.0	4.4	4.1	2.9	3.7	3.1	1.5	3.2	§	4.0	+0.7
Hispanic	4.4	5.6	5.2	3.3	2.6	4.3	3.7	2.6	4.2	3.8	2.8	3.4	2.9	§	3.3	+0.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1976–1988; N is one fifth of N indicated in Table S3. Data based on one of six forms beginning in 1989; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aIn the 1988 questionnaires, a question on the use of wine coolers was added. This change may account for the discontinuity between the 1987 and 1988 use rates for wine.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 117
WINE COOLERS: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days													
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004– 2021
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	—
Total	14.8	15.1	16.1	15.9	15.3	17.5	15.9	15.6	16.5	14.8	15.3	12.9	12.4	—
Gender:														
Male	14.7	13.8	16.2	15.8	14.6	16.1	14.5	15.4	16.7	13.4	14.2	11.9	10.2	—
Female	14.6	16.2	16.0	16.2	15.6	18.7	16.9	15.8	16.3	15.9	15.9	13.3	14.3	—
College Plans:														
None or under 4 years	21.2	23.9	30.6	28.1	25.3	28.7	32.1	27.3	31.9	28.5	31.6	21.1	24.2	—
Complete 4 years	13.7	14.0	14.3	14.5	14.1	15.7	14.0	14.3	14.7	13.3	13.4	12.2	11.1	—
Region:														
Northeast	9.8	11.5	12.7	15.8	15.0	17.7	14.5	15.7	15.4	13.5	17.7	11.8	10.9	—
Midwest	14.5	14.6	15.7	15.0	13.9	18.5	16.1	16.2	17.6	17.0	14.8	12.1	15.5	—
South	17.4	18.1	17.8	16.7	17.0	17.5	16.6	16.2	18.6	14.6	16.8	15.0	12.7	—
West	14.8	14.2	17.3	15.9	13.9	16.0	15.3	14.0	12.7	13.3	11.4	11.3	8.9	—
Population Density:														
Large MSA	14.1	16.4	12.7	14.4	11.1	16.5	14.1	14.0	14.0	12.9	12.3	12.1	10.4	—
Other MSA	14.6	15.4	17.4	17.8	16.1	18.2	15.7	15.4	16.8	14.5	15.4	12.7	12.3	—
Non-MSA	15.8	13.4	17.3	13.2	15.8	17.3	17.9	17.6	18.9	17.5	18.3	14.4	14.7	—
Parental Education: ^a														
1.0–2.0 (Low)	19.3	18.7	18.5	21.1	20.4	21.7	22.2	20.7	30.2	22.1	25.1	22.8	18.2	—
2.5–3.0	16.9	16.6	19.9	17.7	17.6	20.2	19.2	17.8	19.7	18.1	19.7	14.3	15.4	—
3.5–4.0	12.9	16.8	17.2	18.4	15.7	18.4	16.8	16.6	17.5	14.3	19.6	15.7	15.8	—
4.5–5.0	13.9	11.8	13.5	12.0	12.5	15.4	11.9	13.6	13.1	11.8	9.0	9.8	8.1	—
5.5–6.0 (High)	13.5	12.5	11.9	14.7	13.2	13.8	13.5	14.2	11.4	12.1	8.2	9.0	8.7	—
Race/Ethnicity (2-year average): ^b														
White	—	14.4	15.1	15.1	14.9	16.4	16.5	15.4	16.0	15.9	15.0	14.0	12.6	—
African American	—	13.7	13.3	14.4	12.7	12.2	14.7	12.8	12.8	13.1	12.0	9.8	9.6	—
Hispanic	—	21.9	23.2	23.4	24.4	22.9	21.3	21.2	22.4	21.1	18.5	18.9	18.8	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996; N is one half of N indicated in Table S1. Data based on one of four forms beginning in 1997; N is one third of N indicated in Table S1. The category of wine coolers was dropped from the 8th- and 10th-grade questionnaires in 2004 to make space for a more general class of flavored alcoholic beverages.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE 118
WINE COOLERS: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days													
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004– 2021
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	—
Total	24.0	21.5	22.5	21.8	20.8	22.8	22.1	20.9	22.1	21.2	20.8	18.9	17.2	—
Gender:														
Male	23.9	19.8	20.5	21.9	19.4	21.8	20.6	19.0	20.5	19.6	19.8	16.0	14.8	—
Female	24.1	22.9	24.3	21.6	21.9	23.6	23.4	22.0	23.4	22.6	21.7	21.6	19.5	—
College Plans:														
None or under 4 years	31.1	26.7	27.6	28.4	28.9	33.0	28.4	29.3	33.4	32.6	31.6	30.2	23.8	—
Complete 4 years	22.5	20.5	21.4	20.4	19.4	21.0	21.1	19.3	20.3	19.5	19.1	17.2	16.2	—
Region:														
Northeast	23.2	19.0	24.1	21.6	18.5	20.3	21.0	19.1	21.7	18.8	21.5	18.1	18.7	—
Midwest	23.0	20.2	20.4	21.6	20.6	21.9	20.9	19.1	21.1	20.6	20.2	19.2	15.3	—
South	26.0	23.3	23.6	23.0	22.9	25.0	23.4	22.5	24.0	23.3	21.4	17.7	16.9	—
West	23.0	22.7	23.0	20.2	19.2	22.3	22.4	22.0	20.3	20.7	19.6	21.5	18.2	—
Population Density:														
Large MSA	23.1	21.6	21.5	21.2	16.4	19.3	20.2	16.9	23.1	19.8	19.9	17.4	17.0	—
Other MSA	23.8	22.3	22.4	22.6	20.7	23.7	22.2	21.9	21.3	19.7	19.7	18.7	16.4	—
Non-MSA	25.1	20.0	23.6	20.4	23.7	24.6	23.8	23.2	22.6	25.6	23.7	21.2	19.5	—
Parental Education: ^a														
1.0–2.0 (Low)	25.9	24.4	27.4	24.4	27.7	25.4	21.1	22.6	26.3	23.8	22.2	20.3	23.4	—
2.5–3.0	26.8	21.3	22.3	22.4	23.8	25.5	24.3	25.4	25.1	24.2	22.7	22.5	19.8	—
3.5–4.0	24.8	22.5	22.5	22.3	22.1	23.7	23.6	22.0	22.3	23.6	22.2	19.1	17.5	—
4.5–5.0	21.8	20.6	21.9	21.4	19.1	19.9	21.2	15.6	17.5	19.4	17.7	18.2	15.4	—
5.5–6.0 (High)	20.7	20.5	20.2	17.8	14.5	21.3	18.5	18.5	21.2	14.2	20.3	13.7	13.3	—
Race/Ethnicity (2-year average): ^b														
White	—	23.0	21.9	22.1	21.2	21.7	22.6	21.7	21.6	21.9	21.2	20.3	18.1	—
African American	—	19.9	19.3	20.6	20.0	16.6	16.1	17.6	17.0	18.1	18.3	15.3	15.2	—
Hispanic	—	26.3	26.2	24.7	24.9	28.1	28.0	24.1	24.5	25.2	24.5	22.3	21.7	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases.

See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996; N is one half of N indicated in Table S2. Data based on one of four forms beginning in 1997; N is one third of N indicated in Table S2. The category of wine coolers was dropped from the 8th- and 10th-grade questionnaires in 2004 to make space for a more general class of flavored alcoholic beverages.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE 119
WINE COOLERS: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	16,300	16,700	15,200
Total	—	—	—	—	—	—	—	—	—	—	—	—	—	36.6	29.4	26.2
Gender:																
Male	—	—	—	—	—	—	—	—	—	—	—	—	—	30.8	25.5	22.7
Female	—	—	—	—	—	—	—	—	—	—	—	—	—	41.5	33.6	29.8
College Plans:																
None or under 4 years	—	—	—	—	—	—	—	—	—	—	—	—	—	38.2	30.9	29.2
Complete 4 years	—	—	—	—	—	—	—	—	—	—	—	—	—	35.9	28.6	24.9
Region:																
Northeast	—	—	—	—	—	—	—	—	—	—	—	—	—	33.8	29.6	29.2
Midwest	—	—	—	—	—	—	—	—	—	—	—	—	—	37.9	33.0	28.6
South	—	—	—	—	—	—	—	—	—	—	—	—	—	33.6	25.7	22.9
West	—	—	—	—	—	—	—	—	—	—	—	—	—	42.7	31.3	24.7
Population Density:																
Large MSA	—	—	—	—	—	—	—	—	—	—	—	—	—	37.7	28.6	25.9
Other MSA	—	—	—	—	—	—	—	—	—	—	—	—	—	37.1	31.1	28.7
Non-MSA	—	—	—	—	—	—	—	—	—	—	—	—	—	34.6	26.4	21.1
Parental Education: ^a																
1.0–2.0 (Low)	—	—	—	—	—	—	—	—	—	—	—	—	—	28.8	27.7	22.9
2.5–3.0	—	—	—	—	—	—	—	—	—	—	—	—	—	37.8	31.9	28.1
3.5–4.0	—	—	—	—	—	—	—	—	—	—	—	—	—	38.5	31.2	27.1
4.5–5.0	—	—	—	—	—	—	—	—	—	—	—	—	—	35.6	27.6	27.6
5.5–6.0 (High)	—	—	—	—	—	—	—	—	—	—	—	—	—	36.9	23.5	21.5
Race/Ethnicity (2-year average): ^b																
White	—	—	—	—	—	—	—	—	—	—	—	—	—	—	35.4	29.6
African American	—	—	—	—	—	—	—	—	—	—	—	—	—	—	25.3	21.7
Hispanic	—	—	—	—	—	—	—	—	—	—	—	—	—	—	30.2	23.3

Table continued on next page.

TABLE 119 (cont.)
WINE COOLERS: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	24.3	21.9	19.4	22.2	20.6	20.7	20.7	20.1	18.4	17.6	19.0	19.3	14.6	14.4	14.1	13.3
Gender:																
Male	22.1	19.1	15.1	16.9	15.5	17.7	15.9	15.0	15.6	15.6	14.1	12.0	9.4	11.2	8.7	9.2
Female	26.3	24.3	22.8	26.4	25.1	23.6	24.4	23.5	20.6	19.5	22.2	24.2	18.9	17.2	18.7	17.1
College Plans:																
None or under 4 years	29.3	21.9	20.5	22.8	22.7	20.0	17.8	21.8	22.8	21.2	23.1	21.8	17.2	16.7	18.6	17.5
Complete 4 years	22.8	22.0	19.2	22.3	19.8	20.9	21.6	19.6	17.3	16.1	17.7	18.8	14.0	13.7	13.2	12.3
Region:																
Northeast	21.0	19.6	18.9	15.4	14.6	21.1	17.2	20.8	22.1	19.1	18.4	17.8	15.9	14.7	17.3	13.3
Midwest	26.1	23.7	19.4	26.0	19.8	20.9	21.5	18.6	19.2	18.3	19.5	20.9	15.5	11.0	13.7	13.2
South	23.1	21.3	20.6	23.4	24.1	23.2	22.9	20.6	18.7	16.8	19.7	20.6	13.9	16.7	13.5	14.6
West	26.3	22.2	17.8	19.8	20.4	14.9	18.9	20.4	13.7	16.7	17.5	16.6	13.1	14.6	12.7	10.9
Population Density:																
Large MSA	20.9	21.2	20.4	21.2	20.7	22.7	23.9	22.5	14.7	19.8	18.1	20.5	12.8	14.2	13.5	10.7
Other MSA	28.1	21.0	19.0	23.3	19.8	19.3	19.1	19.4	17.5	15.0	18.5	18.4	13.7	13.8	13.2	14.4
Non-MSA	20.6	24.1	19.4	21.3	21.8	21.6	20.1	18.7	23.4	19.4	20.6	19.4	18.0	15.6	16.7	14.7
Parental Education: ^a																
1.0–2.0 (Low)	24.0	25.7	20.8	23.5	21.1	18.2	13.2	19.8	19.0	24.8	19.3	18.4	13.7	21.3	15.6	12.8
2.5–3.0	25.4	21.2	17.1	26.4	20.4	26.8	20.0	21.4	23.2	16.3	22.8	19.6	18.1	17.2	19.8	15.6
3.5–4.0	27.2	22.9	24.4	20.9	22.0	19.5	23.0	23.3	17.3	21.8	18.5	17.8	14.7	16.4	12.6	15.3
4.5–5.0	22.1	18.0	17.2	22.8	21.0	17.0	21.4	17.9	14.3	13.6	19.0	19.4	12.4	9.1	13.6	12.7
5.5–6.0 (High)	18.0	24.5	16.4	14.6	16.7	20.8	21.0	16.6	19.9	16.8	14.6	23.5	10.1	13.3	9.8	9.8
Race/Ethnicity (2-year average): ^b																
White	26.7	23.7	21.3	21.5	21.9	20.8	21.2	21.0	19.6	18.0	18.2	19.7	16.8	13.1	13.6	14.6
African American	20.9	18.2	17.6	18.3	17.6	22.1	21.1	18.4	15.5	12.4	17.0	17.0	15.8	19.4	18.4	10.9
Hispanic	22.3	26.7	22.9	22.9	26.0	22.0	19.8	18.2	17.4	20.7	20.0	16.7	16.0	19.7	19.4	13.8

Table continued on next page.

TABLE 119 (cont.)
WINE COOLERS: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	12.7	12.8	12.2	12.4	10.0	10.7	11.3	11.3	9.8	10.5	11.4	9.7	10.5	§	9.1	-1.4*
Gender:																
Male	9.2	8.3	7.6	9.1	6.1	6.7	6.6	7.0	6.0	6.4	7.5	6.0	8.0	§	4.5	-3.5 s
Female	16.1	16.6	16.0	16.2	13.4	14.6	15.6	15.0	12.9	13.9	15.2	13.0	12.5	§	13.6	+1.1
College Plans:																
None or under 4 years	18.0	16.8	15.1	18.6	12.7	10.8	13.7	12.4	9.3	9.5	15.3	6.9	13.3	§	10.5	-2.8
Complete 4 years	11.5	12.0	11.4	11.3	9.5	10.8	10.8	10.8	10.2	10.5	10.6	10.7	9.8	§	8.9	-0.9
Region:																
Northeast	15.2	11.7	11.7	15.9	14.8	13.5	14.0	12.6	12.4	11.9	11.9	11.6	10.7	§	13.7	+3.0
Midwest	11.9	12.3	14.4	12.3	9.5	10.9	12.7	11.6	8.1	12.3	13.6	13.8	12.2	§	9.0	-3.2
South	13.0	15.1	10.1	12.8	9.8	9.4	9.3	12.6	10.0	9.7	10.8	10.3	11.0	§	8.1	-2.9
West	10.9	10.8	13.1	9.4	7.7	10.6	10.9	7.8	9.4	9.4	9.6	3.9	6.4	§	7.1	+0.8
Population Density:																
Large MSA	11.9	11.4	11.9	12.2	11.6	11.8	12.2	10.7	9.5	8.6	10.3	9.1	9.1	§	10.8	+1.7
Other MSA	12.6	12.8	12.8	12.8	8.5	9.5	10.4	10.1	9.5	9.7	10.1	8.4	9.8	§	5.4	-4.4
Non-MSA	13.9	14.3	11.3	11.8	11.4	12.2	12.3	14.7	11.2	15.1	15.4	13.8	14.6	§	12.8	-1.8
Parental Education: ^a																
1.0–2.0 (Low)	18.4	17.3	17.4	13.5	12.8	10.6	11.0	10.0	13.0	9.9	8.4	5.9	6.3	§	9.8	+3.6
2.5–3.0	13.1	12.8	11.1	16.7	15.5	10.2	13.1	15.0	10.5	8.6	11.1	10.8	9.4	§	9.8	+0.4
3.5–4.0	13.5	12.3	14.5	11.6	7.7	11.9	12.7	11.3	9.8	12.0	14.7	9.2	9.1	§	10.7	+1.6
4.5–5.0	10.8	11.1	11.0	10.9	8.5	12.3	11.0	9.0	8.8	11.2	11.9	11.6	13.3	§	11.4	-1.8
5.5–6.0 (High)	11.7	12.5	7.1	10.7	8.2	6.5	9.6	11.9	9.0	7.5	7.8	11.8	12.1	§	3.6	-8.4 ss
Race/Ethnicity (2-year average): ^b																
White	13.1	11.4	11.1	12.5	12.0	10.8	11.2	11.6	11.1	10.8	12.5	13.7	13.3	§	10.0	-3.3
African American	11.8	16.8	18.5	13.0	8.3	8.9	10.9	10.4	8.9	10.0	8.9	6.8	7.5	§	8.3	+0.9
Hispanic	13.0	12.9	13.2	12.7	10.5	9.7	11.2	11.0	10.0	8.7	8.8	7.0	5.1	§	5.9	+0.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1988; *N* is one fifth of *N* indicated in Table S3. Data based on one of six forms beginning in 1989; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 120
WINE COOLERS: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	16,300	16,700	15,200
Total	—	—	—	—	—	—	—	—	—	—	—	—	—	13.9	12.4	10.5
Gender:																
Male	—	—	—	—	—	—	—	—	—	—	—	—	—	12.1	11.3	9.9
Female	—	—	—	—	—	—	—	—	—	—	—	—	—	15.1	13.5	11.3
College Plans:																
None or under 4 years	—	—	—	—	—	—	—	—	—	—	—	—	—	17.7	16.2	14.2
Complete 4 years	—	—	—	—	—	—	—	—	—	—	—	—	—	12.5	10.7	9.0
Region:																
Northeast	—	—	—	—	—	—	—	—	—	—	—	—	—	14.7	14.3	10.0
Midwest	—	—	—	—	—	—	—	—	—	—	—	—	—	14.2	13.1	12.0
South	—	—	—	—	—	—	—	—	—	—	—	—	—	11.2	11.2	9.3
West	—	—	—	—	—	—	—	—	—	—	—	—	—	17.4	12.0	11.0
Population Density:																
Large MSA	—	—	—	—	—	—	—	—	—	—	—	—	—	14.2	11.7	10.0
Other MSA	—	—	—	—	—	—	—	—	—	—	—	—	—	15.6	12.4	11.0
Non-MSA	—	—	—	—	—	—	—	—	—	—	—	—	—	10.7	13.3	9.9
Parental Education: ^a																
1.0–2.0 (Low)	—	—	—	—	—	—	—	—	—	—	—	—	—	13.9	9.2	10.7
2.5–3.0	—	—	—	—	—	—	—	—	—	—	—	—	—	16.6	13.8	13.3
3.5–4.0	—	—	—	—	—	—	—	—	—	—	—	—	—	13.6	14.9	9.3
4.5–5.0	—	—	—	—	—	—	—	—	—	—	—	—	—	11.7	10.5	11.3
5.5–6.0 (High)	—	—	—	—	—	—	—	—	—	—	—	—	—	11.4	7.4	6.3
Race/Ethnicity (2-year average): ^b																
White	—	—	—	—	—	—	—	—	—	—	—	—	—	—	13.6	12.3
African American	—	—	—	—	—	—	—	—	—	—	—	—	—	—	13.0	9.9
Hispanic	—	—	—	—	—	—	—	—	—	—	—	—	—	—	11.5	6.0

Table continued on next page.

TABLE 120 (cont.)
WINE COOLERS: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	10.2	10.4	8.5	8.9	9.0	8.4	9.4	9.5	7.7	8.8	9.7	10.0	7.1	7.5	7.2	7.1
Gender:																
Male	10.0	10.2	7.3	7.4	6.7	6.9	6.7	7.0	7.0	9.4	6.2	7.2	5.4	7.5	4.2	6.1
Female	10.2	10.3	9.5	10.0	10.8	9.7	11.3	11.0	7.8	8.4	11.4	11.4	8.2	8.0	9.7	8.0
College Plans:																
None or under 4 years	13.5	11.1	10.8	10.9	12.1	8.8	10.1	9.9	10.2	12.1	14.8	13.7	8.9	10.8	11.4	11.9
Complete 4 years	9.2	10.1	8.0	8.6	8.0	8.3	9.2	9.5	7.0	7.4	8.2	9.2	6.7	6.8	6.2	6.0
Region:																
Northeast	7.6	8.8	6.5	5.5	5.5	9.8	8.2	9.6	8.9	10.1	10.2	9.8	8.0	7.5	10.7	6.9
Midwest	10.9	12.0	8.0	9.9	7.0	9.2	9.7	7.3	7.0	10.8	8.9	12.0	8.8	6.4	6.6	7.3
South	8.8	10.3	11.1	9.3	11.5	8.7	9.7	10.5	8.8	5.9	10.9	9.6	6.7	8.6	6.6	8.3
West	13.2	9.7	6.2	10.0	10.4	4.8	9.7	10.2	5.6	10.1	8.4	8.4	5.0	7.1	5.7	5.1
Population Density:																
Large MSA	7.9	11.8	7.9	8.5	11.1	8.8	10.8	12.1	7.5	11.1	10.6	9.6	6.8	7.2	7.0	6.4
Other MSA	12.4	9.2	8.7	8.8	7.5	8.3	9.5	8.7	6.5	8.1	8.2	9.9	6.3	7.0	7.4	6.6
Non-MSA	8.2	11.7	8.6	9.6	9.1	8.2	7.8	7.9	9.7	7.4	11.3	10.7	8.9	8.9	7.1	9.1
Parental Education: ^a																
1.0–2.0 (Low)	12.5	13.1	12.2	11.4	11.7	9.6	6.8	13.2	8.3	16.8	14.1	14.6	9.3	11.9	8.3	12.0
2.5–3.0	10.9	9.5	6.9	10.4	9.5	11.4	9.8	11.0	9.2	8.8	12.6	9.9	9.4	9.5	9.9	8.2
3.5–4.0	12.0	11.1	11.1	8.8	8.5	6.8	10.6	9.9	7.7	8.8	7.3	9.9	6.1	9.1	6.2	7.4
4.5–5.0	9.2	8.8	7.4	9.2	9.2	7.2	8.6	7.7	7.2	8.6	10.7	10.7	5.9	4.2	6.8	7.4
5.5–6.0 (High)	3.7	10.6	5.6	4.8	7.5	7.7	9.5	7.9	5.8	7.0	6.1	8.0	5.9	5.5	5.2	3.4
Race/Ethnicity (2-year average): ^b																
White	10.7	10.3	9.5	8.6	8.6	8.1	8.7	8.8	7.6	7.6	8.9	10.2	8.4	6.4	6.9	7.5
African American	9.2	8.7	8.6	8.6	7.5	10.4	10.0	9.2	6.9	4.2	9.6	10.0	7.8	11.7	10.5	4.6
Hispanic	9.8	14.5	13.5	13.1	13.9	11.1	9.8	12.0	12.6	13.2	11.5	9.1	9.3	10.6	9.4	8.0

Table continued on next page.

TABLE 120 (cont.)
WINE COOLERS: Trends in 2-Week Prevalence of 5 or More Drinks in a Row by Subgroups in Grade 12

	Percentage reporting 5+ drinks in a row on one or more occasions															2019-2021
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	5.8	7.3	6.2	6.4	4.9	5.4	5.3	5.8	5.0	4.3	6.3	5.0	4.9	§	4.2	-0.7*
Gender:																
Male	4.8	4.5	3.5	5.7	3.6	3.0	4.1	4.1	2.5	2.2	4.4	2.8	4.1	§	2.7	-1.4
Female	7.0	9.6	8.7	7.0	6.1	7.2	6.4	7.2	6.8	5.6	8.1	6.5	5.6	§	5.9	+0.3
College Plans:																
None or under 4 years	11.1	8.2	9.4	9.9	6.2	6.7	7.9	6.2	5.4	5.2	8.8	6.9	5.8	§	5.5	-0.3
Complete 4 years	4.7	7.0	5.6	5.6	4.5	5.1	4.6	5.4	5.0	3.9	5.7	4.6	4.7	§	4.0	-0.7
Region:																
Northeast	6.1	7.4	5.9	7.5	9.4	6.1	6.0	5.0	4.5	5.5	8.6	5.2	5.8	§	6.4	+0.6
Midwest	6.0	8.1	7.8	6.1	5.0	4.6	6.9	5.4	4.6	7.7	8.1	6.5	4.8	§	3.6	-1.2
South	5.7	8.4	4.9	7.3	4.6	6.1	3.9	6.5	5.5	2.8	5.2	5.3	5.7	§	3.6	-2.1
West	5.5	4.6	6.7	4.3	2.5	4.9	5.0	5.6	5.1	3.1	4.9	3.2	2.5	§	4.0	+1.4
Population Density:																
Large MSA	5.3	6.3	4.8	7.1	4.9	6.8	6.3	5.3	4.8	3.6	6.7	5.8	4.4	§	5.2	+0.7
Other MSA	6.4	6.7	6.6	6.5	4.0	3.8	4.0	5.1	5.3	3.9	6.0	4.0	4.2	§	2.2	-2.0
Non-MSA	5.3	9.4	7.2	4.9	7.2	6.8	7.0	7.7	4.8	6.3	6.6	6.1	7.7	§	6.3	-1.3
Parental Education: ^a																
1.0–2.0 (Low)	8.8	9.1	9.7	7.5	8.4	4.9	8.3	5.5	7.6	1.0	4.1	7.5	1.6	§	5.3	+3.7
2.5–3.0	8.1	8.0	5.4	8.5	7.9	6.7	4.9	6.4	5.3	3.7	8.0	5.4	5.7	§	3.7	-2.0
3.5–4.0	5.6	6.8	8.6	5.6	4.0	5.0	6.6	6.9	4.6	5.8	8.2	2.7	4.3	§	7.1	+2.8
4.5–5.0	4.7	5.5	4.2	5.3	3.4	5.7	4.3	4.5	4.5	5.6	6.3	4.6	5.7	§	4.3	-1.4
5.5–6.0 (High)	4.1	8.3	3.8	5.2	3.5	3.0	4.6	6.0	5.3	2.5	3.6	7.0	5.0	§	1.2	-3.8 s
Race/Ethnicity (2-year average): ^b																
White	6.4	5.4	5.5	6.1	6.1	5.3	5.4	6.4	5.9	4.7	6.3	6.9	5.6	§	3.8	-1.8
African American	4.9	6.6	8.2	7.2	4.3	3.9	4.7	3.8	4.4	6.1	3.4	1.8	3.4	§	5.4	+2.0
Hispanic	7.9	8.1	8.8	7.4	5.6	5.4	5.8	5.2	5.7	4.1	4.5	5.7	4.0	§	4.2	+0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of five forms in 1988; *N* is one fifth of *N* indicated in Table S3. Data based on one of six forms beginning in 1989; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 121
FLAVORED ALCOHOLIC BEVERAGES:
Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days																			2019– 2021 change
	1991– 2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	14.6	12.9	13.1	12.2	10.2	9.5	9.4	8.6	7.6	6.3	5.7	5.5	4.0	4.4	4.9	4.5	6.6	4.6	+0.1*
Gender:																				
Male	—	12.7	11.4	10.8	10.4	9.2	9.1	8.3	8.3	6.1	5.8	5.1	5.2	3.5	3.7	3.9	3.7	§	3.7	-0.1
Female	—	16.3	14.4	15.3	13.4	10.9	9.7	10.3	8.6	9.1	6.7	5.9	5.8	4.5	5.2	6.1	5.3	§	5.4	+0.1
College Plans:																				
None or under 4 years	—	28.1	24.9	25.9	23.8	24.0	18.3	19.6	14.4	19.9	14.3	13.8	10.4	7.5	9.8	6.0	7.5	§	5.9	-1.6
Complete 4 years	—	13.3	11.8	11.8	11.1	9.0	8.7	8.6	8.3	6.7	5.8	5.1	5.2	3.7	3.8	4.9	4.2	§	4.2	0.0
Region:																				
Northeast	—	14.9	11.0	12.2	9.7	9.2	8.8	9.2	5.4	7.2	4.7	4.9	4.2	2.3	2.0	4.8	2.7	§	3.7	+0.9
Midwest	—	16.2	12.6	12.9	12.0	10.5	9.1	8.2	7.8	5.2	6.3	5.6	5.5	4.0	4.4	4.7	5.1	§	6.3	+1.3
South	—	15.1	14.6	14.3	14.0	12.6	11.8	11.3	9.6	9.2	7.4	4.7	6.6	4.8	4.5	5.1	4.6	§	4.2	-0.4
West	—	11.7	11.9	12.0	10.9	7.2	6.7	7.6	10.4	7.6	5.9	7.8	4.5	3.7	5.9	5.0	5.1	§	4.3	-0.8
Population Density:																				
Large MSA	—	14.4	11.5	12.6	10.6	11.1	7.2	8.5	6.4	6.8	5.7	4.9	4.5	3.2	3.8	4.0	3.2	§	2.4	-0.8
Other MSA	—	14.2	14.0	13.0	12.4	9.7	9.7	10.0	9.7	8.0	7.2	5.3	6.0	4.0	4.5	5.6	4.1	§	5.2	+1.1
Non-MSA	—	15.6	12.8	13.9	14.2	10.2	12.0	9.0	9.6	8.2	5.5	7.4	5.8	5.1	4.9	4.9	7.7	§	6.8	-0.8
Parental Education: ^a																				
1.0–2.0 (Low)	—	19.5	19.5	18.7	21.1	19.3	14.7	11.2	12.7	11.7	12.3	9.5	11.5	10.0	7.7	6.0	4.4	§	2.0	-2.5
2.5–3.0	—	17.5	18.0	18.7	17.5	14.4	13.1	13.4	12.1	11.7	8.7	8.4	5.2	4.4	5.1	6.4	9.1	§	6.0	-3.1
3.5–4.0	—	17.8	16.2	13.4	12.3	11.7	11.4	10.5	10.3	7.8	7.9	7.0	6.4	3.4	4.9	5.6	6.1	§	6.3	+0.2
4.5–5.0	—	11.8	10.3	8.5	9.8	7.2	7.4	8.0	6.2	5.4	5.6	3.3	3.8	3.6	3.4	4.5	3.1	§	4.9	+1.8
5.5–6.0 (High)	—	11.5	7.3	9.9	7.1	6.5	4.7	4.9	6.2	4.7	2.8	4.1	4.0	2.1	3.1	4.5	3.3	§	3.1	-0.2
Race/Ethnicity (2-year average): ^b																				
White	—	—	13.8	12.6	12.1	11.1	9.8	9.0	8.5	7.4	6.4	5.5	5.1	4.4	3.7	4.2	4.9	§	5.7	+0.8
African American	—	—	12.0	11.0	10.1	9.2	8.5	8.0	7.7	8.8	7.3	4.2	4.3	3.9	3.8	3.2	2.3	§	4.8	+2.4
Hispanic	—	—	17.0	17.6	16.7	13.9	11.8	11.7	11.4	10.8	9.1	7.5	7.1	6.6	5.5	5.6	5.4	§	3.3	-2.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S1. In 2011 Sky Blue and Zima were deleted from the list of examples.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 122
FLAVORED ALCOHOLIC BEVERAGES:
Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days																			2019– 2021 change	
	1991– 2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Approximate weighted N =	—	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000		
Total	—	25.1	23.1	24.7	21.8	20.2	19.0	19.4	15.8	16.3	15.5	14.0	12.8	11.0	12.9	11.8	11.1	12.5	7.8	-3.3* ss	
Gender																					
Male	—	24.5	20.3	22.4	18.8	17.0	17.0	18.8	13.8	13.5	13.7	12.4	10.5	8.4	9.9	11.2	10.2	§	5.7	-4.4 sss	
Female	—	25.6	25.8	26.9	24.7	22.9	20.9	19.8	17.7	18.5	16.7	15.7	14.9	13.4	15.7	12.5	11.9	§	9.9	-2.0	
College Plans:																					
None or under 4 years	—	32.5	33.8	37.2	32.0	29.1	30.0	28.6	25.7	21.0	19.1	20.5	18.9	17.2	17.4	14.5	13.8	§	8.1	-5.6 s	
Complete 4 years	—	24.2	21.7	23.1	20.6	19.1	17.8	18.4	14.7	15.8	15.0	13.3	12.3	10.3	12.2	11.6	10.7	§	7.7	-3.0 s	
Region:																					
Northeast	—	26.6	22.4	26.3	23.2	22.2	18.6	18.3	17.5	19.4	15.6	14.4	12.4	11.0	13.9	13.4	12.7	§	10.1	-2.6	
Midwest	—	22.5	23.4	23.5	21.2	20.0	17.6	19.1	14.3	14.5	13.9	12.3	12.6	9.0	11.0	10.6	13.0	§	8.2	-4.7 s	
South	—	25.6	23.7	24.5	20.0	20.3	21.1	19.6	16.6	17.4	17.2	14.5	15.0	12.6	14.1	13.5	10.3	§	7.5	-2.8	
West	—	26.5	22.5	25.4	23.6	19.0	17.9	20.3	15.1	14.0	15.0	14.8	9.7	10.6	12.5	9.1	9.6	§	6.1	-3.5	
Population Density:																					
Large MSA	—	22.4	22.0	23.1	21.5	19.0	17.6	20.2	15.3	17.4	13.8	13.7	12.3	10.5	13.7	11.2	11.0	§	5.2	-5.8 ss	
Other MSA	—	26.1	23.1	25.5	21.4	20.9	19.6	19.0	15.8	16.2	16.4	14.5	12.4	9.7	12.5	12.5	10.7	§	9.5	-1.3	
Non-MSA	—	26.5	24.4	25.5	23.0	20.5	19.7	18.9	16.5	14.7	15.7	13.2	14.5	15.6	13.0	11.2	12.3	§	8.6	-3.7	
Parental Education: ^a																					
1.0–2.0 (Low)	—	27.3	24.9	25.8	25.5	21.8	23.5	26.4	17.6	17.0	16.4	14.6	13.6	9.3	15.5	14.8	13.5	§	11.2	-2.4	
2.5–3.0	—	30.5	28.1	27.9	25.8	23.4	23.3	24.5	19.2	17.3	19.9	16.6	17.7	14.3	13.7	11.2	14.8	§	7.0	-7.8 ss	
3.5–4.0	—	25.0	23.6	28.5	23.1	21.2	18.8	20.2	17.5	15.6	16.3	16.2	13.6	11.4	13.0	12.6	11.1	§	6.4	-4.7 ss	
4.5–5.0	—	24.9	22.6	19.9	19.7	19.0	18.2	15.7	13.6	16.7	14.6	11.5	11.2	12.1	12.5	11.4	9.4	§	7.1	-2.3	
5.5–6.0 (High)	—	20.8	17.3	23.7	16.9	16.2	14.3	15.4	12.3	17.5	10.5	13.3	10.9	7.8	12.3	10.3	10.4	§	11.8	+1.3	
Race/Ethnicity (2-year average): ^b																					
White	—	—	25.7	25.1	24.0	21.8	20.0	19.0	17.4	16.5	16.5	15.8	14.8	13.2	13.2	14.1	13.6	§	11.2	-2.4	
African American	—	—	16.1	15.5	15.9	15.2	14.2	14.8	14.0	10.7	9.6	8.1	8.1	8.4	7.0	6.5	6.2	§	7.6	+1.4	
Hispanic	—	—	29.0	28.6	26.1	23.1	23.0	24.2	22.6	17.7	16.7	16.1	14.3	12.5	12.2	12.5	11.7	§	8.8	-2.9	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S2. In 2011 Skyy Blue and Zima were deleted from the list of examples.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 123
FLAVORED ALCOHOLIC BEVERAGES:
Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days																			2019– 2021 change
	1991– 2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	31.1	30.5	29.3	29.1	27.4	27.4	24.1	23.1	21.8	21.0	19.9	20.8	18.3	20.2	18.1	18.5	§	15.3	-3.2*
Gender:																				
Male	—	28.9	28.3	26.0	25.4	25.8	27.3	22.6	21.8	19.9	20.1	16.8	19.1	17.4	17.7	18.2	17.8	§	12.1	-5.7 s
Female	—	33.2	33.2	32.3	32.8	29.0	27.1	25.0	24.6	23.3	22.0	21.9	22.6	18.6	21.7	18.1	18.5	§	19.5	+1.1
College Plans:																				
None or under 4 years	—	34.7	34.2	32.9	37.0	29.0	35.6	26.8	26.2	22.6	25.8	22.8	24.8	18.1	20.3	20.3	16.6	§	9.3	-7.3 s
Complete 4 years	—	30.3	29.6	28.7	27.5	26.9	25.9	23.9	22.4	21.5	19.9	19.3	20.0	18.1	20.3	17.9	19.2	§	17.2	-2.0
Region:																				
Northeast	—	33.5	38.1	29.4	36.0	33.5	34.0	31.3	25.7	20.6	23.6	22.6	24.9	21.8	22.9	20.1	17.3	§	25.9	+8.6
Midwest	—	35.3	33.4	32.2	28.5	25.8	26.7	23.7	23.1	22.4	18.3	24.3	19.9	18.3	25.1	17.7	21.5	§	13.3	-8.3 s
South	—	25.9	25.9	27.0	28.8	26.9	27.2	23.3	22.1	19.3	19.9	17.6	21.3	16.9	20.0	18.3	16.6	§	11.7	-4.9
West	—	32.1	27.9	29.9	24.9	25.8	23.2	20.9	22.9	24.8	23.9	17.6	18.0	18.5	14.5	16.8	19.9	§	15.1	-4.7
Population Density:																				
Large MSA	—	27.0	29.9	32.3	30.8	28.7	27.0	26.4	25.3	22.5	23.4	18.7	21.9	18.2	17.9	16.7	14.5	§	20.0	+5.5
Other MSA	—	33.4	29.9	27.7	27.8	25.8	28.8	23.8	23.8	22.2	20.0	19.9	19.3	17.0	19.1	17.7	17.9	§	12.1	-5.8
Non-MSA	—	30.9	32.2	28.5	29.5	29.1	24.7	21.8	18.6	19.5	20.1	21.2	22.2	21.2	26.5	21.3	26.4	§	14.8	-11.5 s
Parental Education: ^a																				
1.0–2.0 (Low)	—	28.5	22.2	30.3	34.0	24.9	26.8	13.8	24.8	19.7	21.8	21.0	20.9	11.9	19.0	13.0	13.7	§	13.0	-0.7
2.5–3.0	—	31.0	33.3	30.6	32.4	28.3	28.2	27.8	26.1	24.0	24.8	21.5	19.0	20.5	19.5	18.9	17.1	§	10.6	-6.5
3.5–4.0	—	33.6	31.7	30.9	29.9	28.4	28.7	24.0	25.3	22.9	20.7	22.0	22.6	17.3	21.4	18.0	20.6	§	11.7	-8.9 ss
4.5–5.0	—	29.4	30.2	28.8	27.6	27.6	24.4	26.1	20.0	21.7	20.0	16.1	23.9	21.1	19.1	22.6	19.6	§	20.3	+0.7
5.5–6.0 (High)	—	29.8	30.2	26.4	25.3	26.6	28.4	23.4	20.5	19.7	19.1	19.9	17.8	15.9	24.0	18.3	24.3	§	21.7	-2.6
Race/Ethnicity (2-year average): ^b																				
White	—	—	33.3	31.4	30.6	31.1	29.9	27.0	24.6	22.9	22.2	22.1	22.4	22.7	23.5	23.4	21.9	§	19.7	-2.3
African American	—	—	17.1	20.9	21.6	19.8	19.3	20.0	20.0	17.2	14.2	16.1	14.1	10.9	10.5	13.5	12.8	§	9.6	-3.3
Hispanic	—	—	28.6	32.2	34.5	26.2	25.2	24.7	25.5	26.8	21.9	16.3	18.6	21.4	16.5	13.9	14.6	§	10.8	-3.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; *N* is one sixth of *N* indicated in Table S3. In 2011 Skyy Blue and Zima were deleted from the list of examples.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 124
ALCOHOLIC BEVERAGES CONTAINING CAFFEINE:
Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months												2019– 2021
	1991– 2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	—	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	11.8‡	10.9	10.2	9.5	8.4	6.5	5.6	6.0	7.3	5.7	6.2	-1.1*
Gender:													
Male	—	9.4‡	9.6	8.4	9.2	7.6	6.1	4.6	5.4	7.3	§	4.4	-2.9 s
Female	—	13.8‡	11.9	11.7	9.6	8.9	6.9	6.4	6.7	7.5	§	8.2	+0.6
College Plans:													
None or under 4 years	—	22.3‡	19.1	21.6	22.9	19.3	15.8	10.4	11.9	13.3	§	10.0	-3.3
Complete 4 years	—	11.0‡	10.2	9.4	8.4	7.6	5.7	5.2	5.3	6.6	§	5.5	-1.2
Region:													
Northeast	—	10.4‡	8.7	11.5	6.7	7.4	4.2	2.6	5.5	5.6	§	3.5	-2.1
Midwest	—	10.3‡	8.6	8.7	10.3	8.6	6.7	4.4	5.1	7.0	§	6.9	-0.1
South	—	12.9‡	12.4	11.4	9.0	7.7	6.6	6.1	5.6	7.6	§	7.1	-0.5
West	—	12.5‡	12.3	9.1	11.3	10.3	7.9	7.9	8.2	8.4	§	5.9	-2.4
Population Density:													
Large MSA	—	9.8‡	10.6	9.6	7.8	7.3	5.3	5.7	7.0	5.9	§	4.7	-1.2
Other MSA	—	13.4‡	10.9	10.5	9.6	7.8	6.6	5.3	5.9	7.6	§	6.3	-1.3
Non-MSA	—	11.4‡	11.3	10.5	11.4	11.1	8.4	6.1	4.7	8.6	§	8.3	-0.3
Parental Education: ^a													
1.0–2.0 (Low)	—	19.0‡	13.8	13.4	13.8	12.5	9.9	8.7	8.7	12.7	§	4.4	-8.4 ss
2.5–3.0	—	17.3‡	14.4	11.1	12.6	8.4	7.4	6.6	8.2	12.0	§	10.4	-1.5
3.5–4.0	—	13.8‡	13.3	13.1	10.9	11.5	8.7	7.2	6.6	7.9	§	11.0	+3.1
4.5–5.0	—	8.3‡	8.9	8.2	7.1	6.5	5.4	3.4	4.8	4.7	§	4.3	-0.4
5.5–6.0 (High)	—	7.0‡	6.9	8.8	6.5	4.8	4.8	4.4	4.2	5.3	§	4.8	-0.5
Race/Ethnicity (2-year average): ^b													
White	—	—	—	9.0	9.6	9.2	7.2	5.3	4.8	6.3	§	6.7	+0.4
African American	—	—	—	7.3	5.9	5.6	4.8	3.7	4.2	5.7	§	2.9	-2.7
Hispanic	—	—	—	14.5	11.5	9.4	8.5	7.5	7.1	8.2	§	2.4	-5.7 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '‡' indicates some change in the question. See relevant footnote for that drug. See relevant figure to assess the impact of the wording changes. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. ¹In 2012 the question text was changed from "...had an alcoholic beverage containing caffeine (like Four Loko or Joose)" to "...had an alcoholic beverage mixed with an energy drink (like Red Bull)." The data in 2011 and 2012 are not comparable due to this question change. Data based on one of four forms; *N* is one third of *N* indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 125
ALCOHOLIC BEVERAGES CONTAINING CAFFEINE:
Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months												2019–	
	1991–	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2021	change
	<u>2010</u>													
<i>Approximate weighted N =</i>	—	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000		
Total	—	22.5‡	19.7	16.9	14.3	12.8	10.6	9.9	9.8	8.4	8.3	7.5	-0.9*	
Gender:														
Male	—	20.4‡	19.7	17.0	13.8	12.8	10.6	9.5	11.0	8.9	§	5.4	-3.5 ss	
Female	—	24.7‡	19.8	16.7	14.7	12.7	10.8	10.2	8.3	7.8	§	9.4	+1.6	
College Plans:														
None or under 4 years	—	34.1‡	29.9	26.7	23.9	21.4	19.5	18.2	17.9	12.9	§	10.4	-2.5	
Complete 4 years	—	21.2‡	18.7	15.9	13.2	11.9	9.7	9.0	8.5	7.6	§	7.0	-0.7	
Region:														
Northeast	—	26.4‡	19.9	18.4	12.2	14.2	11.0	9.4	8.8	9.8	§	11.4	+1.7	
Midwest	—	17.4‡	18.5	15.5	13.4	12.2	9.1	8.4	9.6	6.0	§	8.9	+2.9	
South	—	22.9‡	19.5	16.6	14.9	12.4	10.7	8.7	9.0	7.1	§	6.4	-0.6	
West	—	24.7‡	21.1	17.7	16.0	12.9	11.6	13.1	12.2	12.2	§	5.0	-7.2 s	
Population Density:														
Large MSA	—	21.7‡	17.4	14.0	13.8	11.6	10.0	9.1	10.9	6.6	§	4.8	-1.8	
Other MSA	—	23.2‡	20.1	18.2	13.4	13.5	10.4	10.0	9.6	8.8	§	9.1	+0.3	
Non-MSA	—	22.1‡	21.8	17.9	17.3	13.1	12.5	10.6	8.7	10.5	§	8.6	-1.9	
Parental Education: ^a														
1.0–2.0 (Low)	—	33.8‡	23.7	21.7	16.8	18.4	12.4	12.7	13.5	10.5	§	10.9	+0.5	
2.5–3.0	—	25.6‡	23.2	20.9	18.2	14.6	14.7	10.7	12.2	8.5	§	7.4	-1.1	
3.5–4.0	—	25.5‡	22.8	19.7	16.2	14.1	11.6	10.3	10.7	8.8	§	8.7	-0.1	
4.5–5.0	—	17.1‡	18.8	13.4	12.8	11.5	8.8	8.9	7.1	8.6	§	8.0	-0.7	
5.5–6.0 (High)	—	17.0‡	12.4	12.7	9.8	10.7	8.6	8.7	8.5	7.3	§	6.2	-1.2	
Race/Ethnicity (2-year average): ^b														
White	—	—	—	19.3	17.3	15.2	13.5	12.1	11.0	10.2	§	9.4	-0.8	
African American	—	—	—	10.4	7.6	7.2	6.2	5.2	5.2	4.8	§	3.3	-1.5	
Hispanic	—	—	—	20.3	16.6	13.9	11.4	10.7	10.8	9.0	§	8.1	-1.0	

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '‡' indicates some change in the question. See relevant footnote for that drug. See relevant figure to assess the impact of the wording changes. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. ^aIn 2012 the question text was changed from "...had an alcoholic beverage containing caffeine (like Four Loko or Joose)" to "...had an alcoholic beverage mixed with an energy drink (like Red Bull)." The data in 2011 and 2012 are not comparable due to this question change. Data based on one of four forms; N is one third of N indicated in Table S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 126
ALCOHOLIC BEVERAGES CONTAINING CAFFEINE:
Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months												2019–	
	1975–												2020	change
	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>		
<i>Approximate weighted N =</i>	—	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	26.4‡	26.4	23.5	20.0	18.3	17.0	16.9	14.7	12.3	12.3	9.9	-2.4*	
Gender:														
Male	—	26.6‡	28.8	26.9	21.7	19.6	19.0	18.7	15.9	12.4	§	10.3	-2.1	
Female	—	25.9‡	23.7	20.1	18.5	17.4	14.4	15.3	12.9	11.5	§	9.8	-1.8	
College Plans:														
None or under 4 years	—	29.4‡	28.7	29.7	23.7	22.5	18.3	19.9	17.7	16.5	§	9.7	-6.8 ss	
Complete 4 years	—	25.6‡	25.3	22.0	19.1	17.4	16.3	16.0	13.8	11.2	§	10.1	-1.1	
Region:														
Northeast	—	29.7‡	28.7	23.6	22.5	21.1	17.4	21.5	14.5	10.9	§	8.8	-2.1	
Midwest	—	26.6‡	30.2	23.1	23.0	18.5	18.5	18.6	15.2	12.8	§	12.0	-0.8	
South	—	23.8‡	21.9	22.3	17.9	15.5	14.5	15.1	14.4	11.3	§	7.9	-3.4	
West	—	27.8‡	26.4	25.6	18.7	20.6	19.9	15.4	14.9	15.4	§	12.1	-3.3	
Population Density:														
Large MSA	—	27.9‡	27.7	25.2	20.3	18.5	17.5	16.8	13.3	11.1	§	11.4	+0.3	
Other MSA	—	27.4‡	27.0	24.2	20.8	17.7	17.2	16.4	14.9	12.2	§	8.3	-3.9	
Non-MSA	—	21.9‡	22.6	19.3	17.6	19.3	15.7	18.0	16.8	14.7	§	10.9	-3.8	
Parental Education: ^a														
1.0–2.0 (Low)	—	26.5‡	20.3	22.4	21.5	17.0	18.4	14.3	12.0	12.0	§	4.4	-7.7 ss	
2.5–3.0	—	28.1‡	25.9	23.1	19.5	19.2	14.8	17.1	15.9	12.5	§	7.7	-4.8 s	
3.5–4.0	—	27.0‡	30.3	25.3	20.2	18.3	17.9	17.8	15.3	12.3	§	9.6	-2.7	
4.5–5.0	—	24.7‡	26.8	22.8	20.5	19.3	17.6	16.1	15.4	13.1	§	12.5	-0.5	
5.5–6.0 (High)	—	26.3‡	24.2	22.3	20.1	19.0	16.7	17.9	13.0	12.7	§	14.2	+1.5	
Race/Ethnicity (2-year average): ^b														
White	—	—	—	28.1	24.6	22.2	21.0	20.3	19.1	15.8	§	14.2	-1.7	
African American	—	—	—	11.5	11.6	11.2	10.1	8.5	8.0	7.2	§	4.5	-2.8	
Hispanic	—	—	—	23.0	18.9	16.3	15.9	15.4	12.5	10.8	§	7.6	-3.2	

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '‡' indicates some change in the question. See relevant footnote for that drug. See relevant figure to assess the impact of the wording changes. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. In 2012 the question text was changed from "...had an alcoholic beverage containing caffeine (like Four Loko or Joose)" to "...had an alcoholic beverage mixed with an energy drink (like Red Bull)." The data in 2011 and 2012 are not comparable due to this question change. Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 127
CIGARETTES: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>	
Total	14.3	15.5	16.7	18.6	19.1	21.0	19.4	19.1	17.5	14.6	12.2	10.7	10.2	9.2	9.3	
Gender:																
Male	15.5	14.9	17.2	19.3	18.8	20.6	19.1	18.0	16.7	14.3	12.2	11.0	9.6	8.3	8.7	
Female	13.1	15.9	16.3	17.9	19.0	21.1	19.5	19.8	17.7	14.7	12.0	10.4	10.6	9.9	9.7	
College Plans:																
None or under 4 years	29.2	31.9	34.1	36.6	36.5	39.2	40.0	40.1	40.3	34.7	30.0	29.3	27.8	25.6	26.7	
Complete 4 years	11.8	13.1	14.3	16.1	16.8	18.2	16.9	16.5	14.5	12.2	10.0	8.9	8.3	7.4	7.4	
Region:																
Northeast	13.7	14.4	15.0	17.8	18.6	22.1	18.0	15.6	15.7	13.7	11.4	9.1	7.7	7.2	7.8	
Midwest	15.5	16.5	16.3	18.5	20.9	23.2	20.0	22.3	21.3	17.1	12.0	11.0	12.2	10.8	9.5	
South	15.7	17.0	18.2	19.5	19.4	21.1	21.0	21.1	18.7	14.7	14.3	13.0	11.7	10.3	11.6	
West	10.0	12.2	16.4	18.0	16.5	17.1	17.1	15.1	12.1	12.2	9.3	7.5	7.0	7.4	6.3	
Population Density:																
Large MSA	12.8	15.0	14.1	15.5	16.5	19.4	15.8	16.4	12.7	12.1	9.3	7.5	7.7	7.7	6.7	
Other MSA	14.9	15.3	17.8	20.7	19.4	21.4	19.7	17.7	16.0	13.1	11.6	10.6	9.8	8.9	9.8	
Non-MSA	14.8	16.4	17.9	17.8	21.5	22.1	22.8	24.8	26.1	21.1	16.9	14.9	14.4	11.6	11.8	
Parental Education: ^a																
1.0–2.0 (Low)	26.2	24.1	23.3	26.1	25.3	26.5	26.9	26.7	26.6	22.0	20.3	20.3	17.5	15.8	16.6	
2.5–3.0	16.4	16.9	19.8	20.6	22.7	24.4	22.4	23.9	23.5	19.6	16.4	14.5	14.8	12.2	13.4	
3.5–4.0	13.9	14.9	17.4	20.1	20.8	21.4	20.9	21.4	17.0	14.7	12.6	10.5	9.6	9.6	10.5	
4.5–5.0	10.1	13.3	12.5	14.9	14.9	18.4	16.2	14.2	12.3	10.2	8.3	7.8	6.7	6.7	5.9	
5.5–6.0 (High)	11.3	11.5	13.3	15.1	14.5	17.3	15.3	13.8	12.2	9.8	6.9	5.8	6.0	5.2	4.3	
Race/Ethnicity (2-year average): ^b																
White	—	16.2	17.8	18.9	20.7	22.7	22.8	21.5	20.1	17.7	14.7	12.0	10.9	10.0	9.4	
African American	—	5.3	6.6	8.7	8.9	9.6	10.9	10.6	10.7	9.6	8.2	7.7	6.9	6.9	7.1	
Hispanic	—	16.7	18.3	21.3	21.6	19.6	19.1	20.1	20.5	16.6	13.0	12.8	11.9	10.1	9.0	

Table continued on next page.

TABLE 127 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days																2019-2021 change	
<i>Approximate weighted N =</i>	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Total	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	-1.2* sss	
Gender:																		
Male																	0.9	-0.9 ss
Female																	1.1	-1.7 ss
College Plans:																		
None or under 4 years																	3.2	-3.1 ss
Complete 4 years																	0.7	-1.1 sss
Region:																		
Northeast																	0.6	-0.5
Midwest																	1.5	-0.2
South																	1.0	-1.8 ss
West																	1.3	-1.5
Population Density:																		
Large MSA																	0.5	-0.8 sss
Other MSA																	1.4	-0.8
Non-MSA																	1.6	-2.6 s
Parental Education: ^a																		
1.0–2.0 (Low)																	2.6	-0.9
2.5–3.0																	1.6	-2.2 ss
3.5–4.0																	2.1	-0.4
4.5–5.0																	0.7	-0.6 s
5.5–6.0 (High)																	0.6	-0.8
Race/Ethnicity (2-year average): ^b																		
White																	1.8	-0.3
African American																	0.9	-0.3
Hispanic																	0.3	-1.7 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 128
CIGARETTES: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	20.8	21.5	24.7	25.4	27.9	30.4	29.8	27.6	25.7	23.9	21.3	17.7	16.7	16.0	14.9
Gender:															
Male	20.8	20.6	24.6	26.6	27.7	30.1	28.2	26.2	25.2	23.8	20.9	16.7	16.2	16.2	14.5
Female	20.7	22.2	24.5	23.9	27.9	30.8	31.1	29.1	25.8	23.6	21.5	18.6	17.0	15.7	15.1
College Plans:															
None or under 4 years	36.5	35.0	41.9	42.2	46.3	46.2	47.2	45.2	44.0	38.6	38.1	33.3	33.0	33.1	29.4
Complete 4 years	17.3	18.6	21.0	21.7	24.7	27.8	26.8	24.5	22.7	21.5	18.5	15.1	14.0	13.6	12.9
Region:															
Northeast	22.4	21.9	27.1	24.5	27.8	31.7	29.3	30.1	28.0	23.9	18.1	15.9	16.6	16.6	14.9
Midwest	22.9	24.3	26.0	28.8	30.1	32.5	31.7	29.5	30.2	27.1	24.2	19.2	18.4	14.6	16.2
South	21.2	19.8	24.0	25.7	30.8	33.4	32.2	29.8	26.3	25.5	23.5	19.6	18.2	20.4	16.4
West	16.7	20.2	21.2	20.1	19.6	20.8	23.2	19.6	17.5	16.8	15.0	14.1	12.5	10.9	10.9
Population Density:															
Large MSA	19.7	21.6	22.5	22.3	23.3	26.2	26.6	22.5	22.9	23.1	17.3	14.2	13.1	12.0	11.7
Other MSA	20.3	20.3	23.8	26.3	28.9	31.1	28.9	26.6	25.0	21.3	20.5	17.6	16.6	16.6	14.7
Non-MSA	22.7	23.7	28.2	26.7	31.3	33.9	34.9	35.7	30.4	29.4	27.6	22.6	22.4	20.4	19.8
Parental Education: ^a															
1.0–2.0 (Low)	23.5	28.4	29.5	26.4	30.9	28.7	28.2	28.0	30.5	29.3	22.5	21.4	23.4	19.3	16.4
2.5–3.0	24.1	23.3	28.0	29.1	33.2	33.8	33.2	33.0	29.6	26.8	25.7	22.4	21.2	19.9	19.7
3.5–4.0	20.4	20.6	24.8	26.0	27.8	31.6	30.9	27.3	26.0	25.3	21.1	17.4	16.2	17.8	15.7
4.5–5.0	18.5	19.5	20.1	22.6	25.9	28.7	28.5	25.7	22.4	21.2	18.9	15.1	13.4	12.6	12.5
5.5–6.0 (High)	18.5	18.9	21.4	20.7	21.8	27.8	24.6	22.5	21.4	19.1	17.1	12.7	11.6	10.8	10.8
Race/Ethnicity (2-year average): ^b															
White	—	24.1	26.0	27.8	29.7	32.9	34.4	33.2	30.8	28.2	25.7	22.4	20.0	18.7	17.6
African American	—	6.6	7.5	9.8	11.5	12.2	12.8	13.7	12.5	11.1	11.1	9.8	8.9	9.2	8.7
Hispanic	—	18.3	20.5	19.4	21.4	23.7	23.0	21.3	21.1	19.6	16.8	14.3	13.2	13.9	13.5

Table continued on next page.

TABLE 128
CIGARETTES: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	14.5	14.0	12.3	13.1	13.6	11.8	10.8	9.1	7.2	6.3	4.9	5.0	4.2	3.4	3.2	1.8	-1.6* sss
Gender:																	
Male	13.4	14.6	12.7	13.7	15.0	13.4	12.0	10.5	7.7	6.1	5.0	5.0	4.4	3.9	§	1.7	-2.2 sss
Female	15.5	13.3	11.9	12.5	12.1	10.0	9.6	7.5	6.6	6.3	4.4	4.9	3.7	2.9	§	1.2	-1.7 ss
College Plans:																	
None or under 4 years	32.7	30.1	30.1	29.0	32.5	28.5	27.6	23.3	19.4	17.6	14.5	15.0	10.2	8.9	§	4.4	-4.5 sss
Complete 4 years	12.1	11.8	10.0	11.3	11.3	9.9	9.0	7.4	5.8	5.1	3.8	3.7	3.2	2.3	§	1.0	-1.3 sss
Region:																	
Northeast	11.8	13.1	10.1	11.1	12.1	12.2	10.9	7.2	6.7	6.1	4.4	3.9	2.9	2.1	§	1.3	-0.8 s
Midwest	18.1	17.1	14.5	12.9	14.5	11.8	9.7	9.3	8.0	7.0	4.7	5.6	4.2	3.3	§	2.3	-0.9
South	15.6	14.4	14.0	16.7	15.4	13.8	13.6	11.8	7.7	7.0	5.5	4.9	5.5	3.7	§	1.9	-1.8 s
West	10.9	11.1	9.6	10.1	11.1	8.5	7.6	6.9	5.8	4.7	4.5	5.5	3.4	3.9	§	1.6	-2.3
Population Density:																	
Large MSA	11.5	12.1	10.3	11.4	11.1	9.5	8.4	5.8	5.9	4.9	4.4	4.9	3.0	1.5	§	1.2	-0.3
Other MSA	13.3	13.9	12.3	13.0	13.5	11.5	10.2	9.5	6.0	5.7	4.3	3.8	3.7	3.8	§	1.9	-1.9 s
Non-MSA	22.4	17.2	15.4	16.1	17.5	15.7	15.4	13.0	12.3	10.4	7.2	8.4	7.6	5.6	§	2.6	-3.0 sss
Parental Education: ^a																	
1.0–2.0 (Low)	20.4	17.5	15.7	18.2	18.8	14.6	15.4	12.8	9.7	9.3	5.5	7.1	5.5	3.9	§	2.4	-1.5
2.5–3.0	19.7	18.4	15.8	16.5	18.6	16.2	14.4	13.6	9.1	8.7	7.1	7.3	5.5	6.2	§	2.2	-4.0 sss
3.5–4.0	15.5	14.7	13.9	13.8	15.1	12.7	11.2	10.2	8.4	7.8	5.3	5.0	4.7	3.6	§	1.8	-1.7 ss
4.5–5.0	10.8	11.3	9.6	10.9	10.0	8.6	9.6	6.0	5.2	5.0	3.5	3.3	3.0	1.7	§	1.4	-0.3
5.5–6.0 (High)	10.0	9.7	7.8	8.9	7.7	8.0	7.1	4.9	4.9	3.3	3.4	3.6	3.0	2.1	§	0.9	-1.2 s
Race/Ethnicity (2-year average): ^b																	
White	16.6	16.2	15.1	14.4	14.7	14.3	12.9	11.4	9.5	7.8	6.8	6.3	5.8	4.6	§	2.9	-1.7 ss
African American	8.0	7.1	6.5	6.8	6.7	7.1	6.7	5.3	4.2	3.8	2.6	2.1	1.8	1.5	§	1.4	-0.1
Hispanic	12.1	10.5	11.1	12.1	12.2	10.5	8.3	8.3	6.8	5.3	4.0	3.4	3.7	3.1	§	1.1	-2.0 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 129
CIGARETTES: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	36.7	38.8	38.4	36.7	34.4	30.5	29.4	30.0	30.3	29.3	30.1	29.6	29.4	28.7	28.6	29.4
Gender:																
Male	37.2	37.7	36.6	34.5	31.2	26.8	26.5	26.8	28.0	25.9	28.2	27.9	27.0	28.0	27.7	29.1
Female	35.9	39.1	39.6	38.1	37.1	33.4	31.6	32.6	31.6	31.9	31.4	30.6	31.4	28.9	29.0	29.2
College Plans:																
None or under 4 years	—	46.3	46.2	44.6	43.0	39.6	38.1	38.7	38.0	37.9	40.5	38.5	39.7	37.5	38.0	37.5
Complete 4 years	—	29.8	29.4	27.4	26.0	22.3	22.3	22.1	23.3	22.7	22.8	24.0	24.3	24.4	24.1	25.4
Region:																
Northeast	40.1	41.8	43.0	40.6	37.0	34.1	31.5	32.1	34.6	33.5	34.2	35.2	34.1	31.2	29.4	31.9
Midwest	39.5	41.3	40.5	39.0	36.6	31.5	32.4	33.5	33.2	31.4	34.1	32.5	31.7	31.1	34.9	34.0
South	36.2	39.1	37.6	35.7	35.4	31.8	28.9	29.4	28.7	28.6	25.6	26.1	26.0	28.0	26.4	26.1
West	26.3	28.3	27.7	27.3	24.8	21.2	21.8	20.4	21.8	22.9	26.3	23.3	26.6	23.9	22.7	25.1
Population Density:																
Large MSA	39.7	40.4	40.9	37.5	33.4	31.2	30.6	32.1	30.8	31.3	31.9	30.8	29.3	26.9	25.9	27.9
Other MSA	35.1	35.9	36.1	34.3	33.5	29.7	27.4	27.8	29.1	28.2	28.5	28.0	28.2	28.3	28.2	29.6
Non-MSA	36.7	40.9	39.2	39.4	36.4	30.9	30.9	31.2	31.5	29.3	30.8	31.0	31.8	31.4	32.2	30.4
Parental Education: ^a																
1.0–2.0 (Low)	37.2	43.2	39.6	38.1	38.1	32.7	32.5	32.6	32.7	33.6	32.3	28.6	28.8	28.1	25.4	26.3
2.5–3.0	37.0	41.2	40.8	39.3	35.9	34.2	31.7	32.0	32.2	31.8	32.3	32.3	31.4	29.9	30.8	30.8
3.5–4.0	31.9	35.3	37.3	34.0	33.3	28.0	28.2	29.0	28.0	28.1	29.7	29.7	28.8	27.8	29.4	29.3
4.5–5.0	32.3	35.0	33.0	32.6	30.1	25.7	26.0	25.5	27.8	25.2	27.7	26.4	27.6	28.6	27.0	29.1
5.5–6.0 (High)	26.8	30.8	32.8	31.9	29.6	24.0	22.5	25.1	25.5	23.7	22.6	26.7	29.3	27.8	26.3	28.6
Race/Ethnicity (2-year average): ^b																
White	—	—	38.3	37.6	36.0	33.0	30.5	30.7	31.3	31.2	31.3	31.9	32.1	32.2	32.2	32.3
African American	—	—	36.7	32.7	30.2	26.8	23.7	21.8	21.2	19.3	18.1	16.9	14.2	13.3	12.6	12.2
Hispanic	—	—	35.7	32.8	26.8	22.6	23.2	24.7	24.7	25.3	25.5	23.7	22.7	21.9	20.6	21.7

Table continued on next page.

TABLE 129 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	28.3	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	29.5	26.7	24.4	25.0	23.2	21.6
Gender:																
Male	29.0	29.2	30.7	32.9	34.5	34.9	37.3	36.3	35.4	32.8	29.7	27.4	26.2	25.3	24.8	22.4
Female	27.5	26.1	28.7	29.2	32.0	32.4	35.2	33.3	33.5	29.7	28.7	25.5	22.1	24.1	20.7	20.1
College Plans:																
None or under 4 years	38.1	38.6	37.3	40.9	43.5	45.0	45.7	46.7	44.9	43.6	40.8	37.5	36.2	36.8	34.8	32.9
Complete 4 years	24.2	23.8	27.3	28.0	29.9	30.8	33.1	31.3	31.4	27.3	25.9	23.6	20.8	21.6	20.0	18.6
Region:																
Northeast	30.5	29.6	34.2	33.2	34.4	38.5	40.6	35.9	34.2	33.1	30.3	27.3	25.0	25.9	22.0	23.1
Midwest	34.6	31.7	33.2	36.2	37.8	37.7	39.3	40.0	37.8	35.6	35.9	31.7	27.3	28.3	26.8	23.6
South	25.4	26.4	29.0	30.7	33.5	33.2	35.0	34.3	36.2	29.6	25.9	27.2	24.3	24.6	24.6	23.2
West	23.2	22.8	22.9	24.0	26.5	24.4	30.5	29.1	27.6	28.1	25.2	19.4	20.7	20.1	17.5	15.0
Population Density:																
Large MSA	26.2	25.6	29.5	29.0	33.9	32.1	34.9	32.9	30.0	27.4	27.3	24.8	18.9	20.8	20.8	20.7
Other MSA	29.3	26.9	29.8	31.1	31.7	32.6	35.7	34.2	35.0	31.5	28.2	26.2	25.1	26.3	22.6	20.3
Non-MSA	28.6	31.5	30.3	33.8	36.2	38.2	40.0	39.7	38.7	36.3	34.3	30.1	30.4	27.6	27.4	25.3
Parental Education: ^a																
1.0–2.0 (Low)	31.3	27.1	26.5	26.2	31.2	31.5	31.2	32.3	33.0	31.3	24.8	20.9	23.5	21.0	19.1	21.1
2.5–3.0	28.7	30.3	30.4	32.8	35.0	35.5	36.5	36.0	37.3	32.2	31.5	28.9	27.0	28.7	27.3	24.7
3.5–4.0	28.4	27.8	29.9	31.4	33.2	33.2	35.6	36.7	35.0	32.8	30.3	28.6	24.3	26.3	24.8	22.9
4.5–5.0	26.9	25.8	30.1	32.0	32.6	34.5	37.5	34.2	32.4	30.2	29.3	25.0	22.6	23.8	21.8	19.2
5.5–6.0 (High)	27.1	25.5	30.5	30.4	34.0	32.9	38.5	33.1	34.4	27.4	25.0	25.3	21.0	19.9	18.0	17.4
Race/Ethnicity (2-year average): ^b																
White	32.2	31.8	33.2	35.2	36.6	38.1	40.7	41.7	40.1	37.9	35.3	32.5	29.4	28.2	27.6	25.9
African American	10.6	8.7	9.5	10.9	12.9	14.2	14.3	14.9	14.9	14.3	13.3	12.1	10.0	10.1	10.7	10.5
Hispanic	24.0	25.0	24.2	23.6	25.1	25.4	25.9	26.6	27.3	27.7	23.8	21.3	19.0	18.5	17.1	16.0

Table continued on next page.

TABLE 129 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	21.6	20.4	20.1	19.2	18.7	17.1	16.3	13.6	11.4	10.5	9.7	7.6	5.7	7.5	4.1	-1.6*
Gender:																
Male	23.1	21.5	22.1	21.9	21.5	19.3	18.4	15.2	13.0	12.7	10.6	8.4	6.9	§	3.3	-3.7 sss
Female	19.6	19.1	17.6	15.7	15.1	14.5	13.2	11.6	9.1	8.1	8.3	5.9	4.0	§	4.5	+0.5
College Plans:																
None or under 4 years	32.4	32.4	31.2	31.2	32.2	27.9	28.9	25.3	21.9	19.2	16.5	14.6	10.7	§	6.3	-4.4 s
Complete 4 years	19.0	17.6	17.3	16.3	15.6	14.7	13.0	10.8	8.9	8.4	8.0	5.2	4.0	§	2.9	-1.1
Region:																
Northeast	23.8	19.7	18.9	18.2	17.8	18.8	17.0	11.7	10.8	10.9	9.6	6.4	3.7	§	3.3	-0.3
Midwest	24.7	22.1	25.7	21.5	21.8	21.0	16.5	13.8	12.7	12.0	9.6	7.3	5.7	§	2.9	-2.8 ss
South	21.9	22.4	20.0	20.5	18.4	15.6	18.6	15.4	12.5	10.8	11.2	9.2	6.6	§	3.3	-3.3 sss
West	15.8	15.8	14.7	15.3	16.4	13.9	12.1	11.9	8.6	8.3	7.1	6.1	5.8	§	6.9	+1.1
Population Density:																
Large MSA	19.9	18.8	17.7	18.7	17.2	15.8	14.5	11.7	9.3	8.4	7.4	5.3	3.4	§	3.5	+0.1
Other MSA	20.8	19.5	19.1	17.9	18.1	15.6	15.5	12.9	10.5	9.9	9.5	6.8	6.0	§	3.9	-2.2
Non-MSA	25.9	24.3	25.5	23.1	22.3	23.1	20.9	17.8	17.0	15.1	13.8	13.9	9.0	§	5.8	-3.2 s
Parental Education: ^a																
1.0–2.0 (Low)	20.5	18.5	19.9	17.0	18.2	17.0	18.2	14.8	12.9	9.9	7.9	6.9	7.2	§	3.0	-4.3 ss
2.5–3.0	24.7	22.4	23.2	21.6	22.4	19.0	18.2	16.1	13.9	12.0	11.3	8.0	6.6	§	6.0	-0.6
3.5–4.0	21.6	21.3	21.0	20.9	19.6	20.1	18.5	14.2	12.2	11.9	10.4	8.1	5.3	§	5.4	+0.1
4.5–5.0	20.5	18.6	18.0	17.5	15.9	15.7	14.1	12.0	9.7	9.1	8.7	6.6	4.7	§	2.2	-2.5 ss
5.5–6.0 (High)	18.4	19.2	15.3	12.8	14.3	11.2	9.5	9.4	7.5	7.6	8.1	5.9	4.2	§	3.0	-1.2
Race/Ethnicity (2-year average): ^b																
White	24.9	24.7	23.9	22.9	22.2	21.2	19.4	17.5	15.0	13.2	12.7	11.0	7.9	§	5.9	-2.0
African American	10.8	10.3	9.8	10.1	9.7	8.6	9.6	9.0	6.9	6.0	5.0	4.1	3.2	§	3.0	-0.2
Hispanic	15.3	15.0	15.7	15.0	14.3	13.2	12.4	11.4	9.6	8.2	6.9	5.2	4.5	§	3.8	-0.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 130
CIGARETTES: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 8

	Percentage who used daily in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Approximate weighted N =	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	7.2	7.0	8.3	8.8	9.3	10.4	9.0	8.8	8.1	7.4	5.5	5.1	4.5	4.4	4.0
Gender:															
Male	8.1	6.9	8.8	9.5	9.2	10.5	9.0	8.1	7.4	7.0	5.9	5.4	4.4	4.3	3.9
Female	6.2	7.2	7.8	8.0	9.2	10.1	8.7	9.0	8.4	7.5	4.9	4.9	4.5	4.3	4.0
College Plans:															
None or under 4 years	18.5	20.1	21.5	22.6	22.5	26.0	25.4	25.2	25.2	21.7	17.7	17.1	16.1	15.4	14.4
Complete 4 years	5.3	5.1	6.4	6.8	7.5	8.0	6.9	6.6	5.9	5.6	3.9	3.9	3.2	3.1	2.9
Region:															
Northeast	7.2	7.1	7.1	8.6	9.2	11.0	8.8	6.1	7.2	6.9	6.1	3.7	2.9	3.3	3.2
Midwest	7.8	7.6	8.5	9.4	11.0	12.4	10.3	11.2	11.5	9.0	6.4	5.7	5.5	5.7	4.8
South	7.9	7.8	9.3	9.4	9.4	10.4	9.5	10.2	8.5	7.8	6.1	6.6	5.7	4.7	5.0
West	4.6	4.8	7.4	7.4	7.0	7.5	6.8	5.8	3.8	4.9	2.6	2.9	2.4	3.3	2.4
Population Density:															
Large MSA	6.3	6.3	5.7	6.6	7.6	9.5	7.0	6.7	5.4	5.6	4.1	3.2	3.0	3.5	2.3
Other MSA	7.7	7.2	9.1	9.5	9.3	10.2	8.7	7.9	7.4	6.3	5.4	5.1	4.3	4.1	4.1
Non-MSA	7.3	7.8	10.1	9.6	11.1	11.8	11.7	12.7	12.7	11.8	7.2	7.6	7.0	5.9	6.2
Parental Education: ^a															
1.0–2.0 (Low)	15.9	11.9	12.7	13.0	15.8	13.6	14.3	13.0	14.7	13.1	11.1	10.6	9.1	10.1	7.8
2.5–3.0	8.6	8.4	9.7	11.3	11.3	14.0	11.7	12.0	11.4	11.3	7.5	7.1	6.6	5.6	6.3
3.5–4.0	6.5	6.9	8.5	8.9	9.4	10.1	9.2	9.7	8.1	6.7	5.1	5.4	4.4	4.2	4.3
4.5–5.0	4.0	5.2	5.9	6.1	7.2	7.6	6.8	5.7	4.6	3.9	3.0	3.3	2.6	2.8	2.2
5.5–6.0 (High)	4.9	4.2	6.3	5.8	5.7	7.4	5.5	5.2	5.1	4.1	3.1	2.1	2.1	1.9	1.4
Race/Ethnicity (2-year average) ^b															
White	—	7.7	8.8	9.7	10.5	11.7	11.4	10.4	9.7	9.0	7.5	6.0	5.3	4.7	4.6
African American	—	1.4	1.8	2.6	2.8	3.2	3.7	3.8	3.8	3.2	2.8	2.8	2.9	2.7	2.1
Hispanic	—	7.3	7.2	9.0	9.2	8.0	8.1	8.4	8.5	7.1	5.0	4.4	3.7	3.5	3.1

Table continued on next page.

TABLE 130 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 8

	Percentage who used daily in last 30 days																2019-2021 change	
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700		
Total	4.0	3.0	3.1	2.7	2.9	2.4	1.9	1.8	1.4	1.3	0.9	0.6	0.8	0.8	0.8	0.4	-0.4* s	
Gender:																		
Male	4.0	3.4	3.2	2.9	3.5	2.5	2.0	1.7	1.2	1.1	1.0	0.6	0.9	0.6	§	0.3	-0.3	
Female	3.8	2.6	2.9	2.3	2.3	2.2	1.6	1.8	1.3	1.4	0.8	0.5	0.6	0.9	§	0.3	-0.6 s	
College Plans:																		
None or under 4 years	13.2	12.3	10.8	10.9	12.8	8.1	8.0	8.3	4.7	5.9	4.4	1.8	3.3	2.8	§	1.3	-1.5	
Complete 4 years	2.9	2.2	2.3	1.9	2.0	1.9	1.4	1.4	1.0	0.9	0.5	0.4	0.4	0.5	§	0.1	-0.4 ss	
Region:																		
Northeast	3.0	1.7	2.4	2.1	2.4	1.4	1.8	1.3	1.2	0.7	0.6	0.2	0.5	0.4	§	0.2	-0.2	
Midwest	4.7	3.8	3.9	3.0	3.3	2.3	1.3	1.5	0.9	1.6	1.0	0.8	1.0	0.6	§	0.4	-0.2	
South	5.3	3.8	3.7	3.2	3.8	2.9	2.9	2.6	1.9	1.5	1.0	0.6	0.8	0.8	§	0.3	-0.5 s	
West	1.8	1.9	2.0	2.0	1.4	2.3	0.8	1.2	1.1	1.1	0.9	0.6	1.0	1.2	§	0.5	-0.6	
Population Density:																		
Large MSA	2.3	1.6	2.2	1.6	1.7	1.2	1.4	0.8	1.1	0.6	0.4	0.4	0.5	0.4	§	0.1	-0.3 s	
Other MSA	4.1	3.1	3.3	3.0	3.1	3.2	2.0	1.8	1.3	1.2	0.7	0.6	0.9	0.7	§	0.5	-0.2	
Non-MSA	6.3	5.0	4.2	3.7	4.3	2.7	2.8	3.4	2.1	2.7	2.5	0.9	1.4	1.5	§	0.5	-1.0 s	
Parental Education: ^a																		
1.0–2.0 (Low)	7.3	5.5	5.4	6.2	4.2	3.9	2.6	2.7	2.6	1.7	2.2	1.1	1.0	1.1	§	1.4	+0.3	
2.5–3.0	6.6	4.9	4.4	3.4	4.8	3.9	3.3	3.0	1.9	1.9	1.2	0.7	0.8	1.3	§	0.5	-0.8	
3.5–4.0	3.8	3.6	3.0	2.5	2.8	2.9	2.2	2.0	1.6	1.6	0.6	0.6	1.1	0.6	§	0.6	0.0	
4.5–5.0	2.1	1.5	1.9	1.8	1.9	1.4	0.9	1.0	0.8	0.8	0.6	0.3	0.5	0.4	§	0.1	-0.3 s	
5.5–6.0 (High)	1.8	0.7	1.9	1.4	1.0	0.9	1.0	1.0	0.5	0.4	0.5	0.3	0.5	0.7	§	0.1	-0.6 s	
Race/Ethnicity (2-year average) ^b																		
White	4.6	3.9	3.3	3.2	3.2	3.0	2.4	2.0	1.7	1.4	1.2	0.9	0.6	0.7	§	0.6	-0.1	
African American	1.9	2.1	1.9	2.0	1.9	1.5	1.6	1.5	1.2	0.9	0.5	0.4	0.5	0.5	§	0.5	+0.1	
Hispanic	2.8	2.8	2.5	2.2	2.3	2.4	1.8	1.4	1.3	1.0	0.8	0.6	0.4	0.6	§	0.1	-0.4 s	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 131
CIGARETTES: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 10

	Percentage who used daily in last 30 days															
Approximate weighted N =	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
Total	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
	12.6	12.3	14.2	14.6	16.3	18.3	18.0	15.8	15.9	14.0	12.2	10.1	8.9	8.3	7.5	
Gender:																
Male	12.4	12.1	13.8	15.2	16.3	18.1	17.2	14.7	15.6	13.7	12.4	9.4	8.6	8.2	7.2	
Female	12.5	12.4	14.3	13.7	16.1	18.6	18.5	16.8	15.9	14.1	11.9	10.8	9.0	8.2	7.7	
College Plans:																
None or under 4 years	25.7	25.5	28.9	28.9	32.7	34.3	35.4	31.7	32.1	28.8	27.3	22.9	22.1	21.4	19.2	
Complete 4 years	9.6	9.5	11.0	11.5	13.3	15.5	15.0	12.9	13.2	11.6	9.6	7.9	6.7	6.4	5.9	
Region:																
Northeast	14.3	13.1	16.3	14.1	15.8	18.8	18.0	18.7	17.7	14.1	11.0	8.3	8.6	8.5	7.6	
Midwest	14.3	14.3	15.1	16.9	17.6	20.6	19.5	17.3	19.6	16.3	13.2	11.5	10.2	7.4	8.6	
South	12.8	11.4	13.9	15.5	19.3	20.5	20.5	17.1	16.3	15.7	14.3	11.3	10.1	11.0	8.8	
West	9.1	10.7	10.9	9.7	9.4	10.7	11.1	8.8	9.1	7.8	7.0	7.8	6.0	5.2	4.0	
Population Density:																
Large MSA	12.3	11.7	12.3	12.0	12.6	15.3	15.7	12.2	13.2	13.8	9.6	7.4	6.1	5.6	5.5	
Other MSA	11.7	11.6	13.6	15.5	17.5	18.8	16.9	15.1	15.5	12.1	11.6	10.4	8.7	8.1	7.0	
Non-MSA	14.3	14.5	16.9	15.5	18.4	20.8	22.5	21.1	19.7	17.5	16.3	13.1	13.7	12.3	11.4	
Parental Education: ^a																
1.0–2.0 (Low)	16.0	17.8	19.3	15.5	20.0	19.3	17.7	17.4	20.1	18.9	12.9	14.1	15.3	11.4	9.9	
2.5–3.0	15.5	13.9	16.9	17.6	21.6	23.1	22.1	21.3	19.1	17.6	16.2	14.7	12.7	11.5	11.1	
3.5–4.0	12.0	11.8	13.6	15.9	17.0	19.4	18.9	14.9	16.6	14.2	12.2	10.0	8.8	9.3	7.9	
4.5–5.0	10.6	10.5	10.7	11.5	12.6	14.8	15.6	12.9	13.0	11.5	9.7	6.8	5.8	5.4	5.2	
5.5–6.0 (High)	9.6	9.0	10.5	9.6	10.3	13.6	12.0	11.1	11.2	9.8	8.3	6.4	4.5	4.2	4.4	
Race/Ethnicity (2-year average): ^b																
White	—	14.5	15.3	16.5	17.6	20.0	21.4	20.3	19.1	17.7	15.5	13.3	11.4	10.0	9.1	
African American	—	2.8	3.1	3.8	4.7	5.1	5.6	5.8	5.3	5.2	5.0	4.3	4.4	3.9		
Hispanic	—	8.4	8.9	8.1	9.9	11.6	10.8	9.4	9.1	8.8	7.4	6.4	6.0	6.0	5.9	

Table continued on next page.

TABLE 131 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 10

	Percentage who used daily in last 30 days																2019-2021
Approximate weighted N =	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	7.6	7.2	5.9	6.3	6.6	5.5	5.0	4.4	3.2	3.0	1.9	2.2	1.8	1.3	1.2	0.8	-0.6* s
Gender:																	
Male	6.9	7.7	6.2	6.9	7.2	6.4	5.6	5.4	3.5	2.8	2.2	2.3	1.9	1.5	§	0.7	-0.8 ss
Female	8.1	6.6	5.5	5.6	5.9	4.5	4.4	3.4	2.8	2.8	1.5	1.9	1.5	1.2	§	0.4	-0.8 ss
College Plans:																	
None or under 4 years	21.7	18.9	17.9	17.3	19.1	17.1	17.0	13.4	10.5	10.8	7.2	8.9	5.1	3.6	§	1.8	-1.8 s
Complete 4 years	5.6	5.6	4.3	5.0	5.0	4.2	3.7	3.4	2.3	2.0	1.2	1.4	1.2	0.9	§	0.4	-0.5 s
Region:																	
Northeast	6.2	6.8	5.3	5.3	5.7	6.3	5.1	3.6	2.8	2.7	1.6	2.2	1.0	0.4	§	0.7	+0.3
Midwest	10.3	10.0	7.6	5.8	7.3	5.7	4.4	4.9	3.9	2.8	2.3	2.8	1.8	1.6	§	1.1	-0.5
South	8.4	7.2	6.9	8.5	7.9	6.6	6.9	5.7	3.4	3.6	2.2	2.1	2.6	1.5	§	0.7	-0.8 s
West	4.2	4.7	3.4	4.3	4.4	3.0	2.6	3.1	2.4	2.3	1.4	2.1	1.1	1.3	§	0.6	-0.7
Population Density:																	
Large MSA	5.4	5.7	4.3	5.2	4.8	4.3	3.6	2.3	2.5	2.1	1.6	1.7	1.1	0.4	§	0.5	+0.1
Other MSA	6.7	7.2	6.2	6.0	6.3	5.3	4.7	4.8	2.6	2.5	1.6	1.5	1.4	1.6	§	0.6	-1.0 s
Non-MSA	13.2	9.6	7.5	8.5	9.7	7.6	7.4	6.8	5.7	5.5	3.2	4.9	3.8	2.2	§	1.5	-0.7
Parental Education: ^a																	
1.0–2.0 (Low)	12.5	9.8	7.6	9.4	10.0	8.2	8.0	6.8	4.6	5.2	2.4	3.8	3.2	1.5	§	0.7	-0.8
2.5–3.0	11.9	10.8	8.7	8.7	10.1	8.9	7.5	7.2	4.7	4.6	3.0	3.5	2.8	2.5	§	0.8	-1.7 ss
3.5–4.0	7.8	7.2	6.7	6.7	7.1	5.6	4.5	4.7	3.9	3.4	1.9	2.5	1.9	1.5	§	0.9	-0.6
4.5–5.0	4.4	5.5	3.8	4.6	3.9	3.3	4.3	2.8	1.8	2.1	1.2	1.1	1.0	0.4	§	0.5	+0.1
5.5–6.0 (High)	4.3	3.5	3.0	3.5	3.1	2.5	2.8	1.9	1.6	1.0	1.0	1.0	0.8	0.8	§	0.3	-0.6
Race/Ethnicity (2-year average): ^b																	
White	8.7	8.8	8.0	7.1	7.4	7.1	6.2	5.7	4.8	3.5	2.8	2.5	2.4	1.9	§	0.9	-1.0 ss
African American	3.3	3.2	3.0	3.2	3.5	3.5	2.9	2.6	2.3	2.1	1.3	0.9	1.0	0.8	§	1.0	+0.2
Hispanic	5.3	3.8	3.8	4.5	4.4	3.8	3.0	2.6	2.3	2.1	1.9	1.6	1.6	1.1	§	0.3	-0.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 132
CIGARETTES: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	26.9	28.8	28.8	27.5	25.4	21.3	20.3	21.1	21.2	18.7	19.5	18.7	18.7	18.1	18.9	19.1
Gender:																
Male	26.9	28.0	27.1	26.0	22.3	18.5	18.1	18.2	19.2	16.0	17.8	16.9	16.4	17.4	17.9	18.6
Female	26.4	28.8	30.0	28.3	27.8	23.5	21.7	23.2	22.2	20.5	20.6	19.8	20.6	18.1	19.4	19.3
College Plans:																
None or under 4 years	—	36.5	37.2	35.2	33.8	29.7	29.3	29.5	29.3	27.2	29.6	28.2	29.0	27.4	27.9	28.3
Complete 4 years	—	19.8	19.3	18.3	17.0	13.8	12.9	13.2	13.8	11.9	12.4	12.8	13.3	13.4	14.6	14.7
Region:																
Northeast	31.4	32.3	33.8	32.5	28.6	24.1	23.3	23.4	26.1	23.6	24.9	24.9	24.8	21.4	21.3	22.8
Midwest	28.6	30.2	29.4	28.6	27.0	22.0	23.0	24.0	23.4	20.4	22.4	19.9	20.3	19.0	23.0	22.2
South	26.2	29.1	28.7	26.4	25.8	22.6	19.1	20.2	19.4	17.7	16.0	15.8	15.7	17.7	17.1	16.5
West	17.3	19.4	19.2	19.1	17.0	14.0	13.1	12.7	13.0	12.4	14.2	13.4	14.9	14.0	13.8	14.8
Population Density:																
Large MSA	30.8	30.4	30.9	29.2	24.5	21.6	21.9	23.5	22.1	21.5	21.9	20.6	20.3	18.0	16.7	19.0
Other MSA	25.6	27.1	27.2	25.7	25.0	21.3	19.0	19.3	20.2	17.4	17.7	17.0	17.6	17.7	19.0	19.0
Non-MSA	25.8	29.5	29.1	28.7	26.5	21.2	20.7	21.3	21.7	18.2	19.9	19.8	19.3	18.8	20.9	19.5
Parental Education: ^a																
1.0–2.0 (Low)	27.2	32.7	29.6	28.6	29.1	23.7	24.1	24.6	24.0	23.2	22.7	20.4	19.7	19.2	17.1	16.7
2.5–3.0	27.2	31.3	31.5	30.3	26.5	24.7	22.5	23.1	23.2	21.5	21.8	21.4	21.1	19.6	21.5	21.0
3.5–4.0	22.1	25.8	28.1	24.8	24.5	19.4	19.0	19.7	18.8	16.4	19.3	19.4	17.8	17.5	19.0	19.3
4.5–5.0	22.9	24.5	23.7	23.2	21.2	16.6	16.1	16.8	17.5	14.1	16.0	13.9	16.5	16.5	17.2	18.3
5.5–6.0 (High)	17.4	22.8	21.7	22.8	20.6	15.0	13.9	14.5	17.2	14.1	11.2	13.6	16.6	15.1	15.8	16.5
Race/Ethnicity (2-year average): ^b																
White	—	—	28.9	28.3	26.9	23.9	21.4	21.6	22.1	21.0	20.4	20.6	20.5	20.6	21.1	21.8
African American	—	—	24.9	22.7	20.9	17.4	14.6	13.1	12.5	10.7	9.9	9.4	7.9	7.3	6.4	5.8
Hispanic	—	—	22.6	20.4	15.8	12.8	13.6	14.3	14.9	13.9	11.8	11.3	11.0	10.9	10.8	10.9

Table continued on next page.

TABLE 132 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	18.5	17.2	19.0	19.4	21.6	22.2	24.6	22.4	23.1	20.6	19.0	16.9	15.8	15.6	13.6	12.2
Gender:																
Male	18.8	17.2	19.4	20.4	21.7	22.2	24.8	22.7	23.6	20.9	18.4	17.2	17.0	15.4	14.6	12.0
Female	17.9	16.7	18.2	18.1	20.8	21.8	23.6	21.5	22.2	19.7	18.9	16.1	14.0	15.0	11.9	11.8
College Plans:																
None or under 4 years	28.4	28.1	27.8	29.8	33.7	33.2	35.6	34.6	34.2	31.7	30.1	27.6	27.9	26.9	24.9	22.8
Complete 4 years	14.1	12.9	15.9	15.7	17.4	18.9	20.6	18.4	19.5	16.6	15.5	13.8	12.1	12.2	10.5	9.4
Region:																
Northeast	20.9	19.4	23.5	21.3	22.5	27.0	29.4	23.4	23.2	22.8	21.9	18.4	16.4	16.2	13.3	14.3
Midwest	23.0	19.0	21.3	23.8	25.7	26.1	28.0	27.8	25.9	23.6	25.2	22.5	18.2	18.5	16.3	13.2
South	16.4	16.7	18.5	19.3	21.7	20.5	22.6	21.8	24.2	19.4	15.5	16.6	16.3	15.8	15.4	13.3
West	13.9	13.3	13.0	12.4	14.5	13.8	17.5	15.5	17.3	16.9	13.4	9.5	11.8	10.1	7.6	7.0
Population Density:																
Large MSA	16.7	16.6	17.3	17.7	21.3	20.7	23.7	20.6	18.6	16.7	17.4	15.0	11.5	12.6	11.4	10.3
Other MSA	19.0	15.9	19.7	19.2	19.9	21.9	23.9	21.2	22.8	21.1	17.5	16.7	15.9	16.1	12.7	11.5
Non-MSA	19.0	20.3	19.2	21.6	24.8	24.1	26.8	27.2	28.5	24.5	23.9	19.8	21.4	18.0	18.2	16.2
Parental Education: ^a																
1.0–2.0 (Low)	21.2	16.5	17.6	16.9	21.3	21.1	21.9	21.7	23.8	22.8	16.4	12.4	15.1	14.2	11.7	11.4
2.5–3.0	19.8	20.4	20.2	22.4	24.6	24.4	26.0	24.7	26.9	22.9	21.9	19.8	18.3	19.4	18.3	16.1
3.5–4.0	18.5	16.9	18.9	18.9	21.6	21.2	23.8	23.8	23.6	21.2	19.9	19.1	16.5	16.3	14.4	12.3
4.5–5.0	16.2	15.0	18.9	18.7	19.7	22.4	24.9	20.6	20.6	18.6	17.9	14.1	13.0	13.7	11.7	10.4
5.5–6.0 (High)	16.1	12.8	16.6	17.3	18.5	20.0	22.9	17.4	19.0	15.2	13.4	14.3	11.3	10.3	8.1	8.3
Race/Ethnicity (2-year average): ^b																
White	21.5	20.5	21.4	22.9	23.9	25.4	27.8	28.3	26.9	25.7	23.8	21.8	19.5	18.3	17.1	15.3
African American	5.1	4.2	4.1	4.9	6.1	7.0	7.2	7.4	7.7	8.0	7.5	6.4	5.4	5.2	5.6	5.7
Hispanic	11.5	12.5	11.8	10.6	11.6	12.9	14.0	13.6	14.0	15.7	12.0	9.2	8.0	8.2	7.7	7.0

Table continued on next page.

TABLE 132 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	12.3	11.4	11.2	10.7	10.3	9.3	8.5	6.7	5.5	4.8	4.2	3.6	2.4	3.1	2.0	-0.4*
Gender:																
Male	13.0	12.0	11.8	12.3	11.6	10.9	9.7	7.9	6.6	5.6	4.3	3.8	2.8	§	1.2	-1.5 sss
Female	11.2	10.6	9.9	8.7	8.6	7.3	6.5	5.4	3.9	3.7	3.8	2.6	1.6	§	2.4	+0.8
College Plans:																
None or under 4 years	22.5	21.1	20.9	21.6	21.0	18.4	18.8	14.0	14.1	10.3	8.7	8.6	5.6	§	3.6	-2.0
Complete 4 years	9.9	9.0	8.6	8.2	7.9	7.0	5.8	5.0	3.5	3.4	3.1	1.9	1.3	§	1.0	-0.3
Region:																
Northeast	14.3	11.3	10.8	10.3	10.2	11.3	9.8	6.8	5.4	5.9	4.1	2.3	1.8	§	1.3	-0.5
Midwest	14.5	12.9	15.4	12.5	11.7	11.6	8.4	6.6	6.6	5.5	4.3	3.5	2.2	§	1.3	-0.8
South	12.9	13.1	11.4	12.3	10.8	9.0	10.5	8.2	6.1	5.0	5.2	4.9	2.7	§	1.4	-1.3 s
West	7.4	7.0	5.9	6.7	7.9	5.8	4.6	4.5	3.5	3.0	2.7	2.5	2.4	§	4.1	+1.7
Population Density:																
Large MSA	10.3	8.9	8.6	9.9	9.2	8.3	7.2	5.4	4.0	3.4	2.8	2.1	1.2	§	1.2	0.0
Other MSA	11.7	11.2	11.0	9.9	9.5	8.2	7.9	6.3	5.0	4.7	4.0	3.3	2.3	§	2.4	+0.1
Non-MSA	16.6	14.8	15.1	14.0	13.7	13.5	11.8	9.6	9.4	7.2	7.3	7.1	4.5	§	2.3	-2.1 s
Parental Education: ^a																
1.0–2.0 (Low)	14.6	11.9	11.2	8.9	9.9	9.6	10.2	6.9	7.2	5.0	3.8	3.7	2.9	§	2.2	-0.8
2.5–3.0	15.1	13.4	14.7	13.9	13.5	11.6	10.4	8.9	7.8	6.2	5.3	4.6	3.4	§	2.8	-0.6
3.5–4.0	12.3	12.5	11.5	11.6	10.7	11.2	10.0	6.7	5.8	5.2	4.6	3.9	2.4	§	3.1	+0.7
4.5–5.0	11.2	9.0	9.0	8.7	8.2	7.1	6.4	5.8	3.7	3.6	3.6	2.3	1.2	§	0.8	-0.4
5.5–6.0 (High)	7.4	9.1	5.8	6.1	6.7	4.7	3.6	4.7	3.1	2.6	2.6	1.4	1.1	§	0.5	-0.7
Race/Ethnicity (2-year average): ^b																
White	14.5	14.3	13.9	13.5	13.0	12.1	10.9	9.3	7.3	6.1	5.8	5.0	3.5	§	2.5	-1.0
African American	5.8	5.8	5.4	5.3	4.9	4.7	5.3	5.1	4.1	3.5	2.5	2.0	1.8	§	1.6	-0.2
Hispanic	6.6	6.7	6.4	5.7	5.3	4.9	4.7	4.1	3.7	2.9	1.9	1.8	1.9	§	1.6	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 133
CIGARETTES: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by Subgroups in Grade 8

	Percentage who used half pack a day or more in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>
Total	3.1	2.9	3.5	3.6	3.4	4.3	3.5	3.6	3.3	2.8	2.3	2.1	1.8	1.7	1.7
Gender:															
Male	3.7	3.1	4.3	4.2	3.7	4.7	3.7	3.5	3.1	2.8	2.6	2.4	1.8	1.7	2.0
Female	2.4	2.7	2.7	2.9	3.2	3.7	3.1	3.3	3.3	2.6	2.0	1.9	1.7	1.7	1.3
College Plans:															
None or under 4 years	10.1	10.8	11.9	11.7	11.4	13.5	11.6	13.8	13.4	10.3	9.8	8.4	7.5	7.4	7.7
Complete 4 years	1.9	1.7	2.2	2.4	2.3	2.8	2.4	2.2	2.0	1.8	1.4	1.5	1.2	1.1	1.0
Region:															
Northeast	3.3	2.8	2.7	3.7	3.7	4.1	3.7	2.1	3.1	3.3	2.6	1.7	1.0	1.4	1.4
Midwest	3.3	3.2	3.5	3.9	4.4	5.3	4.1	4.6	5.7	3.4	3.1	2.7	2.1	2.3	2.2
South	3.4	3.3	4.6	3.9	3.6	4.5	3.7	4.8	2.9	2.8	2.7	2.5	2.6	1.8	1.9
West	1.9	2.0	2.2	2.6	1.8	2.7	2.2	1.6	1.2	1.3	0.7	0.9	0.8	1.2	0.9
Population Density:															
Large MSA	2.4	2.3	2.1	3.0	2.5	3.7	2.7	2.3	1.8	2.1	1.5	1.3	1.0	1.3	1.0
Other MSA	3.3	3.2	3.7	3.9	3.6	3.8	3.2	3.2	2.8	2.2	2.1	2.2	1.7	1.6	1.5
Non-MSA	3.4	3.3	5.0	3.7	4.3	5.8	4.8	5.6	5.9	4.6	3.9	3.1	3.1	2.4	2.9
Parental Education: ^a															
1.0–2.0 (Low)	7.9	6.5	6.4	5.1	8.2	6.5	6.4	6.2	6.2	7.1	5.2	4.6	3.7	5.6	2.9
2.5–3.0	3.7	3.4	3.9	4.9	4.1	6.4	4.8	5.2	4.8	3.8	3.6	2.8	2.9	2.1	2.8
3.5–4.0	2.5	2.6	3.6	3.4	3.6	3.2	3.5	3.7	3.4	2.3	2.0	2.2	1.8	1.5	1.9
4.5–5.0	1.6	1.8	2.3	2.6	2.0	2.8	2.2	2.0	1.5	1.4	0.9	1.2	0.7	0.8	0.6
5.5–6.0 (High)	1.8	1.5	2.2	2.2	1.5	2.7	1.8	2.1	1.8	1.5	1.4	1.1	0.9	0.6	0.5
Race/Ethnicity (2-year average): ^b															
White	—	3.3	3.8	4.2	4.2	4.6	4.6	4.2	4.0	3.6	3.2	2.8	2.3	1.9	1.9
African American	—	0.4	0.7	1.0	0.9	1.3	1.4	1.2	1.3	1.0	1.1	1.1	1.2	1.1	0.9
Hispanic	—	2.7	2.5	3.1	3.3	2.7	2.4	2.8	3.0	2.3	1.4	1.3	1.5	1.4	1.1

Table continued on next page.

TABLE 133 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by Subgroups in Grade 8

	Percentage who used half pack a day or more in last 30 days																
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2019-2021 change
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	1.5	1.1	1.2	1.0	0.9	0.7	0.6	0.7	0.5	0.4	0.3	0.2	0.3	0.2	0.1	0.2	-0.1*
Gender:																	
Male	1.7	1.3	1.4	1.2	1.1	0.7	0.8	0.9	0.5	0.5	0.4	0.2	0.3	0.2	§	0.2	0.0
Female	1.3	0.8	0.9	0.7	0.7	0.6	0.3	0.5	0.4	0.3	0.2	0.1	0.2	0.2	§	0.0	-0.2 s
College Plans:																	
None or under 4 years	5.8	4.7	4.7	4.7	4.6	3.0	3.0	4.0	1.8	2.4	1.8	0.6	1.1	1.0	§	0.4	-0.6
Complete 4 years	1.1	0.7	0.8	0.6	0.5	0.5	0.4	0.5	0.3	0.2	0.1	0.1	0.1	0.1	§	0.0	+0.0
Region:																	
Northeast	1.2	0.8	0.9	0.8	0.6	0.2	0.7	0.4	0.4	0.3	0.2	0.1	0.3	0.2	§	0.2	-0.1
Midwest	1.9	1.2	1.6	1.1	1.0	0.8	0.5	0.5	0.3	0.4	0.2	0.3	0.6	0.2	§	0.1	-0.1
South	2.1	1.3	1.3	1.1	1.2	0.8	0.8	1.1	0.8	0.6	0.5	0.2	0.2	0.3	§	0.2	-0.1
West	0.6	0.9	0.6	0.7	0.5	1.0	0.3	0.6	0.3	0.3	0.3	0.2	0.4	0.1	§	0.1	0.0
Population Density:																	
Large MSA	0.9	0.7	0.7	0.5	0.4	0.3	0.4	0.5	0.3	0.2	0.2	0.1	0.2	0.2	§	0.1	-0.1
Other MSA	1.6	1.1	1.3	1.1	0.9	1.0	0.6	0.7	0.5	0.4	0.3	0.2	0.3	0.2	§	0.1	0.0
Non-MSA	2.3	1.7	1.6	1.3	1.4	0.8	0.8	1.3	0.9	0.9	0.7	0.4	0.5	0.3	§	0.2	0.0
Parental Education: ^a																	
1.0–2.0 (Low)	3.6	1.9	2.5	2.6	0.9	1.2	0.7	1.0	1.2	0.7	1.0	0.3	0.3	0.3	§	0.7	+0.4
2.5–3.0	2.4	2.0	1.5	1.2	1.7	1.0	1.0	0.9	0.6	0.3	0.3	0.1	0.2	0.1	§	0.3	+0.2
3.5–4.0	1.3	1.3	1.0	0.8	0.9	0.8	0.7	0.7	0.6	0.4	0.1	0.3	0.4	0.1	§	0.1	0.0
4.5–5.0	1.0	0.5	0.7	0.4	0.7	0.4	0.1	0.3	0.2	0.2	0.0	0.2	0.2	0.1	§	0.0	0.0
5.5–6.0 (High)	0.6	0.2	0.9	0.7	0.2	0.2	0.4	0.7	0.1	0.2	0.2	0.0	0.1	0.1	§	0.0	-0.1
Race/Ethnicity (2-year average): ^b																	
White	1.9	1.5	1.2	1.2	1.1	0.9	0.7	0.7	0.6	0.4	0.3	0.3	0.3	0.2	§	0.1	-0.1
African American	0.8	0.9	0.8	0.8	0.6	0.4	0.4	0.6	0.6	0.4	0.3	0.3	0.3	0.3	§	0.2	-0.1
Hispanic	0.7	0.9	0.9	0.7	0.7	0.7	0.5	0.5	0.5	0.4	0.3	0.1	0.1	0.1	§	0.0	-0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 134
CIGARETTES: Trends in 30-Day Prevalence of Use of Half Pack a Day or More
by Subgroups in Grade 10

	Percentage who used half pack a day or more in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
Total	6.5	6.0	7.0	7.6	8.3	9.4	8.6	7.9	7.6	6.2	5.5	4.4	4.1	3.3	3.1	
Gender:																
Male	6.9	6.5	7.8	8.2	8.7	9.9	8.9	8.1	7.8	6.7	6.0	4.1	4.3	3.5	2.9	
Female	6.0	5.1	6.2	6.7	7.7	9.0	8.2	7.8	7.3	5.6	5.1	4.6	3.8	2.9	3.2	
College Plans:																
None or under 4 years	15.9	15.3	18.5	18.5	20.9	22.4	22.0	20.6	18.9	16.0	16.0	11.5	12.3	10.8	9.4	
Complete 4 years	4.4	4.0	4.6	5.2	6.1	7.1	6.3	5.6	5.7	4.5	3.8	3.1	2.8	2.2	2.2	
Region:																
Northeast	7.8	5.9	8.5	7.8	7.7	9.1	8.8	10.0	9.1	6.0	5.0	3.7	4.6	3.6	3.4	
Midwest	7.1	7.3	7.7	8.3	9.5	10.9	9.3	9.1	9.9	8.1	6.0	4.8	5.0	3.0	3.3	
South	7.2	5.5	7.1	8.7	10.3	11.0	10.2	8.9	7.8	7.1	6.5	5.5	4.5	4.3	3.7	
West	4.0	5.0	4.3	4.2	3.4	5.0	4.4	3.0	3.0	2.2	3.3	2.5	2.3	1.8	1.7	
Population Density:																
Large MSA	6.9	5.2	5.9	5.8	6.0	7.6	6.6	5.8	6.0	5.7	4.8	3.0	2.3	1.9	2.3	
Other MSA	5.8	5.8	6.7	8.5	9.2	9.5	8.0	7.7	7.6	5.6	5.0	4.7	4.0	3.3	2.6	
Non-MSA	7.6	6.9	8.7	7.8	9.2	11.5	12.0	11.0	9.4	7.8	7.5	5.7	7.3	5.2	5.4	
Parental Education: ^a																
1.0–2.0 (Low)	9.9	10.7	10.8	8.1	12.0	12.1	10.8	9.0	11.3	10.7	7.6	7.4	8.6	6.6	4.6	
2.5–3.0	8.9	6.9	8.5	10.1	12.2	13.1	11.4	11.6	10.4	8.5	8.3	7.0	6.3	5.3	4.9	
3.5–4.0	5.8	5.4	7.3	8.0	8.6	10.2	8.8	7.4	7.4	5.9	5.3	4.0	4.2	3.8	3.2	
4.5–5.0	4.7	4.7	4.3	5.4	5.0	6.2	6.5	5.9	5.5	4.3	3.9	2.4	2.2	1.4	1.8	
5.5–6.0 (High)	4.5	3.7	3.9	4.0	4.0	5.7	4.8	5.4	4.5	3.6	2.6	2.0	1.4	1.0	1.6	
Race/Ethnicity (2-year average): ^b																
White	—	7.4	7.7	8.6	9.3	10.5	11.0	10.4	9.9	8.5	7.1	6.2	5.5	4.7	3.9	
African American	—	0.8	0.6	1.2	1.8	1.6	1.7	1.8	1.5	1.9	1.7	1.2	1.0	1.0	1.2	
Hispanic	—	3.0	3.0	2.6	3.4	4.3	3.3	3.0	3.1	2.9	2.7	2.2	1.8	1.5	1.9	

Table continued on next page.

TABLE 134 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Use of Half Pack a Day or More by Subgroups in Grade 10

	Percentage who used half pack a day or more in last 30 days																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	3.3	2.7	2.0	2.4	2.4	1.9	1.5	1.5	1.2	1.0	0.6	0.7	0.7	0.5	0.6	0.3	-0.2*
Gender:																	
Male	3.1	2.9	2.4	2.7	3.0	2.5	1.7	1.9	1.5	1.1	0.8	0.7	0.8	0.7	§	0.2	-0.4 s
Female	3.3	2.5	1.7	2.0	1.8	1.2	1.3	1.1	0.9	0.8	0.4	0.4	0.4	0.4	§	0.1	-0.2
College Plans:																	
None or under 4 years	12.0	8.1	7.8	8.6	8.6	7.0	6.8	5.3	4.9	5.0	3.2	3.2	2.1	1.4	§	0.8	-0.6
Complete 4 years	2.1	2.0	1.3	1.7	1.7	1.3	1.0	1.0	0.8	0.5	0.3	0.3	0.4	0.3	§	0.1	-0.2 s
Region:																	
Northeast	3.2	2.5	1.2	2.2	2.6	2.3	1.1	1.4	0.9	0.9	0.5	0.9	0.3	0.1	§	0.5	+0.3
Midwest	4.5	4.2	3.3	2.7	2.7	2.2	1.5	1.8	1.7	1.0	0.9	0.7	0.8	0.7	§	0.4	-0.3
South	3.6	2.8	2.6	3.2	3.1	2.3	2.3	2.0	1.3	1.3	0.7	0.5	0.9	0.5	§	0.4	-0.1
West	1.3	1.3	0.7	1.0	1.1	0.5	0.8	0.7	0.8	0.7	0.4	0.7	0.5	0.7	§	0.1	-0.5
Population Density:																	
Large MSA	1.8	2.0	1.1	1.6	1.6	1.3	0.9	0.8	0.7	0.6	0.5	0.6	0.5	0.2	§	0.4	+0.2 s
Other MSA	3.0	2.7	2.2	2.3	2.4	1.7	1.4	1.4	0.8	0.7	0.4	0.4	0.6	0.6	§	0.2	-0.4
Non-MSA	6.3	4.0	3.1	3.7	3.8	3.0	2.6	2.7	3.0	2.4	1.5	1.4	1.3	1.0	§	0.6	-0.4
Parental Education: ^a																	
1.0–2.0 (Low)	7.1	4.9	3.4	4.8	4.2	3.0	3.9	2.6	1.9	2.0	0.9	0.9	1.3	0.6	§	0.3	-0.4
2.5–3.0	5.6	4.5	3.2	3.7	4.5	3.6	2.5	2.6	1.9	1.3	0.9	1.2	1.1	0.8	§	0.1	-0.7 s
3.5–4.0	3.1	2.6	2.4	2.4	2.3	1.8	1.3	1.5	1.5	1.2	0.4	0.5	0.7	0.6	§	0.3	-0.3
4.5–5.0	1.5	1.6	1.1	1.4	1.1	0.9	1.0	0.9	0.5	0.7	0.4	0.3	0.3	0.1	§	0.2	+0.1
5.5–6.0 (High)	1.6	1.2	0.7	1.1	1.0	0.5	0.5	0.5	0.4	0.1	0.3	0.3	0.3	0.4	§	0.1	-0.3
Race/Ethnicity (2-year average): ^b																	
White	3.8	3.8	3.2	2.8	2.9	2.6	2.1	1.9	1.8	1.2	0.9	0.7	0.8	0.8	§	0.4	-0.4 s
African American	1.3	1.2	0.9	1.1	1.4	1.3	0.9	0.8	1.0	0.9	0.6	0.4	0.4	0.3	§	0.5	+0.2
Hispanic	2.0	1.0	0.9	1.1	1.0	0.9	0.8	0.6	0.5	0.7	0.7	0.5	0.5	0.4	§	0.1	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 135
CIGARETTES: Trends in 30-Day Prevalence of Use of Half Pack a Day or More
by Subgroups in Grade 12

	Percentage who used half pack a day or more in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	17.9	19.2	19.4	18.8	16.5	14.3	13.5	14.2	13.8	12.3	12.5	11.4	11.4	10.6	11.2	11.3
Gender:																
Male	19.6	19.9	19.7	18.9	15.4	13.5	12.8	13.1	13.1	11.0	12.3	10.7	10.1	11.1	11.2	11.6
Female	16.1	18.0	18.9	18.0	17.1	14.7	13.8	14.7	13.6	12.8	12.0	11.6	12.5	9.7	10.7	10.8
College Plans:																
None or under 4 years	—	25.5	26.9	25.5	23.3	21.2	20.8	21.0	20.9	19.6	20.7	19.2	19.5	18.4	18.6	19.2
Complete 4 years	—	11.9	11.2	11.1	9.8	8.2	7.5	7.8	7.6	6.5	6.5	6.4	7.2	6.8	7.5	7.5
Region:																
Northeast	22.0	22.5	24.2	23.6	19.8	17.0	16.6	15.6	16.6	17.4	17.0	15.6	16.5	13.1	13.6	13.8
Midwest	18.8	20.3	20.3	19.8	17.4	15.4	16.0	17.3	17.1	13.0	14.9	12.3	12.3	11.5	14.2	13.7
South	16.8	19.0	18.5	17.0	16.1	14.5	12.0	13.3	12.4	11.3	9.7	10.0	9.4	10.1	9.7	9.4
West	11.3	12.4	11.5	12.2	10.8	8.3	7.3	7.1	6.4	7.4	7.6	6.5	8.1	7.7	6.9	8.3
Population Density:																
Large MSA	21.7	20.1	20.4	19.7	16.2	14.8	15.4	15.9	14.1	14.8	14.4	12.2	13.1	10.8	10.1	11.2
Other MSA	17.4	18.9	18.8	17.9	16.5	13.8	12.4	12.9	13.5	11.4	11.0	9.6	10.0	10.4	11.2	11.0
Non-MSA	15.9	19.0	19.5	19.3	16.7	14.7	13.6	14.2	14.0	11.5	12.9	13.3	12.5	10.7	12.1	12.1
Parental Education: ^a																
1.0–2.0 (Low)	18.6	21.3	20.0	19.2	19.5	16.3	16.0	17.1	17.1	16.4	16.1	15.6	13.8	11.2	11.5	10.2
2.5–3.0	17.7	21.4	22.2	21.0	17.6	16.8	15.6	15.9	15.2	14.8	14.2	13.3	13.9	12.4	13.5	13.4
3.5–4.0	13.9	17.4	18.3	16.9	15.2	12.8	12.5	13.3	11.9	10.5	12.0	11.0	10.5	10.3	10.7	11.6
4.5–5.0	15.9	15.9	14.8	15.4	12.6	10.3	10.1	10.1	10.5	8.0	9.5	6.8	8.9	8.6	9.2	10.2
5.5–6.0 (High)	9.1	15.9	14.6	14.5	13.6	9.8	8.8	9.3	9.3	7.9	5.4	7.4	8.4	8.3	8.4	7.9
Race/Ethnicity (2-year average): ^b																
White	—	—	20.3	20.2	18.6	16.4	15.0	15.1	15.2	14.2	13.6	13.1	12.9	12.9	12.9	13.3
African American	—	—	10.7	9.7	9.1	7.1	5.8	5.4	4.9	4.1	3.9	3.6	2.9	2.3	1.9	1.8
Hispanic	—	—	11.3	9.0	6.4	5.6	6.1	5.6	5.9	6.1	5.3	5.0	4.1	3.5	4.2	4.2

Table continued on next page.

TABLE 135 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Use of Half Pack a Day or More
by Subgroups in Grade 12

	Percentage who used half pack a day or more in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	10.7	10.0	10.9	11.2	12.4	13.0	14.3	12.6	13.2	11.3	10.3	9.1	8.4	8.0	6.9	5.9
Gender:																
Male	11.6	10.4	11.6	12.7	13.2	13.6	15.5	13.5	14.5	11.4	10.2	10.0	9.5	8.0	8.0	6.2
Female	9.5	9.2	9.9	9.5	11.1	12.0	12.6	11.1	11.5	10.8	10.2	7.9	6.6	7.5	5.3	5.2
College Plans:																
None or under 4 years	18.7	19.1	18.7	19.6	22.6	23.1	23.5	23.7	23.2	20.3	19.3	17.5	17.2	16.0	15.2	13.0
Complete 4 years	7.1	6.5	8.1	8.2	8.9	10.0	11.0	8.9	10.1	8.1	7.6	6.7	5.5	5.6	4.6	3.9
Region:																
Northeast	12.9	11.1	14.7	12.2	13.4	17.0	19.4	13.6	13.4	13.6	12.3	10.9	8.7	9.0	7.5	7.9
Midwest	14.1	11.0	12.5	15.3	14.2	15.1	16.9	16.8	15.0	13.7	14.8	13.6	10.1	8.8	8.7	6.7
South	8.9	10.2	10.4	10.8	12.6	12.0	12.3	11.8	13.9	10.6	7.6	8.6	8.9	8.9	7.6	6.0
West	7.2	6.8	6.0	5.9	8.4	6.5	8.2	7.5	9.0	7.5	6.6	3.1	5.0	4.1	3.1	2.9
Population Density:																
Large MSA	10.2	9.9	9.1	10.2	12.3	11.6	12.8	11.0	9.6	8.2	9.4	7.6	5.4	6.0	5.2	4.8
Other MSA	10.7	8.4	11.2	10.5	11.1	12.8	14.3	11.7	12.6	11.6	9.4	8.7	8.4	8.2	6.3	5.0
Non-MSA	11.1	13.1	11.7	13.7	14.7	14.4	16.0	16.5	18.0	14.5	13.3	11.9	12.2	10.0	10.4	9.2
Parental Education: ^a																
1.0–2.0 (Low)	12.5	11.0	10.7	9.5	13.5	13.6	11.9	12.5	15.0	16.1	10.8	7.4	8.7	8.6	6.8	5.4
2.5–3.0	12.4	12.7	12.5	13.7	15.1	14.8	16.4	14.9	16.6	13.8	12.5	12.0	10.5	10.9	10.0	8.4
3.5–4.0	10.7	9.6	10.4	11.0	12.7	12.3	13.9	13.8	13.1	10.7	10.6	10.3	8.7	8.2	7.2	5.8
4.5–5.0	7.9	8.1	10.0	10.4	9.8	12.6	14.5	10.3	10.7	9.3	9.2	6.8	5.7	6.1	5.1	4.6
5.5–6.0 (High)	9.0	5.7	8.3	8.8	9.1	10.8	11.2	7.4	9.5	7.1	6.2	5.4	4.9	4.7	3.2	3.3
Race/Ethnicity (2-year average): ^b																
White	13.1	12.3	12.8	13.7	14.2	15.2	16.9	16.9	15.9	15.1	13.6	12.4	10.9	10.0	9.1	7.9
African American	1.8	1.8	1.5	1.7	2.2	2.5	2.8	2.5	2.4	2.6	2.4	2.1	1.7	1.6	1.8	2.1
Hispanic	4.0	4.6	4.5	4.0	4.1	5.3	5.8	5.2	5.8	7.0	5.3	3.8	3.0	2.9	2.9	2.6

Table continued on next page.

TABLE 135 (cont.)
CIGARETTES: Trends in 30-Day Prevalence of Use of Half Pack a Day or More
by Subgroups in Grade 12

	Percentage who used half pack a day or more in last 30 days															2019-2021 change	
<i>Approximate weighted N =</i>	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Total	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	0.8	-0.1*
Gender:																	
Male															§	0.5	-0.6 s
Female															§	0.8	+0.2
College Plans:																	
None or under 4 years															§	1.9	-0.3
Complete 4 years															§	0.3	-0.2
Region:																	
Northeast															§	0.2	-0.4
Midwest															§	0.9	+0.2
South															§	0.9	-0.3
West															§	1.0	+0.1
Population Density:																	
Large MSA															§	0.6	+0.1
Other MSA															§	0.7	-0.1
Non-MSA															§	1.4	-0.7
Parental Education: ^a																	
1.0–2.0 (Low)															§	1.4	+0.4
2.5–3.0															§	0.6	-0.7
3.5–4.0															§	1.2	+0.2
4.5–5.0															§	0.3	-0.1
5.5–6.0 (High)															§	0.1	-0.5
Race/Ethnicity (2-year average): ^b																	
White															§	0.8	-0.6 s
African American															§	0.7	-0.2
Hispanic															§	1.0	+0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 136
TOBACCO USING A HOOKAH: Trends in Annual
Prevalence of Use by Subgroups in Grade 12

	Percent who used in last 12 months													2019– 2021 change
	1975– 2011	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	17.1	18.5	18.3	21.4	22.9	19.8	13.0	10.1	7.8	5.6	§	2.1	-3.5* sss
Gender:														
Male	—	19.0	20.0	19.9	21.6	24.9	22.2	15.2	9.4	7.0	6.6	§	2.3	-4.3 sss
Female	—	15.2	16.7	17.0	21.0	21.4	17.9	10.7	10.5	7.7	4.4	§	1.4	-3.0 ss
College Plans:														
None or under 4 years	—	19.4	19.8	17.9	21.3	23.7	20.6	18.5	10.8	11.5	8.1	§	3.1	-5.0 s
Complete 4 years	—	16.2	18.6	18.4	21.5	22.8	19.5	11.6	10.2	7.1	5.0	§	1.9	-3.1 sss
Region:														
Northeast	—	15.4	15.1	17.5	23.9	21.6	20.5	18.9	7.3	11.9	4.8	§	1.2	-3.6 s
Midwest	—	17.9	21.3	18.6	18.9	26.2	24.2	13.5	10.5	8.7	4.7	§	1.5	-3.2 s
South	—	15.3	14.5	12.1	16.5	20.1	16.6	8.2	8.7	5.8	5.8	§	2.9	-2.9 s
West	—	20.9	23.9	26.2	29.7	25.9	21.1	17.2	13.7	7.5	7.1	§	2.1	-5.1 s
Population Density:														
Large MSA	—	19.6	18.6	22.7	25.2	24.6	20.1	18.6	9.7	8.3	4.4	§	1.9	-2.5 s
Other MSA	—	18.3	21.2	19.3	23.6	23.5	20.2	11.0	9.7	8.5	5.6	§	2.2	-3.4 ss
Non-MSA	—	10.5	12.3	9.1	10.7	19.3	18.6	9.2	11.5	5.4	7.7	§	2.3	-5.3 ss
Parental Education: ^a														
1.0–2.0 (Low)	—	16.8	19.8	18.1	19.2	17.9	16.4	10.7	8.1	8.3	6.7	§	2.8	-3.9
2.5–3.0	—	16.3	17.6	16.8	20.6	20.2	19.8	14.6	9.3	7.8	9.6	§	2.7	-6.8 ss
3.5–4.0	—	18.7	18.1	19.4	23.7	22.9	21.3	15.2	12.7	10.5	6.3	§	2.5	-3.8 s
4.5–5.0	—	15.0	18.6	19.6	20.8	29.0	19.5	11.2	8.4	6.3	3.6	§	1.5	-2.1
5.5–6.0 (High)	—	21.3	20.6	17.6	20.4	22.0	23.8	10.6	12.7	5.9	3.4	§	2.0	-1.4
Race/Ethnicity (2-year data): ^b														
White	—	—	20.6	20.5	20.2	24.4	25.2	19.7	13.0	9.4	6.7	§	2.6	-4.0 sss
African American	—	—	6.0	5.9	8.6	10.3	9.5	8.7	8.3	7.3	4.8	§	4.4	-0.4
Hispanic	—	—	17.6	18.5	22.8	23.7	19.9	16.3	11.6	8.8	8.4	§	1.7	-6.7 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; *N* is one sixth of *N* indicated in Table S3. Beginning in 2016 data based on three of six forms; *N* is three sixths of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 137
SMALL CIGARS: Trends in Annual Prevalence
of Use by Subgroups in Grade 12

	Percent who used in last 12 months													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	23.1	19.5	19.9	20.4	18.9	15.9	15.6	13.3	9.2	7.8	§	3.4	-4.4* sss
Gender:														
Male	—	29.7	26.7	27.1	26.5	26.4	23.6	22.5	19.3	12.3	11.7	§	5.6	-6.1 ss
Female	—	16.3	12.3	12.9	14.8	11.8	9.3	8.5	7.8	6.5	3.5	§	1.2	-2.3 ss
College Plans:														
None or under 4 years	—	30.7	27.7	24.5	21.8	22.9	20.3	18.8	18.7	14.4	13.2	§	5.6	-7.6 ss
Complete 4 years	—	21.4	17.8	19.0	20.1	18.0	14.9	14.6	12.4	7.8	6.1	§	2.8	-3.3 sss
Region:														
Northeast	—	26.1	21.5	19.2	21.4	21.2	17.4	18.2	13.9	12.6	6.5	§	3.1	-3.5
Midwest	—	28.1	23.7	28.4	23.7	23.6	17.6	15.5	14.6	8.1	7.5	§	3.2	-4.3 s
South	—	19.8	14.2	12.8	19.8	18.2	14.4	14.2	13.0	9.2	8.9	§	3.6	-5.3 sss
West	—	20.8	21.8	20.5	17.1	14.2	15.9	16.7	11.9	7.9	6.3	§	3.3	-3.0
Population Density:														
Large MSA	—	22.7	16.2	21.6	18.5	16.0	10.7	13.6	11.5	4.6	5.3	§	3.9	-1.4
Other MSA	—	23.5	21.9	19.5	22.4	20.3	17.9	16.7	13.8	11.1	8.5	§	2.9	-5.6 sss
Non-MSA	—	22.5	18.4	18.3	18.3	19.4	19.7	16.1	14.9	12.5	9.8	§	3.5	-6.2 s
Parental Education: ^a														
1.0–2.0 (Low)	—	16.6	14.8	12.8	18.3	9.5	11.8	8.5	10.3	6.5	7.9	§	2.8	-5.0
2.5–3.0	—	23.6	16.2	18.6	17.5	17.6	15.9	15.8	11.1	9.8	6.0	§	4.0	-2.0
3.5–4.0	—	22.9	19.9	20.5	22.8	20.6	18.7	16.0	14.3	10.3	8.9	§	3.0	-6.0 sss
4.5–5.0	—	25.9	19.4	23.2	22.1	20.8	14.8	16.3	14.9	9.3	8.1	§	4.2	-4.0
5.5–6.0 (High)	—	23.5	26.5	19.7	18.1	21.5	19.9	19.1	15.5	9.1	9.4	§	2.9	-6.5 ss
Race/Ethnicity (2-year data): ^b														
White	—	—	25.8	24.3	23.9	24.2	22.0	20.6	19.6	15.1	11.3	§	6.6	-4.8 ss
African American	—	—	6.3	5.9	9.1	10.5	9.4	7.8	5.3	4.3	4.1	§	2.9	-1.2
Hispanic	—	—	17.9	13.6	16.0	13.7	10.0	10.4	8.5	5.9	5.2	§	2.3	-2.9 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 138
Flavored Little Cigars or Cigarillos: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days																			
	8th Graders										10th Graders									
	1991– 2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
Approximate weighted N =	—	14600	14,400	16,900	15,300	14,000	13,600	3,100	10,700		—	13000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	4.1	4.1	2.8	2.6	2.6	2.2	2.3	1.0	-1.2* sss	—	6.9	6.1	4.9	4.0	5.3	3.7	3.0	1.5	-2.2* sss
Gender:																				
Male	—	4.8	4.4	2.6	2.8	2.6	1.9	§	1.2	-0.7	—	8.2	7.5	5.8	4.9	6.1	4.3	§	1.9	-2.5 ss
Female	—	3.4	3.7	2.8	2.4	2.3	2.5	§	0.5	-2.0 sss	—	5.8	4.8	3.7	3.0	4.4	3.2	§	0.9	-2.3 sss
College Plans:																				
None or under 4 years	—	10.0	9.8	10.0	8.5	7.1	6.2	§	2.1	-4.1 sss	—	12.9	15.1	10.4	10.9	13.1	8.0	§	3.7	-4.3 ss
Complete 4 years	—	3.5	3.6	2.2	2.1	1.9	1.6	§	0.8	-0.8 s	—	6.5	5.2	4.2	3.0	4.4	2.9	§	1.1	-1.8 ss
Region:																				
Northeast	—	2.3	3.0	3.5	2.5	1.8	0.6	§	0.7	+0.1	—	7.3	5.2	2.9	4.0	3.3	2.0	§	2.6	+0.7
Midwest	—	3.3	3.1	2.2	2.6	2.7	2.4	§	1.5	-0.9	—	7.2	6.2	4.2	3.3	4.9	3.8	§	1.6	-2.2 ss
South	—	4.8	4.8	3.3	2.5	2.6	2.9	§	0.8	-2.1 sss	—	7.7	7.0	6.2	3.5	7.5	5.0	§	1.5	-3.5 sss
West	—	4.7	4.5	1.9	2.8	3.2	2.1	§	1.0	-1.1	—	5.3	5.5	5.4	5.3	3.8	2.6	§	0.7	-1.9 ss
Population Density:																				
Large MSA	—	4.2	3.4	2.2	2.2	2.2	1.9	§	0.7	-1.2 ss	—	7.0	4.9	4.6	4.5	3.3	2.2	§	0.6	-1.6 ss
Other MSA	—	4.4	4.6	2.7	2.7	2.3	2.2	§	1.0	-1.2 s	—	6.3	5.2	4.8	2.8	6.3	4.1	§	1.7	-2.3 ss
Non-MSA	—	3.2	3.9	4.0	3.0	4.0	2.9	§	1.4	-1.6	—	8.5	10.6	6.0	6.0	6.0	5.2	§	2.6	-2.6 s
Parental Education: a																				
1.0–2.0 (Low)	—	9.1	8.0	4.3	3.1	5.0	3.3	§	0.4	-2.9 s	—	10.5	7.9	5.8	5.2	2.9	6.7	§	1.9	-4.9 s
2.5–3.0	—	6.0	4.5	3.1	4.9	2.8	3.2	§	2.9	-0.3	—	7.2	8.5	4.3	6.2	7.6	6.0	§	1.0	-5.0 sss
3.5–4.0	—	3.9	3.1	2.2	2.6	3.7	2.3	§	0.8	-1.5 s	—	10.8	6.2	5.1	4.1	6.7	4.1	§	1.1	-3.0 sss
4.5–5.0	—	2.2	3.4	2.7	1.9	1.5	1.7	§	0.8	-1.0	—	5.1	4.6	4.3	2.6	4.0	1.6	§	2.5	+0.9
5.5–6.0 (High)	—	2.1	2.6	1.8	1.3	1.0	1.3	§	0.8	-0.5	—	3.7	4.8	5.5	1.9	4.7	2.8	§	0.3	-2.5 sss
Race/Ethnicity (2-year data): b																				
White	—	—	2.6	2.7	2.2	2.0	2.0	§	1.3	-0.7	—	—	6.8	5.7	4.1	4.6	4.6	§	2.6	-2.0 sss
African American	—	—	4.2	3.1	2.8	3.5	3.2	§	1.4	-1.9	—	—	6.4	3.9	3.3	3.6	4.2	§	3.0	-1.1
Hispanic	—	—	6.7	4.4	2.9	3.1	2.8	§	2.5	-0.3	—	—	5.9	4.7	4.6	5.1	5.3	§	1.2	-4.1 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on two of four forms; N is one third of N indicated in S1 and S2. For 12th graders only: Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 138 (cont.)
Flavored Little Cigars or Cigarillos: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days									
	<u>12th Graders</u>									
	1976– <u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	2019– 2021 change
Approximate weighted N =	—	12400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	11.9	11.4	9.5	10.1	8.9	7.7	§	1.9	-5.8* sss
Gender:										
Male	—	16.1	15.4	13.4	12.6	12.0	11.3	§	2.7	-8.6 sss
Female	—	8.2	7.6	5.8	7.5	5.2	4.6	§	1.2	-3.4 sss
College Plans:										
None or under 4 years	—	19.6	18.4	12.6	17.4	14.4	11.0	§	2.5	-8.5 sss
Complete 4 years	—	9.9	9.8	8.8	8.4	7.3	6.6	§	1.9	-4.7 sss
Region:										
Northeast	—	15.5	11.7	10.8	12.3	6.0	5.7	§	2.3	-3.4
Midwest	—	14.5	11.5	11.1	11.1	9.3	9.0	§	1.5	-7.6 sss
South	—	10.7	11.7	8.4	10.1	10.0	7.6	§	3.0	-4.6 ss
West	—	9.0	10.6	8.9	7.6	8.4	7.9	§	0.1	-7.8 ss
Population Density:										
Large MSA	—	7.9	9.0	7.4	9.1	6.8	5.5	§	3.0	-2.5
Other MSA	—	12.6	12.5	9.5	9.5	8.7	8.0	§	1.3	-6.8 sss
Non-MSA	—	15.4	12.7	12.2	12.6	12.8	10.5	§	1.8	-8.6 sss
Parental Education: a										
1.0–2.0 (Low)	—	10.7	13.1	8.5	7.4	8.5	9.1	§	3.3	-5.8
2.5–3.0	—	14.8	14.3	9.7	10.0	9.0	8.5	§	1.2	-7.3 sss
3.5–4.0	—	12.4	10.6	9.9	10.5	9.1	8.3	§	2.1	-6.2 sss
4.5–5.0	—	10.4	11.8	10.4	10.7	9.2	5.2	§	1.3	-3.9 ss
5.5–6.0 (High)	—	9.6	6.9	6.9	10.9	7.5	9.0	§	3.2	-5.8
Race/Ethnicity (2-year data): b										
White	—	—	13.4	12.2	12.0	11.6	9.6	§	2.1	-7.6 sss
African American	—	—	8.8	8.0	6.5	5.6	4.7	§	3.1	-1.6
Hispanic	—	—	9.5	8.1	5.9	5.8	7.3	§	1.8	-5.6 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on two of four forms; N is one third of N indicated in S1 and S2. For 12th graders only: Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 139
Regular Little Cigars or Cigarillos: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days																			
	8th Graders										10th Graders									
	1991– 2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
Approximate weighted N =	—	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	—	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	—	—
Total	—	2.5	3.3	1.9	1.6	1.6	1.6	1.4	0.8	-0.8* s	—	4.4	3.8	3.0	3.0	3.1	2.6	2.4	1.2	-1.5* sss
Gender:																				
Male	—	2.9	4.1	1.8	1.7	1.9	1.5	§	1.2	-0.3	—	5.9	4.9	3.8	3.9	3.5	3.3	§	1.3	-2.0 ss
Female	—	2.0	2.5	1.6	1.5	1.2	1.7	§	0.3	-1.4 sss	—	2.9	2.9	2.0	2.0	2.5	2.0	§	0.7	-1.3 ss
College Plans:																				
None or under 4 years	—	8.5	9.5	6.9	5.4	5.5	4.6	§	1.6	-3.0 s	—	9.9	9.1	8.1	9.2	8.3	5.6	§	4.1	-1.5
Complete 4 years	—	2.0	2.8	1.4	1.3	1.1	1.2	§	0.7	-0.5	—	4.0	3.3	2.4	2.1	2.4	2.1	§	0.6	-1.5 sss
Region:																				
Northeast	—	1.7	3.2	2.6	1.4	1.3	0.3	§	0.3	0.0	—	4.7	4.4	2.3	2.9	2.4	1.2	§	0.6	-0.6
Midwest	—	1.5	2.9	1.4	1.5	1.8	1.7	§	1.5	-0.2	—	3.9	4.1	2.7	2.7	3.1	2.0	§	1.8	-0.2
South	—	3.8	4.0	2.3	1.7	1.5	2.4	§	0.6	-1.8 ss	—	6.0	4.5	3.5	3.0	4.4	3.9	§	1.6	-2.3 ss
West	—	2.0	2.6	1.0	1.8	1.7	1.2	§	0.9	-0.3	—	2.5	1.9	3.1	3.3	1.4	2.1	§	0.3	-1.8 ss
Population Density:																				
Large MSA	—	2.2	3.3	1.2	1.5	1.3	1.1	§	0.6	-0.5	—	4.1	2.8	2.7	2.9	2.1	1.4	§	0.9	-0.6
Other MSA	—	2.7	3.5	1.9	1.7	1.5	2.1	§	1.0	-1.1	—	3.6	3.5	2.8	2.1	3.4	3.2	§	1.0	-2.2 sss
Non-MSA	—	2.5	3.1	2.8	1.6	2.1	1.3	§	0.8	-0.5	—	7.1	6.4	4.1	5.2	3.7	3.3	§	2.2	-1.1
Parental Education: a																				
1.0–2.0 (Low)	—	5.4	8.3	2.3	2.7	3.4	2.1	§	0.6	-1.5	—	7.4	5.5	4.3	3.2	3.3	4.6	§	1.3	-3.4 s
2.5–3.0	—	3.3	3.1	2.1	3.1	1.8	1.6	§	1.7	+0.1	—	3.8	4.7	2.1	3.7	3.8	4.8	§	0.6	-4.1 sss
3.5–4.0	—	1.9	2.6	1.4	1.3	2.0	1.5	§	1.1	-0.3	—	6.6	3.3	4.0	3.9	4.3	2.6	§	1.3	-1.3
4.5–5.0	—	1.9	2.7	1.7	1.0	0.8	1.1	§	0.8	-0.3	—	3.6	2.6	2.2	2.0	1.4	0.9	§	1.4	+0.6
5.5–6.0 (High)	—	1.8	2.6	1.2	0.4	0.8	0.9	§	0.5	-0.3	—	2.8	4.2	3.3	1.2	3.2	2.2	§	0.4	-1.7 s
Race/Ethnicity (2-year data): b																				
White	—	—	2.1	2.0	1.3	1.1	1.1	§	0.7	-0.3	—	—	4.2	3.6	2.8	2.9	2.6	§	1.9	-0.7
African American	—	—	3.4	2.4	2.1	2.7	2.0	§	1.8	-0.3	—	—	4.8	3.4	2.9	3.2	3.8	§	2.0	-1.8
Hispanic	—	—	4.0	2.9	1.8	2.0	1.8	§	0.7	-1.1	—	—	3.7	2.9	3.1	3.0	3.2	§	1.5	-1.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on two of four forms; N is one third of N indicated in S1 and S2. For 12th graders only: Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 139 (cont.)
Regular Little Cigars or Cigarillos: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days									
	<u>12th Graders</u>									
	1976– 2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
<i>Approximate weighted N =</i>	—	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	7.0	7.8	6.1	6.6	5.8	4.9	§	1.8	-3.1* sss
Gender:										
Male	—	9.6	10.9	8.7	8.4	7.9	6.6	§	2.4	-4.2 ss
Female	—	4.6	4.6	3.3	4.3	3.1	3.3	§	1.1	-2.2 s
College Plans:										
None or under 4 years	—	13.6	14.6	8.8	12.0	11.1	8.3	§	1.9	-6.4 ss
Complete 4 years	—	5.3	6.1	5.3	5.4	4.3	4.0	§	1.9	-2.1 s
Region:										
Northeast	—	9.0	7.8	7.3	7.3	4.0	2.5	§	2.9	+0.4
Midwest	—	8.7	7.5	6.4	6.5	5.9	5.1	§	1.2	-3.9 ss
South	—	6.8	8.7	6.4	7.8	7.1	5.2	§	2.2	-3.0 s
West	—	4.1	6.6	4.5	4.1	4.9	6.1	§	0.8	-5.4
Population Density:										
Large MSA	—	4.8	6.0	5.4	4.9	5.4	3.7	§	2.9	-0.7
Other MSA	—	7.0	8.1	6.0	5.9	5.5	4.7	§	1.0	-3.7 ss
Non-MSA	—	9.5	9.9	7.3	10.6	7.1	7.4	§	1.8	-5.6 ss
Parental Education: a										
1.0–2.0 (Low)	—	8.4	7.9	3.0	5.2	6.3	8.1	§	2.4	-5.6 s
2.5–3.0	—	7.9	10.9	6.5	7.8	5.6	5.4	§	1.2	-4.2 ss
3.5–4.0	—	7.4	6.4	6.8	7.2	6.3	4.6	§	1.6	-3.0 s
4.5–5.0	—	5.3	7.9	6.3	5.8	5.6	2.0	§	1.9	-0.1
5.5–6.0 (High)	—	5.7	4.7	4.6	5.5	4.1	6.7	§	3.0	-3.6
Race/Ethnicity (2-year data): b										
White	—	—	8.3	7.8	7.3	6.8	5.5	§	1.8	-3.7 sss
African American	—	—	6.5	6.8	5.6	4.9	3.7	§	3.1	-0.6
Hispanic	—	—	6.1	4.9	3.6	4.1	5.3	§	1.1	-4.2 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on two of four forms; *N* is one third of *N* indicated in S1 and S2. For 12th graders only: Data based on two of six forms; *N* is two sixths of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 140
Large Cigars: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days																			
	8th Graders										10th Graders									
	1991– 2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
<i>Approximate weighted N =</i>	—	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	—	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000		
Total	—	1.9	2.4	1.5	1.5	1.7	1.3	1.5	1.1	-0.2*	—	3.9	3.4	2.3	2.6	2.8	2.1	1.2	1.3	-0.8*
Gender:																				
Male	—	2.8	3.6	1.8	1.6	2.0	1.5	§	1.3	-0.2	—	5.8	4.8	3.2	3.9	4.1	3.1	§	1.9	-1.3
Female	—	1.1	1.2	1.1	1.3	1.2	1.2	§	0.4	-0.8 s	—	2.2	2.1	1.2	1.3	1.3	1.2	§	0.7	-0.6
College Plans:																				
None or under 4 years	—	6.2	7.5	6.3	5.3	5.2	3.3	§	2.4	-0.9	—	7.2	10.4	7.1	8.7	6.1	4.2	§	3.3	-0.8
Complete 4 years	—	1.6	1.9	1.1	1.1	1.3	1.1	§	0.9	-0.2	—	3.7	2.7	1.7	1.8	2.3	1.8	§	0.9	-0.9
Region:																				
Northeast	—	1.8	1.6	2.5	1.5	1.6	0.7	§	0.6	-0.1	—	6.1	3.7	2.2	1.2	1.9	1.2	§	2.8	+1.6
Midwest	—	0.8	1.8	0.8	1.8	1.7	1.1	§	1.4	+0.3	—	3.8	3.2	3.0	2.9	3.2	1.7	§	1.8	+0.1
South	—	2.7	2.9	1.9	1.2	1.4	1.6	§	0.7	-1.0	—	4.1	3.7	1.7	2.5	3.7	3.0	§	0.9	-2.1 s
West	—	1.6	2.5	0.8	1.7	2.3	1.4	§	1.7	+0.3	—	2.1	2.8	2.5	3.5	1.5	1.7	§	0.6	-1.2 s
Population Density:																				
Large MSA	—	2.2	1.8	0.8	1.5	0.8	0.9	§	0.4	-0.5	—	4.4	2.8	1.9	2.8	2.0	1.5	§	1.0	-0.5
Other MSA	—	1.9	2.7	1.5	1.5	1.9	1.5	§	1.4	-0.1	—	2.9	2.6	2.1	1.7	2.9	2.2	§	1.5	-0.7
Non-MSA	—	1.3	2.5	2.6	1.3	2.5	1.6	§	1.5	-0.1	—	5.5	6.4	3.4	4.8	3.8	2.9	§	1.6	-1.3
Parental Education: a																				
1.0–2.0 (Low)	—	4.1	4.5	2.4	2.4	3.8	1.5	§	1.9	+0.3	—	4.2	5.7	2.5	1.6	3.0	1.5	§	2.3	+0.8
2.5–3.0	—	2.8	2.7	1.5	2.3	1.7	2.2	§	2.7	+0.4	—	4.6	2.9	1.8	3.3	3.1	2.9	§	0.8	-2.1 s
3.5–4.0	—	1.5	1.6	1.4	1.7	2.0	1.0	§	1.5	+0.4	—	4.9	2.3	1.9	3.7	3.9	3.4	§	0.7	-2.7 ss
4.5–5.0	—	1.2	1.8	1.0	1.0	1.0	0.8	§	0.4	-0.4	—	3.3	3.1	2.2	1.6	1.4	1.1	§	2.3	+1.2
5.5–6.0 (High)	—	1.2	1.5	1.1	0.9	0.6	1.0	§	1.1	0.0	—	3.8	3.3	3.4	1.9	2.9	1.7	§	0.6	-1.1
Race/Ethnicity (2-year data): b																				
White	—	—	1.6	1.5	1.0	0.9	1.1	§	0.8	-0.3	—	—	4.1	3.3	2.8	3.1	2.8	§	1.5	-1.4 ss
African American	—	—	2.0	1.5	0.9	1.5	1.7	§	1.9	+0.2	—	—	3.5	2.2	1.7	1.8	1.3	§	1.4	+0.1
Hispanic	—	—	3.4	2.3	1.7	2.1	1.6	§	0.3	-1.3 s	—	—	3.2	2.3	2.0	2.2	2.6	§	0.3	-2.3 s

Source: The Monitoring the Future study, the University of Michigan.

Notes: '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on two of four forms; N is one third of N indicated in S1 and S2. For 12th graders only: Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 140 (cont.)
Large Cigars: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days									
	12th Graders									
	1976– 2013	2014	2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
Approximate weighted N =	—	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	6.4	7.0	6.5	5.6	5.2	5.3	§	2.3	-3.0* sss
Gender:										
Male	—	10.7	11.9	10.7	8.5	8.3	8.6	§	3.7	-4.9 ss
Female	—	2.4	2.4	2.4	2.6	1.9	2.0	§	1.0	-1.0
College Plans:										
None or under 4 years	—	11.9	12.2	8.3	9.1	8.3	7.4	§	3.1	-4.3
Complete 4 years	—	5.0	5.8	6.0	4.7	4.3	4.6	§	2.2	-2.4 ss
Region:										
Northeast	—	8.7	8.3	6.1	6.9	4.9	3.7	§	3.4	-0.3
Midwest	—	7.4	6.7	7.3	6.1	5.3	5.6	§	0.5	-5.0 sss
South	—	5.9	6.2	5.5	5.2	6.5	5.4	§	3.4	-1.9
West	—	4.4	7.8	7.6	4.9	3.3	6.2	§	0.9	-5.3 sss
Population Density:										
Large MSA	—	4.4	5.9	6.1	4.7	3.8	4.4	§	2.9	-1.5
Other MSA	—	6.4	7.1	6.4	5.4	5.7	5.2	§	1.1	-4.0 sss
Non-MSA	—	8.7	8.6	7.2	7.3	6.7	7.0	§	3.9	-3.1
Parental Education: a										
1.0–2.0 (Low)	—	4.8	7.9	4.9	2.4	3.6	5.0	§	7.0	+2.0
2.5–3.0	—	6.1	5.8	5.8	5.7	4.6	5.4	§	0.4	-5.1 sss
3.5–4.0	—	6.4	7.1	5.7	5.8	5.7	4.4	§	1.1	-3.3 ss
4.5–5.0	—	7.0	8.2	7.5	6.3	5.0	3.4	§	2.8	-0.6
5.5–6.0 (High)	—	6.1	5.6	5.6	6.7	5.9	9.0	§	4.3	-4.7
Race/Ethnicity (2-year data): b										
White	—	—	8.3	8.5	7.5	6.7	6.2	§	2.9	-3.4 sss
African American	—	—	3.2	3.1	2.7	2.5	1.9	§	1.1	-0.8
Hispanic	—	—	5.0	4.7	3.5	3.2	4.1	§	0.6	-3.5 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on two of four forms; N is one third of N indicated in S1 and S2. For 12th graders only: Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 141
Any Vaping:^c Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

Percentage who used in last 30 days

	8th Graders									10th Graders									12th Graders								
	1991–								2019– 2021 change	1991–								2019– 2021 change	1976–								2019– 2021 change
	2014	2015	2016	2017	2018	2019	2020	2021		2014	2015	2016	2017	2018	2019	2020	2021		2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	14,400	16,900	16,900	14,000	13,600	3,100	10,700	—	15,600	14,700	13,500	14,300	14,000	4,800	11,000	—	12,900	11,800	12,600	13,300	12,900	3,500	8,300			
Total	—	8.0	6.2‡	6.6	10.4	12.2	12.5	8.9	-3.3* sss	—	14.2	11.0‡	13.1	21.7	25.0	23.5	15.6	-9.4* sss	—	16.3	12.5‡	16.6	26.7	30.9	28.2	24.0	-6.9* sss
Gender:																											
Male	—	8.6	7.0‡	7.2	10.5	10.9	§	6.9	-4.1 sss	—	17.2	12.1‡	13.3	21.2	24.9	§	12.9	-12.1 sss	—	20.9	16.1‡	21.0	30.0	33.0	§	22.2	-10.7 sss
Female	—	7.4	5.4‡	6.2	10.1	13.4	§	10.1	-3.3 ss	—	11.1	9.7‡	12.7	21.8	24.9	§	17.3	-7.7 sss	—	11.2	9.1‡	12.8	23.8	28.8	§	25.3	-3.5
College Plans:																											
None or under 4 years	—	15.7	18.2‡	14.6	19.2	20.5	§	15.6	-4.9	—	26.0	19.3‡	22.0	25.8	33.8	§	24.3	-9.6 sss	—	25.8	17.7‡	21.8	29.2	36.6	§	27.2	-9.4 sss
Complete 4 years	—	7.3	5.3‡	5.8	9.3	11.1	§	7.6	-3.5 sss	—	13.0	9.9‡	12.0	21.0	23.3	§	13.4	-10.0 sss	—	14.2	11.6‡	15.1	26.4	29.1	§	22.4	-6.7 ss
Region:																											
Northeast	—	6.1	6.5‡	6.9	7.3	10.3	§	4.9	-5.4 sss	—	14.0	8.4‡	12.8	24.0	25.0	§	15.5	-9.5 sss	—	16.0	12.8‡	18.3	32.9	29.4	§	27.2	-2.1
Midwest	—	7.9	7.2‡	7.6	12.5	13.2	§	9.6	-3.6 s	—	14.7	11.3‡	15.9	23.4	25.3	§	16.6	-8.7 sss	—	17.2	14.8‡	17.1	25.6	31.2	§	24.0	-7.2 s
South	—	8.8	5.7‡	5.5	9.9	13.1	§	10.8	-2.2	—	16.0	12.1‡	11.9	20.8	24.6	§	17.7	-6.9 s	—	15.4	10.7‡	15.0	24.7	30.7	§	21.5	-9.2 sss
West	—	8.0	6.1‡	7.5	11.6	11.4	§	8.0	-3.4	—	10.9	11.1‡	12.6	19.4	25.2	§	11.3	-13.9 sss	—	17.3	13.6‡	18.3	26.9	32.5	§	25.5	-7.1
Population Density:																											
Large MSA	—	6.7	5.6‡	5.3	9.5	9.8	§	4.5	-5.3 sss	—	13.9	13.2‡	14.7	22.3	21.6	§	10.8	-10.8 sss	—	13.5	12.8‡	16.6	22.2	25.3	§	23.1	-2.2
Other MSA	—	7.9	5.9‡	7.3	11.0	11.3	§	10.1	-1.2	—	13.8	9.0‡	12.3	21.7	25.5	§	16.4	-9.0 sss	—	17.6	11.8‡	16.1	29.5	32.4	§	22.2	-10.2 sss
Non-MSA	—	10.0	8.0‡	7.1	10.2	18.4	§	13.4	-5.0 s	—	15.9	12.2‡	12.8	20.7	29.2	§	21.8	-7.5 ss	—	17.8	14.0‡	18.5	28.1	36.7	§	29.8	-6.9 s
Parental Education: ^a																											
1.0–2.0 (Low)	—	9.7	10.3‡	10.1	12.7	16.3	§	15.6	-0.7	—	14.2	9.2‡	12.9	20.3	25.5	§	19.7	-5.8	—	11.5	13.2‡	12.2	17.9	23.7	§	20.3	-3.4
2.5–3.0	—	9.8	7.8‡	8.7	11.3	16.3	§	13.6	-2.6	—	16.1	15.1‡	14.4	22.2	28.7	§	18.8	-9.9 sss	—	18.6	12.2‡	17.4	25.7	32.3	§	26.4	-5.8
3.5–4.0	—	10.6	6.7‡	6.9	12.2	13.7	§	12.2	-1.5	—	18.3	11.4‡	13.5	22.9	27.7	§	17.7	-10.0 sss	—	17.2	11.9‡	14.8	29.1	33.7	§	25.6	-8.1 ss
4.5–5.0	—	5.6	5.8‡	5.6	9.7	11.1	§	7.0	-4.1 sss	—	13.8	10.6‡	13.6	23.1	24.0	§	13.7	-10.2 sss	—	18.0	13.1‡	20.7	32.3	30.9	§	22.9	-7.9 ss
5.5–6.0 (High)	—	6.5	4.0‡	5.8	7.5	8.8	§	5.7	-3.0 s	—	9.8	9.4‡	12.0	21.4	21.4	§	12.5	-8.9 sss	—	11.4	12.2‡	17.5	31.0	31.7	§	24.3	-7.4 s
Race/Ethnicity (2-year data): ^b																											
White	—	—	6.4‡	—	8.7	13.0	§	11.2	-1.8	—	—	14.8‡	—	21.5	28.6	§	21.3	-7.3 sss	—	—	17.4‡	—	28.0	35.8	§	30.4	-5.3 sss
African American	—	—	6.2‡	—	3.3	5.3	§	6.6	+1.3	—	—	6.4‡	—	7.0	14.2	§	12.5	-1.6	—	—	6.5‡	—	8.3	12.7	§	16.3	+3.6
Hispanic	—	—	8.4‡	—	9.3	11.5	§	9.0	-2.5	—	—	10.3‡	—	14.1	20.2	§	18.1	-2.1	—	—	11.5‡	—	15.2	21.2	§	20.2	-1.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. * — * indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

‡ indicates some change in the question. See relevant footnote for that drug. See relevant figures to assess the impact of the change.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on one of four forms; N is one third of N indicated in S1 and S2 until 2019. In 2019, N is two thirds of N indicated. For 12th graders only: Data based on two of six forms until 2019N is two sixths of N indicated in Table S3.

In 2019, N is two thirds of N indicated. Beginning in 2020, all estimates for 8th, 10th, and 12th graders based on all forms.

In 2017, the surveys switched from asking about vaping in general to asking separately about vaping nicotine, marijuana, and just flavoring. Beginning in 2017, data presented for any vaping are based on these new questions. The 2017 results by race/ethnicity are based on data from the old version of the question in 2016 combined with the new version in 2017.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data

that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cIn 2017, the surveys switched from asking about vaping in general to asking separately about vaping nicotine, marijuana, and just flavoring. Beginning in 2017, data presented for any vaping are based on these new questions.

TABLE 142
Vaping Nicotine: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days																				
	<u>8th Graders</u>							<u>10th Graders</u>							<u>12th Graders</u>						
	1991– 2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2016	2017	2018	2019	2020	2021	2019– 2021 change	1976– 2016	2017	2018	2019	2020	2021	2019– 2021 change
<i>Approximate weighted N =</i>	—	15,300	14,000	13,600	3,100	10,700		—	13,500	14,300	14,000	4,800	11,000		—	12,600	13,300	12,900	3,500	8,300	
Total	—	3.5	6.1	9.6	10.5	7.6	-2.0* s	—	8.2	16.1	19.9	19.3	13.1	-6.8* sss	—	11.0	20.9	25.5	24.7	19.6	-5.8* ss
Gender:																					
Male	—	3.7	6.8	8.8	§	5.9	-2.8 ss	—	8.2	16.2	20.3	§	10.5	-9.9 sss	—	14.3	24.1	28.1	§	18.6	-9.5 sss
Female	—	3.3	5.3	10.3	§	8.6	-1.7	—	8.1	15.9	19.5	§	14.8	-4.7 s	—	8.0	18.4	22.9	§	20.4	-2.5
College Plans:																					
None or under 4 years	—	9.2	12.8	16.4	§	13.4	-3.0	—	13.5	20.4	26.8	§	21.0	-5.8 s	—	14.5	22.9	29.9	§	23.7	-6.2 s
Complete 4 years	—	2.9	5.3	8.7	§	6.4	-2.3 ss	—	7.6	15.6	18.7	§	11.2	-7.6 sss	—	10.2	20.7	23.9	§	17.9	-6.0 ss
Region:																					
Northeast	—	3.2	4.4	8.4	§	4.1	-4.3 sss	—	7.9	18.1	20.0	§	13.0	-6.9 ss	—	11.3	25.3	24.9	§	21.5	-3.5
Midwest	—	4.1	9.2	11.2	§	8.0	-3.2 s	—	10.3	18.3	20.9	§	14.7	-6.2 ss	—	11.5	20.9	27.4	§	21.2	-6.2
South	—	3.2	5.2	9.6	§	9.2	-0.3	—	7.5	15.7	19.3	§	15.2	-4.2	—	10.1	19.2	25.2	§	18.5	-6.6 s
West	—	3.8	6.1	9.0	§	7.1	-1.9	—	7.5	12.9	20.0	§	8.6	-11.4 sss	—	11.7	20.8	23.9	§	18.5	-5.4
Population Density:																					
Large MSA	—	2.5	6.0	7.4	§	3.5	-3.9 ss	—	9.7	17.9	15.9	§	8.2	-7.7 sss	—	10.0	17.2	19.4	§	17.6	-1.8
Other MSA	—	3.7	6.7	9.1	§	8.7	-0.3	—	7.1	14.8	20.2	§	14.1	-6.1 ss	—	10.7	23.1	26.7	§	17.6	-9.2 sss
Non-MSA	—	4.5	5.0	14.6	§	11.9	-2.7	—	8.5	16.4	26.0	§	19.4	-6.6 s	—	13.0	22.3	32.6	§	27.7	-4.8
Parental Education: a																					
1.0–2.0 (Low)	—	5.9	5.7	11.6	§	12.6	+1.0	—	5.7	16.4	19.3	§	16.6	-2.7	—	7.8	11.4	15.2	§	12.5	-2.6
2.5–3.0	—	3.8	5.8	13.3	§	11.6	-1.7	—	10.1	16.4	22.6	§	16.2	-6.4 ss	—	11.7	17.9	25.8	§	22.6	-3.2
3.5–4.0	—	3.8	8.1	10.7	§	10.7	0.0	—	8.2	14.7	21.8	§	14.7	-7.1 sss	—	9.6	21.5	28.0	§	22.2	-5.8 s
4.5–5.0	—	3.5	5.8	9.4	§	6.2	-3.2 ss	—	8.5	17.9	20.0	§	11.6	-8.4 sss	—	14.7	28.2	27.0	§	18.9	-8.1 ss
5.5–6.0 (High)	—	2.8	4.8	6.9	§	4.9	-2.0	—	7.9	18.1	17.9	§	10.2	-7.7 sss	—	10.5	27.1	28.1	§	19.1	-9.1 sss
Race/Ethnicity (2-year data): b																					
White	—	—	5.5	10.4	§	9.8	-0.6	—	—	16.4	24.2	§	18.8	-5.4 sss	—	—	22.9	31.4	§	27.1	-4.3 s
African American	—	—	1.6	2.9	§	4.7	+1.8	—	—	2.7	7.6	§	9.5	+1.8	—	—	4.6	8.6	§	12.8	+4.1
Hispanic	—	—	4.7	6.8	§	6.9	+0.1	—	—	8.5	14.0	§	12.9	-1.1	—	—	7.7	12.3	§	15.1	+2.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on one of four forms; *N* is one third of *N* indicated in S1 and S2 until 2019. In 2019, *N* is two thirds of *N* indicated. For 12th graders only: Data based on two of six forms until 2019; *N* is two sixths of *N* indicated in Table S3.

In 2019, *N* is two thirds of *N* indicated. Beginning in 2020, all estimates for 8th, 10th, and 12th graders based on all forms.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 143
Vaping Marijuana: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days																				
	<u>8th Graders</u>							<u>10th Graders</u>							<u>12th Graders</u>						
	1991– 2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2016	2017	2018	2019	2020	2021	2019– 2021 change	1976– 2016	2017	2018	2019	2020	2021	2019– 2021 change
Approximate weighted N =	—	15,300	14,000	13,600	3,100	10,700		—	13,500	14,300	14,000	4,800	11,000		—	12,600	13,300	12,900	3,500	8,300	
Total	—	1.6	2.6	3.9	4.2	2.9	-1.0* ss	—	4.3	7.0	12.6	11.3	8.4	-4.2* sss	—	4.9	7.5	14.0	12.2	12.4	-1.6*
Gender:																					
Male	—	1.8	2.9	3.3	§	2.0	-1.3 ss	—	4.7	6.9	13.6	§	7.1	-6.5 sss	—	7.2	9.5	14.7	§	10.8	-3.9 sss
Female	—	1.6	2.3	4.5	§	3.5	-1.0 s	—	3.8	6.9	11.6	§	9.0	-2.6 s	—	2.9	5.5	13.0	§	13.7	+0.7
College Plans:																					
None or under 4 years	—	4.9	8.7	8.2	§	5.3	-2.9 s	—	8.7	9.3	19.5	§	13.6	-5.9 ss	—	5.8	8.3	17.1	§	13.4	-3.6 s
Complete 4 years	—	1.2	1.8	3.3	§	2.3	-0.9 ss	—	3.7	6.6	11.2	§	7.2	-4.0 sss	—	4.5	7.1	12.7	§	11.7	-1.0
Region:																					
Northeast	—	1.3	1.2	3.5	§	2.2	-1.3	—	6.1	9.5	13.9	§	9.8	-4.1 s	—	6.7	13.9	16.4	§	15.3	-1.0
Midwest	—	1.3	1.1	3.5	§	3.6	+0.1	—	4.4	6.1	11.7	§	7.9	-3.8 s	—	4.1	3.8	12.9	§	11.5	-1.4
South	—	1.6	2.7	3.1	§	2.8	-0.4	—	2.6	4.8	11.2	§	9.0	-2.2	—	3.5	5.0	11.6	§	9.1	-2.5 s
West	—	2.5	5.2	5.9	§	3.1	-2.8 sss	—	5.4	9.7	15.0	§	6.9	-8.1 sss	—	6.9	10.7	18.9	§	16.2	-2.7
Population Density:																					
Large MSA	—	0.8	2.7	3.2	§	1.7	-1.4 ss	—	5.7	8.8	14.2	§	6.8	-7.5 sss	—	7.1	9.2	14.1	§	12.8	-1.3
Other MSA	—	1.9	3.1	4.6	§	3.5	-1.1	—	3.8	6.8	13.0	§	9.2	-3.8 s	—	4.3	7.4	15.0	§	12.9	-2.0
Non-MSA	—	2.2	1.3	3.7	§	3.6	0.0	—	3.4	4.6	8.8	§	9.4	+0.5	—	2.8	4.5	11.3	§	10.5	-0.8
Parental Education: a																					
1.0–2.0 (Low)	—	3.0	5.3	6.1	§	6.2	+0.1	—	5.1	8.8	15.7	§	12.2	-3.4	—	4.8	7.7	13.6	§	12.2	-1.4
2.5–3.0	—	2.9	3.9	6.6	§	4.8	-1.8	—	5.5	7.8	14.1	§	11.1	-3.0	—	5.4	7.9	14.8	§	14.5	-0.4
3.5–4.0	—	1.2	3.2	3.9	§	3.9	0.0	—	4.1	7.7	15.1	§	9.7	-5.4 ss	—	3.8	7.4	14.2	§	12.9	-1.4
4.5–5.0	—	1.0	1.4	2.9	§	1.9	-1.0	—	3.7	6.3	10.6	§	7.2	-3.4 ss	—	4.9	7.5	12.3	§	10.6	-1.7
5.5–6.0 (High)	—	1.2	0.6	2.4	§	1.8	-0.6	—	4.5	6.5	10.0	§	5.8	-4.2 ss	—	5.7	8.5	14.5	§	13.1	-1.5
Race/Ethnicity (2-year data): b																					
White	—	—	1.5	2.6	§	2.9	+0.3	—	—	6.1	10.9	§	9.7	-1.1	—	—	6.7	12.0	§	12.7	+0.7
African American	—	—	1.2	2.0	§	1.7	-0.3	—	—	2.6	7.8	§	6.5	-1.3	—	—	2.5	6.5	§	7.7	+1.3
Hispanic	—	—	3.1	5.1	§	3.7	-1.4	—	—	5.7	11.5	§	10.5	-1.1	—	—	6.8	11.8	§	12.3	+0.5

Source: The Monitoring the Future study, the University of Michigan.

Notes: '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on one of four forms; N is one third of N indicated in S1 and S2 until 2019. In 2019, N is two thirds of N indicated. For 12th graders only: Data based on two of six forms until 2019; N is two sixths of N indicated in Table S3.

In 2019, N is two thirds of N indicated. Beginning in 2020, all estimates for 8th, 10th, and 12th graders based on all forms.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 144
Vaping Just Flavoring: Trends in 30-Day
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 30 days																				
	8th Graders							10th Graders							12th Graders						
	1991– 2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2016	2017	2018	2019	2020	2021	2019– 2021 change	1976– 2016	2017	2018	2019	2020	2021	2019– 2021 change
Approximate weighted N =	—	15,300	14,000	13,600	3,100	10,700		—	13,500	14,300	14,000	4,800	11,000		—	12,600	13,300	12,900	3,500	8,300	
Total	—	5.3	8.1	7.7	6.8	4.6	-3.1* sss	—	9.2	13.1	10.5	10.4	6.3	-4.2* sss	—	9.7	13.5	10.7	8.4	7.4	-3.3* sss
Gender:																					
Male	—	6.0	8.1	6.9	§	2.9	-4.0 sss	—	9.4	12.9	10.1	§	4.4	-5.6 sss	—	11.6	15.0	11.0	§	5.5	-5.6 sss
Female	—	4.8	8.0	8.6	§	5.9	-2.6 ss	—	8.9	13.1	11.0	§	8.0	-3.0 s	—	7.9	11.8	10.2	§	9.2	-1.0
College Plans:																					
None or under 4 years	—	11.5	13.2	13.0	§	8.2	-4.9 ss	—	15.4	16.2	15.7	§	10.0	-5.7 sss	—	13.0	15.3	13.5	§	7.7	-5.9 sss
Complete 4 years	—	4.8	7.4	7.1	§	4.0	-3.1 sss	—	8.4	12.6	9.6	§	5.3	-4.2 sss	—	8.6	13.0	9.8	§	7.0	-2.8 ss
Region:																					
Northeast	—	5.2	5.6	6.4	§	2.1	-4.2 sss	—	8.8	13.4	9.1	§	6.1	-3.0	—	10.4	15.2	7.8	§	6.3	-1.5
Midwest	—	7.0	10.3	8.2	§	5.2	-3.0 s	—	10.8	15.2	9.5	§	5.7	-3.8 ss	—	9.8	13.7	10.7	§	8.3	-2.4
South	—	4.1	7.5	8.8	§	6.0	-2.8 ss	—	8.5	12.8	11.4	§	7.8	-3.6 s	—	9.1	13.6	11.3	§	7.9	-3.4 s
West	—	5.8	8.7	6.4	§	3.6	-2.8 s	—	8.8	11.3	11.2	§	4.7	-6.5 sss	—	9.9	11.9	12.0	§	6.6	-5.5 ss
Population Density:																					
Large MSA	—	4.5	7.2	5.4	§	2.1	-3.3 sss	—	8.6	11.9	6.3	§	3.9	-2.4 ss	—	8.8	9.7	7.0	§	6.3	-0.7
Other MSA	—	5.8	8.6	6.8	§	5.3	-1.6	—	9.6	13.7	11.1	§	6.6	-4.5 sss	—	9.7	15.6	11.4	§	6.3	-5.1 sss
Non-MSA	—	5.4	8.3	13.6	§	7.4	-6.2 sss	—	8.8	13.7	16.2	§	9.8	-6.4 sss	—	11.0	15.2	15.3	§	11.9	-3.5
Parental Education: a																					
1.0–2.0 (Low)	—	7.5	8.5	10.9	§	9.7	-1.2	—	8.7	14.9	15.0	§	11.0	-4.0	—	7.5	11.4	11.4	§	7.9	-3.5
2.5–3.0	—	7.4	8.1	10.8	§	7.6	-3.2 s	—	10.3	14.5	13.8	§	7.8	-6.0 sss	—	9.4	13.9	13.0	§	8.4	-4.6 ss
3.5–4.0	—	5.5	9.9	8.2	§	6.1	-2.0	—	10.6	14.5	11.5	§	7.3	-4.2 ss	—	8.1	16.9	12.0	§	9.4	-2.7
4.5–5.0	—	4.6	7.8	7.2	§	3.5	-3.7 sss	—	8.7	12.4	8.7	§	4.6	-4.1 sss	—	12.1	14.5	8.4	§	6.2	-2.2 s
5.5–6.0 (High)	—	4.5	6.9	5.3	§	3.0	-2.3 s	—	8.6	11.7	7.2	§	4.8	-2.4 s	—	11.1	11.3	7.8	§	5.2	-2.6
Race/Ethnicity (2-year data): b																					
White	—	—	7.2	9.0	§	5.9	-3.1 sss	—	—	13.1	13.6	§	8.6	-5.0 sss	—	—	13.5	12.7	§	7.9	-4.8 sss
African American	—	—	2.4	3.9	§	3.1	-0.9	—	—	4.8	6.5	§	6.2	-0.4	—	—	5.6	6.3	§	5.9	-0.4
Hispanic	—	—	7.2	7.8	§	5.3	-2.5	—	—	9.4	10.0	§	8.5	-1.5	—	—	9.6	9.8	§	8.9	-0.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on one of four forms; N is one third of N indicated in S1 and S2 until 2019. In 2019, N is two thirds of N indicated. For 12th graders only: Data based on two of six forms until 2019; N is two sixths of N indicated in Table S3.

In 2019, N is two thirds of N indicated. Beginning in 2020, all estimates for 8th, 10th, and 12th graders based on all forms.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 145
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800
Total	6.9	7.0	6.6	7.7	7.1	7.1	5.5	4.8	4.5	4.2	4.0	3.3	4.1	4.1	3.3
Gender:															
Male	12.7	12.5	10.9	12.8	11.8	11.4	9.9	8.1	6.9	6.7	6.9	5.4	6.7	6.4	5.3
Female	1.4	2.0	2.7	2.4	2.9	2.9	1.5	1.5	2.1	1.8	1.4	1.3	1.8	1.7	1.5
College Plans:															
None or under 4 years	12.7	17.1	15.5	16.7	15.4	16.4	12.6	13.9	13.2	11.4	14.6	10.2	12.8	12.3	10.8
Complete 4 years	6.1	5.5	5.3	6.5	6.0	5.6	4.6	3.8	3.5	3.4	2.9	2.6	3.3	3.2	2.4
Region:															
Northeast	5.0	4.9	3.4	6.1	5.4	4.9	3.2	2.7	2.5	2.7	3.7	2.7	3.1	3.4	1.4
Midwest	7.1	7.5	7.2	7.1	7.6	8.3	6.8	4.3	5.3	4.8	4.0	3.9	3.5	4.2	3.2
South	9.5	9.3	8.0	9.9	8.7	8.1	6.7	6.9	5.9	5.8	5.4	4.1	5.9	5.5	5.3
West	3.5	4.4	6.3	6.0	5.0	5.9	4.1	3.9	2.9	1.9	2.1	1.5	2.5	2.0	1.4
Population Density:															
Large MSA	4.8	4.2	3.3	4.6	4.1	4.2	3.6	2.9	1.8	2.4	2.4	1.5	2.6	1.9	1.3
Other MSA	6.2	6.9	6.8	6.4	6.7	7.1	4.7	4.1	3.9	3.9	3.5	2.9	3.7	3.6	2.8
Non-MSA	10.4	10.3	9.9	13.0	11.2	10.6	9.0	8.5	8.9	7.0	7.0	6.2	6.9	7.7	7.0
Parental Education: ^a															
1.0–2.0 (Low)	11.4	7.8	9.4	8.9	10.6	6.3	8.3	5.4	6.6	7.4	5.0	4.5	6.8	6.9	5.7
2.5–3.0	8.4	8.5	7.5	8.4	9.9	8.8	6.0	5.1	5.7	5.2	5.4	5.1	5.1	6.0	4.9
3.5–4.0	6.7	7.0	7.5	8.7	7.0	7.2	6.5	5.9	4.5	4.5	3.7	3.2	4.1	3.5	3.1
4.5–5.0	4.8	7.0	5.2	6.1	5.0	6.8	4.8	4.4	3.3	2.9	2.5	2.4	3.1	3.2	2.4
5.5–6.0 (High)	6.1	4.6	4.9	6.8	5.8	5.9	3.7	3.9	3.1	3.0	4.2	2.5	2.7	2.9	1.8
Race/Ethnicity (2-year average): ^b															
White	—	8.3	8.0	8.1	8.9	8.8	7.6	6.1	5.4	5.2	4.8	4.1	3.9	4.4	4.2
African American	—	1.8	2.7	3.2	2.6	2.2	2.6	2.3	2.3	2.7	2.2	1.6	2.7	3.0	2.0
Hispanic	—	4.2	4.0	5.0	5.7	5.2	4.6	4.5	4.6	3.7	3.3	4.0	4.7	4.0	2.6

Table continued on next page.

TABLE 145 (cont.)
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 30 days																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	3.7	3.2	3.5	3.7	4.1	3.5	2.8	2.8	3.0	3.2	2.5	1.7	2.1	2.5	2.3	1.6	-0.9* s
Gender:																	
Male	5.3	4.7	5.5	6.3	6.3	4.9	4.0	3.8	3.8	4.0	3.6	2.2	3.0	3.3	§	2.1	-1.2 s
Female	2.1	1.7	1.5	1.4	1.9	1.9	1.5	1.9	2.2	2.3	1.4	1.3	1.1	1.6	§	1.3	-0.3
College Plans:																	
None or under 4 years	11.3	8.9	12.7	11.1	14.0	11.1	11.5	9.2	9.6	11.3	9.1	7.2	6.7	7.5	§	5.1	-2.5
Complete 4 years	2.9	2.7	2.7	3.1	3.2	2.8	2.1	2.4	2.4	2.4	2.0	1.2	1.4	1.8	§	0.9	-0.9 ss
Region:																	
Northeast	2.9	1.5	2.5	3.2	3.9	2.9	2.6	1.8	1.3	1.7	1.9	1.9	2.0	1.0	§	0.3	-0.7 s
Midwest	4.0	2.9	3.4	3.0	4.8	3.3	2.1	1.4	2.7	2.8	2.4	1.4	2.0	2.5	§	2.4	-0.2
South	4.6	4.1	5.2	5.6	5.0	4.3	3.6	4.7	3.7	4.0	3.3	1.9	1.5	3.8	§	2.3	-1.5 s
West	2.2	2.8	1.8	1.8	1.9	2.6	2.0	2.0	3.2	3.0	1.6	1.6	3.1	1.5	§	0.9	-0.6
Population Density:																	
Large MSA	1.9	1.9	2.8	2.5	2.3	1.8	1.4	1.7	1.8	1.8	1.0	1.3	1.2	1.4	§	1.1	-0.2
Other MSA	3.1	3.1	3.1	3.6	4.7	3.7	3.1	2.6	2.7	2.8	2.6	1.6	2.0	2.4	§	1.5	-0.9
Non-MSA	7.1	5.3	5.4	5.8	5.0	5.4	4.2	4.9	5.2	5.8	4.7	2.7	3.5	4.5	§	2.7	-1.8
Parental Education: ^a																	
1.0–2.0 (Low)	5.9	4.9	5.4	5.3	5.2	6.6	4.2	4.1	5.6	4.4	3.4	2.9	1.8	4.0	§	3.2	-0.9
2.5–3.0	3.9	3.7	5.1	5.5	6.0	4.2	3.8	3.5	4.4	4.0	2.8	2.3	3.1	3.5	§	1.4	-2.0 ss
3.5–4.0	3.7	3.1	3.8	4.9	4.1	3.3	3.2	3.7	3.0	3.6	3.2	2.2	2.9	3.0	§	2.7	-0.3
4.5–5.0	3.1	2.5	2.4	2.9	3.3	2.7	2.0	1.6	1.9	2.5	2.0	1.0	1.5	2.0	§	1.4	-0.7
5.5–6.0 (High)	3.4	2.3	2.3	1.4	2.5	1.5	1.4	1.8	1.9	1.9	1.2	0.5	0.9	1.4	§	1.4	+0.1
Race/Ethnicity (2-year average): ^b																	
White	3.7	3.8	3.9	4.4	4.9	4.6	3.6	3.1	3.4	3.5	3.4	2.6	1.9	2.1	§	2.6	+0.6
African American	2.1	2.3	2.4	2.6	2.2	1.8	1.8	2.2	2.0	2.2	2.4	1.9	1.8	2.3	§	1.3	-1.0
Hispanic	3.1	2.9	2.4	2.5	2.6	3.2	3.1	2.4	2.6	2.7	2.2	1.9	2.0	2.3	§	0.4	-1.9 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996 and on two of four forms beginning in 1997; *N* is one half of *N* indicated in Table S1. In 2011 snus and dissolvable tobacco were added to the list of examples.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 146
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	
Total	10.0	9.6	10.4	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	5.6	
Gender:																
Male	18.7	18.1	19.3	19.2	17.2	15.0	14.9	13.8	12.2	11.4	12.7	9.9	9.6	9.0	9.7	
Female	1.3	1.8	2.0	2.1	2.1	2.3	2.7	1.7	1.3	1.3	1.6	2.1	1.3	1.0	1.6	
College Plans:																
None or under 4 years	16.9	17.5	20.2	19.9	20.3	16.3	18.5	17.8	13.2	13.9	16.0	13.6	13.0	12.2	13.3	
Complete 4 years	8.4	8.0	8.4	8.5	7.8	7.2	7.2	5.7	5.4	4.8	5.4	4.8	4.1	3.9	4.5	
Region:																
Northeast	8.6	5.3	8.0	9.0	7.6	6.8	9.3	6.5	5.2	4.6	4.9	4.7	4.5	5.1	4.6	
Midwest	11.0	9.6	10.0	10.0	11.0	9.5	7.1	7.9	8.1	6.2	7.0	4.8	4.9	3.7	5.7	
South	11.6	11.4	11.8	11.7	10.9	10.2	10.2	9.5	7.9	7.7	9.6	8.3	7.5	7.3	7.0	
West	7.8	10.9	11.1	10.9	7.7	6.0	8.2	4.6	4.0	4.5	3.0	5.1	3.5	3.0	4.5	
Population Density:																
Large MSA	5.9	6.4	6.5	6.2	5.9	5.5	4.2	3.7	4.6	5.6	4.1	4.5	3.7	3.0	3.2	
Other MSA	9.2	9.3	10.1	10.9	9.2	8.4	8.3	5.7	5.3	4.3	5.7	6.1	4.8	4.3	5.3	
Non-MSA	14.7	13.3	14.1	13.9	15.0	12.2	14.7	15.1	11.3	9.8	12.5	8.2	9.2	9.0	9.4	
Parental Education: ^a																
1.0–2.0 (Low)	6.6	10.1	10.9	9.4	9.6	8.1	9.0	6.8	7.2	7.4	6.9	6.7	7.4	3.7	4.4	
2.5–3.0	12.1	11.0	12.2	12.5	10.4	9.7	9.4	8.2	7.0	6.4	8.9	8.1	5.0	5.8	6.7	
3.5–4.0	10.6	10.5	10.9	10.2	10.9	8.3	10.3	8.6	7.3	6.3	7.1	5.5	4.9	5.2	6.0	
4.5–5.0	9.3	7.6	9.9	9.8	9.8	8.5	7.2	6.9	6.1	6.2	5.7	5.4	5.7	4.4	5.6	
5.5–6.0 (High)	8.6	8.1	7.0	8.9	6.0	7.7	8.3	5.2	4.8	4.0	4.8	5.2	4.3	4.4	3.8	
Race/Ethnicity (2-year average): ^b																
White	—	11.4	12.0	12.5	12.0	11.0	10.4	10.0	8.7	7.5	7.5	7.8	6.9	6.1	6.6	
African American	—	2.9	2.3	2.3	2.5	2.5	2.8	2.3	1.6	2.0	3.2	2.6	2.5	2.7	2.5	
Hispanic	—	6.2	6.1	4.3	3.6	4.0	4.6	4.8	4.8	4.5	4.0	4.0	4.1	3.3	3.1	

Table continued on next page.

TABLE 146 (cont.)
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	5.7	6.1	5.0	6.5	7.5	6.6	6.4	6.4	5.3	4.9	3.5	3.8	3.9	3.2	3.5	1.7	-1.6* sss
Gender:																	
Male	10.2	10.2	8.2	11.1	13.0	11.5	11.2	11.1	8.9	7.9	5.8	6.1	6.2	5.3	§	2.4	-2.9 sss
Female	1.2	2.0	2.0	2.0	2.0	1.9	1.9	2.0	1.9	1.8	1.3	1.5	1.4	1.4	§	0.8	-0.6
College Plans:																	
None or under 4 years	14.9	14.0	14.6	17.5	21.8	17.9	16.8	18.6	17.9	12.3	10.1	10.2	10.2	6.2	§	4.7	-1.4
Complete 4 years	4.5	5.0	3.8	5.3	5.8	5.2	5.3	5.1	4.1	4.0	2.8	2.9	2.9	2.7	§	1.0	-1.7 sss
Region:																	
Northeast	4.5	5.9	3.3	6.6	6.7	5.6	4.3	6.4	4.1	4.8	2.7	3.0	3.4	1.9	§	0.9	-1.0
Midwest	7.5	6.3	7.2	7.7	9.2	6.4	7.9	6.9	5.7	5.1	3.3	4.4	4.2	3.1	§	1.3	-1.8 ss
South	6.0	8.3	6.5	7.7	9.1	7.9	8.1	6.9	6.8	5.5	4.6	3.8	5.1	3.5	§	2.8	-0.8
West	4.1	3.3	2.2	3.3	4.0	5.4	4.0	5.4	3.6	3.6	2.9	3.8	2.0	3.7	§	0.9	-2.8 s
Population Density:																	
Large MSA	3.9	4.1	3.4	4.8	5.1	4.6	3.6	5.1	4.1	3.5	2.7	3.4	3.3	1.8	§	1.0	-0.8 s
Other MSA	5.3	5.8	4.4	6.3	7.0	6.1	6.3	5.3	4.4	4.3	2.9	2.6	3.4	3.3	§	1.6	-1.7 s
Non-MSA	9.6	9.9	8.7	9.3	12.1	10.2	10.5	11.2	9.5	8.6	6.8	7.2	5.9	5.4	§	3.1	-2.3 s
Parental Education: ^a																	
1.0–2.0 (Low)	7.4	6.1	5.0	7.9	6.5	6.4	7.2	5.6	4.7	4.7	2.4	3.4	3.0	4.6	§	3.6	-1.0
2.5–3.0	7.1	6.9	6.5	7.4	9.1	8.8	7.7	8.4	6.7	7.1	4.9	3.6	4.6	3.6	§	2.3	-1.3
3.5–4.0	5.7	6.4	5.4	7.4	8.7	7.2	7.5	6.9	5.8	4.2	4.0	4.6	3.8	2.5	§	1.4	-1.1
4.5–5.0	4.9	6.1	4.6	5.4	7.0	5.2	6.1	5.9	4.3	5.0	3.0	3.6	4.2	3.5	§	1.1	-2.4 ss
5.5–6.0 (High)	5.3	4.8	2.9	5.6	6.0	5.3	3.8	5.3	4.6	3.8	2.8	3.6	4.0	2.6	§	1.7	-0.8
Race/Ethnicity (2-year average): ^b																	
White	6.9	7.3	7.3	7.5	9.1	9.1	8.0	8.1	7.7	6.5	5.7	5.3	5.5	5.0	§	3.0	-1.9 ss
African American	3.2	3.3	2.8	2.6	2.3	2.2	2.3	2.2	2.5	2.8	2.4	1.9	1.4	1.8	§	3.0	+1.2
Hispanic	3.4	2.7	2.6	2.9	2.8	3.4	4.4	3.4	3.0	3.1	1.9	1.7	2.4	2.7	§	0.6	-2.0 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996 and on two of four forms beginning in 1997; N is one half of N indicated in Table S2. In 2011 snus and dissolvable tobacco were added to the list of examples.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 147
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990 ^a
<i>Approximate weighted N =</i>	—	—	—	—	—	—	—	—	—	—	—	15,200	16,300	16,300	16,700	—
Total	—	—	—	—	—	—	—	—	—	—	—	11.5	11.3	10.3	8.4	—
Gender:																
Male	—	—	—	—	—	—	—	—	—	—	—	22.3	22.8	19.9	15.9	—
Female	—	—	—	—	—	—	—	—	—	—	—	1.6	0.7	1.7	1.2	—
College Plans:																
None or under 4 years	—	—	—	—	—	—	—	—	—	—	—	14.5	15.5	13.1	9.6	—
Complete 4 years	—	—	—	—	—	—	—	—	—	—	—	9.8	9.0	8.8	7.7	—
Region:																
Northeast	—	—	—	—	—	—	—	—	—	—	—	9.5	7.3	5.9	5.0	—
Midwest	—	—	—	—	—	—	—	—	—	—	—	13.5	11.3	10.8	8.3	—
South	—	—	—	—	—	—	—	—	—	—	—	12.2	13.7	12.1	9.8	—
West	—	—	—	—	—	—	—	—	—	—	—	9.3	11.7	10.9	9.1	—
Population Density:																
Large MSA	—	—	—	—	—	—	—	—	—	—	—	9.0	6.4	7.7	6.8	—
Other MSA	—	—	—	—	—	—	—	—	—	—	—	8.9	10.5	8.5	7.6	—
Non-MSA	—	—	—	—	—	—	—	—	—	—	—	17.1	17.5	16.1	11.7	—
Parental Education: ^b																
1.0–2.0 (Low)	—	—	—	—	—	—	—	—	—	—	—	8.6	11.7	10.7	5.3	—
2.5–3.0	—	—	—	—	—	—	—	—	—	—	—	14.4	11.5	10.7	7.0	—
3.5–4.0	—	—	—	—	—	—	—	—	—	—	—	11.5	12.1	10.6	9.0	—
4.5–5.0	—	—	—	—	—	—	—	—	—	—	—	10.4	11.7	11.8	10.2	—
5.5–6.0 (High)	—	—	—	—	—	—	—	—	—	—	—	7.7	8.1	7.2	8.4	—
Race/Ethnicity (2-year average): ^c																
White	—	—	—	—	—	—	—	—	—	—	—	—	12.9	12.0	10.6	—
African American	—	—	—	—	—	—	—	—	—	—	—	—	2.1	4.5	4.5	—
Hispanic	—	—	—	—	—	—	—	—	—	—	—	—	4.4	5.2	5.1	—

Table continued on next page.

TABLE 147 (cont.)
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															
	1991 ^a	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	—	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	—	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	7.6	6.1
Gender:																
Male	—	20.8	19.7	20.3	23.6	19.5	18.7	15.6	15.5	14.4	14.2	12.2	12.5	12.2	12.7	11.0
Female	—	2.0	2.3	2.6	1.8	1.1	1.2	1.5	1.3	1.3	1.6	1.2	1.0	1.6	1.9	1.5
College Plans:																
None or under 4 years	—	18.0	14.9	15.8	18.7	17.6	16.9	14.3	10.5	15.8	13.0	10.8	12.8	11.9	15.2	13.4
Complete 4 years	—	9.4	9.4	9.3	9.9	7.6	7.4	7.1	7.6	5.4	6.1	4.8	4.8	4.9	5.4	4.1
Region:																
Northeast	—	8.2	9.6	12.0	9.6	8.4	6.9	2.6	4.3	5.3	5.4	5.3	6.3	5.1	8.0	5.3
Midwest	—	12.3	13.6	14.7	16.7	12.6	13.4	11.8	8.9	11.1	9.9	7.8	5.7	7.4	10.5	7.0
South	—	12.5	11.1	9.7	11.9	9.2	9.0	10.5	10.7	7.3	8.5	7.9	7.9	8.4	7.5	7.5
West	—	11.1	7.0	8.5	8.6	8.5	9.1	7.3	7.0	6.3	6.2	3.9	6.0	4.3	4.0	3.4
Population Density:																
Large MSA	—	5.9	7.1	7.5	12.5	8.6	6.5	4.7	4.9	4.2	4.4	3.4	3.4	3.8	4.3	4.8
Other MSA	—	11.1	9.9	11.3	9.5	7.4	7.4	7.7	8.5	7.9	8.0	5.7	6.9	5.6	7.8	4.6
Non-MSA	—	16.9	15.0	14.7	16.7	15.3	17.9	16.1	11.7	11.4	11.5	11.9	10.4	12.1	11.3	10.9
Parental Education: ^b																
1.0–2.0 (Low)	—	14.9	7.0	12.3	9.8	6.3	5.8	6.1	5.4	4.3	6.3	4.1	9.3	6.1	5.3	4.9
2.5–3.0	—	12.4	11.6	12.9	11.5	10.4	10.7	9.0	9.1	9.9	7.5	5.6	5.9	8.8	9.7	8.0
3.5–4.0	—	12.4	10.8	9.8	12.8	9.1	10.4	9.8	8.8	8.9	8.6	7.4	6.5	4.6	7.8	6.4
4.5–5.0	—	8.0	13.3	11.1	12.8	11.4	9.1	9.6	8.5	6.2	6.2	7.3	7.6	7.9	8.0	6.1
5.5–6.0 (High)	—	10.6	7.8	10.2	11.6	8.1	9.9	7.4	7.9	5.7	10.3	4.6	6.2	5.4	3.7	3.5
Race/Ethnicity (2-year average): ^c																
White	—	—	13.8	13.8	13.8	13.0	12.2	11.8	11.0	10.5	10.3	9.7	8.5	8.2	8.7	8.8
African American	—	—	2.0	1.9	2.1	2.7	2.2	1.4	1.5	1.5	1.2	1.0	1.0	0.6	0.3	0.5
Hispanic	—	—	6.0	5.4	7.6	8.1	5.3	4.3	3.9	3.8	3.2	2.6	3.1	3.9	4.9	3.8

Table continued on next page.

TABLE 147 (cont.)
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	6.6	6.5	8.4	8.5	8.3	7.89	8.1	8.4	6.1	6.6	4.9	4.2	3.5	§	2.2	-1.3* s
Gender:																
Male	11.9	11.8	15.8	15.7	14.2	13.49	14.6	14.3	10.7	11.9	9.9	7.6	5.7	§	3.6	-2.0
Female	1.2	1.0	1.7	1.7	1.8	1.58	1.4	2.1	1.6	1.5	0.7	1.0	1.1	§	0.7	-0.4
College Plans:																
None or under 4 years	11.7	12.4	14.2	14.4	17.5	14.96	16.1	15.0	14.3	11.7	11.4	8.3	7.1	§	2.9	-4.2 ss
Complete 4 years	5.3	5.0	7.3	7.3	6.4	6.01	6.2	6.5	4.4	5.1	3.4	3.0	2.5	§	2.0	-0.6
Region:																
Northeast	5.3	7.5	7.1	7.6	8.2	7.14	5.6	9.6	5.8	6.7	7.6	3.0	2.5	§	2.1	-0.5
Midwest	9.1	7.2	14.1	12.2	7.8	10.49	11.2	11.8	6.9	7.4	3.5	6.3	3.0	§	1.7	-1.3
South	7.9	6.1	7.1	7.8	8.9	7.73	9.3	5.9	6.1	7.5	5.6	4.9	4.5	§	3.1	-1.4
West	2.8	5.6	4.6	6.7	7.7	5.92	5.2	8.4	5.7	4.3	3.4	1.5	2.7	§	1.4	-1.3
Population Density:																
Large MSA	3.9	2.6	4.4	5.9	5.3	6.35	7.1	3.8	3.1	3.4	2.6	1.7	1.6	§	2.2	+0.7
Other MSA	6.8	6.2	7.7	7.9	8.4	6.39	7.4	9.8	5.1	7.3	5.4	4.3	2.5	§	1.8	-0.7
Non-MSA	10.1	11.8	15.0	13.8	12.4	14.11	11.4	11.1	13.5	10.2	7.8	8.4	9.2	§	3.1	-6.1 sss
Parental Education: ^b																
1.0–2.0 (Low)	5.3	3.4	2.8	6.1	2.2	5.26	7.9	6.3	6.6	5.5	1.9	0.6	2.9	§	0.4	-2.5
2.5–3.0	9.8	7.2	10.4	8.2	9.3	10.04	7.9	9.3	7.1	7.1	5.4	5.0	4.0	§	2.4	-1.7
3.5–4.0	5.3	6.8	9.4	8.7	8.2	8.46	6.7	8.1	6.5	6.7	6.3	5.2	3.9	§	2.9	-1.0
4.5–5.0	6.6	6.5	7.7	9.8	9.9	8.43	10.5	9.5	6.0	7.8	6.0	4.9	3.2	§	2.9	-0.3
5.5–6.0 (High)	5.8	6.3	8.7	7.5	6.3	4.72	7.7	7.9	4.9	6.2	3.6	4.5	3.4	§	1.2	-2.2
Race/Ethnicity (2-year average): ^c																
White	8.3	8.6	10.5	12.1	11.8	11.31	11.0	11.4	10.2	9.6	9.4	7.7	5.6	§	6.3	+0.7
African American	1.8	2.0	1.5	2.2	1.8	1.0	2.4	2.6	1.3	2.0	2.4	1.7	1.3	§	1.3	+0.1
Hispanic	2.7	2.3	2.5	3.7	3.9	2.53	2.3	3.4	3.6	1.8	1.1	1.2	1.5	§	6.2	+4.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; N is one sixth of N indicated in Table S3. In 2011 snus and dissolvable tobacco were added to the list of examples.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aPrevalence of smokeless tobacco use was not asked of 12th graders in 1990 and 1991. Prior to 1990 the prevalence-of-use question on smokeless tobacco was located near the end of one 12th-grade questionnaire form, whereas after 1991 the question was placed earlier and in a different form. This shift could explain the discontinuities between the corresponding data.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 148
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 8

	Percentage who used daily in last 30 days														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>
Total	1.6	1.8	1.5	1.9	1.2	1.5	1.0	1.0	0.9	0.9	1.2	0.8	0.8	1.0	0.7
Gender:															
Male	3.1	3.4	2.9	3.2	2.2	2.9	1.7	1.8	1.6	1.5	2.5	1.5	1.4	1.7	1.2
Female	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.2	0.3	0.1	0.2	0.2	0.3	0.2
College Plans:															
None or under 4 years	4.1	5.6	4.4	5.4	3.5	5.1	3.6	6.1	3.8	2.7	4.5	4.5	4.0	3.8	2.4
Complete 4 years	1.2	1.2	1.1	1.4	0.9	1.0	0.6	0.5	0.5	0.7	0.9	0.5	0.5	0.7	0.5
Region:															
Northeast	1.2	0.9	0.6	0.8	0.6	0.8	0.6	0.4	0.1	0.4	0.7	0.6	0.8	1.0	0.1
Midwest	1.5	1.6	1.6	1.4	1.1	2.0	1.2	1.3	0.9	1.3	1.1	1.1	0.6	0.6	0.3
South	2.4	3.0	2.2	3.3	1.8	2.0	1.5	1.3	1.6	1.3	2.1	1.0	1.3	1.7	1.4
West	0.6	0.8	1.0	0.9	0.8	0.8	0.2	0.7	0.3	0.2	0.1	0.2	0.2	0.3	0.2
Population Density:															
Large MSA	0.5	0.6	0.7	0.7	0.4	0.4	0.5	0.4	0.3	0.2	0.3	0.4	0.7	0.4	*
Other MSA	1.2	1.9	1.5	1.0	0.9	1.2	0.8	0.6	0.8	0.8	0.8	0.6	0.6	0.6	0.5
Non-MSA	3.3	2.8	2.5	4.6	2.6	3.4	1.6	2.6	1.8	2.0	2.9	1.8	1.5	2.6	1.9
Parental Education: ^a															
1.0–2.0 (Low)	2.8	3.5	2.0	3.0	2.2	1.5	3.2	2.6	0.9	1.4	1.8	2.2	1.8	2.8	1.4
2.5–3.0	2.2	2.6	1.9	2.7	1.7	3.1	1.1	1.5	1.6	1.4	2.1	1.0	1.0	1.1	0.5
3.5–4.0	1.4	1.2	1.8	1.9	1.2	1.7	0.9	1.3	0.8	1.3	0.3	1.0	0.8	0.8	0.8
4.5–5.0	0.8	1.3	1.1	1.1	0.9	0.3	0.8	0.5	0.5	0.4	0.5	0.3	0.2	0.7	0.8
5.5–6.0 (High)	1.0	0.9	0.6	0.7	0.8	0.8	0.4	0.5	0.4	0.4	1.7	0.9	1.1	0.8	0.4
Race/Ethnicity (2-year average): ^b															
White	—	2.0	2.0	2.0	1.9	1.7	1.5	1.2	1.1	1.1	1.2	1.0	0.7	0.9	0.9
African American	—	0.3	0.4	0.7	0.6	0.4	0.5	0.4	0.4	0.4	0.5	0.5	0.6	0.5	0.2
Hispanic	—	0.8	0.9	0.9	0.7	1.1	0.9	0.8	1.0	0.8	0.8	0.8	0.9	0.9	0.4

Table continued on next page.

TABLE 148 (cont.)
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 8

	Percentage who used daily in last 30 days																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	0.7	0.8	0.8	0.8	0.9	0.8	0.5	0.5	0.5	0.8	0.6	0.4	0.3	0.5	0.5	0.4	-0.1*
Gender:																	
Male	1.1	1.6	1.2	1.4	1.5	1.5	0.8	0.9	0.9	1.2	1.2	0.5	0.5	0.8	§	0.5	-0.3
Female	0.3	0.1	0.4	0.3	0.3	0.2	0.1	0.1	0.3	0.3	0.1	0.2	0.1	0.2	§	0.4	+0.2
College Plans:																	
None or under 4 years	4.0	3.2	5.0	2.5	4.0	3.6	2.3	1.9	3.1	3.3	2.8	2.3	1.4	1.3	§	1.4	+0.1
Complete 4 years	0.4	0.6	0.4	0.7	0.6	0.6	0.4	0.4	0.3	0.5	0.4	0.2	0.1	0.4	§	0.2	-0.2
Region:																	
Northeast	0.6	0.2	0.9	1.1	0.7	0.7	0.5	0.1	0.1	0.2	0.6	0.8	0.2	0.2	§	0.0	-0.2
Midwest	0.5	0.5	0.4	0.6	0.9	0.6	0.2	0.4	0.3	0.4	0.4	0.2	0.2	0.5	§	0.5	0.0
South	1.1	1.5	1.3	1.3	1.5	1.1	0.7	0.9	0.8	1.1	1.0	0.5	0.4	0.9	§	0.6	-0.2
West	0.4	0.4	0.3	0.1	0.2	0.4	0.5	0.3	0.8	0.8	0.2	0.1	0.5	0.2	§	0.4	+0.2
Population Density:																	
Large MSA	0.3	0.4	0.4	0.5	0.4	0.2	0.2	0.1	0.2	0.4	0.1	0.3	0.1	0.3	§	0.6	+0.4
Other MSA	0.6	0.7	0.7	0.6	1.1	0.8	0.5	0.5	0.2	0.5	0.7	0.3	0.3	0.4	§	0.4	0.0
Non-MSA	1.6	1.7	1.5	1.7	1.1	1.7	1.0	1.2	1.8	1.8	1.3	0.7	0.7	1.2	§	0.2	-1.0 ss
Parental Education: ^a																	
1.0–2.0 (Low)	1.3	1.9	1.5	1.6	1.0	1.6	0.3	0.9	1.5	0.7	0.9	0.3	0.0	0.6	§	1.0	+0.4
2.5–3.0	0.9	1.0	1.3	0.9	1.9	0.9	0.6	0.8	1.3	1.3	0.6	0.8	0.3	0.7	§	0.1	-0.6 s
3.5–4.0	0.7	0.8	0.8	1.3	0.8	0.9	1.0	0.6	0.2	0.3	0.6	0.5	0.6	0.5	§	0.9	+0.3
4.5–5.0	0.6	0.6	0.3	0.4	0.4	0.7	0.3	0.3	0.0	0.7	0.2	0.1	0.0	0.4	§	0.3	-0.1
5.5–6.0 (High)	0.5	0.5	0.6	0.2	0.2	0.1	0.2	0.2	0.6	0.4	0.8	0.3	0.2	0.3	§	0.5	+0.1
Race/Ethnicity (2-year average): ^b																	
White	0.7	0.9	0.9	0.9	1.0	1.0	0.9	0.7	0.8	0.9	0.9	0.7	0.5	0.4	§	0.6	+0.2
African American	0.5	0.5	0.7	1.0	0.7	0.5	0.3	0.3	0.2	0.2	0.3	0.4	0.4	0.6	§	0.5	-0.1
Hispanic	0.5	0.5	0.4	0.3	0.4	0.4	0.3	0.1	0.1	0.3	0.3	0.2	0.3	0.4	§	0.0	-0.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. '*' indicates less than 0.05% but greater than 0%. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.
 1991–1996 and on two of four forms beginning in 1997; N is one half of N indicated in Table S1.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 149
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 10

	Percentage who used daily in last 30 days														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	3.3	3.0	3.3	3.0	2.7	2.2	2.2	2.2	1.5	1.9	2.2	1.7	1.8	1.6	1.9
Gender:															
Male	6.3	6.3	6.4	5.9	5.2	4.2	4.0	4.3	3.2	3.9	4.5	3.0	3.7	3.0	3.4
Female	0.2	0.1	0.3	0.2	0.2	0.2	0.3	0.3	0.1	0.2	0.3	0.2	0.1	0.2	0.3
College Plans:															
None or under 4 years	7.6	8.5	8.8	6.5	7.8	5.4	6.3	6.4	3.6	5.9	6.6	4.6	4.8	5.0	5.4
Complete 4 years	2.3	1.9	2.2	2.2	1.9	1.6	1.5	1.5	1.2	1.2	1.5	1.0	1.3	1.1	1.3
Region:															
Northeast	1.8	1.0	1.7	3.0	2.0	1.3	1.5	1.2	1.2	0.9	1.1	0.8	1.6	2.1	1.3
Midwest	3.1	2.9	2.5	2.4	2.5	2.1	1.4	2.1	2.0	2.1	2.1	1.3	1.3	0.8	1.7
South	4.7	4.5	5.2	3.3	4.1	3.3	3.5	3.8	2.0	2.8	3.6	2.6	3.0	2.6	2.8
West	2.7	2.9	3.1	3.6	1.1	1.0	1.7	0.8	0.7	1.0	0.9	1.1	0.8	0.6	1.2
Population Density:															
Large MSA	1.5	1.6	1.1	1.0	1.5	0.6	0.8	1.0	0.8	1.6	0.8	1.3	1.2	0.8	0.8
Other MSA	3.1	2.8	3.2	3.5	2.3	2.3	1.7	1.5	1.0	1.0	1.7	1.4	1.5	1.3	1.4
Non-MSA	5.0	4.9	5.3	4.2	4.9	3.6	4.6	5.0	3.5	3.8	5.0	2.6	3.5	3.4	4.2
Parental Education: ^a															
1.0–2.0 (Low)	2.5	3.9	4.1	3.2	3.6	1.7	3.8	2.6	1.7	2.4	3.8	1.7	1.7	1.5	2.0
2.5–3.0	4.8	5.0	4.3	3.8	3.4	3.4	2.0	2.8	1.7	2.3	3.2	1.2	2.1	1.8	2.6
3.5–4.0	3.3	2.8	3.1	3.0	2.8	1.4	2.4	2.7	1.6	1.6	1.5	2.0	1.9	2.4	2.0
4.5–5.0	2.5	1.7	2.5	2.7	2.9	2.3	1.7	1.8	1.6	1.9	1.9	1.5	1.7	0.7	1.7
5.5–6.0 (High)	2.5	1.6	2.7	1.7	1.0	1.4	1.9	0.7	0.8	0.8	1.9	1.3	1.4	0.9	1.2
Race/Ethnicity (2-year average): ^b															
White	—	3.8	3.9	3.8	3.3	2.9	2.5	2.7	2.4	2.1	2.4	2.3	2.1	2.1	2.1
African American	—	0.5	0.4	0.6	0.5	0.4	0.4	0.4	0.3	0.4	0.9	0.8	0.4	0.6	0.8
Hispanic	—	1.1	1.0	0.8	1.2	1.2	1.3	1.3	0.8	0.8	1.0	0.5	0.4	0.9	1.0

Table continued on next page.

TABLE 149 (cont.)
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 10

	Percentage who used daily in last 30 days																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	1.7	1.6	1.4	1.9	2.5	1.7	2.0	1.9	1.8	1.6	1.0	0.6	1.0	0.9	0.7	0.4	-0.6* ss
Gender:																	
Male	3.3	2.9	2.6	3.7	4.7	3.3	3.9	3.5	3.4	3.1	1.7	1.2	1.8	1.6	§	0.5	-1.0 ss
Female	*	0.3	0.3	0.2	0.3	0.2	0.3	0.3	0.4	0.2	0.2	0.1	0.2	0.4	§	0.1	-0.2
College Plans:																	
None or under 4 years	6.8	4.4	6.2	7.7	8.2	5.2	6.2	7.6	7.7	5.8	4.3	2.4	3.6	2.4	§	1.6	-0.8
Complete 4 years	1.0	1.2	0.8	1.3	1.8	1.3	1.6	1.2	1.2	1.2	0.7	0.4	0.7	0.7	§	0.1	-0.6 sss
Region:																	
Northeast	0.8	1.4	0.5	1.5	2.3	1.2	1.2	1.7	0.9	1.4	0.5	0.8	1.1	0.8	§	0.1	-0.8 s
Midwest	2.2	1.6	2.4	2.8	2.3	1.7	2.5	1.7	1.9	1.8	0.7	0.4	0.9	0.8	§	0.4	-0.3
South	2.3	2.6	2.1	2.3	3.7	2.4	2.9	2.7	3.1	2.3	1.7	0.8	1.5	1.3	§	0.5	-0.8
West	0.9	0.4	0.2	0.6	0.8	0.9	0.9	1.0	0.5	0.7	0.8	0.4	0.4	0.6	§	0.2	-0.3
Population Density:																	
Large MSA	1.1	1.0	0.7	1.0	1.4	1.2	0.7	1.1	0.9	1.0	0.9	0.6	0.8	0.5	§	0.2	-0.3
Other MSA	1.4	1.5	1.1	1.9	2.6	1.0	1.7	1.5	1.3	1.1	0.7	0.3	0.9	1.0	§	0.3	-0.7 s
Non-MSA	3.2	2.6	3.2	3.1	3.8	3.5	4.5	3.7	4.7	4.1	2.0	1.5	1.7	1.5	§	0.8	-0.8
Parental Education: ^a																	
1.0–2.0 (Low)	3.6	2.0	1.0	2.9	2.5	2.0	1.5	1.1	2.2	1.5	0.5	0.1	0.5	1.3	§	0.3	-1.0
2.5–3.0	2.9	2.1	1.7	2.3	3.4	2.8	3.4	3.0	2.9	3.2	1.7	0.5	1.3	1.2	§	0.8	-0.4
3.5–4.0	1.3	1.6	1.9	2.2	3.0	1.4	2.2	2.0	2.3	0.7	1.3	0.8	1.0	1.3	§	0.2	-1.0 s
4.5–5.0	1.1	1.4	1.3	1.3	1.7	1.4	1.7	1.2	1.2	2.1	0.6	0.6	1.0	0.8	§	0.2	-0.6 s
5.5–6.0 (High)	1.3	1.1	0.1	1.7	2.0	1.4	1.3	1.1	0.9	1.1	0.7	0.7	1.1	0.5	§	0.2	-0.3
Race/Ethnicity (2-year average): ^b																	
White	2.2	2.1	2.1	2.4	3.1	2.8	2.4	2.7	2.6	2.4	1.8	1.2	1.3	1.2	§	0.9	-0.3
African American	0.8	0.6	0.4	0.4	0.6	0.9	0.8	0.7	0.8	1.0	0.9	0.4	0.3	1.0	§	0.3	-0.7
Hispanic	0.7	0.3	0.2	0.4	0.5	0.7	0.9	0.5	0.4	0.3	0.3	0.3	0.2	0.4	§	0.1	-0.3 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. '***' indicates less than 0.05% but greater than 0%. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of two forms in 1991–1996 and on two of four forms beginning in 1997; N is one half of N indicated in Table S2.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 150
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990 ^a
Approximate weighted N =	—	—	—	—	—	—	—	—	—	—	—	15,200	16,300	16,300	16,700	—
Total	—	—	—	—	—	—	—	—	—	—	—	4.7	5.1	4.3	3.3	—
Gender:																
Male	—	—	—	—	—	—	—	—	—	—	—	10.0	10.7	8.6	6.8	—
Female	—	—	—	—	—	—	—	—	—	—	—	0.1	0.1	0.5	0.0	—
College Plans:																
None or under 4 years	—	—	—	—	—	—	—	—	—	—	—	7.1	7.8	5.8	4.2	—
Complete 4 years	—	—	—	—	—	—	—	—	—	—	—	3.3	3.7	3.5	2.7	—
Region:																
Northeast	—	—	—	—	—	—	—	—	—	—	—	4.6	2.1	2.3	1.3	—
Midwest	—	—	—	—	—	—	—	—	—	—	—	4.5	4.5	3.5	2.2	—
South	—	—	—	—	—	—	—	—	—	—	—	6.1	7.4	6.3	4.2	—
West	—	—	—	—	—	—	—	—	—	—	—	2.9	5.5	4.0	4.9	—
Population Density:																
Large MSA	—	—	—	—	—	—	—	—	—	—	—	3.4	3.3	3.0	3.0	—
Other MSA	—	—	—	—	—	—	—	—	—	—	—	3.3	4.3	2.5	2.8	—
Non-MSA	—	—	—	—	—	—	—	—	—	—	—	7.8	8.5	8.9	4.6	—
Parental Education: ^b																
1.0–2.0 (Low)	—	—	—	—	—	—	—	—	—	—	—	1.9	5.6	5.3	1.8	—
2.5–3.0	—	—	—	—	—	—	—	—	—	—	—	7.6	6.9	3.2	3.9	—
3.5–4.0	—	—	—	—	—	—	—	—	—	—	—	3.5	4.7	5.4	3.1	—
4.5–5.0	—	—	—	—	—	—	—	—	—	—	—	3.9	5.0	4.7	4.6	—
5.5–6.0 (High)	—	—	—	—	—	—	—	—	—	—	—	3.3	2.1	3.5	1.2	—
Race/Ethnicity (2-year average): ^c																
White	—	—	—	—	—	—	—	—	—	—	—	—	5.8	5.4	4.5	—
African American	—	—	—	—	—	—	—	—	—	—	—	—	0.6	1.0	0.5	—
Hispanic	—	—	—	—	—	—	—	—	—	—	—	—	0.8	2.1	2.1	—

Table continued on next page.

TABLE 150 (cont.)
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															
	1991 ^a	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Approximate weighted N =	—	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	—	4.3	3.3	3.9	3.6	3.3	4.4	3.2	2.9	3.2	2.8	2.0	2.2	2.8	2.5	2.2
Gender:																
Male	—	7.8	6.4	7.2	7.2	7.1	8.6	6.0	5.7	6.5	5.6	4.3	4.6	5.6	4.7	4.3
Female	—	0.5	0.4	0.3	0.1	0.1	0.2	0.0	0.0	0.4	0.3	0.0	0.0	0.2	0.2	0.1
College Plans:																
None or under 4 years	—	7.4	4.3	6.6	6.5	6.8	9.1	6.5	3.4	7.9	4.7	2.4	5.3	5.0	7.1	6.1
Complete 4 years	—	3.3	3.1	2.8	2.7	2.6	2.7	2.3	2.6	2.0	2.1	1.4	1.4	2.0	1.5	1.2
Region:																
Northeast	—	1.8	1.9	4.5	2.2	3.2	3.5	0.5	1.0	1.3	1.5	1.6	2.7	1.9	2.1	1.9
Midwest	—	4.0	4.4	4.7	4.9	4.1	7.0	4.0	3.4	5.0	2.8	2.2	1.1	3.0	3.8	2.1
South	—	5.4	4.0	3.5	4.2	3.1	3.6	4.6	4.0	4.0	3.8	2.4	2.6	3.7	2.7	3.0
West	—	5.1	1.7	3.2	1.6	2.9	3.0	1.8	1.9	1.6	2.6	1.3	2.2	1.7	1.0	1.0
Population Density:																
Large MSA	—	2.0	1.7	2.0	2.1	2.6	3.3	0.9	0.5	0.9	1.4	0.4	0.5	0.8	1.0	1.4
Other MSA	—	4.2	3.0	3.6	3.2	1.9	3.3	2.4	3.1	3.7	2.6	2.0	2.2	2.2	2.4	1.6
Non-MSA	—	6.5	5.2	6.7	5.8	6.7	7.7	7.6	4.9	5.3	5.0	3.8	4.2	6.2	4.8	4.4
Parental Education: ^b																
1.0–2.0 (Low)	—	6.7	3.9	6.6	2.7	2.2	1.3	2.2	2.1	2.9	4.0	2.2	4.9	3.0	2.2	1.3
2.5–3.0	—	4.8	3.5	3.8	4.7	3.6	5.8	5.2	3.7	4.2	3.2	1.7	1.4	4.2	3.5	3.3
3.5–4.0	—	5.2	3.3	3.3	2.9	3.6	3.7	2.6	3.5	3.9	2.8	2.1	2.0	2.0	3.7	1.7
4.5–5.0	—	2.4	3.7	3.9	3.5	4.6	3.9	3.0	1.9	2.1	2.7	2.4	2.8	2.6	1.8	2.6
5.5–6.0 (High)	—	2.6	1.8	2.7	2.7	1.1	5.0	2.5	2.1	2.5	2.5	0.2	1.8	1.9	0.4	1.3
Race/Ethnicity (2-year average): ^c																
White	—	—	4.8	4.7	4.6	4.1	5.0	5.2	4.3	4.3	4.4	3.5	2.9	3.2	3.5	3.4
African American	—	—	0.3	0.7	0.6	0.3	0.2	0.0	0.0	0.1	0.2	0.1	0.0	0.0	0.0	0.2
Hispanic	—	—	1.6	0.7	1.2	2.2	1.9	0.8	0.4	0.9	0.6	0.3	0.7	1.1	0.7	0.4

Table continued on next page.

TABLE 150 (cont.)
SMOKELESS TOBACCO: Trends in 30-Day Prevalence of Daily Use by Subgroups in Grade 12

	Percentage who used daily in last 30 days															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	2.8	2.7	2.9	3.1	3.1	3.2	3.0	3.4	2.9	2.7	2.0	1.6	1.1	§	0.7	-0.4*
Gender:																
Male	5.6	5.6	5.8	5.9	6.0	5.7	5.7	6.5	5.8	5.1	4.0	3.2	1.9	§	1.2	-0.7
Female	0.2	0.0	0.4	0.2	0.0	0.3	0.0	0.1	0.0	0.3	0.2	0.3	0.2	§	0.3	+0.1
College Plans:																
None or under 4 years	5.3	6.6	6.4	7.5	7.8	8.4	7.8	8.8	8.6	5.1	6.7	3.6	3.0	§	1.2	-1.8
Complete 4 years	2.1	1.9	2.3	2.1	2.2	2.0	1.8	1.9	1.7	1.9	1.0	1.1	0.5	§	0.4	+0.0
Region:																
Northeast	1.9	1.8	2.1	2.4	3.8	3.2	1.6	3.9	2.1	2.1	1.4	1.1	0.4	§	0.6	+0.2
Midwest	3.9	3.1	4.4	5.3	4.1	3.7	4.9	5.6	3.8	1.9	2.0	2.8	1.3	§	1.0	-0.3
South	4.2	2.7	3.7	3.7	2.9	4.3	2.9	2.5	3.3	3.6	3.0	1.6	1.5	§	0.5	-1.0
West	0.1	3.0	0.7	0.6	1.9	1.3	2.2	2.3	2.0	2.2	0.7	0.6	0.6	§	0.7	+0.1
Population Density:																
Large MSA	1.6	0.5	0.9	0.9	1.6	2.1	1.4	0.9	0.4	1.2	0.3	0.5	0.0	§	0.8	+0.8 s
Other MSA	2.4	2.1	2.4	2.0	2.9	1.9	2.3	3.5	2.9	3.1	2.3	1.6	1.0	§	0.4	-0.5
Non-MSA	5.8	6.8	6.7	9.2	5.9	8.1	7.0	6.4	7.1	4.0	4.1	3.6	3.2	§	0.9	-2.3
Parental Education: ^b																
1.0–2.0 (Low)	2.5	0.7	1.2	2.2	0.8	3.1	2.0	0.6	1.2	1.8	0.0	.	1.0	§	0.4	-0.6
2.5–3.0	3.8	3.4	3.5	3.9	4.7	5.9	3.0	3.8	4.1	3.5	3.1	2.3	1.3	§	1.3	0.0
3.5–4.0	2.7	2.9	2.5	2.9	2.9	3.1	3.0	3.8	3.5	2.1	2.6	2.8	0.7	§	0.2	-0.4
4.5–5.0	3.1	2.7	3.5	3.2	3.5	2.3	3.7	4.4	2.7	3.2	2.3	1.6	0.9	§	0.6	-0.3
5.5–6.0 (High)	1.8	2.2	3.1	2.8	0.8	1.9	1.5	2.4	1.8	3.3	0.4	0.9	1.3	§	0.8	-0.5
Race/Ethnicity (2-year average): ^c																
White	3.4	3.8	4.5	4.7	4.6	4.6	4.3	4.6	4.7	4.4	4.1	3.4	2.0	§	3.0	+1.0
African American	0.8	0.9	0.6	1.5	1.4	0.3	0.6	1.3	0.6	0.7	1.1	0.8	0.3	§	0.0	-0.3
Hispanic	0.7	0.6	0.2	0.4	0.5	0.5	0.4	0.5	0.8	0.5	0.0	0.2	0.2	§	0.2	-0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aPrevalence of smokeless tobacco use was not asked of 12th graders in 1990 and 1991. Prior to 1990 the prevalence-of-use question on smokeless tobacco was located near the end of one 12th-grade questionnaire form, whereas after 1991 the question was placed earlier and in a different form. This shift could explain the discontinuities between the corresponding data.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 151
DISSOLVABLE TOBACCO PRODUCTS: Trends in Annual
Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months											2019– 2021 change
	1991– 2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	1.0	1.1	1.1	0.9	0.7	0.6	0.6	1.1	0.6	0.8	-0.3*
Gender:												
Male	—	0.8	1.5	1.3	1.1	0.7	0.8	0.9	1.4	§	0.6	-0.8 s
Female	—	1.0	0.8	1.0	0.8	0.6	0.5	0.3	0.8	§	1.1	+0.3
College Plans:												
None or under 4 years	—	4.4	4.5	6.1	3.3	4.5	3.5	2.1	2.6	§	2.9	+0.4
Complete 4 years	—	0.8	0.9	0.7	0.7	0.3	0.3	0.4	0.8	§	0.5	-0.4
Region:												
Northeast	—	0.3	0.5	1.4	0.5	0.2	0.3	0.6	1.1	§	0.1	-1.0 s
Midwest	—	0.8	0.4	0.5	1.0	0.4	0.2	0.1	0.8	§	1.0	+0.2
South	—	1.4	1.7	1.2	1.2	1.1	1.0	1.0	1.4	§	1.3	-0.1
West	—	1.1	1.3	1.4	0.7	0.5	0.8	0.3	0.7	§	0.3	-0.4
Population Density:												
Large MSA	—	1.2	1.0	1.2	0.8	0.8	0.5	0.8	1.3	§	0.5	-0.7
Other MSA	—	0.7	1.1	0.8	1.0	0.5	0.8	0.4	1.0	§	1.0	0.0
Non-MSA	—	1.3	1.3	1.7	0.9	1.0	0.4	0.4	0.8	§	0.7	-0.1
Parental Education: ^a												
1.0–2.0 (Low)	—	1.2	2.6	2.4	1.3	2.1	1.1	0.6	1.1	§	0.8	-0.3
2.5–3.0	—	1.3	0.7	1.9	1.1	0.7	0.3	0.3	1.7	§	1.8	+0.1
3.5–4.0	—	1.4	1.7	0.8	1.4	0.8	0.7	0.2	1.0	§	1.7	+0.7
4.5–5.0	—	0.7	0.7	0.8	0.6	0.2	0.4	0.5	0.8	§	0.2	-0.6
5.5–6.0 (High)	—	0.6	0.8	0.8	0.2	0.2	0.8	0.8	0.9	§	0.5	-0.4
Race/Ethnicity (2-year data): ^b												
White	—	—	0.8	1.0	0.9	0.4	0.3	0.3	0.5	§	0.8	+0.4
African American	—	—	1.2	1.0	1.0	1.1	0.9	1.4	2.0	§	0.6	-1.4
Hispanic	—	—	1.7	1.6	1.2	0.9	0.9	0.8	0.9	§	0.0	-0.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 152
DISSOLVABLE TOBACCO PRODUCTS: Trends in Annual
Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months											
	1991–											2019–
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2021
												change
<i>Approximate weighted N =</i>	—	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	1.6	1.2	1.3	1.1	0.9	0.6	1.1	0.8	1.3	0.3	-0.4*
Gender:												
Male	—	2.3	2.0	1.9	1.4	1.3	0.9	1.6	1.5	§	0.2	-1.3 s
Female	—	0.9	0.6	0.7	0.6	0.5	0.3	0.5	0.0	§	0.4	+0.3 s
College Plans:												
None or under 4 years	—	5.6	2.8	5.9	4.1	3.0	1.3	1.9	1.4	§	0.7	-0.7
Complete 4 years	—	1.2	1.0	0.8	0.7	0.7	0.5	0.9	0.7	§	0.3	-0.4
Region:												
Northeast	—	0.5	1.0	1.1	0.6	1.1	0.1	1.2	0.3	§	0.3	0.0
Midwest	—	1.7	1.2	0.9	1.0	1.0	0.9	1.2	0.5	§	0.2	-0.3
South	—	1.7	1.6	1.3	0.9	0.9	0.5	0.8	0.4	§	0.5	+0.1
West	—	2.1	0.9	1.9	1.6	0.7	1.0	1.1	1.9	§	0.2	-1.7
Population Density:												
Large MSA	—	1.4	0.8	1.1	0.7	0.7	0.6	1.2	0.2	§	0.1	-0.1
Other MSA	—	1.5	1.0	1.0	1.0	0.9	0.6	0.9	1.0	§	0.4	-0.6
Non-MSA	—	1.9	2.2	2.6	1.7	1.2	0.7	1.2	1.1	§	0.5	-0.6
Parental Education: ^a												
1.0–2.0 (Low)	—	2.2	1.4	0.5	2.8	0.9	0.9	1.5	0.2	§	0.2	0.0
2.5–3.0	—	2.2	2.0	1.2	1.6	1.2	0.1	1.9	0.3	§	0.7	+0.4
3.5–4.0	—	2.3	1.0	2.2	0.7	0.9	0.7	0.7	0.2	§	0.6	+0.3
4.5–5.0	—	0.6	1.1	0.9	0.6	0.6	0.7	0.8	1.9	§	0.0	-1.8
5.5–6.0 (High)	—	1.1	0.9	0.7	0.8	0.7	0.3	0.8	0.7	§	0.1	-0.5
Race/Ethnicity (2-year data): ^b												
White	—	—	1.4	1.3	1.1	0.7	0.6	0.9	1.0	§	0.4	-0.6
African American	—	—	0.8	1.2	1.3	1.1	0.7	0.5	0.4	§	2.5	+2.0
Hispanic	—	—	1.4	1.0	1.0	1.4	1.4	1.2	0.9	§	0.6	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 153
DISSOLVABLE TOBACCO PRODUCTS: Trends in Annual
Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months												
	2009–											2019–	
	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2021
													change
<i>Approximate weighted N =</i>	—	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	1.5	1.6	1.9	1.1	1.4	1.1	1.4	1.3	1.1	§	1.1	-0.1*
Gender:													
Male	—	1.8	2.4	3.3	1.1	1.8	1.2	2.3	1.0	1.8	§	2.3	+0.5
Female	—	1.0	0.4	0.5	0.7	0.7	0.6	0.2	0.7	0.4	§	0.0	-0.4
College Plans:													
None or under 4 years	—	3.0	1.7	3.9	4.6	2.3	4.1	1.0	3.4	2.6	§	2.3	-0.3
Complete 4 years	—	1.1	1.6	1.3	0.3	1.0	0.5	1.2	0.4	0.6	§	0.7	0.0
Region:													
Northeast	—	1.3	1.4	1.3	0.3	1.1	0.7	0.5	0.6	0.4	§	0.0	-0.4
Midwest	—	0.5	0.8	2.0	0.8	0.5	0.2	1.3	2.3	0.4	§	1.6	+1.2
South	—	2.4	2.2	1.5	1.5	2.5	2.0	1.8	1.5	1.4	§	1.5	+0.1
West	—	1.5	2.0	2.7	1.3	0.6	0.6	1.5	0.5	2.3	§	0.5	-1.7
Population Density:													
Large MSA	—	1.7	1.7	1.8	1.9	0.7	0.9	1.0	2.1	1.0	§	0.8	-0.2
Other MSA	—	1.4	1.4	1.6	0.8	1.8	1.3	1.5	0.9	1.1	§	1.0	-0.1
Non-MSA	—	1.5	2.1	2.7	0.7	1.4	1.0	1.9	0.8	1.7	§	1.6	0.0
Parental Education: ^a													
1.0–2.0 (Low)	—	0.6	1.1	3.8	2.5	1.9	1.6	4.0	1.9	0.8	§	1.2	+0.4
2.5–3.0	—	2.7	2.5	1.0	1.9	0.6	1.4	1.5	0.8	1.0	§	0.0	-1.0
3.5–4.0	—	1.1	1.6	2.3	1.0	1.0	2.2	0.7	0.8	1.5	§	1.6	+0.1
4.5–5.0	—	0.5	1.3	1.5	0.1	2.1	0.3	0.4	0.3	0.5	§	1.2	+0.7
5.5–6.0 (High)	—	1.9	0.7	1.7	0.1	0.4	0.0	1.3	1.4	0.3	§	0.6	+0.3
Race/Ethnicity (2-year data): ^b													
White	—	—	1.6	1.8	1.5	1.2	0.9	0.5	0.8	0.8	§	0.6	-0.2
African American	—	—	1.5	1.8	1.9	2.4	2.7	2.5	1.7	2.1	§	2.4	+0.4
Hispanic	—	—	1.0	1.4	1.1	0.5	1.0	1.4	1.1	1.0	§	1.0	0.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 154
SNUS: Trends in Annual Prevalence of Use
by Subgroups in Grade 8

	Percentage who used in last 12 months											
	1991–											2019–
	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	<u>change</u>
<i>Approximate weighted N =</i>	—	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	—	2.4	2.0	2.2	1.9	2.2	1.1	1.3	1.5	1.6	1.2	-0.2*
Gender:												
Male	—	3.3	3.3	3.1	3.0	3.3	1.7	2.1	1.9	§	1.6	-0.3
Female	—	1.5	0.8	1.5	1.0	1.0	0.5	0.7	1.0	§	0.9	-0.2
College Plans:												
None or under 4 years	—	9.0	7.4	8.9	6.0	9.5	4.7	3.7	4.5	§	3.4	-1.1
Complete 4 years	—	2.0	1.6	1.6	1.5	1.5	0.8	1.0	1.1	§	0.9	-0.2
Region:												
Northeast	—	1.8	1.1	1.4	1.2	1.1	0.5	0.7	0.6	§	0.1	-0.5
Midwest	—	2.0	1.2	2.2	2.4	1.3	0.8	0.7	1.5	§	1.0	-0.5
South	—	3.4	3.3	2.4	2.3	3.5	1.6	2.1	2.0	§	2.2	+0.2
West	—	1.6	1.4	2.2	1.0	1.4	1.0	1.1	1.3	§	0.9	-0.5
Population Density:												
Large MSA	—	1.2	1.6	1.1	0.8	1.0	0.5	0.9	1.3	§	0.5	-0.8
Other MSA	—	2.4	1.8	1.7	1.7	2.1	1.0	0.9	1.1	§	1.5	+0.4
Non-MSA	—	4.5	3.0	4.6	3.9	4.0	2.3	2.9	2.8	§	1.9	-0.8
Parental Education: ^a												
1.0–2.0 (Low)	—	2.2	3.0	3.1	1.1	3.2	1.2	0.7	1.8	§	0.2	-1.6 s
2.5–3.0	—	3.9	2.0	3.5	2.7	2.6	1.5	0.8	2.7	§	0.9	-1.7
3.5–4.0	—	3.9	3.0	2.2	1.7	2.4	1.5	1.5	2.1	§	3.6	+1.5
4.5–5.0	—	1.5	1.4	1.4	2.1	1.3	0.7	2.2	1.4	§	0.5	-1.0
5.5–6.0 (High)	—	1.1	1.3	1.2	0.9	0.9	1.0	1.1	0.3	§	1.0	+0.7
Race/Ethnicity (2-year data): ^b												
White	—	—	2.7	2.6	2.7	2.5	1.9	1.4	1.6	§	1.8	+0.2
African American	—	—	0.9	0.7	0.9	1.4	1.4	1.4	1.7	§	0.6	-1.2
Hispanic	—	—	1.8	1.5	1.0	1.3	1.3	0.9	0.9	§	0.9	0.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in S1.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 155
SNUS: Trends in Annual Prevalence of Use
by Subgroups in Grade 10

	Percentage who used in last 12 months											
	1991–											2019–
	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	<u>change</u>
Approximate weighted N =	—	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	—	6.9	5.2	4.5	4.0	3.0	2.6	3.1	2.3	2.2	1.0	-1.3* s
Gender:												
Male	—	12.1	9.3	7.5	6.5	5.1	4.5	5.6	4.4	§	1.4	-3.0 ss
Female	—	2.1	1.3	1.7	1.7	1.1	0.8	0.7	0.5	§	0.5	+0.1
College Plans:												
None or under 4 years	—	17.5	16.0	15.1	8.2	9.1	7.1	7.8	4.0	§	2.3	-1.7
Complete 4 years	—	5.8	4.0	3.4	3.6	2.3	2.1	2.4	2.1	§	0.7	-1.3 s
Region:												
Northeast	—	6.2	5.5	3.3	4.0	2.6	1.8	2.4	1.1	§	0.3	-0.9
Midwest	—	7.8	6.7	5.7	4.8	2.9	3.1	3.4	2.2	§	1.1	-1.1
South	—	8.4	5.0	5.2	4.7	3.6	3.6	4.1	2.5	§	1.7	-0.8
West	—	4.2	3.8	3.5	2.1	2.6	1.4	1.7	2.9	§	0.5	-2.4
Population Density:												
Large MSA	—	4.1	3.1	3.4	2.9	2.0	2.8	2.2	0.7	§	0.3	-0.4
Other MSA	—	7.4	4.8	3.9	3.7	2.5	1.5	2.8	2.4	§	0.8	-1.6 s
Non-MSA	—	9.6	8.9	8.1	6.7	6.2	5.2	5.1	4.7	§	2.9	-1.9
Parental Education: ^a												
1.0–2.0 (Low)	—	4.9	4.8	3.4	3.8	2.1	2.1	2.8	2.0	§	0.2	-1.8
2.5–3.0	—	8.5	6.9	4.3	5.3	4.0	2.6	3.3	1.6	§	1.7	+0.2
3.5–4.0	—	8.3	6.2	5.2	3.8	3.7	3.4	4.3	2.0	§	1.2	-0.9
4.5–5.0	—	6.3	3.4	4.7	3.9	2.9	2.1	3.0	4.4	§	0.9	-3.4 ss
5.5–6.0 (High)	—	5.4	4.7	4.4	3.7	1.8	2.8	2.2	0.9	§	0.6	-0.3
Race/Ethnicity (2-year data): ^b												
White	—	—	8.1	7.1	6.0	4.9	4.1	4.4	4.3	§	1.6	-2.8 sss
African American	—	—	1.6	1.4	2.2	2.1	1.1	0.6	1.0	§	2.4	+1.4
Hispanic	—	—	2.6	1.2	1.1	1.6	1.5	1.3	1.0	§	0.6	-0.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in S2.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 156
SNUS: Trends in Annual Prevalence of Use
by Subgroups in Grade 12

	Percentage who used in last 12 months												2019– 2021 change
	1975– 2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	7.9	7.9	7.7	5.8	5.8	5.8	4.2	4.7	2.7	§	2.6	-0.1*
Gender:													
Male	—	13.9	14.1	14.6	9.9	10.5	9.9	7.2	7.6	4.5	§	4.6	+0.1
Female	—	1.9	1.2	1.4	1.5	1.3	1.4	1.2	1.1	0.7	§	0.9	+0.2
College Plans:													
None or under 4 years	—	15.1	12.2	13.6	16.0	12.8	14.0	7.0	8.0	6.2	§	3.1	-3.1
Complete 4 years	—	6.2	6.6	6.1	3.5	4.4	4.0	3.4	3.5	1.7	§	2.4	+0.8
Region:													
Northeast	—	7.8	8.0	7.3	6.0	7.8	5.1	4.3	2.3	2.1	§	3.6	+1.5
Midwest	—	10.6	8.3	8.3	7.1	5.6	7.3	3.4	8.1	3.0	§	3.6	+0.6
South	—	6.4	8.8	8.9	5.0	6.8	6.2	5.7	5.0	2.4	§	2.5	+0.1
West	—	7.1	6.2	5.5	5.9	2.9	4.1	2.6	2.6	3.4	§	0.8	-2.5
Population Density:													
Large MSA	—	6.9	6.1	6.7	3.7	3.6	3.1	3.1	3.6	1.5	§	3.0	+1.5
Other MSA	—	6.5	7.8	7.9	5.2	6.8	6.4	3.9	4.5	2.6	§	2.2	-0.4
Non-MSA	—	12.2	10.8	8.6	9.6	6.9	8.3	6.4	7.5	4.8	§	2.5	-2.3
Parental Education: ^a													
1.0–2.0 (Low)	—	2.9	4.8	9.2	4.5	4.4	4.0	3.7	2.9	2.8	§	0.7	-2.1
2.5–3.0	—	8.8	8.9	5.7	7.5	4.3	3.4	4.5	4.2	2.6	§	1.1	-1.5
3.5–4.0	—	9.4	6.8	9.7	5.5	6.0	6.4	4.1	4.8	2.8	§	2.0	-0.8
4.5–5.0	—	8.1	10.0	7.5	6.3	7.8	7.7	3.4	4.9	1.7	§	4.1	+2.4
5.5–6.0 (High)	—	5.4	6.2	6.7	2.5	5.3	5.3	5.3	4.4	3.0	§	3.9	+0.8
Race/Ethnicity (2-year data): ^b													
White	—	—	10.2	9.9	9.1	8.1	7.6	6.8	6.2	4.3	§	5.6	+1.2
African American	—	—	3.4	3.3	2.1	2.2	3.0	2.2	1.1	2.1	§	2.2	+0.1
Hispanic	—	—	4.4	4.6	2.9	2.8	3.2	1.9	1.8	1.9	§	0.0	-1.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 157
STEROIDS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
<i>Approximate weighted N =</i>	<i>17,500</i>	<i>18,600</i>	<i>18,300</i>	<i>17,300</i>	<i>17,500</i>	<i>17,800</i>	<i>18,600</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>
Total	1.0	1.1	0.9	1.2	1.0	0.9	1.0	1.2	1.7	1.7	1.6	1.5	1.4	1.1	1.1
Gender:															
Male	1.8	1.7	1.4	1.8	1.3	1.1	1.3	1.6	2.5	2.2	2.3	1.8	1.8	1.3	1.2
Female	0.3	0.5	0.3	0.6	0.8	0.7	0.7	0.7	0.9	1.0	1.0	1.2	1.1	1.0	0.9
College Plans:															
None or under 4 years	2.2	2.4	2.2	2.5	2.2	1.5	2.4	2.8	4.0	3.6	3.1	3.6	2.7	3.2	2.4
Complete 4 years	0.8	0.9	0.7	1.0	0.9	0.8	0.8	1.0	1.4	1.5	1.5	1.3	1.3	0.9	0.9
Region:															
Northeast	0.7	1.1	0.6	1.0	1.0	1.1	1.0	1.1	1.6	1.7	1.8	1.2	1.1	0.8	0.6
Midwest	1.1	1.2	1.0	1.0	1.1	0.8	1.0	1.2	1.6	1.8	1.7	1.6	1.7	1.2	1.1
South	1.2	1.1	1.0	1.6	1.1	0.9	0.9	1.4	1.9	1.8	1.9	1.9	1.6	1.3	1.3
West	0.7	0.9	0.7	1.0	1.0	0.8	1.1	0.9	1.4	1.3	0.9	1.2	1.2	1.0	1.0
Population Density:															
Large MSA	0.8	1.0	0.8	0.9	0.9	0.8	0.9	1.0	1.3	1.6	1.6	1.1	1.0	1.0	0.8
Other MSA	1.2	1.2	0.9	1.2	1.2	0.9	0.9	1.2	1.9	1.5	1.7	1.6	1.5	1.1	1.1
Non-MSA	1.0	0.9	0.9	1.5	0.9	0.9	1.2	1.4	1.7	2.0	1.6	2.1	1.8	1.3	1.3
Parental Education: ^b															
1.0–2.0 (Low)	1.8	1.2	1.2	1.6	1.3	0.9	1.4	1.7	2.2	2.5	2.0	1.8	1.3	1.5	2.1
2.5–3.0	1.1	1.2	0.8	1.6	1.3	0.7	0.9	1.1	1.9	2.1	1.6	1.7	1.9	1.4	1.0
3.5–4.0	1.0	1.0	1.1	1.3	0.8	0.9	1.2	1.4	1.6	1.3	1.9	1.9	1.6	1.0	1.1
4.5–5.0	0.7	0.9	0.8	0.8	0.8	1.2	0.9	1.1	1.4	1.6	1.5	1.0	1.3	1.2	1.1
5.5–6.0 (High)	1.0	1.3	0.6	0.9	1.5	0.9	1.2	1.1	2.0	1.6	1.7	1.6	1.0	0.7	0.7
Race/Ethnicity (2-year average): ^c															
White	—	1.1	1.0	1.0	1.1	0.9	0.9	1.1	1.5	1.9	1.8	1.7	1.6	1.3	1.0
African American	—	0.7	0.6	0.8	0.9	0.7	0.6	0.7	0.8	0.7	0.8	1.2	1.2	0.9	0.9
Hispanic	—	1.2	1.1	1.1	1.3	1.5	1.4	1.4	1.8	1.8	1.5	1.5	1.7	1.7	1.2

Table continued on next page.

TABLE 157 (cont.)
STEROIDS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 8

	Percentage who used in last 12 months																2019-2021
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700	
Total	0.9	0.8	0.9	0.8	0.5	0.7	0.6	0.6	0.6	0.5	0.5	0.6	0.6	0.8	1.1	0.5	-0.3* s
Gender:																	
Male	1.2	1.1	1.2	1.0	0.7	1.0	0.8	0.7	0.6	0.5	0.5	0.6	0.6	0.7	§	0.5	-0.2
Female	0.6	0.4	0.5	0.5	0.3	0.4	0.3	0.4	0.5	0.5	0.5	0.6	0.7	0.9	§	0.4	-0.5 ss
College Plans:																	
None or under 4 years	2.4	2.7	2.3	2.2	1.4	1.4	1.4	1.6	1.4	1.1	1.1	1.3	1.0	1.5	§	0.6	-0.9 s
Complete 4 years	0.7	0.6	0.8	0.7	0.5	0.7	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.7	§	0.4	-0.2
Region:																	
Northeast	0.6	0.5	0.8	0.8	0.5	0.5	0.5	0.6	0.6	0.5	0.2	0.8	0.5	0.6	§	0.1	-0.4 ss
Midwest	0.8	0.7	0.9	0.7	0.7	0.9	0.5	0.3	0.6	0.6	0.6	0.6	0.6	0.8	§	0.7	-0.1
South	1.1	1.0	1.1	0.7	0.5	0.8	0.8	0.8	0.6	0.6	0.6	0.4	0.7	0.7	§	0.6	-0.1
West	0.7	0.8	0.6	0.9	0.4	0.6	0.5	0.5	0.4	0.4	0.6	0.8	0.7	1.0	§	0.3	-0.7 s
Population Density:																	
Large MSA	0.8	0.7	0.8	0.4	0.4	0.3	0.6	0.6	0.3	0.5	0.5	0.4	0.6	0.6	§	0.3	-0.2
Other MSA	0.8	0.8	0.9	0.9	0.7	0.8	0.6	0.6	0.6	0.4	0.4	0.7	0.7	0.8	§	0.5	-0.2
Non-MSA	1.2	1.0	1.0	0.9	0.4	1.2	0.6	0.7	0.7	0.7	0.9	0.9	0.7	1.1	§	0.5	-0.6 s
Parental Education: ^b																	
1.0–2.0 (Low)	1.0	1.7	1.7	2.1	0.7	1.0	0.6	0.6	0.8	0.8	0.8	0.8	0.6	0.6	§	0.9	+0.3
2.5–3.0	0.9	0.9	1.0	0.7	0.5	0.9	0.6	0.7	0.5	0.7	0.5	0.7	1.0	1.2	§	0.4	-0.8
3.5–4.0	1.0	0.8	1.0	0.9	0.5	0.9	0.4	0.6	0.7	0.3	0.5	0.5	0.7	0.6	§	0.7	+0.1
4.5–5.0	0.7	0.4	0.7	0.5	0.6	0.5	0.5	0.3	0.5	0.6	0.5	0.7	0.5	0.9	§	0.4	-0.5
5.5–6.0 (High)	1.1	0.8	0.8	0.6	0.5	0.8	0.9	0.8	0.6	0.5	0.6	0.6	0.4	0.7	§	0.4	-0.2
Race/Ethnicity (2-year average): ^c																	
White	0.9	0.8	0.8	0.7	0.5	0.5	0.6	0.5	0.5	0.5	0.5	0.5	0.6	0.5	§	0.6	0.0
African American	0.8	0.8	0.7	0.7	0.7	0.5	0.7	0.8	0.7	0.6	0.5	0.6	0.6	0.6	§	0.6	-0.1
Hispanic	0.9	0.7	0.7	0.9	0.8	0.7	0.4	0.5	0.6	0.5	0.6	0.6	0.7	0.8	§	1.1	+0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S1 for the number of subgroup cases. See appendix B for definition of variables in table. In 2006 the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2007 the remaining forms were changed in a like manner. In 2008 the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2009 the remaining forms were changed in a like manner.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 158
STEROIDS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200
Total	1.1	1.1	1.0	1.1	1.2	1.2	1.2	1.2	1.7	2.2	2.1	2.2	1.7	1.5	1.3
Gender															
Male	1.9	1.9	1.7	1.9	2.0	1.7	1.8	1.9	2.8	3.6	3.3	3.2	2.3	2.3	1.8
Female	0.3	0.3	0.3	0.4	0.5	0.6	0.6	0.6	0.7	0.8	1.0	1.2	1.1	0.9	0.7
College Plans															
None or under 4 years	1.7	1.3	1.9	2.1	2.1	2.1	2.4	1.9	2.6	3.6	3.2	3.0	2.7	2.9	1.9
Complete 4 years	0.9	1.0	0.8	0.9	1.1	1.0	1.0	1.1	1.6	1.9	1.9	2.1	1.5	1.4	1.2
Region															
Northeast	1.2	0.9	1.0	1.0	1.1	1.0	0.9	1.4	2.0	1.8	1.9	2.0	1.9	1.4	1.5
Midwest	1.0	1.1	1.2	1.1	1.2	1.4	1.2	1.1	1.8	2.1	1.5	1.8	1.9	1.4	1.2
South	1.0	1.2	1.0	1.3	1.3	1.4	1.4	1.4	1.7	2.5	2.5	2.5	1.5	1.9	1.5
West	1.0	1.2	0.8	1.1	1.3	0.6	1.3	0.9	1.4	2.1	2.3	2.4	1.4	1.4	0.9
Population Density															
Large MSA	1.5	0.9	0.8	0.8	0.8	0.8	1.0	0.8	1.2	2.3	1.9	2.1	1.4	1.2	0.9
Other MSA	1.0	1.0	0.9	1.1	1.4	1.2	1.2	1.3	1.9	2.0	2.0	2.3	1.5	1.5	1.4
Non-MSA	0.8	1.4	1.4	1.5	1.4	1.5	1.5	1.5	2.0	2.5	2.4	2.1	2.6	2.1	1.5
Parental Education ^b															
1.0–2.0 (Low)	0.7	0.9	1.5	1.8	1.2	1.7	1.7	1.3	1.2	2.3	1.6	1.6	1.9	1.5	1.8
2.5–3.0	1.3	1.1	1.0	0.9	1.1	0.7	1.1	1.1	1.7	2.2	2.1	2.0	1.9	1.8	1.2
3.5–4.0	1.0	1.2	1.1	0.8	1.6	1.2	1.4	1.7	2.0	2.0	2.3	2.8	1.9	1.7	1.0
4.5–5.0	0.9	1.0	0.8	1.4	1.1	1.2	1.0	0.9	1.5	2.4	2.0	2.0	1.6	1.5	1.5
5.5–6.0 (High)	1.2	1.4	1.1	1.1	1.0	1.1	1.1	1.1	2.4	2.2	2.1	3.0	1.3	1.2	1.3
Race/Ethnicity (2-year average) ^c															
White	—	1.0	1.0	1.0	1.2	1.3	1.3	1.3	1.5	2.1	2.3	2.4	2.3	1.8	1.6
African American	—	0.7	0.8	0.8	0.7	0.7	0.5	0.5	0.7	1.2	1.6	1.2	0.8	0.7	0.9
Hispanic	—	1.2	1.4	1.3	0.9	1.1	1.2	1.2	1.5	1.8	2.1	2.1	1.8	1.6	1.3

Table continued on next page.

TABLE 158 (cont.)
STEROIDS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 12 months																2019-2021 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000	
Total	1.2	1.1	0.9	0.8	1.0	0.9	0.8	0.8	0.8	0.7	0.7	0.7	0.6	0.8	0.9	0.3	-0.5* sss
Gender																	
Male	1.9	1.7	1.4	1.2	1.3	1.4	1.3	1.3	1.1	1.0	0.9	0.8	0.6	1.1	§	0.3	-0.8 sss
Female	0.5	0.4	0.5	0.4	0.5	0.4	0.4	0.5	0.5	0.4	0.5	0.5	0.6	0.6	§	0.2	-0.3 s
College Plans																	
None or under 4 years	2.3	1.9	2.6	1.6	2.6	2.1	2.3	1.2	2.5	1.5	1.2	1.7	0.9	1.0	§	0.5	-0.5
Complete 4 years	1.1	0.9	0.7	0.7	0.8	0.8	0.7	0.8	0.6	0.7	0.7	0.6	0.6	0.8	§	0.2	-0.6 sss
Region																	
Northeast	0.9	1.1	0.8	1.2	1.1	0.8	0.8	1.0	0.5	1.1	0.5	0.7	0.7	0.8	§	0.6	-0.2
Midwest	1.2	1.2	1.0	0.7	0.7	0.8	0.7	0.8	0.7	0.7	0.9	0.8	0.7	0.6	§	0.1	-0.5 sss
South	1.6	1.3	1.0	0.7	1.2	1.2	1.0	0.8	1.2	0.7	0.9	0.7	0.7	1.0	§	0.4	-0.7 s
West	0.9	0.7	0.8	0.6	0.8	0.7	0.7	0.8	0.6	0.6	0.5	0.6	0.4	0.8	§	0.1	-0.6 s
Population Density																	
Large MSA	1.0	0.8	0.7	0.9	0.9	0.6	0.9	0.9	0.8	0.7	0.4	0.6	0.7	0.5	§	0.1	-0.4 ss
Other MSA	1.3	1.1	1.0	0.7	0.9	1.0	0.8	0.9	0.6	0.7	0.8	0.7	0.6	0.9	§	0.2	-0.7 sss
Non-MSA	1.4	1.3	1.1	0.7	1.1	1.1	0.7	0.8	1.2	1.0	1.1	0.8	0.7	1.2	§	0.6	-0.6
Parental Education ^b																	
1.0–2.0 (Low)	1.4	1.0	1.5	1.0	1.5	0.5	1.1	1.0	0.6	0.4	0.7	1.2	0.4	1.0	§	0.4	-0.7
2.5–3.0	1.5	1.4	1.0	0.9	1.0	0.9	0.9	0.9	0.6	0.6	0.7	0.6	0.5	0.6	§	0.1	-0.5 s
3.5–4.0	1.3	1.0	0.9	0.7	0.9	0.8	0.7	1.2	1.0	0.9	0.9	0.9	1.0	§	0.2	-0.8 ss	
4.5–5.0	0.9	1.1	0.8	0.7	0.7	1.1	0.9	0.5	0.7	0.9	0.5	0.6	0.5	§	0.3	-0.2	
5.5–6.0 (High)	1.4	0.9	1.0	0.9	1.2	1.0	0.7	0.7	0.8	0.5	1.0	0.4	0.6	§	0.2	-1.0 ss	
Race/Ethnicity (2-year average) ^c																	
White	1.3	1.2	1.0	0.8	0.7	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.7	0.8	§	0.5	-0.3
African American	1.1	0.9	0.8	0.7	1.0	1.3	1.0	0.9	1.0	1.0	0.7	0.7	0.6	0.4	§	0.8	+0.4
Hispanic	1.0	0.7	1.1	1.3	0.8	0.6	0.8	0.9	0.5	0.6	0.4	0.4	0.6	0.7	§	0.2	-0.5 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S2 for the number of subgroup cases. See appendix B for definition of variables in table. In 2006 the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2007 the remaining forms were changed in a like manner. In 2008 the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2009 the remaining forms were changed in a like manner.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^bOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 159
STEROIDS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months																	
	1975– 1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<i>Approximate weighted N =</i>	—	16,700	15,200	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700
Total	—	1.9	1.7	1.4	1.1	1.2	1.3	1.5	1.4	1.4	1.7	1.8	1.7	2.4	2.5	2.1	2.5	1.5
Gender																		
Male	—	2.8	2.6	2.4	2.1	2.5	2.1	2.4	2.2	2.5	2.8	3.1	2.5	3.8	3.8	3.2	3.3	2.6
Female	—	0.9	0.3	0.2	0.1	0.1	0.5	0.6	0.4	0.5	0.3	0.6	0.9	1.1	1.3	1.1	1.7	0.4
College Plans																		
None or under 4 years	—	2.3	2.2	2.1	2.1	2.0	1.9	2.0	2.3	2.3	2.1	3.6	2.0	4.6	4.1	2.6	2.8	2.1
Complete 4 years	—	1.6	1.3	1.2	0.8	0.9	1.1	1.2	1.1	1.2	1.4	1.3	1.6	1.8	2.0	2.0	2.4	1.2
Region																		
Northeast	—	2.0	1.3	1.2	0.6	1.5	1.5	1.6	1.3	2.1	0.9	1.3	1.9	4.5	2.1	3.0	3.8	1.6
Midwest	—	1.5	1.3	1.4	1.4	0.8	2.2	1.5	2.1	2.1	2.3	1.6	2.3	1.7	2.8	1.8	2.4	1.6
South	—	2.1	2.2	1.7	0.6	1.6	1.0	1.7	1.3	0.5	1.6	2.6	1.5	2.1	2.5	1.8	2.2	1.6
West	—	1.9	1.7	1.0	2.3	1.1	0.8	1.0	0.3	1.6	1.7	1.2	1.1	2.1	2.4	2.2	2.0	1.0
Population Density																		
Large MSA	—	1.6	1.3	1.1	1.0	0.7	1.1	1.4	1.3	0.8	2.1	1.3	1.7	1.9	2.0	2.1	2.6	1.4
Other MSA	—	2.3	1.5	1.4	1.4	0.9	1.5	1.3	1.3	1.7	1.5	2.0	2.0	2.9	3.0	2.1	2.9	1.4
Non-MSA	—	1.1	2.4	1.6	0.8	2.2	1.3	2.1	1.5	1.6	1.6	1.9	1.2	2.0	2.1	2.3	1.8	1.7
Parental Education ^b																		
1.0–2.0 (Low)	—	0.6	1.1	2.0	2.1	1.1	2.8	1.1	1.5	3.4	3.0	1.5	1.3	2.1	3.6	2.9	4.9	2.1
2.5–3.0	—	2.1	2.0	0.6	0.9	1.3	1.7	1.3	1.9	1.4	1.4	0.8	1.6	3.2	3.0	1.3	2.9	1.3
3.5–4.0	—	2.6	2.3	1.6	1.3	1.5	1.1	1.1	1.3	1.1	1.1	2.5	1.9	1.9	2.8	2.7	2.5	1.7
4.5–5.0	—	1.3	1.1	1.6	0.9	1.3	0.6	2.0	1.1	1.2	1.9	2.2	1.4	1.9	2.3	2.3	1.8	1.3
5.5–6.0 (High)	—	2.1	1.0	0.5	0.8	0.8	1.2	1.4	1.1	1.3	1.5	1.1	2.4	2.2	1.0	1.4	2.4	1.2
Race/Ethnicity (2-year average) ^c																		
White	—	—	1.6	1.3	1.3	1.2	1.1	1.4	1.3	1.2	1.5	1.7	1.8	2.1	2.5	2.4	2.4	2.1
African American	—	—	1.8	1.2	0.5	1.1	1.8	1.2	1.4	1.5	0.9	0.7	1.0	1.2	1.0	1.1	1.3	1.6
Hispanic	—	—	2.3	3.3	1.8	0.9	1.7	1.3	0.6	1.6	2.4	2.9	2.4	2.1	2.2	1.8	2.4	2.0

Table continued on next page.

TABLE 159 (cont.)
STEROIDS: ^a Trends in Annual Prevalence of Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021	
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	change
<i>Approximate weighted N =</i>	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	1.8	1.4	1.5	1.5	1.5	1.2	1.3	1.5	1.5	1.7	1.0	1.1	1.1	1.0	1.2	0.5	-0.5* s
Gender																	
Male	2.7	2.3	2.5	2.5	2.5	1.8	1.7	2.2	2.0	2.5	1.3	1.4	1.5	1.4	§	1.0	-0.4
Female	0.7	0.6	0.4	0.4	0.3	0.5	0.7	0.7	0.7	0.7	0.6	0.5	0.4	0.6	§	0.1	-0.5 s
College Plans																	
None or under 4 years	2.7	2.1	3.1	2.9	3.3	1.6	1.4	2.7	1.8	1.8	2.1	2.3	2.2	3.0	§	1.2	-1.8
Complete 4 years	1.5	1.3	1.1	1.2	1.1	1.1	1.1	1.1	1.3	1.5	0.8	0.7	0.7	0.5	§	0.3	-0.2
Region																	
Northeast	2.0	1.6	2.2	1.3	1.7	1.2	1.6	1.0	2.1	1.1	0.9	0.8	0.6	1.5	§	0.0	-1.5
Midwest	1.5	1.3	1.7	1.4	1.6	0.6	0.9	1.6	0.9	1.6	1.1	0.9	1.4	0.8	§	1.3	+0.5
South	2.0	2.0	1.4	1.7	1.5	1.5	1.6	1.9	1.4	1.9	1.1	1.4	1.4	1.1	§	0.4	-0.7 s
West	1.5	0.5	0.9	1.4	1.1	1.5	1.0	0.9	1.5	2.0	1.0	0.8	0.7	0.7	§	0.5	-0.2
Population Density																	
Large MSA	1.6	1.0	0.9	1.3	1.2	1.0	1.4	1.1	0.8	1.8	0.7	0.6	1.4	1.1	§	0.5	-0.6
Other MSA	1.6	1.5	1.4	1.7	1.5	1.4	1.2	1.1	1.9	1.9	1.3	1.0	1.0	0.9	§	0.2	-0.7
Non-MSA	2.2	1.9	2.4	1.1	1.6	1.2	1.0	2.8	1.1	1.2	0.9	1.8	0.9	1.3	§	1.2	-0.1
Parental Education ^b																	
1.0–2.0 (Low)	2.8	3.0	2.1	1.7	1.6	2.2	1.5	3.3	1.9	1.1	1.5	1.4	1.6	1.1	§	0.0	-1.1
2.5–3.0	1.8	1.6	1.6	1.8	2.2	1.3	1.0	1.5	1.4	1.1	0.9	1.4	1.3	0.5	§	0.0	-0.5
3.5–4.0	1.7	1.1	1.3	1.6	1.6	1.5	1.1	0.8	1.1	2.1	1.1	0.7	0.7	1.8	§	1.1	-0.7
4.5–5.0	1.3	1.1	1.4	1.3	0.7	0.6	1.0	1.7	1.2	1.9	0.7	1.0	1.3	0.6	§	0.8	+0.2
5.5–6.0 (High)	1.5	1.5	1.5	0.9	1.4	0.8	1.8	0.5	1.4	1.9	1.2	0.6	0.7	0.8	§	0.3	-0.5
Race/Ethnicity (2-year average) ^c																	
White	1.6	1.5	1.5	1.5	1.5	1.3	1.2	1.3	1.1	1.3	1.4	0.9	0.8	0.8	§	0.5	-0.3
African American	1.7	1.6	1.4	1.5	1.7	1.8	1.6	1.4	2.7	2.2	1.3	2.2	2.4	1.8	§	1.9	+0.1
Hispanic	2.3	2.3	1.3	1.0	1.3	1.4	1.0	1.1	1.4	1.3	1.0	0.7	1.1	1.4	§	0.5	-0.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms in 1989–1990; *N* is one sixth of *N* indicated in Table S3. Data based on two of six forms in 1991–2005; *N* is two sixths of *N* indicated in Table S3. Data based on three of six forms beginning in 2006; *N* is three sixths of *N* indicated in Table S3. In 2006 there were no changes in the two forms containing the original version of the question text. A slightly altered version of the question was added to a third form. An examination of the data did not show any effect from the wording change. In 2007 the forms containing the original version were changed in a like manner. In 2008 the question wording was changed slightly in two of the questionnaire forms. An examination of the data did not show any effect from the wording changes. In 2009 the remaining form was changed in a like manner.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aOnly drug use not under a doctor's orders is included here.

^bParental education is an average score of mother's education and father's education. See appendix B for details.

^cEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 160
MARIJUANA^c: Trends in Lifetime Prevalence of Daily Use for a Month or More by Subgroups in Grade 12

	Percentage who used daily for month or more in lifetime															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Approximate weighted N =	—	—	—	—	—	—	—	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total ^d	—	—	—	—	—	—	—	20.5	16.8	16.3	15.6	14.9	14.7	12.8	11.5	10.0
Gender:																
Male	—	—	—	—	—	—	—	20.1	18.1	17.2	17.7	16.6	16.2	14.8	12.7	10.6
Female	—	—	—	—	—	—	—	18.0	13.5	12.9	12.0	11.6	12.2	9.6	9.7	7.9
College Plans:																
None or under 4 years	—	—	—	—	—	—	—	22.5	20.3	18.9	19.6	17.2	18.0	14.5	15.3	12.8
Complete 4 years	—	—	—	—	—	—	—	13.8	10.5	10.7	10.6	11.0	11.1	9.8	9.1	7.4
Region:																
Northeast	—	—	—	—	—	—	—	25.1	20.4	24.1	20.9	21.5	17.0	13.1	14.6	10.4
Midwest	—	—	—	—	—	—	—	21.1	15.9	12.8	16.3	11.3	12.7	10.3	13.4	10.8
South	—	—	—	—	—	—	—	15.7	12.7	14.0	8.9	11.3	11.9	10.9	8.1	8.7
West	—	—	—	—	—	—	—	20.8	21.4	17.6	18.5	18.3	19.7	19.0	12.3	11.0
Population Density:																
Large MSA	—	—	—	—	—	—	—	23.8	20.0	19.4	18.1	17.0	16.7	14.0	10.6	8.3
Other MSA	—	—	—	—	—	—	—	20.3	18.2	16.6	16.0	14.9	15.0	14.9	12.4	11.7
Non-MSA	—	—	—	—	—	—	—	17.9	12.6	13.2	12.8	13.2	12.2	7.6	10.4	8.2
Parental Education: ^a																
1.0–2.0 (Low)	—	—	—	—	—	—	—	20.9	15.5	15.4	15.0	17.2	11.2	11.9	8.4	9.3
2.5–3.0	—	—	—	—	—	—	—	20.1	15.5	16.8	15.3	13.3	15.6	10.0	10.4	10.4
3.5–4.0	—	—	—	—	—	—	—	20.1	14.7	15.9	15.2	14.5	14.2	12.8	14.0	8.0
4.5–5.0	—	—	—	—	—	—	—	16.8	17.0	10.8	12.8	12.4	14.2	14.5	8.5	10.3
5.5–6.0 (High)	—	—	—	—	—	—	—	12.8	15.3	14.8	11.8	14.6	11.4	10.0	13.5	7.5
Race/Ethnicity (2-year average): ^b																
White	—	—	—	—	—	—	—	18.2	16.2	15.8	15.0	15.0	15.1	14.6	13.0	11.4
African American	—	—	—	—	—	—	—	11.3	9.9	9.8	9.2	9.2	8.6	6.9	6.2	4.2
Hispanic	—	—	—	—	—	—	—	19.5	14.8	11.0	11.8	11.8	11.3	11.4	9.4	8.3

Table continued on next page.

TABLE 160 (cont.)
MARIJUANA^c: Trends in Lifetime Prevalence of Daily Use for a Month or More by Subgroups in Grade 12

	Percentage who used daily for month or more in lifetime															
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total ^d	9.0	8.4	9.6	11.3	12.1	15.7	18.8	18.0	17.9	17.0	18.0	15.5	16.4	17.8	14.5	16.6
Gender:																
Male	10.5	8.3	10.7	13.3	12.9	18.7	19.7	19.5	18.5	18.8	20.3	17.2	17.1	19.9	15.8	17.0
Female	6.4	7.5	7.2	8.5	7.9	10.7	15.2	13.9	14.4	13.7	13.8	11.7	12.5	12.3	11.1	13.3
College Plans:																
None or under 4 years	11.5	11.2	11.6	16.1	14.2	21.5	22.6	22.1	22.1	19.1	22.8	20.5	22.2	24.5	20.0	24.8
Complete 4 years	6.5	5.9	7.7	8.6	9.2	11.9	14.9	13.4	14.2	13.7	13.8	11.7	11.9	13.4	11.4	12.2
Region:																
Northeast	10.3	8.7	12.0	12.2	12.8	21.3	24.6	22.7	17.9	19.8	23.4	20.7	20.8	19.5	15.5	17.7
Midwest	8.4	8.0	9.3	11.0	13.6	14.6	16.5	16.1	14.3	13.8	18.4	16.3	15.0	17.9	16.6	16.3
South	7.4	5.9	8.3	11.8	11.2	12.7	14.9	15.6	19.1	14.7	12.7	14.6	15.5	18.6	15.8	17.1
West	11.3	13.4	10.4	10.2	10.6	17.0	23.0	20.6	20.4	21.9	21.2	11.7	15.4	14.3	8.9	15.2
Population Density:																
Large MSA	7.2	8.4	8.6	10.3	13.9	15.3	18.8	18.0	16.3	18.4	19.7	15.2	14.0	17.0	13.7	18.7
Other MSA	11.1	8.9	10.2	13.6	11.3	18.2	20.1	19.7	19.2	18.3	17.5	15.1	19.0	19.5	14.6	14.2
Non-MSA	7.1	7.6	9.6	8.4	11.2	11.6	16.2	14.4	17.1	13.0	17.1	16.8	14.8	15.5	15.3	18.2
Parental Education: ^a																
1.0–2.0 (Low)	9.6	12.4	7.2	10.3	10.0	14.1	16.1	18.0	13.9	20.9	16.7	13.7	16.4	17.2	11.1	26.5
2.5–3.0	7.1	8.3	10.8	8.8	11.2	16.3	20.2	16.5	15.6	16.6	16.7	17.7	15.0	17.5	18.0	15.4
3.5–4.0	9.3	8.6	9.8	12.6	11.5	15.1	14.9	17.6	18.0	16.0	19.4	13.3	14.1	18.4	12.9	16.5
4.5–5.0	8.2	4.9	8.0	10.2	9.4	13.2	18.1	14.7	18.7	12.7	15.2	16.0	14.6	16.1	11.1	12.6
5.5–6.0 (High)	7.9	6.5	6.4	9.6	10.1	14.3	14.1	16.0	9.1	17.5	9.8	6.9	10.6	11.4	11.6	9.9
Race/Ethnicity (2-year average): ^b																
White	9.6	8.7	9.4	10.8	11.0	13.0	16.8	17.7	16.7	16.2	17.1	16.7	16.2	17.1	16.9	16.3
African American	2.6	2.9	4.2	7.3	9.7	10.3	11.2	13.3	14.5	13.6	12.4	10.2	8.7	11.0	11.9	12.7
Hispanic	10.7	11.7	7.9	7.5	9.1	13.0	18.3	18.2	21.1	24.3	16.8	13.2	10.8	10.9	11.0	11.6

Table continued on next page.

TABLE 160 (cont.)
MARIJUANA^c: Trends in Lifetime Prevalence of Daily Use for a Month or More by Subgroups in Grade 12

	Percentage who used daily for month or more in lifetime															2019-2021 change
	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total ^d	15.7	15.1	14.9	15.5	17.4	18.2	15.8	13.7	12.4	14.3	13.9	12.3	14.9	§	12.4	-2.5*
Gender:																
Male	17.5	15.3	16.8	17.4	20.4	21.6	18.4	15.5	13.1	17.2	15.7	12.3	16.2	§	13.0	-3.2
Female	12.6	12.8	10.7	11.5	12.0	11.9	11.7	10.9	9.8	10.3	10.5	9.3	11.7	§	11.8	+0.2
College Plans:																
None or under 4 years	21.8	23.4	18.6	21.9	26.4	28.0	24.3	22.9	19.0	19.2	22.5	18.5	16.4	§	19.4	+3.1
Complete 4 years	13.3	11.0	11.8	12.5	13.9	14.1	12.5	10.3	9.7	11.5	10.4	8.1	12.3	§	8.5	-3.8 s
Region:																
Northeast	19.0	19.7	14.2	16.0	18.6	17.6	17.5	15.9	13.6	12.3	13.6	9.7	15.4	§	10.4	-5.0
Midwest	17.1	13.8	17.0	15.4	15.9	15.7	14.5	11.3	10.8	11.1	13.5	14.6	14.3	§	9.8	-4.5
South	14.2	13.3	15.9	14.8	17.5	16.7	14.2	12.8	13.8	15.5	13.6	11.0	14.2	§	9.2	-5.0 s
West	13.9	15.6	11.2	16.4	17.8	23.2	18.2	15.6	10.4	16.6	14.9	14.0	17.2	§	20.4	+3.3
Population Density:																
Large MSA	14.2	18.3	13.8	16.6	18.6	20.2	16.8	13.8	12.5	17.3	14.1	12.2	15.6	§	11.9	-3.8
Other MSA	16.2	13.5	15.4	16.5	17.1	18.2	16.3	14.0	11.6	10.9	14.0	11.5	15.2	§	12.7	-2.5
Non-MSA	16.8	14.8	15.1	11.5	16.1	14.9	13.3	13.0	14.2	17.8	13.2	14.3	12.6	§	12.5	-0.1
Parental Education: ^a																
1.0–2.0 (Low)	15.8	17.7	13.8	13.6	16.4	20.6	21.4	12.5	12.9	13.7	17.4	10.9	15.8	§	14.1	-1.7
2.5–3.0	18.0	18.5	16.2	14.5	15.2	18.3	18.0	17.2	15.2	13.3	15.0	10.3	15.7	§	16.2	+0.5
3.5–4.0	15.7	12.3	13.2	17.2	16.5	16.5	16.9	13.6	11.9	16.2	13.3	11.7	15.1	§	13.3	-1.8
4.5–5.0	13.3	12.1	11.5	10.8	16.5	16.4	12.7	11.5	10.2	12.3	9.1	9.8	11.8	§	8.1	-3.7
5.5–6.0 (High)	12.2	9.2	11.0	10.7	15.6	13.9	8.1	5.4	6.6	9.3	7.5	8.4	7.7	§	8.1	+0.4
Race/Ethnicity (2-year average): ^b																
White	15.9	15.0	14.3	14.3	15.3	16.8	16.2	13.7	11.2	11.5	12.5	11.6	11.2	§	12.3	+1.1
African American	13.9	10.4	10.8	14.7	14.6	14.5	13.7	14.0	14.1	14.3	14.9	12.8	13.3	§	16.6	+3.3
Hispanic	13.8	13.9	13.0	12.8	15.4	19.0	18.5	15.8	13.6	12.8	14.7	12.2	12.6	§	10.2	-2.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. Data based on one of six forms; N is one sixth of N indicated in Table S3. See appendix B for definition of variables in table.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^dTotal prevalence includes respondents who are missing data on some subgroups, and thus may appear inconsistent with subgroup prevalence values.

TABLE 161
NONPRESCRIPTION DIET PILLS: Trends in Annual Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	—	—	—	—	—	—	—	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	—	—	—	—	—	—	20.5	20.5	18.8	16.9	15.3	13.9	12.2	10.9	10.4
Gender:																
Male	—	—	—	—	—	—	—	10.7	10.6	9.2	9.0	6.9	6.4	4.9	4.3	4.3
Female	—	—	—	—	—	—	—	29.6	30.0	27.5	24.4	23.2	21.1	18.8	17.2	16.7
College Plans:																
None or under 4 years	—	—	—	—	—	—	—	23.3	21.4	18.2	20.7	16.1	14.9	13.1	11.9	13.2
Complete 4 years	—	—	—	—	—	—	—	17.5	19.0	18.8	14.7	15.0	13.3	11.7	10.9	9.7
Region:																
Northeast	—	—	—	—	—	—	—	19.1	18.5	18.4	16.5	14.9	14.3	10.5	10.4	11.5
Midwest	—	—	—	—	—	—	—	24.6	23.3	20.2	19.2	16.6	15.0	13.7	15.0	11.1
South	—	—	—	—	—	—	—	18.2	19.2	19.6	14.9	13.9	13.1	12.0	9.3	10.0
West	—	—	—	—	—	—	—	18.9	21.1	15.8	17.3	16.4	13.5	12.1	8.7	8.9
Population Density:																
Large MSA	—	—	—	—	—	—	—	19.7	18.7	17.3	17.1	15.0	13.0	12.1	10.3	7.4
Other MSA	—	—	—	—	—	—	—	20.0	22.8	18.6	17.1	15.6	13.7	12.4	10.9	11.2
Non-MSA	—	—	—	—	—	—	—	21.7	19.2	20.5	16.5	15.2	15.2	11.9	11.7	11.7
Parental Education: ^a																
1.0–2.0 (Low)	—	—	—	—	—	—	—	19.5	15.9	19.2	14.6	12.0	13.5	13.4	9.5	5.1
2.5–3.0	—	—	—	—	—	—	—	21.6	21.3	18.2	17.8	15.6	13.1	12.0	9.9	12.3
3.5–4.0	—	—	—	—	—	—	—	20.6	20.2	20.6	18.0	16.6	14.5	11.5	11.8	9.2
4.5–5.0	—	—	—	—	—	—	—	19.3	22.4	17.4	16.8	15.0	15.9	12.0	10.4	12.0
5.5–6.0 (High)	—	—	—	—	—	—	—	21.0	22.3	19.1	17.1	15.7	11.6	13.6	13.4	12.2
Race/Ethnicity (2-year average): ^b																
White	—	—	—	—	—	—	—	23.2	22.2	20.4	18.4	16.3	14.7	13.4	12.5	
African American	—	—	—	—	—	—	—	6.6	8.1	6.4	5.5	7.5	6.9	4.3	2.9	
Hispanic	—	—	—	—	—	—	—	11.6	12.6	14.8	10.8	7.8	7.9	9.6	9.8	

Table continued on next page.

TABLE 161 (cont.)
NONPRESCRIPTION DIET PILLS: Trends in Annual Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	8.8	8.4	8.0	9.3	9.8	9.3	9.8	9.6	10.2	11.1	11.8	15.1	13.0	10.7	10.0	9.4
Gender:																
Male	3.0	4.3	3.2	2.5	3.5	3.7	4.9	4.3	4.0	4.9	6.2	8.1	6.9	6.5	6.0	5.7
Female	14.2	12.2	12.3	15.0	15.1	14.1	14.6	15.4	15.7	17.2	15.6	20.0	17.5	14.1	13.2	12.5
College Plans:																
None or under 4 years	8.9	10.5	9.7	10.8	11.5	10.8	12.0	10.7	9.1	10.0	9.4	17.8	13.9	10.4	11.5	12.1
Complete 4 years	8.6	8.0	7.3	9.3	9.3	8.6	9.2	10.1	10.4	11.5	11.6	13.8	12.6	10.5	9.3	8.6
Region:																
Northeast	5.7	6.3	7.6	8.6	8.2	10.1	8.1	8.6	8.6	8.0	12.8	16.7	11.6	11.9	9.1	12.0
Midwest	10.7	9.3	8.4	11.8	11.8	9.5	10.2	10.4	10.4	12.6	15.5	14.4	14.4	9.1	10.0	9.9
South	9.0	7.7	9.2	8.9	10.8	9.4	11.5	10.1	11.2	12.9	9.9	16.7	13.6	10.5	10.4	9.2
West	8.8	10.3	5.4	7.4	6.3	7.9	7.8	8.6	9.3	9.3	8.6	12.4	11.7	12.2	10.1	6.5
Population Density:																
Large MSA	7.7	7.4	7.3	8.5	10.7	8.5	9.4	9.5	9.0	8.3	12.3	13.9	12.5	9.8	7.9	8.0
Other MSA	9.2	8.4	6.8	9.9	8.9	9.3	8.7	9.1	9.5	11.5	11.5	13.6	10.8	11.5	10.7	9.9
Non-MSA	9.1	9.2	10.5	9.1	10.1	10.0	12.3	11.0	12.6	13.6	11.7	19.8	17.7	10.3	11.1	10.3
Parental Education: ^a																
1.0–2.0 (Low)	9.4	10.6	6.9	11.0	8.9	8.3	14.4	11.8	9.9	9.8	5.6	12.7	15.8	7.0	10.5	6.9
2.5–3.0	8.6	8.9	8.9	11.0	10.9	8.5	10.5	8.9	11.1	13.1	15.0	17.7	13.8	14.1	11.1	13.3
3.5–4.0	8.2	7.9	7.8	10.6	11.5	9.7	8.8	9.9	10.5	11.6	14.1	15.6	11.1	10.7	10.2	9.0
4.5–5.0	9.3	6.6	8.0	8.5	7.5	8.8	10.7	11.6	11.2	12.7	9.0	14.8	11.5	9.0	8.9	8.6
5.5–6.0 (High)	8.5	8.2	6.2	5.3	9.2	8.4	9.4	9.6	7.4	7.4	7.9	8.5	13.4	10.1	7.5	6.4
Race/Ethnicity (2-year average): ^b																
White	11.0	9.8	9.2	9.7	10.9	10.9	10.8	11.2	11.2	11.8	12.9	14.8	15.4	13.1	11.3	10.7
African American	3.5	3.0	4.5	6.1	5.3	4.3	5.3	4.9	4.2	5.5	6.5	5.5	4.6	4.8	4.5	4.2
Hispanic	5.6	4.6	7.1	7.1	7.5	7.0	8.2	9.3	9.8	10.0	9.1	13.2	12.7	7.9	7.5	7.7

Table continued on next page.

TABLE 161 (cont.)
NONPRESCRIPTION DIET PILLS: Trends in Annual Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	6.7	7.2	6.1	4.3	4.9	5.5	5.3	6.4	5.1	4.5	4.0	3.5	3.1	§	2.5	-0.6*
Gender:																
Male	3.4	4.2	3.0	2.4	3.0	3.7	2.6	3.7	1.6	1.6	2.7	2.0	1.6	§	1.6	0.0
Female	9.2	9.9	9.3	6.0	6.1	7.0	7.3	8.6	8.3	7.1	4.5	4.5	3.9	§	3.4	-0.5
College Plans:																
None or under 4 years	4.4	10.2	9.4	4.3	6.1	5.2	5.2	12.0	4.8	4.7	3.4	3.8	3.5	§	2.0	-1.5
Complete 4 years	7.1	6.8	5.6	4.5	4.4	5.5	4.8	5.0	5.1	4.6	3.5	3.6	2.9	§	2.7	-0.2
Region:																
Northeast	7.2	6.3	4.6	3.5	3.2	4.8	5.7	8.1	5.0	4.6	3.6	3.8	2.5	§	3.6	+1.1
Midwest	7.4	8.0	6.0	5.4	5.8	4.3	5.8	6.4	6.3	4.4	3.6	3.9	2.1	§	3.3	+1.1
South	6.9	6.5	6.8	3.8	4.7	7.4	6.0	6.5	4.5	4.9	5.0	3.7	3.1	§	1.5	-1.6
West	5.3	8.0	6.6	4.7	5.4	4.8	3.3	4.6	5.3	3.9	2.8	2.8	5.5	§	2.6	-2.9
Population Density:																
Large MSA	4.8	6.8	5.2	3.9	4.2	6.1	5.7	4.4	4.6	3.2	3.3	2.8	3.1	§	4.0	+0.9
Other MSA	7.4	7.0	5.8	5.0	4.3	5.6	4.7	7.2	5.0	5.2	3.9	3.5	3.3	§	2.0	-1.2
Non-MSA	8.3	7.9	8.0	3.1	7.3	4.6	6.1	7.1	6.3	4.9	5.2	4.9	2.8	§	1.1	-1.6
Parental Education: ^a																
1.0–2.0 (Low)	5.8	7.4	5.9	4.4	5.5	8.9	5.9	6.1	4.4	4.2	2.4	2.1	5.0	§	1.8	-3.2
2.5–3.0	8.8	7.8	8.2	4.6	4.7	4.5	6.9	6.1	4.9	5.0	5.6	3.6	2.2	§	0.5	-1.7
3.5–4.0	6.0	7.5	5.9	5.0	6.0	4.9	5.4	7.9	8.0	5.1	3.1	4.8	2.0	§	2.6	+0.5
4.5–5.0	5.0	8.7	4.9	3.7	4.7	5.7	3.1	6.3	4.4	4.7	4.1	3.7	3.2	§	3.1	-0.1
5.5–6.0 (High)	7.4	3.0	6.5	4.9	2.7	3.7	3.4	4.2	3.0	3.0	3.3	4.0	2.1	§	4.7	+2.6
Race/Ethnicity (2-year average): ^b																
White	9.3	8.1	7.2	4.8	4.3	4.9	4.8	5.6	6.0	5.7	5.0	4.4	3.4	§	1.7	-1.7 ss
African American	3.2	2.5	3.4	4.4	3.0	3.3	3.6	4.3	3.6	1.8	2.4	3.0	2.4	§	1.4	-1.0
Hispanic	4.9	5.2	7.2	6.4	5.5	5.6	5.6	5.1	5.0	4.4	2.6	1.9	2.9	§	1.3	-1.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. Data based on one of six forms; N is one sixth of N indicated in Table S3. See appendix B for definition of variables in table.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 162
NONPRESCRIPTION STAY-AWAKE PILLS: Trends in Annual Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
<i>Approximate weighted N =</i>	—	—	—	—	—	—	—	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Total	—	—	—	—	—	—	—	11.8	12.3	13.9	18.2	22.3	25.2	26.4	23.0	23.4
Gender:																
Male	—	—	—	—	—	—	—	12.8	13.8	15.4	19.7	22.3	25.5	27.6	24.8	22.3
Female	—	—	—	—	—	—	—	10.0	10.5	12.5	17.0	22.2	25.0	25.2	21.7	24.5
College Plans:																
None or under 4 years	—	—	—	—	—	—	—	11.4	10.8	12.5	15.5	18.1	23.1	23.3	21.8	22.9
Complete 4 years	—	—	—	—	—	—	—	10.5	12.6	14.0	20.4	24.9	26.5	27.5	24.1	24.1
Region:																
Northeast	—	—	—	—	—	—	—	9.6	9.5	11.9	18.2	20.4	26.4	23.8	18.4	22.0
Midwest	—	—	—	—	—	—	—	15.1	16.8	16.3	18.4	24.5	26.8	27.5	29.1	28.4
South	—	—	—	—	—	—	—	9.6	10.7	12.0	13.3	19.8	20.9	25.6	20.4	20.6
West	—	—	—	—	—	—	—	13.5	11.5	16.0	25.6	25.5	28.9	28.9	24.0	22.9
Population Density:																
Large MSA	—	—	—	—	—	—	—	11.5	11.1	12.2	17.7	19.0	25.6	23.2	20.9	19.0
Other MSA	—	—	—	—	—	—	—	12.4	14.5	14.0	19.1	24.1	24.1	27.7	22.9	25.1
Non-MSA	—	—	—	—	—	—	—	11.3	10.5	15.4	17.4	22.1	27.0	27.4	25.2	24.5
Parental Education: ^a																
1.0–2.0 (Low)	—	—	—	—	—	—	—	8.8	8.2	8.3	10.9	12.0	15.0	17.1	18.1	15.3
2.5–3.0	—	—	—	—	—	—	—	10.6	10.1	13.9	15.5	19.5	22.8	22.5	21.6	23.2
3.5–4.0	—	—	—	—	—	—	—	12.6	13.5	13.5	21.3	26.5	30.0	28.4	26.0	25.6
4.5–5.0	—	—	—	—	—	—	—	13.2	15.3	16.1	24.0	23.7	29.9	30.3	24.0	28.0
5.5–6.0 (High)	—	—	—	—	—	—	—	13.0	16.7	18.0	20.2	28.7	24.9	32.3	25.1	22.3
Race/Ethnicity (2-year average): ^b																
White	—	—	—	—	—	—	—	—	13.6	15.0	18.7	23.7	27.7	30.4	29.8	27.7
African American	—	—	—	—	—	—	—	—	2.0	3.0	3.6	4.4	5.5	6.2	6.0	6.4
Hispanic	—	—	—	—	—	—	—	—	5.7	8.3	8.4	9.7	13.8	15.6	16.5	14.1

Table continued on next page.

TABLE 162 (cont.)
NONPRESCRIPTION STAY-AWAKE PILLS: Trends in Annual Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who used in last 12 months															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<i>Approximate weighted N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Total	22.2	20.4	19.2	20.7	20.3	19.0	19.7	19.0	15.7	15.0	17.3	14.9	12.5	11.8	10.4	10.0
Gender:																
Male	22.3	20.9	19.7	20.3	19.7	18.2	17.4	19.5	14.5	14.0	17.8	13.9	9.3	11.1	9.4	10.2
Female	22.0	20.2	17.6	20.4	20.1	18.7	21.0	18.0	15.9	15.9	16.5	14.6	14.3	12.3	10.3	9.4
College Plans:																
None or under 4 years	21.0	20.5	18.6	20.1	21.1	18.2	21.8	18.5	14.3	15.5	18.9	16.1	14.3	13.0	11.2	9.1
Complete 4 years	22.3	21.0	18.7	20.6	19.7	18.3	19.1	18.4	15.1	14.7	16.5	14.0	11.9	11.4	9.3	9.8
Region:																
Northeast	18.3	18.2	20.2	21.2	18.4	22.5	19.1	17.6	16.0	10.2	14.7	13.9	10.9	14.0	8.0	8.3
Midwest	31.8	25.7	22.0	26.2	24.2	19.8	23.8	22.0	17.3	19.3	24.4	18.9	12.9	12.3	13.4	14.1
South	16.1	17.6	18.7	20.2	18.8	17.5	20.1	18.8	15.6	13.8	15.4	13.5	11.6	9.7	11.0	9.7
West	23.4	20.0	14.9	13.7	19.1	16.5	13.3	16.8	13.3	16.3	12.4	13.2	14.9	12.5	8.0	7.0
Population Density:																
Large MSA	16.7	15.2	16.7	18.4	18.9	14.4	15.5	15.3	9.6	11.0	15.7	11.9	9.2	8.3	7.4	7.7
Other MSA	25.3	21.2	19.8	21.1	19.3	20.2	18.4	21.1	18.4	15.2	14.3	14.7	12.6	13.6	11.7	10.2
Non-MSA	21.7	23.4	19.9	22.3	23.6	20.7	26.8	18.9	17.3	19.3	24.3	19.3	16.5	12.2	11.4	12.7
Parental Education: ^a																
1.0–2.0 (Low)	14.9	16.2	11.0	16.6	13.2	11.5	18.2	11.7	7.9	7.3	15.3	8.9	10.3	5.9	4.2	6.1
2.5–3.0	20.2	22.8	19.0	18.1	18.1	19.0	21.0	16.5	13.4	15.1	17.1	13.4	12.0	13.9	9.7	9.0
3.5–4.0	23.9	22.4	18.6	21.6	24.3	17.4	17.6	19.9	18.3	17.0	20.3	16.5	11.4	13.5	12.8	11.4
4.5–5.0	25.1	20.0	21.1	24.4	20.4	23.2	20.2	20.3	15.6	16.7	16.2	14.9	12.7	11.3	9.7	10.0
5.5–6.0 (High)	25.8	17.8	20.2	18.4	17.3	17.4	19.3	22.6	14.9	13.4	13.6	15.7	14.5	11.4	8.3	8.9
Race/Ethnicity (2-year average): ^b																
White	27.4	25.8	24.0	23.7	23.9	23.3	23.1	23.2	20.7	18.2	19.9	19.5	16.0	14.2	12.8	11.6
African American	5.1	3.5	3.5	3.6	4.8	4.4	5.3	6.2	3.6	3.0	4.2	3.2	2.2	1.9	2.3	2.6
Hispanic	11.6	11.9	13.3	14.2	12.3	9.5	9.6	10.1	12.3	11.8	10.6	12.3	9.4	5.6	5.7	6.9

Table continued on next page.

TABLE 162 (cont.)
NONPRESCRIPTION STAY-AWAKE PILLS: Trends in Annual Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who used in last 12 months															2019-2021 change
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	7.6	6.3	4.8	3.2	3.9	3.8	3.2	3.5	2.7	2.5	2.5	2.4	1.8	§	1.5	-0.3*
Gender:																
Male	7.8	5.2	5.9	3.2	2.8	3.9	2.9	2.8	2.1	2.3	2.6	1.9	1.8	§	0.4	-1.5 ss
Female	7.1	7.0	3.5	3.1	4.6	3.8	3.4	3.6	2.7	2.5	1.7	2.7	1.3	§	2.0	+0.8
College Plans:																
None or under 4 years	9.5	9.5	3.9	3.0	5.1	4.4	3.0	4.4	3.1	1.1	2.1	5.0	2.6	§	2.6	0.0
Complete 4 years	7.1	5.3	4.7	3.3	3.6	3.5	2.9	3.3	2.4	2.8	2.0	1.4	1.4	§	0.8	-0.5
Region:																
Northeast	5.7	6.7	5.6	3.0	2.7	3.0	4.0	2.6	2.9	1.9	1.5	1.8	1.4	§	1.6	+0.2
Midwest	10.9	6.6	6.1	3.1	4.2	3.1	3.1	4.2	4.6	2.8	1.7	2.9	1.6	§	2.5	+0.9
South	7.4	5.9	4.6	3.3	2.8	4.2	2.7	3.9	2.3	2.0	3.7	1.9	2.1	§	0.7	-1.4 s
West	6.3	6.2	3.0	3.2	6.2	4.5	3.4	2.8	1.5	3.3	1.8	3.2	1.9	§	1.8	-0.1
Population Density:																
Large MSA	5.3	7.2	4.9	2.8	3.9	2.5	2.6	2.0	2.0	1.2	1.8	1.5	1.6	§	1.5	-0.1
Other MSA	8.9	5.0	4.4	3.8	3.8	4.1	3.3	3.9	2.8	3.0	2.7	2.3	2.0	§	1.8	-0.2
Non-MSA	8.3	8.1	5.9	2.3	4.1	5.1	3.7	4.6	3.5	3.1	3.1	4.2	1.7	§	0.9	-0.8
Parental Education: ^a																
1.0–2.0 (Low)	3.7	6.0	3.3	1.8	3.5	6.5	2.8	3.8	3.6	1.4	0.7	1.9	2.3	§	2.0	-0.3
2.5–3.0	9.5	6.7	4.7	1.6	4.3	4.7	3.1	2.0	2.2	2.0	3.9	3.0	2.2	§	0.5	-1.8 s
3.5–4.0	7.6	7.4	3.7	3.9	5.1	3.3	4.8	3.2	2.4	3.0	1.8	3.5	1.5	§	0.3	-1.2
4.5–5.0	6.2	4.9	4.5	3.3	3.1	3.3	2.0	3.5	2.8	3.0	2.6	0.9	1.2	§	1.7	+0.5
5.5–6.0 (High)	9.7	5.4	5.7	5.5	2.9	2.4	2.3	6.2	1.7	1.4	1.5	1.9	1.6	§	1.8	+0.2
Race/Ethnicity (2-year average): ^b																
White	10.6	8.4	6.3	4.4	3.8	4.0	3.6	3.3	3.3	3.3	2.5	2.2	2.0	§	1.0	-0.9
African American	1.2	1.2	3.0	3.3	2.0	2.2	2.3	2.1	1.8	1.2	1.1	1.1	2.5	§	1.6	-0.9
Hispanic	5.2	4.4	4.4	2.7	2.7	4.5	4.0	3.3	2.5	1.4	1.3	1.8	1.5	§	0.9	-0.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. Data based on one of six forms; N is one sixth of N indicated in Table S3. See appendix B for definition of variables in table.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 163
STIMULANT-TYPE ADHD DRUGS:
Trends in Lifetime Prevalence of Legal Use by Subgroups in Grade 8

	Percentage who used in lifetime																		2019– 2021 change
	1991– 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	8.3	9.3	8.3	8.1	7.8	8.2	7.6	7.7	7.1	7.2	7.1	7.5	6.6	7.1	6.5	5.0	9.0	+2.4* s
Gender:																			
Male	—	9.3	11.2	9.4	9.9	9.2	9.3	8.6	9.1	8.8	7.9	8.3	8.9	7.6	8.6	7.9	§	11.1	+3.2 s
Female	—	7.2	7.4	7.3	6.5	6.8	7.1	6.4	6.4	5.4	6.7	6.0	6.0	5.2	6.1	4.9	§	6.2	+1.3
College Plans:																			
None or under 4 years	—	14.4	16.6	10.8	10.8	7.7	11.1	11.3	10.3	12.4	12.2	9.6	13.5	8.6	11.1	10.2	§	11.7	+1.4
Complete 4 years	—	7.6	8.6	8.0	7.8	7.9	7.9	7.2	7.5	6.7	6.9	6.8	7.0	6.4	6.6	6.1	§	8.6	+2.5 s
Region:																			
Northeast	—	7.8	7.9	7.2	9.0	8.5	6.3	6.0	5.6	7.1	7.2	5.7	5.6	5.8	8.1	6.7	§	8.0	+1.4
Midwest	—	8.5	8.6	8.8	6.7	7.0	8.9	8.2	7.6	6.7	7.1	7.5	8.7	7.7	7.0	5.7	§	9.6	+3.9 ss
South	—	9.3	10.2	9.3	9.9	9.5	8.9	7.5	9.4	8.8	7.7	8.6	9.4	7.7	9.3	7.1	§	9.0	+1.8
West	—	6.7	9.7	6.8	6.6	5.6	7.4	8.4	6.0	4.6	6.7	4.9	4.2	4.5	3.2	6.2	§	9.0	+2.8
Population Density:																			
Large MSA	—	7.2	7.6	7.1	7.2	6.5	7.2	7.5	6.4	6.1	6.4	6.3	6.2	6.9	6.3	5.9	§	8.3	+2.5
Other MSA	—	10.0	10.3	9.0	9.32	8.2	8.9	7.7	8.1	7.1	8.2	7.4	8.3	6.5	8.8	6.3	§	10.0	+3.7 ss
Non-MSA	—	6.9	10.0	8.5	7.0	8.6	7.9	7.6	8.7	8.4	6.4	7.4	7.5	6.7	4.7	7.9	§	7.7	-0.3
Parental Education: ^a																			
1.0–2.0 (Low)	—	12.1	12.0	9.0	11.2	11.0	7.2	8.6	7.7	4.2	4.9	7.4	7.2	4.3	7.8	4.7	§	7.1	+2.3
2.5–3.0	—	7.9	10.6	9.1	7.8	7.5	8.9	8.0	8.8	6.0	7.2	7.6	8.5	6.4	5.9	8.4	§	9.4	+0.9
3.5–4.0	—	9.7	7.9	8.7	6.8	8.0	7.3	8.4	7.4	8.3	7.4	7.4	7.0	6.2	6.9	7.3	§	11.1	+3.9
4.5–5.0	—	6.9	9.0	7.5	8.6	7.5	8.3	7.1	7.8	7.8	7.6	7.0	7.4	6.9	8.4	6.6	§	8.4	+1.8
5.5–6.0 (High)	—	8.2	8.5	8.8	9.7	8.1	9.2	6.9	6.2	6.7	6.6	6.7	6.4	8.2	7.9	4.9	§	9.0	+4.1 s
Race/Ethnicity (2-year average): ^b																			
White	—	—	9.3	9.5	9.1	9.0	9.0	8.3	8.1	8.3	8.3	8.2	8.5	8.4	8.4	8.5	§	9.0	+0.5
African American	—	—	6.4	5.8	6.7	7.2	7.0	6.4	6.1	6.6	6.0	5.7	6.1	6.9	7.2	4.7	§	2.2	-2.5 s
Hispanic	—	—	7.5	7.4	6.6	6.1	5.8	7.1	6.4	4.1	4.9	5.5	5.2	4.2	3.9	4.4	§	2.8	-1.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 164
STIMULANT-TYPE ADHD DRUGS:
Trends in Lifetime Prevalence of Legal Use by Subgroups in Grade 10

	Percentage who used in lifetime																		2019– 2021 change
	1991– 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	8.7	8.5	8.4	7.8	8.2	8.6	7.2	8.0	8.3	6.8	8.8	7.1	6.5	8.2	6.6	6.0	7.0	+0.4*
Gender:																			
Male	—	10.4	9.2	9.6	9.3	9.1	10.0	8.1	8.8	10.9	8.3	9.9	7.4	7.4	9.4	7.7	§	6.5	-1.2
Female	—	6.9	7.7	7.4	6.4	7.3	7.2	6.4	7.2	5.7	5.3	7.7	6.8	5.6	6.9	5.5	§	7.1	+1.6
College Plans:																			
None or under 4 years	—	12.6	14.7	10.6	13.3	14.4	12.8	10.2	11.7	11.7	12.5	15.0	12.4	9.6	11.7	11.6	§	11.0	-0.5
Complete 4 years	—	8.2	7.6	8.1	7.2	7.6	8.2	6.9	7.6	8.0	6.1	8.2	6.6	6.2	7.5	5.6	§	6.2	+0.5
Region:																			
Northeast	—	9.1	6.4	7.0	6.8	7.5	7.4	6.1	8.0	9.2	6.2	8.9	6.5	5.6	5.7	5.7	§	7.1	+1.4
Midwest	—	8.0	7.9	8.6	8.9	9.9	10.3	9.0	6.7	9.1	7.8	8.5	6.3	5.4	11.2	6.8	§	8.5	+1.7
South	—	10.2	10.8	9.3	8.7	9.7	9.9	8.8	10.5	9.2	8.6	10.3	9.0	8.0	8.2	5.9	§	7.5	+1.6
West	—	6.8	7.6	8.0	6.4	5.2	5.8	3.6	6.0	6.0	3.5	6.6	5.6	6.2	6.6	8.0	§	4.8	-3.2
Population Density:																			
Large MSA	—	7.5	7.5	6.8	6.7	8.5	8.1	6.4	6.9	8.2	5.8	8.6	6.1	6.1	7.1	3.6	§	6.5	+2.9 s
Other MSA	—	9.4	9.2	9.2	7.3	8.3	8.7	7.3	8.7	8.2	6.9	9.0	7.0	6.9	8.4	8.6	§	7.7	-0.9
Non-MSA	—	8.6	8.5	9.0	10.5	7.7	9.3	8.1	8.1	8.8	8.1	8.8	9.2	6.4	9.0	6.2	§	5.9	-0.3
Parental Education: ^a																			
1.0–2.0 (Low)	—	6.6	8.7	8.0	5.8	7.3	9.0	5.4	7.2	7.8	4.0	7.4	5.5	6.0	7.4	4.1	§	4.0	-0.1
2.5–3.0	—	8.6	9.0	8.8	8.2	7.9	10.6	8.3	9.6	8.5	5.2	8.0	7.1	6.8	7.5	5.9	§	8.7	+2.9
3.5–4.0	—	8.2	9.8	7.7	8.3	7.6	8.4	7.5	6.9	9.3	7.0	10.0	6.4	6.3	7.8	8.1	§	6.4	-1.6
4.5–5.0	—	8.7	7.4	8.8	7.6	8.5	7.9	7.0	8.4	8.5	6.8	7.6	6.8	7.2	9.7	7.2	§	7.0	-0.3
5.5–6.0 (High)	—	10.0	8.0	8.8	7.9	10.3	8.8	6.0	8.4	8.9	8.6	10.4	8.9	6.4	8.6	6.7	§	7.8	+1.0
Race/Ethnicity (2-year average): ^b																			
White	—	—	9.8	9.4	9.2	8.9	9.4	9.1	8.7	9.4	9.0	9.4	9.7	8.5	9.3	9.6	§	8.2	-1.4
African American	—	—	5.2	6.7	6.0	4.8	6.1	6.8	5.7	5.9	5.6	6.3	6.7	5.1	4.1	3.9	§	5.0	+1.1
Hispanic	—	—	5.6	5.4	4.8	5.3	5.9	5.1	5.9	5.0	3.7	4.5	4.5	4.6	5.5	4.4	§	3.1	-1.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 165
STIMULANT-TYPE ADHD DRUGS:
Trends in Lifetime Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who used in lifetime																		2019– 2021 change
	1991– 2004	2005 ^c	2006 ^c	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	8.5	7.8	7.6	8.6	8.2	8.3	8.4	9.0	9.6	9.1	9.9	8.4	8.6	8.6	7.9	7.5	8.0	+0.1*
Gender:																			
Male	—	12.0	10.2	10.1	10.3	8.7	9.8	10.3	10.5	10.9	9.7	9.6	9.8	10.1	10.3	9.6	§	8.4	-1.2
Female	—	5.3	5.4	4.9	7.0	6.5	6.3	6.4	7.4	7.9	8.0	8.7	6.2	6.7	6.2	6.2	§	7.4	+1.3
College Plans:																			
None or under 4 years	—	12.3	11.1	10.8	13.6	13.4	11.2	13.2	11.4	12.7	12.4	14.8	10.8	10.0	10.4	10.6	§	7.6	-3.0
Complete 4 years	—	7.8	7.0	6.9	7.5	7.2	7.6	7.4	8.4	8.8	8.1	9.1	7.6	7.9	7.7	7.1	§	8.0	+0.9
Region:																			
Northeast	—	8.8	9.8	7.8	9.5	7.4	7.7	8.3	8.0	6.7	7.4	7.5	7.5	9.7	8.3	4.5	§	8.0	+3.5
Midwest	—	9.3	6.0	7.8	8.5	8.9	9.6	8.9	9.0	9.1	9.6	11.2	9.1	9.5	9.0	8.1	§	7.1	-1.0
South	—	9.2	9.5	8.5	8.4	10.3	9.1	9.2	10.3	10.8	11.3	12.9	9.4	9.1	9.0	9.3	§	8.6	-0.7
West	—	6.2	5.2	5.5	8.3	4.5	5.8	6.7	8.2	10.2	6.1	5.4	6.5	6.3	7.6	7.5	§	8.1	+0.6
Population Density:																			
Large MSA	—	7.1	8.7	8.5	10.2	7.1	7.2	7.5	9.8	10.6	8.1	9.2	8.8	7.6	8.4	6.2	§	7.3	+1.1
Other MSA	—	9.9	7.1	7.6	8.6	8.1	9.4	9.8	9.1	9.9	9.3	10.4	8.2	9.0	8.6	8.6	§	7.9	-0.7
Non-MSA	—	7.5	7.8	6.3	6.9	9.7	7.0	6.4	7.7	7.6	9.7	9.8	8.3	9.4	8.9	9.1	§	9.5	+0.4
Parental Education: ^a																			
1.0–2.0 (Low)	—	7.7	7.0	5.8	5.4	7.5	8.5	6.9	5.9	10.4	8.6	8.3	6.6	5.9	3.2	6.6	§	3.5	-3.0
2.5–3.0	—	7.1	8.4	6.3	6.5	6.5	7.2	7.1	9.6	8.8	8.1	7.1	7.1	7.9	7.5	6.1	§	7.1	+1.1
3.5–4.0	—	9.0	6.2	7.6	10.3	8.1	8.8	7.8	8.7	9.2	8.5	10.3	7.5	7.7	9.1	7.3	§	7.0	-0.3
4.5–5.0	—	8.1	8.5	7.2	8.3	8.4	8.6	8.7	9.1	9.3	8.6	11.0	9.4	11.7	9.2	8.5	§	12.5	+4.0 s
5.5–6.0 (High)	—	11.3	8.7	11.2	10.2	11.1	8.2	10.7	11.1	12.4	12.2	11.0	11.2	7.8	11.9	8.7	§	7.6	-1.2
Race/Ethnicity (2-year average): ^b																			
White	—	—	9.1	8.5	9.2	9.9	9.5	9.6	10.4	10.7	10.5	11.0	11.1	10.3	10.8	9.8	§	9.1	-0.7
African American	—	—	5.4	4.6	4.6	5.5	6.5	6.2	6.5	6.8	7.1	7.9	7.5	7.6	7.9	8.1	§	7.6	-0.6
Hispanic	—	—	6.0	5.8	5.6	5.1	5.7	6.1	5.4	5.8	5.7	5.1	4.9	5.0	4.5	3.8	§	3.2	-0.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cData from one of the questionnaire forms was omitted in 2005 due to an error in the skip pattern in the questionnaire. For the 2006 results by race/ethnicity, data from the remaining form in 2005 were double-weighted to account for the omission of the other form.

TABLE 166
STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 8

	Percentage who said "Yes, I take them now."																		2019– 2021 change
	1991- 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	3.9	3.5	3.1	3.5	3.7	3.4	3.3	3.5	3.4	3.2	3.6	3.7	3.4	3.7	2.8	2.0	4.2	+1.5* ss
Gender:																			
Male	—	4.7	4.8	4.0	4.5	4.6	3.7	4.3	4.6	4.0	4.0	4.6	4.6	4.0	4.7	3.9	§	5.3	+1.4
Female	—	3.0	2.1	2.2	2.4	3.0	3.1	2.4	2.5	2.8	2.6	2.8	2.7	2.5	2.9	1.6	§	2.5	+0.9
College Plans:																			
None or under 4 years	—	4.9	6.0	3.9	3.6	3.0	5.0	2.3	4.1	5.5	2.6	5.9	6.3	3.0	5.9	3.5	§	6.1	+2.6
Complete 4 years	—	3.7	3.2	3.0	3.4	3.7	3.3	3.4	3.4	3.3	3.3	3.3	3.5	3.4	3.4	2.7	§	4.0	+1.3 s
Region:																			
Northeast	—	4.2	3.0	3.1	3.1	5.0	2.1	2.6	2.9	3.8	2.9	3.6	2.5	3.6	4.1	4.2	§	4.0	-0.1
Midwest	—	4.4	3.1	3.3	2.7	2.7	3.8	4.0	4.1	3.6	3.8	4.1	4.6	4.0	3.9	2.6	§	5.4	+2.8 s
South	—	3.7	4.1	3.3	4.8	4.3	3.9	3.1	4.4	3.9	3.5	3.8	4.8	4.1	4.8	2.5	§	3.5	+1.0
West	—	3.3	3.0	2.5	2.4	2.7	3.0	3.7	1.7	2.1	2.3	2.6	1.6	1.4	1.2	2.4	§	4.5	+2.1
Population Density:																			
Large MSA	—	3.7	2.5	2.7	3.8	4.0	3.0	3.2	3.0	3.4	2.5	3.2	2.5	3.5	3.3	2.2	§	3.9	+1.6
Other MSA	—	4.5	4.1	3.0	3.3	3.8	3.8	3.4	4.0	3.4	3.9	4.1	4.3	3.2	4.6	2.7	§	5.0	+2.3 ss
Non-MSA	—	3.0	3.6	3.9	3.3	3.0	3.0	3.4	3.0	3.7	2.7	2.9	3.9	3.6	2.1	3.7	§	3.2	-0.6
Parental Education: ^a																			
1.0–2.0 (Low)	—	4.3	3.9	2.8	3.1	3.5	2.2	2.0	2.5	1.9	2.0	2.0	1.7	2.2	4.1	0.9	§	3.2	+2.3
2.5–3.0	—	3.2	3.7	3.1	2.2	3.2	2.8	3.0	3.6	2.6	2.8	3.1	2.8	3.1	1.5	3.0	§	4.6	+1.6
3.5–4.0	—	3.7	2.7	2.5	2.5	3.3	3.3	2.9	3.1	3.3	3.6	3.3	4.2	2.9	2.6	2.8	§	5.2	+2.5
4.5–5.0	—	3.9	2.9	3.2	4.2	3.6	3.3	3.9	3.6	4.0	3.8	4.0	3.9	4.1	5.5	3.5	§	3.8	+0.3
5.5–6.0 (High)	—	4.7	4.0	3.9	5.8	5.3	4.1	3.5	3.5	3.8	3.1	4.4	3.9	4.5	4.3	2.9	§	4.6	+1.7
Race/Ethnicity (2-year average): ^b																			
White	—	—	4.2	3.7	4.0	4.6	4.5	4.1	4.3	4.3	4.2	4.5	5.2	5.2	5.2	4.6	§	4.0	-0.6
African American	—	—	2.1	2.2	2.5	2.7	1.8	1.9	2.4	2.6	2.7	2.7	2.0	1.9	2.0	0.9	§	0.2	-0.7 s
Hispanic	—	—	2.5	1.9	1.3	1.5	1.7	2.0	1.7	1.3	1.4	1.6	1.6	1.0	0.9	1.6	§	1.5	-0.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; N is one third of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 167
STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 10

	Percentage who said "Yes, I take them now."																		2019– 2021 change	
	1991– 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
<i>Approximate weighted N =</i>	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	3.4	2.8	2.8	2.9	3.3	3.1	2.8	3.8	3.7	3.4	4.2	3.0	3.0	3.9	2.9	2.5	3.6	+0.7*	
Gender:																				
Male	—	4.1	3.5	3.2	3.4	3.6	4.1	3.0	4.4	4.8	4.2	5.0	3.1	3.5	4.7	3.1	§	3.2	+0.1	
Female	—	2.7	2.1	2.5	2.4	3.1	2.1	2.7	3.4	2.5	2.7	3.3	2.8	2.5	3.1	2.7	§	3.8	+1.1	
College Plans:																				
None or under 4 years	—	4.8	4.1	3.5	3.2	6.9	4.7	3.3	2.4	5.0	6.1	6.6	4.4	4.1	4.8	4.3	§	5.1	+0.8	
Complete 4 years	—	3.2	2.6	2.7	2.9	3.0	2.9	2.8	4.0	3.6	3.1	3.9	2.9	3.0	3.7	2.7	§	3.3	+0.6	
Region:																				
Northeast	—	3.3	1.9	2.4	2.9	2.1	2.7	2.9	3.3	5.8	3.3	4.0	2.9	2.5	2.6	2.0	§	4.5	+2.5	
Midwest	—	3.5	2.9	3.1	3.9	5.2	3.3	3.1	4.1	4.5	3.7	4.6	3.4	2.3	5.2	3.3	§	4.0	+0.7	
South	—	4.2	3.4	3.5	2.9	4.1	3.6	3.5	5.3	3.3	4.3	5.0	3.7	4.1	4.5	2.4	§	3.3	+0.9	
West	—	2.0	2.5	2.0	2.1	1.4	2.5	1.4	1.8	2.0	2.0	2.4	1.8	2.7	2.4	3.9	§	2.9	-1.0	
Population Density:																				
Large MSA	—	2.9	2.5	2.2	3.2	3.4	3.3	3.2	3.4	4.6	3.7	4.0	2.6	2.8	3.8	1.6	§	3.6	+2.1 sss	
Other MSA	—	3.9	3.0	3.1	2.7	3.2	2.9	2.5	4.1	3.4	3.1	4.7	2.5	3.5	4.1	3.9	§	3.8	-0.1	
Non-MSA	—	3.1	2.6	3.1	3.2	3.4	3.2	3.0	3.9	3.3	3.5	3.1	5.3	2.2	3.5	2.5	§	2.9	+0.4	
Parental Education: ^a																				
1.0–2.0 (Low)	—	2.6	3.6	1.8	1.7	1.5	1.4	1.8	1.9	1.4	1.1	0.9	1.0	1.7	2.3	1.5	§	2.0	+0.5	
2.5–3.0	—	2.7	1.7	2.1	3.1	2.7	3.2	3.1	4.0	3.1	2.1	2.8	1.3	2.8	2.7	2.4	§	3.5	+1.1	
3.5–4.0	—	3.7	3.4	3.1	2.3	2.9	3.5	1.8	4.3	3.2	3.1	4.4	2.5	3.3	3.1	2.9	§	3.4	+0.5	
4.5–5.0	—	2.9	2.6	2.9	2.8	3.6	2.9	3.6	3.6	4.5	3.4	4.2	4.3	3.1	5.3	3.4	§	3.5	+0.1	
5.5–6.0 (High)	—	5.1	3.1	3.3	4.2	5.5	4.6	3.4	4.1	6.1	6.2	6.0	3.9	3.7	5.5	4.3	§	4.7	+0.3	
Race/Ethnicity (2-year average): ^b																				
White	—	—	3.8	3.3	3.5	3.7	3.8	3.6	4.0	4.7	4.7	5.1	5.1	4.1	4.7	4.8	§	4.2	-0.5	
African American	—	—	1.2	1.9	1.8	1.3	1.2	1.9	2.8	2.2	1.0	1.4	1.4	1.2	1.5	1.9	§	1.2	-0.7	
Hispanic	—	—	1.3	1.3	1.1	1.6	1.6	1.3	1.7	1.4	0.9	1.0	0.9	1.5	2.1	1.1	§	0.7	-0.4	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 168
STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who said "Yes, I take them now."																		2019– 2021 change	
	1991– 2004	2005 ^c	2006 ^c	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	2.9	2.3	2.6	2.9	2.9	3.0	3.3	3.8	4.4	3.8	4.0	3.9	3.4	3.5	3.2	3.1	3.4	+0.2*	
Gender:																				
Male	—	4.1	2.6	3.7	3.0	2.6	3.6	3.5	3.9	4.5	3.4	4.4	4.7	3.6	3.9	4.2	§	3.0	-1.2	
Female	—	1.8	2.0	2.0	2.6	3.2	2.5	3.2	3.8	4.2	4.0	2.9	3.0	3.3	3.1	2.4	§	3.5	+1.1	
College Plans:																				
None or under 4 years	—	5.3	2.4	3.6	3.3	1.8	3.9	3.9	3.4	5.1	3.4	4.0	2.9	2.9	3.3	3.2	§	2.2	-1.0	
Complete 4 years	—	2.5	2.3	2.4	2.9	3.0	2.9	3.2	3.8	4.2	3.8	4.0	3.8	3.7	3.3	3.3	§	3.7	+0.4	
Region:																				
Northeast	—	3.5	3.0	2.7	3.2	2.0	3.2	3.3	3.1	3.1	3.6	2.5	3.3	4.8	3.2	1.7	§	3.4	+1.7	
Midwest	—	2.7	2.0	3.3	2.9	2.7	4.0	4.6	5.0	3.7	4.5	5.0	3.8	4.0	3.8	4.2	§	3.2	-1.0	
South	—	2.7	2.6	2.8	2.8	4.3	3.3	3.2	3.7	5.4	4.8	5.2	4.1	3.1	3.8	3.3	§	3.0	-0.3	
West	—	3.0	1.6	1.2	2.9	1.6	1.4	2.3	3.1	4.5	1.6	2.1	4.0	2.6	2.7	3.2	§	4.6	+1.4	
Population Density:																				
Large MSA	—	2.3	2.2	2.9	3.7	2.9	2.7	3.0	3.9	5.0	3.5	3.9	4.7	3.6	3.3	2.5	§	2.7	+0.2	
Other MSA	—	3.5	2.4	2.4	2.9	2.7	3.5	4.5	4.5	4.9	3.9	4.2	4.1	3.5	3.4	3.6	§	3.6	+0.1	
Non-MSA	—	2.3	2.3	2.4	2.1	3.1	2.3	1.2	2.0	2.3	3.9	3.3	2.1	3.1	3.9	3.7	§	4.2	+0.6	
Parental Education: ^a																				
1.0–2.0 (Low)	—	2.2	1.0	1.4	1.9	1.3	2.6	2.8	2.5	3.7	1.9	2.0	2.7	1.0	1.0	0.8	§	2.2	+1.4	
2.5–3.0	—	2.0	1.4	2.4	1.9	1.6	2.3	2.2	3.2	4.1	2.9	1.9	2.0	3.0	2.6	2.0	§	2.7	+0.7	
3.5–4.0	—	2.9	2.1	2.4	2.8	2.8	3.2	3.6	3.2	3.9	2.8	3.1	3.4	2.5	3.4	3.2	§	1.8	-1.4	
4.5–5.0	—	3.3	3.2	2.1	3.6	3.0	2.9	3.6	4.2	4.3	4.2	5.2	5.1	5.8	3.9	3.7	§	6.6	+3.0	
5.5–6.0 (High)	—	3.8	3.0	4.3	3.8	5.9	4.7	4.8	6.5	6.9	8.0	6.9	5.2	4.4	6.9	4.9	§	3.7	-1.2	
Race/Ethnicity (2-year average): ^b																				
White	—	—	3.1	2.9	3.3	3.5	3.7	4.3	4.8	4.8	4.7	4.9	5.2	4.8	4.6	4.4	§	4.5	+0.1	
African American	—	—	0.6	1.1	1.2	1.3	1.5	1.3	1.8	2.8	2.3	1.5	1.4	2.4	2.8	2.4	§	2.8	+0.4	
Hispanic	—	—	1.1	0.9	1.2	1.6	1.8	1.7	1.5	1.7	2.0	1.4	1.4	2.1	1.6	1.1	§	0.5	-0.6	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cData from one of the questionnaire forms was omitted in 2005 due to an error in the skip pattern in the questionnaire. For the 2006 results by race/ethnicity, data from the remaining form in 2005 were double-weighted to account for the omission of the other form.

TABLE 169
NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Lifetime Prevalence of Legal Use by Subgroups in Grade 8

	Percentage who used in lifetime																		2019– 2021 change	
	1991– 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
<i>Approximate weighted N =</i>	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	7.3	7.9	6.3	6.3	5.8	5.8	6.1	5.1	5.1	4.8	5.1	5.7	4.9	4.4	4.5	4.2	2.8	-1.7* ss	
Gender:																				
Male	—	7.6	8.7	6.7	6.8	6.5	6.6	6.7	5.2	5.7	6.1	5.7	6.7	5.3	5.1	5.6	§	3.3	-2.2 s	
Female	—	7.0	6.8	5.9	5.7	5.1	4.9	5.4	4.9	4.4	3.6	4.6	4.8	4.2	3.6	3.5	§	2.1	-1.4	
College Plans:																				
None or under 4 years	—	11.1	12.9	8.1	7.3	8.9	8.8	11.7	8.8	11.5	7.7	7.8	10.0	7.6	6.3	10.2	§	3.5	-6.7 sss	
Complete 4 years	—	6.9	7.4	6.0	6.2	5.5	5.4	5.7	4.9	4.6	4.6	4.8	5.3	4.6	4.2	3.8	§	2.6	-1.2 s	
Region:																				
Northeast	—	7.4	6.0	4.3	5.1	6.1	4.6	5.0	4.2	4.7	5.6	5.2	7.1	4.7	5.5	4.8	§	2.8	-2.0	
Midwest	—	6.9	8.8	7.6	6.7	5.3	6.2	5.5	5.5	3.8	4.7	4.8	5.4	5.3	4.4	4.1	§	2.6	-1.5	
South	—	7.9	7.5	6.6	7.2	5.8	6.4	7.0	5.7	5.6	5.1	4.8	5.2	5.3	4.4	5.6	§	3.8	-1.9 s	
West	—	6.5	9.3	5.3	5.3	6.0	5.0	6.2	4.1	6.0	4.0	5.9	5.8	3.9	3.5	3.0	§	1.4	-1.6	
Population Density:																				
Large MSA	—	5.3	7.8	4.5	6.0	4.0	5.3	5.6	4.4	4.3	4.3	4.6	6.0	4.2	3.9	4.5	§	2.1	-2.5 s	
Other MSA	—	8.0	7.8	7.7	6.7	6.8	6.6	6.7	4.7	5.3	5.4	4.9	5.3	5.7	4.7	5.0	§	3.4	-1.6	
Non-MSA	—	8.6	8.4	5.9	5.9	5.8	4.4	5.7	6.9	5.9	4.3	6.2	6.0	4.0	4.4	3.3	§	2.4	-0.9	
Parental Education: ^a																				
1.0–2.0 (Low)	—	8.7	8.1	4.5	7.0	7.7	7.1	9.1	8.0	5.4	3.4	6.2	4.6	3.6	4.2	4.1	§	3.4	-0.7	
2.5–3.0	—	9.6	6.7	7.4	5.5	6.0	5.8	6.2	4.4	4.4	4.1	4.9	6.0	6.0	4.1	4.8	§	1.7	-3.1 s	
3.5–4.0	—	8.1	10.1	6.5	6.4	5.5	5.4	7.1	5.9	6.4	5.9	4.7	5.3	4.6	5.8	4.7	§	3.7	-1.0	
4.5–5.0	—	5.5	8.0	5.6	6.9	5.0	5.2	5.3	5.0	4.1	4.8	5.2	6.4	6.2	4.6	4.9	§	3.1	-1.8	
5.5–6.0 (High)	—	6.6	5.4	6.6	6.3	7.1	5.9	5.4	4.6	5.5	5.1	4.7	6.2	3.4	3.7	4.1	§	2.8	-1.3	
Race/Ethnicity (2-year average): ^b																				
White	—	—	7.7	7.6	7.3	6.7	5.8	5.6	5.5	5.0	5.2	5.6	6.6	6.4	5.0	4.7	§	4.9	+0.2	
African American	—	—	6.0	5.0	4.3	4.8	4.9	5.6	4.8	4.0	4.1	3.6	3.7	4.9	5.2	4.6	§	2.0	-2.5 s	
Hispanic	—	—	6.5	5.3	3.7	4.7	6.0	6.6	6.0	5.1	4.7	4.1	4.2	3.8	3.1	3.2	§	1.9	-1.3	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 170
NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Lifetime Prevalence of Legal Use by Subgroups in Grade 10

	Percentage who used in lifetime																			2019– 2021 change
	1991– 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
<i>Approximate weighted N =</i>	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	8.3	8.3	6.7	6.8	6.8	6.1	6.4	5.2	4.9	5.8	5.8	5.2	4.6	5.1	5.2	5.1	3.0	-2.2* sss	
Gender:																				
Male	—	9.5	9.2	8.0	7.4	8.9	6.6	6.7	6.4	5.7	7.1	6.4	5.8	5.6	6.1	6.4	§	2.8	-3.7 sss	
Female	—	7.0	7.5	5.2	6.2	4.8	5.5	6.2	4.1	4.0	4.5	5.1	4.7	3.7	4.0	3.8	§	2.9	-0.9	
College Plans:																				
None or under 4 years	—	12.6	10.1	9.5	10.5	14.0	10.1	10.6	11.2	7.1	8.7	12.1	6.4	6.2	7.9	7.9	§	5.4	-2.5	
Complete 4 years	—	7.7	8.0	6.3	6.3	6.1	5.6	5.9	4.5	4.7	5.4	5.1	5.0	4.5	4.5	4.5	§	2.5	-2.1 sss	
Region:																				
Northeast	—	8.6	8.8	6.5	6.1	7.0	5.7	7.2	5.3	4.7	5.2	4.9	6.1	5.3	4.1	5.6	§	5.4	-0.2	
Midwest	—	6.9	7.4	6.8	7.7	7.7	5.9	6.7	3.8	4.8	6.8	5.4	4.6	3.8	5.7	5.3	§	2.7	-2.6 s	
South	—	10.6	9.6	7.2	8.2	7.2	7.0	6.6	6.1	5.0	6.1	6.9	5.5	4.4	5.6	3.9	§	2.3	-1.6 s	
West	—	6.6	7.0	6.1	4.6	5.4	5.3	5.1	5.2	5.1	4.7	5.1	4.8	5.4	4.6	6.9	§	2.3	-4.6 s	
Population Density:																				
Large MSA	—	6.0	7.2	6.1	4.3	6.2	6.1	5.1	5.3	5.5	5.0	5.4	5.0	4.5	4.2	4.1	§	2.6	-1.5	
Other MSA	—	9.2	8.4	6.5	7.1	7.5	5.7	6.8	4.9	4.8	5.6	5.4	5.6	5.0	4.9	5.6	§	3.6	-2.0	
Non-MSA	—	9.3	9.7	8.2	9.6	6.4	6.8	7.4	5.6	4.4	7.6	7.3	4.6	3.9	6.9	5.8	§	2.1	-3.7 ss	
Parental Education: ^a																				
1.0–2.0 (Low)	—	8.2	10.5	6.6	5.1	7.0	6.2	6.3	7.5	5.6	4.7	7.5	2.8	5.9	4.9	2.0	§	1.9	-0.2	
2.5–3.0	—	8.7	7.6	6.0	8.3	6.9	6.5	7.7	4.3	4.3	5.7	5.3	4.9	2.2	4.2	5.4	§	3.2	-2.1	
3.5–4.0	—	7.4	9.7	7.4	7.1	5.3	6.4	5.6	5.3	5.8	7.1	5.1	5.6	4.7	5.8	4.3	§	2.9	-1.3	
4.5–5.0	—	9.1	7.4	6.3	7.4	7.2	5.5	5.9	5.9	4.4	4.5	6.1	5.5	5.2	4.3	5.9	§	3.6	-2.3	
5.5–6.0 (High)	—	8.6	8.3	6.3	5.1	7.8	6.4	6.8	4.0	5.7	6.3	6.1	5.5	5.4	6.3	7.5	§	3.2	-4.3	
Race/Ethnicity (2-year average): ^b																				
White	—	—	9.4	8.3	7.4	7.4	6.9	6.8	6.2	5.4	5.9	6.3	5.9	5.1	5.7	6.6	§	4.2	-2.5 ss	
African American	—	—	4.7	5.0	4.2	4.6	6.1	4.9	3.2	4.0	5.3	5.1	4.3	5.0	3.6	2.0	§	4.9	+2.9 s	
Hispanic	—	—	4.9	4.5	5.0	5.7	4.8	5.6	6.4	4.3	3.5	4.7	5.4	4.9	4.4	3.8	§	2.8	-0.9	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 171
NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Lifetime Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who used in lifetime																		2019– 2021 change
	1991– 2004	2005 ^c	2006 ^c	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	6.2	6.1	7.0	6.4	5.4	6.7	5.8	5.9	5.4	5.6	5.6	5.8	6.4	6.1	5.7	4.8	4.5	-1.2*
Gender:																			
Male	—	7.0	6.1	8.3	7.0	6.5	7.7	6.4	5.8	7.1	6.0	6.1	6.8	7.6	6.4	6.2	§	3.6	-2.6 s
Female	—	5.0	5.7	5.6	6.5	4.0	5.6	5.2	5.6	3.8	4.8	5.1	4.8	5.3	4.7	4.8	§	4.6	-0.2
College Plans:																			
None or under 4 years	—	9.2	7.6	9.4	9.9	6.7	9.0	8.0	8.1	6.3	7.4	7.3	7.3	9.0	6.4	7.9	§	4.4	-3.6 s
Complete 4 years	—	5.3	5.8	6.4	5.5	5.0	6.1	5.2	5.4	4.9	5.0	5.4	5.4	5.7	5.9	5.0	§	4.6	-0.5
Region:																			
Northeast	—	6.0	5.9	7.6	8.0	5.1	4.9	5.4	6.5	5.5	6.5	5.8	6.8	7.7	6.6	4.0	§	4.7	+0.7
Midwest	—	5.8	5.5	7.1	6.8	6.1	8.4	5.0	6.3	5.6	6.9	6.4	6.7	7.2	6.5	6.6	§	3.3	-3.2 s
South	—	7.6	7.4	7.6	5.2	5.7	7.0	6.8	5.8	5.9	6.2	6.9	5.7	7.0	6.3	5.6	§	4.7	-0.9
West	—	4.5	4.4	5.1	6.7	4.4	5.7	5.6	5.3	4.3	3.0	2.8	4.6	4.2	5.0	6.1	§	5.2	-0.9
Population Density:																			
Large MSA	—	5.1	6.1	6.9	6.6	5.6	7.3	5.4	6.1	5.3	6.7	4.5	6.3	6.3	5.8	5.6	§	4.5	-1.1
Other MSA	—	7.9	6.2	7.3	6.3	5.0	6.2	6.3	6.2	4.7	5.5	6.2	5.4	6.8	6.5	5.4	§	4.4	-1.0
Non-MSA	—	3.8	5.8	6.3	6.4	6.2	7.2	5.5	4.9	7.0	4.6	6.2	6.2	5.8	5.8	6.3	§	4.6	-1.7
Parental Education: ^a																			
1.0–2.0 (Low)	—	5.2	4.3	5.5	6.6	6.0	5.1	4.9	5.8	4.7	6.2	5.0	5.7	6.0	3.8	8.3	§	1.6	-6.7 sss
2.5–3.0	—	4.1	6.2	5.7	6.4	3.4	6.1	4.5	4.8	5.2	4.5	5.4	3.8	7.7	4.1	4.7	§	2.2	-2.5 s
3.5–4.0	—	7.6	5.9	6.9	7.0	5.2	6.6	5.9	5.6	4.8	5.0	4.1	6.0	5.8	5.3	4.9	§	4.2	-0.7
4.5–5.0	—	6.8	6.4	8.1	4.7	6.5	7.9	6.5	6.1	6.2	5.7	6.5	6.4	6.3	8.1	5.2	§	8.2	+3.1
5.5–6.0 (High)	—	6.4	6.8	7.8	7.5	6.4	6.6	6.8	7.0	6.3	8.5	7.4	6.3	6.7	9.8	6.8	§	4.3	-2.4
Race/Ethnicity (2-year average): ^b																			
White	—	—	6.6	7.3	7.3	6.3	6.7	6.9	6.1	5.9	6.1	6.5	6.9	7.1	7.3	6.5	§	5.5	-1.0
African American	—	—	4.9	4.9	5.0	4.7	5.8	5.5	5.9	5.5	5.6	5.5	4.5	6.3	7.7	9.1	§	4.2	-4.9 ss
Hispanic	—	—	4.2	4.0	4.7	4.5	4.5	5.2	5.1	4.3	3.7	3.8	3.9	4.4	3.3	2.6	§	1.6	-1.0 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cData from one of the questionnaire forms was omitted in 2005 due to an error in the skip pattern in the questionnaire. For the 2006 results by race/ethnicity, data from the remaining form in 2005 were double-weighted to account for the omission of the other form.

TABLE 172
NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 8

	Percentage who said "Yes, I take them now."																		2019– 2021 change
	1991– 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	2.2	1.9	1.4	1.6	1.2	1.4	1.5	1.2	1.4	1.2	1.2	2.0	1.1	1.2	1.4	1.4	0.9	-0.5*
Gender:																			
Male	—	2.4	2.3	1.6	1.7	1.2	1.5	1.9	1.1	1.6	2.0	1.2	2.0	1.1	1.6	2.0	§	0.9	-1.1 s
Female	—	1.9	1.5	1.3	1.5	1.1	1.3	0.9	1.1	1.0	0.6	1.0	1.8	1.0	0.9	0.9	§	0.7	-0.2
College Plans:																			
None or under 4 years	—	2.5	3.8	1.3	3.1	1.6	2.1	3.9	1.6	4.2	1.1	3.4	4.0	2.8	3.5	2.8	§	1.0	-1.8 s
Complete 4 years	—	2.1	1.7	1.4	1.5	1.1	1.3	1.2	1.2	1.2	1.3	0.9	1.6	1.0	0.9	1.3	§	0.8	-0.5
Region:																			
Northeast	—	2.4	2.5	1.1	1.8	1.0	1.1	1.1	1.5	1.3	0.6	0.8	1.8	1.3	1.4	1.0	§	0.5	-0.6
Midwest	—	2.0	2.7	1.9	1.7	0.7	1.8	1.7	1.3	0.7	1.5	1.6	1.4	1.2	1.0	1.2	§	0.4	-0.8
South	—	2.2	1.6	1.5	1.8	1.3	1.4	1.4	1.2	1.8	1.3	1.1	2.4	1.3	1.6	1.6	§	1.6	-0.1
West	—	2.0	1.3	0.8	1.2	1.5	1.0	1.4	1.0	1.7	1.2	1.2	1.9	0.6	0.6	1.5	§	0.6	-0.9
Population Density:																			
Large MSA	—	1.4	1.7	0.9	1.9	1.0	1.2	1.3	1.3	1.4	1.1	0.6	2.2	1.2	1.0	0.9	§	0.2	-0.7
Other MSA	—	2.3	2.1	1.9	1.5	1.4	1.6	1.7	1.2	1.5	1.4	1.2	1.8	1.0	1.3	1.5	§	1.4	-0.1
Non-MSA	—	2.9	1.9	1.1	1.6	0.9	1.1	1.2	0.9	1.2	1.0	1.9	2.0	1.1	1.3	2.0	§	0.9	-1.1
Parental Education: ^a																			
1.0–2.0 (Low)	—	2.8	1.2	1.2	1.3	2.1	1.1	1.7	0.4	0.7	0.9	2.4	0.7	0.7	0.4	1.6	§	0.4	-1.1
2.5–3.0	—	2.4	1.4	1.1	0.9	0.5	0.7	1.4	0.8	1.0	0.5	1.1	1.9	1.1	0.7	0.8	§	0.2	-0.6
3.5–4.0	—	2.3	2.8	0.9	1.9	0.4	1.2	1.7	1.0	1.9	2.4	0.8	1.4	1.2	1.6	2.0	§	1.3	-0.7
4.5–5.0	—	1.8	1.8	1.8	1.6	1.9	1.6	1.3	2.0	1.3	1.2	1.3	1.7	1.7	2.1	1.3	§	0.9	-0.4
5.5–6.0 (High)	—	2.3	1.5	2.2	2.3	1.2	1.6	1.4	1.4	2.0	0.9	0.3	3.3	0.7	0.5	1.8	§	1.0	-0.8
Race/Ethnicity (2-year average): ^b																			
White	—	—	2.3	2.0	1.9	1.6	1.4	1.6	1.5	1.2	1.2	1.4	1.9	1.9	1.5	1.6	§	1.8	+0.1
African American	—	—	1.3	0.9	0.9	1.1	1.5	1.5	1.0	1.4	1.3	0.7	1.0	1.7	1.7	0.9	§	0.6	-0.4
Hispanic	—	—	1.0	0.8	0.7	0.6	0.5	1.0	1.0	0.9	1.2	0.8	1.1	1.2	0.5	0.5	§	0.5	-0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 173
NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 10

	Percentage who said "Yes, I take them now."																		2019– 2021 change	
	1991- 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
<i>Approximate weighted N =</i>	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	2.3	2.3	1.6	1.7	1.9	1.6	1.3	1.3	1.3	1.4	1.7	1.2	1.0	1.4	1.8	1.8	1.5	-0.2*	
Gender:																				
Male	—	2.6	2.7	1.8	1.9	2.2	2.1	1.3	1.6	1.2	1.9	1.8	1.4	0.9	1.3	1.9	§	1.4	-0.5	
Female	—	1.9	2.0	1.4	1.4	1.5	1.2	1.3	0.9	1.2	1.0	1.6	1.0	1.0	1.4	1.6	§	1.5	-0.1	
College Plans:																				
None or under 4 years	—	4.4	2.6	2.5	1.8	4.4	3.5	2.8	2.3	2.5	1.0	3.2	1.8	1.5	1.7	2.1	§	2.9	+0.7	
Complete 4 years	—	2.0	2.2	1.4	1.6	1.7	1.4	1.0	1.2	1.2	1.4	1.6	1.1	0.9	1.2	1.7	§	1.3	-0.4	
Region:																				
Northeast	—	3.1	2.6	1.6	1.6	2.9	1.4	1.0	1.4	2.0	1.2	1.6	1.2	1.2	1.2	2.0	§	2.7	+0.7	
Midwest	—	2.2	2.2	2.3	2.6	2.5	1.2	1.3	0.7	1.1	1.7	2.4	1.3	1.4	1.5	1.6	§	1.2	-0.4	
South	—	2.2	2.5	1.4	1.9	1.4	2.4	1.3	1.7	1.1	1.9	1.5	1.2	0.7	1.6	1.1	§	1.1	-0.1	
West	—	1.8	2.0	1.0	0.7	1.2	1.1	1.4	1.1	1.3	0.5	1.4	1.1	1.0	1.3	2.9	§	1.5	-1.4	
Population Density:																				
Large MSA	—	1.5	1.9	1.5	0.9	2.3	2.1	1.2	1.3	1.5	1.2	1.6	1.1	0.9	1.3	1.4	§	1.1	-0.3	
Other MSA	—	2.7	2.7	1.6	1.9	2.0	1.2	1.0	1.3	1.5	1.5	1.7	1.3	1.1	1.4	2.1	§	2.1	-0.1	
Non-MSA	—	2.3	2.2	1.7	2.2	1.2	1.8	1.9	1.2	0.8	1.6	2.0	1.3	1.1	1.8	1.5	§	1.0	-0.5	
Parental Education: ^a																				
1.0–2.0 (Low)	—	1.4	3.2	2.5	1.2	1.3	1.5	1.5	0.5	1.4	0.7	0.3	1.0	1.8	1.2	0.7	§	0.6	-0.2	
2.5–3.0	—	1.6	1.5	0.6	1.8	1.9	1.6	1.4	1.0	0.8	0.6	1.1	0.8	0.3	1.1	1.9	§	2.0	0.0	
3.5–4.0	—	2.5	2.5	1.5	1.4	1.0	2.0	1.2	0.9	2.2	2.0	1.6	0.8	1.3	1.2	1.0	§	1.0	0.0	
4.5–5.0	—	2.9	2.6	1.7	2.2	2.7	1.4	1.3	1.8	0.9	1.5	2.2	1.3	0.7	1.4	1.8	§	1.7	-0.1	
5.5–6.0 (High)	—	2.2	2.5	2.0	1.6	2.8	1.6	1.0	1.9	1.4	1.4	2.3	1.9	1.3	2.1	3.2	§	2.3	-0.9	
Race/Ethnicity (2-year average): ^b																				
White	—	—	2.6	2.3	1.9	2.0	2.0	1.6	1.3	1.3	1.5	2.1	2.0	1.3	1.4	2.0	§	2.1	+0.1	
African American	—	—	1.7	1.2	0.7	0.7	1.0	1.0	0.8	1.6	1.7	0.9	0.5	0.6	0.7	0.5	§	1.1	+0.6	
Hispanic	—	—	1.3	0.8	1.1	1.6	1.1	1.0	1.3	0.7	0.2	0.5	0.4	0.5	0.7	1.1	§	0.2	-0.9 s	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 174
NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who said "Yes, I take them now."																			2019– 2021 change
	1991– 2004	2005 ^c	2006 ^c	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	1.6	1.6	1.7	1.9	1.5	2.3	1.9	1.8	1.8	2.2	1.5	2.1	2.5	2.6	2.3	1.7	2.3	0.0*	
Gender:																				
Male	—	1.3	1.4	1.3	1.1	1.5	2.3	1.7	1.6	2.4	2.2	1.8	2.5	2.5	2.3	2.7	§	1.3	-1.4 s	
Female	—	1.6	1.8	1.6	2.2	1.6	2.3	1.9	2.0	1.3	2.2	1.0	1.6	2.3	2.1	2.0	§	2.9	+1.0	
College Plans:																				
None or under 4 years	—	2.9	0.8	2.5	2.2	1.5	3.2	3.3	2.7	1.7	3.6	2.0	1.7	3.3	3.3	3.0	§	1.7	-1.2	
Complete 4 years	—	1.2	1.7	1.6	1.9	1.5	2.1	1.5	1.6	1.6	1.9	1.4	2.0	2.3	2.4	2.2	§	2.5	+0.3	
Region:																				
Northeast	—	2.5	1.5	2.4	3.9	1.1	1.9	2.8	1.5	2.2	2.2	1.5	3.2	2.1	2.6	1.4	§	2.8	+1.4	
Midwest	—	1.6	1.5	1.7	2.0	1.9	2.8	1.5	2.5	1.6	2.7	1.8	2.4	3.1	2.8	2.9	§	1.4	-1.5	
South	—	1.5	1.9	1.9	1.1	1.9	2.4	2.3	1.4	2.1	2.3	1.7	1.9	2.7	2.5	1.9	§	1.6	-0.3	
West	—	0.8	1.1	0.8	1.8	0.7	1.9	1.1	1.8	1.3	1.8	0.8	1.3	1.8	2.6	3.5	§	4.0	+0.5	
Population Density:																				
Large MSA	—	1.4	1.9	1.7	2.4	1.8	1.7	1.9	1.8	2.6	2.4	1.2	1.9	2.5	3.0	2.1	§	2.3	+0.1	
Other MSA	—	1.9	1.3	2.1	1.7	1.2	2.5	2.0	2.1	1.3	2.5	1.6	2.4	3.0	2.2	2.4	§	2.4	0.0	
Non-MSA	—	1.0	1.5	1.1	1.9	1.7	2.7	1.6	1.1	1.9	1.5	1.6	1.7	1.4	2.7	2.5	§	2.2	-0.3	
Parental Education: ^a																				
1.0–2.0 (Low)	—	0.3	2.2	1.0	2.2	1.3	3.3	1.8	2.3	2.2	3.8	1.6	2.2	2.1	1.6	3.4	§	0.3	-3.1 ss	
2.5–3.0	—	1.5	0.9	1.5	1.9	0.9	2.2	1.6	1.6	1.8	1.7	0.9	1.8	3.4	1.3	1.8	§	0.6	-1.2	
3.5–4.0	—	2.4	1.3	1.9	1.8	1.4	2.4	1.6	1.8	1.3	1.7	0.9	2.0	2.3	2.5	2.1	§	1.5	-0.5	
4.5–5.0	—	1.7	1.6	1.6	1.5	2.1	2.3	2.2	1.6	1.7	2.4	1.5	1.7	2.5	3.3	2.1	§	4.9	+2.8	
5.5–6.0 (High)	—	0.9	2.8	2.3	2.7	1.2	1.8	1.6	2.1	2.8	2.8	2.6	2.5	2.5	3.9	2.8	§	3.3	+0.4	
Race/Ethnicity (2-year average): ^b																				
White	—	—	1.7	1.7	1.9	1.7	2.1	2.4	1.9	1.7	2.0	2.1	2.2	2.7	3.0	2.8	§	2.5	-0.3	
African American	—	—	2.0	1.8	2.0	1.5	1.5	1.9	1.9	1.8	2.4	2.0	1.1	1.7	2.5	3.0	§	2.4	-0.6	
Hispanic	—	—	1.2	1.3	1.2	1.4	1.7	1.1	1.4	2.0	1.8	1.1	0.9	1.8	1.2	0.7	§	0.3	-0.4	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cData from one of the questionnaire forms was omitted in 2005 due to an error in the skip pattern in the questionnaire. For the 2006 results by race/ethnicity, data from the remaining form in 2005 were double-weighted to account for the omission of the other form.

TABLE 175
EITHER STIMULANT-TYPE OR NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Lifetime Prevalence of Legal Use by Subgroups in Grade 8

	1991- 2004	Percentage who used in lifetime																	2019- 2021 change
		2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	13.7	15.8	13.4	13.1	12.8	12.8	12.4	11.6	11.5	11.2	11.4	12.1	10.9	11.0	9.8	7.3	11.5	+1.8*
Gender:																			
Male	—	14.9	18.0	14.8	14.7	14.5	14.5	13.7	13.3	13.6	12.7	12.8	14.2	12.5	13.1	11.9	§	14.3	+2.4
Female	—	12.4	13.2	12.1	11.5	11.4	11.3	10.9	10.0	9.4	9.9	10.2	10.0	8.6	9.2	7.5	§	8.2	+0.6
College Plans:																			
None or under 4 years	—	20.6	24.2	17.9	17.1	15.8	17.2	18.8	16.7	21.5	17.2	14.2	20.6	15.7	16.5	17.0	§	14.3	-2.7
Complete 4 years	—	12.9	14.9	12.9	12.7	12.6	12.4	11.9	11.3	10.8	10.8	11.1	11.4	10.3	10.2	8.9	§	11.1	+2.2
Region:																			
Northeast	—	12.9	12.0	10.6	13.7	14.3	9.9	11.0	9.7	11.0	11.5	10.1	11.6	10.2	12.5	10.2	§	11.0	+0.9
Midwest	—	14.2	15.3	14.7	12.5	11.6	13.8	12.0	11.4	10.0	11.0	11.8	12.3	12.4	10.6	8.8	§	11.9	+3.0
South	—	15.1	16.5	14.7	15.0	14.2	14.1	13.2	13.9	13.6	11.8	13.0	13.4	12.2	13.1	10.9	§	11.9	+1.0
West	—	11.3	17.8	11.3	10.8	11.0	11.6	12.7	8.9	10.0	10.1	9.0	9.6	7.7	6.6	8.5	§	11.0	+2.4
Population Density:																			
Large MSA	—	11.4	14.1	11.1	12.3	10.1	11.3	11.9	10.6	10.1	9.8	10.3	11.0	10.8	9.9	9.7	§	11.0	+1.3
Other MSA	—	15.7	16.2	14.7	14.3	14.2	14.3	13.2	11.7	11.4	12.5	11.5	12.6	11.1	12.8	9.8	§	12.8	+3.0
Non-MSA	—	13.1	17.6	13.9	11.6	13.2	11.4	11.7	13.1	13.7	10.3	12.6	12.3	10.4	8.7	9.9	§	9.7	-0.2
Parental Education: ^a																			
1.0–2.0 (Low)	—	18.8	17.8	12.6	17.5	17.0	13.9	15.3	12.5	8.6	7.8	12.7	10.9	7.4	11.5	8.7	§	9.5	+0.8
2.5–3.0	—	14.3	16.3	15.6	11.9	12.9	13.9	13.0	12.4	9.4	10.6	11.1	14.2	12.3	9.3	11.6	§	11.5	-0.1
3.5–4.0	—	15.7	16.2	14.2	12.6	12.6	11.7	14.0	12.1	13.4	11.5	11.4	10.9	9.8	11.4	10.0	§	13.5	+3.6
4.5–5.0	—	11.3	15.5	11.7	13.3	12.0	12.5	11.3	12.0	11.5	11.9	12.2	12.6	12.0	12.9	10.2	§	11.0	+0.8
5.5–6.0 (High)	—	13.0	13.4	13.5	14.8	13.7	13.7	11.5	9.6	11.8	11.1	10.8	11.7	11.5	10.7	7.9	§	12.3	+4.4 s
Race/Ethnicity (2-year average): ^b																			
White	—	—	15.2	15.7	15.0	14.4	13.8	13.1	12.7	12.3	12.7	13.4	14.4	13.9	12.8	12.3	§	11.8	-0.5
African American	—	—	11.0	9.5	9.8	11.3	11.5	10.3	9.4	10.1	9.6	8.8	9.2	10.9	11.2	8.0	§	4.0	-4.1 ss
Hispanic	—	—	12.4	11.7	9.7	10.4	11.2	12.2	10.7	8.2	8.5	8.2	8.2	7.4	6.7	7.1	§	4.8	-2.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 177
EITHER STIMULANT-TYPE OR NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Lifetime Prevalence of Legal Use by Subgroups in Grade 12

	1991- 2004	Percentage who used in lifetime																		2019- 2021 change
		2005 ^c	2006 ^c	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	12.4	11.7	12.1	13.1	11.0	12.7	12.2	12.7	13.2	12.6	13.7	12.7	13.0	12.7	11.1	9.9	10.9	-0.2*	
Gender:																				
Male	—	15.8	13.2	15.7	14.6	13.0	14.7	14.3	14.4	15.5	13.5	13.4	14.5	15.3	14.6	13.0	§	11.4	-1.5	
Female	—	8.8	9.6	8.6	11.5	9.1	10.1	9.9	10.8	10.6	11.0	12.6	10.2	10.2	9.7	8.8	§	9.8	+0.9	
College Plans:																				
None or under 4 years	—	16.7	16.3	15.7	18.8	17.3	16.1	17.4	16.6	16.3	16.9	18.9	15.5	15.1	14.8	14.4	§	9.9	-4.6	
Complete 4 years	—	11.3	10.6	11.2	11.2	10.6	11.8	11.0	11.8	12.2	11.4	13.0	11.6	12.1	11.8	10.1	§	11.1	+0.9	
Region:																				
Northeast	—	13.1	13.8	12.2	14.2	11.2	11.1	12.0	13.1	10.5	12.0	11.5	13.1	15.4	12.5	7.9	§	11.2	+3.3	
Midwest	—	12.7	9.7	12.5	12.0	12.9	14.7	12.3	12.9	12.5	13.7	15.5	13.4	14.1	13.3	12.4	§	9.7	-2.7	
South	—	13.9	14.0	13.3	12.3	14.0	13.6	13.0	13.0	14.8	14.7	17.3	13.7	13.7	12.9	12.1	§	11.9	-0.2	
West	—	8.6	8.1	9.2	12.5	7.2	10.3	10.9	12.0	13.2	8.5	7.9	9.7	9.4	11.8	10.0	§	10.2	+0.2	
Population Density:																				
Large MSA	—	10.4	12.6	12.4	14.5	10.9	12.4	10.6	13.4	13.9	12.3	12.4	13.6	12.0	12.2	10.0	§	10.5	+0.4	
Other MSA	—	14.6	11.1	12.6	12.5	11.2	13.3	14.1	13.2	13.3	12.6	14.5	12.0	13.4	13.0	11.3	§	10.6	-0.7	
Non-MSA	—	9.9	11.6	10.5	10.6	14.2	11.8	9.8	10.5	11.8	12.9	14.0	12.9	13.5	12.9	12.4	§	12.3	-0.1	
Parental Education: ^a																				
1.0–2.0 (Low)	—	10.1	10.3	9.5	10.7	11.1	11.5	10.4	9.7	13.1	12.1	11.1	11.9	10.2	6.3	11.0	§	4.0	-7.0 ss	
2.5–3.0	—	10.0	12.5	10.8	10.1	8.9	10.5	10.0	12.5	12.2	11.6	11.5	9.6	13.4	10.5	8.9	§	9.0	+0.2	
3.5–4.0	—	13.6	10.4	12.2	14.4	11.8	13.1	11.7	12.5	12.4	11.4	13.2	11.9	11.4	12.4	10.4	§	10.1	-0.3	
4.5–5.0	—	12.1	12.3	12.2	11.2	12.8	14.4	13.0	12.8	13.4	12.5	15.4	13.8	15.7	14.7	11.3	§	17.1	+5.8 s	
5.5–6.0 (High)	—	15.3	12.2	15.2	15.2	14.9	13.4	14.9	14.8	17.0	17.5	16.0	15.8	13.7	19.0	12.9	§	10.1	-2.8	
Race/Ethnicity (2-year average): ^b																				
White	—	—	13.0	13.1	13.6	13.8	14.0	14.4	14.5	14.5	14.3	15.3	15.8	15.3	15.8	14.0	§	12.3	-1.7	
African American	—	—	8.3	8.0	8.0	8.8	10.4	9.7	9.7	10.1	10.6	11.1	11.0	11.7	11.7	13.0	§	9.6	-3.4	
Hispanic	—	—	9.0	8.7	8.9	8.0	8.1	8.9	8.2	8.4	8.4	8.1	8.0	8.4	7.2	5.8	§	4.5	-1.3	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cData from one of the questionnaire forms was omitted in 2005 due to an error in the skip pattern in the questionnaire. For the 2006 results by race/ethnicity, data from the remaining form in 2005 were double-weighted to account for the omission of the other form.

TABLE 178
EITHER STIMULANT-TYPE OR NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 8

	Percentage who said "Yes, I take them now."																		2019– 2021 change	
	1991- 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	6.1	5.2	4.5	5.1	4.9	4.7	4.9	4.7	5.0	4.6	4.9	5.6	4.7	5.2	3.8	2.7	5.5	+1.7* ss	
Gender:																				
Male	—	7.1	6.9	5.6	6.3	5.9	5.0	6.3	5.8	5.7	6.1	5.9	6.6	5.6	6.8	5.0	§	6.8	+1.7	
Female	—	5.0	3.3	3.3	3.9	4.1	4.4	3.5	3.6	4.0	3.4	4.0	4.5	3.6	4.0	2.6	§	3.4	+0.8	
College Plans:																				
None or under 4 years	—	7.1	8.6	5.3	6.9	4.6	6.5	6.3	5.5	9.0	3.7	9.0	9.6	6.3	9.8	5.3	§	7.8	+2.5	
Complete 4 years	—	5.9	4.8	4.4	4.9	4.9	4.5	4.8	4.7	4.7	4.8	4.5	5.2	4.5	4.6	3.6	§	5.1	+1.5 s	
Region:																				
Northeast	—	6.2	5.0	4.1	4.8	6.1	3.5	4.1	4.4	5.0	3.7	4.5	4.4	4.8	5.9	5.1	§	5.0	-0.1	
Midwest	—	6.9	5.2	5.5	4.8	3.6	5.3	5.8	5.3	4.7	5.3	5.9	6.2	5.6	5.1	3.6	§	6.3	+2.6	
South	—	5.9	5.7	4.6	6.5	5.7	5.3	4.7	5.6	5.9	5.1	5.1	6.7	5.6	6.9	3.5	§	5.1	+1.7	
West	—	5.6	4.4	3.5	3.6	4.2	3.8	5.0	2.8	3.8	3.7	3.6	3.7	2.3	2.0	3.5	§	5.7	+2.2	
Population Density:																				
Large MSA	—	5.5	4.1	3.9	5.4	4.9	4.1	4.9	4.7	5.0	3.9	4.1	4.9	5.0	4.6	2.9	§	4.7	+1.8	
Other MSA	—	6.7	5.8	4.7	5.1	5.3	5.4	5.2	5.2	5.0	5.4	5.6	6.0	4.4	6.5	3.9	§	6.6	+2.7 ss	
Non-MSA	—	5.7	5.6	5.0	4.6	4.1	3.9	4.4	3.8	5.0	3.9	4.5	5.6	5.1	3.4	4.8	§	4.3	-0.6	
Parental Education: ^a																				
1.0–2.0 (Low)	—	7.0	4.5	3.7	4.9	4.7	3.3	3.7	2.9	2.9	2.9	4.4	2.7	3.0	4.9	2.1	§	3.5	+1.4	
2.5–3.0	—	5.3	5.0	4.5	3.0	3.9	3.6	4.5	4.3	3.8	3.4	4.0	5.0	4.7	2.3	4.0	§	5.3	+1.3	
3.5–4.0	—	6.1	5.3	3.4	4.4	4.1	4.3	4.7	4.2	5.1	5.8	4.4	5.4	4.1	4.5	4.1	§	7.0	+2.9	
4.5–5.0	—	5.9	4.5	4.7	5.8	5.6	4.7	5.2	5.8	5.3	5.5	5.7	5.9	5.7	8.2	4.3	§	5.1	+0.8	
5.5–6.0 (High)	—	7.1	5.3	6.2	8.4	6.2	6.1	5.3	4.5	6.1	4.3	5.3	6.5	5.7	5.4	4.0	§	5.8	+1.8	
Race/Ethnicity (2-year average): ^b																				
White	—	—	6.4	5.5	5.9	6.3	5.9	5.8	6.0	5.8	5.7	6.3	7.5	7.3	7.0	6.2	§	5.1	-1.1	
African American	—	—	3.3	3.2	3.5	3.8	3.2	3.1	3.1	4.0	4.2	3.5	3.1	3.8	3.9	1.9	§	0.8	-1.1	
Hispanic	—	—	3.4	2.7	2.2	2.3	2.4	3.0	2.7	2.2	2.7	2.6	2.5	2.1	1.6	2.3	§	2.1	-0.2	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; N is one third of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 179
EITHER STIMULANT-TYPE OR NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 10

	Percentage who said "Yes, I take them now."																		2019– 2021 change
	1991– 2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	5.6	4.8	4.2	4.5	5.0	4.6	4.2	5.1	5.0	4.8	5.8	4.3	4.0	5.1	4.4	4.0	4.8	+0.4*
Gender:																			
Male	—	6.4	5.6	4.8	5.2	5.7	5.8	4.5	5.8	6.3	5.9	6.8	4.7	4.5	6.2	5.0	§	4.3	-0.7
Female	—	4.5	3.9	3.7	3.8	4.3	3.2	4.0	4.5	3.6	3.7	4.9	3.8	3.5	4.2	3.8	§	5.0	+1.2
College Plans:																			
None or under 4 years	—	8.7	6.2	5.4	4.8	9.9	7.7	6.2	3.9	7.5	7.7	9.3	6.5	5.4	5.6	6.3	§	7.1	+0.7
Complete 4 years	—	5.2	4.5	4.0	4.4	4.5	4.2	4.0	5.2	4.7	4.4	5.5	4.1	3.9	4.9	4.0	§	4.4	+0.3
Region:																			
Northeast	—	6.0	3.8	4.0	4.3	4.4	4.0	4.1	4.8	7.4	4.6	5.8	4.1	3.2	3.5	3.4	§	7.0	+3.5 ss
Midwest	—	5.7	4.7	5.1	6.2	7.3	4.1	4.5	4.6	5.6	5.3	6.5	4.7	3.4	6.5	4.9	§	5.3	+0.3
South	—	6.2	5.7	4.6	4.8	5.5	6.0	4.9	7.2	4.5	5.9	6.7	5.1	5.1	5.8	3.4	§	4.3	+0.9
West	—	3.9	4.2	2.9	2.6	2.4	3.4	2.9	2.9	3.3	2.7	3.5	3.1	3.6	3.6	6.2	§	3.6	-2.6
Population Density:																			
Large MSA	—	4.6	4.0	3.2	4.1	5.5	4.9	4.5	4.8	5.8	4.9	5.6	3.7	3.6	4.9	2.7	§	4.6	+1.9 s
Other MSA	—	6.4	5.4	4.8	4.3	5.0	4.2	3.8	5.2	5.0	4.5	6.3	3.9	4.6	5.2	5.6	§	5.3	-0.3
Non-MSA	—	4.9	4.4	4.2	5.5	4.3	5.0	4.8	5.3	4.0	5.3	4.7	6.5	3.3	5.1	4.0	§	4.1	+0.1
Parental Education: ^a																			
1.0–2.0 (Low)	—	3.5	5.7	3.6	2.6	2.3	3.0	3.5	2.3	2.7	1.7	1.3	2.2	3.1	3.3	2.4	§	2.1	-0.3
2.5–3.0	—	4.2	3.2	2.8	4.6	4.4	4.7	4.7	5.1	3.7	2.8	3.9	2.3	3.3	3.4	4.2	§	5.0	+0.8
3.5–4.0	—	5.9	5.1	4.5	3.7	3.8	5.3	3.1	5.2	5.3	5.1	6.0	3.5	4.4	4.2	3.7	§	4.2	+0.5
4.5–5.0	—	5.6	4.9	4.5	5.0	5.8	4.2	4.9	5.2	5.6	4.8	5.9	5.7	3.9	6.8	4.6	§	5.2	+0.6
5.5–6.0 (High)	—	7.6	5.3	5.1	5.2	8.1	5.7	4.7	6.1	7.5	7.7	8.4	5.4	4.9	7.2	7.1	§	6.0	-1.1
Race/Ethnicity (2-year average): ^b																			
White	—	—	6.1	5.3	5.2	5.5	5.4	5.1	5.4	6.1	6.2	7.0	7.0	5.4	6.0	6.5	§	5.9	-0.7
African American	—	—	2.6	2.7	2.6	2.1	2.3	3.1	3.8	4.0	2.8	2.2	1.9	1.8	2.2	2.6	§	2.3	-0.3
Hispanic	—	—	2.2	1.7	1.9	2.8	2.7	2.3	2.8	1.9	1.3	1.7	1.4	1.9	2.6	2.1	§	0.7	-1.4 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on one of four forms; N is one third of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 180
EITHER STIMULANT-TYPE OR NON-STIMULANT-TYPE ADHD DRUGS:
Trends in Current Prevalence of Legal Use by Subgroups in Grade 12

	Percentage who said "Yes, I take them now."																		2019– 2021 change	
	1991– 2004	2005 ^c	2006 ^c	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
Approximate weighted N =	—	14,500	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300		
Total	—	4.5	3.7	4.1	4.4	4.3	5.2	5.1	5.5	6.0	5.5	5.3	5.6	5.7	5.9	5.0	4.2	5.2	+0.2*	
Gender:																				
Male	—	5.5	3.9	5.2	3.9	3.9	5.8	5.2	5.5	6.5	5.1	5.7	6.7	5.9	6.1	6.0	§	4.2	-1.7	
Female	—	3.4	3.5	3.4	4.6	4.5	4.6	4.9	5.4	5.6	5.7	4.0	4.4	5.3	5.1	4.0	§	5.5	+1.5	
College Plans:																				
None or under 4 years	—	8.2	3.2	5.9	5.2	3.1	6.5	6.9	5.8	6.9	6.3	5.6	4.0	5.5	6.5	5.3	§	3.8	-1.5	
Complete 4 years	—	3.7	3.7	3.8	4.4	4.3	4.9	4.6	5.3	5.7	5.2	5.4	5.6	5.9	5.4	4.9	§	5.5	+0.6	
Region:																				
Northeast	—	6.3	4.1	4.7	5.9	3.0	5.0	6.3	4.8	5.1	5.5	3.8	6.0	7.1	5.4	3.0	§	5.9	+2.9 s	
Midwest	—	4.0	3.5	5.0	4.5	4.5	6.7	6.3	7.2	5.0	6.1	6.8	5.7	7.1	6.3	6.4	§	4.8	-1.6	
South	—	4.2	4.2	4.5	3.9	5.8	5.4	4.9	4.8	7.4	6.6	6.6	5.9	5.6	6.2	4.9	§	4.6	-0.3	
West	—	4.0	2.7	2.1	4.5	2.3	3.3	3.3	4.8	5.7	3.2	3.0	4.7	4.0	5.3	4.8	§	6.0	+1.2	
Population Density:																				
Large MSA	—	3.5	3.7	4.5	5.7	4.5	4.3	4.6	5.6	6.9	5.4	5.1	6.4	6.1	6.2	4.3	§	4.9	+0.6	
Other MSA	—	5.6	3.6	4.2	4.3	3.7	6.0	6.4	6.3	6.2	5.7	5.7	6.0	6.2	5.4	5.2	§	5.0	-0.1	
Non-MSA	—	3.5	3.8	3.4	3.5	4.9	4.4	2.6	3.0	4.3	5.3	4.8	3.6	4.2	6.4	5.5	§	6.0	+0.5	
Parental Education: ^a																				
1.0–2.0 (Low)	—	2.7	3.3	2.3	4.2	2.3	5.1	4.8	3.7	5.7	4.8	3.3	4.8	3.0	2.6	4.1	§	2.6	-1.5	
2.5–3.0	—	3.7	2.0	3.8	3.2	2.5	4.4	3.6	4.8	5.7	4.4	2.8	3.4	5.9	4.1	3.4	§	3.5	+0.1	
3.5–4.0	—	5.0	3.3	4.1	4.1	3.9	5.2	5.3	5.0	5.1	4.2	4.1	5.2	4.3	5.6	4.6	§	2.9	-1.7	
4.5–5.0	—	5.0	4.6	3.7	4.8	4.8	5.4	5.6	5.7	5.8	5.7	6.5	6.8	8.3	6.8	5.0	§	9.6	+4.6 s	
5.5–6.0 (High)	—	4.8	5.3	6.2	6.5	7.4	6.5	6.3	8.3	9.9	11.0	9.0	6.9	7.0	10.4	7.1	§	6.8	-0.4	
Race/Ethnicity (2-year average): ^b																				
White	—	—	4.6	4.5	4.9	4.9	5.6	6.6	6.6	6.4	6.3	6.4	7.0	7.2	7.5	6.7	§	6.1	-0.6	
African American	—	—	2.3	2.5	3.0	2.7	3.0	3.2	3.2	4.1	4.3	3.2	2.5	3.8	4.8	4.9	§	5.4	+0.4	
Hispanic	—	—	2.3	2.2	2.4	2.8	3.1	2.5	2.7	3.3	3.5	2.5	2.3	3.5	2.6	1.8	§	0.7	-1.1 s	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data are based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cData from one of the questionnaire forms was omitted in 2005 due to an error in the skip pattern in the questionnaire. For the 2006 results by race/ethnicity, data from the remaining form in 2005 were double-weighted to account for the omission of the other form.

TABLE 181
Any Nicotine Use^c: Trends in 30-Day
Prevalence of Use by Subgroups in Grade 12

	Percentage who used in past 30 days									
	12th Graders									
	1976– <u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	2019– 2021 change
<i>Approximate weighted N =</i>	—	—	—	—	12,600	13,300	12,900	3,500	8,300	
Total	—	—	—	—	25.6	32.5	33.6	§	24.6	-9.0* ss
Gender:										
Male	—	—	—	—	29.9	36.5	37.9	§	22.6	-15.3 sss
Female	—	—	—	—	20.7	28.0	28.9	§	25.5	-3.4
College Plans:										
None or under 4 years	—	—	—	—	37.0	42.7	37.6	§	26.5	-11.1 s
Complete 4 years	—	—	—	—	22.8	28.5	30.8	§	21.8	-9.0 ss
Region:										
Northeast	—	—	—	—	29.8	37.9	33.6	§	29.5	-4.1
Midwest	—	—	—	—	24.5	30.9	36.1	§	28.5	-7.6
South	—	—	—	—	27.2	35.1	32.6	§	24.9	-7.7 s
West	—	—	—	—	21.2	26.1	32.4	§	15.8	-16.6
Population Density:										
Large MSA	—	—	—	—	23.4	25.4	28.8	§	20.5	-8.3
Other MSA	—	—	—	—	24.0	35.3	34.6	§	22.7	-11.9 s
Non-MSA	—	—	—	—	32.2	38.0	38.6	§	34.9	-3.7
Parental Education: ^a										
1.0–2.0 (Low)	—	—	—	—	19.5	23.6	30.9	§	24.9	-6.0
2.5–3.0	—	—	—	—	28.5	32.6	37.1	§	20.4	-16.7 sss
3.5–4.0	—	—	—	—	25.5	32.9	34.2	§	27.7	-6.5
4.5–5.0	—	—	—	—	28.1	36.0	30.0	§	25.0	-5.0
5.5–6.0 (High)	—	—	—	—	24.1	32.5	32.3	§	23.1	-9.2
Race/Ethnicity (2-year data): ^b										
White	—	—	—	—	—	36.1	37.8	§	31.1	-6.7 s
African American	—	—	—	—	—	13.4	15.2	§	20.6	+5.4
Hispanic	—	—	—	—	—	17.9	24.4	§	12.1	-12.3 s

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cIncludes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, smokeless tobacco, or vaping nicotine.

TABLE 182
Any Nicotine Use other than Vaping^c: Trends in 30-Day
Prevalence of Use by Subgroups in Grade 12

	Percentage who used in past 30 days									
	12th Graders									
	1976– <u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>	2019– 2021 change
<i>Approximate weighted N =</i>	—	—	—	—	12,600	13,300	12,900	3,500	8,300	
Total	—	—	—	—	20.6	18.5	15.7	§	7.7	-8.0* sss
Gender:										
Male	—	—	—	—	24.5	23.9	20.0	§	7.8	-12.2 sss
Female	—	—	—	—	16.1	12.9	11.1	§	6.5	-4.5
College Plans:										
None or under 4 years	—	—	—	—	32.2	29.7	20.4	§	9.6	-10.8 ss
Complete 4 years	—	—	—	—	17.6	14.4	13.2	§	6.6	-6.6 sss
Region:										
Northeast	—	—	—	—	24.4	14.9	11.7	§	6.6	-5.1 s
Midwest	—	—	—	—	18.5	17.1	15.7	§	7.5	-8.2 s
South	—	—	—	—	23.5	23.1	16.3	§	7.9	-8.4 ss
West	—	—	—	—	15.4	14.5	17.8	§	8.5	-9.2
Population Density:										
Large MSA	—	—	—	—	18.6	13.5	13.4	§	7.8	-5.6 s
Other MSA	—	—	—	—	19.3	19.2	15.9	§	5.9	-10.0 ss
Non-MSA	—	—	—	—	26.4	24.9	18.8	§	11.4	-7.4 s
Parental Education: ^a										
1.0–2.0 (Low)	—	—	—	—	17.0	14.2	20.5	§	14.6	-5.9
2.5–3.0	—	—	—	—	23.7	21.9	16.2	§	4.5	-11.7 sss
3.5–4.0	—	—	—	—	19.9	21.4	14.1	§	7.7	-6.4
4.5–5.0	—	—	—	—	22.5	16.2	11.7	§	5.6	-6.1 s
5.5–6.0 (High)	—	—	—	—	18.4	13.2	16.9	§	9.4	-7.4 s
Race/Ethnicity (2-year data): ^b										
White	—	—	—	—	—	23.4	18.2	§	10.1	-8.1 sss
African American	—	—	—	—	—	10.1	8.1	§	11.3	+3.2
Hispanic	—	—	—	—	—	13.2	15.6	§	3.5	-12.1 sss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; N is one sixth of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

^cIncludes use of any of the following: cigarettes, large cigars, flavored small cigars, regular small cigars, tobacco using a hookah, or smokeless tobacco.

TABLE 183
Use of Marijuana Under a Doctor's Orders: Trends in Lifetime
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in lifetime																				
	8th Graders							10th Graders							12th Graders						
	1991– 2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2016	2017	2018	2019	2020	2021	2019– 2021 change	1976– 2016	2017	2018	2019	2020	2021	2019– 2021 change
<i>Approximate weighted N =</i>	—	15,300	14,000	13,600	3,100	10,700		—	13,500	14,300	14,000	4,800	11,000		—	12,600	13,300	12,900	3,500	8,300	
Total	—	1.1	1.1	1.3	1.0	1.3	+0.1*	—	1.1	1.3	2.0	2.0	1.4	-0.7*	—	1.5	1.2	2.0	§	2.3	+0.3*
Gender:																					
Male	—	1.1	1.1	1.3	§	1.4	0.0	—	1.0	1.6	2.2	§	1.1	-1.1	—	1.9	1.3	1.8	§	1.6	-0.2
Female	—	0.9	1.0	1.2	§	1.2	0.0	—	1.2	0.9	1.7	§	1.5	-0.2	—	1.2	0.8	1.2	§	2.7	+1.5
College Plans:																					
None or under 4 years	—	2.2	1.1	3.0	§	2.1	-0.9	—	2.8	1.2	3.2	§	1.8	-1.4	—	4.1	1.7	3.2	§	3.5	+0.3
Complete 4 years	—	0.8	1.0	1.0	§	1.2	+0.3	—	0.8	1.3	1.6	§	1.2	-0.5	—	0.8	0.8	1.4	§	1.2	-0.2
Region:																					
Northeast	—	0.4	0.8	0.9	§	1.4	+0.5	—	0.7	1.0	1.8	§	0.8	-1.0	—	0.7	1.2	1.4	§	2.3	+0.9
Midwest	—	0.4	0.7	1.0	§	1.1	+0.1	—	1.1	1.0	1.5	§	0.7	-0.8	—	0.2	0.7	0.8	§	1.6	+0.8
South	—	1.6	1.2	1.4	§	1.1	-0.4	—	0.8	1.6	2.3	§	2.0	-0.3	—	1.4	1.4	1.7	§	2.0	+0.3
West	—	1.2	1.5	1.5	§	1.9	+0.4	—	1.8	1.2	2.2	§	1.3	-0.9	—	3.2	1.5	4.8	§	3.3	-1.5
Population Density:																					
Large MSA	—	1.0	0.8	1.5	§	1.4	-0.1	—	0.8	0.9	1.8	§	1.1	-0.7	—	1.6	1.4	1.7	§	2.8	+1.1
Other MSA	—	1.4	1.1	1.1	§	1.2	+0.2	—	1.4	1.5	2.3	§	1.0	-1.2 s	—	1.0	1.1	2.4	§	1.8	-0.6
Non-MSA	—	0.3	1.7	1.4	§	1.4	+0.1	—	0.7	1.4	1.8	§	2.7	+1.0	—	2.3	1.1	1.5	§	2.4	+0.9
Parental Education: a																					
1.0–2.0 (Low)	—	1.0	2.5	1.5	§	4.0	+2.5	—	1.8	1.5	2.1	§	2.7	+0.5	—	2.5	1.2	2.4	§	3.6	+1.2
2.5–3.0	—	0.9	1.3	2.1	§	1.7	-0.3	—	2.4	2.2	2.8	§	1.4	-1.4	—	2.4	1.0	2.8	§	2.5	-0.3
3.5–4.0	—	1.0	1.1	0.7	§	2.3	+1.5 s	—	0.8	0.6	2.1	§	0.8	-1.3 s	—	1.2	1.5	1.7	§	2.1	+0.4
4.5–5.0	—	0.5	0.4	1.1	§	0.2	-0.9 ss	—	0.3	1.3	1.3	§	1.1	-0.3	—	0.6	1.0	1.3	§	1.0	-0.3
5.5–6.0 (High)	—	1.0	0.7	1.2	§	0.8	-0.3	—	0.7	0.7	1.8	§	1.0	-0.8	—	1.0	1.0	1.1	§	1.7	+0.6
Race/Ethnicity (2-year data): b																					
White	—	—	0.7	0.9	§	1.1	+0.2	—	—	0.6	1.2	§	1.0	-0.1	—	—	0.8	0.9	§	3.0	+2.1 ss
African American	—	—	1.3	1.1	§	1.2	+0.1	—	—	2.1	2.2	§	1.9	-0.3	—	—	1.8	1.6	§	2.2	+0.5
Hispanic	—	—	1.1	1.6	§	0.8	-0.8	—	—	1.8	1.7	§	1.9	+0.2	—	—	1.8	2.4	§	2.3	-0.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only; Data based on one of four forms; N is one third of N indicated in S1 and S2. For 12th graders only; Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 184
Powdered Alcohol: Trends in Annual
Prevalence of Use by Subgroups in Grades 8, 10, and 12

	Percentage who used in last 12 months																							
	8th Graders							10th Graders							12th Graders									
	1991– 2015	2016	2017	2018	2019	2020	2021	2019– 2021 change	1991– 2015	2016	2017	2018	2019	2020	2021	2019– 2021 change	1976– 2015	2016	2017	2018	2019	2020	2021	2019– 2021 change
Approximate weighted N =	—	16,900	15,300	14,000	13,600	—	—	—	—	14,700	13,500	14,300	14,000	—	—	—	—	11,800	12,600	13,300	12,900	—	—	—
Total	—	1.0	0.8	0.8	1.2	—	—	—	—	1.3	0.8	1.2	1.0	—	—	—	—	1.7	1.0	1.3	1.4	—	—	—
Gender:																								
Male	—	1.3	0.9	0.8	1.2	—	—	—	—	1.8	1.0	1.5	1.4	—	—	—	—	1.9	1.3	1.4	2.2	—	—	—
Female	—	0.8	0.6	0.7	1.3	—	—	—	—	0.8	0.7	0.9	0.5	—	—	—	—	1.3	0.5	0.7	0.5	—	—	—
College Plans:																								
None or under 4 years	—	4.5	2.4	2.1	2.7	—	—	—	—	3.2	2.3	1.6	1.8	—	—	—	—	4.1	1.2	2.8	2.9	—	—	—
Complete 4 years	—	0.7	0.6	0.6	1.0	—	—	—	—	1.0	0.6	1.1	0.8	—	—	—	—	1.1	0.7	0.7	0.9	—	—	—
Region:																								
Northeast	—	0.4	0.6	1.1	1.2	—	—	—	—	1.4	0.7	2.2	0.4	—	—	—	—	1.2	0.7	0.8	1.3	—	—	—
Midwest	—	0.8	0.6	0.1	0.7	—	—	—	—	1.2	0.4	1.4	0.7	—	—	—	—	1.8	1.0	1.5	0.5	—	—	—
South	—	1.2	0.8	0.7	1.4	—	—	—	—	1.3	0.9	0.7	0.6	—	—	—	—	2.1	1.3	1.5	1.6	—	—	—
West	—	1.2	1.1	1.2	1.5	—	—	—	—	1.3	1.1	1.2	2.4	—	—	—	—	1.3	0.6	1.3	2.1	—	—	—
Population Density:																								
Large MSA	—	1.1	0.9	0.9	1.3	—	—	—	—	0.9	0.7	1.7	0.6	—	—	—	—	1.7	1.1	1.6	2.2	—	—	—
Other MSA	—	0.9	0.9	0.8	1.4	—	—	—	—	1.3	0.7	1.2	1.1	—	—	—	—	1.8	1.1	1.1	1.0	—	—	—
Non-MSA	—	0.9	0.5	0.4	0.9	—	—	—	—	2.0	1.4	0.8	1.5	—	—	—	—	1.6	0.6	1.5	1.1	—	—	—
Parental Education: a																								
1.0–2.0 (Low)	—	2.0	1.5	1.8	0.2	—	—	—	—	1.3	2.2	2.1	0.6	—	—	—	—	3.8	1.7	2.4	2.7	—	—	—
2.5–3.0	—	1.1	0.1	1.1	1.9	—	—	—	—	2.3	0.3	1.7	0.3	—	—	—	—	1.6	1.7	1.7	1.1	—	—	—
3.5–4.0	—	1.4	0.8	0.5	1.2	—	—	—	—	1.0	1.1	1.1	1.8	—	—	—	—	2.1	0.5	0.9	1.2	—	—	—
4.5–5.0	—	0.7	0.9	0.5	1.0	—	—	—	—	0.6	0.3	0.7	1.2	—	—	—	—	0.8	0.4	0.5	0.7	—	—	—
5.5–6.0 (High)	—	0.4	0.7	0.3	1.1	—	—	—	—	1.4	0.2	1.1	0.2	—	—	—	—	1.2	0.7	1.0	0.8	—	—	—
Race/Ethnicity (2-year data): ^b																								
White	—	—	0.6	0.6	0.8	—	—	—	—	—	0.8	0.7	1.0	—	—	—	—	—	1.0	0.7	0.7	—	—	—
African American	—	—	1.3	1.2	1.5	—	—	—	—	—	1.3	0.9	1.1	—	—	—	—	—	1.5	0.9	2.6	—	—	—
Hispanic	—	—	1.3	1.1	1.3	—	—	—	—	—	1.6	2.1	1.6	—	—	—	—	—	1.3	1.2	1.8	—	—	—

Source: The Monitoring the Future study, the University of Michigan.

Notes: '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

See Tables S1, S2, and S3 for the number of subgroup cases. See appendix B for definition of variables in table. For 8th and 10th graders only: Data based on one of four forms; N is one third of N indicated in S1 and S2. For 12th graders only: Data based on two of six forms; N is two sixths of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE 185
ENERGY DRINKS: Trends in Daily Prevalence
of Use by Subgroups in Grade 8

	Percent who used daily													2019– 2020 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	18.6	17.7	16.3	14.2	12.8	12.1	11.3	10.1	10.3	10.5	§	13.8	+3.3* ss
Gender:														
Male	—	22.5	20.3	17.6	16.4	14.9	14.0	12.8	12.5	13.8	12.2	§	11.7	-0.5
Female	—	15.2	15.0	14.9	11.9	10.5	10.2	9.8	6.9	6.4	8.8	§	14.9	+6.1 sss
College Plans:														
None or under 4 years	—	32.2	37.5	36.9	32.2	25.9	27.7	25.0	20.2	20.4	21.0	§	21.0	0.0
Complete 4 years	—	17.1	16.1	14.9	12.8	11.5	10.5	9.9	8.9	8.8	9.0	§	12.5	+3.5 ss
Region:														
Northeast	—	19.7	14.9	12.3	12.5	11.5	8.2	8.8	6.6	11.2	9.5	§	9.3	-0.2
Midwest	—	18.0	17.3	13.8	15.4	12.5	11.5	11.6	8.6	8.4	10.6	§	16.8	+6.1 ss
South	—	20.4	19.6	18.2	16.7	14.1	14.7	12.4	11.2	10.1	10.9	§	15.9	+4.9 ss
West	—	15.5	17.0	18.7	9.9	11.7	10.5	11.0	11.8	11.8	10.6	§	11.1	+0.5
Population Density:														
Large MSA	—	14.1	15.8	13.6	12.4	12.4	9.4	10.4	8.7	8.8	9.3	§	10.0	+0.7
Other MSA	—	21.3	18.1	16.3	13.0	12.6	11.7	11.3	11.2	11.0	10.0	§	15.5	+5.5 sss
Non-MSA	—	18.5	19.8	20.9	19.5	13.9	16.7	13.0	9.4	10.9	13.9	§	16.3	+2.4
Parental Education: ^a														
1.0–2.0 (Low)	—	23.5	26.5	22.9	20.8	17.7	15.4	15.8	11.6	12.2	15.3	§	22.9	+7.6
2.5–3.0	—	23.2	23.4	21.7	17.4	15.1	15.8	13.0	13.1	10.9	13.9	§	19.2	+5.4
3.5–4.0	—	18.8	18.6	18.9	13.7	12.3	11.2	10.9	11.1	11.4	11.6	§	18.8	+7.2 ss
4.5–5.0	—	15.5	12.4	11.8	11.4	9.7	8.9	9.1	7.9	8.3	9.0	§	12.1	+3.1
5.5–6.0 (High)	—	11.5	10.8	9.0	9.9	9.4	6.9	8.3	5.3	9.6	6.9	§	7.7	+0.8
Race/Ethnicity (2-year data): ^b														
White	—	—	16.5	14.7	12.8	10.3	10.7	10.4	9.0	8.3	9.5	§	12.9	+3.3 s
African American	—	—	19.1	18.2	15.8	14.3	13.3	13.1	14.0	13.1	9.8	§	12.0	+2.1
Hispanic	—	—	22.6	22.6	18.5	15.2	14.5	13.4	11.9	10.7	10.8	§	8.9	-1.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 186
ENERGY DRINKS: Trends in Daily Prevalence
of Use by Subgroups in Grade 10

	Percent who used daily													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	13.6	11.4	10.8	10.3	9.6	7.8	9.2	8.8	9.1	10.5	§	12.6	+2.2*
Gender:														
Male	—	17.0	14.1	13.8	12.5	11.4	9.0	10.8	9.9	10.9	12.8	§	11.3	-1.5
Female	—	10.0	8.9	7.8	8.1	7.8	6.3	6.8	7.3	7.3	8.2	§	13.2	+5.1 sss
College Plans:														
None or under 4 years	—	28.1	23.9	27.2	18.7	20.4	15.3	17.4	21.6	17.5	18.9	§	22.6	+3.7
Complete 4 years	—	11.8	9.9	9.1	9.3	8.2	7.0	8.0	7.1	7.7	8.9	§	10.7	+1.8
Region:														
Northeast	—	11.0	9.3	7.1	8.1	6.7	7.0	7.2	8.1	9.8	8.1	§	14.0	+5.9 s
Midwest	—	12.0	11.1	12.1	8.6	7.8	6.7	8.7	9.8	10.0	10.0	§	12.4	+2.4
South	—	15.8	14.3	11.6	11.2	10.5	7.8	10.0	6.8	8.7	10.0	§	12.9	+2.9
West	—	14.3	9.1	11.5	12.5	12.3	9.6	10.0	11.2	8.3	13.3	§	11.5	-1.8
Population Density:														
Large MSA	—	11.0	10.4	10.4	9.3	8.5	7.5	7.6	8.3	8.6	7.3	§	8.8	+1.5
Other MSA	—	13.6	10.0	9.6	9.9	7.8	7.8	9.3	8.3	8.8	11.2	§	13.5	+2.3
Non-MSA	—	17.4	15.5	14.1	12.7	15.7	8.5	11.9	11.0	10.6	13.9	§	17.6	+3.7
Parental Education: ^a														
1.0–2.0 (Low)	—	18.8	16.5	15.7	19.4	13.4	10.7	12.2	14.3	14.9	9.5	§	19.5	#### sss
2.5–3.0	—	18.1	16.5	15.2	15.3	14.0	10.6	12.6	9.2	9.3	12.3	§	15.3	+3.0
3.5–4.0	—	14.0	10.7	10.6	10.2	9.0	7.4	8.7	8.7	9.8	10.6	§	13.8	+3.2
4.5–5.0	—	8.3	7.6	8.0	5.5	7.5	5.7	7.3	6.7	6.7	8.5	§	9.2	+0.7
5.5–6.0 (High)	—	8.9	8.3	6.9	6.3	5.4	5.4	5.4	4.9	6.9	8.6	§	10.7	+2.1
Race/Ethnicity (2-year data): ^b														
White	—	—	11.7	10.0	9.1	9.0	8.0	7.8	8.6	8.8	9.9	§	13.7	+3.8 ss
African American	—	—	13.6	11.5	10.2	10.2	10.3	8.7	7.6	8.0	8.7	§	9.7	+1.0
Hispanic	—	—	14.2	14.1	14.7	11.8	9.2	9.3	9.8	10.1	9.7	§	10.1	+0.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 187
ENERGY DRINKS: Trends in Daily Prevalence
of Use by Subgroups in Grade 12

	Percent who used daily													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	12.3	9.5	9.2	8.2	8.3	7.8	9.8	9.4	10.1	11.6	§	13.1	+1.4*
Gender:														
Male	—	14.8	11.6	10.7	10.7	10.1	9.0	11.2	12.7	12.2	15.0	§	12.5	-2.6
Female	—	8.9	7.5	7.5	5.8	6.4	6.7	7.8	6.3	7.6	7.9	§	13.6	+5.7 s
College Plans:														
None or under 4 years	—	18.5	14.1	16.5	15.3	18.8	13.3	19.7	15.1	18.4	22.7	§	18.8	-4.0
Complete 4 years	—	11.1	8.4	7.6	6.3	5.8	6.4	7.8	8.0	7.8	8.2	§	11.2	+3.0
Region:														
Northeast	—	15.7	7.2	8.6	9.7	9.5	4.5	11.2	7.3	7.7	8.4	§	7.6	-0.8
Midwest	—	10.7	8.9	7.2	7.5	9.1	10.6	8.3	10.9	12.4	12.7	§	11.6	-1.2
South	—	11.9	10.3	12.4	8.2	7.4	6.5	12.4	10.8	9.8	11.7	§	15.0	+3.3
West	—	12.4	10.4	7.8	7.8	8.3	9.5	5.5	7.2	10.0	13.0	§	15.2	+2.3
Population Density:														
Large MSA	—	10.6	8.2	9.6	8.7	6.9	8.4	9.1	10.5	9.4	9.1	§	9.8	+0.7
Other MSA	—	11.7	8.8	9.0	7.0	8.3	7.5	10.1	8.4	8.7	11.2	§	12.6	+1.4
Non-MSA	—	16.3	12.9	9.4	10.2	10.2	7.4	10.3	9.8	14.8	16.9	§	19.3	+2.4
Parental Education: ^a														
1.0–2.0 (Low)	—	19.0	10.8	14.1	17.4	12.3	10.6	12.8	16.0	11.1	9.4	§	9.3	-0.1
2.5–3.0	—	16.9	12.9	10.9	7.6	14.1	9.0	11.5	10.3	12.4	11.7	§	11.3	-0.4
3.5–4.0	—	11.2	10.5	11.8	8.2	7.0	6.9	11.8	9.3	9.7	14.7	§	17.4	+2.7
4.5–5.0	—	10.1	5.9	5.3	5.7	4.9	6.8	4.7	6.2	7.7	10.0	§	10.3	+0.3
5.5–6.0 (High)	—	3.5	8.4	5.6	5.3	4.8	6.2	9.2	6.4	9.8	6.4	§	14.5	+8.1 s
Race/Ethnicity (2-year data): ^b														
White	—	—	10.4	8.1	7.6	8.5	9.1	9.3	9.5	10.1	11.5	§	12.4	+1.0
African American	—	—	8.9	10.7	10.4	7.9	7.6	8.9	10.2	8.1	9.2	§	7.9	-1.3
Hispanic	—	—	11.8	12.8	10.9	7.9	6.2	5.8	9.1	8.7	7.9	§	6.3	-1.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 188
ENERGY SHOTS: Trends in Daily Prevalence
of Use by Subgroups in Grade 8

	Percent who used daily													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	6.4	6.8	5.7	5.6	4.2	5.3	4.4	4.0	3.7	4.6	§	3.7	-0.9*
Gender:														
Male	—	7.6	8.5	6.6	6.8	4.6	6.6	5.1	5.4	5.2	5.5	§	3.8	-1.7
Female	—	5.3	5.2	4.7	4.3	3.7	4.2	3.7	2.4	1.9	3.6	§	3.6	0.0
College Plans:														
None or under 4 years	—	12.9	17.5	17.7	18.7	12.3	14.2	13.4	6.9	8.2	9.5	§	7.2	-2.3
Complete 4 years	—	5.7	6.0	4.8	4.7	3.4	4.4	3.4	3.5	3.0	3.9	§	3.1	-0.7
Region:														
Northeast	—	7.8	5.8	4.0	3.4	2.8	2.5	3.1	2.9	4.0	3.9	§	1.1	-2.8 ss
Midwest	—	5.1	6.0	5.6	6.2	3.6	5.1	3.9	2.9	2.7	3.2	§	3.1	-0.1
South	—	7.4	7.9	6.5	7.6	5.4	7.0	5.0	5.1	4.3	6.3	§	6.3	0.0
West	—	5.4	6.7	5.7	3.4	4.0	4.4	4.8	3.7	3.7	3.9	§	2.1	-1.8
Population Density:														
Large MSA	—	5.3	6.7	5.5	5.0	4.6	4.6	4.6	3.8	4.1	4.4	§	3.0	-1.4
Other MSA	—	7.4	7.6	5.2	4.7	4.1	4.8	4.1	4.7	3.2	4.4	§	3.6	-0.8
Non-MSA	—	5.8	5.3	7.1	8.7	4.0	7.5	4.7	2.5	4.3	5.5	§	5.2	-0.3
Parental Education: ^a														
1.0–2.0 (Low)	—	9.6	14.0	9.0	8.6	8.0	6.8	8.7	5.6	6.6	7.9	§	12.4	+4.5
2.5–3.0	—	8.1	8.7	7.2	6.3	5.4	6.2	4.0	5.1	3.3	5.5	§	4.7	-0.8
3.5–4.0	—	5.1	6.8	6.8	6.0	3.5	4.8	3.6	4.0	3.4	4.0	§	4.8	+0.8
4.5–5.0	—	5.5	4.7	3.4	4.4	2.8	3.8	3.7	3.2	2.7	3.5	§	2.5	-1.0
5.5–6.0 (High)	—	4.6	4.1	3.2	3.7	2.8	4.0	2.2	2.2	3.7	3.3	§	1.8	-1.6
Race/Ethnicity (2-year data): ^b														
White	—	—	5.2	4.4	3.4	3.2	2.9	3.2	2.7	2.5	3.0	§	3.1	+0.1
African American	—	—	7.0	7.7	8.1	7.1	6.6	6.5	6.4	5.8	5.9	§	5.5	-0.4
Hispanic	—	—	10.4	10.1	7.1	5.2	6.3	6.4	5.6	4.6	4.9	§	6.3	+1.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 189
ENERGY SHOTS: Trends in Daily Prevalence
of Use by Subgroups in Grade 10

	Percent who used daily													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	4.3	4.6	4.0	4.0	3.4	2.6	3.3	3.3	3.8	4.1	§	2.6	-1.5* s
Gender:														
Male	—	5.7	5.8	4.8	5.0	4.3	3.0	3.7	4.3	4.8	5.9	§	2.6	-3.3 sss
Female	—	2.8	3.4	2.9	2.8	2.5	2.1	2.8	2.1	2.7	2.2	§	1.9	-0.2
College Plans:														
None or under 4 years	—	10.0	8.6	9.9	6.0	9.9	7.0	5.9	9.8	6.5	7.4	§	5.2	-2.2
Complete 4 years	—	3.6	4.1	3.4	3.6	2.7	2.1	2.9	2.5	3.3	3.5	§	2.1	-1.4 ss
Region:														
Northeast	—	3.8	5.0	3.2	2.9	2.8	2.1	3.0	3.3	3.1	2.4	§	1.3	-1.1
Midwest	—	3.8	3.5	4.6	3.1	2.5	2.5	3.1	3.0	4.0	2.9	§	1.7	-1.3
South	—	5.5	5.9	3.5	4.6	4.0	2.2	3.8	2.8	4.4	4.8	§	4.0	-0.8
West	—	3.4	3.6	4.4	4.9	3.9	3.9	3.1	4.2	3.1	5.1	§	2.3	-2.8 s
Population Density:														
Large MSA	—	4.0	4.4	4.3	4.1	3.1	2.8	3.2	3.5	2.8	2.4	§	1.7	-0.7
Other MSA	—	3.9	4.2	3.5	3.8	2.4	2.2	3.2	3.0	4.1	4.7	§	2.9	-1.9
Non-MSA	—	5.4	5.7	4.4	4.2	6.3	3.4	4.0	3.6	4.7	5.2	§	3.5	-1.7
Parental Education: ^a														
1.0–2.0 (Low)	—	6.5	7.5	8.9	7.4	4.2	3.6	5.4	5.4	6.8	5.0	§	5.0	0.0
2.5–3.0	—	5.8	5.8	5.0	5.8	5.0	3.6	3.3	2.9	3.3	5.1	§	2.9	-2.2
3.5–4.0	—	4.5	3.9	3.7	3.8	2.9	2.0	3.5	3.1	4.6	4.1	§	3.0	-1.1
4.5–5.0	—	2.4	3.1	2.4	2.2	2.6	2.4	2.3	2.6	2.0	2.2	§	1.9	-0.3
5.5–6.0 (High)	—	3.1	4.1	2.3	2.7	2.0	0.9	2.3	1.6	3.0	3.3	§	1.5	-1.9 s
Race/Ethnicity (2-year data): ^b														
White	—	—	3.6	3.3	2.6	2.4	2.4	2.4	2.2	2.7	3.4	§	2.8	-0.6
African American	—	—	7.4	6.9	6.7	6.4	4.6	4.7	5.1	4.9	5.7	§	7.2	+1.5
Hispanic	—	—	5.2	5.7	5.4	4.5	3.4	3.9	4.7	4.8	4.2	§	3.4	-0.8

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 190
ENERGY SHOTS: Trends in Daily Prevalence
of Use by Subgroups in Grade 12

	Percent who used daily													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	4.3	4.0	2.7	2.5	2.1	3.1	4.1	3.8	4.2	4.1	§	2.9	-1.1*
Gender:														
Male	—	5.9	4.5	2.9	3.4	1.6	2.9	4.7	5.1	5.2	5.0	§	4.2	-0.8
Female	—	2.2	3.4	2.7	1.7	2.1	3.0	3.2	2.4	2.3	2.9	§	1.5	-1.4
College Plans:														
None or under 4 years	—	8.9	5.9	4.3	4.1	5.7	4.5	8.4	6.8	7.8	11.3	§	6.9	-4.4
Complete 4 years	—	3.4	3.4	2.4	1.9	1.2	2.6	2.9	2.9	3.0	2.0	§	1.6	-0.4
Region:														
Northeast	—	4.7	5.4	2.9	4.3	2.4	1.5	4.7	3.4	5.1	4.1	§	0.7	-3.4
Midwest	—	3.3	4.3	2.4	1.2	2.2	2.9	2.5	3.8	4.0	3.1	§	5.7	+2.7
South	—	4.8	4.5	3.1	2.3	2.4	4.0	4.7	5.2	4.4	5.1	§	2.7	-2.4
West	—	4.2	2.0	2.6	3.1	1.5	3.0	4.2	2.0	3.4	3.0	§	2.6	-0.5
Population Density:														
Large MSA	—	2.8	2.4	3.1	3.2	2.6	3.3	4.5	4.2	4.8	4.3	§	3.1	-1.2
Other MSA	—	4.7	4.6	2.2	2.0	1.5	3.5	3.6	3.4	3.7	4.0	§	2.3	-1.6
Non-MSA	—	5.2	4.8	3.6	2.8	3.0	1.8	4.7	4.1	4.0	3.9	§	3.9	+0.1
Parental Education: ^a														
1.0–2.0 (Low)	—	9.0	5.0	5.2	6.7	4.0	5.6	8.1	8.5	8.4	7.7	§	4.7	-3.1
2.5–3.0	—	6.5	4.8	3.3	1.9	2.5	3.6	5.0	4.5	5.2	3.0	§	2.6	-0.4
3.5–4.0	—	3.6	5.2	3.4	2.2	2.3	3.2	3.2	2.9	2.9	5.3	§	3.4	-1.9
4.5–5.0	—	2.3	2.9	1.3	2.1	0.6	2.7	2.5	2.6	2.9	2.1	§	1.9	-0.2
5.5–6.0 (High)	—	2.1	2.2	1.9	1.7	1.0	1.9	3.1	1.9	2.1	1.1	§	0.8	-0.3
Race/Ethnicity (2-year data): ^b														
White	—	—	3.1	2.8	1.9	1.8	2.2	2.6	2.7	3.3	3.3	§	1.9	-1.5 s
African American	—	—	6.6	6.9	4.6	3.7	4.5	6.6	8.1	6.0	6.1	§	8.7	+2.7
Hispanic	—	—	5.7	4.0	3.5	2.6	3.0	4.2	4.1	3.9	3.9	§	1.3	-2.6 ss

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 191
EITHER ENERGY DRINKS OR ENERGY SHOTS: Trends in Daily Prevalence
of Use by Subgroups in Grade 8

	Percent who used daily													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	19.5	18.9	17.2	15.4	13.5	13.0	12.3	11.1	11.4	11.7	§	14.5	+2.8* s
Gender:														
Male	—	23.2	22.0	18.8	18.2	15.7	15.2	14.1	14.0	15.2	13.4	§	12.6	-0.8
Female	—	16.0	15.7	15.4	12.4	11.2	10.9	10.5	7.5	7.0	9.8	§	15.4	+5.6 sss
College Plans:														
None or under 4 years	—	34.0	40.9	39.9	36.4	27.5	29.1	28.6	21.8	22.1	23.0	§	21.9	-1.0
Complete 4 years	—	17.9	17.1	15.5	13.8	12.0	11.4	10.7	9.9	9.8	9.9	§	13.1	+3.2 ss
Region:														
Northeast	—	20.9	16.2	12.4	12.9	11.7	8.9	9.7	7.7	12.6	10.2	§	9.5	-0.6
Midwest	—	18.7	18.3	14.6	16.3	13.3	12.1	12.1	9.9	8.9	11.2	§	17.4	+6.2 ss
South	—	21.5	21.0	19.4	19.0	15.2	16.2	13.3	12.2	11.5	12.9	§	17.1	+4.2 s
West	—	16.0	18.1	19.5	10.4	12.2	11.1	12.8	12.8	12.7	11.5	§	11.3	-0.3
Population Density:														
Large MSA	—	15.6	17.0	14.4	13.7	13.3	10.1	11.4	9.7	10.2	10.6	§	10.4	-0.2
Other MSA	—	21.9	19.4	17.3	14.0	13.3	12.6	12.2	12.6	11.7	11.3	§	16.1	+4.8 ss
Non-MSA	—	19.2	20.8	21.5	21.3	14.3	18.0	14.3	9.7	12.5	14.6	§	17.7	+3.1
Parental Education: ^a														
1.0–2.0 (Low)	—	24.9	28.7	24.6	22.5	18.7	16.2	18.2	13.2	14.2	17.6	§	26.2	+8.6
2.5–3.0	—	24.0	24.6	22.4	18.5	15.9	17.0	14.2	14.4	12.3	15.2	§	20.5	+5.3
3.5–4.0	—	19.4	20.3	19.9	15.0	12.8	12.3	11.4	11.8	12.0	12.3	§	20.0	+7.7 ss
4.5–5.0	—	16.3	13.4	12.0	12.8	10.3	9.5	9.9	9.2	9.0	9.6	§	12.2	+2.5
5.5–6.0 (High)	—	12.4	11.7	9.5	10.9	10.2	7.8	8.8	5.7	11.0	7.7	§	7.7	+0.1
Race/Ethnicity (2-year data): ^b														
White	—	—	17.3	15.6	13.4	11.8	11.3	11.1	9.7	9.1	10.4	§	13.6	+3.2 s
African American	—	—	20.4	19.3	17.5	16.9	14.7	14.6	15.1	14.0	11.3	§	13.3	+1.9
Hispanic	—	—	24.0	24.2	20.1	16.0	15.7	14.9	13.4	12.0	12.3	§	12.3	0.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; N is one third of N indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 192
EITHER ENERGY DRINKS OR ENERGY SHOTS: Trends in Daily Prevalence
of Use by Subgroups in Grade 10

	Percent who used daily													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	14.4	12.4	11.8	11.3	10.1	8.4	10.0	9.5	9.9	11.6	§	13.2	+1.6*
Gender:														
Male	—	17.8	15.4	14.8	13.7	12.3	9.8	11.6	10.8	12.0	14.4	§	12.0	-2.3
Female	—	10.7	9.6	8.6	8.8	8.0	6.7	7.7	7.8	7.7	8.7	§	13.6	+4.9 sss
College Plans:														
None or under 4 years	—	29.0	25.8	28.8	20.7	22.4	17.2	19.0	22.8	18.0	20.5	§	23.6	+3.1
Complete 4 years	—	12.5	10.9	10.0	10.0	8.6	7.6	8.7	7.7	8.5	9.9	§	11.2	+1.3
Region:														
Northeast	—	11.8	10.9	7.9	8.2	7.5	7.3	8.0	8.8	10.2	9.6	§	14.2	+4.6
Midwest	—	12.6	11.7	12.9	9.6	8.2	7.2	9.6	10.4	10.7	10.9	§	12.7	+1.8
South	—	16.8	15.3	12.8	12.7	11.0	8.4	10.9	7.4	9.8	11.5	§	14.2	+2.7
West	—	14.8	10.1	12.3	13.3	12.9	10.8	10.7	12.0	9.0	13.8	§	11.6	-2.3
Population Density:														
Large MSA	—	11.8	11.4	11.5	10.2	9.4	8.4	8.3	9.1	9.2	8.1	§	9.3	+1.3
Other MSA	—	14.5	11.1	10.5	10.9	8.1	8.1	10.0	8.8	9.8	12.6	§	14.1	+1.5
Non-MSA	—	17.9	16.3	15.0	13.8	16.2	9.2	13.1	11.8	11.1	15.0	§	18.4	+3.4
Parental Education: ^a														
1.0–2.0 (Low)	—	20.4	17.3	18.1	21.4	13.9	11.5	14.1	15.1	15.9	11.8	§	20.4	+8.7 ss
2.5–3.0	—	19.3	17.3	16.5	16.1	14.9	11.2	13.9	10.1	9.9	14.0	§	16.1	+2.1
3.5–4.0	—	14.6	11.8	11.3	11.5	9.4	8.1	9.6	9.5	11.0	11.4	§	14.1	+2.7
4.5–5.0	—	8.7	8.4	8.3	5.8	8.1	6.4	7.7	7.1	6.9	8.7	§	9.8	+1.1
5.5–6.0 (High)	—	9.9	9.8	7.6	7.1	5.7	5.6	5.9	5.4	7.8	9.7	§	11.3	+1.6
Race/Ethnicity (2-year data): ^b														
White	—	—	12.4	10.7	9.7	9.3	8.4	8.4	9.0	9.3	10.6	§	14.3	+3.6 ss
African American	—	—	15.8	13.4	11.8	11.9	11.7	9.9	9.4	9.7	10.3	§	13.1	+2.7
Hispanic	—	—	15.2	15.6	16.3	10.6	9.9	10.4	10.9	10.8	10.6	§	11.8	+1.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of four forms; *N* is one third of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019-2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE 193
EITHER ENERGY DRINKS OR ENERGY SHOTS: Trends in Daily Prevalence
of Use by Subgroups in Grade 12

	Percent who used daily													2019– 2021 change
	1975– 2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
<i>Approximate weighted N =</i>		14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300	
Total	—	13.5	11.0	9.9	9.1	9.3	9.0	10.9	10.9	11.2	12.8	§	14.3	+1.6*
Gender:														
Male	—	16.0	13.7	11.1	11.9	10.7	10.2	12.6	14.8	14.1	16.7	§	14.2	-2.5
Female	—	9.9	8.5	8.4	6.6	7.5	7.8	8.5	7.1	7.9	8.6	§	14.1	+5.6
College Plans:														
None or under 4 years	—	21.5	16.9	17.7	16.1	20.9	15.7	22.3	16.4	20.1	25.5	§	20.7	-4.8
Complete 4 years	—	11.8	9.7	8.2	7.2	6.3	7.3	8.4	9.2	8.7	8.7	§	12.1	+3.4
Region:														
Northeast	—	16.2	9.8	9.0	11.2	10.9	5.5	13.2	8.2	10.7	10.0	§	7.6	-2.4
Midwest	—	11.7	10.7	7.7	8.1	9.3	12.3	8.8	13.0	12.8	13.8	§	13.5	-0.3
South	—	13.4	11.9	13.6	9.2	8.6	8.3	13.6	12.7	10.8	13.1	§	16.4	+3.3
West	—	13.5	10.8	8.3	8.7	9.1	9.6	6.2	7.9	10.8	13.3	§	16.7	+3.5
Population Density:														
Large MSA	—	11.7	9.2	9.9	10.1	8.7	10.1	10.1	11.8	11.2	10.8	§	11.9	+1.1
Other MSA	—	12.6	10.9	9.5	7.8	8.8	8.6	11.0	9.3	9.7	12.1	§	13.2	+1.2
Non-MSA	—	18.1	13.6	10.8	10.8	11.1	8.2	11.6	12.7	14.8	17.7	§	20.7	+3.0
Parental Education: ^a														
1.0–2.0 (Low)	—	21.4	13.0	16.6	20.0	14.5	13.5	13.4	18.0	13.0	13.3	§	12.2	-1.2
2.5–3.0	—	19.0	13.9	11.3	8.1	14.8	10.3	12.5	12.7	14.1	12.3	§	13.1	+0.8
3.5–4.0	—	12.2	12.8	12.6	9.2	7.7	7.9	12.6	10.3	9.9	16.5	§	18.4	+1.9
4.5–5.0	—	10.7	7.4	5.7	6.3	5.2	8.0	6.3	7.7	9.1	10.3	§	11.2	+1.0
5.5–6.0 (High)	—	3.9	8.7	5.7	6.3	5.1	6.9	9.2	6.4	10.7	6.4	§	14.5	+8.1 s
Race/Ethnicity (2-year data): ^b														
White	—	—	11.6	9.1	8.1	10.0	9.7	10.0	10.4	11.1	12.4	§	13.0	+0.6
African American	—	—	11.1	13.3	12.2	11.8	10.3	11.2	12.8	10.4	10.6	§	14.3	+3.7
Hispanic	—	—	13.8	13.7	12.4	7.3	7.8	7.3	9.9	9.3	9.2	§	7.0	-2.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table S3 for the number of subgroup cases. See appendix B for definition of variables in table. Data based on one of six forms; *N* is one sixth of *N* indicated in Table S3.

Caution: Limited sample sizes (see Notes above). Use caution in interpreting subgroup trends.

§ Insufficient data for 2020 estimate.

*Changes across the 2019–2021 total samples provided here for comparison with subgroup changes for the same years.

^aParental education is an average score of mother's education and father's education. See appendix B for details.

^bEstimates for racial/ethnic subgroups combine data from current year with data from previous year to augment sample sizes and produce more reliable estimates. In 2021 the data are combined with 2020 data that are upweighted so that the two years have equal influence on the results (in 2020 MTF collected only about 25% the size of its usual sample because of pandemic-curtailed data collection).

TABLE S1
Approximate Weighted Ns by Subgroups in Grade 8

	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>
Total	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500
Gender																
Male	8,600	8,800	8,600	8,300	8,100	8,400	8,600	8,600	7,800	7,900	7,500	7,000	7,600	8,100	8,000	7,800
Female	8,600	9,300	9,200	8,600	8,700	8,800	9,300	8,900	8,400	8,300	8,300	7,600	8,400	8,500	8,400	8,200
College Plans																
None or under 4 years	2,300	2,400	2,100	2,000	1,900	2,200	1,900	1,800	1,700	1,600	1,600	1,300	1,600	1,600	1,600	1,500
Complete 4 years	14,600	15,400	15,400	14,700	14,800	14,800	15,800	15,600	14,500	14,500	14,100	13,400	14,500	15,000	14,800	14,600
Region																
Northeast	3,000	3,700	3,900	3,400	3,100	3,200	3,400	3,300	3,000	2,800	2,900	2,800	3,200	3,200	3,200	2,900
Midwest	5,300	5,300	4,700	4,200	4,300	4,600	4,100	4,300	4,200	4,300	4,000	4,000	4,100	4,000	3,700	3,500
South	6,300	6,200	6,400	6,300	6,600	6,300	7,200	6,600	6,100	6,300	5,900	5,400	6,300	6,300	6,300	6,300
West	2,900	3,400	3,300	3,400	3,500	3,700	3,900	3,900	3,400	3,300	3,400	2,900	2,900	3,500	3,600	3,800
Population Density																
Large MSA	4,500	5,700	5,500	4,400	5,200	5,200	5,000	4,800	4,800	4,900	4,700	4,500	4,900	5,000	5,400	5,400
Other MSA	8,400	8,300	8,800	8,300	7,800	8,400	9,000	8,800	7,900	7,900	7,500	6,900	7,700	7,900	7,400	7,300
Non-MSA	4,600	4,600	4,000	4,600	4,500	4,200	4,600	4,500	4,000	3,900	4,000	3,700	3,900	4,100	4,000	3,800
Parental Education																
1.0–2.0 (Low)	1,400	1,700	1,700	1,600	1,500	1,500	1,500	1,300	1,300	1,300	1,300	1,100	1,100	1,200	1,400	1,400
2.5–3.0	4,400	4,600	4,500	4,100	3,900	4,300	4,000	3,900	3,800	3,700	3,400	3,200	3,400	3,400	3,400	3,300
3.5–4.0	4,100	4,300	4,300	4,200	4,000	4,100	4,300	4,100	3,800	3,900	4,000	3,500	3,700	4,000	3,700	3,800
4.5–5.0	4,100	4,100	4,100	3,900	3,900	3,900	4,500	4,500	4,000	3,900	3,900	3,800	4,200	4,300	4,200	4,000
5.5–6.0 (High)	2,200	2,300	2,300	2,200	2,300	2,200	2,600	2,700	2,200	2,200	2,100	2,100	2,400	2,500	2,600	2,300
Race/Ethnicity (2-year average) ^a																
White	—	21,900	22,000	20,900	19,800	20,200	21,400	21,300	19,800	18,900	18,600	17,600	18,400	20,400	20,500	19,500
African American	—	4,200	4,800	5,500	5,600	5,300	4,700	4,900	5,000	4,800	4,500	4,500	4,400	3,900	3,800	4,000
Hispanic	—	3,400	3,600	4,000	4,000	4,000	4,200	4,100	4,100	4,000	4,100	3,900	3,400	3,200	3,600	4,700

Table continued on next page.

TABLE S1 (cont.)
Approximate Weighted Ns by Subgroups in Grade 8

	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>
Total	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900	15,300	14,000	13,600	3,100	10,700
Gender															
Male	7,800	7,600	7,100	7,100	7,600	7,200	7,000	6,800	6,700	8,000	7,200	6,500	6,500	§	5,100
Female	7,900	7,600	7,400	7,800	7,900	7,400	7,100	7,200	7,200	8,200	7,500	6,800	6,600	§	4,800
College Plans															
None or under 4 years	1,300	1,200	1,100	1,200	1,100	980	900	1,000	1,000	1,200	1,300	1,400	1,500	§	1,500
Complete 4 years	14,400	14,000	13,500	13,800	14,500	13,700	13,300	13,000	12,900	15,100	13,300	11,800	11,400	§	8,800
Region															
Northeast	2,400	2,600	2,500	2,500	2,900	2,800	2,500	2,400	2,200	2,800	2,600	2,500	2,400	§	1,800
Midwest	3,600	3,400	3,400	3,700	3,400	3,200	3,400	3,200	3,000	3,700	3,200	3,100	2,800	§	2,300
South	6,400	5,600	5,700	5,600	6,200	5,800	5,400	5,600	5,900	6,500	6,000	5,100	5,100	§	4,100
West	3,700	4,100	3,400	3,500	3,500	3,300	3,300	3,400	3,300	3,900	3,500	3,300	3,300	§	2,500
Population Density															
Large MSA	5,000	5,100	4,500	4,600	5,400	5,200	4,800	4,700	4,300	5,500	4,600	4,500	4,300	§	3,600
Other MSA	7,800	7,500	7,300	7,500	7,300	6,900	6,800	6,800	7,100	8,200	7,600	6,700	6,800	§	5,000
Non-MSA	3,300	3,100	3,200	3,200	3,300	3,000	3,000	3,100	3,000	3,200	3,100	2,700	2,500	§	2,100
Parental Education															
1.0–2.0 (Low)	1,400	1,300	1,200	1,400	1,300	1,300	1,400	1,300	1,400	1,600	1,600	1,400	1,200	§	600
2.5–3.0	3,100	2,800	2,700	2,800	2,700	2,500	2,400	2,400	2,500	2,800	2,600	2,400	2,100	§	1,300
3.5–4.0	3,700	3,500	3,300	3,400	3,500	3,000	2,900	3,100	2,900	3,300	3,100	2,700	2,500	§	1,900
4.5–5.0	3,900	4,100	3,900	3,800	4,100	3,900	3,900	3,600	3,500	4,300	3,600	3,400	3,500	§	2,900
5.5–6.0 (High)	2,300	2,300	2,100	2,200	2,700	2,600	2,400	2,300	2,200	2,700	2,400	2,100	2,600	§	2,300
Race/Ethnicity (2-year average) ^a															
White	18,300	17,300	16,400	16,000	16,700	16,300	14,800	13,100	12,700	13,800	13,700	12,200	11,900	§	11,600
African American	3,900	4,000	4,100	4,000	3,700	3,500	3,500	4,100	4,300	4,100	3,600	2,900	3,000	§	2,200
Hispanic	5,400	5,100	4,700	5,000	5,200	5,200	5,500	5,700	6,200	7,200	8,000	7,300	5,600	§	2,700

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

Caution: The Ns in this table are based on the entire sample at each grade level. Some drug use questions are asked only in some of the questionnaire forms rather than in all, in which case these Ns need to be adjusted appropriately. Look under Notes in each table to see if only a fraction of the sample was asked about that drug. If there is no such indication, the entire sample received the question.

^aNs for each racial subgroup represent the combination of the specified year and the previous year. Data have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE S2
Approximate Weighted Ns by Subgroups in Grade 10

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	16,200
Gender																
Male	7,200	7,000	7,300	7,700	8,300	7,500	7,400	7,100	6,300	6,800	6,600	6,900	7,500	7,900	7,900	7,900
Female	7,400	7,400	7,800	7,900	8,400	7,800	7,800	7,700	7,000	7,200	7,100	7,100	8,000	8,300	8,000	8,000
College Plans																
None or under 4 years	2,600	2,400	2,500	2,700	2,500	2,300	2,200	2,200	1,900	1,900	1,900	2,000	2,100	1,900	1,800	1,800
Complete 4 years	11,900	12,000	12,400	12,800	14,200	13,000	13,000	12,500	11,500	12,100	11,900	12,100	13,400	14,300	14,100	14,100
Region																
Northeast	2,700	3,000	2,900	3,100	3,300	3,100	3,300	3,100	3,000	2,800	2,700	2,600	3,400	3,600	3,500	3,200
Midwest	3,700	3,800	4,800	4,700	4,400	3,900	3,900	3,600	3,100	3,700	4,100	3,700	4,000	4,600	4,500	4,300
South	4,900	5,000	4,900	5,200	6,100	5,600	5,500	5,200	4,700	5,000	5,000	5,100	4,900	4,900	5,000	5,300
West	3,500	3,000	2,700	2,800	3,200	3,000	2,800	3,100	2,800	2,800	2,200	2,900	3,500	3,300	3,200	3,400
Population Density																
Large MSA	3,400	3,700	3,500	4,100	4,700	4,300	4,300	4,300	3,700	4,000	3,900	4,300	4,900	5,000	4,900	5,100
Other MSA	7,400	7,300	7,600	7,500	8,200	7,500	7,300	7,000	6,700	6,700	6,700	6,800	7,800	7,800	7,600	7,900
Non-MSA	4,000	3,800	4,200	4,200	4,100	3,800	3,900	3,700	3,200	3,600	3,400	3,200	3,100	3,600	3,700	3,200
Parental Education																
1.0–2.0 (Low)	1,300	1,300	1,300	1,300	1,200	1,100	1,300	1,300	1,100	1,200	1,200	1,300	1,200	1,300	1,100	1,100
2.5–3.0	3,900	3,900	4,100	4,100	4,100	3,600	3,700	3,700	3,200	3,200	3,200	3,300	3,500	3,400	3,200	3,200
3.5–4.0	3,900	3,900	4,100	4,300	4,600	4,300	4,100	4,000	3,600	3,700	3,700	3,700	4,200	4,200	4,100	4,300
4.5–5.0	3,500	3,400	3,500	3,700	4,000	3,900	3,700	3,500	3,300	3,500	3,400	3,500	3,900	4,300	4,400	4,500
5.5–6.0 (High)	1,800	1,700	1,700	1,800	2,300	1,900	1,900	1,800	1,700	1,900	1,800	1,700	2,100	2,400	2,400	2,300
Race/Ethnicity (2-year average) ^a																
White	—	19,600	20,700	22,000	22,900	22,400	20,900	19,800	18,400	18,200	18,600	18,000	18,500	19,900	20,400	21,100
African American	—	3,900	3,600	3,300	3,300	3,100	3,200	3,600	3,600	3,100	2,800	3,400	4,600	4,600	4,300	3,900
Hispanic	—	2,600	2,700	2,800	2,900	3,000	3,200	3,500	3,200	3,100	3,400	3,600	3,600	3,500	3,500	3,300

Table continued on next page.

TABLE S2 (cont.)
Approximate Weighted Ns by Subgroups in Grade 10

	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Total	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700	13,500	14,300	14,000	4,800	11,000
Gender															
Male	7,800	7,000	7,600	7,200	7,200	7,200	6,100	6,200	7,600	7,100	6,500	6,800	6,600	§	5,300
Female	7,900	7,700	8,000	7,700	7,400	7,500	6,500	6,400	7,600	7,200	6,600	7,000	7,100	§	5,000
College Plans															
None or under 4 years	1,800	1,600	1,500	1,600	1,500	1,300	1,300	1,200	1,400	1,400	1,400	1,800	2,000	§	1,900
Complete 4 years	14,000	13,200	14,100	13,400	13,200	13,400	11,400	11,500	13,900	12,900	11,700	12,000	11,600	§	8,700
Region															
Northeast	3,200	2,800	3,400	3,000	2,600	2,900	2,400	2,500	3,100	2,800	2,300	2,600	2,300	§	1,900
Midwest	3,900	3,600	3,700	3,700	3,800	3,800	3,200	3,000	3,600	3,200	3,100	3,300	3,200	§	2,400
South	5,100	4,800	5,200	5,200	5,100	4,900	4,100	4,400	5,500	5,100	4,800	5,200	5,300	§	4,100
West	3,900	3,900	3,600	3,300	3,400	3,400	3,200	3,100	3,400	3,600	3,300	3,100	3,200	§	2,600
Population Density															
Large MSA	5,200	4,700	5,200	5,200	4,900	4,700	3,800	4,500	5,100	4,800	4,200	4,600	4,400	§	3,800
Other MSA	7,600	7,300	7,300	6,600	6,600	7,100	6,400	6,000	7,600	7,300	6,600	6,800	7,100	§	5,000
Non-MSA	3,300	3,100	3,400	3,400	3,400	3,200	2,700	2,500	2,900	2,600	2,700	2,900	2,500	§	2,200
Parental Education															
1.0–2.0 (Low)	1,200	1,300	1,300	1,400	1,200	1,100	1,100	1,100	1,300	1,400	1,300	1,500	1,400	§	800
2.5–3.0	3,300	3,000	3,000	3,000	3,000	2,600	2,400	2,300	2,400	2,400	2,300	2,500	2,500	§	1,600
3.5–4.0	4,400	3,800	4,000	3,800	3,900	3,800	3,100	3,100	3,600	3,400	3,000	3,200	3,200	§	2,300
4.5–5.0	4,300	4,000	4,300	3,900	3,800	4,100	3,500	3,600	4,300	3,800	3,400	3,500	3,600	§	3,100
5.5–6.0 (High)	2,100	2,000	2,400	2,200	2,000	2,500	2,000	2,100	2,800	2,500	3,000	2,200	2,200	§	1,900
Race/Ethnicity (2-year average) ^a															
White	20,800	18,300	17,900	18,400	17,600	17,600	16,000	13,900	15,700	16,000	13,500	13,100	13,200	§	10,900
African American	3,400	3,600	3,600	3,400	3,200	3,100	2,900	2,900	3,000	3,300	3,700	3,800	3,700	§	2,600
Hispanic	3,800	4,500	4,500	4,600	4,500	4,100	4,100	4,300	4,400	5,100	5,300	5,400	5,700	§	4,000

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

Caution: The Ns in this table are based on the entire sample at each grade level. Some drug use questions are asked only in some of the questionnaire forms rather than in all, in which case these Ns need to be adjusted appropriately. Look under Notes in each table to see if only a fraction of the sample was asked about that drug. If there is no such indication, the entire sample received the question.

^aNs for each racial subgroup represent the combination of the specified year and the previous year. Data have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.

TABLE S3
Approximate Weighted Ns by Subgroups in Grade 12

	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
Total	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Gender																
Male	4,300	6,900	7,100	8,500	7,500	7,500	8,400	8,500	7,800	7,600	7,600	7,100	7,700	7,700	8,000	7,700
Female	5,200	7,000	7,600	9,000	8,000	7,800	8,600	8,600	8,000	7,800	8,000	7,700	8,200	8,200	8,300	7,100
College Plans																
None or under 4 years	—	6,500	6,700	8,100	6,800	6,300	6,700	7,200	6,300	5,900	5,600	5,100	5,000	4,700	4,800	4,200
Complete 4 years	—	6,800	7,200	8,600	8,000	8,500	9,700	9,200	8,800	8,900	9,300	9,100	10,300	10,600	11,000	10,100
Region																
Northeast	2,200	3,400	3,700	4,400	3,800	3,600	4,100	4,600	3,900	3,200	3,700	3,600	3,500	3,200	3,200	3,300
Midwest	2,900	4,500	4,600	5,200	4,800	4,700	5,300	5,200	4,600	4,500	4,400	4,300	4,400	4,300	4,500	4,200
South	3,000	4,300	4,600	6,000	4,800	4,800	5,300	5,300	5,200	5,300	4,900	4,700	5,200	5,600	6,100	5,000
West	1,400	2,200	2,200	2,500	2,600	2,700	2,800	2,600	2,600	2,900	3,000	2,600	3,200	3,200	2,900	2,700
Population Density																
Large MSA	2,100	3,700	4,000	4,600	4,000	3,900	4,500	4,800	4,200	4,100	4,200	3,700	4,200	4,400	4,000	3,800
Other MSA	4,000	5,700	6,200	8,000	6,800	6,700	7,100	7,300	6,800	6,900	6,900	7,000	8,000	7,700	8,800	7,700
Non-MSA	3,400	5,000	4,900	5,500	5,200	5,200	5,900	5,600	5,300	4,900	4,900	4,500	4,100	4,200	3,900	3,700
Parental Education																
1.0–2.0 (Low)	1,700	2,200	2,600	3,100	2,500	2,300	2,400	2,700	2,200	1,900	1,800	1,800	1,700	1,600	1,700	1,600
2.5–3.0	3,000	4,300	5,400	6,200	5,600	5,300	5,800	5,900	5,500	5,100	5,100	4,600	4,500	4,500	4,600	4,300
3.5–4.0	1,600	2,500	3,200	4,000	3,600	3,600	4,200	4,200	3,900	4,000	4,000	3,800	4,300	4,400	4,500	4,100
4.5–5.0	1,100	1,600	2,200	2,800	2,600	2,700	3,100	2,900	2,800	2,900	3,000	2,900	3,400	3,500	3,500	3,100
5.5–6.0 (High)	440	710	1,100	1,200	1,200	1,300	1,500	1,300	1,200	1,400	1,500	1,500	1,800	1,900	1,700	1,600
Race/Ethnicity (2-year average) ^a																
White	—	—	23,400	26,500	27,500	25,600	26,300	27,300	26,200	24,700	24,200	23,600	23,800	24,200	24,000	23,400
African American	—	—	3,300	3,700	3,500	3,500	4,000	4,000	3,900	4,000	4,000	3,500	3,200	3,600	3,900	3,500
Hispanic	—	—	890	1,000	940	740	930	1,300	1,300	1,200	1,200	1,500	1,900	2,100	2,400	2,500

Table continued on next page.

TABLE S3 (cont.)
Approximate Weighted Ns by Subgroups in Grade 12

	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>
Total	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200
Gender																
Male	7,400	7,400	7,500	6,900	7,200	6,700	7,100	7,100	6,300	5,800	5,800	5,800	6,600	6,800	6,800	6,600
Female	7,200	7,900	8,200	8,000	7,800	7,100	7,700	7,500	6,700	6,400	6,500	6,600	7,400	7,200	7,300	7,100
College Plans																
None or under 4 years	4,000	3,700	3,700	3,400	3,300	2,600	3,200	3,100	2,800	2,600	2,500	2,400	2,800	2,800	2,600	2,500
Complete 4 years	10,300	11,200	11,600	11,100	11,200	10,800	11,000	11,100	10,200	9,300	9,600	9,700	11,100	11,000	11,300	11,000
Region																
Northeast	2,800	2,800	2,700	2,700	2,800	3,000	3,300	2,800	2,500	2,500	2,400	2,500	3,100	3,100	2,900	2,600
Midwest	4,000	4,400	4,600	4,000	4,300	3,800	4,100	3,800	3,600	3,100	3,700	3,300	3,600	3,800	3,600	3,500
South	5,100	5,600	5,800	5,700	5,400	5,100	5,300	5,700	4,900	4,500	4,100	4,300	4,900	5,000	5,200	5,200
West	3,100	3,000	3,200	3,000	2,900	2,400	2,700	2,900	2,600	2,700	2,600	2,800	3,000	2,700	3,000	2,900
Population Density																
Large MSA	3,600	3,600	3,700	4,300	4,400	3,400	4,100	4,300	3,800	3,800	3,800	4,000	4,600	4,200	4,300	4,700
Other MSA	7,200	8,200	7,800	7,100	7,000	7,000	7,500	7,500	6,200	5,800	5,800	5,900	6,500	6,800	7,000	6,200
Non-MSA	4,200	4,000	4,800	4,000	4,000	3,900	3,800	3,400	3,600	3,200	3,200	3,000	3,500	3,600	3,400	3,300
Parental Education																
1.0–2.0 (Low)	1,500	1,400	1,600	1,400	1,200	1,100	1,300	1,200	960	860	1,000	980	1,200	1,000	1,100	1,100
2.5–3.0	4,100	4,100	4,300	3,700	3,700	3,300	3,600	3,700	3,200	3,000	2,900	2,800	3,400	3,400	3,200	3,100
3.5–4.0	4,200	4,600	4,500	4,300	4,400	3,800	4,100	4,300	3,900	3,600	3,600	3,800	4,200	4,000	3,900	4,000
4.5–5.0	3,100	3,400	3,600	3,500	3,700	3,500	3,500	3,300	3,200	3,100	3,200	3,100	3,400	3,600	3,600	3,600
5.5–6.0 (High)	1,500	1,700	1,700	1,800	1,800	2,100	2,100	2,000	1,800	1,600	1,600	1,500	1,800	2,000	2,100	1,900
Race/Ethnicity (2-year average) ^a																
White	21,900	21,500	22,000	21,800	21,600	20,700	19,800	20,200	19,500	17,700	16,200	16,300	17,800	19,600	19,600	18,600
African American	3,200	3,900	4,200	3,600	3,300	3,200	3,600	3,700	3,400	3,300	3,100	2,900	3,000	3,200	3,000	3,000
Hispanic	2,400	2,600	2,900	3,100	2,700	2,600	2,800	3,000	2,500	2,200	2,600	3,100	3,100	2,800	2,900	3,500

Table continued on next page.

TABLE S3 (cont.)
Approximate Weighted *N*s by Subgroups in Grade 12

	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>2020</u>	<u>2021</u>
Total	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800	12,600	13,300	12,900	3,500	8,300
Gender:															
Male	6,500	6,400	6,300	6,700	6,800	6,600	5,900	5,700	5,700	5,400	5,600	5,800	5,900	§	3,600
Female	7,400	6,900	6,700	7,100	6,700	6,600	6,100	6,100	6,300	5,600	6,100	6,400	6,300	§	4,100
College Plans															
None or under 4 years	2,400	2,300	2,100	2,100	2,100	2,100	2,200	2,000	1,900	1,900	2,000	2,300	2,500	§	1,900
Complete 4 years	11,300	10,800	10,800	11,300	11,200	10,900	9,700	9,600	10,200	9,100	9,700	9,700	9,400	§	5,900
Region															
Northeast	2,700	2,400	2,600	2,700	2,400	2,300	2,200	2,400	2,200	1,900	2,000	2,100	2,300	§	1,500
Midwest	3,200	3,500	3,500	3,200	3,400	3,600	3,100	2,500	2,700	2,500	2,800	3,000	3,000	§	1,700
South	5,500	5,000	4,800	5,400	5,100	4,400	4,300	4,700	5,100	4,700	4,800	5,100	5,400	§	3,100
West	3,100	3,100	2,800	3,100	3,200	3,400	3,000	2,800	2,900	2,700	3,000	3,100	2,200	§	2,000
Population Density															
Large MSA	4,700	3,700	3,900	4,200	4,300	4,300	3,800	3,500	4,100	3,700	4,300	4,600	4,200	§	2,800
Other MSA	6,600	7,100	6,900	7,300	6,900	6,700	6,200	6,300	6,300	5,700	5,700	6,000	6,200	§	3,800
Non-MSA	3,200	3,200	2,900	2,900	2,900	2,700	2,600	2,600	2,500	2,400	2,600	2,600	2,500	§	1,700
Parental Education															
1.0–2.0 (Low)	1,100	1,200	1,400	1,400	1,200	1,200	1,200	1,200	1,300	1,200	1,400	1,600	1,500	§	900
2.5–3.0	3,200	3,000	2,900	3,000	2,800	2,800	2,500	2,400	2,500	2,300	2,500	2,600	2,500	§	1,600
3.5–4.0	4,200	3,900	3,700	4,000	3,900	3,700	3,500	3,400	3,200	3,000	3,200	3,200	3,200	§	1,900
4.5–5.0	3,600	3,500	3,300	3,500	3,600	3,600	3,300	3,200	3,400	3,000	2,900	3,100	3,100	§	2,100
5.5–6.0 (High)	1,800	1,700	1,600	1,600	1,900	1,800	1,500	1,300	1,600	1,400	1,600	1,600	1,700	§	1,200
Race/Ethnicity (2-year average) ^a															
White	18,200	17,300	16,300	16,400	16,500	16,300	15,600	14,100	13,400	12,800	12,200	12,500	13,000	§	8,400
African American	3,200	3,500	3,100	3,000	3,400	3,200	2,700	2,800	3,300	3,400	3,200	2,800	2,500	§	1,700
Hispanic	3,700	4,000	4,400	4,500	4,000	3,700	3,900	4,000	4,200	4,000	4,500	5,700	5,700	§	3,500

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' — ' indicates data not available. See appendix B for definition of variables in table.

§ Insufficient data for 2020 estimate.

Caution: The *N*s in this table are based on the entire sample at each grade level. Some drug use questions are asked only in some of the questionnaire forms rather than in all, in which case these *N*s need to be adjusted appropriately. Look under Notes in each table to see if only a fraction of the sample was asked about that drug. If there is no such indication, the entire sample received the question.

^a*N*s for each racial subgroup represent the combination of the specified year and the previous year. Data have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B for details on how race/ethnicity is defined.