

## Catalytical Photocyclization of Arylamines with $\alpha$ -olefins in the Synthesis of 2-Alkylquinoline

Aynur Makhmutov<sup>a</sup>, Salavat Usmanov<sup>a</sup> and Ahat Mustafin<sup>b</sup>

<sup>a</sup>Birsk branch of Bashkir State University, Birsk, RUSSIA; <sup>b</sup>Bashkir State University, Ufa, RUSSIA

### ABSTRACT

The article deals with the results of investigation of the process of catalytical photocyclization of aniline and aniline hydrochloride with  $\alpha$ -olefins (hexene-1, heptene-1 and octene-1). The following compounds of d- and f-metals are tested as probable catalysts: CuSO<sub>4</sub>•5H<sub>2</sub>O, EuCl<sub>3</sub>•6H<sub>2</sub>O, PrCl<sub>3</sub>•6H<sub>2</sub>O, TbCl<sub>3</sub>•6H<sub>2</sub>O, La<sub>2</sub>O<sub>3</sub>, MnO<sub>2</sub>, NiSO<sub>4</sub>•6H<sub>2</sub>O, NiCl<sub>2</sub>•6H<sub>2</sub>O, Ni(NO<sub>3</sub>)<sub>2</sub>•6H<sub>2</sub>O, [Ni(NH<sub>3</sub>)<sub>4</sub>]Cl<sub>2</sub> and [(C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>P]<sub>2</sub>NiCl<sub>2</sub>. Catalytic effect in the reaction of photocyclization is only detected for nickel compounds. The products of catalytical photocyclization are 2-alkylquinolines. Maximum output is observed for 2-propylquinoline and is equal to 45 % at conversion of aniline of 57 %. Photosynthesis of quinolinic bases proceeds under the influence of UV-radiation at room temperature. Catalytical activity of nickel ions in a series of investigated compounds marginally depends on ligand environment. The influence of solvents' nature on the efficiency of catalytical photocyclization is detected. The output of the target product decreased in the following series: EtOH > DMSO > DMFA > Toluol > Hexane > solventless system. Catalytical photocyclization of hydrochloride aniline with  $\alpha$ -olefine in water environment relates to ecologically friendly way of 2-alkylquinolines synthesis.

### KEYWORDS

Photocyclization, catalysis, aniline, aniline hydrochloride, alkene, 2-alkylquinoline

### ARTICLE HISTORY

Received 20 August 2016  
Revised 28 September 2016  
Accepted 18 November 2016

## Introduction

Quinolinic bases are widely applied in pharmacy as active constituents in the medical products with antimalarial, antibacterial, antidiabetic and anti-inflammatory activity (Soldatenkov, Koljadina & Shendrik, 2001; Jenekhe, Lu & Alam, 2001; Liu, Tang & Zeng, 2016; Zeng & Zhou, 2016). The materials on the basis of polyquinolines synthesized from substituted quinolone bases are

**CORRESPONDENCE** Aynur Makhmutov ✉ [ainurmax@mail.ru](mailto:ainurmax@mail.ru)

© 2016 Makhmutov, Usmanov and Mustafin. Open Access terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>) apply. The license permits unrestricted use, distribution, and reproduction in any medium, on the condition that users give exact credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if they made any changes.




promising for creation of composite and electroluminescent structures (Rusanov et al., 2005), nano- and meso-structural compounds for organic electronics and nonlinear optics (García-López, Çetin & Greaney, 2015; Gulevich et al., 2012; Sarkar et al., 2015;). Some derivatives of quinolines are antioxidants (age resistors) of rubber and inhibitors of corrosion, and are consisted in cyanine dyes, extractants and sorbents (Biswas & Greaney, 2011; Li & Shi, 2016; Pintori & Greaney, 2011).

There are the methods of traditional synthesis of quinolinic nitroheterocyclic compounds (Selimov, Ptashko & Dzhemilev, 2002) and modern methods with the application of metal complex catalysts (Bulgakov et al., 2005).

In our laboratory we developed a brand new way of synthesis of alkyl-substituted quinolines by means of catalytical photocyclization (Mahmutov et al., 2016a; 2016b).

Photocatalytic transformations are of particular interest both for the science and for the practice. The most significant of them is the process of photosynthesis. The possibility of reacting under exposure to light, for example, photocatalytical destruction of water by sunlight, is one of the hottest issues of the modern times (Corrigan et al., 2016; Huang, et al., 2016; Martin, 2015; Parasram, Iaroshenko & Gevorgyan, 2014; Parasram et al., 2016).

The first example of successful photocatalytical synthesis of nitroheterocyclic compounds is the work by A. Sagadevan, A. Ragupathi & K.C. Hwang (2015). According to the authors, they managed to perform a unique reaction of synthesis of complex indole by means of catalytical photocyclization of arylamines and terminal alkynes with quinone (Figure 1).



**Figure 1.** The Scheme of Catalytical Photocyclization of Arylamines and Terminal Alkyne Benzoquinone

Detected reaction of photocyclization is catalyzed by copper chloride (I) during exposure to visible light at room temperature.

The research of catalytical photocyclization of arylamines with  $\alpha$ -olefine for direct synthesis of 2-alkylquinolines was the objective of our work. As a model system we chose the following reaction mixture: Aniline – Hexene-1 – EtOH. As catalysts we tested the following compounds of d- and f-metals:  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ,  $\text{EuCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{PrCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{TbCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{La}_2\text{O}_3$ ,  $\text{MnO}_2$ ,  $\text{NiSO}_4 \cdot 6\text{H}_2\text{O}$ ,  $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ ,  $\text{Ni}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ ,  $[\text{Ni}(\text{NH}_3)_4]\text{Cl}_2$  and  $[(\text{C}_6\text{H}_5)_3\text{P}]_2\text{NiCl}_2$ .

Influence of the solvent nature on the process of catalytical photocyclization is considered with the examples of various types of solvents: bipolar, polar and apolar. The main emphasis is laid on the choice of the water environment which

is preconditioned by the availability and cheapness of such solvent as water. Besides, the synthesis in water environment is more ecologically friendly in comparison with the application of toxic solvents.

### **Experimental Part**

The researches were conducted in the physical and chemical testing laboratory of the Birk Branch of the Bashkir State University.

Catalytical photocyclization was carried out in a photocatalytical apparatus Photo Catalytic Reactor Lelesil Innovative Systems with silica reactor having the capacity of 250 ml (photoreactor of Stromeyer's type with magnetic stir bar).

Original reagents (aniline, hexene-1, heptene-1, octene-1), catalysts ( $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ,  $\text{EuCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{PrCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{TbCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{La}_2\text{O}_3$ ,  $\text{MnO}_2$ ,  $\text{NiSO}_4 \cdot 6\text{H}_2\text{O}$ ,  $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ ,  $\text{Ni}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ ) and solvents (EtOH, DMSO, DMFA, Toluol, Hexane) are CP (chemically pure) commercial products.

Ammoniac complex  $[\text{Ni}(\text{NH}_3)_4]\text{Cl}_2$  is synthesized in accordance with the methods (Koldica, 1984) Triphenylphosphinic complex  $[(\text{C}_6\text{H}_5)_3\text{P}]_2\text{NiCl}_2$  is synthesized in glacial acetic acid in accordance with the methods (Experiment 3, 2013).

Hydrochloride aniline (phenylammonium chloride,  $\text{C}_6\text{H}_5\text{NH}_2 \cdot \text{HCl}$ ) is synthesized in accordance with the well-known methods based on the reaction of aniline with the concentrated chlorine hydride (Agranomov, 1974).

Preparation of catalysts and conduction of the process of catalytical photocyclization were performed according to the methods described in (Mahmutov et al., 2016a).

The source of ultraviolet emission was DRT-125-1 low-pressure mercury-vapor lamp. The sphere of spectral range is 230 – 400 nm at the maximum at 315 nm. Nominal light flow is 1850 lm. Ultraviolet emission reaches the reaction environment passing through the water layer. The temperature of the thermostat is 20 °C. The time of conduction of the catalytical photocyclization process is 6 hours.

For identification of the structure of the reaction products we used SHIMADZU GCMS-QP2010S gas chromatograph/mass spectrometer.

### **Evaluation**

Based on the results of the conducted researches we discovered that catalytical activity in the reaction of photocyclization of aniline (1) with hexene-1 (2) is peculiar only to nickel compounds. Catalytical activity of the other tested compounds such as:  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ,  $\text{EuCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{PrCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{TbCl}_3 \cdot 6\text{H}_2\text{O}$ ,  $\text{La}_2\text{O}_3$  and  $\text{MnO}_2$  in reaction of photocyclization is not found.

According to the data of the chromato/mass spectral analysis the main product of catalytical photocyclization in the model system of Aniline – Hexene-1 – EtOH is 2-propylquinoline (3).

General scheme of reaction of catalytical photocyclization 1 and 2 under the influence of nickel catalyst may be represented as follows (Figure 2).



The highest output of product 3 and conversion of initial 1 is observed at conduction of the reaction in ethanol and DMSO, while the lowest one is in apolar solvents.


Influence of polar solvents EtOH > DMSO > DMFA on catalytic effect is probably connected with presence of electrodonor centers (according to Pearson's classification) (Pearson, 1971). In the effective solvents there is only one such center – atom O, while in less effective DMFA there are two centers: atoms N and O.

In apolar solvents decrease in output and conversion is first connected with the disturbance of the system homogeneity. In the absence of solvent the reaction mass becomes resinified turning into the viscous mass and the general catalytic effect thus significantly decreases the output of the target product.

With the purpose of synthesis of 2-alkyl-substituted quinolines in water environment we studied catalytical photocyclization of water-soluble hydrochloride aniline (4) with alkenes: hexene-1 (2), heptene-1 (5) and octene-1 (6). As catalyst we applied crystallohydrates of nickel salts: NiCl<sub>2</sub>·6H<sub>2</sub>O, NiSO<sub>4</sub>·6H<sub>2</sub>O and Ni(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O.

The main photochemical reaction proceeds in the boundary of two liquid phases, because the reaction system is initially heterogeneous. Lower reaction layer is water solution of nickel salts and hydrochloride aniline, and the upper organic layer is the corresponding liquid alkene. The moment of the end of the process of catalytical photocyclization is clearly seen as the upper organic phase is disappearing. In the average the consumption of alkene in the chemical reaction lasts for 6-8 hours. This phenomenon will probably be used for investigation of kinetics of the catalytical photocyclization process.

The main photochemical process proceeds according to figure 3 with formation of water soluble hydrochloride 2-alkylquinoline 7-9.



**Figure 3.** Catalytical Photocyclization of Hydrochloride Aniline with Alkenes: R = C<sub>4</sub>H<sub>9</sub>- (2), C<sub>5</sub>H<sub>11</sub>- (5) and C<sub>6</sub>H<sub>13</sub>- (6); R' = C<sub>3</sub>H<sub>7</sub>- (7), C<sub>4</sub>H<sub>9</sub>- (8) and C<sub>5</sub>H<sub>11</sub>- (9),  $\gamma\Phi = UV$

The synthesized 2-alkyl-substituted quinolines 2-propylquinoline (3), 2-bulylquinoline (10) and 2-amylquinoline (11) were extracted from the reaction mixture according to the reaction of neutralization with 20% sodium carbonate solution. The output of reaction products 3, 10 and 11 according to the results of the initial vacuum fractionating is represented in table 3.

**Table 3.** Output of Substituted Quinolones at Catalytical Photocyclization of Hydrochloride Aniline with Alkenes in Water Environment(Catalyst - NiCl<sub>2</sub>·6H<sub>2</sub>O)

No	Alkene	Product of Photocyclization	Output of Product, %
----	--------	-----------------------------	----------------------



1	Hexene-1 (2)	 2-propylquinoline (3)	36
2	Heptene-1 (5)	 2-butylquinoline (10)	31
3	Octene-1 (6)	 2-amylquinoline (11)	24

The highest output is observed at the synthesis of product 3 – 36%, while the lowest – 24% for product 11. Thus the output of the photocyclization product decreases with the increase in the length of hydrocarbyl alkene.

We studied the influence of catalyst's nature, i.e. ligand environment of nickel ion on output 3 in water environment. The results are represented in table 4.

**Table 4.** Influence of Catalyst's Nature on the Output of 2-Propylquinoline in Water Environment

No	Catalyst	Output of 2-propylquinoline, %
1	NiCl <sub>2</sub>	36
2	NiSO <sub>4</sub>	34
3	Ni(NO <sub>3</sub> ) <sub>2</sub>	31
4	without catalyst	0

## Conclusions

We found that the studied series of nickel ions salts in water environment have no significant influence on the output of product 3. It is probably connected with formation of stable hydrated complexes of nickel ions in water solution free from the ligand influence of salt-forming ions. In the absence of catalyst there is no reaction between photocyclization hydrochloride aniline and hexene-1.

The studied photosynthesis of 2-alkyl-substituted quinolines in water environment is acceptable and ecologically friendly way of synthesis of substituted quinolines.

Thus, we found and studied a new way of synthesis of 2-alkylquinolines in catalytical reaction of arylamines with  $\alpha$ -olefines, proceeding according to photocyclization type during exposure to ultraviolet light. The catalysts of photocyclization are available crystallohydrates of nickel salts.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Notes on contributors

**Aynur Makhmutov**, PhD, assistant professor, Birsk branch of Bashkir state university, Birsk, Russia.

**Salavat Usmanov**, PhD, professor, Birsk branch of Bashkir state university, Birsk, Russia.

**Ahat Mustafin**, PhD, professor, Bashkir state university, Ufa, Russia.

## References

- Agranomov, A.E. (1974). *Laboratory researches in organic practicum*. Moscow: Himija.
- Biswas, K. & Greaney M.F. (2011). Insertion of arynes into thioureas: a new amidine synthesis. *Org. Lett.*, *13*(18), 4946-4949.
- Bulgakov, R.G., Kuleshov, S.P., Makhmutov, A.R. & Dzhemilev, U.M. (2006). Crystal hydrates  $\text{LnCl}_3 \cdot 6\text{H}_2\text{O}$  as highly effective catalysis in the synthesis of alkyl-substituted quinolines and phenanthrolines. *Russian Journal of Organic Chemistry*, *42*(10), 1573-1575.
- Corrigan, N., Shanmugam, S., Xu, J. & Boyer, C. (2016). Photocatalysis in organic and polymer synthesis. *Chem. Soc. Rev.*, *45*(22), 6165-6212.
- Experiment 3. (2013). *Magnetochemistry of Some Ni (II) Complexes*. Direct access: [http://www.chem.mun.ca/homes/cmkhome/Experiment\\_3\\_W2013.pdf](http://www.chem.mun.ca/homes/cmkhome/Experiment_3_W2013.pdf)
- García-López, J.A., Çetin, M. & Greaney M.F. (2015). Synthesis of hindered biaryls via aryne addition and in situ dimerization. *Org. Lett.*, *17*(11), 2649-2651.
- Gulevich, A.V., Melkonyan, F.S., Sarkar, D. & Gevorgyan, V. (2012). Double-fold c-h oxygenation of arenes using pyridine: a general and efficient traceless/modifiable silicon-tethered directing group. *J. Am. Chem. Soc.*, *134*(12), 5528-5531.
- Huang, X., Webster, R.D., Harms, K. & Meggers, E. (2016). Asymmetric catalysis with organic azides and diazo compounds initiated by photoinduced electron transfer. *Journal of the American Chemical Society*, *138*(38), 12636-12642.
- Jenekhe, S.A., Lu, L. & Alam, M.M. (2001). New conjugated polymers with donor-acceptor architectures: synthesis and photophysics of carbazole-quinoline and phenothiazine-quinoline copolymers and oligomers exhibiting large intramolecular charge transfer. *Macromolecules*, *34*, 7315-7324.
- Koldica, L. (1984). *Anorganikum*. Moscow: Mir.
- Li, Xin & Shi, Zhang-Jie. (2016). Aliphatic C-H azidation through a peroxydisulfate induced radical pathway. *Org. Chem. Front*, *3*(10), 1326-1330.
- Liu, P., Tang, J., Zeng, X. (2016). Hydrogen transfer: synthesis of  $\alpha$ -silyl benzamides. *Organic Letters*, *18*(21), 5536-5539.
- Makhmutov, A.R., Kamatov, A.Ju., Nurtdinova, R.R. & Usmanov, S.M. (2016a). Photocatalytic Condensation of Aniline with Hexene-1. *Bashkirskij himicheskij zhurnal*, *23*(1), 46-48.
- Makhmutov, A.R., Kamatov, A.Ju., Nurtdinova, R.R. & Usmanov, S.M. (2016b). Research of photocyclization arylamines with  $\alpha$ -olefins. "Natural and Mathematical Science in the modern world ": a collection of articles based on XLI international scientific-practical conference. *Novosibirsk: ANS "SibAK"*, *4*(39), 196-201.
- Martin, D.J. (2015). *Investigation into high efficiency visible light photocatalysts for water reduction and oxidation*. London: Springer.
- Parasram, M., Chuentragool, P., Sarkar, D. & Gevorgyan, V. (2016). Photoinduced formation of hybrid aryl pd-radical species capable of 1,5-hat: selective catalytic oxidation of silyl ethers into silyl enol ethers. *J. Am. Chem. Soc.*, *138*(20), 6340-6343.
- Parasram, M., Iaroshenko, V.O. & Gevorgyan, V. (2014). Endo-selective pd-catalyzed silyl methyl heck reaction. *J. Am. Chem. Soc.*, *136*(52), 17926-17929.
- Pearson, R.Dzh. (1971). Hard and soft acids and bases. *Russian chemical reviews*, *40*(7), 1259-1282.
- Pintori, D.G. & Greaney M.F. (2011). Oxidative C-H homodimerization of phenylacetamides. *Org. Lett.*, *13*(21), 5713-5715.
- Rusanov, A.L., Komarova, L.G., Prigozhina, M.P. & Lihachev, D.Ju. (2005). Polyquinolines and polyanthrazolines: synthesis and properties. *Russ. Chem. Rev.*, *74*(7), 671-683.



- Sagadevan, A., Ragupathi, A. & Hwang, K.C. (2015). Photoinduced copper-catalyzed regioselective synthesis of indoles: three-component coupling of arylamines, terminal alkynes, and quinones. *Angew. Chem.*, *54*(47), 13896-13901.
- Sarkar, D., Gulevich, A.V., Melkonyan, F.S. & Gevorgyan, V. (2015). Synthesis of multisubstituted arenes via pyrdipsi-directed unsymmetrical iterative c-h functionalizations. *ACS Catal.*, *5*(11), 6792-6801.
- Selimov, F.A., Ptashko, O.A. & Dzhemilev, U.M. (2002). *Synthesis of Natural Bases under the Influence of Metal Complex Catalysts*. Ufa: "Gilem".
- Soldatenkov, A.T., Koljadina, N.M. & Shendrik, I.V. (2001). *Basics of organic chemistry and medications*. Moscow: "Himiija".
- Zeng, X.P. & Zhou, J. (2016). Me<sub>2</sub>(CH<sub>2</sub>Cl)SiCN: bifunctional cyanating reagent for the synthesis of tertiary alcohols with a chloromethyl ketone moiety via ketone cyanosilylation. *Journal of the American Chemical Society*, *138*(28), 8730-8733.