

Testing Impacts of Goal-Oriented Outreach With the Girl Scouts: Can a Single Activity Change Attitudes Toward Insects?

Andrew J. Mongue and Kaila L. Colyott

Abstract

Most people meet insects with fear and disgust, even though few insects pose health risks; in fact, many are beneficial and their absence would adversely affect human life. Such misunderstandings lead to insect conservation being important but unpopular. We addressed these concerns as part of a broader effort to establish an ongoing outreach partnership between graduate students at the University of Kansas and the Girl Scouts of Northeast Kansas/Northwest Missouri. To explore ways to advocate for insect conservation, we held an insect-collecting activity at a Girl Scout summer camp and surveyed changes in attitudes toward insects. This activity positively changed reactions to insect encounters and increased confidence in identifying harmful insects but did not strongly reduce fears or increase curiosity toward insects. Beyond these proximate results, this project highlights the potential of Girl Scout troops as targets for informal science education that can benefit both academics and the broader community.

Keywords: university–community partnership, sustained collaboration, women in science, entomology



Insects are among the most abundant and diverse groups of organisms, accounting for over half of modern animal life on the planet (Footitt & Adler, 2009). Because of this, humans have encountered insects perhaps more than any other animal (Robinson, 1996), as shown by ancient insect remains among prehistoric settlements (Overgaard Nielsen et al., 2000; Panagiotakopulu, 2003). Further, even though we may think that modern ways of life have separated us from natural ecosystems, many insects are very successful in urban environments. Unlike larger animals that need substantial tracts of undisturbed habitats, insects can thrive in small, fragmented urban environments (e.g., parks and lawns). And unlike other urban species that are associated with low-income areas (e.g., mice; Cohn et al., 2004), insect diversity actually increases in affluent areas (Leong et al., 2016), making them a ubiquitous sight in and around homes in all communities.

In spite of, or more likely because of, this familiarity, insects are profoundly unpopular with the general public in the United States. One study found less than 10% of people enjoy insect encounters in nature, and even fewer (<1%) enjoy encountering insects in their home (Byrne et al., 1984). Another survey found that over 85% of people dislike or are afraid of arthropods (the animal phylum including insects, spiders, and crustaceans) found in the house (Hahn & Ascerno, 1991). Some of this fear and disgust may be justified in the case of disease-vector species, including mosquitoes (Beerntsen et al., 2000; Ledesma & Harrington, 2011) and kissing bugs (Prata, 2001). Researchers have in fact proposed that the disgust that is so often generalized to all insects is an evolutionary behavioral adaptation to avoid the parasitic or disease-spreading species (Curtis et al., 2004).

Although the instinct to fear and be repulsed by insects may have served our species well

historically, it is also demonstrably an over-reaction to most commonly encountered insects, the vast majority of which pose no threat to humans. A survey of biodiversity of arthropods in North Carolina households showed that most species encountered in the American home are benign (Bertone et al., 2016). For instance, only a small minority (~10%) of mosquito species are known disease transmitters (Rueda, 2008); in other words, disease vectoring is less common than one might imagine among insects that feed on humans. In fact, in other cultures, the opposite relationship exists: Some insects are seen as suitable human food. People in Japan (Mitsuhashi, 1997), Australia (Conway, 1991), and even some Native American tribes (Navarro et al., 2010), to name but a few, have historically eaten insects as a regular part of their diet; indeed, some continue to do so. Food scientists are beginning to recognize the potential of insects as a way to provide more sustainable nutrition to growing global populations (Gahukar, 2011; Van Huis, 2013). But among Europeans (Hartmann & Siegrist, 2017) and Americans (Ruby et al., 2015), there is a strong cultural disgust with the idea of eating insects, especially in an unprocessed form. Both this disgust and the generalized dislike of insects are unfounded and ignore the many beneficial services insects provide.

A Case for Insects

Setting aside the potential for insects as human food, nonpest insects play vital roles in ecosystem health and stability, most commonly by breaking down organic matter and facilitating nutrient cycling (Samways, 1994). In forests, for instance, presence of insect herbivores significantly increases available nutrients like nitrogen and phosphorus in the soil (Chapman et al., 2003), and up to 20% of wood degradation can be attributed to insects like termites and wood-boring beetles (Ulyshen, 2016). Even in human-made ecosystems, insects fill human-benefiting niches in waste decomposition, like the removal of dung from livestock pastures (Jones & Snyder, 2018), which helps promote grass growth and reduce habitat for insects that parasitize livestock (Fincher, 1981; Gillard, 1967). Even more important for the limiting of disease spread, many insects assist in the decomposition of carcasses (Matuszewski et al., 2008), a fact that also provides clues to fo-

rensic analysts in criminal cases (Buckland, 1988; Byrd, 2002).

Finally, and most popularly known, insects also pollinate many plants, including agricultural crops. The majority of crops are at least partially insect-dependent for pollination and fruit production, and crops like almonds, hay, and blueberries are completely dependent on insect pollinators (Morse & Calderone, 2000). The economic value of bee pollination alone in the United States provides services worth upwards of \$5 billion (Southwick & Southwick, 1992). Add to this the other benefits, including those described above, and the total value of all insect services to society is estimated at \$57 billion in the United States alone (Losey & Vaughn, 2006). Thus, the overwhelming fear and dislike of insects and their relatives is not only unfounded but also problematic from an economic point of view.

Generating Public Support

For all of the above reasons, there is a great need to promote acceptance and conservation of insects, but little has been done to advocate for this group of animals. Most conservation efforts focus on charismatic species, typically large mammals (e.g., whales; Scott & Parsons, 2005). Only a few well-known insects, most prominently the monarch butterfly, have received comparable attention (Diffendorfer et al., 2014; Missrie & Nelson, 2005; Oberhauser & Solensky, 2004). Given the scale of insect diversity, rather than attempting to generate case-by-case popularity, a greater ecosystem-level and, indeed, human benefit could be obtained with conservation of the broader group of arthropods, focusing on their positive contribution as members of an ecological community (Hughes et al., 2000; Panzer & Schwartz, 1998; Samways, 2007). Before tackling more comprehensive conservation efforts, however, public attitudes toward insects must be improved to ensure the success of those efforts. In this study, we sought to quantify the efficacy of single-intervention teaching in changing attitudes and reactions toward insects.

Reaction to insects, especially in the West, is a gendered response. The abovementioned study of American reactions to eating insects found women less willing to consider insects as food than men were (Ruby et al., 2015). More generally, both adult women and girls have stronger negative reactions to insects and associated invertebrates

(especially spiders; Borgi & Cirulli, 2015; Cornelius & Averill, 1983). Some research suggests that young girls are more inclined to generalize learned fear cues for invertebrates than boys are (Rakison, 2009), but both this research and the observation that both gender and cultural attitudes vary around the world suggest that these fears are learned. Thus in our attempt to change attitudes toward insects, we focused on a demographic with perhaps the worst perceptions of insects: grade-school-aged children socialized as girls.

Outreach Partnership With the Girl Scouts

Our efforts to change insect popularity grew from a broad partnership with the Girl Scouts of Northeast Kansas and Northwest Missouri. After initial successful volunteer events with local troops, we were approached by Girl Scout program managers to expand involvement and increase outreach teaching opportunities for graduate students at the University of Kansas. With a formal community partnership, graduate students designed five single-activity modules based on both their research interests and relevance to teaching objectives for Girl Scout badges. These activities were hosted on the Girl Scouts' community partner webpage, and troop leaders could then contact these graduate students to schedule an activity for their troop. This arrangement benefited both parties, as troop leaders could select the most relevant activity for the needs or interests of their Girl Scouts, and graduate students offered activities most directly relevant to their own interests and expertise. Over the course of 2 years, these programs have reached roughly 500 Girl Scouts and resulted in our programs receiving a 2018 Community Collaboration Award.

For a specific example, we, the authors, began by offering an activity to help Brownie Scouts meet requirements to earn their Bugs Badge. The badge has multiple components, ranging from insect-themed arts and crafts to exploration of insect habitats. We focused on the latter, showing Girl Scouts where and how to collect local insects. This activity was one of the more popular, being requested by 166 Girl Scouts in total. Owing to the qualitative change in attitudes we noticed in Girl Scouts who participated in these activities in the first year of our partnership, we designed a simple survey to test whether a single activity interacting with arthropods

could reduce fear and increase appreciation of local insect species.

We obtained Institutional Review Board approval for study design and consent language from the University of Kansas (IRB ID: 00141007) and carried out the survey at a Girl Scout summer camp in summer 2017. We informed the parent or guardian of each participating child upon their arrival at camp that their child was in a camp group connected to a research study. We gave the parent or guardian a verbal summary of the project and a paper copy of the survey to review before asking them to sign a consent form allowing their child to participate in the study. Parents had the option of opting out of the study by not signing the consent form, without affecting their child's ability to participate in camp activities, including insect collection. Children without parental or guardian consent were not given a survey to complete, and no identifying information was collected for any child during the survey process.

We administered the survey to groups of Junior and Cadette rank Girl Scouts for 2 months in summer 2017 at Camp Daisy Hindman, in rural Dover, Kansas ($n = 88$ total respondents). To minimize identifiable information collected, we did not record ages of participants, but these ranks correspond to fourth to eighth grade students. To be clear, we did not ask, nor do we make proscriptive assumptions about the gender of any individual scout in our study, but the Girl Scouts as a group is surely enriched for individuals socialized as girls. Throughout the summer camp season (June–July), we visited the camp each week and collected data from two groups of Girl Scouts. Each week, one group worked with us on an insect-collecting activity before taking a survey of attitudes and reactions toward insects. The second (control) group had no interaction with us prior to the survey. Collecting activities varied by week (blacklight trapping, stream collecting, or open field sweep netting) depending on the camp program and weather, but in each activity Girl Scouts collected insects and transferred them from a net to a mesh cage by hand. Throughout the activity, we encouraged Girl Scouts to share their discoveries and help each other with collecting. With the Girl Scouts' consent, we saved representative specimens of collected species to be pinned and spread by us as part of a display kept at the camp.

For each activity, we used a teaching collection of pinned insects to facilitate a short discussion (~10 minutes) that included an overview of stinging insects and an open-ended question and answer session. Immediately after the hands-on collecting session, we spent a short time (~5 minutes) asking Girl Scouts to share their favorite catches. The Girl Scouts for whom we had obtained prior parental consent were then given a survey to fill out. For the control group, Girl Scouts were given surveys immediately after completing their regularly scheduled camp activities (e.g., tie-dyeing or horseback riding) with no collecting activity or discussion of insects. Camp groups that were chosen as control groups were selected to keep the number of participants and ages roughly equal between the treatment and control.

Survey Content and Analysis

The anonymous surveys consisted of 15 questions, with three background questions and 12 retrospective before/after questions that asked participants to answer how they felt both before and after their time at the summer camp (full survey shown in Table 1). The use of a retrospective pretest-posttest design (i.e., administering both the pre- and posttest questions after the intervention) provides a more accurate assessment of change than a conventional pretest-posttest design (i.e., administering pretest before and posttest after) because it enables the respondent to use a consistent scale when answering questions about both the present and past (Nakonezny & Rodgers, 2005). Additionally, scheduling a single block of time for survey completion minimized the disruption to the children's other camp activities and facilitated scheduling with the counselors.

Most questions were based on a Likert-like scale of responses (e.g., *always, often, sometimes, never*), but the reactions to the encounter questions were subjectively ranked from least desirable to most: killing the insect or spider (fearful and destructive), running away (fearful and passive), ignoring it (neutral/nondestructive), moving it outside (active and unafraid). Background questions were implemented as a check to ensure no systematic differences existed in everyday exposure to arthropods between our treatment and control groups. The 12 retrospective before/after questions were also designed in pairs for control: One

set asked about attitudes toward insects ("bugs" in the survey) and the other asked about spiders, which were not a part of the hands-on teaching or open-ended discussion, but are often cited in popular culture and scientific literature as an invertebrate that women fear more than men do. The spider questions thus created an additional check that time spent at camp was not changing attitudes about invertebrates in general by virtue of bringing campers closer to nature than they would be at home.

Prior to downstream analyses, we curated the data for irregularities. A small minority of Girl Scouts skipped questions, chose multiple answers to a single question, or answered in a manner seemingly contrary to the design of the experiment (i.e., individuals from the control group indicating that they worked with us, despite no interaction at camp prior to the survey). This last class of problems was rare but potentially confounding, as we had been doing community outreach workshops for the 2 preceding years in the area, so some Girl Scouts in our control groups may have had previous experience with our teaching outside the scope of this project. To be conservative in analyses, we discarded both of the surveys that had the uncertain treatment status described above; this curation brought our sample size down from 88 to 86 (45 treatment, 41 control). For the remaining irregularities, answers were excluded on a case-by-case basis (e.g., a Girl Scout who skipped or gave multiple answers to Question 3 would still have their answers to Questions 4–15 included in analyses), resulting in slight differences in sample sizes between questions. We coded each potential response to a question as a number from 0 to 3 for analysis. Although these data are not continuous and not necessarily normally distributed, parametric tests should be robust to these assumptions, given our sample sizes surveyed. Thus we assessed simple differences in the treatment and control groups with *t*-tests for the background questions.

For the focal questions, however, we needed to consider the compound effects of both our direct intervention and the Girl Scouts' broader camp experiences changing attitudes on their own; this design defies a simple *t*-test approach. To parse the more complicated effect of our lessons while controlling for time at camp, we analyzed results in a permutation framework that

Table 1. Survey Questions Presented to Girl Scouts

1. Did you spend time working with the bug people at camp?	YES	NO		
2. How often do you encounter bugs at home?	Always	Often	Sometimes	Never
3. How often do you encounter spiders at home?	Always	Often	Sometimes	Never
4. How afraid were you when encountering bugs at home BEFORE coming to camp?	Very afraid	Somewhat	Not very	Not afraid
5. How afraid were you when encountering spiders at home BEFORE coming to camp?	Very afraid	Somewhat	Not very	Not afraid
6. BEFORE coming to camp, when you encountered a bug at home, what would you do?	Kill it	Run away	Ignore it	Move it outside
7. BEFORE coming to camp, when you encountered a spider at home, what would you do?	Kill it	Run away	Ignore it	Move it outside
8. How likely were you to pick up a bug and be curious about it at home, BEFORE coming to camp?	Always	Often	Sometimes	Never
9. How good do you think you were at determining if a bug was dangerous or not, BEFORE coming to camp?	Great	Good	OK	Not good
10. How afraid are you of encountering bugs at home AFTER coming to camp?	Very afraid	Somewhat	Not very	Not afraid
11. How afraid are you of encountering spiders at home AFTER coming to camp?	Very afraid	Somewhat	Not very	Not afraid
12. AFTER coming to camp, when you encountered a bug at home, what will you do?	Kill it	Run away	Ignore it	Move it outside
13. AFTER coming to camp, when you encountered a spider at home, what will you do?	Kill it	Run away	Ignore it	Move it outside
14. How often do you think you will pick up a bug and be curious about it, AFTER coming to camp?	Always	Often	Sometimes	Never
15. How good do you think you are at determining if a bug is dangerous or not, AFTER coming to camp?	Great	Good	OK	Not good

Note. Participants were asked to circle an answer to each question either after the insect activity (treatment) or immediately upon gathering after a non-insect-related activity (controls).

made no assumptions about underlying data distributions. First we calculated the difference in means before and after time at camp in the treatment and control groups separately. Then we calculated the difference of these differences to get a measure of how dissimilar the treatment and control groups were while controlling for general experiences at camp. To assess significance of these differences, we then carried out permutations randomly assigning Girl Scouts

to treatment or control groups of sizes equal to the true groups. As before, we calculated the difference of differences between our pseudotreatment and pseudocontrol groups. By repeating this for 1,000 permutations, we generated an expectation of differences between groups that could occur by chance. We then compared our true value to this distribution; the p -value was taken as the proportion of times that the true difference was more extreme than the randomly gen-

erated differences. Consequently, there are no test statistics per se to report for these analyses, only p -values. Questions with significant shifts in attitude are shown in the figures that follow, with counts for each answer in both treatment and control groups. All analyses and data visualizations were carried out using custom scripts written in R version 3.4.1.

Findings

The control and treatment groups did not differ in exposure to insects at home (Question 2; $t_{78.1} = 0.19$, $p = 0.85$), but, oddly, they reported a difference in spider encounters, with the control group encountering fewer spiders (Question 3; $t_{80.5} = 2.41$, $p = 0.02$). This starting difference is less relevant for our focus on insects, and moreover appears to have no biasing effect, as groups did not differ from each other in their reaction to ($p = 0.555$, Questions 7 & 13) or fear of ($p = 0.293$, Questions 5 & 11) spiders while controlling for time at camp.

With regard to insects, two of our metrics showed significant changes in our treatment group after the activity. Girl Scouts became more confident in being able to identify dangerous insects ($p = 0.018$, Questions 9 & 15) and became less likely to kill or run away from an insect encountered at home ($p = 0.041$, Questions 6 & 12). Results with sample sizes can be seen in Figure 1A and Figure 1B respectively. Our two other metrics, curiosity (Questions 8 & 14) and fear of insects (Questions 4 & 10) did not significantly change after our lesson, but did trend in the direction of more curiosity ($p = 0.099$) and less fear ($p = 0.180$). In the latter case, both treatment and control groups reported marginal decreases in fear after their time at camp. These results, including sample sizes, are summarized in Figure 2A and Figure 2B.

Discussion

The public in general dislikes insects more than most animals (Byrne et al., 1984), and women more than men find them disgusting (Curtis et al., 2004). In a targeted effort to change these attitudes, we designed a set of activities to carry out with the Girl Scouts of Camp Daisy Hindman. Afterward, we surveyed Girl Scouts to assess how effective activities involving live insects can be in changing perceptions. In brief, we found that we increased confidence in differenti-

ating dangerous and harmless insects and positively changed self-reported reactions to an insect encounter. Girl Scouts became less likely to kill insects encountered in the home and more likely to move them outside or ignore them. Both of these results are encouraging for the goal of increasing acceptance of insects. Decreasing instances of immediately killing an insect found in the home can only help conserve insects, and learning to confidently differentiate dangerous and harmless insects should lead to fewer perceived threats from encounters with harmless insects. Whether or not this effect persists in the long term would be an obvious target for future study.

On the other hand, we failed to strongly increase curiosity or decrease fear of insects in the span of this activity. However, both of these metrics showed small changes in the desired direction after our lesson, so it is possible that this hands-on approach could be effective but would require more engagement time to generate strong changes. As with the positive results, it would also be fruitful to examine the fear and curiosity components after repeated interactions.

Another possibility is that we failed to identify fears in a precise enough way to notice a change. Girl Scouts who participated in our activity reported becoming more confident in identifying dangerous insects but did not report a decrease in fear of “bugs” as a blanket category. Considering these two outcomes together, it would be interesting to ask about fears of specific groups of insects to see if fears become less generalized and more concentrated on groups that can cause harm, like ants and wasps. We avoided such detailed questions in this initial survey out of a desire to keep the survey portion short and easy to complete, but it would be appropriate for a more targeted follow-up study.

More generally, our activity sparked engagement in spite of using an unpopular group of animals, suggesting a great potential to stimulate excitement with this age group. The surveyed scouts were late elementary school to middle school aged, the time when girls become less likely than boys to pursue interests in sciences (Blickenstaff, 2005); similarly, hands-on approaches with Girl Scouts make an obvious target for promoting women in STEM (science, technology, engineering, and math). Finally, we recognize that our work as well as the work cited here is limited to





Figure 1A and 1B. Metrics Showing Significant Change in Treatment Group

Note. 1A. Girl Scouts who participated in the insect-catching activity became more confident ($p = 0.018$) in identifying dangerous insects (top panel) than their control counterparts (bottom panel). 1B. Likewise, Girl Scouts from the treatment group became more likely to move an insect outside and less likely to kill it or run away ($p = 0.041$) after the lesson. Each response category is color coded per the legend, and numbers in each category represent counts of respondents.





Figure 2A and 2B. Metrics Showing Lack of Significant Change in Treatment Group

Note. 2A. The difference in curiosity and willingness to pick up insects for both treatment and control groups after treatment was not significant ($p = 0.099$) but trended toward an increase in curiosity, as can be seen in the top panel. 2B. Our activity did not decrease respondents' fears of insects ($p = 0.180$) in any meaningful way. As with the other figure, each response category is color coded per the legend, and numbers in each category represent counts of respondents.

binary gender categories (i.e., women and men, girls and boys). We suggest that future research would benefit from being more gender inclusive (i.e., nonbinary and other minority gender individuals) in surveying, especially given that these groups have been historically overlooked when promoting inclusion in STEM.

Lessons Learned: Graduate Student-Girl Scouts Partnership as a Mutually Beneficial Relationship

We found that our regional Girl Scout organization offers a receptive audience for informal STEM education, and we suggest that they make an excellent venue for outreach across the sciences. By advertising our STEM expertise and taking requests for outreach teaching, we were able to match our science skillset with a demand in the community. This two-way interaction sparked our initial interest in formally assessing the effectiveness of our outreach activities, and we submit that this approach can serve as a useful model for goal-oriented outreach among academic researchers.

Although such outreach may be more common among other educational groups, for many research-focused scientists, outreach remains an unorganized endeavor. Developing broad community impacts is an important component of many academic positions, but it often receives less attention than research or formal (i.e., classroom-based) teaching. We submit that outreach

can and should be approached in the same manner as the rest of the scientific process: with concrete objectives and empirical validation to assess how successfully these objectives are met. Under this paradigm, outreach events are more beneficial to both the researchers and the public. Researchers can have meaningful interactions and encourage interest in science, as we saw with scouts' confidence in insect identification and decreased inclination to kill insects in this study.

Next Steps

The established structure and persistent groups of Girl Scout troops make excellent partners for repeated scientific engagement across multiple years. Anecdotally, we have seen some of the same children at multiple outreach events, but, due to our limited time in graduate school, we are not able to formally track the longer term impacts of our activities on either interest in science or attitudes toward insects. Although we, the authors, have since graduated, we are happy to report that the Girl Scout partnership still exists with current graduate students at the University of Kansas and continues to offer a platform for informal STEM teaching. In its current incarnation, the partnership consists of the continued independent modules as well as an annual STEM activity day at one of the camps (Camp Tongawood); current graduate students have plans to use this venue for outreach outcome surveying.



About the Authors

Andrew J. Mongue is a postdoctoral researcher in the Institute of Evolutionary Biology at the University of Edinburgh.

Kaila L. Colyott is a program coordinator in the Center for Teaching Excellence at the University of Kansas.

References

- Beerntsen, B. T., James, A. A., & Christensen, B. M. (2000). Genetics of mosquito vector competence. *Microbiology and Molecular Biology Reviews*, 64(1), 115–137. <https://doi.org/10.1128/MMBR.64.1.115-137.2000>
- Bertone, M. A., Leong, M., Bayless, K. M., Malow, T. L. F., Dunn, R. R., & Trautwein, M. D. (2016). Arthropods of the great indoors: Characterizing diversity inside urban and suburban homes. *PeerJ*, 4, Article e1582. <https://doi.org/10.7717/peerj.1582>
- Blickenstaff, J. C. (2005). Women and science careers: Leaky pipeline or gender filter? *Gender and Education*, 17(4), 369–386. <https://doi.org/10.1080/09540250500145072>
- Borgi, M., & Cirulli, F. (2015). Attitudes toward animals among kindergarten children: Species preferences. *Anthrozoös*, 28(1), 45–59. <https://doi.org/10.2752/089279315X14129350721939>
- Buckland, P. C. (1988). [Review of *A manual of forensic entomology*, by K. G. V. Smith]. *American Journal of Archaeology*, 92(2), 287. <https://doi.org/10.2307/505635>
- Byrd, J. H. (2002). *Forensic entomology: The utility of arthropods in legal investigations*. CRC Press.
- Byrne, D. N., Carpenter, E. H., Thoms, E. M., & Cotty, S. T. (1984). Public attitudes toward urban arthropods. *Bulletin of the ESA*, 30(2), 40–44. <https://doi.org/10.1093/besa/30.2.40>
- Chapman, S. K., Hart, S. C., Cobb, N. S., Whitham, T. G., & Koch, G. W. (2003). Insect herbivory increases litter quality and decomposition: An extension of the acceleration hypothesis. *Ecology*, 84(11), 2867–2876. <https://doi.org/10.1890/02-0046>
- Cohn, R. D., Arbes, S. J., Yin, M., Jaramillo, R., & Zeldin, D. C. (2004). National prevalence and exposure risk for mouse allergen in US households. *Journal of Allergy and Clinical Immunology*, 113(6), 1167–1171. <https://doi.org/10.1016/j.jaci.2003.12.592>
- Conway, J. R. (1991). The biology and aboriginal use of the honeypot ant, “*Camponotus inflatus*” Lubbock, in Northern Territory, Australia. *Australian Entomologist*, 18(2), 49–56.
- Cornelius, R. R., & Averill, J. R. (1983). Sex differences in fear of spiders. *Journal of Personality and Social Psychology*, 45(2), 377–383. <https://doi.org/10.1037/0022-3514.45.2.377>
- Curtis, V., Aunger, R., & Rabie, T. (2004). Evidence that disgust evolved to protect from risk of disease. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271(suppl_4), S131–S133. <https://doi.org/10.1098/rsbl.2003.0144>
- Diffendorfer, J. E., Loomis, J. B., Ries, L., Oberhauser, K., Lopez-Hoffman, L., Semmens, D., Semmens, B., Butterfield, B., Bagstad, K., Goldstein, J., Wiederholt, R., Mattsson, B., & Thogmartin, W. E. (2014). National valuation of monarch butterflies indicates an untapped potential for incentive-based conservation. *Conservation Letters*, 7(3), 253–262. <https://doi.org/10.1111/conl.12065>
- Fincher, G. T. (1981). The potential value of dung beetles in pasture ecosystems. *Journal of the Georgia Entomological Society*, 16, 316–333.
- Footitt, R. G., & Adler, P. H. (Eds.). (2009). *Insect biodiversity: Science and society*. Wiley Online Library. <https://doi.org/10.1002/9781444308211>
- Gahukar, R. (2011). Entomophagy and human food security. *International Journal of Tropical Insect Science*, 31(3), 129–144. <https://doi.org/10.1017/S1742758411000257>
- Gillard, P. (1967). Coprophagous beetles in pasture ecosystems. *Journal of the Australian Institute of Agricultural Science*, 33(1), 30–34. <http://hdl.handle.net/102.100.100/324003?index=1>
- Hahn, J. D., & Ascerno, M. E. (1991). Public attitudes toward urban arthropods. *American Entomologist*, 37(3), 179–185. <https://doi.org/10.1093/ae/37.3.179>
- Hartmann, C., & Siegrist, M. (2017). Insects as food: Perception and acceptance: Findings from current research. *Ernährungs Umschau*, 64(3), 44–50. <https://doi.org/10.4455/eu.2017.010>

- Hughes, J. B., Daily, G. C., & Ehrlich, P. R. (2000). Conservation of insect diversity: A habitat approach. *Conservation Biology*, 14(6), 1788–1797. <https://doi.org/10.1111/j.1523-1739.2000.99187.x>
- Jones, M. S., & Snyder, W. E. (2018). Beneficial insects in agriculture: Enhancement of biodiversity and ecosystem services. *Insect biodiversity: Science and society* (vol. 2, pp. 105–122). Wiley Blackwell.
- Ledesma, N., & Harrington, L. (2011). Mosquito vectors of dog heartworm in the United States: Vector status and factors influencing transmission efficiency. *Topics in Companion Animal Medicine*, 26(4), 178–185. <https://doi.org/10.1053/j.tcam.2011.09.005>
- Leong, M., Bertone, M. A., Bayless, K. M., Dunn, R. R., & Trautwein, M. D. (2016). Exoskeletons and economics: Indoor arthropod diversity increases in affluent neighbourhoods. *Biology Letters*, 12(8). <https://doi.org/10.1098/rsbl.2016.0322>
- Losey, J. E., & Vaughn, M. (2006). The economic value of ecological services provided by insects. *BioScience*, 56(4), 311–323. [https://doi.org/10.1641/0006-3568\(2006\)56\[311:TEVOES\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)56[311:TEVOES]2.0.CO;2)
- Matuszewski, S., Bajerlein, D., Konwerski, S., & Szpila, K. (2008). An initial study of insect succession and carrion decomposition in various forest habitats of Central Europe. *Forensic Science International*, 180(2–3), 61–69. <https://doi.org/10.1016/j.forsciint.2008.06.015>
- Missrie, M., & Nelson, K. (2005). Direct payments for conservation: Lessons from the Monarch Butterfly Conservation Fund. *Economics*, 3(88), 339–353.
- Mitsuhashi, J. (1997). Insects as traditional foods in Japan. *Ecology of Food and Nutrition*, 36(2–4), 187–199. <https://doi.org/10.1080/03670244.1997.9991514>
- Morse, R. A., & Calderone, N. W. (2000, March). The value of honey bees as pollinators of US crops in 2000. *Bee Culture*, 128, 1–15.
- Nakonezny, P. A., & Rodgers, J. L. (2005). An empirical evaluation of the retrospective pretest: Are there advantages to looking back? *Journal of Modern Applied Statistical Methods*, 4(1), 240–250. <https://doi.org/10.22237/jmasm/1114906920>
- Navarro, J. C. A., Prado, S. M. C., Cardenas, P. A., Santos, R. D., & Caramelli, B. (2010). Pre-historic eating patterns in Latin America and protective effects of plant-based diets on cardiovascular risk factors. *Clinics*, 65(10), 1049–1054. <https://doi.org/10.1590/S1807-59322010001000022>
- Oberhauser, K. S., & Solensky, M. J. (2004). *The Monarch butterfly: Biology & conservation*. Cornell University Press.
- Overgaard Nielsen, B., Mahler, V., & Rasmussen, P. (2000). An arthropod assemblage and the ecological conditions in a byre at the Neolithic settlement of Weier, Switzerland. *Journal of Archaeological Science*, 27(3), 209–218. <https://doi.org/10.1006/jasc.1999.0448>
- Panagiotakopulu, E. (2003). Insect remains from the collections in the Egyptian Museum of Turin. *Archaeometry*, 45(2), 355–362. <https://doi.org/10.1111/1475-4754.00113>
- Panzer, R., & Schwartz, M. W. (1998). Effectiveness of a vegetation-based approach to insect conservation. *Conservation Biology*, 12(3), 693–702. <https://doi.org/10.1111/j.1523-1739.1998.97051.x>
- Prata, A. (2001). Clinical and epidemiological aspects of Chagas disease. *The Lancet: Infectious Diseases*, 1(2), 92–100. [https://doi.org/10.1016/S1473-3099\(01\)00065-2](https://doi.org/10.1016/S1473-3099(01)00065-2)
- Rakison, D. H. (2009). Does women's greater fear of snakes and spiders originate in infancy? *Evolution and Human Behavior*, 30(6), 438–444. <https://doi.org/10.1016/j.evolhumbehav.2009.06.002>
- Robinson, W. H. (1996). *Urban entomology: Insect and mite pests in the human environment*. Chapman & Hall.
- Ruby, M. B., Rozin, P., & Chan, C. (2015). Determinants of willingness to eat insects in the USA and India. *Journal of Insects as Food and Feed*, 1(3), 215–225. <https://doi.org/10.3920/JIFF2015.0029>
- Rueda, L. M. (2008). Global diversity of mosquitoes (Insecta: Diptera: Culicidae) in freshwater. *Hydrobiologia*, 595(1), 477–487. <https://doi.org/10.1007/s10750-007-9037-x>

- Samways, M. J. (1994). *Insect conservation biology* (Conservation Biology No. 2). Springer Science & Business Media.
- Samways, M. J. (2007). Insect conservation: A synthetic management approach. *Annual Review of Entomology*, 52, 465–487. <https://doi.org/10.1146/annurev.ento.52.110405.091317>
- Scott, N. J., & Parsons, E. C. M. (2005). A survey of public opinion in south-west Scotland on cetacean conservation issues. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 15(3), 299–312. <https://doi.org/10.1002/aqc.662>
- Southwick, E. E., & Southwick, L. (1992). Estimating the economic value of honey bees (Hymenoptera: Apidae) as agricultural pollinators in the United States. *Journal of Economic Entomology*, 85(3), 621–633. <https://doi.org/10.1093/jee/85.3.621>
- Ulyshen, M. D. (2016). Wood decomposition as influenced by invertebrates. *Biological Reviews*, 91(1), 70–85. <https://doi.org/10.1111/brv.12158>
- Van Huis, A. (2013). Potential of insects as food and feed in assuring food security. *Annual Review of Entomology*, 58, 563–583. <https://doi.org/10.1146/annurev-ento-120811-153704>