

Potential of morphology-based Pterydophyta diversity in supporting field-based practicum of low plant botany learning

Y. Yudhastian^{a,1}, Tabitha Sri Hartati Wulandari^{a,2,*}

^a Department of Biology Education, Faculty of Teacher Training and Education, Universitas PGRI Ronggolawe, Jl. Manunggal 61 Tuban, East Java 62391. Indonesia

¹yudhastianadis@gmail.com; ²tabithawulandari7@gmail.com*

Abstract: Field-Based Practicum (FBP) about Pterydophyta diversity material in Low plant botany learning is very important, even though the facts in the field show that FBP is still minimally carried out. This research aims to utilize the potential diversity of Pterydophyta in the Tuban-Lamongan Pantura area as a support for FBP about low plant botany learning. The type of research is descriptive exploratory with a transect method, which divides the research area into five plots. Data collection techniques involve observing abiotic parameters and fern morphology in each plot and counting the number of individuals of each species. Data analysis used the Shannon & Wiener index for diversity and qualitative descriptive analysis for morphology. The research results showed 19 species of ferns with 210 individuals divided into two classes, namely Polypodiopsida and Psilotopsida. The fern diversity index is in the medium category ($H' = 2.10$). The diversity of ferns obtained shows that the Tuban-Lamongan Pantura area has a high potential to support FBP about low plant botany learning to provide insight and direct experience to students about the diversity of ferns in the surrounding environment.

Keywords: diversity; field-based practicum; morphology; Pterydophyta

*For correspondence:

tabithawulandari7@gmail.com

Article history:

Received: 26 April 2023

Revised: 9 July 2024

Accepted: 13 July 2024

Published: 14 July 2024

 10.22219/jpbi.v10i2.33263

© Copyright Yudhastian *et al.*

This article is distributed under the terms of the Creative Commons Attribution License



p-ISSN: 2442-3750

e-ISSN: 2537-6204

How to cite:

Yudhastian, Wulandari, T.S.H. (2024). Potential of morphology-based Pterydophyta diversity in supporting field-based practicum of low plant botany learning. *JPBI (Jurnal Pendidikan Biologi Indonesia)*, 10(2), 497-511. <https://doi.org/10.22219/jpbi.v10i2.33263>

Introduction

Forest vegetation ecosystems in Indonesia have abundant flora diversity, one of which is the diversity of Pterydophyta (Azrai et al., 2024; Deng et al., 2024; Juliasih & Adnyana, 2023b; Qian et al., 2023). Pterydophyta is highly diverse in Indonesia and is widely studied in terms of morphology (Azrai et al., 2024; Pramudita et al., 2023). Pterydophyta (ferns) is the oldest dominant vascular plant group without seeds and flowers that have free spores with a diversity of more than 13.500 species worldwide (De-Villa & Lagat, 2023; Di-Michele et al., 2024; Guillen-Otero et al., 2024; Ibiye et al., 2023; Katayama et al., 2024; Marasinghe & Daulagala, 2024; Michalak et al., 2024; Randrianarimanana et al., 2024; Xie et al., 2023). Ferns are classified in Equisetales, Psilotopsida, Marattiopsida, and Polypodiopsida (Kumar et al., 2022). The diversity of ferns in the form of morphology, global distribution, and phylogenetic distribution is mainly found on the top of tropical mountains with humidity and temperature ranging from 21-27°C (Fujiwara et al., 2023; Juliasih & Adnyana, 2023a; Khan et al., 2024; Michel et al., 2023; Moulatlet et al., 2024; Qian et al., 2024). Ferns are known as cosmopolitan plants that can proliferate horizontally and vertically in water (hydrophytes), attach to other plants (epiphytes), and grow on the rest of dead plants (saprophytes) (Afifah et al., 2023; Rahmawati & Santhyami, 2023; Wheeler et al., 2024). The body parts of ferns are divided into roots, stems, and leaves (Khotimah et al., 2023). Ferns are an essential component of the ecosystem due to their rapid photosynthetic ability and their ability to increase microbial diversity in the soil (Lu et al., 2023; Riyanto et al., 2024; Yi et al., 2023; Zeng et al., 2024). In its development, the diversity of Pterydophyta can be utilized to support 21st-century learning models, one of which is to support Field-Based Practicum (FBP) learning of Low plant botany. This is in alignment with the research findings of Boileau and O'Donoghue (2024) who emphasized the importance of providing students with outdoor learning experiences. Practicum is one of the learning methods that can improve student understanding (Yuhanafia et al., 2024). Practicum is considered the best innovation for educators to apply theory into practice and is

essential in improving the quality of learning outcomes and processes (Li et al., 2023; Lolowang et al., 2023). In its implementation, practicum requires cognitive, interpersonal, creativity, collaboration, and critical thinking skills (Mubai et al., 2023). However, several factors cause practicum to be rarely implemented, such as the relatively long preparation time for practicum, the lack of availability of practicum equipment, and the high level of risk of practicum in the laboratory (Juniar et al., 2024). This is because the quality of practicum learning in Indonesia still needs to catch up to the development of existing industries (Suputra & Susanto, 2023). Therefore, FBP is one of the innovative learning methods that can be given to students to improve learning outcomes; this is because students understand the material directly and more meaningfully (Das et al., 2024; Sejati et al., 2023; Tariq et al., 2023). Through observation, FBP is conducted outside the laboratory, which benefits students by providing learning flexibility (Kandriasari et al., 2023; Nguyen & Nguyen, 2024). In addition, FBP can help students apply theoretical knowledge in the classroom in actual practice through skill development (Hande & Sharma, 2021; Joseph & Jamal, 2023; Lee & Davis, 2023).

So far, the FBP of Low plant botany learning material on Pterydophyta diversity is minimal. This is evidenced by several studies in various countries that only identify the diversity of Pterydophyta without integrating it into 21st-century learning, such as Ibrahim et al. (2024) in the Gunung Masigit Hunting Park area of Bandung; Dubuisson et al., (2024) in Africa; Fischer and Lobin (2024) in Rwanda; Michel et al. (2023) in the lowland tropical rainforest of Ecuador; Yoneoka et al. (2023) in regional Japan and Zhang et al. (2023) in Zhejiang Province. As a result, learning resources to support the FBP of learning Low plant botany are rarely found, so students also need help finding contextual learning resources that present the diversity of Pterydophyta. This directly impacts students' low understanding of Pterydophyta diversity and their inability to solve contextual problems critically. Therefore, the novelty of this research is to explore the potential diversity of Pterydophyta in the Tuban-Lamongan Pantura area based on morphology. This is because, so far, research has yet to be conducted to study the diversity of ferns in the Tuban-Lamongan coastal area with a morphological approach. In comparison, the study of morphology-based pteridophyta diversity is beneficial to support FBP in learning low-level plant botany. Our next novelty also utilizes the potential diversity of Pterydophyta to support the FBP of low-level plant botany learning as well as a form of integration in 21st-century learning. This FBP is expected to give students an in-depth understanding of Pterydophyta diversity and improve their critical contextual problem-solving skills.

Based on the description related to the diversity of Pterydophyta and the need for field-based practicum to support 21st-century learning models, the formulation of the problem in this study is how to utilize the potential diversity of Pterydophyta in the Tuban-Lamongan Pantura area based on morphology in supporting field-based practicum learning of low plant botany? The research objectives are to utilize the potential diversity of Pterydophyta in the Pantura Tuban-Lamongan area based on morphology in supporting field-based practicum learning of low plant botany.

Method

This is an explorative and descriptive research that describes the level of Pterydophyta diversity in terms of morphological characteristics directly in the Tuban-Lamongan Pantura area, as shown in Figure 1.



Figure 1. Location of the research plots in the Tuban-Lamongan Pantura area

The research was conducted on March 8, 2024. The research method used was the transect method,

which divided the area into five plots. The transect method has the advantage of covering a large area quickly (Mwampeta et al., 2024). Instruments used include wooden stakes measuring 1 meter (4 pieces), raffia, stationery, a camera, a meter, a soil meter, a hygrometer, and a lux meter. The material used is the diversity of fern species found in the plot. Data collection techniques were carried out using exploratory techniques by directly observing the abiotic parameters of each plot, observing ferns in the plot based on their morphological characteristics, and counting the number of individuals of each species found in the plot. Data in this study are divided into two types: (1) quantitative in the form of the number of individuals of each species; (2) qualitative in the form of morphological characteristics of each species. The data analysis technique for the number of individuals of each species was carried out descriptively quantitatively using the Shannon & Wiener (1963) Index equation, which refers to Khalmuratova et al. (2021) in Equation 1, which is then interpreted into Table 1, which refers to Baderan et al. (2021). Meanwhile, data from morphological observations were analyzed descriptively and qualitatively to describe the morphological characteristics of each species.

$$H' = - \sum_{i=1}^s p_i \cdot \ln p_i \quad (1)$$

Description: H' = Shannon-Wiener diversity index

p_i = number of individuals found in i genus in a population

$p_i = \frac{n_i}{N}$ with; n_i = number of individuals in one species;

N = total number of individuals of the species found

\ln = Natural logarithm

Table 1. Diversity index value criteria

H' Value	Diversity Criteria
$H' < 1$	Low level
$1 \leq H' \leq 3$	Medium level
$H' > 3$	High level

Results and Discussion

Identifying Morphology of Pterydophyta

Abiotic factors strongly influence the growth of fern diversity. Therefore, abiotic parameters were observed in each research plot in the Tuban-Lamongan Pantura area, as shown in Table 2. Abiotic factors include soil pH, soil moisture, air temperature, light intensity, and air humidity. According to Umair et al. (2023), air temperature and rainfall are significant factors affecting fern diversity growth.

Table 2. Abiotic parameters for each plot

Plot Numbers	Abiotic Parameters				
	Soil pH	Soil Moisture	Air Temperature (°C)	Light Intensity (lux)	Air Humidity (%)
1	6,8	8	27	3072	96.1
2	6,2	4	26	2464	85.3
3	6,1	6	24	4966	81.6
4	6,2	2.5	30	1701	76.0
5	6,8	1	28	2163	74.1

The results of measuring abiotic parameters in various plots show that the highest soil pH value is in plots 1 and 5. which is 6.8. In contrast, The lowest pH value is in Plot 3. 6.1. However, the soil pH is considered ideal for fern growth. This is by Khairunisa & Wisanti (2023), who state that ferns thrive with soil pH between 5.5-6.5. Measurement of soil and air moisture parameters showed that plot 1 had the highest soil and air moisture compared to other plots, while plot 5 had the lowest soil and air moisture. Pramaisela et al. (2024) explained that ferns can grow and develop well with 60-90% humidity. Soil properties, including pH and soil moisture, play an essential role in the abundance and diversity of ferns (Cicuzza & Mammides, 2022). In addition, ferns tend to be in shady places, have shade, are relaxed, and have high levels of air humidity (Ibrahim et al., 2024; Singh & Chauhan, 2024). Air temperature measurements also show that Plot 4 has the highest air temperature, while Plot 3 has the lowest. Ferns thrive in tropical areas with a temperature range of 21-27°C due to these ideal temperature conditions (Rahmawati & Santhyami, 2023). Light intensity measurements showed that plot 1 had the lowest light intensity compared to the other plots, while plot 3 had the highest light intensity. Sianturi et al. (2021) explained that low light intensity is influenced by the absence of canopy and cloud cover; this condition is very suitable for fern growth habitat.

The results of research related to the diversity of ferns found in the Tuban-Lamongan Pantura area in the five plots showed that 19 species of ferns were classified into two classes, Polypodiopsida and Psilotopsida, with 210 individuals in total, as shown in [Table 3](#).

Table 3. Diversity of ferns on each plot

Plot Numbers	Class	Species Name	Number of Individuals (ni)		
1	Polypodiopsida	<i>Adiantum capillus-veneris</i>	26		
		<i>Pteris vittata</i>	3		
		<i>Pellaea viridis</i>	1		
	Psilotopsida	<i>Helminthostachys zeylanica</i>	1		
		<i>Pteris vittata</i>	12		
		<i>Pellaea atropurpurea</i>	2		
		<i>Cheilanthes micropetris</i>	5		
		<i>Adiantum latifolium</i>	1		
		<i>Asplenium pinnatifidum</i>	6		
2	Polypodiopsida	<i>Drynaria quercifolia</i>	4		
		<i>Botrychium jenmanii</i>	1		
		<i>Elaphoglossum stenoglossum</i>	1		
		<i>Christella dentata</i>	1		
		<i>Drynaria fortunei</i>	3		
		<i>Drynaria sparsisora</i>	4		
		<i>Platycterium bifurcatum</i>	9		
		3	Polypodiopsida	<i>Pteris vittata</i>	1
				<i>Asplenium pinnatifidum</i>	1
				<i>Drynaria quercifolia</i>	1
Psilotopsida	<i>Asplenium trichomanes</i>			3	
	<i>Botrychium jenmanii</i>			5	
4	Polypodiopsida	<i>Adiantum capillus-veneris</i>	18		
		<i>Pteris vittata</i>	4		
		<i>Helminthostachys zeylanica</i>	3		
		<i>Amblovenatum terminans</i>	6		
		<i>Pteris vittata</i>	58		
		<i>Elaphoglossum stenoglossum</i>	1		
		5	Polypodiopsida	<i>Drynaria sparsisora</i>	13
<i>Oleandra distenta</i>	4				
<i>Asplenium pinnatifidum</i>	9				
<i>Microsorium pustulatum</i>	3				

[Table 3](#) shows many ferns found in the plot 2 areas with ten species, with the most found being the *Pteris vittata* species. *Pteris vittata* species are found because of their ability to adapt to various types of soil and live in extreme environments. Besides that, *Pteris vittata* can also extract heavy metals in the soil, commonly known as phytoremediation ([Widyastuti & Surya, 2023](#)). Plot 1 is the tiniest plot where ferns are found, namely, only four types of species. However, in plot one, many species of *Adiantum capillus-veneris* were found, with a total of 26 individuals. *Adiantum capillus-veneris* species is a cosmopolitan fern species that lives in various niches ([Rachmalia et al., 2023](#)). In plots 3 and 4, fern species were found with the same number of species, namely five species with different types. In plot 3, the most species found were *Platycterium bifurcatum*, while in plot 4, the most species found were *Adiantum capillus-veneris*. *Platycterium bifurcatum* species are found attached to trees as their host. This is by [Wu & Brock \(2023\)](#), who state that *Platycterium bifurcatum* is an epiphytic plant.

In plot 5, 7 types of fern species were found. The most common fern species found was *Pteris vittata* in the plot 5 area with 58 individuals. In addition, *Pteris vittata* species were also found in all plots. This is per the opinion of [Sukmawati et al. \(2024\)](#) that *Pteris vittata* is known for its ability to live in various types of substrates and ecosystems so that it is widely distributed. *Pteris vittata* is also known for its ability to hyperaccumulate arsenic ([Kohda et al., 2024](#)). In addition, many fern species are only found in one individual in several plots, such as *Pellaea viridis*, *Helminthostachys zeylanica*, *Adiantum latifolium*, *Botrychium jenmanii*, *Elaphoglossum stenoglossum*, *Christella dentata*, *Asplenium pinnatifidum*, *Drynaria quercifolia* and *Elaphoglossum stenoglossum*. The ferns found were the Polypodiopsida class found in all plots with a small number of species, while the Psilotopsida class was only found in plot 1 and plot 4 with a small number of species. This is under [Azzahro \(2022\)](#), who states that Polypodiopsida is a class of ferns with the most members. In addition, almost all species of the Polypodiopsida class have been identified. They are widely distributed in the tropics and subtropics, with the majority of species living epiphytically and in modern fern groups, so this type is included in a group that has high adaptation, which causes the distribution of this group to expand ([Inelia et al., 2023](#); [Isa et al., 2023](#)). [Hidayah, \(2019\)](#)

explains that the Psilotopsida class is an ancient fern; many of these ferns have become fossils, while Polypodiopsida ferns are accurate or actual ferns.

Figure 2, Figure 3, Figure 4, Figure 5, and Figure 6 present images of the morphological characteristics of ferns found in the Tuban-Lamongan Pantura area according to each plot. The results of observations of the morphological characteristics of the ferns found are presented in Table 4.



Figure 2. Fern species in plot 1: a) *Adiantum capillus-veneris*; b) *Pteris vittata*; c) *Pellaea viridis*; d) *Helminthostachys zeylanica*



Figure 3. Fern species in plot 2: a) *Pteris vittata*; b) *Pellaea atropurpurea*; c) *Cheilanthes micropetris*; d) *Adiantum latifolium*; e) *Asplenium pinnatifidum*; f) *Drynaria quercifolia*; g) *Botrychium jenmanii*; h) *Elaphoglossum stenoglossum*; i) *Christella dentata*; j) *Drynaria fortune*




Figure 4. Fern species in plot 3: a) *Drynaria sparsisora*; b) *Platycerium bifurcatum*; c) *Pteris vittata*; d) *Asplenium pinnatifidum*; e) *Drynaria quercifolia*




Figure 5. Fern species in plot 4: a) *Asplenium trichomanes*; b) *Botrychium jenmanii*; c) *Adiantum capillus-veneris*; d) *Pteris vittata*; e) *Helminthostachys zeylanica*




Figure 6. Fern species in plot 5: a) *Amblovenatum terminans*; b) *Pteris vittata*; c) *Elaphoglossum stenoglossum*; d) *Drynaria sparsisora*; e) *Oleandra distenta*; f) *Asplenium pinnatifidum*; g) *Microsorium pustulatum*

Table 4. Observation of morphological characteristics of each species

Species Name	Morphological Characteristics				
	Root	Stem	Leaf	Sorus	Rhizoma
<i>Adiantum capillus-veneris</i>	Rhizoid	Black, Spreading, smooth, shiny	Sterile leaves, macrophyllous, shiny light green, curved round, serrated edge	On the edge of the fertile leaf, brownish yellow, small round, brown	Branched, small short, scaly
<i>Pteris vittata</i>	Fibrous, calyptra	Brown, spreading, hard	Sterile leaves, macrophyllous, long-pinnate compound, dark green, sharp tip and longer	Elongated at the edge of fertile leaves, small round, brown	Short, stiff, spreading, green when young and light brown when old
<i>Pellaea viridis</i>	Rhizoid	Light green, spreading	Sterile leaves, macrophyllous, round, serrated edges, dark green above and light green below	Below along the edge of fertile leaves, small round, brown	Small, short, spreading, scaly
<i>Elaphoglossum stenoglossum</i>	Rhizoid	Black, smooth	Sterile leaves, macrophyllous, dark green, shiny, wavy, elongated	Below along the edge of fertile leaves, small round, brown	Small, short, scaly
<i>Helminthostachys zeylanica</i>	Rhizoid	Brown, black, hard	Sterile leaves, macrophyll, pinnate, pointed tip, dark green	On stem at base of midrib sterile, small round, brown	Short, stiff, spreading
<i>Drynaria sparsisora</i>	Fibers (sporophytes), calyptra	Hard, brown	Sterile leaves, macrophyll, dark green, fingered, wavy, smooth	Located on the underside of fertile leaves, small round, brown	Long, tight, attached to the host, branched, scaly
<i>Platynerium bifurcatum</i>	Fibers (sporophytes), calyptra	Light brown, round	Fertile leaves, macrophyll, green, branched like deer antlers, curled tips	Located on the underside of fertile leaves, small round, brown	Spreading, flat, attached to the host
<i>Pellaea atropurpurea</i>	Rhizoid	Hard, light green	Sterile leaves, macrophyll, light green, oval, serrated edges	Located on the underside of fertile leaves, small round, brown	Short, spreading, scaly, curly, reddish brown
<i>Cheilanthes micropetris</i>	Rhizoid	Hard, spreading, smooth-haired	Sterile leaves, macrophyllous, pinnately oval, wavy, light green, smooth-haired	Located on the underside of fertile leaves, round and oval	Spreading, hard, upright, scaly, short, tight
<i>Adiantum latifolium</i>	Rhizoid	Light green, soft, long, slender, smooth-haired	Sterile leaves, microphyllous, round, curled, hairy, curved, thin	Located on the underside of fertile leaves, brownish black	Short, spreading, small-branched, brown
<i>Amblovenatum terminans</i>	Fibers (sporophytes)	Dark green, hard	Sterile leaves, macrophyll, parallel, long, dark green, smooth	Located on the lower edge of fertile leaves, small round, brown	Long, tight
<i>Asplenium pinnatifidum</i>	Rhizoid	Green, soft, shiny	Sterile leaves, macrophyll, round, serrated edge, pointed tip, oval	Located on the underside of the fertile leaf edge, small round, brown	Short, spreading, tight
<i>Drynaria quercifolia</i>	Fibrous (sporophyte), calyptra	Brown, hard, long	Sterile leaves, macrophyll, light green, fingered, pinnate, blunt tip	Located on the lower edge of the fertile leaves, arranged in rows, small round, brown	Tight, long, creeping, scaly
<i>Christella dentata</i>	Rhizoid	Light green, long, slender, smooth-haired	Sterile leaves, macrophyllous, compound, fingered, sharp tip, rounded edge	Below along the edge of the fertile leaf, small round, brown	Short, small, scaly
<i>Drynaria fortunei</i>	Rhizoid	Light green, long, round	Sterile leaves, microphyllous, parallel, compound, wavy, tip rounded	Below along edge of fertile leaf, small round, brown	Round, hairy, scaly, brown
<i>Botrychium jenmanii</i>	Rhizoid	Soft, light green	Sterile leaves, macrophyllous, compound, rounded, wide rounded tips	Below surface fertile leaves, small round, brown	Short, small

Species Name	Morphological Characteristics				
	Root	Stem	Leaf	Sorus	Rhizoma
<i>Asplenium trichomanes</i>	Fibrous (sporophyte), smooth	Keras, coklat	ard, brown Leaves sterile, macrophyllous, compound, small round, serrated edge, rounded leaf tip	Below surface edge of fertile leaf, small round, brown	Long, creeping, hairy
<i>Microsorium pustulatum</i>	Fibers (sporophytes), smooth	Hard, light green	Sterile leaves, macrophyll, light green, fingered, wide	Below surface edge of fertile leaves, small round, brown	Short, round hairy
<i>Oleandra distenta</i>	Fibers (sporophytes), smooth	Hard, dark green, smooth	Leaves sterile, macrophyllous, dark green, wide smooth, parallel	Crosswise below surface of fertile leaves, small round, brown	Creeping, round, hairy

Almost all of the 19 fern species found have root morphology in the form of saprophytes and rhizoids. Ferns have rhizoid roots in the gametophyte generation, while ferns will show the characteristics of fibrous roots in the sporotiv generation (Kurnia, 2023). Observations also show that the stems of ferns have a rigid structure that is green when the plant is young and turns brown when the plant is old, such as in the species *Pteris vittata*. *Pteris vittata* has a dark brown rhizome base that spreads (Ramndana et al., 2023). Observations of fern stem morphology also show the presence of fine hairs covering part of the stem surface, such as in the species *Cheilanthes micropetris*, *Adiantum latifolium*, and *Christella dentata*. This is per Ulfa et al. (2023a), which states that the stems of ferns are covered mainly by several light brown fine hairs that are sparsely distributed along the stem, but more and more fine hairs are found near the roots. In addition, the species *Christella dentata* has brown scales on its rhizome (Majid et al., 2022). The species *Adiantum capillus-veneris* has a shiny black stem and looks slippery; this is what distinguishes it from the other fern species found, in accordance with the opinion of Nurcahyani (2021), which states that *Adiantum capillus-veneris*, commonly known as supliir, has leaf stalks and curved stems that are shiny, black and smooth. In Unayah & Widodo (2023), it is explained that the morphological characteristics of ferns are influenced by the genetic characteristics of these plants and environmental factors, such as climate change, temperature, soil type, soil conditions, altitude, and humidity.

The leaf morphology of ferns found has a variety of diversity; there are leaves that have curved, pinnate, fingered, and parallel leaf bones. Most fern leaves grow in compounds and include macrophilous leaves. The findings Sofiyanti et al. (2020) show that most ferns have linear and alternate compound leaves. In addition, compound leaves are a common form of fern leaf morphology (Ulfa et al., 2023b). The leaves of ferns are also classified as simple and still need to be in the form of strands (Anggraini et al., 2023). Ulfa, et al. (2023a) also explained that, in general, the leaves of ferns can be classified into two categories based on their size, namely large macrophil leaves and small microfil leaves, usually in the form of scales. The results of identifying the leaf morphology of ferns also show that most fern leaf morphology is in the form of sterile leaves, evidenced by the absence of sorus on the edge or bottom of the leaf. In addition, the tip of the leaf also rolls at a young age. Young leaves of fern plants have characteristics where the leaves form rolls and do not produce spores (Lestari & Indriyani, 2023; Ridhwan et al., 2023). Sterile leaves indicate that the age of the fern is still young. However, there are identification results where fertile leaves are found in *Platyserium bifurcatum* species. *Platyserium bifurcatum* species grow by attaching to their host tree with brown rhizomes covered by many supporting leaves (Nafisah et al., 2023).

Most sori on ferns are located at the bottom of fertile leaves with a transverse position or on the edge of the leaf. This is by Pradipta et al. (2023), which states that most Polyodiopsida class ferns have the sorus's location and position on the leaf's lower surface. The location of the sorus against the leaf bone is a crucial trait in the classification of ferns (Naiym & Munir, 2024). Sorus is a characteristic of ferns because of its location in a grouped spore (Aulia et al., 2024). The results of morphological identification of rhizomes of ferns showed that overall rhizomes of ferns with hair of various sizes, ranging from long and short depending on the size of the ferns; besides the rhizomes of ferns, are also many creeping and creeping to attach to the host, such as *Drynaria sparsisora*, *Platyserium bifurcatum* and *Drynaria quercifolia*. This suggests that some ferns are epiphytes. *Drynaria quercifolia* species has scaly rhizomes resembling brown hair with large rhizome leaves and creeping (Afriana et al., 2021). *Drynaria quercifolia* species are also found attached to the trunk of oil palm plants with creeping stems and have ovoid supporting leaves that cover the roots and rhizomes (Jusri et al., 2022; Listiyanti et al., 2022). In addition, *Drynaria quercifolia* species have sorus in the form of brown round dots scattered on the lower surface of the leaves (Tnunay & Hanas, 2020).

Identification of Pterydophyta Diversity Index

Table 5 presents the results of the analysis related to the diversity index in each fern species found in the Tuban-Lamongan Pantura area.

Table 5. Diversity index of each species

Species Name	ni	$pi \left(\frac{ni}{N} \right)$	$\ln pi$	$pi \ln pi$
<i>Adiantum capillus-veneris</i>	44	0.21	-1.57	-0.33
<i>Pteris vittata</i>	78	0.37	-1.00	-0.37
<i>Pellaea viridis</i>	1	0.00	-5.35	-0.03
<i>Elaphoglossum stenoglossum</i>	2	0.01	-4.66	-0.04
<i>Helminthostachys zeylanica</i>	4	0.02	-3.97	-0.08
<i>Drynaria sparsisora</i>	17	0.08	-2.52	-0.20
<i>Platyserium bifurcatum</i>	9	0.04	-3.15	-0.13
<i>Pellaea atropurpurea</i>	2	0.01	-4.66	-0.04
<i>Cheilanthes micropetris</i>	5	0.02	-3.74	-0.09
<i>Adiantum latifolium</i>	1	0.00	-5.35	-0.03
<i>Amblovenatum terminans</i>	6	0.03	-3.56	-0.10
<i>Asplenium pinnatifidum</i>	16	0.08	-2.58	-0.20
<i>Drynaria quercifolia</i>	5	0.02	-3.74	-0.09
<i>Christella dentata</i>	1	0.00	-5.35	-0.03
<i>Drynaria fortune</i>	3	0.01	-4.25	-0.06
<i>Botrychium jenmanii</i>	6	0.03	-3.56	-0.10
<i>Asplenium trichomanes</i>	3	0.01	-4.25	-0.06
<i>Microsorium pustulatum</i>	3	0.01	-4.25	-0.06
<i>Oleandra distenta</i>	4	0.02	-3.97	-0.08
Total	210	1.00	-71.48	2.10

Based on the data analysis in Table 5, it is known that the fern diversity index in the Pantura Tuban-Lamongan area obtained a result of $H'=2.10$, which is classified as in the medium category with the criteria $1 \leq H' \leq 3$ with a total of 19 types of species and a total of 210 individuals. In [Laeto & Taharu \(2021\)](#) it is explained that the moderate diversity index shows that the number of species among the total number of individuals of all existing species is moderate, meaning that the ratio of the number of individuals of a species to the total number of individuals of all species is moderate. The diversity index value is relatively high, considering the Tuban-Lamongan coastal area has high air temperature and light intensity. In addition, the number of individuals of each species also varies and has an uneven distribution pattern. The acquisition of the number of individuals of each different species is due to several abiotic factors that influence it, such as soil pH and air temperature ([Zhang et al., 2021](#)). This is in accordance with the results of research by [Liu et al. \(2021\)](#), which states that functional richness, maximum air temperature, soil pH, and their interactions show a strong influence on species diversity. Based on the diversity index value obtained also shows that the existing fern species community is relatively stable, as evidenced by the diversity index value obtained, which is classified as medium. This is considering that the higher the diversity index value, the more stable the existing species community ([Inama et al., 2023](#); [Leki et al., 2022](#)). In addition, the high and low diversity index of a plant community depends on the number of species, individuals of each type, and existing environmental factors ([Ballo et al., 2024](#); [Windari et al., 2021](#)). If the number of identified species increases, the diversity index will increase. However, conversely, if the number of identified species is only small, then the diversity index will be low ([Yolla et al., 2022](#)). The fern species with the highest diversity value is *Pteris vittata*, 0.37. Meanwhile, the fern species with the lowest diversity values are *Elaphoglossum stenoglossum*, *Adiantum latifolium*, and *Christella dentata*. The *Christella dentata* species is usually found in soil near water sources ([Puspa et al., 2023](#)). This is because there is only one individual in each of the three species of ferns. Ferns can spread quickly and form a diversity that can be identified based on morphological characteristics. The diversity in question is the richness of fern species found in an area ([Adlini et al., 2021](#)).

Analysis of Potential Research Results in Support of FBP

The results of research that has been carried out regarding morphology-based Pterydophyta diversity in the Pantura Tuban-Lamongan area can potentially be used in the world of education to support the implementation of FBP in Low plant botany courses for biology education students with the contribution of providing knowledge and understanding of morphology-based Pterydophyta diversity learning materials comprehensively, directly through experience and application of theories and research methods used in everyday life.

FBP is one of the activities in Project Learning (PjBL) courses that aim to make students independent through direct learning in nature, the field, or the community ([Nurdiansyah, 2023](#)). FBP plays a role in students integrating and applying knowledge values into practice in a complex manner involving several skills ([Arundel et al., 2022](#)). [Matsna et al. \(2023\)](#) explained that the application of FBP can improve students' Science Process Skills (KPS) by allowing students to experience or do it themselves and be able to determine problems, observe, analyze, hypothesize, conclude, and apply the theories they have

according to their needs, apart from that FBP It is also able to arouse learning motivation and encourage students' curiosity so that these principles will support students to discover their knowledge through field exploration. Therefore, FBP is widely applied to support students in determining the effectiveness of ongoing learning (Petra et al., 2020; Tedam et al., 2021).

PjBL-based FBP has been applied in learning activities for students, and the results of the learning process show positive changes in students because students experience themselves actively in their activities. Student activities through FBP on morphology-based Pteridophyta diversity can increase student knowledge, understanding, learning outcomes, motivation, and critical thinking skills. This is in accordance with the results of research by Aleman et al. (2021) which states that students who use field-based practicum show a positive increase in the benefits, value and student engagement with learning. Pratomo & Nur (2024) added that field practicum activities are an effort to broaden cognitive insights while honing students' psychomotor abilities. In addition, Alamri (2018) stated that practicum problems are a big critical concern and essential to learning. FBP is an innovation in learning considering that practicum in the laboratory has several serious concerns, including classroom management, non-existent laboratory space, lack of tools and materials, teacher training on laboratory management is very lacking, practicum duration, lack of teacher regulations and experience, and inefficient feedback (Bahtiar et al., 2022).

The application of FBP in Low plant botany courses for biology education students regarding morphology-based Pteridophyta diversity in the Pantura Tuban-Lamongan area requires further research with developing field-based practical instructions.

Conclusion

Based on the research results, we conclude that the level of fern diversity in the Pantura Tuban-Lamongan area is classified as medium with diverse morphological characteristics. The results of the research that has been carried out have the potential to be used in the world of education to support field-based practicums in Low plant botany courses with the contribution of providing direct knowledge and experience regarding the diversity of ferns in the surrounding environment.

Acknowledgment

The author would like to thank the Biology Education study program at Universitas PGRI Ronggolawe for providing the guidance and direction that made this article successful. The author is also very grateful to all parties who have assisted technically in taking research samples.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

Author Contributions

Y. Yudhastian: data analysis, methodology, writing original draft preparation, observation and sampling; and **T. S. H. Wulandari:** review and editing.

References

- Adlini, M. N., Hartono, A., Khairani, M., Tanjung, I. F., & Khairuna. (2021). Identification of ferns (Pteridophyta) in Universitas Islam Negeri (UIN) Sumatera Utara. *Biota: Jurnal Ilmiah Ilmu-Ilmu Hayati*, 6(2), 1–7. <https://doi.org/10.24002/biota.v6i1.3023>
- Afifah, F. N., Putri, N. R. A., Hartanti, A. N., Pramudita, D. A., Armando, M. F., Rahmayani, D., Indrawan, M., Safira, R. N., Jr, I. E. B., & Setyawan, A. D. (2023). Fern diversity and conservation status in the South Gombong karst forest, Kebumen District, Indonesia. *Pros Sem Nas Masy Biodiv Indon*, 9(1), 186–195. <https://doi.org/10.13057/psnmbi/m090128>
- Afriana, M., Darwin, C., Lubis, R., & Saroni, S. (2021). Keanekaragaman jenis tumbuhan paku (Pteridophyta) di Kecamatan Ketahun Kabupaten Bengkulu Utara. *Jurnal Riset Dan Inovasi Pendidikan Sains (JRIPS)*, 1(1), 1–18. <https://doi.org/10.36085/jrips.v1i1.2785>
- Alamri, H. R. H. (2018). Challenges in practicum: Views and perceptions of EFL pre-service teachers towards field experience skills in real classrooms. *Arab World English Journal (AWEJ)*, 9(1), 146–162. <https://doi.org/10.24093/awej/vol9no1.11>
- Aleman, R., Duball, C., Schwyter, A., & Vaughan, K. (2021). Remote delivery of field experiences in

- soil sciences. *Natural Sciences Education*, 50(1), e20049. <https://doi.org/10.1002/nse2.20049>
- Anggraini, N., Nugroho, A. S., & Dzakiy, M. A. (2023). Identifikasi keanekaragaman Pteridophyta di kawasan wisata Curug Lawe Secepit Kendal. *Prosiding Webinar Biofair 2023*. 478–487. <https://conference.upgris.ac.id/index.php/biofair/article/view/4219>
- Arundel, M. K., Morrison, S., Mantulak, A., & Csiernik, R. (2022). Social work field practicum instruction during COVID-19: Facilitation of the remote learning plan. *Fieldeducator Simmons Edu*, 12(April), 1–19. <https://fieldeducator.simmons.edu/wp-content/uploads/2022/02/21-285.pdf>
- Aulia, A. A., Fatma, E. T. R., Nurcholis, N. A., Fadilah, N., Dini, T. A., & Fardhani, I. (2024). Keanekaragaman jenis paku-pakuan (Pteridophyta) di Coban Putri Kota Batu beserta potensi kebermanfaatannya. *Jurnal Pendidikan MIPA*, 14(1), 187–193. <https://doi.org/10.37630/jpm.v14i1.1439>
- Azrai, E. P., Miarsyah, M., Widarsih, K., Gusverizon, A. N. R., Muslimin, S. I., Syahrani, Y., & Rojak. (2024). Association study of various species of epiphyte fern with host tree in Srengseng city forest. *AIP Conference Proceedings*, 2982(1), 50031. <https://doi.org/10.1063/5.0184896>
- Azzahro, F. (2022). Inventarisasi tumbuhan paku di Desa Cengkehan, Kabupaten Batul, Yogyakarta. *Jurnal Tropika Mozaika*, 1(1), 1–6. <https://ejournal.uin-suka.net/index.php/mozaika/article/view/2>
- Baderan, D. W. K., Rahim, S., Angio, M., & Salim, A. I. Bin. (2021). Keanekaragaman, pemerataan, dan kekayaan spesies tumbuhan dari Geosite Potensial Benteng Otanaha sebagai rintisan pengembangan Geopark Provinsi Gorontalo. *Al-Kauniah: Jurnal Biologi*, 14(2), 264–274. <https://doi.org/10.15408/kauniah.v14i2.16746>
- Bahtiar, B., Maimun, M., & Baiq Lily Anggriani W. (2022). Pengaruh model discovery learning melalui kegiatan praktikum IPA terpadu terhadap kemampuan berpikir kritis siswa. *Jurnal Pendidikan Mipa*, 12(2), 134–142. <https://doi.org/10.37630/jpm.v12i2.564>
- Ballo, A., Zacharia, J., & Nge, S. T. M. (2024). Keanekaragaman tumbuhan paku (Pteridophyta) di kawasan air terjun oesusu Kabupaten Kupang. *Journal of Biology Education and Biotechnology*, 1(1), 18–26. <https://www.nafatimahpustaka.org/jbeb/article/view/343>
- Boileau, E., & O'Donoghue, L. (2024). Learning to embrace outdoor pedagogy: early childhood education student experiences of a nature-focused practicum. *Journal of Outdoor and Environmental Education*. <https://doi.org/10.1007/s42322-023-00153-1>
- Cicuzza, D., & Mammides, C. (2022). Soil, topography and forest structure shape the abundance, richness and composition of fern species in the fragmented tropical landscape of Xishuangbanna, Yunnan, China. *Forests*, 13(9), 1–15. <https://doi.org/10.3390/f13091453>
- Das, T. K., Paul, K., & Bhattacharjee, P. (2024). What happened to field practicum in social work education during COVID-19? evidence from Bangladesh. *Asian Social Work and Policy Review*, 18(1), e12294. <https://doi.org/10.1111/aswp.12294>
- De-Villa, K. R., & Lagat, R. D. (2023). Species diversity and habitat association of ferns and Lycophytes in Mts. Palay-Palay Mataas na Gulod protected landscape BT - plant diversity in Biocultural landscapes. *Springer Singapore*, 135–161. https://doi.org/10.1007/978-981-19-8649-9_7
- Deng, W., Lu, Y., Lyu, M., Deng, C., Li, X., Jiang, Y., Zhu, H., Yang, Y., & Xie, J. (2024). Chemical composition of soil carbon is governed by microbial diversity during understory fern removal in subtropical pine forests. *Science of The Total Environment*, 914, 169904. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2024.169904>
- Di-Michele, W. A., Lucas, S. G., Eble, C. F., Kerp, H., Reynolds, S. J., May, P., & Pigg, K. B. (2024). A detailed stratigraphic and taphonomic reassessment of the late Paleozoic fossil flora from Promontory Butte, Arizona. *Review of Palaeobotany and Palynology*, 320, 105004. <https://doi.org/10.1016/j.revpalbo.2023.105004>
- Dubuisson, J.-Y., Nivart, A., Bidault, E., Deblauwe, V., Droissart, V., Kamdem, N. G., Rouhan, G., Ebihara, A., & le Péchon, T. (2024). Diversity, taxonomy, and history of the tropical fern genus *Didymoglossum* Desv. (Hymenophyllaceae, Polypodiidae) in Africa. *Journal of Systematics and Evolution*, 62(1), 84–101. <https://doi.org/10.1111/jse.12951>
- Fischer, E., & Lobin, W. (2024). Checklist of Lycopodiopsida (clubmosses and quillworts) and Polypodiopsida (ferns) of Rwanda. *Willdenowia*, 53(3), 149–172. <https://doi.org/10.3372/wi.53.53302>
- Fujiwara, T., Liu, H., Meza-Torres, E. I., Morero, R. E., Vega, A. J., Liang, Z., Ebihara, A., Leitch, I. J., & Schneider, H. (2023). Evolution of genome space occupation in ferns: Linking genome diversity and species richness. *Annals of Botany*, 131(1), 59–70. <https://doi.org/10.1093/aob/mcab094>
- Guillen-Otero, T., Lee, S. J., Hertel, D., & Kessler, M. (2024). Facultative mycorrhization in a fern (*Struthiopteris spicant* L. Weiss) is bound to light intensity. *BMC Plant Biology*, 24(1), 103. <https://doi.org/10.1186/s12870-024-04782-6>
- Hande, S. S., & Sharma, L. K. (2021). Impact of supervision styles on field practicum experiences. *International Journal for Advanced Research in Science & Technology*, 11(12), 1323–1328. <https://www.ijarst.in/public/uploads/paper/244521706193030.pdf>

- Hidayah, F. (2019). Kekayaan jenis tumbuhan paku (Pteridophyta) di kawasan Cagar Alam Watangan Puger Jember serta pemanfaatannya sebagai Booklet [Universitas Jember]. In *Universitas Jember*. <https://doi.org/10.1515/9783110949995.29>
- Ibiye, A., Green, B. O., & Ajuru, M. G. (2023). Short communication: Comparative foliar epidermal study of some species of pteridophytes in Rivers State University, Nigeria. *Cell Biology & Development*, 7(2), 51–55. <https://doi.org/10.13057/cellbioldev/v070201>
- Ibrahim, Y., Gurnita, & Karima, E. M. (2024). Identification of the diversity of fern plant types in Mount Masigit Kareumbi Buru Park, Bandung, Indonesia. *S&S: Journal of Law and Sustainable Development*, 12(2), 1–17. <https://doi.org/10.55908/sdgs.v12i2.3174>
- Inama, Santoso, N. P., Raihandhany, R., & Kurniawan, F. H. (2023). Inventarisasi jenis-jenis tumbuhan yang tumbuh meliar di Eks pabrik gula Kadipaten, Jatiwangi, dan Parungjaya, Kabupaten Majalengka. *Jurnal Biologica Samudra*, 5(2), 174–189. <https://doi.org/10.33059/jbs.v2i1.6897>
- Inelia, S., Nurhadi, & Safitri, E. (2023). Pteridophyta yang ditemukan di air terjun Batu Basurek Kecamatan Sungai Geringging Kabupaten Padang Pariaman. *Jurnal Pendidikan Tambusai*, 7(3), 20317–20322. <https://doi.org/10.31004/jptam.v7i3.9477>
- Isa, D., Turnip, M., & Linda, R. (2023). Jenis tumbuhan paku terestrial (Pteridophyta) di kawasan Universitas Tanjungpura Pontianak Kalimantan Barat. *Biologica Samudra*, 5(1). <https://doi.org/10.33059/jbs.v5i1.6200>
- Joseph, M. K., & Jamal, A. (2023). Social work field practicum in government-initiated projects. *Journal of Social Work Education and Practice*, 8(2), 1–7. <https://jswep.in/index.php/jswep/article/view/153>
- Juliasih, N. K. A., & Adnyana, I. M. D. M. (2023a). Ethnopharmacology and species diversity of Pteridophyta in Cyathea Park, Bali: A Field study and Literature Review. *Research Square*, 1–21. <https://doi.org/10.21203/rs.3.rs-2497775/v2>
- Juliasih, N. K. A., & Adnyana, I. M. D. M. (2023b). Utilization of Pteridophytaspecies in Cyathea Park, Bali, as traditional medicine agents: A field study and as traditional medicine agents: A field study and field study and meta-synthesis review. *Notulae Scientia Biologicae (NSB)*, 13(1), 1–24. <https://doi.org/10.15835/nsb15211522>
- Juniar, A., Monalisa, C., & Tarigan, B. (2024). The effect of implementing practicum-based contextual teaching and learning models to improve learning outcomes on the subject of Colloids. *PENDIPA Journal of Science Education*, 8(1), 57–64. <https://doi.org/10.33369/pendipa.8.1.57-64>
- Jusri, Nildayanti, Poerwanti, H., & Sofyan. (2022). Identification of epifite potential spikes on stems palm oil plant (*Elaeis Guineensis* Jacq.). *Jurnal Agroplantae: Jurnal Ilmiah Terapan Budidaya Dan Pengelolaan Tanaman Pertanian Dan Perkebunan*, 11(2), 59–71. <https://doi.org/10.51978/agro.v11i2.453>
- Kandriasari, A., Yulianti, Y., Febriana, R., & Mohd Nordin, R. (2023). Development of practicum learning models assisted with practice guide videos. *JTP - Jurnal Teknologi Pendidikan*, 25(2), 246–256. <https://doi.org/10.21009/jtp.v25i2.34678>
- Katayama, N., Yamamoto, T., Aiuchi, S., Watano, Y., & Fujiwara, T. (2024). Subgenome evolutionary dynamics in allotetraploid ferns: insights from the gene expression patterns in the allotetraploid species *Phegopteris decursivopinnata* (Thelypteridaceae, Polypodiales). *Frontiers in Plant Science*, 14(January), 1–15. <https://doi.org/10.3389/fpls.2023.1286320>
- Khairunisa, & Wisanti. (2023). Species diversity Pteridophyta in the tourism area of Supit Urang waterfall Pujon Malang Regency. *LenteraBio*, 13(3), 343–353. <https://doi.org/10.26740/lenterabio.v12n3.p343-353>
- Khalmuratova, I., Choi, D. H., Yoon, H. J., Yoon, T. M., & Kim, J. G. (2021). Diversity and plant growth promotion of fungal endophytes in five halophytes from the buan salt marsh. *Journal of Microbiology and Biotechnology*, 31(3), 408–418. <https://doi.org/10.4014/JMB.2012.12041>
- Khan, N., Ullah, R., Okla, M. K., Abdel-maksoud, M. A., Saleh, I. A., Abu-harirah, H. A., Alramadneh, T. N., & Abdelgawad, H. (2024). Ecological factors affecting minerals and nutritional quality of “*Dryopteris filix-mas* (L.) Schott”: an underutilized wild leafy vegetable in rural communities. *Frontiers in Nutrition*, 11. 1276307. <https://doi.org/10.3389/fnut.2024.1276307>
- Khotimah, M. H., Sutarno, M., Parlidungan, D., Uliyandari, M., & Defianti, A. (2023). Terrestrial fern diversity in Palak Siring waterfall and Its potential to support Junior High School science learning. *Jurnal Penelitian Pendidikan IPA*, 9(12), 10832–10844. <https://doi.org/10.29303/jppipa.v9i12.4844>
- Kohda, Y. H.-T., Hamai, T., Horiuchi, K., Semoto, Y., & Inoue, C. (2024). New evidence of the arsenic uptake and translocation in As-hyperaccumulator fern *Pteris cretica* using a modified hydroponic system. *Journal of Hazardous Materials*, 463. 132855. <https://doi.org/10.1016/j.jhazmat.2023.132855>
- Kumar, A., Giri, P., & Uniyal, P. L. (2022). *Genome Evolution in Ferns: Molecular Phylogenomics – A Review BT- Ferns: Biotechnology, Propagation, Medicinal Uses and Environmental Regulation* (J. Marimuthu, H. Fernández, A. Kumar, & S. Thangaiyah (eds.); pp. 13–37). Springer Nature Singapore. https://doi.org/10.1007/978-981-16-6170-9_2

- Kurnia, N. (2023). Studi morfologi dan hubungan kekerabatan familia polypodiaceae di kawasan gunung galunggung tasikmalaya sebagai sumber belajar biologi. Universitas Siliwangi. <http://repositori.unsil.ac.id/id/eprint/8583>
- Laeto, A. Bin, & Taharu, F. I. (2021). Keanekaragaman jenis tumbuhan paku (Pteridophyta) berdasarkan topografi di kawasan hutan suaka margasatwa Lambusango Kabupaten Buton. *Sang Pencerah: Jurnal Ilmiah Universitas Muhammadiyah Buton*, 7(1), 71–86. <https://doi.org/10.35326/pencerah.v7i1.961>
- Lee, Y.-J., & Davis, R. (2023). Preservice teachers' self-efficacy through hybrid field practicum in a Korean teacher education program. *International Journal of Instruction*, 16(4), 349–366. <https://doi.org/10.29333/iji.2023.16421a>
- Leki, P. T., Makaborang, Y., & Ndjeroemana, Y. (2022). Keanekaragaman tumbuhan paku (Pteridophyta) di daerah aliran sungai Pepuwatu Desa Prai Paha Kabupaten Sumba Timur sebagai sumber belajar biologi. *BIOEDUKASI (Jurnal Pendidikan Biologi)*, 13(1), 42. <https://doi.org/10.24127/bioedukasi.v13i1.5304>
- Lestari, I., & Indriyani, S. (2023). Jenis paku-pakuan di desa Jelapat II berdasarkan karakter morfologi. *Al Kawnu: Science and Local Wisdom Journal*, 3(1), 39–48. <https://doi.org/10.18592/alkawnu.v3i1.8792>
- Li, M., Kuang, F., & Dan, W. (2023). Exploring the characteristics of pre-service EFL teachers' practicum experiences: a complexity theory-based case study in China. *Asian-Pacific Journal of Second and Foreign Language Education*, 8(13). <https://doi.org/10.1186/s40862-023-00187-4>
- Listiyanti, R., Indriyani, S., & Ilmiah, N. (2022). Karakteristik morfologi jenis-jenis paku epifit pada tanaman kelapa sawit di Desa Tegalrejo. *Al Kawnu: Science and Local Wisdom Journal*, 2(1), 99–106. <https://doi.org/10.18592/ak.v1i3.7281>
- Liu, J. J., Xu, Y., Shan, Y. X., Burgess, K. S., & Ge, X. J. (2021). Biotic and abiotic factors determine species diversity-productivity relationships in mountain meadows. *Journal of Plant Ecology*, 14(6), 1175–1188. <https://doi.org/10.1093/jpe/rtab064>
- Lolowang, Y., Arrijani, & Ogi, N. L. I. M. (2023). Effectiveness of practicum activities as an effort to improve student learning outcomes at SMA Negeri 2 Tondano. *Jurnal Arjuna: Publikasi Ilmu Pendidikan, Bahasa Dan Matematika*, 1(5), 160–170. <https://doi.org/10.61132/arjuna.v1i4.219>
- Lu, Y., Lyu, M., Xiong, X., Deng, C., Jiang, Y., Zeng, M., & Xie, J. (2023). Understory ferns promote the restoration of soil microbial diversity and function in previously degraded lands. *Science of The Total Environment*, 870. 161934. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2023.161934>
- Majid, A., Ajizah, A., & Amintarti, S. (2022). Keragaman tumbuhan paku (Pteridophyta) di taman biodiversitas hutan hujan tropis Mandiangin. *Jurnal Al-Azhar Indonesia Seri Sains Dan Teknologi*, 7(2), 102. <https://doi.org/10.36722/sst.v7i2.1117>
- Marasinghe, L. D. K., & Daulagala, P. W. H. K. P. (2024). Assessment of diversity and distribution pattern of Pteridophytic flora in Dunumadalawa Forest Reserve in Kandy District, Sri Lanka. *Asian Journal of Research in Botany*, 7(1), 36–46. <http://archive.sdpublishers.com/id/eprint/2456>
- Matsna, F. U., Rokhimawan, M. A., & Rahmawan, S. (2023). Analisis keterampilan proses sains siswa melalui pembelajaran berbasis praktikum pada materi Titrasi Asam-Basa kelas XI SMA/MA. *Dalton: Jurnal Pendidikan Kimia Dan Ilmu Kimia*, 6(1), 21. <https://doi.org/10.31602/dl.v6i1.9187>
- Michalak, K. M., Wojciechowska, N., Marzec-Schmidt, K., & Bagniewska-Zadworna, A. (2024). Conserved autophagy and diverse cell wall composition: Unifying features of vascular tissues in evolutionarily distinct plants. *Annals of Botany*, 1–22. <https://doi.org/10.1093/aob/mcae015>
- Michel, J., Lehnert, M., & Quandt, D. (2023). Elevation and cation exchange capacity determine diversity of ferns in a low-montane tropical rainforest in Ecuador. *Journal of Tropical Ecology*, 39(e20), 1–11. <https://doi.org/10.1017/S0266467423000081>
- Moulatlet, G. M., Riaño, K., & Capparelli, M. V. (2024). New register of ferns in soils in contact with hot springs from an active volcano in the Ecuadorian Andes. *Caldasia*, 46(2), 1–27. <https://doi.org/10.15446/caldasia.v46n2.105722>
- Mubai, A., Ambiyar, Irfan, D., & Rasul, M. S. (2023). Flipped direct instruction (FDI): A new practicum learning model in vocational education. *International Journal of Learning, Teaching and Educational Research*, 22(7), 547–565. <https://doi.org/10.26803/ijlter.22.7.29>
- Mwampeta, S. B., Røskaft, E., Realfagbygget, Fyumagwa, R., & Belant, J. L. (2024). A review of field techniques to estimate lion presence and abundance. *Mammal Review*, 54(1), 47–62. <https://doi.org/10.1111/mam.12328>
- Nafisah, J., Amintarti, S., & Rezeki, A. (2023). Identifikasi tumbuhan makro epifit di taman biodiversitas hutan hujan tropis Mandiangin. *Jurnal Jeumpa*, 10(2), 293–302. <https://doi.org/10.33059/jj.v10i2.8625>
- Naiym, J., & Munir, A. (2024). Eksplorasi tumbuhan paku (Pteridophyta) pada tiga daerah topografi berbeda di kawasan taman wisata alam Tirta Rimba Baubau. *Jurnal Penelitian Inovatif*, 4(1), 159–164. <https://doi.org/10.54082/jupin.284>
- Nguyen, T. D. Q., & Nguyen, T. P. L. (2024). Developing social work education in Vietnam: the student field practicum during the pandemic. *Inter-Asia Cultural Studies*, 25(1), 111–124.

- <https://doi.org/10.1080/14649373.2024.2293539>
Nurchayani, P. (2021). Identifikasi jenis dan potensi tumbuhan paku di sekitar curug lontar Desa Karyasari Kecamatan Leuwiliang Kabupaten Bogor. Universitas Islam Negeri Syarif Hidayatullah. <https://repository.uinjkt.ac.id/dspace/handle/123456789/55775>
- Nurdiansyah, S. I. (2023). Sosialisasi dan implementasi standar operasional prosedur kesehatan keselamatan kerja pada praktikum lapangan. *Jurnal Pengabdian Kepada Masyarakat Nusantara (JPKMN)*, 4(4), 4915–4919. <https://doi.org/10.55338/jpkmn.v4i4.2276>
- Petra, M. M., Tripepi, S., & Guardiola, L. (2020). The effects of changes in field practicum hours on student preparedness for social work. *Fieldeducator Simmons Edu*, 10(May), 1–21. <https://fieldeducator.simmons.edu/wp-content/uploads/2020/05/20-235-1.pdf>
- Pradipta, A. R., Hariani, S. A., & Novenda, I. L. (2023). Identification of ferns based on the location and position of the sorus with different elevation in Bondowoso Regency. *Jurnal Biologi Edukasi*, 15(1), 18–28. <https://doi.org/10.24815/jbe.v15i1.30490>
- Pramaisela, D., Abizar, & Risdawati, R. (2024). Pteridophyta yang terdapat di sepanjang pinggir aliran batang kuantan kawasan geopark Silokek Kabupaten Sijunjung sebagai sumbangsih pada pembelajaran biologi SMA/MA. *Jurnal Pendidikan Tambusai*, 8(1), 12360–12369. <https://www.jptam.org/index.php/jptam/article/view/14233>
- Pramudita, D. A., Armando, M. F., Rahmayani, D., Afifah, F. N., Putri, N. R. A., Hartanti, A. N., Safira, R. N., Mahajoeno, E., Indrawan, M., Nazar, I. A., Inocencio E. Buot, J., & Setyawan, A. D. (2023). Species diversity, richness, and conservation status of Pteridophyta in the karst ecosystem of Donorejo Forest, Kaligesing, Purworejo, Indonesia. *International Journal of Tropical Drylands*, 7(1), 16–25. <https://doi.org/10.13057/tropdrylands/t070103>
- Pratomo, R. H. S., & Nur, S. M. (2024). Pengembangan penuntun praktikum zoologi vertebrata berbasis model pembelajaran kooperatif tipe jigsaw terhadap kemampuan berfikir kritis mahasiswa. *Biogenerasi: Jurnal Pendidikan Biologi*, 9(1), 618–627. <https://doi.org/10.30605/biogenerasi.v9i1.2964>
- Puspa, V. R., Fathiya, N., & Muhammad, N. (2023). Inventarisasi dan potensi tumbuhan paku (Pteridophyta) di wisata alam Brayeun sebagai tanaman hias dan obat. *Jurnal Jeumpa*, 10(2), 345–358. <https://doi.org/10.33059/jj.v10i2.8792>
- Qian, H., Kessler, M., & Jin, Y. (2023). Spatial patterns and climatic drivers of phylogenetic structure for ferns along the longest elevational gradient in the world. *Ecography*, 2023(1), 1–13. <https://doi.org/10.1111/ecog.06516>
- Qian, H., Kessler, M., Qian, S., & Zhang, J. (2024). Patterns and drivers of taxonomic and phylogenetic endemism in regional fern floras across the world. *Biological Conservation*, 291, 110506. <https://doi.org/https://doi.org/10.1016/j.biocon.2024.110506>
- Rachmalia, F., Fauziyyah, M. D., Faturrahman, A. D., Ramadhan, M. F., Indriyani, S., Aurina, D. M., Septiasari, A., & Setyawan, A. D. (2023). Studi potensi pemanfaatan tumbuhan vaskuler non-timber di Desa Donorejo, Kecamatan Kaligesing, Purworejo, Jawa Tengah, Indonesia. *Pros Sem Nas Masy Biodiv Indon*, 9(2), 16–28. <https://doi.org/10.13057/psnmbi/m090203>
- Rahmawati, I., & Santhyami. (2023). Diversity of ferns (Pteridophyta) in the Montana Zone, Samiran Village, Selo District, Boyolali Regency, Central Java Province. *International Conference on Biology Education, Natural Science, and Technology (INCOBEST)*, 1(1), 84–90. <https://proceedings.ums.ac.id/index.php/incobest/article/view/3421>
- Ramdana, Mas'anah, M., Aini, Q., Sumarni, T., Marlina, I., Az-Zahra, M., Kamullah, K., Maemunah, M., Nurgamala, N., Juliati, E., Intan, I., Tusa'diah, N., Irwanto, D., Arham, A., Ernawati, E., Azmin, N., & Nasir, M. (2023). Identifikasi tumbuhan paku (Pteridophyta) di wisata air terjun Desa Riamau Kabupaten Bima. *JUSTER: Jurnal Sains Dan Terapan*, 2(1), 61–68. <https://doi.org/10.57218/juster.v2i1.424>
- Randrianarimanana, R., Rakotondrainibe, F., Boucheron-Dubuisson, E., Marline, L., Rakotoarivivo, M., & Reeb, C. (2024). Diversity and distribution of ferns and clubmosses in the eastern canyons of Isalo National Park, Madagascar. *Plant Ecology and Evolution*, 157(1), 3–19. <https://doi.org/10.5091/plecevo.101827>
- Ridhwan, M., Mulyah, E., Hapsari, A. K., & Welani, D. (2023). Study of existence and biodiversity of ferns (Pteridophyta) in the Situ Gintung area, South Tangerang City. *NUCLEUS*, 3(2), 203–209. <https://doi.org/10.37010/nuc.v3i2.1063>
- Riyanto, M., Asbari, M., & Latif, D. (2024). Keefektifan problem based learning terhadap kemampuan berpikir kritis. *JISMA: Journal of Information Systems and Management*, 3(1), 1–5. <https://doi.org/10.4444/jisma.v3i1.744>
- Sejati, A. E., Ihsan, N. H. L., Sugiarto, A., Anasi, P. T., Hariyadi, E., Nasruddin, & Kasmianti, S. (2023). The effect of practicum methods on geographic information system learning outcomes and students' activeness in Senior High School. *E3S Web of Conferences*, 400, 11. <https://doi.org/10.1051/e3sconf/202340001012>
- Sianturi, A., Ridlo, S., & Retnoningsih, A. (2021). Diversity and distribution of ferns at different altitudes in Central Java. *Journal of Physics: Conference Series*, 1918(5), 52016.

- <https://doi.org/10.1088/1742-6596/1918/5/052016>
Singh, S., & Chauhan, R. K. (2024). Investigating IAA'S effects On *Ceratopteris thalictroides*: Implications for spore germination and gametophyte development in Sitamata Wildlife Sanctuary. *American Journal of Agriculture and Environmental Sciences*, 12(1), 51–59.
<https://doi.org/10.5281/zenodo.10623221>
- Sofiyanti, N., Marpaung, A. A., Suriatno, R., & Pranata, S. (2020). Jenis-jenis tumbuhan paku di Pulau Rangsang, Kepulauan Meranti, Riau dan karakteristik morfologi-palinologi. *Jurnal Biologi Tropis*, 20(1), 102–110. <https://doi.org/10.29303/jbt.v20i1.1711>
- Sukmawati, D., Thaharah, A. K., Komariah, E., Theodora, V., Supiyani, A., Nida, S., Husna, A., Ratnaningtyas, I., Enshasy, H. El, & Dailin, D. J. (2024). Isolation of dark septate endophyte (DSE) from ferns (*Pteris vittata*) roots. *International Conference On Mathematics And Science Education (ICMScE)*, 2024. 97–106. <https://doi.org/10.18502/ikls.v8i1.15403>
- Suputra, I. N., & Susanto, H. (2023). *Development of Edutech Startup Learning Kit Industry Lab Integrated Cloud Learning in Learning Practicum SMK and Higher Education Based on Partnership Sharing System in Indonesia*. Atlantis Press International BV.
https://doi.org/10.2991/978-94-6463-178-4_20
- Tariq, A., Butt, B. I., & Javed, Y. (2023). Social work field practicum: From signature pedagogy to the field workers challenges. *Journal of Development and Social Sciences (JDSS)*, 4(IV), 1–11.
[https://doi.org/10.47205/jdss.2023\(4-iv\)01](https://doi.org/10.47205/jdss.2023(4-iv)01)
- Tedam, P., Powell, H., Alfalasi, S. J. H., & Almarqab, E. (2021). Enhancing the social work field Practicum experience through the practice development forum (PDF) at the United Arab Emirates University. *Journal of Teaching in Social Work*, 41(4), 430–442.
<https://doi.org/10.1080/08841233.2021.1954578>
- Tnunay, I. M. Y., & Hanas, D. F. (2020). Keragaman tumbuhan paku sebagai pendukung objek wisata di hutan wisata alam Oeluan, Timor Tengah Utara. *Jurnal Saintek Lahan Kering*, 3(1), 10–12.
<https://doi.org/10.32938/slk.v3i1.1045>
- Ulfa, S. W., Ayunda, D. K., Hasibuan, S. A., & Harahap, A. A. (2023a). Identifikasi spora (warna, bentuk) pada tumbuhan paku yang ada di beberapa kecamatan di Kota Medan. *Jurnal Ilmiah Wahana Pendidikan*, 9(17), 610–621. <https://doi.org/10.5281/zenodo.8320899>
- Ulfa, S. W., Rahmah, S., Latama, F., Boban, A., & Hartady. (2023b). Keanekaragaman tumbuhan paku yang ada di kota Medan. *Ar-Raudah: Jurnal Pendidikan Dan Keagamaan*, 2(1), 61–69.
<https://doi.org/10.30821/ar-raudah.v2i1.69>
- Umair, M., Hu, X., Cheng, Q., Ali, S., & Ni, J. (2023). Distribution patterns of fern species richness along elevations the Tibetan Plateau in China: regional differences and effects of climate change variables. *Frontiers in Plant Science*, 14(May), 1–15. <https://doi.org/10.3389/fpls.2023.1178603>
- Unayah, H., & Widodo. (2023). Karakteristik morfologi paku pohon (*Cyathea contaminans*) dari hutan Kaliurang. *Jurnal Tropika Mozaika*, 2(2), 71–80.
<http://sunankalijaga.org/jurnal/index.php/jtm/article/view/52>
- Wheeler, G. S., Lake, E. C., Mattison, E., & Sutton, G. F. (2024). Host range, biology, and climate suitability of *Callopietia exotica*, a potential biological control agent of Old World climbing fern (*Lygodium microphyllum*) in the USA. *Biological Control*, 188(November 2023), 105410.
<https://doi.org/10.1016/j.biocontrol.2023.105410>
- Widyastuti, G., & Surya, G. (2023). Phytoremediation strategies of arsenic-contaminated surface water near former mining pit 'Kolong' using arsenic hyperaccumulator *Pteris vittata* as mitigation measurement in Bangka Island, Indonesia. In *Arsenic in the Environment: Bridging Science to Practice for Sustainable Development As2021* (1st Editio, pp. 475–476). CRC Press.
<https://doi.org/10.5772/57469.Gonzaga>
- Windari, Perwati, L. K., & Murningsih. (2021). Diversity of fern (Pteridophyta) species in the Jurang Nganten waterfall tourism area, Jepara Regency. *Bioma: Berkala Ilmiah Biologi*, 23(2), 107–111.
<https://doi.org/10.14710/bioma.23.2.107-111>
- Wu, J., & Brock, J. (2023). The invasion of non-native epiphyte *Platyterium bifurcatum* in Auckland's urban forest canopy. *New Zealand Journal of Ecology*, 41(1).
<https://doi.org/10.20417/nzjecol.47.3542>
- Xie, P. J., Ke, Y. T., & Kuo, L. Y. (2023). Modified CTAB protocols for high-molecular-weight DNA extractions from ferns. *Applications in Plant Sciences*, 11(3), 1–9.
<https://doi.org/10.1002/aps3.11526>
- Yi, H., Dong, S., Yang, L., Wang, J., Kidner, C., & Kang, M. (2023). Genome-wide data reveal cryptic diversity and hybridization in a group of tree ferns. *Molecular Phylogenetics and Evolution*, 184. 107801. <https://doi.org/10.1016/j.ympev.2023.107801>
- Yolla, A. S., Damayanti, F., & Gresinta, E. (2022). Keanekaragaman tumbuhan paku terestrial di kawasan hutan pinus gunung Pancar, Bogor. *EduBiologia: Biological Science and Education Journal*, 2(1), 63. <https://doi.org/10.30998/edubiologia.v2i1.11844>
- Yoneoka, K., Hori, K., Kataoka, T., Fujiwara, T., Ebihara, A., & Murakami, N. (2023). Hidden diversity of ferns: *Haplopteris mediosora* have survived as independent gametophytes in Japan. *Acta*

- Phytotaxonomica et Geobotanica*, 74(1), 1–15. <https://doi.org/10.18942/apg.202217>
- Yuhanafia, N., Widodo, A., Budi, L., & Julianto, E. N. (2024). The relevance of practicum implementation on student interests, understanding and learning outcomes in the land surveying subject, Civil Engineering Department, Universitas Negeri Semarang (issue veic 2023). Atlantis Press SARL. https://doi.org/10.2991/978-2-38476-198-2_148
- Zeng, K., Huang, D., Zhang, X., Liu, S., Huang, X., & Xin, G. (2024). Fern species and seasonal variation alter arbuscular mycorrhizal fungal colonization and co-occurrence patterns in the Heishiding Natural Reserve, South China. *Applied Soil Ecology*, 193(February 2023), 105172. <https://doi.org/10.1016/j.apsoil.2023.105172>
- Zhang, K., Zhang, H., Zhang, H., Mei, X., Mas, D., & Jin, S. (2023). Study on species diversity of lycophytes and ferns in Zhejiang province. *Journal of Zhejiang University (Science Edition)*, 50(2), 195–203. <https://doi.org/10.3785/j.issn.1008-9497.2023.02.009>
- Zhang, Q., Wang, J., & Wang, Q. (2021). Effects of abiotic factors on plant diversity and species distribution of alpine meadow plants. *Ecological Informatics*, 61. 101210. <https://doi.org/10.1016/j.ecoinf.2021.101210>