Ethnic disparities in preventable hospital admissions in England

Supplementary materials

Table S1 ICD10 code list for ACSC conditions adopted from Bardsley et al. (2013)

Acute ACS conditions

Cellulitis L03, L04, L08, L88, L980, L983 Dehydration E86 Dental conditions A690, K02-K06, K08, K098, K099, K12, K13 Ear, nose and throat infections H66, H67, J02, J03, J06, J312 Gangrene R02 Gastroentereritis K522, K528, K529 Nutritional deficiencies E40-E43, E55, E643 Pelvic inflammatory disease N70, N73, N74 Perforated/bleeding ulcer K250-K252, K254-K256, K260-K262, K264-K266, K270-K272, K274-K276, K280-K282, 284-K286 Urinary tract infection/Pyelonephritis N10, N11, N12, N136, N390

Chronic ACS conditions

Angina I20, I240, I248, I249 Asthma J45, J46 Chronic obstructive pulmonary disease J20, J41-J44, J47 Congestive heart failure I110, I50, J81 Convulsions and epilepsy G40, G41, 015, R56 Diabetes complications E100-E108, E110-E118, E120-E128, E130-E138, E140-E148 Hypertension I10, I119 Iron deficiency anaemia D501, D508, D509

Vaccine preventable ACS conditions

Influenza J10, J11 Pneumonia J13, J14, J153, J154, J157, J159, J168, J181, J188 Tuberculosis A15, A16, A19 Other vaccine preventable A35-A37, A80, B05, B06, B161, B169, B180, B181, B26, G000, M014

Table S2 Age-, sex-, and deprivation adjusted odds ratios (OR 95% CI) for the combined ACSC outcome. Ethnicity group as recorded by NHS.

| Ethnic group | Age-sex adj. OR (95% CI) | р | Age-sex-deprivation adj. OR (95% CI) | р |
|-----------------|--------------------------|-------|--------------------------------------|-------|
| White British | Ref | - | Ref | - |
| Asian Other | 1.04 (1.02-1.07) | <.001 | 1.00 (0.98-1.03) | .777 |
| Bangladeshi | 1.11 (1.08-1.15) | <.001 | 0.97 (0.94-1.00) | .027 |
| Chinese | 0.49 (0.47-0.52) | <.001 | 0.47 (0.45-0.49) | <.001 |
| Indian | 1.02 (1.00-1.04) | .022 | 0.96 (0.94-0.98) | <.001 |
| Pakistani | 1.70 (1.67-1.72) | <.001 | 1.50 (1.47-1.52) | <.001 |
| Black African | 0.91 (0.89-0.93) | <.001 | 0.89 (0.87-0.92) | <.001 |
| Black Caribbean | 1.01 (0.98-1.03) | .549 | 0.89 (0.87-0.92) | <.001 |
| Other | 1.98 (1.95-2.01) | <.001 | 1.81 (1.78-1.84) | <.001 |
| White Other | 1.15 (1.14-1.17) | <.001 | 1.12 (1.10-1.13) | <.001 |
| White Irish | 0.71 (0.69-0.65) | <.001 | 0.68 (0.66-0.70) | <.001 |
| Mixed | 0.63 (0.62-0.65) | <.001 | 0.60 (0.59-0.62) | <.001 |

Figure S1 Incidence odds ratios (OR) of ACSC admission by ethnic group (as recorded by NHS) relative to White British adjusted for age, sex, and area deprivation. Top: Acute ACSC. Middle: Chronic ACSC. Bottom: Vaccine-preventable diseases. Open circles: risk adjusted for age group and sex. Filled circles: adjusted for age group, sex, and area deprivation.

