

Different strategies for screening and prevention of type 2 diabetes in adults: cost effectiveness analysis

Clare L Gillies, lecturer in medical statistics,¹ Paul C Lambert, senior lecturer in medical statistics,¹ Keith R Abrams, professor of medical statistics,¹ Alex J Sutton, reader in medical statistics,¹ Nicola J Cooper, MRC training fellow in health services research,¹ Ron T Hsu, senior clinical teaching fellow in epidemiology and public health,¹ Melanie J Davies, professor of diabetes medicine,² Kamlesh Khunti, professor of primary care diabetes and vascular medicine³

¹Centre for Biostatistics and Genetic Epidemiology, Department of Health Sciences, University of Leicester, Leicester LE1 7RH

²Department of Cardiovascular Sciences, University of Leicester

³Division of General Practice and Primary Health Care, Department of Health Sciences, University of Leicester

Correspondence to: C L Gillies
clg13@le.ac.uk

doi:10.1136/bmj.39545.585289.25

ABSTRACT

Objective To compare four potential screening strategies, and subsequent interventions, for the prevention and treatment of type 2 diabetes: (a) screening for type 2 diabetes to enable early detection and treatment, (b) screening for type 2 diabetes and impaired glucose tolerance, intervening with lifestyle interventions in those with a diagnosis of impaired glucose tolerance to delay or prevent diabetes, (c) as for (b) but with pharmacological interventions, and (d) no screening.

Design Cost effectiveness analysis based on development and evaluation of probabilistic, comprehensive economic decision analytic model, from screening to death.

Setting A hypothetical population, aged 45 at time of screening, with above average risk of diabetes.

Data sources Published clinical trials and epidemiological studies retrieved from electronic bibliographic databases; supplementary data obtained from the Department of Health statistics for England and Wales, the screening those at risk (STAR) study, and the Leicester division of the ADDITION study.

Methods A hybrid decision tree/Markov model was developed to simulate the long term effects of each screening strategy, in terms of both clinical and cost effectiveness outcomes. The base case model assumed a 50 year time horizon with discounting of both costs and benefits at 3.5%. Sensitivity analyses were carried out to investigate assumptions of the model and to identify which model inputs had most impact on the results.

Results Estimated costs for each quality adjusted life year (QALY) gained (discounted at 3.5% a year for both costs and benefits) were £14 150 (€17 560; \$27 860) for screening for type 2 diabetes, £6242 for screening for diabetes and impaired glucose tolerance followed by lifestyle interventions, and £7023 for screening for diabetes and impaired glucose tolerance followed by pharmacological interventions, all compared with no screening. At a willingness-to-pay threshold of £20 000 the probability of the intervention being cost effective was 49%, 93%, and 85% for each of the active screening strategies respectively.

Conclusions Screening for type 2 diabetes and impaired glucose tolerance, with appropriate intervention for those with impaired glucose tolerance, in an above average risk population aged 45, seems to be cost effective. The cost effectiveness of a policy of screening for diabetes alone, which offered no intervention to those with impaired glucose tolerance, is still uncertain.

INTRODUCTION

In 2000, an estimated 171 million people worldwide had diabetes and numbers are projected to double by 2030.¹ Life expectancy in people with diabetes might be shortened by as much as 15 years.² Currently there is no systematic or structured screening policy for type 2 diabetes in the United Kingdom, though some general guidance has recently been issued by the National Screening Committee.³ One approach to screening would be to screen only for type 2 diabetes, which will allow for early diagnosis and treatment. This might be important as early detection and treatment could prevent future associated microvascular and macrovascular complications. An estimated 50% of people with diabetes are currently undiagnosed,⁴ and at presentation around 20-30% have already developed complications.⁵ An alternative screening approach would be to lower the threshold of the screening test and to screen for impaired glucose tolerance and type 2 diabetes together. As well as allowing for earlier diagnosis of type 2 diabetes, interventions can be administered to those identified with impaired glucose tolerance to attempt to delay the onset of type 2 diabetes. A recent systematic review and meta-analysis of intervention trials for prevention of type 2 diabetes⁶ found both lifestyle and pharmacological interventions significantly reduced the risk of type 2 diabetes in people with impaired glucose tolerance.

As no definitive trials have examined the effectiveness of screening for type 2 diabetes or impaired glucose tolerance,^{7,8} assessment of such policies has so far been conducted through simulation studies. Several decision models have been compiled that have assessed either the clinical and cost effectiveness of

interventions to prevent type 2 diabetes⁹⁻¹⁶ or strategies for screening and early detection of diabetes.^{7,17-20} Previous models of screening for type 2 diabetes alone have generally assessed the impact of early treatment on cardiovascular events, though some additionally included microvascular events such as retinopathy. Overall most of the models produced favourable results for screening, but cost effectiveness varied with age group screened and the population targeted for screening. Only two studies reported costs for a UK setting,^{7,19} one of which had a limited time

horizon of five years.¹⁹ Both of these studies concluded there was still uncertainty concerning the cost effectiveness of screening for diabetes.

Of the eight models assessing cost effectiveness of interventions for prevention of diabetes, only three included costs of identifying individuals with impaired glucose tolerance.^{10,12,16} The time horizon over which the models were run ranged from just three years after the intervention up to the expected lifetime of the population. Models used data from various sources from published trials, epidemiological studies, and national statistics. In general data were limited to a few sources. All models compared a strategy of interventions against no interventions, rather than screening for impaired glucose tolerance followed by interventions, compared with no screening. All but one model simulated populations where all individuals had impaired glucose tolerance at the start of the model and the end state was development of diabetes, or death, hence only a limited section of the disease pathway was modelled. Also the models did not take into account that screening for impaired glucose tolerance will at the same time allow individuals with undiagnosed diabetes to be identified, thus allowing for early treatment and possibly reducing rates of complications. Hence, while these studies offer an assessment of the cost effectiveness of interventions for prevention of diabetes, none assessed the impact of screening followed by interventions on the whole disease pathway. In 2007 Waugh et al assessed screening or intervention strategies for type 2 diabetes in a thorough review of previous decision models.⁷

We compared three active screening strategies: (a) a one-off screening for type 2 diabetes; (b) screening for impaired glucose tolerance and type 2 diabetes and intervening with lifestyle interventions in those with impaired glucose tolerance; and (c) as for (b) but with pharmacological interventions. We compared these three active screening strategies against a fourth strategy of no screening (current practice). The full pathway from screening, to interventions and treatment for type 2 diabetes, all the way through to death, was modelled. This model directly compares the two alternative approaches of screening for type 2 diabetes alone or screening for impaired glucose tolerance and type 2 diabetes together. When modelling the effectiveness of interventions, we used all data from relevant randomised controlled trials⁶ and included uncertainty around model inputs when appropriate. By carrying out several sensitivity analyses we investigated the essential elements that affect the cost and clinical effectiveness of different screening policies.

METHODS

The hybrid model consists of a decision tree and a Markov model (fig 1). The decision tree comprises three main arms, representing no screening, screening for undiagnosed type 2 diabetes, and screening for impaired glucose tolerance and undiagnosed diabetes, with either lifestyle or pharmacological interventions applied in those with impaired glucose tolerance.




Fig 1 | Decision model comparing no screening, screening for type 2 diabetes, and screening for impaired glucose tolerance and diabetes and intervening to delay or prevent type 2 diabetes with either lifestyle or pharmacological interventions

Table 1 | Estimates used to determine parameters for decision model

Parameter	Distribution	Value (SE)	Source(s)
Data for decision tree			
Prevalences	Dirichlet	Normal glucose tolerance 83%; impaired glucose tolerance 12%; type 2 diabetes 5%	STAR study ²⁴
Screening test efficiency	Multi-nominal	For type 2 diabetes: sensitivity 89.5%, specificity 91.3%; for impaired glucose tolerance and type 2 diabetes: sensitivity 59.4%, specificity 88.0%	STAR study ²⁴
Transition rates (per 100 person years)			
Normal to impaired glucose tolerance:			
<65 years	Log normal	1.66 (0.08)	Baltimore study ³³
≥65 years	Log normal	2.49 (0.11)	
Impaired glucose tolerance to type 2 diabetes	Log normal	1.96 (0.25)	12 studies ²⁵⁻³⁶
Time spent with undetected diabetes (years)	Log normal	1.65 (0.68)	Harris ³⁷
Mortality rates (per 100 person years)			
45-54 years	—	0.32	DoH statistics (2000)
55-64 years	—	0.84	
65-74 years	—	2.36	
75-84 years	—	6.09	
≥85 years	—	15.68	
Increased risk of death with diabetes (hazard ratio)	Log normal	0.756 (0.087)	DECODE ³⁸
Increased risk of death for 1% increase in HbA _{1c} (hazard ratio)	Log normal	0.104 (0.039)	Rossing ³⁹
Intervention effects on risk of developing type 2 diabetes (hazard ratio)			
Lifestyle v standard treatment	Log normal	−0.646 (0.099)	12 studies ⁶
Antidiabetic drugs v placebo	Log normal	−0.425 (0.141)	9 studies ⁶
HbA _{1c}			
Undiagnosed diabetes	Normal	9.0% (0.056)	UKPDS ⁴⁰
Screen detected diabetes	Normal	7.0% (0.028)	UKPDS ⁴¹
Clinically detected diabetes	Normal	7.9% (0.042)	UKPDS ⁴¹
Utilities			
Undiagnosed diabetes	Normal	0.788 (0.020)†	ADDITION ³⁴
Screen detected diabetes	Normal	0.788 (0.020)‡	ADDITION ³⁴
Clinically detected diabetes	Normal	0.771 (0.035)‡	UKPDS ^{43 44}
Costs*			
Screening tests:			
FPG test	—	£0.40/person	NHS (2006)
OGTT test	—	£1.30/person	
Nurse cost	—	£26/hour	Curtis ⁴⁵
Metformin intervention	—	£16.10/year	NHS (2006)
Lifestyle intervention:			
Year 1	—	£398/year	Avenell ⁹
Subsequent years	—	£280/year	
Undiagnosed diabetes:			
Year before diagnosis	—	£114/year	Gulliford, ⁴⁶ Curtis ⁴⁵
Years 2-5 before diagnosis	—	£22/year	
Diagnosed diabetes:			
Screen detected	Normal	£2490 (53.3)/year	UKPDS ⁴⁷
Clinically detected	Normal	£2756 (63.1)/year	

FPG=fasting plasma glucose; OGTT=oral glucose tolerance test.

*Costs are standardised to 2006.⁴⁸

†Constant for all time spent with undetected type 2 diabetes.

‡Starting utility, which was then decreased for each year spent with diabetes because of predicted increases in complications, based on UKPDS data.^{44 45}

Individuals who have already been identified as having type 2 diabetes are excluded from the screening process. The decision tree uses prevalence of impaired glucose tolerance and undiagnosed type 2 diabetes and

estimates sensitivity and specificity of a screening test to determine how many individuals from the population start in each state of the Markov model. The Markov model consists of seven states: normal glucose

tolerance, undiagnosed impaired glucose tolerance, diagnosed impaired glucose tolerance, death, and three states for people with diabetes (undiagnosed, diagnosed clinically, or diagnosed through screening, either from a screening test or because they are diagnosed with impaired glucose tolerance initially and hence enter a surveillance programme). We ran four Markov models simultaneously, one for each of the screening strategies. Whether type 2 diabetes and impaired glucose tolerance are diagnosed or undiagnosed determines whether the patients receive relevant treatments or interventions, and they are modelled accordingly in terms of transition rates to other states. For example, individuals identified with impaired glucose tolerance receive an intervention and the estimated intervention effect slows their progression to development of diabetes. Each model cycle represents one year and the model is run for a time horizon of 50 years. Table 1 summarised all the model inputs. When more than one estimate was available for a parameter, we pooled estimates using a Bayesian random effects meta-analysis within the comprehensive decision model. Model results include both clinical and cost effectiveness outcomes, with cost per quality adjusted life year (QALY) being the primary outcome. We investigated both an undiscounted model and a model with costs and benefits discounted at 3.5%

annually, as recommended by the National Institute for Health and Clinical Excellence.²¹

The hybrid model was implemented within WinBUGS using a Bayesian comprehensive decision modelling approach.²² We adopted this approach because of its flexibility in terms of statistical modelling and it enabled us to include and propagate all uncertainty in parameters throughout the model.²² We assumed non-informative prior distributions for all model parameters. Model parameters were estimated by using Markov chain Monte Carlo simulation methods.²³ Results are based on a sample of 20 000 simulations, following a “burn in” of 10 000, and we assessed convergence of the Markov chain by visually inspecting trace plots and by running multiple chains with different initial values.²³ We have reported the results from the decision model with 95% credibility intervals, which are analogous to confidence intervals.

Data for the decision tree

The base case scenario for the model was a one-off screening for a population aged 45, in whom type 2 diabetes had not previously been diagnosed. Data for the decision tree—that is, test sensitivity and specificity and prevalence of impaired glucose tolerance and type 2 diabetes—were taken from the screening those at risk (STAR) study.²⁴ For this study, individuals aged 40–75

Table 2 | Clinical and cost outcomes from decision model, where prevalence of impaired glucose tolerance was 15% and type 2 diabetes 7.5%, and sensitivity and specificity of screening tests was 85% and 80%, respectively. Figures are mean values per person (95% credible intervals) for no screening and mean difference from or compared with no screening (95% credible intervals) for all other strategies

			Screening for diabetes and impaired glucose tolerance	
	No screening	Screening for diabetes only	Lifestyle interventions	Pharmacological interventions
Undiscounted				
Total life years	30.34 (27.75 to 32.86)	0.06 (0.02 to 0.12)	0.15 (0.08 to 0.22)	0.13 (0.06 to 0.20)
QALYs	28.06 (23.49 to 32.01)	0.07 (−0.03 to 0.18)	0.22 (0.08 to 0.36)	0.17 (0.03 to 0.32)
Years spent without diabetes	20.85 (10.36 to 29.45)	—	0.33 (0.21 to 0.43)	0.20 (0.10 to 0.37)
Lifetime risk of diabetes (%)	64.55 (18.02 to 91.83)	—	−0.98 (−0.50 to −1.42)	−0.54 (−0.21 to −1.17)
Total cost	17 290 (5746 to 39580)	730 (−9 to 2341)	610 (−373 to 2693)	579 (−428 to 2658)
Cost per life year gained	—	11 460	4179	4768
Cost per QALY gained	—	8681	2863	3429
Cost per case prevented	—	—	62 810	105 000
Probability of cost effectiveness at willingness to pay threshold per QALY (%):				
£20 000		68.1	98.6	94.7
£30 000		76.5	99.6	97.3
Discounted at 3.5% a year for both costs and benefits				
Total life years	18.19 (17.25 to 18.98)	0.02 (−0.01 to 0.05)	0.05 (0.03 to 0.08)	0.05 (0.02 to 0.07)
QALYs	17.13 (15.02 to 18.49)	0.03 (−0.02 to 0.09)	0.09 (0.03 to 0.17)	0.07 (0.01 to 0.15)
Years spent diabetes free	13.69 (7.99 to 17.08)	—	0.17 (0.11 to 0.23)	0.11 (0.06 to 0.19)
Total cost	7636 (2636 to 19 370)	587 (61 to 1525)	580 (−103 to 1760)	528 (−163 to 1719)
Cost per life year gained	—	23 710	10 900	11 690
Cost per QALY gained	—	14 150	6242	7023
Probability of cost effectiveness at willingness to pay threshold per QALY (%):				
£20 000		48.6	93.0	85.0
£30 000		60.8	97.4	91.6

(white) or 25-75 (non-white) from 15 general practices in Leicestershire who had at least one recognised risk factor for type 2 diabetes were invited for screening. Risk factors included a known history of coronary heart disease, hypertension, dyslipidaemia, cerebrovascular disease, a first degree relative with type 2 diabetes, and a body mass index (BMI) >25. Therefore the screening data included in the primary model were from a population considered to be “at risk” of type 2 diabetes. For the base case model we used data only from white patients, though we used the data on South Asians for sensitivity analyses to assess results for different ethnic groups.

Transition rates and HbA_{1c} concentrations

To estimate annual transition rates we used several sources, including epidemiological studies and clinical trials.²⁵⁻³⁶ To estimate the annual transition rate from undiagnosed to clinically diagnosed diabetes, we used the estimated average time people have diabetes before being diagnosed.³⁷ We estimated the effects of interventions on the transition from impaired glucose tolerance to diabetes using studies identified in a recent meta-analysis of lifestyle and pharmacological intervention trials.⁶ Death rates were taken from Department of Health statistics for England and Wales for 2000 and were increased for people with diabetes compared with those without.³⁸ For the three diabetic states (undiagnosed, clinically diagnosed, and screen detected) death rates varied depending on predicted HbA_{1c} (haemoglobin A_{1c}) concentrations.³⁹ HbA_{1c} was predicted to be highest in people with undiagnosed diabetes, as they are yet to receive any interventions, and was estimated by using HbA_{1c} concentrations at entry to the UK prospective diabetes study⁴⁰ before




Fig 2 Cost effectiveness acceptability curves for each of three active screening strategies compared with no screening (discounted estimates)

treatment began. We expected HbA_{1c} concentrations to be the best controlled in people with diabetes detected by screening because of early detection, and estimated levels using the 10 year average from the intensively treated group in the UK prospective diabetes study.⁴¹ For people with clinically diagnosed diabetes, we used the HbA_{1c} concentrations of the group receiving conventional treatment in the UK prospective diabetes study.⁴¹

Quality of life variables

For the states of normal glucose tolerance, undiagnosed impaired glucose tolerance, and diagnosed impaired glucose tolerance, we assumed the utility value to be that of full health and set at 1. We calculated

Table 3 | Results (undiscounted) of sensitivity analyses for varying prevalence rates of impaired glucose tolerance, normal glucose tolerance, and type 2 diabetes

			Screening for type 2 diabetes and impaired glucose tolerance	
Prevalence*	No screening	Screening for type 2 diabetes only	Lifestyle interventions	Pharmacological interventions
QALY				
83/12/5	28.06 (23.49 to 32.01)	28.12 (23.58 to 32.08)	28.26 (23.74 to 32.23)	28.22 (23.69 to 32.18)
70/20/10	28.26 (24.72 to 31.18)	28.26 (24.79 to 31.14)	28.47 (25.02 to 31.34)	28.41 (24.96 to 31.29)
10/60/30	23.75 (21.82 to 25.58)	24.16 (22.40 to 25.85)	24.91 (23.15 to 26.55)	24.67 (22.89 to 26.35)
Total cost (£)				
83/12/5	17 290 (5746 to 39 580)	18 040 (7083 to 39 970)	17 910 (7124 to 39 740)	17 900 (7061 to 39 710)
70/20/10	21 320 (9132 to 41 270)	22 780 (12 470 to 41 840)	22 620 (12 650 to 41 370)	22 560 (12 540 to 41 420)
10/60/30	38 440 (19 740 to 49 690)	42 580 (32 660 to 51 190)	41 980 (33 990 to 49 980)	41 830 (33 530 to 50 090)
Cost per QALY gained (£)				
83/12/5	—	8681	2863	3429
70/20/10	—	8617	3203	3809
10/60/30	—	8464	3148	3781
Probability (%) of being cost effective at willingness to pay threshold of £20 000/£30 000 per QALY				
83/12/5	—	68/76	99/100	95/97
70/20/10	—	68/76	98/99	93/96
10/60/30	—	68/76	98/99	93/96

*Impaired glucose tolerance/normal glucose tolerance/type 2 diabetes.

Table 4 | Results (undiscounted) of sensitivity analyses for varying compliance rates

			Screening for type 2 diabetes and impaired glucose tolerance	
Compliance	No screening	Screening for type 2 diabetes only	Lifestyle interventions	Pharmacological interventions
Compliance with screening (%)				
QALY:				
100	28.06 (23.49 to 32.01)	28.12 (23.58 to 32.08)	28.26 (23.74 to 32.23)	28.22 (23.69 to 32.18)
70	28.06 (23.49 to 32.01)	28.07 (23.52 to 32.05)	28.17 (23.64 to 32.16)	28.14 (23.60 to 32.13)
50	28.06 (23.49 to 32.01)	28.04 (23.51 to 32.04)	28.13 (23.61 to 32.13)	28.10 (23.59 to 32.11)
Total cost (£):				
100	17 290 (5746 to 39 580)	18 040 (7083 to 39 970)	17 910 (7124 to 39 740)	17 900 (7061 to 39 710)
70	17 290 (5746 to 39 580)	18 070 (6777 to 39 800)	18 080 (6957 to 39 620)	18 070 (6907 to 39 710)
50	17 290 (5746 to 39 580)	17 870 (6409 to 39 750)	17 930 (6705 to 39 680)	17 910 (6671 to 39 690)
Cost (£) per QALY gained:				
100	—	8681	2863	3429
70	—	8732	3112	3800
50	—	8743	3515	4192
Probability of being cost effective at willingness to pay threshold of £20 000/£30 000 per QALY (%):				
100	—	68/76	99/100	95/97
70	—	69/77	98/99	93/96
50	—	68/77	97/98	92/95
Compliance with interventions (%)				
QALY:				
100	28.06 (23.49 to 32.01)	28.12 (23.58 to 32.08)	28.26 (23.74 to 32.23)	28.22 (23.69 to 32.18)
70	28.06 (23.49 to 32.01)	28.12 (23.58 to 32.08)	28.22 (23.69 to 32.18)	28.19 (23.66 to 32.15)
50	28.06 (23.49 to 32.01)	28.12 (23.58 to 32.08)	28.19 (23.66 to 32.15)	28.17 (23.64 to 32.13)
Total cost (£):				
100	17 290 (5746 to 39 580)	18 040 (7083 to 39 970)	17 910 (7124 to 39 740)	17 900 (7061 to 39 710)
70	17 290 (5746 to 39 580)	18 040 (7083 to 39 970)	18 140 (7343 to 39 950)	18 040 (7209 to 39 880)
50	17 290 (5746 to 39 580)	18 040 (7083 to 39 970)	18 261 (7455 to 40 050)	18 120 (7302 to 39 960)
Cost (£) per QALY gained:				
100	—		2863	3429
70	—		4947	5039
50	—		6634	6243
Probability of being cost effective at willingness to pay threshold of £20 000/£30 000 per QALY (%):				
100	—		99/100	95/97
70	—		94/97	89/94
50	—		88/93	84/90

utilities for those with undiagnosed and screen detected diabetes from EQ-5D data, using data on individual patients made available by the Leicester arm of the ADDITION study.⁴² The data were of a screen detected sample population with type 2 diabetes at baseline. For people with clinically diagnosed diabetes, utilities were taken from those reported by the UK prospective diabetes study as this comprised a clinically detected sample.⁴³ The utility for undiagnosed diabetes was kept constant for the whole duration spent in this state as we assumed that if complications developed, which reduced the quality of life, then a diagnosis would be made. For the states of clinically and screen detected diabetes we needed to account for the fact that duration of diabetes would lead to an increased number of complications and hence a reduction in the utility value. This was done by using reported complication rates, modelled for duration of diabetes and adjusted for estimated HbA_{1c}

concentrations in each group and their estimated effect on utility values.^{43 44} Hence, utilities decreased for each year of duration of diabetes, to reflect increasing incidence of complications. Because of a higher predicted HbA_{1c} concentration, the utility value was lower at diagnosis and decreased marginally more rapidly in individuals clinically diagnosed compared with those who were screen detected.

Economic variables

We estimated costs from various sources. Screening costs included the costs of an initial screening test of fasting plasma glucose and a confirmatory oral glucose tolerance test in those who tested positive. We estimated the cost of nurse time of 5 minutes for the screening test and 25 minutes for the oral glucose tolerance test.⁴⁵ People with undiagnosed diabetes incur costs before diagnosis because of increased visits to the general practitioner and prescriptions,⁴⁶ with a

reported average of three additional visits the year before diagnosis and an average of 1.4 additional visits in the two to five years before diagnosis. An estimation of these costs was included.⁴⁵ For lifestyle interventions we included dietitian costs and costs of twice weekly group exercise sessions, as detailed in a previous study.⁹ Costs of pharmacological interventions were based on 250 mg of metformin three times a day, the standard dose used by most intervention studies. For people with diagnosed diabetes, we took average annual costs of antidiabetic treatment, implementation of treatment, and costs of complications from the UK prospective diabetes study.⁴⁷ For the people with diabetes detected at screening, in whom we would expect costs of complications to be lower, we used costs from the intensively treated arm of the UK prospective diabetes study. For those with clinically diagnosed diabetes, which represents how individuals are diagnosed currently, we used the reported costs of the conventionally treated group. All costs are reported in 2006 UK £, standardised by using inflation indices.⁴⁵

Sensitivity analyses and model extensions

We carried out sensitivity analyses using a range of values of prevalence of disease, as well as compliance levels to both screening and interventions. Changing prevalence allows us to assess the effectiveness of the screening strategies for different “at risk” populations. The effects of compliance to both screening and interventions were also important as we assumed 100% compliance to both in the base case model, which could never be achieved in practice.

To evaluate the robustness of the model we also carried out sensitivity analyses on model inputs, particularly those that were estimated from only one or two sources or were thought to be important drivers in the model. These were sensitivities of screening tests,

costs of interventions, costs of diabetes, effectiveness of interventions, previous distributions on the standard deviations between studies of the four meta-analyses run within the model, and the time horizon the model was run for.

For the base case scenario we considered only a one-off screening at age 45. The model was extended further to assess the impact of having one or two additional screenings, at age 50 and 60. This was done by applying the test sensitivities from the STAR study to the numbers in the states of undiagnosed impaired glucose tolerance and type 2 diabetes at the corresponding model cycle and moving the individuals to the relevant diagnosed state.

Though the base case model used prevalences and test sensitivities and specificities of a white population, the effect of screening a South Asian or a mixed race population is also relevant in the UK. South Asians are thought to have a greater risk of type 2 diabetes, with a greater prevalence of impaired glucose tolerance and a higher transition rate to type 2 diabetes. We extended this model with data from the STAR study and estimated the transition rate from impaired glucose tolerance to type 2 diabetes from the Indian diabetes prevention programme.⁴⁸

RESULTS

Table 2 shows clinical and cost effectiveness outcomes for an undiscounted model and a model discounted for both costs and benefits at 3.5% a year. Discounted costs for each QALY gained, compared with no screening, were £14 150 (€17 560; \$27 860) for type 2 diabetes screening, £6242 for screening for diabetes and impaired glucose tolerance with lifestyle interventions, and £7023 for screening for both diabetes and impaired glucose tolerance with pharmacological interventions. Costs were lower in the undiscounted model: £8681,

Table 5 | Results of model extensions for number of screens (undiscounted)

			Screening for type 2 diabetes and impaired glucose tolerance	
No of screens	No screening	Screening for type 2 diabetes only	Lifestyle interventions	Pharmacological interventions
QALY				
1	28.06 (23.49 to 32.01)	28.12 (23.58 to 32.08)	28.26 (23.74 to 32.23)	28.22 (23.69 to 32.18)
2	28.06 (23.49 to 32.01)	28.13 (23.74 to 32.06)	28.56 (24.74 to 32.30)	28.44 (24.45 to 32.24)
3	28.06 (23.49 to 32.01)	28.15 (23.86 to 32.16)	28.80 (25.04 to 32.32)	28.62 (24.70 to 32.26)
Total cost (£)				
1	17 290 (5746 to 39 580)	18 040 (7083 to 39 970)	17 910 (7124 to 39 740)	17 900 (7061 to 39 710)
2	17 290 (5746 to 39 580)	18 850 (7491 to 40 980)	19 300 (7570 to 41 160)	19 150 (7468 to 41 150)
3	17 290 (5746 to 39 580)	19 670 (7735 to 42 110)	20 220 (7740 to 42 210)	19 860 (7621 to 42 210)
Cost per QALY gained (£)				
1	—	8681	2863	3429
2	—	9544	2777	3317
3	—	10 360	2966	3517
Probability of being cost effective at willingness to pay threshold of £20 000/£30 000 per QALY (%)				
1	—	68/76	99/100	95/97
2	—	57/66	99/100	96/98
3	—	54/64	99/100	97/99

Table 6 | Results of model extensions for different ethnic groups (undiscounted)

			Screening for type 2 diabetes and impaired glucose tolerance	
Ethnic group	No screening	Screening for type 2 diabetes only	Lifestyle interventions	Pharmacological interventions
QALY				
White	28.06 (23.49 to 32.01)	28.12 (23.58 to 32.08)	28.26 (23.74 to 32.23)	28.22 (23.69 to 32.18)
South Asian	25.24 (20.65 to 30.79)	25.35 (20.83 to 30.91)	25.47 (20.96 to 31.02)	25.43 (20.92 to 30.98)
Mixed*	27.10 (23.79 to 30.31)	27.18 (23.88 to 30.39)	27.32 (24.02 to 30.53)	27.27 (23.99 to 30.53)
Total cost (£)				
White	17 290 (5746 to 39 580)	18 040 (7083 to 39 970)	17 910 (7124 to 39 740)	17 900 (7061 to 39 710)
South Asian	28 250 (10 170 to 55 120)	29 390 (12 270 to 55 490)	29 420 (12 500 to 55 220)	29 480 (12 550 to 55 270)
Mixed*	22 145 (8345 to 41 657)	23 051 (9820 to 42 131)	22 973 (9809 to 41 962)	22 976 (11 885 to 42 006)
Cost per QALY gained (£)				
White	—	8681	2863	3429
South Asian	—	8168	4657	5643
Mixed*	—	8523	3555	4497
Probability of being cost effective at willingness to pay threshold of £20 000/£30 000 per QALY (%)				
White	—	68/76	99/100	95/97
South Asian	—	68/75	89/94	83/88
Mixed*	—	69/77	98/99	96/98
*Modelled as 30% South Asian and 70% white.				

*Modelled as 30% South Asian and 70% white.

£2863, and £3429 for every QALY gained, respectively. At a willingness to pay threshold of £20 000 per QALY the probability of each strategy being cost effective was 49% for screening for type 2 diabetes only, 93% for screening for both diabetes and impaired glucose tolerance and lifestyle interventions, and 85% for screening for both diabetes and impaired glucose tolerance and pharmacological intervention. Figure 2 shows cost effectiveness acceptability curves, illustrating the probability of cost effectiveness over a range of willingness to pay thresholds.

Discounted QALYs gained compared with no screening were 0.03 (−0.02 to 0.09) for diabetes screening, 0.09 (0.03 to 0.17) for screening and lifestyle interventions, and 0.07 (0.01 to 0.15) for screening with pharmacological interventions. Both the intervention strategies showed potential benefits in terms of average years spent without diabetes and cases of diabetes prevented. Although clinical effects seem small, it must be remembered they are average gains across a population, in which only 17% had either impaired glucose tolerance or undiagnosed type 2 diabetes at the time of screening.

Tables 3 and 4 show the results of the more important sensitivity analyses (undiscounted). Increasing the prevalence of impaired glucose tolerance and type 2 diabetes decreased the QALYs and increased total costs of each screening strategy. The comparisons of the three active screening/intervention strategies compared with no screening remained fairly constant in terms of costs per QALY and probability of cost effectiveness (table 3). When we lowered compliance with screening, the impact on results was also minimal (table 4). Reducing compliance with interventions, however, had a greater impact in that the total costs and cost per QALY gained increased for both the

screening/intervention strategies. The probability that these strategies were cost effective compared with no screening still remained high, with an estimated probability of 88% for screening with lifestyle interventions and 84% for screening with pharmacological interventions at the willingness to pay threshold of £20 000.

Other sensitivity analyses did not change the results enough to alter the conclusions of the model. Increasing the costs of both lifestyle and pharmacological interventions by a factor of 10 reduced the probabilities of cost effectiveness of their respective screening strategies to 73% and 93%, at the willingness to pay threshold of £20 000. Increasing the costs of diabetes by a factor of two reduced the probability of cost effectiveness to 49% for screening for type 2 diabetes only, 93% for screening with lifestyle interventions, and 85% for screening with pharmacological interventions at the same threshold. As we increased the time horizon the model was run for, the probability of the three active screening strategies being cost effective compared with no screening increased. This is because the benefits of screening or interventions are not all immediate and most occur in later years of the model, when type 2 diabetes is either delayed or complications are reduced through early diagnosis and treatment. The intervention strategies became cost effective when we considered a time horizon of at least 30 years (probability of being cost effective of 0.97 for lifestyle and 0.91 for pharmacological interventions at the willingness to pay threshold of £20 000). Overall, the model's conclusions were robust to changes made to the sensitivity analyses, giving strength to the conclusions.

Tables 5 and 6 give the results of the model extensions as undiscounted estimates. Increasing the

WHAT IS ALREADY KNOWN ON THIS TOPIC

In people with impaired glucose tolerance interventions are clinically and cost effective
Screening for type 2 diabetes to allow early detection might be cost effective in certain groups

WHAT THIS STUDY ADDS

Modelling the whole screening and intervention pathway from screening to death shows that screening for type 2 diabetes and impaired glucose tolerance, followed by interventions, seems to be cost effective compared with no screening

Uncertainty still exists concerning the cost effectiveness of screening for type 2 diabetes alone

Screening populations with a higher prevalence of glucose intolerance might result in better clinical outcomes, although cost effectiveness seems unaffected

number of screenings of the population increased both total costs and QALYs, which resulted in minimal increases in the cost per QALY for each of the three active strategies (table 5). When we ran the model for a South Asian cohort, results for QALYs were lower because of a higher prevalence of type 2 diabetes at the start of the model and an increased rate of transition to diabetes (table 6). Neither increasing the number of screens nor considering different ethnic cohorts led to a change in the overall model conclusions, in that both the strategies involving interventions for prevention of diabetes seem to be cost effective compared with no screening in an “at risk” population.

DISCUSSION

Screening for impaired glucose tolerance in people at risk of diabetes and intervening with either lifestyle or pharmacological interventions is a cost effective health policy. Although screening for type 2 diabetes alone gave a relatively low predicted incremental cost per QALY of £14 150, because of uncertainty in the model the probability of this strategy being cost effective was only 49% at the willingness to pay threshold of £20 000.

Strengths and weaknesses

Previous studies have compared the cost and clinical effectiveness of intervening in people with impaired glucose to delay onset of type 2 diabetes.^{9–16} Results were all favourable in terms of cost and clinical effectiveness but as the models were designed to assess the effectiveness of interventions rather than screening and intervening, none of the models included a state of undiagnosed diabetes and assumed management of diabetes started as soon as the disease developed. Our model considered the whole screening and intervention pathway from screening to death and a comparison of different approaches to diabetes screening and prevention.

Differences in clinical outcomes between the no screening strategy and the three active screening strategies were small, partly because they were reported as an average for a screened population with mixed glucose tolerance. Also microvascular and macrovascular outcomes were not measured individually in this model, which might show benefits from the early detection or delay of type 2 diabetes.

Our model makes several assumptions. No transition was allowed from normal glucose tolerance to diabetes without first passing through impaired glucose tolerance. This is because it is clinically unlikely that an individual would change from normal glucose tolerance to diabetes within a year, which is one model cycle. No transition was allowed from diabetes back to impaired glucose tolerance or from impaired to normal glucose tolerance. This is clinically accurate because once an individual has a diagnosis of type 2 diabetes, even if their glucose tolerance improves, they are still clinically defined as having diabetes. Also once an individual has had impaired glucose tolerance, even if their glucose tolerance improves their future risk of diabetes is probably more similar to that in individuals with impaired glucose tolerance rather than those who have always had normal glucose tolerance.

Another assumption was that the HbA_{1c} concentration of those with diabetes who were clinically diagnosed would be similar to the 10 year average of an intensively treated group of people with diabetes from the UK prospective diabetes study.⁴¹ This assumption was made in the absence of long term clinical data on individuals whose diabetes was detected by screening. Although 10 year averages of HbA_{1c} concentrations were used for people with diabetes, when we ran our model for longer time horizons the HbA_{1c} concentrations were potentially underestimated, which means complication rates and their effects on utilities and mortality might also be moderately underestimated. Further data are needed on how HbA_{1c} concentration could be expected to increase over time to allow more accurate modelling.

Screening costs incorporated within the model included only costs of the test and the nurse's time, therefore representing the costs of opportunistic screening. We did not include further costs of establishing systematic screening, such as the identification of eligible patients, the issuing of invitations to screening, and the chasing up of non-attenders. In practice, these additional costs would be small for each individual screened, particularly if screening was incorporated into current health checks. When modelling costs of treatment and complications associated with diabetes, we used the average yearly costs taken from the UK prospective diabetes study. As costs would be expected to start off low and then increase, this means that costs of diabetes might be initially overestimated when an individual receives the diagnosis and eventually underestimated by this model. In addition, as average costs were used, we did not account for issues of competing risks of complications associated with diabetes. Unfortunately, yearly data on costs of diabetes, or how the occurrence of complications impacted on the probability of other complications occurring, were not available to enable us to model costs more accurately. The issue of competing risks arises not just for costs but also for the annual probabilities of complications. Ideally, we need data on individual patients to enable the correlation structure

in both the probabilities and costs to be appropriately accounted for.

As we ran the model for a time horizon of 50 years, the screened population (aged 45 at the start) aged with each cycle of the model, thus, when possible, we incorporated time dependent model parameters. For some parameters, such as the treatment intervention effects, however, we assumed that the effect was constant over time. Additionally, although compliance was high in the intervention trials from which estimates of their effectiveness were obtained, it is still to be determined whether compliance could be maintained outside a trial setting. Therefore long term compliance with interventions is an important consideration. Sensitivity analyses of compliance with interventions found that even with compliance rates as low as 50%, the screening strategies involving either lifestyle or pharmacological interventions were still cost effective when compared with a strategy of no screening.

Conclusions

A policy of a one-off screening for type 2 diabetes and impaired glucose tolerance, with appropriate intervention for those identified with impaired glucose tolerance, seems to be cost effective in an "at risk" population. Changing compliance with screening or interventions or increasing the number of screenings did not change the conclusions of the model. Given the uncertainty in the results presented here, particularly for the assessment of screening for type 2 diabetes, further research is needed on the long term clinical effects of early diagnosis. Furthermore, to model the two strategies that involved interventions more accurately, we require additional information on long term compliance with interventions and their potential harms and benefits.

We thank the STAR study, in particular Jenny Tringham, and the Leicester arm of the ADDITION study, for providing the data on individual patients that were used for the analyses. We also thank Philip Clarke for advice on the UKPDS outcomes model.

Contributors: CLG performed the data extraction and analyses, wrote the first draft of the article, and is guarantor. KRA and PCL gave detailed advice at all stages of the analyses. All authors contributed to the writing of the paper and gave substantial advice and input into the study. KRA and KK had the initial idea for this project.

Funding: CLG is funded jointly by the UK Medical Research Council and the Economic and Social Research Council, under an interdisciplinary postgraduate research studentship in the social and medical sciences. NJC is funded by a Medical Research Council training fellowship in health services research.

Competing interests: MJD and KK have received sponsorship for attending conferences and small honorariums and funding for research from pharmaceutical companies that manufacture hypoglycaemic and anti-obesity drugs. KRA has also received funding for research from pharmaceutical companies that manufacture hypoglycaemic and anti-obesity drugs and has acted as a paid consultant to consultancy companies who undertake work for the healthcare industry generally.

Ethical approval: Not required.

Provenance and peer review: Not commissioned; externally peer reviewed.

- 1 Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004;27:1047-53.

- 2 Donnelly R, Emslie-Smith AM, Gardner ID, Morris AD. ABC of arterial and venous vascular disease: Vascular complications of diabetes. *BMJ* 2000;320:1062-6.
- 3 UK National Screening Committee. *Handbook for vascular risk assessment, risk reduction and risk management*. Leicester: University of Leicester, 2008.
- 4 King H, Aubert RE, Herman WH. Global burden of diabetes, 1995-2025: prevalence numerical estimates, and projections. *Diabetes Care* 1998;21:1414-31.
- 5 DECODE study group, European Diabetes Epidemiology Group. Glucose tolerance and mortality: comparison of WHO and American Diabetes Association diagnostic criteria. Diabetes epidemiology: collaborative analysis of diagnostic criteria in Europe. *Lancet* 1999;354:617-21.
- 6 Gillies CL, Abrams KR, Lambert PC, Cooper NJ, Sutton AJ, Hsu RT, et al. Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance. *BMJ* 2007;334:299-302.
- 7 Waugh N, Scotland G, McNamee P, Gillett M, Brennan A, Goyder E, et al. Screening for type 2 diabetes: literature review and economic modelling. *Health Technol Assess* 2007;11:1-125.
- 8 Davies MJ, Tringham JR, Troughton J, Khunti KK. Prevention of type 2 diabetes mellitus. A review of the evidence and its application in a UK setting. *Diabet Med* 2004;21:403-14.
- 9 Avenell A, Broom J, Brown TJ, Poobalan A, Aucott L, Stearns SC, et al. Systematic review of the long-term effects and economic consequences of treatments for obesity and implications for health improvement. *Health Technol Assess* 2004;8:1-182.
- 10 Caro JJ, Getsios D, Caros I, Klittich WS, O'Brien JA. Economic evaluation of therapeutic interventions to prevent type 2 diabetes in Canada. *Diabet Med* 2004;21:1229-36.
- 11 Eddy DM, Schlessinger L, Kahn R. Clinical outcomes and cost-effectiveness of strategies for managing people at high risk for diabetes. *Ann Intern Med* 2005;143:251-64.
- 12 Herman WH, Hoerger TJ, Brandle M, Hicks K, Sorensen S, Zhang P, et al. The cost-effectiveness of lifestyle modification or metformin in preventing type 2 diabetes in adults with impaired glucose tolerance. *Ann Intern Med* 2005;142:323-32.
- 13 Icks A, Rathmann W, Haastert B, Gandjour A, Holle R, John J, et al. Clinical and cost-effectiveness of primary prevention of type 2 diabetes in a "real world" routine healthcare setting: model based on the KORA Survey 2000. *Diabet Med* 2007;24:473-80.
- 14 Jacobs-van der Bruggen MAM, Bos G, Bemelmans WJ, Hoogenveen RT, Vijgen SM, Baan CA. Lifestyle interventions are cost-effective in people with different levels of diabetes risk. *Diabetes Care* 2007;30:128-34.
- 15 Palmer AJ, Roze S, Valentine WJ, Spinaz GA, Shaw JE, Zimmet PZ. Intensive lifestyle changes or metformin in patients with impaired glucose tolerance: modeling the long-term health economic implications of the diabetes prevention program in Australia, France, Germany, Switzerland, and the United Kingdom. *Clin Ther* 2004;26:304-21.
- 16 Segal L, Dalton AC, Richardson J. Cost-effectiveness of the primary prevention of non-insulin dependent diabetes mellitus. *Health Promot Int* 1998;13:197-209.
- 17 CDC Diabetes Cost-Effectiveness Study Group. The cost-effectiveness of screening for type 2 diabetes. *JAMA* 1998;280:1757-63.
- 18 Chen THH, Yen MF, Tung TH. A computer simulated model for cost-effectiveness analysis of mass screening for type 2 diabetes mellitus. *Diabetes Res Clin Pract* 2001;54:S37-42.
- 19 Glumer C, Yuyun M, Griffin S, Farewell D, Spiegelhalter D, Kinmonth AL, et al. What determines the cost-effectiveness of diabetes screening? *Diabetologia* 2006;49:1536-44.
- 20 Hoerger TJ, Harris R, Hicks KA, Donahue K, Sorensen S, Engelgau M. Screening for type 2 diabetes mellitus: a cost-effectiveness analysis. *Ann Intern Med* 2004;140:756-8.
- 21 National Institute for Clinical Excellence. *Guide to the methods of technology appraisal*. London: NICE, 2004.
- 22 Cooper NJ, Sutton AJ, Abrams KR, Turner D, Wailoo A. Comprehensive decision analytical modelling in economic evaluation: a Bayesian approach. *Health Econ* 2004;13:203-26.
- 23 Spiegelhalter DJ, Thomas A, Best NG, Gilks W. *BUGS: Bayesian inference using Gibbs sampling, version 1.4.1*. Cambridge: MRC Biostatistics Unit, 2003.
- 24 Davies MJ, Tringham JR, Jarvis J, Skinner TC, Farooqi AM, Khunti K. Systematic screening for type 2 diabetes mellitus: results of a large population based study targeting those with conventional risk factors. *Diabet Med* 2005;22(suppl 2):99.
- 25 Bonora E, Kiechl S, Willeit J, Oberhollenzer F, Egger G, Meigs JB, et al. Population-based incidence rates and risk factors for type 2 diabetes in white individuals: the Bruneck study. *Diabetes* 2004;53:1782-9.
- 26 Chiasson JL, Josse RG, Gomis R, Hanefeld M, Karasik A, Laakso M. Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial. *Lancet* 2002;359:2072-7.

- 27 De Vegt F, Dekker JM, Jager A, Hienkens E, Kostense PJ, Stehouwer CD, et al. Relation of impaired fasting and postload glucose with incident type 2 diabetes in a Dutch population: the Hoorn study. *JAMA* 2001;285:2109-13.
- 28 Edelstein SL, Knowler WC, Bain RP, Andres R, Barrett-Connor EL, Dowse GK, et al. Predictors of progression from impaired glucose tolerance to NIDDM: an analysis of six prospective studies. *Diabetes* 1997;46:701-10.
- 29 Eriksson J, Lindstrom J, Valle T, Aunola S, Hamalainen H, Ilanne-Parikka P, et al. Prevention of type II diabetes in subjects with impaired glucose tolerance: the diabetes prevention study (DPS) in Finland. Study design and 1-year interim report on the feasibility of the lifestyle intervention programme. *Diabetologia* 1999;42:793-801.
- 30 Eriksson JG, Lehtovirta M, Ehrstrom B, Salmela S, Groop L. Long-term beneficial effects of glipizide treatment on glucose tolerance in subjects with impaired glucose tolerance. *J Intern Med* 2006;259:553-60.
- 31 Jarrett RJ, Keen H, Fuller JH, McCartney M. Worsening to diabetes in men with impaired glucose tolerance ("borderline diabetes"). *Diabetologia* 1979;16:25-30.
- 32 Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393-403.
- 33 Meigs JB, Muller DC, Nathan DM, Blake DR, Andres R, Baltimore Longitudinal Study. The natural history of progression from normal glucose tolerance to type 2 diabetes in the Baltimore longitudinal study of aging. *Diabetes* 2003;52:1475-84.
- 34 Rasmussen SS, Glumer C, Sandbaek A, Lauritzen T, Borch-Johnsen K. Progression from impaired fasting glucose and impaired glucose tolerance to diabetes in a high-risk screening programme in general practice: the ADDITION study, Denmark. *Diabetologia* 2006;50:293-7.
- 35 Wareham NJ, Byrne CD, Williams R, Day NE, Hales CN. Fasting proinsulin concentrations predict the development of type 2 diabetes. *Diabetes Care* 1999;22:262-70.
- 36 Wein P, Beischer N, Harris C, Permezel M. A trial of simple versus intensified dietary modification for prevention of progression to diabetes mellitus in women with impaired glucose tolerance. *Aust NZ J Obstet Gynaecol* 1999;39:162-6.
- 37 Harris MI, Klein R, Welborn TA, Knudman MW, Harris MI, Klein R, et al. Onset of NIDDM occurs at least 4-7 yr before clinical diagnosis. *Diabetes Care* 1992;15:815-9.
- 38 DECODE Study Group, European Diabetes Epidemiology Group. Glucose tolerance and mortality: comparison of WHO and American Diabetes Association diagnostic criteria. Diabetes epidemiology: collaborative analysis of diagnostic criteria in Europe. *Lancet* 1999;354:617-21.
- 39 Rossing P, Hougaard P, Borch-Johnsen K, Parving HH, Rossing P, Hougaard P, et al. Predictors of mortality in insulin dependent diabetes: 10 year observational follow up study. *BMJ* 1996;313:779-84.
- 40 Turner RC, Holman RR, Matthews DR, Oakes SF, Bassett PA, Stratton IM, et al. Study design, progress and performance (UKPDS VIII). *Diabetologia* 1991;34:877-90.
- 41 UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837-53.
- 42 Srinivasan BT, Davies MJ, Webb D, Healey E, Farooqi A, Hiles S, et al. Baseline characteristics and risk of progression from prediabetes to type 2 diabetes in a multiethnic, population based screening. *Diabetic Med* 2007;24(suppl 1):51.
- 43 Clarke P, Gray A, Holman R. Estimating utility values for health states of type 2 diabetic patients using the EQ-5D (UKPDS 62). *Med Decis Making* 2002;4:340-9.
- 44 Clarke PM, Gray AM, Briggs A, Farmer AJ, Fenn P, Stevens RJ, et al. A model to estimate the lifetime health outcomes of patients with type 2 diabetes: the United Kingdom prospective diabetes study (UKPDS) outcomes model (UKPDS 68). *Diabetologia* 2004;47:1747-59.
- 45 Curtis L, Netten A. *Unit costs of health and social care*. Canterbury: Personal Social Services Research Unit, University of Kent, 2006.
- 46 Gulliford MC, Charlton J, Latinovic R. Increased utilization of primary care 5 years before diagnosis of type 2 diabetes. *Diabetes Care* 2005;28:47-52.
- 47 Clarke PM, Gray AM, Briggs A, Stevens RJ, Matthews DR, Holman RR. Cost-utility analyses of intensive blood glucose and tight blood pressure control in type 2 diabetes. *Diabetologia* 2004;48:868-77.
- 48 Ramachandran A, Snehalatha C, Mary S, Mukesh B, Bhaskar AD, Vijay V. The Indian diabetes prevention programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia* 2006;49:289-97.

Accepted: 25 March 2008