

Patient information from BMJ

Last published: Jul 10, 2020

How can I tell when my child's asthma is getting out of control?

You should see your doctor if any of the following is true:

- Your child is waking up at night more than usual with asthma symptoms
- The inhaler you use to treat your child's symptoms quickly (the reliever) doesn't help their breathing for very long, or doesn't help at all, or you are using it more often
- Your child can't breathe out as much air as usual, as indicated by their PEFR meter (PEFR is short for peak expiratory flow rate). Doctors sometimes recommend that older children use these meters at home to regularly check their breathing.

You should seek emergency treatment if:

- It's hard for your child to talk because of asthma
- Your child cannot eat or drink because of asthma
- Your child is breathing very quickly
- Your child's neck muscles are straining and moving quickly
- Your child's heartbeat or pulse is very fast
- Your child finds it hard to walk
- Your child's lips or fingernails turn grey or blue
- Your child is exhausted for no apparent reason
- Your child seems agitated or confused and is losing consciousness (blacking out).

Your child's doctor, or a practice nurse with special training in asthma, should regularly check that your child's asthma is under control. They may ask you these types of questions about your child's symptoms:

- Has your child had difficulty sleeping because of their asthma symptoms (including coughing)?
- Has your child had any symptoms during the day (coughing, wheezing, chest tightness, or breathlessness)?

How can I tell when my child's asthma is getting out of control?

- Has your child's asthma stopped any activities (such as going to school)?

Depending on your answers, they may decide to change your child's treatment.

The patient information from *BMJ Best Practice* is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2024. All rights reserved.

What did you think about this patient information guide?

Complete the [online survey](#) or scan the QR code to help us to ensure our content is of the highest quality and relevant for patients. The survey is anonymous and will take around 5 minutes to complete.



BMJ Group