

Patient information from BMJ

Last published: Sep 27, 2021

Lyme disease

Lyme disease is an infection that you can get from the bite of a tiny creature called a tick. It can be serious, but most people recover completely with treatment.

You can use our information to talk to your doctor and decide which treatments are best for you.

What is Lyme disease?

Lyme disease is an illness that's spread by ticks. Ticks are tiny creatures that live on the skin of other animals and feed on their blood. But they also sometimes bite and feed on people. They look a bit like tiny spiders.

Ticks can carry the bacteria that cause Lyme disease and, if they bite you, the bacteria can get into your body and make you sick.

Not all ticks carry the Lyme disease bacteria (called *Borrelia*). And a tick usually has to stay attached to your skin for a long time (at least one or two days) for there to be much risk of it making you sick.

If you spot a tick on your skin you need to remove it. Grip the tick with fine-tipped tweezers as close to your skin as you can. Pull the tick gently but firmly off your skin, then wash the area with soap and water.

What are the symptoms?

The most common symptom of Lyme disease is a pink or red rash. It will probably appear one to two weeks after you were bitten, spreading out from the bite. Some people get a bull's-eye pattern, with no red in the center.

But not everyone gets a rash. You may get flu-like symptoms, such as:

- a headache
- a fever, or
- muscle pains.

Lyme disease

An untreated infection can, after a few weeks or months, spread to your nerves and joints. You may lose control of some of the muscles in your face. This is called **neurological Lyme disease**.

If this happens you may get pain or numbness, or have difficulties moving parts of your body.

If you're not treated you could also get swollen, painful joints. This is called **Lyme arthritis**.

A few people with Lyme disease get **meningitis**. Meningitis means that the tissue covering your brain has become inflamed.

Lyme disease can also sometimes cause **heart problems**. These can be treated but you may need to have treatment in the hospital.

Doctors often diagnose Lyme disease based on people's symptoms and on whether they may have been exposed to ticks carrying Lyme disease. A blood test is also sometimes used. But it isn't effective until two to six weeks after a tick bite, so it isn't helpful for diagnosing early infections.

What treatments are available?

Preventing tick bites

If you're spending time in places where ticks are common, such as grassy or wooded areas, you can:

- wear long-sleeved tops and long pants tucked into your socks
- use an insect repellent called DEET on any exposed skin and your clothes
- spray your clothes, and any tents and camping gear, with a repellent called permethrin
- wear light-colored clothing to make it easier to find and remove ticks.

If you've been in an area where there may be ticks, brush off or wash your clothes as soon as you can. Check your body for ticks, especially in skin folds and hairy areas.

Young ticks are about the size of a poppy seed, so they're hard to spot. Adult ticks are bigger (about the size of a sesame seed) and may reach the size of a coffee bean if they are full of blood. If you find a tick, carefully remove it right away.

Treating Lyme disease

If you have Lyme disease you'll be treated with **antibiotics**. Antibiotics are drugs that kill bacteria. If you take them soon after getting bitten you're very likely to make a complete recovery.

Antibiotics may also help to prevent Lyme disease. If you are in an area with Lyme disease and you've had a tick attached to your skin for at least 36 hours, your doctor may prescribe an antibiotic. But you would need to have treatment within three days of removing the tick.

Antibiotics are also used to treat more serious symptoms of Lyme disease, such as arthritis, and nerve and heart problems. You may need to be given antibiotics as a drip (an intravenous infusion, or IV for short).

Lyme disease

To relieve joint pain your doctor may recommend taking a **nonsteroidal anti-inflammatory drug** (NSAID), such as diclofenac or ibuprofen.

Lyme disease in pregnancy

It is possible, but rare, for pregnant women with Lyme disease to pass it onto their babies.

There is some evidence that untreated Lyme disease can cause problems during pregnancy, as well as birth defects. But women who get treatment should not have problems. If you are concerned about Lyme disease in pregnancy, talk to your doctor.

What to expect in the future?

If Lyme disease is treated early you're very likely to recover completely. If you get a rash or flu-like illness and think you could have been bitten by a tick (even if you didn't notice a bite), see your doctor as soon as you can.

Getting treatment early helps prevent later-stage Lyme disease, which can be more serious. If you don't have treatment, your early symptoms (such as the rash) may clear up. But you have a strong chance of then getting later-stage problems, such as joint problems and nerve damage.

Some people do have lasting pain and tiredness even after they've been treated for Lyme disease. This is sometimes called "post-Lyme syndrome." People with this syndrome usually get better with time. But it can take months to feel completely better.

The patient information from *BMJ Best Practice* is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2024. All rights reserved.

What did you think about this patient information guide?

Complete the [online survey](#) or scan the QR code to help us to ensure our content is of the highest quality and relevant for patients. The survey is anonymous and will take around 5 minutes to complete.

