


Neonatal frequent premature ventricular contraction

Rie Chida,¹ Kazuki Iio ,^{1,2} Yu Ishida,¹ Gaku Yamanaka³

¹Department of Pediatrics, Tokyo Medical University Hachioji Medical Center, Hachioji, Tokyo, Japan

²Division of Pediatric Emergency Medicine, Tokyo Metropolitan Children's Medical Center, Fuchu, Tokyo, Japan

³Department of Pediatrics and Adolescent Medicine, Tokyo Medical University, Shinjuku-ku, Tokyo, Japan

Correspondence to

Dr Kazuki Iio;
kazukiio1026@gmail.com

Accepted 21 February 2023

DESCRIPTION

A female baby was born at term by emergency caesarean section, due to non-reassuring fetal status on the cardiotocography and chorioamnionitis. Her Apgar score was 8 (1 min)/9 (5 min), and no heart murmur or irregular beats were heard on physical examination soon after the birth. One hour after the birth, the nurse noticed that the patient's heartbeats were irregular during the routine examination.

The patient appeared normal without any abnormal physical findings other than irregular heartbeats. Her vital signs were all normal. Electronic cardiac monitoring revealed frequent monomorphic premature ventricular contractions (PVCs), frequently converted to bigeminy or trigeminy ([figure 1](#)). The patient's blood test and echocardiography results were normal. The 12-lead ECG detected frequent monomorphic PVCs ([figure 2](#)). Holter monitoring for 24 hours revealed no lethal ventricular tachycardias. PVCs accounted for 9.3% (675 beats/hour) of the entire heartbeats. In the second 24-hour continuous Holter monitoring performed on day 5, her PVC had decreased significantly to 1.9% (155 beats/hour). She was asymptomatic without any treatment and discharged from the neonatal intensive care unit on day 7 with her mother. We repeated 24-hour continuous Holter monitoring 1 month after birth, which revealed no PVCs.

Isolated PVCs are a common phenomenon observed in about 20% of healthy neonates.¹ However, frequent PVCs are quite rare in neonates. Because of the intrinsically higher heart rates in neonates, the threshold for 'frequent' PVCs in neonates has been set at more than 60 beats/hour, compared with more




Figure 2 The 12-lead ECG detected frequent monomorphic premature ventricular contractions.

than 30 beats/hour in adults.¹ Exclusion of underlying conditions is important in evaluating frequent neonatal PVCs, including congenital heart disease, inflammatory myocardial disease, metabolic disease, electrolyte disorders or long QT syndrome.² If neonates are otherwise healthy, a simple blood test with ECG and echocardiography would be sufficient for this purpose, as metabolic disorders presenting with cardiac symptoms in neonates usually present with heart failure rather than isolated arrhythmia.³

Even in the absence of underlying disease, frequent PVCs may rarely develop in otherwise healthy neonates.¹ In these cases, PVCs usually resolve within a few months of birth,¹ which is much earlier than in adolescent patients.⁴ This rapid resolution may be explained by the pathogenesis of frequent neonatal PVCs, which is thought to be due to immature cardiac conduction tissue and autonomic nervous system.⁵




Figure 1 Electronic cardiac monitoring revealed frequent bigeminy or trigeminy.



© BMJ Publishing Group Limited 2023. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Chida R, Iio K, Ishida Y, et al. *BMJ Case Rep* 2023;**16**:e254837. doi:10.1136/bcr-2023-254837

Learning points

- ▶ Neonatal frequent premature ventricular contractions (PVCs) are defined as PVCs over 60 beats/hour. The underlying cardiac or systemic illnesses should be evaluated.
- ▶ If there were no underlying illnesses, neonatal frequent PVCs would follow a self-limiting course, which usually resolves spontaneously within a few months of birth.

Antiarrhythmic medication would not be needed unless other types of arrhythmia developed during follow-up.

Contributors RC and KI drafted the initial manuscript, and reviewed and revised the manuscript. YI and GY critically reviewed and revised the manuscript. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

Kazuki Iio <http://orcid.org/0000-0002-6430-4957>

REFERENCES

- 1 De Rosa G, Butera G, Chessa M, *et al.* Outcome of newborns with asymptomatic monomorphic ventricular arrhythmia. *Archives of Disease in Childhood - Fetal and Neonatal Edition* 2006;91:F419–22.
- 2 Ban JE. Neonatal arrhythmias: diagnosis, treatment, and clinical outcome. *Korean J Pediatr* 2017;60:344.
- 3 Saudubray JM, Garcia-Cazorla A. Inborn errors of metabolism overview: pathophysiology, manifestations evaluation, and management. *Pediatr Clin North Am* 2018;65:179–208.
- 4 Tsuji A, Nagashima M, Hasegawa S, *et al.* Long-term follow-up of idiopathic ventricular arrhythmias in otherwise normal children. *Jpn Circ J* 1995;59:654–62.
- 5 Jaeggi E, Öhman A. Fetal and neonatal arrhythmias. *Clin Perinatol* 2016;43:99–112.

Copyright 2023 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow