

# Live 'Baby dance' among the red blood cells

Praveen Sharma,<sup>1</sup> Pulkit Rastogi,<sup>1</sup> Vikarn Vishwajeet,<sup>2</sup> Man Updesh Singh Sachdeva<sup>1</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bcr-2017-222321>).

<sup>1</sup>Department of Hematology, Post Graduate Institute of Medical Education and Research, Chandigarh, Punjab, India

<sup>2</sup>Department of Histopathology, Postgraduate Institute of Medical Education and Research, Chandigarh, Punjab, India

## Correspondence to

Dr Man Updesh Singh Sachdeva, [drmanupdeshpgi@yahoo.co.in](mailto:drmanupdeshpgi@yahoo.co.in)

Accepted 29 September 2017

## DESCRIPTION

A 35-year-old woman presented to outpatients with a 1-week history of high-grade fever and generalised malaise. General and systemic examination was unremarkable. On evaluation, her complete haemogram was within normal limits; however peripheral smear revealed the presence of thread-like larval forms of filaria, belonging to the species *Wuchereria bancrofti*. A wet mount preparation of the centrifuged whole blood sample was done, which demonstrated 'dancing' microfilaria, recorded in real time (see online supplementary video 1).

Dancing adult worms can be visualised within the dilated lymphatic channels on high-resolution ultrasound by their characteristic wriggling type movements, known as the 'filarial dance sign'.<sup>1</sup> A wet mount of blood from a patient with microfilaraemia can demonstrate 'dance sign' of the baby worms under a microscope, identical to the parental wriggling movements. Microfilariae are larval forms, released into the blood from adult female worms inhabiting the regional lymphatic vessels. Circulating microfilariae are taken up by vector mosquitoes, only to infect another individual and develop as adult worms in a subsequent

host. Although wet mount preparations are used routinely for the diagnosis of intestinal parasitic infections, such preparations can also be used to see circulating microfilaria, which can be seen up to several months of acquiring infection and serve as a source of infection to others. Various methods like direct wet mount preparation, stained blood smear examination, citrate saponin acid method and modified Knott's method can be used to detect microfilariasis.<sup>2</sup> Studies have shown that the efficacy of wet film examination is up to 65% and it is considered superior for microfilaria detection compared to Giemsa stained blood smears. Modified Knott's method demonstrates greater efficacy than the above mentioned methods.<sup>2</sup> A centrifuged wet mount preparation might increase the yield of microfilariae in patients with low parasitaemia, however, in the index case it was done to appreciate the characteristic 'microfilarial dance sign'.

**Contributors** PS drafted the manuscript; PR prepared the wet mount and recorded the video; VV picked up the diagnosis on blood smear; MUSS edited the manuscript.

**Competing interests** None declared.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

## REFERENCES

- 1 Chaubal NG, Pradhan GM, Chaubal JN, et al. Dance of live adult filarial worms is a reliable sign of scrotal filarial infection. *J Ultrasound Med* 2003;22:765–9.
- 2 Sundar B, Ravindran R. Comparison of various methods for the detection of microfilaria of setaria in the blood of cattle. *TN/VAS* 2000;6:45–8.

## Learning points

- Wet mount smears of the centrifuged blood sample can also be helpful in diagnosis of filariasis.
- 'Microfilarial dance sign' is the characteristic motility of the larval form and is observed only in wet mount smears.



CrossMark

**To cite:** Sharma P, Rastogi P, Vishwajeet V, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-222321

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact [consortiasales@bmjgroup.com](mailto:consortiasales@bmjgroup.com)

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow