

Evidence-Based Medicine

for Primary Care and Internal Medicine

Editors: Brian Haynes & Paul Glasziou



www.evidence-basedmedicine.com

Contents

Purpose and procedure	33	high risk patients after the programme was stopped	50
EBM notebook		Community based occupational therapy improved daily functioning in older patients with dementia	51
Jottings...	34	Review: Inhaled insulin provided better glycaemic control than oral hypoglycaemic agents but not subcutaneous insulin	52
Editor's choices	34	Review: pioglitazone does not reduce risk of mortality or cardiovascular events in type 2 diabetes	53
Evidently	35		
Teaching tip: using the "Who wants to be a millionaire?" game to teach searching skills	36	Diagnosis	
Physiotherapy for tennis elbow	37	Clinical examination had poor sensitivity for detecting non-cephalic presentation in late pregnancy	54
Methods in research: key principles of graph construction	38	Review: alarm features in dyspepsia do not have high diagnostic accuracy for predicting upper gastrointestinal cancer	55
		Review: Sensitivities and specificities vary for different diagnostic tests of gout	56
Therapeutics		Clinical prediction guide	
Corticosteroid injections or physiotherapy were not more effective than wait and see for tennis elbow at 1 year	39	A risk model predicted major bleeding in older patients with atrial fibrillation receiving warfarin therapy	57
Review: evidence from 2 low quality screening studies does not show a reduction in death from prostate cancer	40		
Vaccinating infants with a wide long needle was noninferior to a narrow short needle for immune response	41	Economics	
Review: high dose statins reduce risk of non-fatal cardiovascular events more than standard dose statins	42	Simvastatin was cost effective across a broad range of risk and age groups	58
Atorvastatin at 80 mg/day reduced cerebrovascular events more than atorvastatin at 10 mg/day in stable coronary heart disease	43		
Physical counterpressure manoeuvres reduced vasovagal syncope	44	Resource review	59
An abnormal D-dimer test result indicated that anticoagulant therapy should be continued	45		
Use of a 2 hour partogram action line instead of a 4 hour action line did not reduce caesarean delivery rate	46	Letter	61
Review: children <2 years of age with bilateral acute otitis media and children with otorrhoea benefit most from antibiotics	47		
The Air-Stirrup brace with elastic wrap improved function in first incident grade 1 or 2 lateral ankle ligament sprain	48	Additional articles abstracted in ACP Journal Club	62
Addition of peak flow monitoring to symptom monitoring did not improve healthcare visits, quality of life, or lung function in older adults with moderate-to-severe asthma	49		
A lifestyle intervention continued to prevent type 2 diabetes in		Other articles noted	63
		Glossary	Inside back cover