Evidence-Based Medicine



Contents

Editorial

151 Treating elevated LDL cholesterol in patients with low short-term risk: Decision making at the limits of EBM

Perspective

154 Superiority trials: raising the bar of null hypothesis statistical testing

Original EBM Research

156 How does evidence affect clinical decision-making?

Systematic review

A review of the clinical evidence related to early treatment of elevated LDL for cardiovascular primary prevention

Therapeutics/Prevention

- 170 Restrictive red blood cell transfusion strategies appear safe in most clinical settings
- 171 α-channel and calcium-channel blockers are ineffective as medical expulsive therapy for ureteral stones regardless of size and location
- 172 Personalised chronic care planning shows limited but promising effects on physical and psychological health and self-management capabilities
- 173 In knee osteoarthritis, pharmacological interventions, with the exception of acetaminophen, significantly improve pain; with intra-articular administration being more effective
- 174 An IVC filter and anticoagulation for 3 months is unlikely to show a benefit over anticoagulation alone for high-risk patients with acute pulmonary embolism
- 175 In patients with persistent atrial fibrillation complex ablation techniques were no better than pulmonary vein isolation alone in preventing recurrent atrial fibrillation
- 176 Ezetimibe provides incremental reduction in risk for cardiovascular events and need for revascularisation following an acute coronary syndrome
- 177 A supplementation of DHA and AA to human milk-fed VLBW infants has no significant cognitive improvement or measurable neuroanatomical effects when evaluated at 8 years of age

October 2015 Volume 20 No 5

- 178 Antivirals in combination with corticosteroids compared to corticosteroids alone improve facial recovery in severe Bell's palsy... and possibly more?
- 179 The RTS,S/AS01 vaccine continues to show modest protection against malaria in African infants and children
- 180 Directly observed treatment is not the only solution for poor adherence and low cure of tuberculosis
- 181 Percutaneous vertebroplasty does not reduce pain and disability from osteoporotic vertebral compression fracture
- 182 Short-course antimicrobial therapy may be clinically similar to a longer course for complicated intraabdominal infections
- 183 Blood pressure lowering in patients with type 2 diabetes improves cardiovascular events including mortality, but more intensive lowering to systolic blood pressure less than 130 mm Hg is associated with further reduction in stroke and albuminuria without further reduction in cardiac events
- 184 Immediate delivery in women with non-severe hypertensive disorders at 34–37 weeks' gestation does not reduce maternal complications, and increases neonatal risks more than under expectant management
- 185 Perioperative use of chewing gum affects the inflammatory response and reduces postoperative ileus following major colorectal surgery

Diagnosis

187 CT coronary angiography increases diagnostic certainty in patients with stable chest pain

Aetiology/Harm

- **188** NSAIDs increase bleeding risk in patients with atrial fibrillation on antithrombotic therapy
- 189 Combined oral contraceptives and risk of venous thromboembolism: there is higher risk in new generations compared to second generations, but paradoxically not in norgestimate-containing-pills

Letter

190 Informed dissent: can we handle internet medicine?

EDITOR'S CHOICE This article has been chosen by the Editor to be of special interest or importance and is freely available online



This article has been made freely available online under the BMJ Journals Open Access scheme. See http://ebm.bmj.com/site/about/guidelines.xhtml#open



This journal is a member of and subscribes to the principles of the Committee on Publication

Ethics



When you have finished with this please recycle

Receive regular table of contents by email.

Register using this QR code.

