

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Protocol for a feasibility study of OnTrack: a digital system for upper-limb rehabilitation after stroke.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034936
Article Type:	Protocol
Date Submitted by the Author:	16-Oct-2019
Complete List of Authors:	Fusari, Gianpaolo; Royal College of Art, Helix Centre; Imperial College London, Helix Centre Gibbs, Ella; Imperial College London, Helix Centre Hoskin, Lily; Imperial College London, Helix Centre Dickens, Daniel; Imperial College London, Helix Centre Leis, Melanie; Imperial College London, Big Data and Analytical Unit, Institute of Global Health Innovation Taylor, Elizabeth; Kingston and St George's Faculty of Health Social Care and Education Jones, Fiona; Kingston and St George's Faculty of Health Social Care and Education Darzi, Ara; Imperial College London, Helix Centre
Keywords:	Stroke < NEUROLOGY, REHABILITATION MEDICINE, NEUROLOGY, PUBLIC HEALTH

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Title

Protocol for a feasibility study of OnTrack: a digital system for upper-limb rehabilitation after stroke.

Authors and affiliations

Gianpaolo Fusari¹, Ella Gibbs¹, Lily Hoskin¹, Daniel Dickens¹, Melanie Leis², Elizabeth Taylor³, Fiona Jones³, Ara Darzi¹

¹Helix Centre, Imperial College London and the Royal College of Art, London, United Kingdom

²Big Data and Analytical Unit, Institute of Global Health Innovation, Imperial College London, London, United Kingdom

³Kingston University and St George's, University of London, London, United Kingdom

Correspondence to:

Gianpaolo Fusari

HELIX Centre, Institute of Global Health Innovation

Imperial College London, 3rd Floor, Paterson Building, St Mary's Hospital, 16 South Wharf Road, London W2 1PF, UK

gianpaolo@helixcentre.com

+44 (0)77 6779 2770

Key words

Stroke, rehabilitation medicine, neurology, public health, digital health

Word count

3806

Abstract

Introduction

Arm weakness is a common problem after stroke (affecting 450,000 people in the UK) leading to loss of independence. Repetitive activity is critical for recovery but research shows people struggle with knowing what or how much to do, and keeping track of progress. Working with >100 therapists and patients, we co-developed the OnTrack intervention - consisting of software for smart-devices and coaching support - that has the potential to address this problem. This is a protocol to assess the feasibility of OnTrack for evaluation in a randomised control trial.

Methods and analysis

A mixed methods, single-arm study design will be used to evaluate the feasibility of OnTrack for hospital and community use. Participants from a stroke unit will be recruited (n=24) into the study and will be involved for 14 weeks. During week 1, 8 and 14 participants will complete assessments relating to their arm function, arm impairment, and activation. During weeks 2-13 participants will use OnTrack to track their arm movement in real time, receive motivational messages, and face-to-face sessions to address problems, gain feedback on activity, and receive self-management skills coaching. All equipment will be loaned to study participants. A parallel process evaluation will be conducted to assess the intervention's fidelity, dose and reach, using a mixed methods approach. A Public and Patient Involvement (PPI) group will oversee the study and help with interpretation of qualitative and quantitative data findings.

Ethics and dissemination

Ethical approval granted by the NHS Health Research Authority, Health and Care Research Wales, and the London - Surrey Research Ethics Committee (ref.19/LO/0881). Trial results will be submitted for publication in peer review journals, presented at international conferences and disseminated amongst stroke communities. The results of this trial will inform development of a definitive trial.

Trial registration details

ClinicalTrials.gov (NCT03944486), pre-results.

Strengths and limitations

- This is a feasibility trial of a novel intervention which employs an integrated approach for tracking arm activity and coaching with the aim of increasing stroke survivors' confidence and ability to use their impaired arm in daily activities, increasing the opportunities for repetitive rehabilitation.
- PPI involvement from more than 100 stroke survivors, carers, and clinicians have contributed to our needs-finding phase, co-designed OnTrack and informed the feasibility study. A new PPI group will oversee the running of the study and help with interpretation of qualitative and quantitative data findings.
- An independent process evaluation will provide detailed information about implementation, context, and the mechanisms of impact of the intervention. Findings will help in the understanding of intervention fidelity and training needs required for a definitive trial.
- For pragmatic reasons the study uses a non-randomised design carried out at a single site- this will limit understanding about randomisation and recruitment
- Participants will not be followed-up after intervention; however participant views will be sought regarding appropriate follow-up times in a subsequent definitive trial.

Introduction

Every year around the world over 15 million people experience a stroke, leaving 5 million people with a permanent disability.(1) Stroke is the leading cause of disability in the UK; half of the nearly 1.2 million stroke survivors who live in the country have some form of disability, significantly contributing to the loss of independence and feeling of isolation that they experience. (2,3) Furthermore, stroke is estimated to cost UK society £26 billion every year, with the vast majority of these costs borne by the informal care sector.(2)

Upper-limb (arm) weakness is the main cause of physical impairment affecting 75% of disabled stroke survivors; this equates to around 450,000 people in the UK.(2) Dose-intensive repetitive rehabilitation is widely accepted as the 'gold-standard' for regaining ability after stroke, however, NHS resources are often limited and unable to provide this.(4,5) A recent Cochrane review of over 500 trials failed to yield high-quality practice recommendations.(6) Arm recovery after stroke is a national research priority,(7) nonetheless, studies suggest that the actual time patients spend exercising is minimal.(8,9) Many current approaches to solving this problem focus on improving the prescribed rehabilitation sessions, often employing gamification techniques.(10,11) Whilst this is important, there is untapped potential to increase repetitive rehabilitation by targeting the large proportion of the day where patients are going about their daily activities and can use their arm movement (however small) to a greater extent. Capacity for activity could be increased further by using self-management methods as demonstrated by several different programmes in stroke and other long-term conditions.(12-15) This has informed the

development of OnTrack which aims to increase opportunities for activity by improving individuals' self-management skills through tailored support and real-time activity feedback on their arm movement.

An ethnographic study conducted by the Helix Centre(16) (funded by Innovate UK) revealed that patients struggle to see and keep track of improvements, this impacts their motivation and leaves them dependent on therapists for feedback. Stroke survivors often report feeling unsupported after leaving hospital and not knowing how to best help themselves improve their arm function, confirming views documented by other studies.(17-19) Feedback gathered from over 100 stroke survivors and clinicians was the basis for developing the OnTrack intervention.

A proof-of-concept test of OnTrack gathered data from a small group of patients (n=7) and confirmed that the intervention was safe and generally users could understand how and when to use it. Participants reported they were more aware of their impaired arm and had increased confidence in using it for new tasks. A 20% mean increase in activity was observed. The work conducted to date suggests that OnTrack has the potential to be a scalable solution that requires minimal training and could be used in conjunction with NHS services to help increase the overall amount of arm rehabilitation received. This study will assess the feasibility of the OnTrack intervention and inform the design of a definitive randomised controlled trial (RCT) to evaluate its clinical effectiveness, and follows the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) guidelines.(20)

Methods and analysis

Aims and objectives

The primary aim is to evaluate the feasibility of an RCT to test the effectiveness of the OnTrack intervention for upper limb rehabilitation after stroke.

The objectives are to:

- Assess the feasibility of recruitment from hyper-acute and acute stroke units, and rehabilitation wards to ascertain strategy and recruitment rates.
- Assess dropout rates by observing adherence and compliance with the intervention.
- Understand the acceptability and usability of the intervention by stroke survivors.
- Understand the acceptability of study procedures by healthcare professionals.
- Explore implementation fidelity, dose and reach of the OnTrack intervention.

The study will also collect clinical outcomes regarding arm function, impairment and activation to identify an appropriate primary outcome, and to estimate parameters for a sample size calculation for an RCT ([Table 1](#)).

Study design

A feasibility study with a nested process evaluation ([Figure 1](#)). The study is a single-site, non-randomised intervention trial. The design of the study was developed through a collaborative approach between the study researchers, a PPI steering group, front-line therapists, and the Research Design Service at the National Institute for Health Research.

An independent process evaluation will be conducted in parallel to learn about usage and engagement mechanisms of participants, therapists and other frontline staff, providing critical information for implementation fidelity and impact mechanisms necessary for scale-up.

Study setting

The study will be conducted at an inner city NHS hospital Trust in London. Recruited participants will be able to continue to receive the intervention at home if discharged from hospital prior to ending the intervention period (14 weeks).

Participants

The inclusion criteria encompasses:

- Adults (aged 18 or over).
- Stroke diagnosis less than 6 months previously (first or recurrent).
- Arm impairment of any type or level (including weakness - including dense hemiplegia, neglect, and sensory deficits).
- Ability to provide informed consent.
- Reliability to communicate (verbally or nonverbally) and understand English.
- Ability to read a predefined short message.

Potential participants who at the time of recruitment present with any of the following will be excluded:

- Unstable medical condition.
- Severe pain in the arm affected either at rest or during movement.
- Severe oedema in the arm affected by their stroke.
- Known discharge plans to a hospital other than the site Trust or residential care in less than 7 weeks.

Recruitment

Participants will be recruited from the Hyperacute Stroke Unit (HASU), Acute Stroke Unit (ASU), and Clinical Neurorehabilitation Unit (CNRU) at an inner city NHS hospital Trust in London.

Stroke clinicians will be responsible for screening and identifying suitable patients. They will introduce the study to potential participants and provide information documents. Potential

participants will be given a minimum of 24 hours to consider the advantages and disadvantages of participating in the study and to formulate questions. Therapists will be able to answer questions or will liaise with the research team to provide an answer. Once all questions are answered and a potential participant is willing to participate, consent will be taken by the therapist. Only at this stage will patient information be shared with the research team. There may be situations where a therapist is only able to take verbal consent from a participant due to time or material constraints, in such cases the researchers will be able to take written consent from the participant upon first meeting them.

Sample size calculation

Guidelines advocate a sample size of 12-30 participants for feasibility studies.(21) Experienced clinical academics and clinicians at the trial site have advised to expect about 50% of eligible patients to agree to participation and a 50% completion rate. This has informed a recruitment plan to identify at least 60 potential participants in a period of 30 weeks to reach the minimum sample size.

Intervention

The OnTrack system consists of smart-devices (smartphone and smartwatch), software (OnTrack app), and coaching support. Smart-devices are used to track arm movement. Motivational messages and a real-time display of completed arm activity are presented to the user via the OnTrack app. Coaching support is provided through fortnightly consultations by the researchers. During consultations, participants will receive self-management training informed by the Bridges(22) and TaCAS(23) self-management programs. Data gathered by the OnTrack system can be accessed by the researchers via a digital dashboard to inform consultations.

Participants will be loaned all equipment necessary for the trial and no previous experience with using smart-devices is required to participate. Technical support will be provided only in cases where the hardware and/or software fail to perform the required functions to deliver the intervention.

[Table 2](#) provides a participation schedule and a summary of the intervention procedures.

Outcomes

Feasibility of trial design and procedures

- Recruitment strategy and rates (feasibility of recruitment from HASU, ASU, CNRU wards) - percentage of patients: screened; eligible; approached; consented; excluded after screening. Participants consented and recruited will be logged in DOCUMAS(24)
- Compliance and adherence to intervention - percentage of participants who start OnTrack daily for the duration of the intervention period, measure of minutes of activity

per participant as recorded by the OnTrack app, engagement with OnTrack app as measured by system analytics.

- Completion rates - percentage of participants who complete the intervention
- Acceptability and reasons for decline/withdrawal - number of participants who withdraw or decline the intervention and reasons why

Clinical assessments

As a secondary objective, clinical outcomes will be collected at different time points to identify an appropriate primary outcome, and to estimate parameters for a sample size calculation for an RCT ([Table 1](#)). The outcome measures and assessments are listed below.

Patient activation

Patient activation is a concept recognised by the NHS that describes the knowledge, skills and confidence a person has in managing their own health and health care.(25) This will be measured using the Patient Activation Measure (PAM)(26) which has been validated in stroke populations in the UK.(27) The PAM survey measures patients on a 0–100 scale and can categorise patients into one of four activation levels along an empirically derived continuum.(26) Activation levels will be used to allocate participants one of three different OnTrack coaching tiers.

Arm impairment

Arm impairment will be measured objectively using the Fugl-Meyer Assessment for upper extremity (FMA-UE).(28) The FMA-UE has been tested extensively, and is found to have excellent psychometric properties and is recommended as core measures to be used in every stroke recovery and rehabilitation trial.(29)

Arm function

Arm function will be assessed using the Upper-Extremity Motor Activity Log-14 (MAL).(30) The MAL is a scripted, structured interview developed to self-report the amount and quality of use of the impaired arm in individuals with stroke in 14 different activities of daily living.

Gross level of disability

The modified Rankin Scale (mRS)(31) is the most prevalent functional outcome measure in contemporary stroke trials. The mRS quantifies disability using an ordinal hierarchical grading from zero (no symptoms) to 5 (severe disability).

Arm pain

Pain will be assessed using a visual analogue scale (VAS) from 0 (no pain) to 10 (excruciating pain) over the last 24 hours. VAS is a valid measure of pain intensity and is responsive to change.(32) Individuals scoring 3/10 or more in the affected arm will be excluded/withdrawn from the study unless their pain is only on movements that are not part of their usual everyday activities (e.g. arm pain when doing overhead reaching).

Cognitive impairment

Cognitive impairment will be assessed using the Montreal Cognitive Assessment (MoCA). The MoCA is a brief cognitive screening tool with high sensitivity and specificity for detecting mild cognitive impairment.(33)

Perceptual neglect

Albert's Test (AT) is a simple test where participants are asked to cross out lines ruled in a standard fashion on a sheet of paper. The test is very easy to administer and is a good predictor of functional activity six months after stroke onset.(34)

Quality of life

The EQ-5D-5L is a widely used standardised preference based measure of health status developed by the EuroQol Group in order to provide a simple, generic measure of health for clinical and economic appraisal.(35)

Additional assessments

A Lap-to-Table (LTT) timed test will be performed where the researchers measure the time it takes a participant to move their hand three times from resting on their lap to a table positioned in front of them. This test is performed to assess its potential to use as part of the inclusion criteria for an RCT.

The NHS Friends and Family Test (FFT)(36) will be used to obtain feedback on the overall experience of using OnTrack and participating in the trial. Participants will be asked: "How likely are you to recommend OnTrack to friends and family if they needed similar care or treatment?" with answers provided in a Likert 5-point scale ranging from "extremely likely" to "extremely unlikely" and an "I don't know" option.

The System Usability Scale (SUS) will be used to subjectively assess the usability of the OnTrack intervention. The test is a simple, ten-item scale covering a variety of aspects of system usability, such as the need for support, training, and complexity, and thus have a high level of face validity for measuring the usability of a system.(37)

Process evaluation

A process evaluation will be carried out by researchers working independently to the intervention team and in parallel to the trial to determine whether the OnTrack intervention was delivered as intended and to understand the mechanisms of impact. The aim of the process evaluation at the feasibility stage is mainly to understand how the trial design and intervention could be optimised ahead of an RCT.(38) A logic model(39,40) that defines the intervention in terms of inputs, outputs, causal assumptions and expected outcomes has been developed to help identify core questions for the evaluation team to explore (Figure 2). The evaluation team will observe a percentage of all intervention sessions with the objective of documenting fidelity, dose and reach of the intervention.

Interim results will be shared with the intervention team at the half-way point with the objective to review some of the procedures and make minor adjustments as necessary.

In-depth semi-structured interviews will be conducted with patients at the end of their participation, a minimum sample of 12 is anticipated. A topic guide with themes drawing from the logic model will be used. Interviews will focus on participants' experiences using OnTrack, their perceptions of arm tracking, motivational messaging and the researcher consultations. Additionally, the interviews will explore participants' perceptions of the impact OnTrack had on them in terms of progress, awareness, participation, and confidence in self-management. Participants' responses will be compared against activity data collected from the OnTrack app.

NHS therapists caring for participants taking part will be invited to complete a short online survey to gather their feedback regarding acceptability of study procedures.

Data analysis

Analysis will be completed on the parameters and implementation of the study in addition to the usability of OnTrack.

Data collected for the process evaluation will capture changes over time and will be a combination of qualitative data from interviews with stroke participants and therapists to explore their experiences of using OnTrack, as well as quantitative data on usage of OnTrack and the self-reported SUS. OnTrack therapy support sessions will be monitored through a fidelity checklist and observations. Interview data will undergo thematic analysis by the evaluation team. Data will be entered into NVIVO, (41) line by line coding and analysis will be informed by Braun and Clark's approach to thematic analysis.(42)

The team will analyse users' activity patterns by day and hour of day. This will allow an understanding of how usage varies for each user over time, as well as how patterns of usage vary from one user to another. [Figure 3](#) includes examples of visualisations created using aggregated data captured by OnTrack from beta testers (all healthy individuals) between June-August 2019. By adding self-reported SUS data to the analyses, the team will be able to explore the potential correlation between SUS and OnTrack usage.

Subgroup analyses are planned based on patient demographics, stroke disability, stroke subtype and the care pathway patients go through during the intervention period.

All data will be stored and accessed in accordance with GDPR guidance.

Clinical trial support will be provided by the Big Data and Analytical Unit (BDAU) at Imperial College London's Institute of Global Health Innovation (IGHI).

Patient and public involvement (PPI)

To date, over 100 stroke survivors, carers and therapists have been involved in the design of OnTrack. Participants and have been instrumental in highlighting areas for improvement in upper-limb stroke rehabilitation. They have contributed to a co-design process (including workshops, interviews, observations and surveys) resulting in the design, development and initial testing of OnTrack.

A steering group comprising of four stroke survivors was formed for the purpose of this feasibility study. Diversity within the group - both in terms of demographics and stroke severity - was considered. The group has supervised the development of all patient-facing material ensuring its clarity. They will also participate in data analysis by helping to refine themes and key messages arising from qualitative interviews. Participants will be trained by experienced researchers for this purpose.

The steering group will meet five times over the duration of the study, including an initial briefing session at the start to outline their involvement. Steering group members will be key members of the research team and their time and travel will be reimbursed according to INVOLVE(43) guidelines.

The PPI involvement plan was shared with Imperial College London's PPI 'Research Partners Group' on 21.02.19 who felt that the needs of the steering group have been accounted for.

Ethics and dissemination

The OnTrack study will be conducted in accordance with the recommendations for physicians involved in research on human subjects adopted by the 18th World Medical Assembly, Helsinki 1964 and later revisions; and in compliance with the relevant UK and European legislation including the NHS Health Research Authority (HRA) policy frameworks and the General Data Protection Regulation 2018 (GDPR).

The study was granted ethical approval by the HRA, Health and Care Research Wales, and the London - Surrey Research Ethics Committee (ref. 19/LO/0881). Local site capacity and capability approval has been granted by the hospital Trust.

The current approved protocol version is V1.3 dated 19.06.2019. Protocol amendments will be submitted for approval to the NHS HRA in the first instance and to the local site thereafter ahead of implementation.

The Chief Investigator is responsible for preserving the confidentiality of participants taking part in the study. Researchers will have patients' names, contact numbers, emails and home addresses for the purposes of arranging visits. This information will be stored in accordance with GDPR legislation. Participants are free to withdraw from the study at any time. However, anonymised activity data collected may still be used for data analysis as this is unlinked of any patient identifiable information.

The day-to-day management of the study will be coordinated by the Helix Centre. A study steering committee formed by the intervention team, evaluation team, PPI group, and representatives from the local site will meet at regular intervals throughout the study.

Regular updates about the trial will be made available through social media, blog posts, newsletters and the Helix Centre website (www.helixcentre.com). Trial results will be submitted for publication in journals, presented at national and international stroke meetings and conferences and disseminated amongst stroke communities.

Trial status

The first participant was enrolled on 09.09.2019 and recruitment is expected to complete by the end of March 2020. Enrolment and data collection was continuing as planned at the time of submission of this protocol.

Author statement

AD is grant holder and has project oversight along with DD. GF, EG and LH developed the intervention and conceived of the study. GF, EG, and FJ initiated the study design and ET and ML helped with further refinement. EG and GF are responsible for delivering the intervention and data collection. FJ and ET are responsible for the process evaluation. ML provides statistical expertise in trial design and is conducting the primary statistical analysis. All authors contributed to the refinement of the study protocol and approved the final manuscript.

Funding

This study is funded by the NIHR Imperial Biomedical Research Centre (BRC), grant 1215-20013. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Data statement

Anonymised data will be made available in a public repository once the data have obtained validation through publication.

Conflict of interests

FJ is the founder of the social enterprise Bridges Self-Management. She has not received any financial support for this work that could have influenced the design.

Acknowledgments

The authors would like to acknowledge the contributions of the members of the PPI group; Jennifer Crow and Sarah Daniels for their input on recruitment and sample size calculations; and Dr Gaby Judah for her help in defining some aspects of the intervention.

Figure legends

Figure 1

Trial diagram

Figure 2

Logic model

Figure 3

Examples of visualisations created using aggregated data captured by OnTrack from healthy beta testers between June-August 2019

Table 1

Outcome measures

Concept	Assessment	Week of administration
Patient Activation / Engagement	Patient Activation Measure (PAM)	1, 8, 14
Arm impairment	Fugl-Meyer Assessment for upper extremity (FMA-UE)	1, 8, 14
Arm function	Upper-Extremity Motor Activity Log-14 (MAL)	1, 8, 14
Gross level of disability	modified Rankin Scale (mRS)	1, 8, 14
Arm pain	Visual Analogue Scale (VAS)	1, 8, 14
Cognitive impairment	Montreal Cognitive Assessment (MoCA)	1, 8, 14
Arm neglect	Albert's Test (AT)	1, 8, 14
Quality of life	EQ-5D-5L	1, 8, 14
Arm function	Lap-to-Table (LTT)	1, 8, 14
Service experience	Friends and Family Test (FTT)	8, 14
System usability	System Usability Scale (SUS)	14

Table 2

Participation schedule

Week	Phase	Description	OnTrack consultation	Assessments
0	Information and consent	NHS therapists screen for eligible patients, provide information and consent participants		Screening, information, and consent
1	Baseline assessment (initial)	Participants complete outcome measures and wear activity trackers (Axivity AX3) on both arms for one week to gather a baseline of activity allowing left-to-right usage comparison		PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT
2	OnTrack intervention	Participants wear a smartwatch (Apple Watch Series 3 or 4) on their impaired arm only. They will receive real-time feedback on the amount of movement completed (measured in minutes) and daily motivational messages. Participants will receive fortnightly consultations with a researcher to troubleshoot and receive self-management skills training Baseline assessments are repeated during week 8 (halfway)	Onboarding	
3			Check-in & self-management skills training (Problem Solving)	
4				
5			Check-in & self-management skills training (Self-Discovery)	
6				
7			Check-in & self-management skills training (Goal Setting)	
8			Halfway assessment Check-in & self-management skills training (Goal Setting cont.)	PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT, FFT
9				

10			Check-in & self-management skills training (Reflection)	
11				
12			Check-in & self-management skills training (Sign-posting)	
13				
14	Baseline assessment (exit)	Participants complete outcome measures and wear activity trackers (Axivity AX3) on both arms for one week to gather a baseline of activity allowing left-to-right usage comparison		PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT, FFT, SUS
15	Feedback	Independent evaluator leads feedback sessions with participants who have completed the intervention End of participation		Semi-structured interview, online survey (therapists)

References

- (1) WHO. *The Atlas of Heart Disease and Stroke*. Available from: https://www.who.int/cardiovascular_diseases/resources/atlas/en/ [Accessed 04 Oct 2019].
- (2) Stroke Association. *State of the Nation: stroke statistics*. Stroke Association. 2018.
- (3) Salter K, Hellings C, Foley N, Teasell R. The Experience of Living with Stroke: A Qualitative Meta-Synthesis. *Journal of Rehabilitation Medicine*. 2008; 40 (8): 595-602. Available from: doi: 10.2340/16501977-0238 Available from: <http://www.ingentaconnect.com/content/mjl/sreh/2008/00000040/00000008/art00003> .
- (4) Pollock A, Farmer SE, Brady MC, Langhorne P, Mead GE, Mehrholz J, et al. Interventions for improving upper limb function after stroke. *The Cochrane database of systematic reviews*. 2014; 2014 (11): CD010820. Available from: doi: 10.1002/14651858.CD010820.pub2 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25387001> .
- (5) Care Quality Commission. *Supporting life after stroke: a review of services for people who have had a stroke and their carers*. London: Care Quality Commission. 2011.
- (6) Van Peppen RP, Kwakkel G, Wood-Dauphinee S, Hendriks HJ, Van der Wees, Ph J, Dekker J. The impact of physical therapy on functional outcomes after stroke: what's the evidence? *Clinical Rehabilitation*. 2004; 18 (8): 833-862. Available from: doi: 10.1191/0269215504cr8430a Available from: <https://journals.sagepub.com/doi/full/10.1191/0269215504cr8430a> .
- (7) Pollock A, St George B, Fenton M, Firkins L. Top 10 Research Priorities Relating to Life after

- Stroke – Consensus from Stroke Survivors, Caregivers, and Health Professionals. *International Journal of Stroke*. 2014; 9 (3): 313-320. Available from: doi: 10.1111/j.1747-4949.2012.00942.x Available from: <https://journals.sagepub.com/doi/full/10.1111/j.1747-4949.2012.00942.x> .
- (8) Kunkel D, Fitton C, Burnett M, Ashburn A. Physical inactivity post-stroke: a 3-year longitudinal study. *Disability & Rehabilitation*. 2015; 37 (4): 304-310. Available from: doi: 10.3109/09638288.2014.918190 Available from: <http://www.tandfonline.com/doi/abs/10.3109/09638288.2014.918190> .
- (9) Bernhardt J, Dewey H, Thrift A, Donnan G. Inactive and Alone: Physical Activity Within the First 14 Days of Acute Stroke Unit Care. *Stroke*. 2004; 35 (4): 1005-1009. Available from: doi: 10.1161/01.STR.0000120727.40792.40 Available from: <http://stroke.ahajournals.org/cgi/content/abstract/35/4/1005> .
- (10) Laver KE, Lange B, George S, Deutsch JE, Saposnik G, Crotty M. Virtual reality for stroke rehabilitation. *The Cochrane database of systematic reviews*. 2017; 11 (11): CD008349. Available from: doi: 10.1002/14651858.CD008349.pub4 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29156493> .
- (11) Loureiro R, Harwin W, Nagai K, Johnson M. Advances in upper limb stroke rehabilitation: a technology push. *Medical & Biological Engineering & Computing*. 2011; 49 (10): 1103-1118. Available from: doi: 10.1007/s11517-011-0797-0 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21773806> .
- (12) Jones F, Gage H, Drummond A, Bhalla A, Grant R, Lennon S, et al. Feasibility study of an integrated stroke self-management programme: a cluster-randomised controlled trial. *BMJ Open*. 2016; 6 (1): e008900. Available from: doi: 10.1136/bmjopen-2015-008900 Available from: <http://dx.doi.org/10.1136/bmjopen-2015-008900> .
- (13) Harwood M, Weatherall M, Talemaitoga A, Barber PA, Gommans J, Taylor W, et al. Taking charge after stroke: promoting self-directed rehabilitation to improve quality of life – a randomized controlled trial. *Clinical Rehabilitation*. 2012; 26 (6): 493-501. Available from: doi: 10.1177/0269215511426017 Available from: <https://journals.sagepub.com/doi/full/10.1177/0269215511426017> .
- (14) Kendall E, Catalano T, Kuipers P, Posner N, Buys N, Charker J. Recovery following stroke: The role of self-management education. *Social Science & Medicine*. 2007; 64 (3): 735-746. Available from: doi: 10.1016/j.socscimed.2006.09.012 Available from: <https://www.sciencedirect.com/science/article/pii/S0277953606004801> .
- (15) Grady PA, Gough LL. Self-Management: A Comprehensive Approach to Management of Chronic Conditions. *American journal of public health*. 2014; 104 (8): e25-e31. Available from: doi: 10.2105/AJPH.2014.302041 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24922170> .
- (16) Helix Centre. *Project OnTrack*. Available from: <https://helixcentre.com/project-ontrack> [Accessed 04 Oct 2019].
- (17) Buckland D. Leaving hospital is 'like falling off a cliff' for stroke survivors. *Raconteur*. APRIL 27, 2017 . Available from: <https://www.raconteur.net/healthcare/leaving-hospital-is-like-falling-off-a-cliff-for-stroke-survivors> [Accessed Oct 04 2019].
- (18) McKeivitt C, Fudge N, Redfern J, Sheldenkar A, Crichton S, Rudd AR, et al. Self-Reported Long-Term Needs After Stroke. *Stroke*. 2011; 42 (5): 1398-1403. Available from: doi: 10.1161/STROKEAHA.110.598839 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21441153> .
- (19) Woodman P, Riaz A, Pereira C, Jones F. Social participation post stroke: a meta-ethnographic review of the experiences and views of community-dwelling stroke survivors. *Disability & Rehabilitation*. 2014; 36 (24): 2031-2043. Available from: doi: 10.3109/09638288.2014.887796 Available from: <http://www.tandfonline.com/doi/abs/10.3109/09638288.2014.887796> .
- (20) Chan A, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Kraljević Jerić K, et al. SPIRIT

- 2013 Statement: Defining Standard Protocol Items for Clinical Trials. *Annals of internal medicine*. 2013; 158 (3): 200-207. Available from: doi: 10.7326/0003-4819-158-3-201302050-00583 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23295957> .
- (21) Billingham SAM, Whitehead AL, Julious SA. An audit of sample sizes for pilot and feasibility trials being undertaken in the United Kingdom registered in the United Kingdom Clinical Research Network database. *BMC medical research methodology*. 2013; 13 (1): 104. Available from: doi: 10.1186/1471-2288-13-104 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23961782> .
- (22) Bridges Self-Management Limited. *Bridges Self-Management*. Available from: <http://www.bridgesselfmanagement.org.uk/> [Accessed 07 Oct 2019].
- (23) Medical Research Institute of New Zealand. *Taking Charge After Stroke (TaCAS)*. Available from: <https://mrinz.ac.nz/programmes/stroke/> [Accessed 07 oct 2019].
- (24) DOCUMAS. *DOCUMAS: efficient health research management*. Available from: <https://www.documas.eu/> [Accessed 04 Oct 2019].
- (25) Hibbard J, Gilbert H. *Supporting people to manage their health An introduction to patient activation*. London: The Kings Fund. 2014.
- (26) Hibbard JH, Stockard J, Mahoney ER, Tusler M. Development of the Patient Activation Measure (PAM): Conceptualizing and Measuring Activation in Patients and Consumers. *Health Services Research*. 2004; 39 (4p1): 1005-1026. Available from: doi: 10.1111/j.1475-6773.2004.00269.x Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1475-6773.2004.00269.x> .
- (27) Roberts NJ, Kidd L, Dougall N, Patel IS, McNarry S, Nixon C. Measuring patient activation: The utility of the Patient Activation Measure within a UK context—Results from four exemplar studies and potential future applications. *Patient Education and Counseling*. 2016; 99 (10): 1739-1746. Available from: doi: 10.1016/j.pec.2016.05.006 Available from: <https://www.clinicalkey.es/playcontent/1-s2.0-S0738399116301938> .
- (28) Sahlgrenska Academy Institute of Neuroscience and Physiology, University of Gothenburg. *Fugl-Meyer*. Available from: https://neurophys.gu.se/english/departments/clinical_neuroscience_and_rehabilitation/medicin/fugl-meyer [Accessed 04 Oct 2019].
- (29) Pamela W Duncan, Martha Propst, Steven G Nelson. Reliability of the Fugl-Meyer Assessment of Sensorimotor Recovery Following Cerebrovascular Accident. *Physical Therapy*. 1983; 63 (10): 1606-1610. Available from: doi: 10.1093/ptj/63.10.1606 Available from: <http://ptjournal.apta.org/content/63/10/1606.abstract> .
- (30) Uswatte G, Taub E, Morris D, Vignolo M, McCulloch K. Reliability and Validity of the Upper-Extremity Motor Activity Log-14 for Measuring Real-World Arm Use. *Stroke*. 2005; 36 (11): 2493-2496. Available from: doi: 10.1161/01.STR.0000185928.90848.2e Available from: <http://stroke.ahajournals.org/cgi/content/abstract/36/11/2493> .
- (31) Quinn TJ, Dawson J, Walters MR, Lees KR. Exploring the Reliability of the Modified Rankin Scale. *Stroke*. 2009; 40 (3): 762-766. Available from: doi: 10.1161/STROKEAHA.108.522516 Available from: <http://stroke.ahajournals.org/cgi/content/abstract/40/3/762> .
- (32) Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison of six methods. *Pain*. 1986; 27 (1): 117-126. Available from: doi: 10.1016/0304-3959(86)90228-9 Available from: <https://www.sciencedirect.com/science/article/pii/0304395986902289> .
- (33) Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment. *Journal of the American Geriatrics Society*. 2005; 53 (4): 695-699. Available from: doi: 10.1111/j.1532-5415.2005.53221.x Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1532-5415.2005.53221.x> .
- (34) Fullerton KJ, McSherry D, Stout RW. Albert's test: a neglected test of perceptual neglect. *Lancet (London, England)*. 1986; 1 (8478): 430. Available from:

1
2
3 <https://www.ncbi.nlm.nih.gov/pubmed/2868349> .

4 (35) EuroQol - a new facility for the measurement of health-related quality of life. *Health policy*.
5 1990; 16 (3): 199-208. Available from: doi: 10.1016/0168-8510(90)90421-9 Available from:
6 <https://www.sciencedirect.com/science/article/pii/0168851090904219> .

7 (36) NHS England. *The Friends and Family Test*. NHS England. 2014.

8 (37) SUS: A 'Quick and Dirty' Usability Scale. *Usability Evaluation In Industry*. CRC Press; 1996.
9 pp. 207-212.

10 (38) Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W, et al. Process evaluation
11 of complex interventions: Medical Research Council guidance. *BMJ : British Medical Journal*.
12 2015; 350 (mar19 6): h1258. Available from: doi: 10.1136/bmj.h1258 Available from:
13 <http://dx.doi.org/10.1136/bmj.h1258> .

14 (39) W.K. Kellogg Foundation. *Logic Model Development Guide*. Battle Creek, Michigan: W.K.
15 Kellogg Foundation. 2004.

16 (40) The Scottish Government. *Designing and Evaluating Behaviour Change Interventions*. The
17 Scottish Government. 2015.

18 (41) QSR International Ltd. *NVIVO*. Available from:
19 <https://www.qsrinternational.com/nvivo/nvivo-products> [Accessed 07 Oct 2019].

20 (42) Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in*
21 *Psychology*. 2006; 3 (2): 77-101. Available from: doi: 10.1191/1478088706qp063oa Available
22 from: <http://www.tandfonline.com/doi/abs/10.1191/1478088706qp063oa> .

23 (43) National Institute for Health Research. *About INVOLVE*. Available from:
24 <https://www.invo.org.uk/about-involve/> [Accessed 04 Oct 2019].
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1

Trial diagram




Figure 2


Logic model



Figure 3


Examples of visualisations created using aggregated data captured by OnTrack from healthy beta testers between June-August 2019

Active minutes per hour of day (0-23) by user, aggregated over time



Source: Sample OnTrack usage data collected from healthy beta testers, June-August 2019

Active minutes per day by user



Source: Sample OnTrack usage data collected from healthy beta testers, June-August 2019

Reporting checklist for protocol of a clinical trial.

Based on the SPIRIT guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the SPIRIT reporting guidelines, and cite them as:

Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, Hróbjartsson A, Mann H, Dickersin K, Berlin J, Doré C, Parulekar W, Summerskill W, Groves T, Schulz K, Sox H, Rockhold FW, Rennie D, Moher D. SPIRIT 2013 Statement: Defining standard protocol items for clinical trials. *Ann Intern Med.* 2013;158(3):200-207

		Reporting Item	Page Number
Administrative information			
Title	#1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1
Trial registration	#2a	Trial identifier and registry name. If not yet registered, name of intended registry	2
Trial registration: data set	#2b	All items from the World Health Organization Trial Registration Data Set	n/a
Protocol version	#3	Date and version identifier	11
Funding	#4	Sources and types of financial, material, and other support	12
Roles and responsibilities: contributorship	#5a	Names, affiliations, and roles of protocol contributors	11

1	Roles and	#5b	Name and contact information for the trial sponsor	n/a
2	responsibilities:			
3	sponsor contact			
4	information			
5				
6				
7				
8	Roles and	#5c	Role of study sponsor and funders, if any, in study design;	11
9	responsibilities:		collection, management, analysis, and interpretation of data;	
10	sponsor and funder		writing of the report; and the decision to submit the report for	
11			publication, including whether they will have ultimate authority	
12			over any of these activities	
13				
14				
15				
16	Roles and	#5d	Composition, roles, and responsibilities of the coordinating centre,	n/a
17	responsibilities:		steering committee, endpoint adjudication committee, data	
18	committees		management team, and other individuals or groups overseeing the	
19			trial, if applicable (see Item 21a for data monitoring committee)	
20				
21				
22				
23	Introduction			
24				
25	Background and	#6a	Description of research question and justification for undertaking	3
26	rationale		the trial, including summary of relevant studies (published and	
27			unpublished) examining benefits and harms for each intervention	
28				
29				
30				
31	Background and	#6b	Explanation for choice of comparators	n/a
32	rationale: choice of			
33	comparators			
34				
35				
36	Objectives	#7	Specific objectives or hypotheses	4
37				
38	Trial design	#8	Description of trial design including type of trial (eg, parallel	5
39			group, crossover, factorial, single group), allocation ratio, and	
40			framework (eg, superiority, equivalence, non-inferiority,	
41			exploratory)	
42				
43				
44				
45	Methods:			
46	Participants,			
47	interventions, and			
48	outcomes			
49				
50				
51				
52	Study setting	#9	Description of study settings (eg, community clinic, academic	5
53			hospital) and list of countries where data will be collected.	
54			Reference to where list of study sites can be obtained	
55				
56				
57	Eligibility criteria	#10	Inclusion and exclusion criteria for participants. If applicable,	5
58			eligibility criteria for study centres and individuals who will	
59				
60				

		perform the interventions (eg, surgeons, psychotherapists)	
Interventions: description	#11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	6
Interventions: modifications	#11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving / worsening disease)	n/a
Interventions: adherence	#11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return; laboratory tests)	n/a
Interventions: concomitant care	#11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	n/a
Outcomes	#12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	7
Participant timeline	#13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	Table 2
Sample size	#14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	6
Recruitment	#15	Strategies for achieving adequate participant enrolment to reach target sample size	6
Methods: Assignment of interventions (for controlled trials)			
Allocation: sequence generation	#16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	n/a

1	Allocation concealment	#16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	n/a
2	mechanism			
3				
4				
5				
6				
7				
8	Allocation:	#16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	n/a
9	implementation			
10				
11	Blinding (masking)	#17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	n/a
12				
13				
14				
15				
16				
17	Blinding (masking):	#17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	n/a
18	emergency unblinding			
19				
20				
21				
22	Methods: Data			
23	collection,			
24	management, and			
25	analysis			
26				
27				
28				
29	Data collection plan	#18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	9
30				
31				
32				
33				
34				
35				
36				
37				
38				
39	Data collection plan:	#18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	n/a
40	retention			
41				
42				
43				
44	Data management	#19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	9
45				
46				
47				
48				
49				
50				
51	Statistics: outcomes	#20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	9
52				
53				
54				
55				
56	Statistics: additional	#20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	9
57	analyses			
58				
59				
60				

1	Statistics: analysis	#20c	Definition of analysis population relating to protocol non-	n/a
2	population and missing		adherence (eg, as randomised analysis), and any statistical methods	
3	data		to handle missing data (eg, multiple imputation)	
4				
5				
6	Methods: Monitoring			
7				
8	Data monitoring:	#21a	Composition of data monitoring committee (DMC); summary of its	n/a
9	formal committee		role and reporting structure; statement of whether it is independent	
10			from the sponsor and competing interests; and reference to where	
11			further details about its charter can be found, if not in the protocol.	
12			Alternatively, an explanation of why a DMC is not needed	
13				
14	Data monitoring:	#21b	Description of any interim analyses and stopping guidelines,	n/a
15	interim analysis		including who will have access to these interim results and make	
16			the final decision to terminate the trial	
17				
18	Harms	#22	Plans for collecting, assessing, reporting, and managing solicited	n/a
19			and spontaneously reported adverse events and other unintended	
20			effects of trial interventions or trial conduct	
21				
22	Auditing	#23	Frequency and procedures for auditing trial conduct, if any, and	n/a
23			whether the process will be independent from investigators and the	
24			sponsor	
25				
26				
27				
28	Ethics and			
29	dissemination			
30				
31	Research ethics	#24	Plans for seeking research ethics committee / institutional review	10
32	approval		board (REC / IRB) approval	
33				
34	Protocol amendments	#25	Plans for communicating important protocol modifications (eg,	10
35			changes to eligibility criteria, outcomes, analyses) to relevant	
36			parties (eg, investigators, REC / IRBs, trial participants, trial	
37			registries, journals, regulators)	
38				
39	Consent or assent	#26a	Who will obtain informed consent or assent from potential trial	6
40			participants or authorised surrogates, and how (see Item 32)	
41				
42	Consent or assent:	#26b	Additional consent provisions for collection and use of participant	n/a
43	ancillary studies		data and biological specimens in ancillary studies, if applicable	
44				
45	Confidentiality	#27	How personal information about potential and enrolled participants	10
46			will be collected, shared, and maintained in order to protect	
47			confidentiality before, during, and after the trial	
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				

1	Declaration of interests	#28	Financial and other competing interests for principal investigators for the overall trial and each study site	12
2				
3				
4	Data access	#29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	10
5				
6				
7				
8				
9				
10	Ancillary and post trial care	#30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	n/a
11				
12				
13				
14	Dissemination policy: trial results	#31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	10
15				
16				
17				
18				
19				
20				
21	Dissemination policy: authorship	#31b	Authorship eligibility guidelines and any intended use of professional writers	n/a
22				
23				
24	Dissemination policy: reproducible research	#31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	n/a
25				
26				
27				
28	Appendices			
29				
30				
31	Informed consent materials	#32	Model consent form and other related documentation given to participants and authorised surrogates	n/a
32				
33				
34	Biological specimens	#33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	n/a
35				
36				
37				
38				
39				

The SPIRIT checklist is distributed under the terms of the Creative Commons Attribution License CC-BY-ND 3.0. This checklist was completed on 07. October 2019 using <https://www.goodreports.org/>, a tool made by the [EQUATOR Network](#) in collaboration with [Penelope.ai](#)

BMJ Open

Protocol for a feasibility study of OnTrack: a digital system for upper-limb rehabilitation after stroke.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034936.R1
Article Type:	Protocol
Date Submitted by the Author:	26-Dec-2019
Complete List of Authors:	Fusari, Gianpaolo; Royal College of Art, Helix Centre; Imperial College London, Helix Centre Gibbs, Ella; Imperial College London, Helix Centre Hoskin, Lily; Imperial College London, Helix Centre Dickens, Daniel; Imperial College London, Helix Centre Leis, Melanie; Imperial College London, Big Data and Analytical Unit, Institute of Global Health Innovation Taylor, Elizabeth; Kingston and St George's Faculty of Health Social Care and Education Jones, Fiona; Kingston and St George's Faculty of Health Social Care and Education Darzi, Ara; Imperial College London, Helix Centre
Primary Subject Heading:	Neurology
Secondary Subject Heading:	Rehabilitation medicine, Neurology, Public health
Keywords:	Stroke < NEUROLOGY, REHABILITATION MEDICINE, NEUROLOGY, PUBLIC HEALTH

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Title

Protocol for a feasibility study of OnTrack: a digital system for upper-limb rehabilitation after stroke.

Authors and affiliations

Gianpaolo Fusari¹, Ella Gibbs¹, Lily Hoskin¹, Daniel Dickens¹, Melanie Leis², Elizabeth Taylor³, Fiona Jones³, Ara Darzi¹

¹Helix Centre, Imperial College London and the Royal College of Art, London, United Kingdom

²Big Data and Analytical Unit, Institute of Global Health Innovation, Imperial College London, London, United Kingdom

³Kingston University and St George's, University of London, London, United Kingdom

Correspondence to:

Gianpaolo Fusari

HELIX Centre, Institute of Global Health Innovation

Imperial College London, 3rd Floor, Paterson Building, St Mary's Hospital, 16 South Wharf Road, London W2 1PF, UK

gianpaolo@helixcentre.com

+44 (0)77 6779 2770

Key words

Stroke, rehabilitation medicine, neurology, public health, digital health

Word count

4223

Abstract

Introduction

Arm weakness is a common problem after stroke (affecting 450,000 people in the UK) leading to loss of independence. Repetitive activity is critical for recovery but research shows people struggle with knowing what or how much to do, and keeping track of progress. Working with more than 100 therapists (occupational therapists and physiotherapists) and patients with stroke, we co-developed the OnTrack intervention - consisting of software for smart-devices and coaching support - that has the potential to address this problem. This is a protocol to assess the feasibility of OnTrack for evaluation in a randomised control trial.

Methods and analysis

A mixed methods, single-arm study design will be used to evaluate the feasibility of OnTrack for hospital and community use. A minimum sample of 12 participants from a stroke unit will be involved in the study for 14 weeks. During week 1, 8 and 14 participants will complete assessments relating to their arm function, arm impairment, and activation. During weeks 2-13 participants will use OnTrack to track their arm movement in real time, receive motivational messages, and face-to-face sessions to address problems, gain feedback on activity, and receive self-management skills coaching. All equipment will be loaned to study participants. A parallel process evaluation will be conducted to assess the intervention's fidelity, dose and reach, using a mixed methods approach. A Public and Patient Involvement (PPI) group will oversee the study and help with interpretation and dissemination of qualitative and quantitative data findings.

Ethics and dissemination

Ethical approval granted by the NHS Health Research Authority, Health and Care Research Wales, and the London - Surrey Research Ethics Committee (ref.19/LO/0881). Trial results will be submitted for publication in peer review journals, presented at international conferences and disseminated amongst stroke communities. The results of this trial will inform development of a definitive trial.

Trial registration details

ClinicalTrials.gov (NCT03944486), pre-results.

Strengths and limitations

- This is a feasibility trial of a novel intervention which employs an integrated approach for tracking arm activity and coaching with the aim of increasing stroke survivors' confidence and ability to use their impaired arm in daily activities, increasing the opportunities for repetitive rehabilitation (repeating a movement or series of movements with a rehabilitative or functional aim).
- Patient and Public Involvement (PPI) with more than 100 stroke survivors, carers, and clinicians have contributed to our needs-finding phase, co-designed OnTrack and informed the feasibility study. A new PPI group will oversee the running of the study and help with interpretation of qualitative and quantitative data findings.
- An independent process evaluation will provide detailed information about implementation, context, and the mechanisms of impact of the intervention. Findings will help in the understanding of intervention fidelity and training needs required for a definitive trial.
- For pragmatic reasons the study uses a non-randomised design carried out at a single site- this will limit understanding about randomisation and recruitment
- Participants will not be followed-up after intervention period; however participant views will be sought regarding appropriate follow-up times in a subsequent definitive trial.

Introduction

Every year around the world over 15 million people experience a stroke, leaving 5 million people with a permanent disability.(1) Stroke is the leading cause of disability in the UK; half of the nearly 1.2 million stroke survivors who live in the country have some form of disability, significantly contributing to the loss of independence and feeling of isolation that they experience. (2),(3) Furthermore, stroke is estimated to cost UK society £26 billion every year, with the vast majority of these costs borne by the informal care sector.(2)

Upper-limb (arm) weakness is the main cause of physical impairment affecting 75% of disabled stroke survivors; this equates to around 450,000 people in the UK.(2) Dose-intensive repetitive rehabilitation is widely accepted as the 'gold-standard' for regaining ability after stroke, however, NHS resources are often limited and unable to provide this.(4) A recent Cochrane review of over 500 trials failed to yield high-quality practice recommendations for interventions for the upper-limb.(5) Arm recovery after stroke is a national research priority.(6) There is a correlation between physical activity after stroke and the ability to perform activities of daily living (most of which involve the use of the arm) nonetheless, studies suggest that the actual time patients are active is minimal.(7,8) Many current approaches to solving this problem focus on improving the prescribed rehabilitation sessions, often employing gamification techniques.(9,10) Whilst this is important, there is untapped potential to increase repetitive rehabilitation by targeting the large proportion of the day where patients are going about their daily activities and can use their arm movement

(however small) to a greater extent. Capacity for activity could be increased further by using self-management methods as demonstrated by several different programmes in stroke and other long-term conditions.(11-14) This has informed the development of OnTrack which aims to increase opportunities for activity by improving individuals' self-management skills through tailored support and real-time activity feedback on their arm movement.

An ethnographic study conducted by the Helix Centre (funded by Innovate UK) confirmed what other studies have shown (7,8,15) that patients struggle to see and keep track of improvements, this impacts their motivation and leaves them dependent on therapists for feedback. Stroke survivors often report feeling unsupported after leaving hospital and not knowing how to best help themselves improve their arm function.(16-18) Feedback gathered from over 100 stroke survivors and clinicians was the basis for developing the OnTrack intervention.

A proof-of-concept test of OnTrack gathered data from a small group of patients (n=7) and confirmed that the intervention was safe and generally users could understand how and when to use it. Participants reported they were more aware of their impaired arm and had increased confidence in using it for new tasks. A 20% mean increase in minutes of activity on the impaired arm was observed. The work conducted to date is unpublished and has some limitations however it has shaped the intervention and suggests that OnTrack has the potential to be a scalable solution that requires minimal training and could be used in conjunction with NHS services to help increase the overall amount of activity performed with the impaired arm. This study will assess the feasibility of the OnTrack intervention and inform the design of a definitive randomised controlled trial (RCT) to evaluate its clinical effectiveness, and follows the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) guidelines.(19)

Methods and analysis

Aims and objectives

The primary aim is to evaluate the feasibility of an RCT to test the effectiveness of the OnTrack intervention for upper limb rehabilitation after stroke.

The objectives are to:

- Assess the feasibility of recruitment from hyper-acute and acute stroke units, and rehabilitation wards to ascertain strategy and recruitment rates.
- Assess dropout rates by observing adherence and compliance with the intervention.
- Understand the acceptability and usability of the intervention by stroke survivors.
- Understand the acceptability of study procedures by healthcare professionals.
- Explore implementation fidelity, dose and reach of the OnTrack intervention.

The study will also collect clinical outcomes regarding arm function, impairment and activation to identify an appropriate primary outcome, and to estimate parameters for a sample size calculation for an RCT ([Table 1](#)).

Study design

A feasibility study with a nested process evaluation ([Figure 1](#)). The study is a single-site, non-randomised intervention trial. The design of the study was developed through a collaborative approach between the study researchers, a PPI steering group, front-line therapists, and the Research Design Service at the National Institute for Health Research.

An independent process evaluation will be conducted in parallel to learn about usage and engagement mechanisms of participants, therapists and other frontline staff, providing critical information for implementation fidelity and impact mechanisms necessary for scale-up.

Study setting

The study will be conducted at an inner city NHS hospital Trust in London. Recruited participants will be able to continue to receive the intervention at home if discharged from hospital prior to ending the intervention period (14 weeks).

Participants

The inclusion criteria encompasses:

- Adults (aged 18 or over).
- Stroke diagnosis less than 6 months previously (first or recurrent). Some participants will be recruited from an in-patient rehabilitation ward, hence the 6 month post-stroke limit.
- Arm impairment of any type or level (including weakness - including dense hemiplegia, neglect, and sensory deficits). This to enable better understanding of which impairment level groups would benefit *or not* from using the intervention, especially considering the impact it may have on people's motivation regardless of their level of impairment.
- Ability to provide informed consent.
- Reliability to communicate (verbally or nonverbally) and understand English.
- Ability to read a predefined short message.

Potential participants who at the time of recruitment present with any of the following will be excluded:

- Unstable medical condition.
- Self reported "severe" pain in the arm affected either at rest or during movement.
- Severe oedema in the arm affected by their stroke, judged by the consenting therapist.
- Known discharge plans to a hospital other than the site Trust or residential care in less than 7 weeks (a small proportion of patients staying at CNRU may be in hospital for up to 12 weeks).

Recruitment

Participants will be recruited from the Hyperacute Stroke Unit (HASU), Acute Stroke Unit (ASU), and Clinical Neurorehabilitation Unit (CNRU) at an inner city NHS hospital Trust in London.

Stroke therapists (occupational therapists, physiotherapists) will be responsible for screening and identifying suitable patients. They will introduce the study to potential participants and provide information documents. Potential participants will be given a minimum of 24 hours to consider the advantages and disadvantages of participating in the study and to formulate questions. Therapists will be able to answer questions or will liaise with the research team to provide an answer. Once all questions are answered and a potential participant is willing to participate, consent will be taken by the therapist. Only at this stage will patient information be shared with the research team. There may be situations where a therapist is only able to take verbal consent from a participant due to time or material constraints, in such cases the researchers will be able to take written consent from the participant upon first meeting them.

Sample size calculation

Guidelines advocate a sample size of 12-30 participants for feasibility studies.(20) Experienced clinical academics and clinicians at the trial site have advised to expect about 50% of eligible patients to agree to participation and a 50% completion rate. This has informed a recruitment plan to identify at least 60 potential participants in a period of 30 weeks to reach the minimum sample size.

Intervention

The intervention is the OnTrack system as a whole. The system consists of smart-devices (smartphone and smartwatch), software (OnTrack app), and coaching support. Smart-devices are used to track arm movement. Motivational messages and a real-time display of completed arm activity (in minutes) are presented to the user via the OnTrack app. Coaching support is provided through fortnightly consultations by the researchers. During consultations, participants will receive self-management training informed by the Bridges Self-Management (21) and TaCAS (22) self-management programs. Coaching sessions are themed around principles of self-management (see Table 2, OnTrack Consultation column).

Data gathered by the OnTrack system can be accessed by the researchers via a digital dashboard to inform consultations.

Participants will be loaned all equipment necessary for the trial and no previous experience with using smart-devices is required to participate. Technical support will be provided only in cases where the hardware and/or software fail to perform the required functions to deliver the intervention.

[Table 2](#) provides a participation schedule and a summary of the intervention procedures.

Outcomes

Feasibility of trial design and procedures

- Recruitment strategy and rates (feasibility of recruitment from HASU, ASU, CNRU wards) - percentage of patients: screened; eligible; approached; consented; excluded after screening. Participants consented and recruited will be logged by the research team in DOCUMAS(23)
- Compliance and adherence to intervention - measure of minutes of activity per participant as recorded by the OnTrack app, engagement with OnTrack app as measured by system analytics (for example: compliance with starting tracking arm activity daily, number of times and times of the day a particular screen is visited, the number of messages read and replied to, etc.)
- Completion rates - percentage of participants who complete the 14-week intervention period
- Acceptability and reasons for decline/withdrawal - number of participants who withdraw or decline the intervention and reasons why. A record of reasons for withdrawal and declining will be kept by the researchers. Reasons will be categorised in order of most common; this information will help the research team understand the reasons why someone might drop out or decline to participate in the study.

Clinical assessments

As a secondary objective, clinical outcomes will be collected at different time points by a qualified member of the research team to identify an appropriate primary outcome, and to estimate parameters for a sample size calculation for an RCT ([Table 1](#)). The outcome measures and assessments are listed below.

Patient activation

Patient activation is a concept recognised by the NHS that describes the knowledge, skills and confidence a person has in managing their own health and health care.(24) This will be measured using the Patient Activation Measure (PAM)(25) which has been validated in stroke populations in the UK.(26) The PAM survey measures patients on a 0–100 scale and can categorise patients into one of four activation levels along an empirically derived continuum.(25) Activation levels will be used to allocate participants one of three different OnTrack coaching tiers. The tiers aim to make the different aspects of the coaching more relevant and meaningful for the individual participant and their stage of recovery and self-management.

Arm impairment

Arm impairment will be measured objectively using the Fugl-Meyer Assessment for upper extremity (FMA-UE).(27) The FMA-UE has been tested extensively, and is found to have excellent psychometric properties and is recommended as core measures to be used in every stroke recovery and rehabilitation trial.(28)

Arm function

Arm function will be assessed using the Upper-Extremity Motor Activity Log-14 (MAL).(29) The MAL is a scripted, structured interview developed to self-report the amount and quality of use of the impaired arm in individuals with stroke in 14 different activities of daily living.

Gross level of disability

The modified Rankin Scale (mRS)(30) is the most prevalent functional outcome measure in contemporary stroke trials. The mRS quantifies disability using an ordinal hierarchical grading from zero (no symptoms) to 5 (severe disability).

Arm pain

Pain will be assessed using a visual analogue scale (VAS) from 0 (no pain) to 10 (excruciating pain) over the last 24 hours. VAS is a valid measure of pain intensity and is responsive to change.(31) Individuals scoring 3/10 or more in the affected arm will be withdrawn from the study unless their pain is only on movements that are not part of their usual everyday activities (e.g. arm pain when doing overhead reaching).

Cognitive impairment

Cognitive impairment will be assessed using the Montreal Cognitive Assessment (MoCA). The MoCA is a brief cognitive screening tool with high sensitivity and specificity for detecting mild cognitive impairment.(32) The MoCA defines impairment as follows: score of 18-25 = mild, 10-17 = moderate, <10 = severe.(32) Participants' scores will be used to look for associations between the use of OnTrack and any cognitive impairment.

Perceptual neglect

Albert's Test (AT) is being used to assess for unilateral spatial neglect (USN). This a simple test where participants are asked to cross out lines ruled in a standard fashion on a sheet of paper. If any lines are left uncrossed, and more than 70% of uncrossed lines are on the same side as motor deficit, USN is indicated. This may be quantified in terms of the percentage of lines left uncrossed. The test is very easy to administer and is a good predictor of functional activity six months after stroke onset.(33)

Quality of life

The EQ-5D-5L is a widely used standardised preference based measure of health status developed by the EuroQol Group in order to provide a simple, generic measure of health for clinical and economic appraisal.(34)

Additional assessments

A Lap-to-Table (LTT) timed test will be performed where the researchers measure the time it takes a participant to move their hand three times from resting on their lap to a table

1
2
3 positioned in front of them. This test is performed to assess its potential to use as part of the
4 inclusion criteria for an RCT.
5

6
7 The NHS Friends and Family Test (FFT)(35) will be used to obtain feedback on the overall
8 experience of using OnTrack and participating in the trial. Participants will be asked: “How
9 likely are you to recommend OnTrack to friends and family if they needed similar care or
10 treatment?” with answers provided in a Likert 5-point scale ranging from “extremely likely” to
11 “extremely unlikely” and an “I don’t know” option.
12

13 The System Usability Scale (SUS) will be used to subjectively assess the usability of the
14 OnTrack intervention. The test is a simple, ten-item scale covering a variety of aspects of
15 system usability, such as the need for support, training, and complexity, and thus have a
16 high level of face validity for measuring the usability of a system.(36)
17
18

19 **Process evaluation**

20
21 A process evaluation will be carried out by researchers working independently to the
22 intervention team and in parallel to the trial to determine whether the OnTrack intervention
23 was delivered as intended and to understand the mechanisms of impact. The aim of the
24 process evaluation at the feasibility stage is mainly to understand how the trial design and
25 intervention could be optimised ahead of an RCT.(37) A logic model(38,39) that defines the
26 intervention in terms of inputs, outputs, causal assumptions and expected outcomes has
27 been developed to help identify core questions for the evaluation team to explore (Figure 2).
28 The evaluation team will observe 10% of all intervention sessions with the objective of
29 documenting fidelity, dose and reach of the intervention.
30
31

32
33 Critical reflection and the process evaluation will help refine the intervention, as shown by
34 mid-range theories (i.e. theories that help understand implementation).(40) Interim results
35 will be shared with the intervention team at the half-way point with the objective to review
36 some of the procedures and make minor adjustments as necessary.
37

38
39 In-depth semi-structured interviews will be conducted with patients at the end of their
40 participation, a minimum sample of 12 is anticipated. A topic guide with themes drawing
41 from the logic model will be used. Interviews will focus on participants’ experiences using
42 OnTrack, their perceptions of arm tracking, motivational messaging and the researcher
43 consultations. Additionally, the interviews will explore participants’ perceptions of the impact
44 OnTrack had on them in terms of progress, awareness, participation, and confidence in self-
45 management. Participants’ responses will be compared against activity data collected from
46 the OnTrack app.
47
48

49
50 NHS therapists caring for participants taking part will be consented and invited to complete a
51 short online survey to gather their feedback regarding acceptability of study procedures,
52 they have the option to respond anonymously. The total number of therapists involved is
53 difficult to predict as there may be team changes and staff movement during the course of
54 the study. The survey will ask questions around three themes: 1) participation, relevance,
55 quality and time spent in study procedures; 2) opinions on the benefit/detriment OnTrack
56
57
58
59

1
2
3 may have for patients; 3) opinions on how the intervention may or may not fit with service
4 provision and their workflow.
5

6 7 **Data analysis**

8
9 Analysis will be completed on the parameters and implementation of the study in addition to
10 the usability of OnTrack.
11

12 Data collected for the process evaluation will be a combination of qualitative data from
13 interviews with stroke participants and therapists to explore their experiences of using
14 OnTrack, as well as quantitative data on usage of OnTrack and the self-reported SUS.
15 OnTrack therapy support sessions will be monitored through a fidelity checklist and
16 observations (10 live sessions will be observed in total. In addition, the evaluation team will
17 have access to recorded sessions that can be observed at their discretion). Interview data
18 will undergo thematic analysis by the evaluation team. Data will be entered into NVIVO,(41)
19 line by line coding and analysis will be informed by Braun and Clark's approach to thematic
20 analysis.(42)
21

22 Changes over time will be evaluated in both OnTrack usage and outcome measures.
23

24 For OnTrack usage, the team will analyse users' activity patterns by day and hour of day.
25 [Figure 3](#) illustrates examples of visualisations created using aggregated data captured by
26 OnTrack from healthy beta testers between June-August 2019. It compares users on active
27 minutes per hour of day (aggregated over time) and active minutes per day.
28

29 OnTrack also captures specific usage metrics, including:
30

- 31 ● Number of times OnTrack messages were opened
- 32 ● Number of times daily and weekly activity were viewed on the phone
- 33 ● Number of swipes on watch to reveal activity graph

34 These values will be plotted against the users' minutes of activity to better understand the
35 potential impact of the app on activity over time.
36

37 The self-reported PAM will be captured at weeks 1, 8 and 14 for each user. It will be
38 analysed in relation to the minutes of activity of each user over time to better understand the
39 potential impact of the app on their levels of activation. SUS will be captured at weeks 8 and
40 14 and will be compared against actual usage metrics (described above) to assess usability.
41

42 Subgroup analyses are planned based on patient demographics, stroke disability (measured
43 by mRankin scale at start of participation), stroke subtype and the care pathway patients go
44 through during the intervention period.
45

46 The number of subgroups that will be available for analyses will depend on the
47 characteristics of the participants. Whilst it's clear that the sample size will be relatively
48 small, it is valuable to understand how we might approach subgroup analysis in a definitive
49 trial with a larger sample size.
50

All data will be stored and accessed in accordance with GDPR guidance.

Clinical trial support will be provided by the Big Data and Analytical Unit (BDAU) at Imperial College London's Institute of Global Health Innovation (IGHI).

Patient and public involvement (PPI)

To date, over 100 stroke survivors, carers and therapists have been involved in the design of OnTrack. Participants and have been instrumental in highlighting areas for improvement in upper-limb stroke rehabilitation. They have contributed to a co-design process (including workshops, interviews, observations and surveys) resulting in the design, development and initial testing of OnTrack.

A steering group comprising of four stroke survivors was formed for the purpose of this feasibility study. Diversity within the group - both in terms of demographics and stroke severity - was considered. The group has supervised the development of all patient-facing material ensuring its clarity. They will also participate in data analysis by helping to refine themes and key messages arising from qualitative interviews. Participants will be trained by experienced researchers for this purpose.

The steering group will meet five times over the duration of the study, including an initial briefing session at the start to outline their involvement. Steering group members will be key members of the research team and their time and travel will be reimbursed according to INVOLVE(43) guidelines.

The PPI involvement plan was shared with Imperial College London's PPI 'Research Partners Group' on 21.02.19 who felt that the needs of the steering group have been accounted for.

Ethics and dissemination

The OnTrack study will be conducted in accordance with the recommendations for physicians involved in research on human subjects adopted by the 18th World Medical Assembly, Helsinki 1964 and later revisions; and in compliance with the relevant UK and European legislation including the NHS Health Research Authority (HRA) policy frameworks and the General Data Protection Regulation 2018 (GDPR).

The study was granted ethical approval by the HRA, Health and Care Research Wales, and the London - Surrey Research Ethics Committee (ref. 19/LO/0881). Local site capacity and capability approval has been granted by the hospital Trust.

The current approved protocol version is V1.3 dated 19.06.2019. Protocol amendments will be submitted for approval to the NHS HRA in the first instance and to the local site thereafter ahead of implementation.

1
2
3 The Chief Investigator is responsible for preserving the confidentiality of participants taking
4 part in the study. Researchers will have patients' names, contact numbers, emails and home
5 addresses for the purposes of arranging visits. This information will be stored in accordance
6 with GDPR legislation. Participants are free to withdraw from the study at any time.
7 However, anonymised activity data collected may still be used for data analysis as this is
8 unlinked of any patient identifiable information.
9

10
11 The day-to-day management of the study will be coordinated by the Helix Centre. A study
12 steering committee formed by the intervention team, evaluation team, PPI group, and
13 representatives from the local site will meet at regular intervals throughout the study.
14

15
16 Regular updates about the trial will be made available through social media, blog posts,
17 newsletters and the Helix Centre website (www.helixcentre.com). Trial results will be
18 submitted for publication in journals, presented at national and international stroke meetings
19 and conferences and disseminated amongst stroke communities.
20
21

22 Trial status

23
24 The first participant was enrolled on 09.09.2019 and recruitment is expected to complete by
25 the end of March 2020. Enrolment and data collection was continuing as planned at the time
26 of submission of this protocol.
27
28

29 Author statement

30
31 AD is grant holder and has project oversight along with DD. GF, EG and LH developed the
32 intervention and conceived of the study. GF, EG, and FJ initiated the study design and ET
33 and ML helped with further refinement. EG and GF are responsible for delivering the
34 intervention and data collection. FJ and ET are responsible for the process evaluation. ML
35 provides statistical expertise in trial design and is conducting the primary statistical analysis.
36 All authors contributed to the refinement of the study protocol and approved the final
37 manuscript.
38
39
40

41 Funding

42
43 This study is funded by the NIHR Imperial Biomedical Research Centre (BRC), grant 1215-
44 20013. The views expressed are those of the author(s) and not necessarily those of the
45 NIHR or the Department of Health and Social Care.
46
47

48
49 All intellectual property associated with the OnTrack system is owned by the Helix Centre
50 which is a collaboration between Imperial College London and the Royal College of Art.
51

52 Data statement

53
54 Anonymised data will be made available in a public repository once the data have obtained
55 validation through publication.
56
57
58
59

Conflict of interests

FJ is the founder of the social enterprise Bridges Self-Management. She has not received any financial support for this work that could have influenced the design.

Acknowledgments

The authors would like to acknowledge the contributions of the members of the PPI group; Jennifer Crow and Sarah Daniels for their input on recruitment and sample size calculations; and Dr Gaby Judah for her help in defining some aspects of the intervention.

Figure legends

Figure 1

Trial diagram

Figure 2

Logic model

Figure 3

Examples of visualisations created using aggregated data captured by OnTrack from healthy beta testers. Data for a minimum of 5 and a maximum of 18 days were aggregated for the period between June and August 2019.

Table 1

Outcome measures

Concept	Assessment	Week of administration
Patient Activation / Engagement	Patient Activation Measure (PAM)	1, 8, 14
Arm impairment	Fugl-Meyer Assessment for upper extremity (FMA-UE)	1, 8, 14
Arm function	Upper-Extremity Motor Activity Log-14 (MAL)	1, 8, 14
Gross level of disability	modified Rankin Scale (mRS)	1, 8, 14
Arm pain	Visual Analogue Scale (VAS)	1, 8, 14
Cognitive impairment	Montreal Cognitive Assessment (MoCA)	1, 8, 14
Arm neglect	Albert's Test (AT)	1, 8, 14
Quality of life	EQ-5D-5L	1, 8, 14
Arm function	Lap-to-Table (LTT)	1, 8, 14
Service experience	Friends and Family Test (FTT)	8, 14
System usability	System Usability Scale (SUS)	14

Table 2

Intervention and Participation schedule

Week	Phase	Description	OnTrack consultation	Assessments
0	Information and consent	NHS therapists screen for eligible patients, provide information and consent participants		Screening, information, and consent
1	Baseline assessment (initial)	Participants complete outcome measures and wear activity trackers (Axivity AX3) on both arms during waking hours (typically 12 hours per day) for one week to gather accelerometer data which is translated into minutes of activity. This data creates a baseline of activity allowing left-to-right arm usage comparison		PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT
2	OnTrack intervention	Participants wear a smartwatch (Apple Watch Series 3 or 4) on their impaired arm only during waking hours (typically 12 hours per day). They will receive real-time feedback on the amount of movement completed (measured in minutes) and daily motivational messages. Participants will receive fortnightly consultations with a researcher to troubleshoot and receive self-management skills training	Onboarding	
3			Check-in & self-management skills training (Problem Solving)	
4				
5			Check-in & self-management skills training (Self-Discovery)	
6			Baseline assessments are repeated during week 8 (halfway)	
7			Check-in & self-management skills training (Goal Setting)	
8			Halfway assessment Check-in & self-management skills training (Goal Setting)	PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT, FFT

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

BMJ Open: first published as 10.1136/bmjopen-2019-034936 on 23 March 2020. Downloaded from <http://bmjopen.bmj.com/> on May 2, 2025 at Department GEZ-LTA

Erasmus Hogeschool

			cont.)	
9				
10			Check-in & self-management skills training (Reflection)	
11				
12			Check-in & self-management skills training (Sign-posting)	
13				
14	Baseline assessment (exit)	Participants complete outcome measures and wear activity trackers (Axivity AX3) on both arms during waking hours (typically 12 hours per day) for one week to gather accelerometer data which is translated into minutes of activity. This data creates a baseline of activity allowing left-to-right arm usage comparison		PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT, FFT, SUS
15	Feedback	Independent evaluator leads feedback sessions with participants who have completed the intervention End of participation		Semi-structured interview, online survey (therapists)

References

References

- (1) WHO. *The Atlas of Heart Disease and Stroke*. Available from: https://www.who.int/cardiovascular_diseases/resources/atlas/en/ [Accessed 04 Oct 2019].
- (2) Stroke Association. *State of the Nation: stroke statistics*. Stroke Association. 2018.
- (3) Salter K, Hellings C, Foley N, Teasell R. The Experience of Living with Stroke: A Qualitative Meta-Synthesis. *Journal of Rehabilitation Medicine*. 2008; 40 (8): 595-602. Available from: doi: 10.2340/16501977-0238 Available from: <http://www.ingentaconnect.com/content/mjl/sreh/2008/00000040/00000008/art00003>.

- 1 (4) Care Quality Commission. *Supporting life after stroke: a review of services for people who*
2 *have had a stroke and their carers*. London: Care Quality Commission. 2011.
- 3 (5) Pollock A, Farmer SE, Brady MC, Langhorne P, Mead GE, Mehrholz J, et al. Interventions
4 for improving upper limb function after stroke. *The Cochrane database of systematic reviews*.
5 2014; 2014 (11): CD010820. Available from: doi: 10.1002/14651858.CD010820.pub2 Available
6 from: <https://www.ncbi.nlm.nih.gov/pubmed/25387001> .
- 7 (6) Pollock A, St George B, Fenton M, Firkins L. Top 10 Research Priorities Relating to Life after
8 Stroke – Consensus from Stroke Survivors, Caregivers, and Health Professionals. *International*
9 *Journal of Stroke*. 2014; 9 (3): 313-320. Available from: doi: 10.1111/j.1747-4949.2012.00942.x
10 Available from: <https://journals.sagepub.com/doi/full/10.1111/j.1747-4949.2012.00942.x> .
- 11 (7) Kunkel D, Fitton C, Burnett M, Ashburn A. Physical inactivity post-stroke: a 3-year
12 longitudinal study. *Disability & Rehabilitation*. 2015; 37 (4): 304-310. Available from: doi:
13 10.3109/09638288.2014.918190 Available from:
14 <http://www.tandfonline.com/doi/abs/10.3109/09638288.2014.918190> .
- 15 (8) Bernhardt J, Dewey H, Thrift A, Donnan G. Inactive and Alone: Physical Activity Within the
16 First 14 Days of Acute Stroke Unit Care. *Stroke*. 2004; 35 (4): 1005-1009. Available from: doi:
17 10.1161/01.STR.0000120727.40792.40 Available from:
18 <http://stroke.ahajournals.org/cgi/content/abstract/35/4/1005> .
- 19 (9) Laver KE, Lange B, George S, Deutsch JE, Saposnik G, Crotty M. Virtual reality for stroke
20 rehabilitation. *The Cochrane database of systematic reviews*. 2017; 11 (11): CD008349.
21 Available from: doi: 10.1002/14651858.CD008349.pub4 Available from:
22 <https://www.ncbi.nlm.nih.gov/pubmed/29156493> .
- 23 (10) Loureiro R, Harwin W, Nagai K, Johnson M. Advances in upper limb stroke rehabilitation: a
24 technology push. *Medical & Biological Engineering & Computing*. 2011; 49 (10): 1103-1118.
25 Available from: doi: 10.1007/s11517-011-0797-0 Available from:
26 <https://www.ncbi.nlm.nih.gov/pubmed/21773806> .
- 27 (11) Jones F, Gage H, Drummond A, Bhalla A, Grant R, Lennon S, et al. Feasibility study of an
28 integrated stroke self-management programme: a cluster-randomised controlled trial. *BMJ*
29 *Open*. 2016; 6 (1): e008900. Available from: doi: 10.1136/bmjopen-2015-008900 Available from:
30 <http://dx.doi.org/10.1136/bmjopen-2015-008900> .
- 31 (12) Harwood M, Weatherall M, Talemaitoga A, Barber PA, Gommans J, Taylor W, et al. Taking
32 charge after stroke: promoting self-directed rehabilitation to improve quality of life – a
33 randomized controlled trial. *Clinical Rehabilitation*. 2012; 26 (6): 493-501. Available from: doi:
34 10.1177/0269215511426017 Available from:
35 <https://journals.sagepub.com/doi/full/10.1177/0269215511426017> .
- 36 (13) Kendall E, Catalano T, Kuipers P, Posner N, Buys N, Charker J. Recovery following stroke:
37 The role of self-management education. *Social Science & Medicine*. 2007; 64 (3): 735-746.
38 Available from: doi: 10.1016/j.socscimed.2006.09.012 Available from:
39 <https://www.sciencedirect.com/science/article/pii/S0277953606004801> .
- 40 (14) Grady PA, Gough LL. Self-Management: A Comprehensive Approach to Management of
41 Chronic Conditions. *American journal of public health*. 2014; 104 (8): e25-e31. Available from:
42 doi: 10.2105/AJPH.2014.302041 Available from:
43 <https://www.ncbi.nlm.nih.gov/pubmed/24922170> .
- 44 (15) Esmonde T, McGinley J, Wittwer J, Goldie P, Martin C. Stroke rehabilitation: patient activity
45 during non-therapy time. *Australian Journal of Physiotherapy*. 1997; 43 (1): 43-51. Available
46 from: doi: 10.1016/S0004-9514(14)60401-3 Available from:
47 <https://www.sciencedirect.com/science/article/pii/S0004951414604013> .
- 48 (16) Buckland D. Leaving hospital is 'like falling off a cliff' for stroke survivors. *Raconteur*. APRIL
49 27, 2017 . Available from: <https://www.raconteur.net/healthcare/leaving-hospital-is-like-falling-off-a-cliff-for-stroke-survivors> [Accessed Oct 04 2019].
- 50 (17) McKeivitt C, Fudge N, Redfern J, Sheldenkar A, Crichton S, Rudd AR, et al. Self-Reported
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Long-Term Needs After Stroke. *Stroke*. 2011; 42 (5): 1398-1403. Available from: doi:
4 10.1161/STROKEAHA.110.598839 Available from:
5 <https://www.ncbi.nlm.nih.gov/pubmed/21441153> .
6 (18) Woodman P, Riazi A, Pereira C, Jones F. Social participation post stroke: a meta-
7 ethnographic review of the experiences and views of community-dwelling stroke survivors.
8 *Disability & Rehabilitation*. 2014; 36 (24): 2031-2043. Available from: doi:
9 10.3109/09638288.2014.887796 Available from:
10 <http://www.tandfonline.com/doi/abs/10.3109/09638288.2014.887796> .
11 (19) Chan A, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Kraljević K, et al. SPIRIT
12 2013 Statement: Defining Standard Protocol Items for Clinical Trials. *Annals of internal*
13 *medicine*. 2013; 158 (3): 200-207. Available from: doi: 10.7326/0003-4819-158-3-201302050-
14 00583 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23295957> .
15 (20) Billingham SAM, Whitehead AL, Julious SA. An audit of sample sizes for pilot and feasibility
16 trials being undertaken in the United Kingdom registered in the United Kingdom Clinical
17 Research Network database. *BMC medical research methodology*. 2013; 13 (1): 104. Available
18 from: doi: 10.1186/1471-2288-13-104 Available from:
19 <https://www.ncbi.nlm.nih.gov/pubmed/23961782> .
20 (21) Bridges Self-Management Limited. *Bridges Self-Management*. Available from:
21 <http://www.bridgesselfmanagement.org.uk/> [Accessed 07 Oct 2019].
22 (22) Medical Research Institute of New Zealand. *Taking Charge After Stroke (TaCAS)*.
23 Available from: <https://mrinz.ac.nz/programmes/stroke/> [Accessed 07 oct 2019].
24 (23) DOCUMAS. *DOCUMAS: efficient health research management*. Available from:
25 <https://www.documas.eu/> [Accessed 04 Oct 2019].
26 (24) Hibbard J, Gilbert H. *Supporting people to manage their health An introduction to patient*
27 *activation*. London: The Kings Fund. 2014.
28 (25) Hibbard JH, Stockard J, Mahoney ER, Tusler M. Development of the Patient Activation
29 Measure (PAM): Conceptualizing and Measuring Activation in Patients and Consumers. *Health*
30 *Services Research*. 2004; 39 (4p1): 1005-1026. Available from: doi: 10.1111/j.1475-
31 6773.2004.00269.x Available from: [https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1475-](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1475-6773.2004.00269.x)
32 [6773.2004.00269.x](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1475-6773.2004.00269.x) .
33 (26) Roberts NJ, Kidd L, Dougall N, Patel IS, McNarry S, Nixon C. Measuring patient activation:
34 The utility of the Patient Activation Measure within a UK context—Results from four exemplar
35 studies and potential future applications. *Patient Education and Counseling*. 2016; 99 (10):
36 1739-1746. Available from: doi: 10.1016/j.pec.2016.05.006 Available from:
37 <https://www.clinicalkey.es/playcontent/1-s2.0-S0738399116301938> .
38 (27) Sahlgrenska Academy Institute of Neuroscience and Physiology, University of Gothenburg.
39 *Fugl-Meyer*. Available from:
40 [https://neurophys.gu.se/english/departments/clinical_neuroscience_and_rehabilitation/rehabilitat](https://neurophys.gu.se/english/departments/clinical_neuroscience_and_rehabilitation/rehabilitation-medicine/fugl-meyer)
41 [ion-medicine/fugl-meyer](https://neurophys.gu.se/english/departments/clinical_neuroscience_and_rehabilitation/rehabilitation-medicine/fugl-meyer) [Accessed 04 Oct 2019].
42 (28) Pamela W Duncan, Martha Propst, Steven G Nelson. Reliability of the Fugl-Meyer
43 Assessment of Sensorimotor Recovery Following Cerebrovascular Accident. *Physical Therapy*.
44 1983; 63 (10): 1606-1610. Available from: doi: 10.1093/ptj/63.10.1606 Available from:
45 <http://ptjournal.apta.org/content/63/10/1606.abstract> .
46 (29) Uswatte G, Taub E, Morris D, Vignolo M, McCulloch K. Reliability and Validity of the Upper-
47 Extremity Motor Activity Log-14 for Measuring Real-World Arm Use. *Stroke*. 2005; 36 (11):
48 2493-2496. Available from: doi: 10.1161/01.STR.0000185928.90848.2e Available from:
49 <http://stroke.ahajournals.org/cgi/content/abstract/36/11/2493> .
50 (30) Quinn TJ, Dawson J, Walters MR, Lees KR. Exploring the Reliability of the Modified Rankin
51 Scale. *Stroke*. 2009; 40 (3): 762-766. Available from: doi: 10.1161/STROKEAHA.108.522516
52 Available from: <http://stroke.ahajournals.org/cgi/content/abstract/40/3/762> .
53 (31) Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison
54
55
56
57
58
59

- of six methods. *Pain*. 1986; 27 (1): 117-126. Available from: doi: 10.1016/0304-3959(86)90228-9 Available from: <https://www.sciencedirect.com/science/article/pii/0304395986902289> .
- (32) Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment. *Journal of the American Geriatrics Society*. 2005; 53 (4): 695-699. Available from: doi: 10.1111/j.1532-5415.2005.53221.x Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1532-5415.2005.53221.x> .
- (33) Fullerton KJ, McSherry D, Stout RW. Albert's test: a neglected test of perceptual neglect. *Lancet (London, England)*. 1986; 1 (8478): 430. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/2868349> .
- (34) EuroQol - a new facility for the measurement of health-related quality of life. *Health policy*. 1990; 16 (3): 199-208. Available from: doi: 10.1016/0168-8510(90)90421-9 Available from: <https://www.sciencedirect.com/science/article/pii/0168851090904219> .
- (35) NHS England. *The Friends and Family Test*. NHS England. 2014.
- (36) SUS: A 'Quick and Dirty' Usability Scale. *Usability Evaluation In Industry*. CRC Press; 1996. pp. 207-212.
- (37) Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W, et al. Process evaluation of complex interventions: Medical Research Council guidance. *BMJ : British Medical Journal*. 2015; 350 (mar19 6): h1258. Available from: doi: 10.1136/bmj.h1258 Available from: <http://dx.doi.org/10.1136/bmj.h1258> .
- (38) W.K. Kellogg Foundation. *Logic Model Development Guide*. Battle Creek, Michigan: W.K. Kellogg Foundation. 2004.
- (39) The Scottish Government. *Designing and Evaluating Behaviour Change Interventions*. The Scottish Government. 2015.
- (40) Kislov R, Pope C, Martin GP, Wilson PM. Harnessing the power of theorising in implementation science. *Implementation Science*. 2019; 14 (1): 103. Available from: doi: 10.1186/s13012-019-0957-4 Available from: <https://search.proquest.com/docview/2324921962> .
- (41) QSR International Ltd. *NVIVO*. Available from: <https://www.qsrinternational.com/nvivo/nvivo-products> [Accessed 07 Oct 2019].
- (42) Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in Psychology*. 2006; 3 (2): 77-101. Available from: doi: 10.1191/1478088706qp063oa Available from: <http://www.tandfonline.com/doi/abs/10.1191/1478088706qp063oa> .
- (43) National Institute for Health Research. *About INVOLVE*. Available from: <https://www.invo.org.uk/about-involve/> [Accessed 04 Oct 2019].

Figure 1

Trial diagram




Figure 2


Logic model



Figure 3

Examples of visualisations created using aggregated data captured by OnTrack from healthy beta testers between June-August 2019

Active minutes per hour of day (0-23) by user, aggregated over time



Source: Sample OnTrack usage data collected from healthy beta testers, June-August 2019

Active minutes per day by user



Source: Sample OnTrack usage data collected from healthy beta testers, June-August 2019

Reporting checklist for protocol of a clinical trial.

Based on the SPIRIT guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the SPIRIT reporting guidelines, and cite them as:

Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, Hróbjartsson A, Mann H, Dickersin K, Berlin J, Doré C, Parulekar W, Summerskill W, Groves T, Schulz K, Sox H, Rockhold FW, Rennie D, Moher D. SPIRIT 2013 Statement: Defining standard protocol items for clinical trials. *Ann Intern Med.* 2013;158(3):200-207

		Reporting Item	Page Number
Administrative information			
Title	#1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1
Trial registration	#2a	Trial identifier and registry name. If not yet registered, name of intended registry	2
Trial registration: data set	#2b	All items from the World Health Organization Trial Registration Data Set	n/a
Protocol version	#3	Date and version identifier	11
Funding	#4	Sources and types of financial, material, and other support	12
Roles and responsibilities: contributorship	#5a	Names, affiliations, and roles of protocol contributors	11

1	Roles and	#5b	Name and contact information for the trial sponsor	n/a
2	responsibilities:			
3	sponsor contact			
4	information			
5				
6				
7				
8	Roles and	#5c	Role of study sponsor and funders, if any, in study design;	11
9	responsibilities:		collection, management, analysis, and interpretation of data;	
10	sponsor and funder		writing of the report; and the decision to submit the report for	
11			publication, including whether they will have ultimate authority	
12			over any of these activities	
13				
14				
15				
16	Roles and	#5d	Composition, roles, and responsibilities of the coordinating centre,	n/a
17	responsibilities:		steering committee, endpoint adjudication committee, data	
18	committees		management team, and other individuals or groups overseeing the	
19			trial, if applicable (see Item 21a for data monitoring committee)	
20				
21				
22				
23	Introduction			
24				
25	Background and	#6a	Description of research question and justification for undertaking	3
26	rationale		the trial, including summary of relevant studies (published and	
27			unpublished) examining benefits and harms for each intervention	
28				
29				
30	Background and	#6b	Explanation for choice of comparators	n/a
31	rationale: choice of			
32	comparators			
33				
34				
35				
36	Objectives	#7	Specific objectives or hypotheses	4
37				
38	Trial design	#8	Description of trial design including type of trial (eg, parallel	5
39			group, crossover, factorial, single group), allocation ratio, and	
40			framework (eg, superiority, equivalence, non-inferiority,	
41			exploratory)	
42				
43				
44				
45	Methods:			
46	Participants,			
47	interventions, and			
48	outcomes			
49				
50				
51				
52	Study setting	#9	Description of study settings (eg, community clinic, academic	5
53			hospital) and list of countries where data will be collected.	
54			Reference to where list of study sites can be obtained	
55				
56				
57	Eligibility criteria	#10	Inclusion and exclusion criteria for participants. If applicable,	5
58			eligibility criteria for study centres and individuals who will	
59				
60				

		perform the interventions (eg, surgeons, psychotherapists)	
Interventions: description	#11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	6
Interventions: modifications	#11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving / worsening disease)	n/a
Interventions: adherence	#11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return; laboratory tests)	n/a
Interventions: concomitant care	#11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	n/a
Outcomes	#12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	7
Participant timeline	#13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	Table 2
Sample size	#14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	6
Recruitment	#15	Strategies for achieving adequate participant enrolment to reach target sample size	6
Methods: Assignment of interventions (for controlled trials)			
Allocation: sequence generation	#16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	n/a

1	Allocation concealment	#16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	n/a
2	mechanism			
3				
4				
5				
6				
7				
8	Allocation:	#16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	n/a
9	implementation			
10				
11				
12	Blinding (masking)	#17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	n/a
13				
14				
15				
16				
17	Blinding (masking):	#17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	n/a
18	emergency unblinding			
19				
20				
21				
22	Methods: Data			
23	collection,			
24	management, and			
25	analysis			
26				
27				
28				
29	Data collection plan	#18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	9
30				
31				
32				
33				
34				
35				
36				
37				
38				
39	Data collection plan:	#18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	n/a
40	retention			
41				
42				
43				
44	Data management	#19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	9
45				
46				
47				
48				
49				
50				
51	Statistics: outcomes	#20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	9
52				
53				
54				
55				
56	Statistics: additional	#20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	9
57	analyses			
58				
59				
60				

1	Statistics: analysis	#20c	Definition of analysis population relating to protocol non-	n/a
2	population and missing		adherence (eg, as randomised analysis), and any statistical methods	
3	data		to handle missing data (eg, multiple imputation)	
4				
5				
6	Methods: Monitoring			
7				
8	Data monitoring:	#21a	Composition of data monitoring committee (DMC); summary of its	n/a
9	formal committee		role and reporting structure; statement of whether it is independent	
10			from the sponsor and competing interests; and reference to where	
11			further details about its charter can be found, if not in the protocol.	
12			Alternatively, an explanation of why a DMC is not needed	
13				
14	Data monitoring:	#21b	Description of any interim analyses and stopping guidelines,	n/a
15	interim analysis		including who will have access to these interim results and make	
16			the final decision to terminate the trial	
17				
18	Harms	#22	Plans for collecting, assessing, reporting, and managing solicited	n/a
19			and spontaneously reported adverse events and other unintended	
20			effects of trial interventions or trial conduct	
21				
22	Auditing	#23	Frequency and procedures for auditing trial conduct, if any, and	n/a
23			whether the process will be independent from investigators and the	
24			sponsor	
25				
26				
27	Ethics and			
28	dissemination			
29				
30	Research ethics	#24	Plans for seeking research ethics committee / institutional review	10
31	approval		board (REC / IRB) approval	
32				
33	Protocol amendments	#25	Plans for communicating important protocol modifications (eg,	10
34			changes to eligibility criteria, outcomes, analyses) to relevant	
35			parties (eg, investigators, REC / IRBs, trial participants, trial	
36			registries, journals, regulators)	
37	Consent or assent	#26a	Who will obtain informed consent or assent from potential trial	6
38			participants or authorised surrogates, and how (see Item 32)	
39				
40	Consent or assent:	#26b	Additional consent provisions for collection and use of participant	n/a
41	ancillary studies		data and biological specimens in ancillary studies, if applicable	
42				
43	Confidentiality	#27	How personal information about potential and enrolled participants	10
44			will be collected, shared, and maintained in order to protect	
45			confidentiality before, during, and after the trial	
46				
47				
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				

1	Declaration of interests	#28	Financial and other competing interests for principal investigators for the overall trial and each study site	12
2				
3				
4	Data access	#29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	10
5				
6				
7				
8				
9				
10	Ancillary and post trial care	#30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	n/a
11				
12				
13				
14	Dissemination policy: trial results	#31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	10
15				
16				
17				
18				
19				
20				
21	Dissemination policy: authorship	#31b	Authorship eligibility guidelines and any intended use of professional writers	n/a
22				
23				
24	Dissemination policy: reproducible research	#31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	n/a
25				
26				
27				
28	Appendices			
29				
30				
31	Informed consent materials	#32	Model consent form and other related documentation given to participants and authorised surrogates	n/a
32				
33				
34				
35	Biological specimens	#33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	n/a
36				
37				
38				
39				

40 The SPIRIT checklist is distributed under the terms of the Creative Commons Attribution License CC-BY-ND
 41 3.0. This checklist was completed on 07. October 2019 using <https://www.goodreports.org/>, a tool made by the
 42 [EQUATOR Network](#) in collaboration with [Penelope.ai](#)
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59

BMJ Open

Protocol for a feasibility study of OnTrack: a digital system for upper-limb rehabilitation after stroke.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034936.R2
Article Type:	Protocol
Date Submitted by the Author:	25-Feb-2020
Complete List of Authors:	Fusari, Gianpaolo; Royal College of Art, Helix Centre; Imperial College London, Helix Centre Gibbs, Ella; Imperial College London, Helix Centre Hoskin, Lily; Imperial College London, Helix Centre Dickens, Daniel; Imperial College London, Helix Centre Leis, Melanie; Imperial College London, Big Data and Analytical Unit, Institute of Global Health Innovation Taylor, Elizabeth; Kingston and St George's Faculty of Health Social Care and Education Jones, Fiona; Kingston and St George's Faculty of Health Social Care and Education Darzi, Ara; Imperial College London, Helix Centre
Primary Subject Heading:	Neurology
Secondary Subject Heading:	Rehabilitation medicine, Neurology, Public health
Keywords:	Stroke < NEUROLOGY, REHABILITATION MEDICINE, NEUROLOGY, PUBLIC HEALTH

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Title

Protocol for a feasibility study of OnTrack: a digital system for upper-limb rehabilitation after stroke.

Authors and affiliations

Gianpaolo Fusari¹, Ella Gibbs¹, Lily Hoskin¹, Daniel Dickens¹, Melanie Leis², Elizabeth Taylor³, Fiona Jones³, Ara Darzi¹

¹Helix Centre, Imperial College London and the Royal College of Art, London, United Kingdom

²Big Data and Analytical Unit, Institute of Global Health Innovation, Imperial College London, London, United Kingdom

³Kingston University and St George's, University of London, London, United Kingdom

Correspondence to:

Gianpaolo Fusari

HELIX Centre, Institute of Global Health Innovation

Imperial College London, 3rd Floor, Paterson Building, St Mary's Hospital, 16 South Wharf Road, London W2 1PF, UK

gianpaolo@helixcentre.com

+44 (0)77 6779 2770

Key words

Stroke, rehabilitation medicine, neurology, public health, digital health

Word count

4198

Abstract

Introduction

Arm weakness is a common problem after stroke (affecting 450,000 people in the UK) leading to loss of independence. Repetitive activity is critical for recovery but research shows people struggle with knowing what or how much to do, and keeping track of progress. Working with more than 100 therapists (occupational therapists and physiotherapists) and patients with stroke, we co-developed the OnTrack intervention - consisting of software for smart-devices and coaching support - that has the potential to address this problem. This is a protocol to assess the feasibility of OnTrack for evaluation in a randomised control trial.

Methods and analysis

A mixed methods, single-arm study design will be used to evaluate the feasibility of OnTrack for hospital and community use. A minimum sample of 12 participants from a stroke unit will be involved in the study for 14 weeks. During week 1, 8 and 14 participants will complete assessments relating to their arm function, arm impairment, and activation. During weeks 2-13 participants will use OnTrack to track their arm movement in real time, receive motivational messages, and face-to-face sessions to address problems, gain feedback on activity, and receive self-management skills coaching. All equipment will be loaned to study participants. A parallel process evaluation will be conducted to assess the intervention's fidelity, dose and reach, using a mixed methods approach. A Public and Patient Involvement (PPI) group will oversee the study and help with interpretation and dissemination of qualitative and quantitative data findings.

Ethics and dissemination

Ethical approval granted by the NHS Health Research Authority, Health and Care Research Wales, and the London - Surrey Research Ethics Committee (ref.19/LO/0881). Trial results will be submitted for publication in peer review journals, presented at international conferences and disseminated amongst stroke communities. The results of this trial will inform development of a definitive trial.

Trial registration details

ClinicalTrials.gov (NCT03944486), pre-results.

Strengths and limitations

- This is a feasibility trial of a novel intervention which employs an integrated approach for tracking arm activity and coaching with the aim of increasing stroke survivors' confidence and ability to use their impaired arm in daily activities, increasing the opportunities for repetitive rehabilitation (repeating a movement or series of movements with a rehabilitative or functional aim).
- Patient and Public Involvement (PPI) with more than 100 stroke survivors, carers, and clinicians have contributed to our needs-finding phase, co-designed OnTrack and informed the feasibility study. A new PPI group will oversee the running of the study and help with interpretation of qualitative and quantitative data findings.
- An independent process evaluation will provide detailed information about implementation, context, and the mechanisms of impact of the intervention. Findings will help in the understanding of intervention fidelity and training needs required for a definitive trial.
- For pragmatic reasons the study uses a non-randomised design carried out at a single site- this will limit understanding about randomisation and recruitment
- Participants will not be followed-up after intervention period; however participant views will be sought regarding appropriate follow-up times in a subsequent definitive trial.

Introduction

Every year around the world over 15 million people experience a stroke, leaving 5 million people with a permanent disability.(1) Stroke is the leading cause of disability in the UK; half of the nearly 1.2 million stroke survivors who live in the country have some form of disability, significantly contributing to the loss of independence and feeling of isolation that they experience. (2),(3) Furthermore, stroke is estimated to cost UK society £26 billion every year, with the vast majority of these costs borne by the informal care sector.(2)

Upper-limb (arm) weakness is the main cause of physical impairment affecting 75% of disabled stroke survivors; this equates to around 450,000 people in the UK.(2) Dose-intensive repetitive rehabilitation is widely accepted as the 'gold-standard' for regaining ability after stroke, however, NHS resources are often limited and unable to provide this.(4) A recent Cochrane review of over 500 trials failed to yield high-quality practice recommendations for interventions for the upper-limb.(5) Arm recovery after stroke is a national research priority.(6) There is a correlation between physical activity after stroke and the ability to perform activities of daily living (most of which involve the use of the arm).(7) Despite this evidence, studies suggest that the actual time patients are active is minimal.(8) Many current approaches to increasing repetitive rehabilitation focus on improving the prescribed rehabilitation sessions (typically lasting 45-60 minutes), often by employing gamification techniques.(9,10)

1
2
3 Whilst this is important, there is untapped potential to increase repetitive rehabilitation by
4 targeting the large proportion of the day where patients are going about their daily activities
5 and can use their arm movement (however small) to a greater extent. Capacity for activity
6 could be increased further by using self-management methods as demonstrated by several
7 different programmes in stroke and other long-term conditions.(11-14) This has informed the
8 development of OnTrack which aims to increase opportunities for activity by improving
9 individuals' self-management skills through tailored support and real-time activity feedback
10 on their arm movement.
11
12

13
14 An unpublished ethnographic study conducted by the Helix Centre (funded by Innovate UK)
15 confirmed what other studies have shown (7,8,15) that patients struggle to see and keep
16 track of improvements, this impacts their motivation and leaves them dependent on
17 therapists for feedback. Stroke survivors often report feeling unsupported after leaving
18 hospital and not knowing how to best help themselves improve their arm function.(16-18)
19 Feedback gathered from over 100 stroke survivors and clinicians was the basis for
20 developing the OnTrack intervention.
21
22

23 A proof-of-concept test of OnTrack gathered data from a small group of patients (n=7) and
24 confirmed that the intervention was safe and generally users could understand how and
25 when to use it. Participants reported they were more aware of their impaired arm and had
26 increased confidence in using it for new tasks. A 20% mean increase in minutes of activity
27 on the impaired arm was observed. The work conducted to date is unpublished and has
28 some limitations however it has shaped the intervention and suggests that OnTrack has the
29 potential to be a scalable solution that requires minimal training and could be used in
30 conjunction with NHS services to help increase the overall amount of activity performed with
31 the impaired arm. This study will assess the feasibility of the OnTrack intervention and
32 inform the design of a definitive randomised controlled trial (RCT) to evaluate its clinical
33 effectiveness, and follows the Standard Protocol Items: Recommendations for Interventional
34 Trials (SPIRIT) guidelines.(19)
35
36
37
38

39 **Methods and analysis**

40 **Aims and objectives**

41
42 The primary aim is to evaluate the feasibility of an RCT to test the effectiveness of the
43 OnTrack intervention for upper limb rehabilitation after stroke.
44
45

46 The objectives are to:

- 47 ● Assess the feasibility of recruitment from hyper-acute and acute stroke units, and
48 rehabilitation wards to ascertain strategy and recruitment rates.
 - 49 ● Assess dropout rates by observing adherence and compliance with the intervention.
 - 50 ● Understand the acceptability and usability of the intervention by stroke survivors.
 - 51 ● Understand the acceptability of study procedures by healthcare professionals.
 - 52 ● Explore implementation fidelity, dose and reach of the OnTrack intervention.
- 53
54
55
56
57
58
59
60

The study will also collect clinical outcomes regarding arm function, impairment and activation to identify an appropriate primary outcome, and to estimate parameters for a sample size calculation for an RCT ([Table 1](#)).

Study design

A feasibility study with a nested process evaluation ([Figure 1](#)). The study is a single-site, non-randomised intervention trial. The design of the study was developed through a collaborative approach between the study researchers, a PPI steering group, front-line therapists, and the Research Design Service at the National Institute for Health Research.

An independent process evaluation will be conducted in parallel to learn about usage and engagement mechanisms of participants, therapists and other frontline staff, providing critical information for implementation fidelity and impact mechanisms necessary for scale-up.

Study setting

The study will be conducted at an inner city NHS hospital Trust in London. Recruited participants will be able to continue to receive the intervention at home if discharged from hospital prior to ending the intervention period (14 weeks).

Participants

The inclusion criteria encompasses:

- Adults (aged 18 or over).
- Stroke diagnosis less than 6 months previously (first or recurrent). Some participants will be recruited from an in-patient rehabilitation ward, hence the 6 month post-stroke limit.
- Arm impairment of any type or level (including weakness - including dense hemiplegia, neglect, and sensory deficits). This to enable better understanding of which impairment level groups could benefit *or not* from using the intervention, especially considering the impact it may have on people's motivation regardless of their level of impairment.
- Ability to provide informed consent.
- Reliability to communicate (verbally or nonverbally) and understand English.
- Ability to read a predefined short message.

Potential participants who at the time of recruitment (or during participation) present with any of the following will be excluded:

- Unstable medical condition.
- Self reported "severe" pain in the arm affected either at rest or during movement.
- Severe oedema in the arm affected by their stroke, judged by the consenting therapist.
- Known discharge plans to a hospital other than the site Trust or residential care in less than 7 weeks (a small proportion of patients staying at CNRU may be in hospital for up to 12 weeks).

- Participants who are unable to engage with the intervention for a period of more than 7 consecutive days will be reviewed in a case-by-case basis by the members of the team responsible for delivering the intervention to determine if study continuation is appropriate

Recruitment

Participants will be recruited from the Hyperacute Stroke Unit (HASU), Acute Stroke Unit (ASU), and Clinical Neurorehabilitation Unit (CNRU) at an inner city NHS hospital Trust in London.

Stroke therapists (occupational therapists, physiotherapists) will be responsible for screening and identifying suitable patients. They will introduce the study to potential participants and provide information documents. Potential participants will be given a minimum of 24 hours to consider the advantages and disadvantages of participating in the study and to formulate questions. Therapists will be able to answer questions or will liaise with the research team to provide an answer. Once all questions are answered and a potential participant is willing to participate, consent will be taken by the therapist. Only at this stage will patient information be shared with the research team. There may be situations where a therapist is only able to take verbal consent from a participant due to time or material constraints, in such cases the researchers will be able to take written consent from the participant upon first meeting them.

Sample size calculation

Guidelines advocate a sample size of 12-30 participants for feasibility studies.⁽²⁰⁾ Experienced clinical academics and clinicians at the trial site have advised to expect about 50% of eligible patients to agree to participation and a 50% completion rate. This has informed a recruitment plan to identify at least 60 potential participants in a period of 30 weeks to reach the minimum sample size.

Intervention

The intervention is the OnTrack system as a whole. The system consists of smart-devices (smartphone and smartwatch), software (OnTrack app), and coaching support. Smart-devices are used to track arm movement. Motivational messages and a real-time display of completed arm activity (in minutes) are presented to the user via the OnTrack app. Coaching support is provided through fortnightly consultations by the researchers. During consultations, participants will receive self-management training informed by the Bridges Self-Management (21) and TaCAS (22) self-management programs. Coaching sessions are themed around principles of self-management (see Table 2, OnTrack Consultation column).

Data gathered by the OnTrack system can be accessed by the researchers via a digital dashboard to inform consultations.

Participants will be loaned all equipment necessary for the trial and no previous experience with using smart-devices is required to participate. Technical support will be provided only in cases where the hardware and/or software fail to perform the required functions to deliver the intervention.

[Table 2](#) provides a participation schedule and a summary of the intervention procedures.

Outcomes

Feasibility of trial design and procedures

- Recruitment strategy and rates (feasibility of recruitment from HASU, ASU, CNRU wards) - percentage of patients: screened; eligible; approached; consented; excluded after screening. Participants consented and recruited will be logged by the research team in DOCUMAS(23)
- Compliance and adherence to intervention - measure of minutes of activity per participant as recorded by the OnTrack app, engagement with OnTrack app as measured by system analytics (for example: compliance with starting tracking arm activity daily, number of times and times of the day a particular screen is visited, the number of messages read and replied to, etc.)
- Completion rates - percentage of participants who complete the 14-week intervention period (not dropping out or being withdrawn from the study)
- Acceptability and reasons for decline/withdrawal - number of participants who withdraw or decline the intervention and reasons why. A record of reasons for withdrawal and declining will be kept by the researchers. Reasons will be categorised in order of most common; this information will help the research team understand the reasons why someone might drop out or decline to participate in the study.

Clinical assessments

As a secondary objective, clinical outcomes will be collected at different time points by a qualified member of the research team to identify an appropriate primary outcome, and to estimate parameters for a sample size calculation for an RCT ([Table 1](#)). The outcome measures and assessments are listed below.

Patient activation

Patient activation is a concept recognised by the NHS that describes the knowledge, skills and confidence a person has in managing their own health and health care.(24) This will be measured using the Patient Activation Measure (PAM)(25) which has been validated in stroke populations in the UK.(26) The PAM survey measures patients on a 0–100 scale and can categorise patients into one of four activation levels along an empirically derived continuum.(25) Activation levels will be used to allocate participants one of three different OnTrack coaching tiers. The tiers aim to make the different aspects of the coaching more relevant and meaningful for the individual participant and their stage of recovery and self-management.

Arm impairment

Arm impairment will be measured objectively using the Fugl-Meyer Assessment for upper extremity (FMA-UE).(27) The FMA-UE has been tested extensively, and is found to have excellent psychometric properties and is recommended as core measures to be used in every stroke recovery and rehabilitation trial.(28)

Arm function

Arm function will be assessed using the Upper-Extremity Motor Activity Log-14 (MAL).(29) The MAL is a scripted, structured interview developed to self-report the amount and quality of use of the impaired arm in individuals with stroke in 14 different activities of daily living.

Gross level of disability

The modified Rankin Scale (mRS)(30) is the most prevalent functional outcome measure in contemporary stroke trials. The mRS quantifies disability using an ordinal hierarchical grading from zero (no symptoms) to 5 (severe disability).

Arm pain

Pain will be assessed using a visual analogue scale (VAS) from 0 (no pain) to 10 (excruciating pain) over the last 24 hours. VAS is a valid measure of pain intensity and is responsive to change.(31) Individuals scoring 3/10 or more in the affected arm will be withdrawn from the study unless their pain is only on movements that are not part of their usual everyday activities (e.g. arm pain when doing overhead reaching).

Cognitive impairment

Cognitive impairment will be assessed using the Montreal Cognitive Assessment (MoCA). The MoCA is a brief cognitive screening tool with high sensitivity and specificity for detecting mild cognitive impairment.(32) The MoCA defines impairment as follows: score of 18-25 = mild, 10-17 = moderate, <10 = severe.(32) Participants' scores will be used to look for associations between the use of OnTrack and any cognitive impairment.

Perceptual neglect

Albert's Test (AT) is being used to assess for unilateral spatial neglect (USN). This a simple test where participants are asked to cross out lines ruled in a standard fashion on a sheet of paper. If any lines are left uncrossed, and more than 70% of uncrossed lines are on the same side as motor deficit, USN is indicated. This may be quantified in terms of the percentage of lines left uncrossed. The test is very easy to administer and is a good predictor of functional activity six months after stroke onset.(33)

Quality of life

The EQ-5D-5L is a widely used standardised preference based measure of health status developed by the EuroQol Group in order to provide a simple, generic measure of health for clinical and economic appraisal.(34)

Additional assessments

A Lap-to-Table (LTT) timed test will be performed where the researchers measure the time it takes a participant to move their hand three times from resting on their lap to a table positioned in front of them. This test is performed to assess its potential to use as part of the inclusion criteria for an RCT.

The NHS Friends and Family Test (FFT)(35) will be used to obtain feedback on the overall experience of using OnTrack and participating in the trial. Participants will be asked: “How likely are you to recommend OnTrack to friends and family if they needed similar care or treatment?” with answers provided in a Likert 5-point scale ranging from “extremely likely” to “extremely unlikely” and an “I don’t know” option.

The System Usability Scale (SUS) will be used to subjectively assess the usability of the OnTrack intervention. The test is a simple, ten-item scale covering a variety of aspects of system usability, such as the need for support, training, and complexity, and thus have a high level of face validity for measuring the usability of a system.(36)

Process evaluation

A process evaluation will be carried out by researchers working independently to the intervention team and in parallel to the trial to determine whether the OnTrack intervention was delivered as intended and to understand the mechanisms of impact. The aim of the process evaluation at the feasibility stage is mainly to understand how the trial design and intervention could be optimised ahead of an RCT.(37) A logic model(38,39) that defines the intervention in terms of inputs, outputs, causal assumptions and expected outcomes has been developed to help identify core questions for the evaluation team to explore (Figure 2). The evaluation team will observe 10% of all intervention sessions with the objective of documenting fidelity, dose and reach of the intervention.

Critical reflection and the process evaluation will help refine the intervention, as shown by mid-range theories (i.e. theories that help understand implementation).(40) Interim results will be shared with the intervention team at the half-way point with the objective to review some of the procedures and make minor adjustments as necessary.

In-depth semi-structured interviews will be conducted with patients at the end of their participation, a minimum sample of 12 is anticipated. A topic guide with themes drawing from the logic model will be used. Interviews will focus on participants’ experiences using OnTrack, their perceptions of arm tracking, motivational messaging and the researcher consultations. Additionally, the interviews will explore participants’ perceptions of the impact OnTrack had on them in terms of progress, awareness, participation, and confidence in self-

management. Participants' responses will be compared against activity data collected from the OnTrack app.

NHS therapists caring for participants taking part will be consented and invited to complete a short online survey to gather their feedback regarding acceptability of study procedures, they have the option to respond anonymously. The total number of therapists involved is difficult to predict as there may be team changes and staff movement during the course of the study. The survey will ask questions around three themes: 1) participation, relevance, quality and time spent in study procedures; 2) opinions on the benefit/detriment OnTrack may have for patients; 3) opinions on how the intervention may or may not fit with service provision and their workflow.

Data analysis

Analysis will be completed on the parameters and implementation of the study in addition to the usability of OnTrack.

Data collected for the process evaluation will be a combination of qualitative data from interviews with stroke participants and therapists to explore their experiences of using OnTrack, as well as quantitative data on usage of OnTrack and the self-reported SUS. OnTrack therapy support sessions will be monitored through a fidelity checklist and observations (10 live sessions will be observed in total. In addition, the evaluation team will have access to recorded sessions that can be observed at their discretion). Interview data will undergo thematic analysis by the evaluation team. Data will be entered into NVIVO,(41) line by line coding and analysis will be informed by Braun and Clark's approach to thematic analysis.(42)

Changes over time will be evaluated in both OnTrack usage and outcome measures.

For OnTrack usage, the team will analyse users' activity patterns by day and hour of day. [Figure 3](#) illustrates examples of visualisations created using aggregated data captured by OnTrack from healthy beta testers between June-August 2019. It compares users on active minutes per hour of day (aggregated over time) and active minutes per day.

OnTrack also captures specific usage metrics, including:

- Number of times OnTrack messages were opened
- Number of times daily and weekly activity were viewed on the phone
- Number of swipes on watch to reveal activity graph

For each patient, we will plot the values above against their minutes of activity to better understand the potential impact of the app on activity over time.

The self-reported PAM will be captured at weeks 1, 8 and 14 for each user. It will be analysed in relation to the minutes of activity of each user over time to better understand the potential impact of the app on their levels of activation. SUS will be captured at weeks 8 and 14 and will be compared against actual usage metrics (described above) to assess usability.

1
2
3 Whilst conducting meaningful significant subgroup analyses would be difficult given the
4 relatively small sample size, we believe that outputs from this study could potentially inform
5 the subgroups that might be considered for inclusion in a larger trial.
6

7 All data will be stored and accessed in accordance with GDPR guidance.
8

9
10 Clinical trial support will be provided by the Big Data and Analytical Unit (BDAU) at Imperial
11 College London's Institute of Global Health Innovation (IGHI).
12

13 Patient and public involvement (PPI)

14
15 To date, over 100 stroke survivors, carers and therapists have been involved in the design
16 of OnTrack. Participants and have been instrumental in highlighting areas for improvement
17 in upper-limb stroke rehabilitation. They have contributed to a co-design process (including
18 workshops, interviews, observations and surveys) resulting in the design, development and
19 initial testing of OnTrack.
20

21
22 A steering group comprising of four stroke survivors was formed for the purpose of this
23 feasibility study. Diversity within the group - both in terms of demographics and stroke
24 severity - was considered. The group has supervised the development of all patient-facing
25 material ensuring its clarity. They will also participate in data analysis by helping to refine
26 themes and key messages arising from qualitative interviews. Participants will be trained by
27 experienced researchers for this purpose.
28

29
30 The steering group will meet five times over the duration of the study, including an initial
31 briefing session at the start to outline their involvement. Steering group members will be key
32 members of the research team and their time and travel will be reimbursed according to
33 INVOLVE(43) guidelines.
34

35
36 The PPI involvement plan was shared with Imperial College London's PPI 'Research
37 Partners Group' on 21.02.19 who felt that the needs of the steering group have been
38 accounted for.
39

40 Ethics and dissemination

41
42 The OnTrack study will be conducted in accordance with the recommendations for
43 physicians involved in research on human subjects adopted by the 18th World Medical
44 Assembly, Helsinki 1964 and later revisions; and in compliance with the relevant UK and
45 European legislation including the NHS Health Research Authority (HRA) policy frameworks
46 and the General Data Protection Regulation 2018 (GDPR).
47

48
49 The study was granted ethical approval by the HRA, Health and Care Research Wales, and
50 the London - Surrey Research Ethics Committee (ref. 19/LO/0881). Local site capacity and
51 capability approval has been granted by the hospital Trust.
52
53
54
55
56
57
58
59
60

1
2
3 The current approved protocol version is V1.3 dated 19.06.2019. Protocol amendments will
4 be submitted for approval to the NHS HRA in the first instance and to the local site
5 thereafter ahead of implementation.
6

7
8 The Chief Investigator is responsible for preserving the confidentiality of participants taking
9 part in the study. Researchers will have patients' names, contact numbers, emails and home
10 addresses for the purposes of arranging visits. This information will be stored in accordance
11 with GDPR legislation. Participants are free to withdraw from the study at any time.
12 However, anonymised activity data collected may still be used for data analysis as this is
13 unlinked of any patient identifiable information.
14

15
16 The day-to-day management of the study will be coordinated by the Helix Centre. A study
17 steering committee formed by the intervention team, evaluation team, PPI group, and
18 representatives from the local site will meet at regular intervals throughout the study.
19

20
21 Regular updates about the trial will be made available through social media, blog posts,
22 newsletters and the Helix Centre website (www.helixcentre.com). Trial results will be
23 submitted for publication in journals, presented at national and international stroke meetings
24 and conferences and disseminated amongst stroke communities.
25

26 27 **Trial status**

28
29 The first participant was enrolled on 09.09.2019 and recruitment is expected to complete by
30 the end of March 2020. Enrolment and data collection was continuing as planned at the time
31 of submission of this protocol.
32

33 34 **Author statement**

35
36 AD is grant holder and has project oversight along with DD. GF, EG and LH developed the
37 intervention and conceived of the study. GF, EG, and FJ initiated the study design and ET
38 and ML helped with further refinement. EG and GF are responsible for delivering the
39 intervention and data collection. FJ and ET are responsible for the process evaluation. ML
40 provides statistical expertise in trial design and is conducting the primary statistical analysis.
41 All authors contributed to the refinement of the study protocol and approved the final
42 manuscript.
43
44

45 46 **Funding**

47
48 This study is funded by the NIHR Imperial Biomedical Research Centre (BRC), grant 1215-
49 20013. The views expressed are those of the author(s) and not necessarily those of the
50 NIHR or the Department of Health and Social Care.
51

52
53 All intellectual property associated with the OnTrack system is owned by the Helix Centre
54 which is a collaboration between Imperial College London and the Royal College of Art.
55
56
57
58
59

Data statement

Anonymised data will be made available in a public repository once the data have obtained validation through publication.

Conflict of interests

FJ is the founder of the social enterprise Bridges Self-Management. She has not received any financial support for this work that could have influenced the design.

Acknowledgments

The authors would like to acknowledge the contributions of the members of the PPI group; Jennifer Crow and Sarah Daniels for their input on recruitment and sample size calculations; and Dr Gaby Judah for her help in defining some aspects of the intervention.

Figure legends

Figure 1

Trial diagram

Figure 2

Logic model

Figure 3

Examples of visualisations created using aggregated data captured by OnTrack from healthy beta testers. Data for a minimum of 5 and a maximum of 18 days were aggregated for the period between June and August 2019.

Table 1

Outcome measures

Concept	Assessment	Week of administration
Patient Activation / Engagement	Patient Activation Measure (PAM)	1, 8, 14
Arm impairment	Fugl-Meyer Assessment for upper extremity (FMA-UE)	1, 8, 14
Arm function	Upper-Extremity Motor Activity Log-14 (MAL)	1, 8, 14
Gross level of disability	modified Rankin Scale (mRS)	1, 8, 14
Arm pain	Visual Analogue Scale (VAS)	1, 8, 14
Cognitive impairment	Montreal Cognitive Assessment (MoCA)	1, 8, 14
Arm neglect	Albert's Test (AT)	1, 8, 14
Quality of life	EQ-5D-5L	1, 8, 14
Arm function	Lap-to-Table (LTT)	1, 8, 14
Service experience	Friends and Family Test (FTT)	8, 14
System usability	System Usability Scale (SUS)	14

Table 2

Intervention and Participation schedule

Week	Phase	Description	OnTrack consultation	Assessments
0	Information and consent	NHS therapists screen for eligible patients, provide information and consent participants		Screening, information, and consent
1	Baseline assessment (initial)	Participants complete outcome measures and wear activity trackers (Axivity AX3) on both arms during waking hours (typically 12 hours per day) for one week to gather accelerometer data which is translated into minutes of activity. This data creates a baseline of activity allowing left-to-right arm usage comparison		PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT
2	OnTrack intervention	Participants wear a smartwatch (Apple Watch Series 3 or 4) on their impaired arm only during waking hours (typically 12 hours per day). They will receive real-time feedback on the amount of movement completed (measured in minutes) and daily motivational messages. Participants will receive fortnightly consultations with a researcher to troubleshoot and receive self-management skills training	Onboarding	
3			Check-in & self-management skills training (Problem Solving)	
4				
5			Check-in & self-management skills training (Self-Discovery)	
6			Baseline assessments are repeated during week 8 (halfway)	
7			Check-in & self-management skills training (Goal Setting)	
8			Halfway assessment Check-in & self-management skills training (Goal Setting)	PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT, FFT

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

BMJ Open: first published as 10.1136/bmjopen-2019-034936 on 23 March 2020. Downloaded from <http://bmjopen.bmj.com/> on May 2, 2025 at Department GEZ-LTA

Erasmus Hogeschool

			cont.)	
9				
10			Check-in & self-management skills training (Reflection)	
11				
12			Check-in & self-management skills training (Sign-posting)	
13				
14	Baseline assessment (exit)	Participants complete outcome measures and wear activity trackers (Axivity AX3) on both arms during waking hours (typically 12 hours per day) for one week to gather accelerometer data which is translated into minutes of activity. This data creates a baseline of activity allowing left-to-right arm usage comparison		PAM, FMA-UE, MAL, mRS, VAS, MoCA, AT, EQ-5D-5L, LTT, FFT, SUS
15	Feedback	Independent evaluator leads feedback sessions with participants who have completed the intervention End of participation		Semi-structured interview, online survey (therapists)

References

References

- (1) WHO. *The Atlas of Heart Disease and Stroke*. Available from: https://www.who.int/cardiovascular_diseases/resources/atlas/en/ [Accessed 04 Oct 2019].
- (2) Stroke Association. *State of the Nation: stroke statistics*. Stroke Association. 2018.
- (3) Salter K, Hellings C, Foley N, Teasell R. The Experience of Living with Stroke: A Qualitative Meta-Synthesis. *Journal of Rehabilitation Medicine*. 2008; 40 (8): 595-602. Available from: doi: 10.2340/16501977-0238 Available from: <http://www.ingentaconnect.com/content/mjl/sreh/2008/00000040/00000008/art00003> .
- (4) Care Quality Commission. *Supporting life after stroke: a review of services for people who have had a stroke and their carers*. London: Care Quality Commission. 2011.
- (5) Pollock A, Farmer SE, Brady MC, Langhorne P, Mead GE, Mehrholz J, et al. Interventions for improving upper limb function after stroke. *The Cochrane database of systematic reviews*. 2014; 2014 (11): CD010820. Available from: doi: 10.1002/14651858.CD010820.pub2 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25387001> .
- (6) Pollock A, St George B, Fenton M, Firkins L. Top 10 Research Priorities Relating to Life after Stroke – Consensus from Stroke Survivors, Caregivers, and Health Professionals. *International Journal of Stroke*. 2014; 9 (3): 313-320. Available from: doi: 10.1111/j.1747-4949.2012.00942.x Available from: <https://journals.sagepub.com/doi/full/10.1111/j.1747-4949.2012.00942.x> .
- (7) Kunkel D, Fitton C, Burnett M, Ashburn A. Physical inactivity post-stroke: a 3-year longitudinal study. *Disability & Rehabilitation*. 2015; 37 (4): 304-310. Available from: doi: 10.3109/09638288.2014.918190 Available from: <http://www.tandfonline.com/doi/abs/10.3109/09638288.2014.918190> .
- (8) Bernhardt J, Dewey H, Thrift A, Donnan G. Inactive and Alone: Physical Activity Within the First 14 Days of Acute Stroke Unit Care. *Stroke*. 2004; 35 (4): 1005-1009. Available from: doi: 10.1161/01.STR.0000120727.40792.40 Available from: <http://stroke.ahajournals.org/cgi/content/abstract/35/4/1005> .
- (9) Laver KE, Lange B, George S, Deutsch JE, Saposnik G, Crotty M. Virtual reality for stroke rehabilitation. *The Cochrane database of systematic reviews*. 2017; 11 (11): CD008349. Available from: doi: 10.1002/14651858.CD008349.pub4 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29156493> .
- (10) Loureiro R, Harwin W, Nagai K, Johnson M. Advances in upper limb stroke rehabilitation: a technology push. *Medical & Biological Engineering & Computing*. 2011; 49 (10): 1103-1118. Available from: doi: 10.1007/s11517-011-0797-0 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21773806> .
- (11) Jones F, Gage H, Drummond A, Bhalla A, Grant R, Lennon S, et al. Feasibility study of an integrated stroke self-management programme: a cluster-randomised controlled trial. *BMJ Open*. 2016; 6 (1): e008900. Available from: doi: 10.1136/bmjopen-2015-008900 Available from: <http://dx.doi.org/10.1136/bmjopen-2015-008900> .
- (12) Harwood M, Weatherall M, Talemaitoga A, Barber PA, Gommans J, Taylor W, et al. Taking charge after stroke: promoting self-directed rehabilitation to improve quality of life – a randomized controlled trial. *Clinical Rehabilitation*. 2012; 26 (6): 493-501. Available from: doi: 10.1177/0269215511426017 Available from: <https://journals.sagepub.com/doi/full/10.1177/0269215511426017> .
- (13) Kendall E, Catalano T, Kuipers P, Posner N, Buys N, Charker J. Recovery following stroke: The role of self-management education. *Social Science & Medicine*. 2007; 64 (3): 735-746. Available from: doi: 10.1016/j.socscimed.2006.09.012 Available from: <https://www.sciencedirect.com/science/article/pii/S0277953606004801> .
- (14) Grady PA, Gough LL. Self-Management: A Comprehensive Approach to Management of Chronic Conditions. *American journal of public health*. 2014; 104 (8): e25-e31. Available from:

- doi: 10.2105/AJPH.2014.302041 Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/24922170> .
- (15) Esmonde T, McGinley J, Wittwer J, Goldie P, Martin C. Stroke rehabilitation: patient activity during non-therapy time. *Australian Journal of Physiotherapy*. 1997; 43 (1): 43-51. Available from: doi: 10.1016/S0004-9514(14)60401-3 Available from:
<https://www.sciencedirect.com/science/article/pii/S0004951414604013> .
- (16) Buckland D. Leaving hospital is 'like falling off a cliff' for stroke survivors. *Raconteur*. APRIL 27, 2017 . Available from: <https://www.raconteur.net/healthcare/leaving-hospital-is-like-falling-off-a-cliff-for-stroke-survivors> [Accessed Oct 04 2019].
- (17) McKeivitt C, Fudge N, Redfern J, Sheldenkar A, Crichton S, Rudd AR, et al. Self-Reported Long-Term Needs After Stroke. *Stroke*. 2011; 42 (5): 1398-1403. Available from: doi: 10.1161/STROKEAHA.110.598839 Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/21441153> .
- (18) Woodman P, Riazzi A, Pereira C, Jones F. Social participation post stroke: a meta-ethnographic review of the experiences and views of community-dwelling stroke survivors. *Disability & Rehabilitation*. 2014; 36 (24): 2031-2043. Available from: doi: 10.3109/09638288.2014.887796 Available from:
<http://www.tandfonline.com/doi/abs/10.3109/09638288.2014.887796> .
- (19) Chan A, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, et al. SPIRIT 2013 Statement: Defining Standard Protocol Items for Clinical Trials. *Annals of internal medicine*. 2013; 158 (3): 200-207. Available from: doi: 10.7326/0003-4819-158-3-201302050-00583 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23295957> .
- (20) Billingham SAM, Whitehead AL, Julious SA. An audit of sample sizes for pilot and feasibility trials being undertaken in the United Kingdom registered in the United Kingdom Clinical Research Network database. *BMC medical research methodology*. 2013; 13 (1): 104. Available from: doi: 10.1186/1471-2288-13-104 Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/23961782> .
- (21) Bridges Self-Management Limited. *Bridges Self-Management*. Available from:
<http://www.bridgesselfmanagement.org.uk/> [Accessed 07 Oct 2019].
- (22) Medical Research Institute of New Zealand. *Taking Charge After Stroke (TaCAS)*. Available from: <https://mrinz.ac.nz/programmes/stroke/> [Accessed 07 oct 2019].
- (23) DOCUMAS. *DOCUMAS: efficient health research management*. Available from:
<https://www.documas.eu/> [Accessed 04 Oct 2019].
- (24) Hibbard J, Gilbert H. *Supporting people to manage their health An introduction to patient activation*. London: The Kings Fund. 2014.
- (25) Hibbard JH, Stockard J, Mahoney ER, Tusler M. Development of the Patient Activation Measure (PAM): Conceptualizing and Measuring Activation in Patients and Consumers. *Health Services Research*. 2004; 39 (4p1): 1005-1026. Available from: doi: 10.1111/j.1475-6773.2004.00269.x Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1475-6773.2004.00269.x> .
- (26) Roberts NJ, Kidd L, Dougall N, Patel IS, McNarry S, Nixon C. Measuring patient activation: The utility of the Patient Activation Measure within a UK context—Results from four exemplar studies and potential future applications. *Patient Education and Counseling*. 2016; 99 (10): 1739-1746. Available from: doi: 10.1016/j.pec.2016.05.006 Available from:
<https://www.clinicalkey.es/playcontent/1-s2.0-S0738399116301938> .
- (27) Sahlgrenska Academy Institute of Neuroscience and Physiology, University of Gothenburg. *Fugl-Meyer*. Available from:
https://neurophys.gu.se/english/departments/clinical_neuroscience_and_rehabilitation/rehabilitation-medicine/fugl-meyer [Accessed 04 Oct 2019].
- (28) Pamela W Duncan, Martha Propst, Steven G Nelson. Reliability of the Fugl-Meyer Assessment of Sensorimotor Recovery Following Cerebrovascular Accident. *Physical Therapy*.

- 1983; 63 (10): 1606-1610. Available from: doi: 10.1093/ptj/63.10.1606 Available from: <http://ptjournal.apta.org/content/63/10/1606.abstract> .
- (29) Uswatte G, Taub E, Morris D, Vignolo M, McCulloch K. Reliability and Validity of the Upper-Extremity Motor Activity Log-14 for Measuring Real-World Arm Use. *Stroke*. 2005; 36 (11): 2493-2496. Available from: doi: 10.1161/01.STR.0000185928.90848.2e Available from: <http://stroke.ahajournals.org/cgi/content/abstract/36/11/2493> .
- (30) Quinn TJ, Dawson J, Walters MR, Lees KR. Exploring the Reliability of the Modified Rankin Scale. *Stroke*. 2009; 40 (3): 762-766. Available from: doi: 10.1161/STROKEAHA.108.522516 Available from: <http://stroke.ahajournals.org/cgi/content/abstract/40/3/762> .
- (31) Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison of six methods. *Pain*. 1986; 27 (1): 117-126. Available from: doi: 10.1016/0304-3959(86)90228-9 Available from: <https://www.sciencedirect.com/science/article/pii/0304395986902289> .
- (32) Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment. *Journal of the American Geriatrics Society*. 2005; 53 (4): 695-699. Available from: doi: 10.1111/j.1532-5415.2005.53221.x Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1532-5415.2005.53221.x> .
- (33) Fullerton KJ, McSherry D, Stout RW. Albert's test: a neglected test of perceptual neglect. *Lancet (London, England)*. 1986; 1 (8478): 430. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/2868349> .
- (34) EuroQol - a new facility for the measurement of health-related quality of life. *Health policy*. 1990; 16 (3): 199-208. Available from: doi: 10.1016/0168-8510(90)90421-9 Available from: <https://www.sciencedirect.com/science/article/pii/0168851090904219> .
- (35) NHS England. *The Friends and Family Test*. NHS England. 2014.
- (36) SUS: A 'Quick and Dirty' Usability Scale. *Usability Evaluation In Industry*. CRC Press; 1996. pp. 207-212.
- (37) Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W, et al. Process evaluation of complex interventions: Medical Research Council guidance. *BMJ : British Medical Journal*. 2015; 350 (mar19 6): h1258. Available from: doi: 10.1136/bmj.h1258 Available from: <http://dx.doi.org/10.1136/bmj.h1258> .
- (38) W.K. Kellogg Foundation. *Logic Model Development Guide*. Battle Creek, Michigan: W.K. Kellogg Foundation. 2004.
- (39) The Scottish Government. *Designing and Evaluating Behaviour Change Interventions*. The Scottish Government. 2015.
- (40) Kislov R, Pope C, Martin GP, Wilson PM. Harnessing the power of theorising in implementation science. *Implementation Science*. 2019; 14 (1): 103. Available from: doi: 10.1186/s13012-019-0957-4 Available from: <https://search.proquest.com/docview/2324921962> .
- (41) QSR International Ltd. *NVIVO*. Available from: <https://www.qsrinternational.com/nvivo/nvivo-products> [Accessed 07 Oct 2019].
- (42) Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in Psychology*. 2006; 3 (2): 77-101. Available from: doi: 10.1191/1478088706qp063oa Available from: <http://www.tandfonline.com/doi/abs/10.1191/1478088706qp063oa> .
- (43) National Institute for Health Research. *About INVOLVE*. Available from: <https://www.invo.org.uk/about-involve/> [Accessed 04 Oct 2019].

Figure 1

Trial diagram




Figure 2

Logic model





Figure 3


Examples of visualisations created using aggregated data captured by OnTrack from healthy beta testers between June-August 2019

Active minutes per hour of day (0-23) by user, aggregated over time



Source: Sample OnTrack usage data collected from healthy beta testers, June-August 2019

Active minutes per day by user



Source: Sample OnTrack usage data collected from healthy beta testers, June-August 2019

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies. Erasmushogeschool

Reporting checklist for protocol of a clinical trial.

Based on the SPIRIT guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the SPIRIT reporting guidelines, and cite them as:

Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, Hróbjartsson A, Mann H, Dickersin K, Berlin J, Doré C, Parulekar W, Summerskill W, Groves T, Schulz K, Sox H, Rockhold FW, Rennie D, Moher D. SPIRIT 2013 Statement: Defining standard protocol items for clinical trials. *Ann Intern Med.* 2013;158(3):200-207

		Reporting Item	Page Number
Administrative information			
Title	#1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1
Trial registration	#2a	Trial identifier and registry name. If not yet registered, name of intended registry	2
Trial registration: data set	#2b	All items from the World Health Organization Trial Registration Data Set	n/a
Protocol version	#3	Date and version identifier	11
Funding	#4	Sources and types of financial, material, and other support	12
Roles and responsibilities: contributorship	#5a	Names, affiliations, and roles of protocol contributors	11

1	Roles and	#5b	Name and contact information for the trial sponsor	n/a
2	responsibilities:			
3	sponsor contact			
4	information			
5				
6				
7				
8	Roles and	#5c	Role of study sponsor and funders, if any, in study design;	11
9	responsibilities:		collection, management, analysis, and interpretation of data;	
10	sponsor and funder		writing of the report; and the decision to submit the report for	
11			publication, including whether they will have ultimate authority	
12			over any of these activities	
13				
14				
15				
16	Roles and	#5d	Composition, roles, and responsibilities of the coordinating centre,	n/a
17	responsibilities:		steering committee, endpoint adjudication committee, data	
18	committees		management team, and other individuals or groups overseeing the	
19			trial, if applicable (see Item 21a for data monitoring committee)	
20				
21				
22				
23	Introduction			
24				
25	Background and	#6a	Description of research question and justification for undertaking	3
26	rationale		the trial, including summary of relevant studies (published and	
27			unpublished) examining benefits and harms for each intervention	
28				
29				
30	Background and	#6b	Explanation for choice of comparators	n/a
31	rationale: choice of			
32	comparators			
33				
34				
35				
36	Objectives	#7	Specific objectives or hypotheses	4
37				
38	Trial design	#8	Description of trial design including type of trial (eg, parallel	5
39			group, crossover, factorial, single group), allocation ratio, and	
40			framework (eg, superiority, equivalence, non-inferiority,	
41			exploratory)	
42				
43				
44				
45	Methods:			
46	Participants,			
47	interventions, and			
48	outcomes			
49				
50				
51				
52	Study setting	#9	Description of study settings (eg, community clinic, academic	5
53			hospital) and list of countries where data will be collected.	
54			Reference to where list of study sites can be obtained	
55				
56				
57	Eligibility criteria	#10	Inclusion and exclusion criteria for participants. If applicable,	5
58			eligibility criteria for study centres and individuals who will	
59				
60				

		perform the interventions (eg, surgeons, psychotherapists)	
Interventions: description	#11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	6
Interventions: modifications	#11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving / worsening disease)	n/a
Interventions: adherence	#11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return; laboratory tests)	n/a
Interventions: concomitant care	#11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	n/a
Outcomes	#12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	7
Participant timeline	#13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	Table 2
Sample size	#14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	6
Recruitment	#15	Strategies for achieving adequate participant enrolment to reach target sample size	6
Methods: Assignment of interventions (for controlled trials)			
Allocation: sequence generation	#16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	n/a

1	Allocation concealment	#16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	n/a
2	mechanism			
3				
4				
5				
6				
7				
8	Allocation:	#16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	n/a
9	implementation			
10				
11				
12	Blinding (masking)	#17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	n/a
13				
14				
15				
16				
17	Blinding (masking):	#17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	n/a
18	emergency unblinding			
19				
20				
21				
22	Methods: Data			
23	collection,			
24	management, and			
25	analysis			
26				
27				
28				
29	Data collection plan	#18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	9
30				
31				
32				
33				
34				
35				
36				
37				
38				
39	Data collection plan:	#18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	n/a
40	retention			
41				
42				
43				
44	Data management	#19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	9
45				
46				
47				
48				
49				
50				
51	Statistics: outcomes	#20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	9
52				
53				
54				
55				
56	Statistics: additional	#20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	9
57	analyses			
58				
59				
60				

1	Statistics: analysis	#20c	Definition of analysis population relating to protocol non-	n/a
2	population and missing		adherence (eg, as randomised analysis), and any statistical methods	
3	data		to handle missing data (eg, multiple imputation)	
4				
5				
6	Methods: Monitoring			
7				
8	Data monitoring:	#21a	Composition of data monitoring committee (DMC); summary of its	n/a
9	formal committee		role and reporting structure; statement of whether it is independent	
10			from the sponsor and competing interests; and reference to where	
11			further details about its charter can be found, if not in the protocol.	
12			Alternatively, an explanation of why a DMC is not needed	
13				
14	Data monitoring:	#21b	Description of any interim analyses and stopping guidelines,	n/a
15	interim analysis		including who will have access to these interim results and make	
16			the final decision to terminate the trial	
17				
18	Harms	#22	Plans for collecting, assessing, reporting, and managing solicited	n/a
19			and spontaneously reported adverse events and other unintended	
20			effects of trial interventions or trial conduct	
21				
22	Auditing	#23	Frequency and procedures for auditing trial conduct, if any, and	n/a
23			whether the process will be independent from investigators and the	
24			sponsor	
25				
26				
27				
28	Ethics and			
29	dissemination			
30				
31	Research ethics	#24	Plans for seeking research ethics committee / institutional review	10
32	approval		board (REC / IRB) approval	
33				
34	Protocol amendments	#25	Plans for communicating important protocol modifications (eg,	10
35			changes to eligibility criteria, outcomes, analyses) to relevant	
36			parties (eg, investigators, REC / IRBs, trial participants, trial	
37			registries, journals, regulators)	
38				
39	Consent or assent	#26a	Who will obtain informed consent or assent from potential trial	6
40			participants or authorised surrogates, and how (see Item 32)	
41				
42	Consent or assent:	#26b	Additional consent provisions for collection and use of participant	n/a
43	ancillary studies		data and biological specimens in ancillary studies, if applicable	
44				
45	Confidentiality	#27	How personal information about potential and enrolled participants	10
46			will be collected, shared, and maintained in order to protect	
47			confidentiality before, during, and after the trial	
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				

1	Declaration of interests	#28	Financial and other competing interests for principal investigators for the overall trial and each study site	12
2				
3				
4	Data access	#29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	10
5				
6				
7				
8				
9				
10	Ancillary and post trial care	#30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	n/a
11				
12				
13				
14	Dissemination policy: trial results	#31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	10
15				
16				
17				
18				
19				
20				
21	Dissemination policy: authorship	#31b	Authorship eligibility guidelines and any intended use of professional writers	n/a
22				
23				
24	Dissemination policy: reproducible research	#31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	n/a
25				
26				
27				
28	Appendices			
29				
30				
31	Informed consent materials	#32	Model consent form and other related documentation given to participants and authorised surrogates	n/a
32				
33				
34				
35	Biological specimens	#33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	n/a
36				
37				
38				
39				

The SPIRIT checklist is distributed under the terms of the Creative Commons Attribution License CC-BY-ND 3.0. This checklist was completed on 07. October 2019 using <https://www.goodreports.org/>, a tool made by the [EQUATOR Network](#) in collaboration with [Penelope.ai](#)