



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Reinforcing Informed Medication prescription for low back pain in the Emergency department (RIME): a controlled interrupted time-series implementation study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-082668
Article Type:	Protocol
Date Submitted by the Author:	30-Nov-2023
Complete List of Authors:	<p>O'Leary, Shaun; University of Queensland; Royal Brisbane and Women's Hospital, Physiotherapy Department</p> <p>Heine, Janelle; Royal Brisbane and Women's Hospital Emergency and Trauma Centre; Royal Brisbane and Women's Hospital, Physiotherapy Department</p> <p>Warren, Jacelle; Metro North Hospital and Health Service, Jamieson Trauma Institute, Royal Brisbane & Women's Hospital (RBWH)</p> <p>Smyth, Tanya; Metro North Hospital and Health Service, Jamieson Trauma Institute, Royal Brisbane & Women's Hospital (RBWH)</p> <p>Ballard, Emma ; QIMR Berghofer Medical Research Institute, Mitchell, Gary; Royal Brisbane and Women's Hospital Emergency and Trauma Centre</p> <p>See, William; The Prince Charles Hospital ED</p> <p>Barlas, Panos; Metro North Hospital and Health Service, Jamieson Trauma Institute, Royal Brisbane & Women's Hospital (RBWH)</p> <p>Machado, Gustavo; University of Sydney - Camperdown and Darlington Campus, School of Public Health</p> <p>Cottrell, Michelle; Royal Brisbane and Women's Hospital, Physiotherapy Department</p> <p>Comans, Tracy; 8. Centre for Health Services Research, The University of Queensland</p> <p>Foster, Nadine; The University of Queensland, Surgical Treatment and Rehabilitation Service</p>
Keywords:	<p>Back pain < ORTHOPAEDIC & TRAUMA SURGERY, ACCIDENT & EMERGENCY MEDICINE, CLINICAL PHARMACOLOGY, Implementation Science, Protocols & guidelines < HEALTH SERVICES ADMINISTRATION & MANAGEMENT</p>

SCHOLARONE™
Manuscripts

Reinforcing Informed Medication prescription for low back pain in the Emergency department (RIME): a controlled interrupted time-series implementation study

Authors: Shaun O’Leary,^{1,2} Janelle Heine,^{1,3} Jacelle Warren,⁴ Tanya Smyth,⁴ Emma Ballard,⁵ Gary Mitchell,³ William See⁶, Panagiotis Barlas,⁴ Gustavo C Machado,⁷ Michelle Cottrell,¹ Tracy Comans,⁸ Nadine E Foster^{2,9}

Institutional addresses:

1. Physiotherapy Department, Royal Brisbane and Women’s Hospital, Metro North Health
2. School of Health and Rehabilitation Sciences, The University of Queensland
3. Emergency and Trauma Centre, Royal Brisbane and Women’s Hospital, Metro North Health
4. Jamieson Trauma Institute, Royal Brisbane and Women’s Hospital, Metro North Health
5. Statistics Unit, QIMR Berghofer Medical Research Institute, Herston
6. Emergency Department, The Prince Charles Hospital, Metro North Health
7. Institute for Musculoskeletal Health, University of Sydney and Sydney Local Health District
8. Centre for Health Services Research, The University of Queensland
9. Surgical Treatment and Rehabilitation Service (STARS) Education and Research Alliance, The University of Queensland and Metro North Health

Corresponding author: Associate Professor Shaun O’Leary Postal Address: Physiotherapy Department, School of Health and Rehabilitation Sciences, University of Queensland, St Lucia, Queensland, 4072, Australia. Institute Email Address: s.oleary@uq.edu.au
Phone : +61 4 07746299.

Word Count: 3999

ABSTRACT

Introduction: Management guidelines for low back pain (LBP) recommend exclusion of serious pathology, followed by simple analgesics, superficial heat therapy, early mobilisation, and patient education. An audit in a large metropolitan hospital Emergency Department (ED) revealed high rates of non-recommended medication prescription for LBP (65% of patients prescribed opioids, 17% prescribed benzodiazepines), high inpatient admission rates (20% of ED LBP patients), delayed patient mobilisation (on average 6 hours), and inadequate patient education (48% of patients). This study aims to improve medication prescription for LBP in this ED by implementing an intervention shown previously to improve guideline-based management of LBP in other Australian EDs.

Methods and Analysis: A controlled interrupted time series study will evaluate the intervention in the ED before and after implementation, additionally comparing findings with another ED in the same health service. The multicomponent implementation strategy uses a formalised clinical flow chart to support clinical decision-making and aims to change clinician behaviour, through clinician education, provision of alternative treatments, educational resources, audit and feedback, supported by implementation champions. The primary outcome is the percentage of LBP patients prescribed non-recommended medications (opioids, benzodiazepines, gabapentinoids), assessed via routinely collected ED data. Anticipated sample size is 2000 patients (n=1000 intervention, n=1000 control) based on average monthly admissions of LBP presentations in the EDs. Secondary outcomes include inpatient admission rate, time to mobilisation, provision of patient education, imaging requests, re-presentation to the ED within 6 months, and healthcare costs. In nested qualitative research, we will understand ED clinicians' perceptions of the implementation and determine how benefits can be sustained over time.

ARTICLE SUMMARY

Strengths and limitations of this study

- Controlled interrupted time series design will test the effects of the intervention on relevant outcomes while observing and gathering information on implementation in a real-world ED setting.
- Analysis plan will evaluate outcomes over a 6-month period before and after the implementation of the intervention, and compare the change in outcomes between the intervention ED and control ED.
- Process evaluation will explore components of the implementation that clinicians found useful, as well as perceived barriers to adoption and sustainability of improved LBP management in the ED.
- Both participating EDs are within the same health service and therefore results may lack generalisability to other settings.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Low back pain (LBP) is the fifth most common presentation to Australian Emergency Departments (EDs).[1] Management guidelines for LBP recommend an initial exclusion of serious pathology (e.g. spinal fracture, infection, cancer), followed by the use of simple analgesics, superficial heat therapy, early mobilisation to improve function, and the provision of patient education to enable self-management.[2] Research to date has evidenced inconsistent use of best practice guidelines for LBP in Australian EDs.[3]

In line with previous research a recent audit of ED routinely collected data at a large metropolitan tertiary hospital in Queensland, Australia, identified a high rate of medication prescription for LBP, particularly opioids (prescribed for 65% of LBP patients) but also muscle relaxants (benzodiazepine, 17%). These are not recommended first line medications for LBP and have known side-effects including dependence, poisoning and death.[4] Other concerning audit findings were that 20% of patients with LBP were admitted as an inpatient (to either the Short Stay Unit or hospital wards), had long waits until they received support to mobilise (6 hours on average until they were supported to walk), and may not have received patient education (48% of patients' records had no documented evidence of education/advice).[5] In response, an intervention was sought to promote guideline based management of LBP in this metropolitan hospital ED to improve patient care, particularly with regard to medication prescription.

A recent randomised trial conducted in other Australian EDs was identified that showed improvement in the use of LBP best practice guidelines by clinicians.[6] The Sydney Health Partners Emergency Department (SHaPED) trial successfully reduced opioid prescription by over 12% (50.5% in intervention EDs v 62.8% in control) (OR=0.57, 95% CI 0.38-0.85), with no adverse effects on patients' pain scores or satisfaction with care.[6] Improvements were also observed regarding clinicians' beliefs and knowledge regarding the management of LBP.[6]

The SHaPED trial specifically intervened to change ED clinician behaviour regarding the use of LBP guidelines.[6] The SHaPED trial used a multicomponent implementation strategy that was informed not only by a review of practice, but also included locally driven strategies that clinicians identified as potential facilitators and barriers to the uptake of LBP guidelines in the participating Sydney EDs.[7]

This protocol paper describes the Reinforcing Informed Medication prescription for low back pain in the Emergency department (RIME) study. The aim of the RIME study is to improve the management of LBP in our Queensland-based ED, by implementing and evaluating the intervention shown to be effective in the SHaPED trial in Sydney EDs. The primary and secondary objectives of the study are as follows.

Primary objective

Objective 1: To evaluate the impact of the intervention on the percentage of patients with LBP who are prescribed non-recommended medications (i.e. opioids/ benzodiazepines/ gabapentinoids) in the ED (primary outcome).

Secondary objectives

Objective 2: Evaluate the impact of the intervention on the percentage of patients with LBP who are admitted to hospital, or who receive support to mobilise (including time to support to walk), appropriate advice/education, lumbar spine imaging, and who re-present to ED within 6 months (secondary outcomes).

Process evaluation

Objective 3: Undertake a cost analysis of implementing the intervention from a hospital resources perspective.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Objective 4: Understand the perspectives (e.g. acceptability, adoption, appropriateness, feasibility, sustainability) of ED clinicians towards the intervention, the implementation strategy, and sustainability of changes in LBP management in the ED.

Objective 5: Determine the fidelity to which each component of the implementation strategy was delivered, including the percentage of appropriate clinician encounters with the RIME training team during implementation.

METHODS AND ANALYSIS

Study design

RIME is a controlled Interrupted Time Series (ITS) study design evaluating outcomes across study phases with nested process evaluation. This design permits evaluation of changes in outcomes both pre- and post-implementation in the intervention ED, as well as intervention and control ED comparisons to exclude confounders due to co-interventions or other events occurring during the study timeline.[8] The study is a prospective effectiveness-implementation hybrid model (Type 1), testing intervention effects on relevant measurable outcomes (Objectives 1-3) while observing and gathering information on implementation (Objectives 4-5) in a real-world setting.[9]

Scheduled activity over the three study phases are shown in Table 1 and include; Phase 1/ Pre-implementation usual care - during which primary and secondary clinical outcome data will be collected over a 6-month period for adult patients (aged 18 years and over) presenting with LBP to either the intervention or control EDs; Phase 2/ Implementation - during which the intervention will be implemented with clinicians over a 12-week period at the intervention ED only; and Phase 3/ Post-implementation care – which repeats the identical data collection procedure of Phase 1 over a 6-month period at both intervention and control EDs.

Table 1 Schedule of procedures during the RIME study with week (W) 1-24 representing the pre-implementation phase (Pre), week 25-36 the implementation phase, and week 37-60 the post-implementation phase (Post). Implementation weeks and implementation strategies are shaded grey. Black arrows denote continuous periods of outcome measurements recorded fortnightly, thus initial recordings are W2 and W38 of the pre-and post-implementation phases, respectively.

	Pre			Implementation							Post					
Procedures/Measurements	W2	W24	W25	W26	W27	W28	W29	W30	W31	W32	W33	W34	W35	W36	W38	W60
Measurement of Primary and Secondary Outcomes (fortnightly)	✓	➡		✓		✓		✓		✓		✓		✓	✓	➡
Study Information Sheets provided			✓													
Educational seminars delivered			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓		
Educational materials provided			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓		
Provision of alternative LBP treatment options			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓		
Fast-track referral to outpatient services			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓		
Audit and feedback						✓				✓				✓		
Implementation Champion actioned			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Cost data collection

√ √ √ √ √ √ √ √ √ √ √ √ √ √ √

Audit & Observation
Measures Recorded

√ √ √ √

Qualitative Interviews
commence

√

For peer review only

36/bmjopen-2023-082668 on 12 March 2024. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2025 at Department GEZ-LTA
Erasmus Hogeschool .
d by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Patient and public involvement

No patients were involved in the study design; as this study focuses only on the use of routinely collected data (ie. no data are being collected directly from ED patients).

Study setting

The study will be conducted in two large metropolitan tertiary hospital EDs in Queensland, Australia. One ED will serve as the intervention site and the other the control site. Both EDs are governed by the same Hospital and Health Service, and the catchment populations of both are almost identical, projected to be approximately 368,000 persons by 2026.[10] The median age of residents in the intervention ED catchment area is slightly younger (approximately 34 years) than for the control ED (approximately 37 years),[11] however both catchment areas have over 1,000 LBP patients presenting to their ED each year (January to December 2021 data; intervention ED 1,046 patients, control ED 1,485 patients), and each hospital has a similar skill-mix in their ED clinical teams. These clinical teams include senior and junior ED doctors and nurses, as well as Emergency Physiotherapy Practitioners (EPPs). The EPPs work relatively independently within the ED and are usually responsible for the assessment, investigation, and management of LBP patients.

Participants

There are three participant groups in this study.

Patients with LBP: Data from adult patients (18+ years) presenting to either the intervention or control ED with LBP will be included in the study. A waiver of consent has been granted for the use of the anonymous (de-identified) ED patient data. Only routine ED service data will be

required for the study, there is no foreseeable risk or additional burden placed on patients, and it is impractical to gain individual consent for the use of these routine service data. It was also not deemed necessary to burden patients with patient-reported outcome data collection, given the previous SHaPED trial already showed the intervention to be effective in reducing opioid prescription without compromising patient reported pain or satisfaction.[6]

Potentially eligible LBP patient cases will be identified at the two sites using the Electronic Medical Record (EMR) system - Emergency Department Information System (EDIS) - with the following parameters:

- Seen in a low-acuity area of the ED, AND
- Presenting complaint related to ‘Back Pain’, ‘Altered Sensation’, ‘Mechanical Strain’, ‘Requesting Investigation’ or ‘Requesting Medication/Certificate/Results’, OR
- Primary diagnosis code related to LBP (coding system ICD-10-AM)

Exclusions are patient cases with spinal fracture, infections, cauda equina syndrome, major trauma, malignancy, and other non-spinal causes of LBP. These exclusions will be identified through screening EDIS information (see Supplementary Table 1 for a full list of EDIS extraction parameters). Potentially eligible LBP patients will then be manually screened by site-based study staff who will review the presenting problem free text in the ED data to further refine the study sample to those with LBP who do not present with one of the exclusions. Reasons for exclusion during manual review will be collected and summarised.

ED Clinicians: All ED clinicians at the intervention ED will be informed about the intervention as a service improvement activity but their level of involvement will be at their discretion. Individual ED clinician consent to participate will not be sought. This approach ensures these ED clinicians do not feel pressured to undertake activities or practices they do not wish to. The

research team will attempt to facilitate uptake by offering all ED clinicians all components of the implementation strategy. Only the intervention ED clinicians will be given a RIME study information sheet to minimise the risk of potential contamination at the control ED.

ED Clinicians who take part in the nested interviews: Following the implementation phase, up to 20 ED clinicians will be invited to participate in a qualitative semi-structured interview exploring their views of the intervention, its implementation and potential sustainability of improvements in practice. This sample size will enable representation from the three ED clinician disciplines (ie. EPPs, nurses, medical) and is similar to our earlier ED clinician qualitative study.[5] Consent from these participants will be implied by their voluntary participation in the interviews.

Sample Size

Based on recommendations[12, 13] and given the number of observations within each time-period and effect on statistical power, the study will use 12 time-periods spaced 2 weeks apart (6 months total) for all outcomes in both the pre- and post-implementation phases (Phases 1 and 3). Simulations based on the SHaPED trial findings[6] using SAS v9.4 suggests the power to detect at least a 12% reduction in non-recommended medication prescription (the primary outcome) at the intervention ED is between 70% and 80%.[14] Anticipating 500 LBP ED presentations in 6 months at the intervention ED allows for approximately 45 patients in each 2-week time-period before and after implementation. The total anticipated sample size is 2000 (500 in each of the two EDs, in each of the pre- and post-implementation periods); (n=1000 intervention, n=1000 control).

Intervention and Implementation Plan

The intervention addresses ED clinician behaviour to improve their use of best-practice LBP management guidelines, specifically the NSW Agency for Clinical Innovation (ACI) model of care for acute LBP[2] and the more recent Australian LBP Clinical Care Standard.[15]

To best inform the implementation approach for the local ED context, five focus groups with 18 ED clinicians (medical, nursing, EPPs) were previously conducted.[5] These interviews gained the local perspective on current practice, LBP guidelines, and barriers and enablers potentially impacting the uptake of guidelines for LBP management at the intervention ED.[5] Findings indicated that adherence to guidelines is primarily driven by (i) clinician beliefs and behaviours; (ii) patient expectations and behaviours; and (iii) workflow processes. Clinicians' beliefs around role scope and accountability were perceived to serve as barriers to providing recommended treatments such as prescription of simple analgesics and early mobilisation. Patients' expectations of care (e.g., specifically requesting stronger analgesics) were perceived to influence clinical decisions regarding medication prescription. Limited access to out-of-hours physiotherapy, along with National Emergency Access Target pressures, were also felt to influence clinical decisions, particularly inpatient admissions to the Short Stay Unit (SSU).[5]

Through methodological triangulation (audit, survey, focus groups), the focus group data provided a deeper understanding of the previous ED audit results.[5] Strategies suggested to improve best practice within the intervention ED context included materials such as a formalised clinical flow chart (to support clinical decision-making that aligns with best practice), targeted education to medical and nursing staff around safe mobilisation practices and best practice management of LBP patients, and patient education resources to better support patient self-management. Additionally, it was identified that the multifaceted intervention would require organisational support to embed the strategies within clinical

management systems including the EMR, to facilitate clinicians to adopt best practice routinely.[5]

Implementation of the intervention will be underpinned by the Knowledge-to-Action framework (as per the SHaPED trial),[16] incorporating evidence-based implementation strategies specifically targeting the behaviour of ED clinicians at the intervention ED. The Capability-Opportunity-Motivation=Behaviour (COM-B) behaviour change theoretical framework[17] has been used to shape the implementation plans to support behaviour change in ED clinicians to enhance capability, opportunity, and motivation to improve uptake of guidelines when managing LBP patients in the ED.

Figure 1 depicts the multicomponent implementation strategy, including:

1. Educational Seminars: These will comprise structured best practice updates from experienced ED clinicians (i.e. Emergency Physiotherapy Practitioners, Emergency Medical Consultants) that focus on knowledge and skills for assessing, managing, educating, and referring patients in line with LBP guidelines.[2] Preparation of these best practice updates will be assisted by the investigator team. These best practice sessions will be offered on numerous occasions throughout the 12-week implementation period in protected ED teaching time. All ED clinicians will be invited to participate and clinician participation in the education sessions will be tracked through a logbook, with reminders sent and personal communication from the study team and ED clinical leads where needed.

2. Educational materials: ED clinicians will receive a hard copy of the LBP guidelines and LBP Clinical Standards,[2][15] a link to an established, contemporary evidence-based patient-facing website (<https://mybackpain.org.au/>), and the formalised clinical flow chart to support clinical decision-making such as the appropriate use of analgesic medicines. Posters highlighting key messages about benefits and harms of opioid medicines, lumbar imaging, and

inpatient admission will be displayed throughout the intervention ED. Anonymised patient cases from study phase 1 will be discussed, illustrating examples of non-recommended practice and best practice. Patient educational materials (based on the LBP guidelines) and scripts to guide conversations with patients will be provided so that clinicians can use these with patients.

3. Provision of alternative treatment options for LBP: Non-opioid pain medicines will be made more easily accessible to clinicians as an evidence-based alternative to opioid medicines or muscle relaxants. Heat wraps (used as a non-pharmacological modality for pain relief) will also be made available to clinicians with encouragement to use these as alternatives to non-recommended pharmacological treatments.

4. Fast-track referral to outpatient services: Clinicians will be educated on available referral pathways, for whom these are appropriate, and how to collaboratively facilitate the referral process with the patient. Referral pathway options include; private physiotherapy services within the primary care setting, onsite public physiotherapy outpatient services, or onsite advanced-practice musculoskeletal physiotherapy screening services.

5. Audit and feedback: During the implementation phase, routinely collected ED data will be audited and clinicians provided with structured ED-level- and discipline-level audit and feedback reports, including medication prescription data. These reports are planned to be provided at 4, 8 and 12 weeks after the start of the implementation phase.

6. Support from an ED 'Implementation Champion': A key feature of the implementation is the inclusion of a dedicated 'Implementation Champion'. An experienced ED clinician (investigator Heine) will undertake this role, as champions are considered vital to successful implementation and change in their own sphere of influence, particularly when intrinsically motivated and enthusiastic about the practices they promote.[18] The implementation champion will support reinforcement of implementation aims with ED staff, provide personal feedback sessions, and offer one-to-one or small group discussions with ED staff as needed.

Insert Figure 1 Here

Study Outcomes

Study outcomes include primary and secondary clinical outcomes, as well as a nested process evaluation.

Data related to the primary and secondary clinical outcomes, in addition to key demographic characteristics and ED management data, will be obtained using either routine electronic data from EDIS, or via manual extraction from the medical records. Data validation rules will be implemented to reduce data transcription errors and improve data quality. Following completion of the data extraction, patient identifying information will be removed and replaced with an anonymised ID code.

Data obtained from EDIS include patient presentation date/time, demographic characteristics (age, gender, postcode) and ED management details (triage category, arrival mode [ambulance, walk in etc], primary clinician, consultations performed, ED discharge destination, time to treatment etc). Data obtained via manual extraction from the medical record include previous history of LBP, pain severity, ED treatment details (attempted mobilisation, provision of education/advice, heat pack used, imaging requested etc), and medications prescribed while in the ED and on discharge. Data relating to re-presentations within 6 months will also be obtained from EDIS.

Primary Clinical Outcome (Objective 1)

- Percentage (%) of patients presenting to the ED with LBP prescribed non-recommended medications in the ED. Non-recommended medications are grouped under the Anatomical

Therapeutic Chemical (ATC) classification[19] of N02A (Opioids), N05B (Benzodiazepines) or N02BF (Gabapentinoids).[6, 7]

Secondary Clinical Outcomes (Objective 2)

- Percentage (%) of patients presenting to the ED with LBP prescribed analgesic medications in the following ATC categories: N02BE01 (Paracetamol) or M01A (Anti-inflammatory and Antirheumatic Products, non-steroidal).
- Mean/Median time to mobilisation (time to be supported to walk).
- Percentage (%) of patients receiving education about LBP care.
- Percentage (%) of patients receiving lumbar spine imaging.
- Percentage (%) of patients admitted to hospital or short stay.
- Percentage (%) of patients re-presenting to the same ED facility within 6 months.

Process Evaluation

The process evaluation of the implementation is underpinned by the framework developed by Proctor et al.[20] This framework consists of eight distinct implementation outcomes which serve to understand the implementation process and determine implementation success. Implementation outcomes will be assessed by the approaches shown in Table 2.

Cost Evaluation (Objective 3)

Costs will be estimated based on project staff time to conduct the implementation components/activities, clinical staff time to attend the educational seminars and cost of developing materials. Research specific costs (e.g. semi-structured interviews) will be excluded from this analysis, however, will be reported separately. Direct health system costs

will be estimated using actual cost data obtained from each hospital's clinical costings department for each eligible ED patient presentation and any related inpatient episodes of care. The cost categories collected include medical, nursing, allied health, pathology, imaging, consumables, hotel and medications. These data will be used to evaluate and compare the cost of treating LBP patients pre- to post-implementation at the intervention ED, and between intervention and control EDs.

Semi-structured interviews (Objective 4)

The semi-structured interviews will incorporate questions constructed to address Proctor et al.'s [20] implementation outcome domains of: acceptability, adoption, appropriateness, feasibility, and sustainability (Table 2).

Audit & Observation (Objective 5)

Both penetration and fidelity outcomes, as defined by Proctor et al.[20] (Table 2) of each proposed implementation strategy component will be determined by calculating the percentage (%) of encounters where the strategy was actually used, as well as the extent to which the strategy was delivered as intended (%). As feasible, regular monitoring throughout the 12-week implementation period (Phase 2) will occur through both chart audits (reviewed treatment notes) and peer observation led by the implementation champion. Engagement with the implementation components will be additionally captured by recording attendance to educational sessions and use of resources (recorded within a logbook at a department and discipline level, not individual clinician level) as accurately as possible.

Table 2 Implementation outcomes based on Proctor et al.[20] and adapted for RIME. Time points of measurement include Phase 2 Implementation or Phase 3 Post-implementation

Outcome	Definition from Proctor et al.	Definition as applied to RIME	Time point of Measurement	Measurement Source
Acceptability	Perception among implementation stakeholders that a given treatment, service, practice, or innovation is agreeable, palatable, or satisfactory.	ED clinicians’ perceptions of the RIME intervention components, and implementation strategies, as agreeable, palatable, or satisfactory.	Phase 3 Post-implementation	Clinician interviews
Adoption	The intention, initial decision, or action to try, or employ, an innovation or evidence-based practice. Adoption also may be referred to as ‘uptake’.	The intention and behavioural uptake of the RIME intervention components by ED clinicians to facilitate adherence to LBP Clinical Standards.	Phase 3 Post-implementation	Clinician Interviews
Appropriateness	The perceived fit, relevance, or compatibility of the innovation or evidence-based practice for a given practice setting, provider, or consumer; and/or perceived fit of the innovation to address a particular issue or problem.	The extent to which ED clinicians perceive the fit or relevance of the RIME intervention components within their clinical setting to facilitate adherence to LBP Clinical Standards.	Phase 3 Post-implementation	Clinician Interviews
Feasibility	The extent to which a new treatment, or an innovation, can be successfully used or carried out within a given agency or setting.	The extent to which implementation of the RIME intervention, and implementation strategies, is considered feasible for the clinical setting under study, allowing for individual tailoring given the differing needs of clinicians.	Phase 3 Post-implementation	Clinician Interviews

Fidelity	The degree to which an intervention was implemented as it was prescribed in the original protocol or as it was intended by the program developers.	The degree to which each RIME intervention component was implemented as described within the original protocol.	Phase 2 Implementation	Audit & Observation
Implementation Cost	The cost impact of an implementation effort. The cost of implementing an intervention includes the costs of the intervention, the implementation strategy used, and the service delivery.	The costs involved in the development and implementation of the RIME intervention within the ED setting under study.	Phase 2 Implementation	Health Economic Evaluation
Penetration	The integration of a practice within a service setting and its subsystems. Can also be referred to as 'reach' (using the RE-AIM framework).	The extent to which each RIME intervention component was accessed/used by the ED clinicians.	Phase 2 Implementation	Audit & Observation
Sustainability	The extent to which a newly implemented treatment is maintained or institutionalised within a service setting's ongoing, stable operations.	The extent to which the RIME intervention components could be embedded as standard practice within the ED setting, following the completion of the RIME study.	Phase 3 Post implementation	Clinician Interviews

Data Analysis

Primary and Secondary Clinical Outcomes: Two analytic approaches will be considered to assess primary and secondary outcomes: 1) a controlled interrupted time series (ITS) analysis, where data are aggregated on a fortnightly basis, and 2) a segmented regression analysis on all data. The change in trend (slope) between pre- and post-implementation periods and the change in level between the end of the pre-implementation period and the beginning of the post-implementation periods and between the beginning of the pre-implementation period and at the end of the post-implementation period will be described for separate models for each hospital and models including hospital ED as a fixed effect. Additionally for models including hospital ED as a fixed effect, the difference between the two EDs for the change in slope and change in level will be described. All estimates will be reported with 95% confidence intervals (CI). Data will be presented visually, overlaid with the regression line for pre- and post-implementation periods for each outcome. Assumptions relating to linearity, autocorrelation, stationarity and heteroscedasticity will be examined, and consideration given to the influence of outliers and the normality of residuals in developing a robust model.

Patient characteristics (e.g age, sex, Socio-Economic Indexes for Area (SEIFA), ED arrival mode, severity (i.e. Australian Triage Scale (ATS) 1 – 5), type of treating clinician and day/time of presentation will be compared between EDs during the pre-implementation period and between pre and post-implementation periods within EDs. Associations between these characteristics and outcomes will also be examined. These comparisons will be done using the Pearson Chi-squared test or Fisher’s exact test for categorical data and Student t-test or Mann-Whitney U test for continuous data. Categorical data will be presented as frequency and percentage and continuous data as mean and standard deviation or median and interquartile range if not normally distributed. The distribution of these characteristics overtime will be

examined visually and reported. If these characteristics are found to be associated with the intervention or they are associated with the outcome and change over time in both EDs, these variables will be adjusted for in the segmented regression. Stepwise backward elimination will be used to select the most parsimonious model.

Planned exploratory analyses include; a three period (pre, implementation, post) analysis; a two period analysis where data from the implementation period is combined with the post implementation period. Planned sensitivity analyses include; analysis of index presentation only, multilevel segmented regression model to account for clustering of observations within patients and observations within clinician groups, and controlling for differences in the patient base by including the value of the outcome denominator in the ITS model. The planned subgroup analysis is to understand the ED medication usage in patients who are transported to hospital via ambulance (given ambulance staff may provide medications to patients with LBP before their arrival at the ED).

Process Evaluation:

Cost Evaluation: To assess the value for money of the intervention, total healthcare costs will be compared between pre and post-implementation phases, and between intervention and control EDs. The costs will be compared using models appropriate for clustered data and the non-normal nature of the cost and variable (e.g., generalised linear model with robust standard errors). Confounders will be included in the model if assessed as necessary in the statistical analysis. A final parsimonious model will be selected following data examination and measures of model fit. Following the base case analysis, sub-group analysis will be conducted of the main categories of cost (e.g. medication only) to assess where differences have accrued.

Semi-structured interviews: Interviews will be transcribed and analysed using NVivo software. The Framework Method, as described by Gale et al.[21] will be employed to deductively analyse interview data based upon Proctor et al.’s pre-defined implementation outcomes[20] introduced in Table 2. In addition, ‘open-coding’ will be performed for information that does not readily adhere to the pre-defined framework, where themes/concepts will be generated using an inductive thematic analysis approach.

Audit & Observation: Simple descriptive statistics will be employed to measure fidelity and penetration of each intervention component. Fidelity will be measured at regular intervals (approximately every 2 weeks) and descriptive analysis undertaken to explore any changes in fidelity over the implementation period.

ETHICS AND DISSEMINATION

No risks to patients within this study are foreseeable given its intention to improve the management of LBP in the ED, based on best-practice guidelines. The study is aligned with concerted efforts nationally and globally to reduce the use of opioids which continues to be a major health concern.[4, 22-25] It is hypothesised the intervention will reduce the rate of non-recommended medication prescriptions for patients with LBP in the intervention ED (Objective 1); reduce inpatient admissions, improve early mobilisation, improve the provision of patient education, and reduce re-presentation to the intervention ED (Objective 2); and improve the cost of care from the hospital perspective (Objective 3), all through behavioural change in the clinical practice of ED clinicians (Objective 4). Furthermore, it will provide patients with the resources to better self-manage LBP, reducing recurrence and healthcare dependency.

It is planned to disseminate findings widely including engagement with relevant stakeholders regarding implementation of the intervention at respective sites locally (i.e., the control ED),

1
2
3 nationally and internationally. This will be facilitated by planned publications (results paper,
4 process evaluation paper) in open access journals, presentations at high impact ED and LBP
5 conferences, stakeholder engagement via a national ED online symposium and workshops.
6
7 This includes the provision of training on implementation of the intervention (consistent with
8 the information delivered at the stakeholder engagement), engaging clinicians, and ensuring
9 sustainability in change of practice. It is also planned to produce media resources that will
10 permit the widespread dissemination of the study's findings.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements: The investigators would like to thank the clinical and administrative staff of the Emergency Departments at the participating EDs. We also thank the Physiotherapy Department and Jamieson Trauma Centre at the Royal Brisbane and Women’s Hospital.

Author Contributions: All authors (SO, NEF, GM, TS, JW, TC, GCM, PB, MC, JH, WS, EB) contributed to the study design and preparation, data collection planning and process, intervention design and implementation, and data analysis and interpretation. All authors contributed to and approved the final manuscript preparation.

Funding: This study has been funded by a Hospitals Contribution Fund (HCF) Research Foundation Translational Research Grant Program. This funding body had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Competing interests: None declared.

Patient consent for publication: Not required.

Ethics approval: This study received ethical approval from the Metro North Human Research Ethics Committee (HREC/2022/MNHA/87995). Subsequent site specific approval was then gained from each participating hospital site.

Provenance and peer review: Not commissioned; externally peer reviewed.

Open access: This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
ErasmusHogeschool

given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Emergency department care. Australian Institute of Health and Welfare (AIHW), Australian Government: Canberra, 2021.
- 2 Management of people with acute low back pain: model of care. NSW Agency for Clinical Innovation. NSW Health Chatswood, NSW, 2016. p. 39.
- 3 Machado GC, Rogan E, Maher CG. Managing non-serious low back pain in the emergency department: Time for a change? *Emerg Med Australas* 2018;30(2):279-82.
- 4 Makary MA, Overton HN, Wang P. Overprescribing is major contributor to opioid crisis. *BMJ* 2017;359:j4792.
- 5 Heine J, Window P, Hacker S, et al. Adherence to recommended guidelines for low back pain presentations to an Australian emergency department: Barriers and enablers. *Australas Emerg Care* 2023;26(4):326-32.
- 6 Coombs DM, Machado GC, Richards B, et al. Effectiveness of a multifaceted intervention to improve emergency department care of low back pain: a stepped-wedge, cluster-randomised trial. *BMJ Qual Saf* 2021;30(10):825-35.
- 7 Machado GC, Richards B, Needs C, et al. Implementation of an evidence-based model of care for low back pain in emergency departments: protocol for the Sydney Health Partners Emergency Department (SHaPED) trial. *BMJ Open* 2018;8(4):e019052.
- 8 Lopez Bernal J, Cummins S, Gasparrini A. The use of controls in interrupted time series studies of public health interventions. *Int J Epidemiol* 2018;47(6):2082-93.

- 9 Curran GM, Bauer M, Mittman B, et al. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care* 2012;50(3):217-26.
- 10 Health Service Strategy 2021-26. Available from:
11 <https://metronorth.health.qld.gov.au/about-us/publications/health-service-strategy-planning>.
- 11 Brisbane North PHN and Metro North Hospital and Health Service. Population Health Report. Brisbane: Department of Health; 2019.
- 12 Penfold RB, Zhang F. Use of interrupted time series analysis in evaluating health care quality improvements. *Acad Pediatr* 2013;13(6 Suppl):S38-44.
- 13 Wagner AK, Soumerai SB, Zhang F, et al. Segmented regression analysis of interrupted time series studies in medication use research. *J Clin Pharm Ther* 2002;27(4):299-309.
- 14 Interrupted Time Series Power Calculation using DO Loop Simulations (sas.com).
- 15 Care ACoSaQiH. Low Back Pain Clinical Care Standard. Sydney: ACSQHC; 2022.
- 16 Graham ID, Logan J, Harrison MB, et al. Lost in knowledge translation: time for a map? *J Contin Educ Health Prof* 2006;26(1):13-24.
- 17 Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011;6:42.
- 18 Damschroder LJ, Banaszak-Holl J, Kowalski CP, et al. The role of the champion in infection prevention: results from a multisite qualitative study. *Qual Saf Health Care* 2009;18(6):434-40.
- 19 WHO Collaborating Centre for Drug Statistics Methodology. ATC/DDD Index. Oslo: Norwegian Institute of Public Health; 2023.

- 20 Proctor E, Silmere H, Raghavan R, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Adm Policy Ment Health* 2011;38(2):65-76.
- 21 Gale NK, Heath G, Cameron E, et al. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol* 2013;13:117.
- 22 The L. Managing the opioid crisis in North America and beyond. *Lancet* 2022;399(10324):495.
- 23 Lalic S, Ilomaki J, Bell JS, et al. Prevalence and incidence of prescription opioid analgesic use in Australia. *Br J Clin Pharmacol* 2019;85(1):202-15.
- 24 Donovan PJ, Arroyo D, Pattullo C, et al. Trends in opioid prescribing in Australia: a systematic review. *Aust Health Rev* 2020;44(2):277-87.
- 25 Caughey GE, Gadzhanova S, Shakib S, et al. Concomitant prescribing of opioids and benzodiazepines in Australia, 2012-2017. *Med J Aust* 2019;210(1):39-40.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGENDS

Figure 1 Diagram depicting the 6 components of the implementation plan designed to change clinician behaviour towards the use of best-practice LBP management guidelines, specifically the NSW Agency for Clinical Innovation (ACI) model of care for acute LBP[2] and the more recent Australian LBP Clinical Care Standard.[15]

For peer review only

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Erasmus Hogeschool



Figure 1 Diagram depicting the 6 components of the implementation plan designed to change clinician behaviour towards the use of best-practice LBP management guidelines, specifically the NSW Agency for Clinical Innovation (ACI) model of care for acute LBP[2] and the more recent Australian LBP Clinical Care Standard.[15]

338x190mm (300 x 300 DPI)

Supplementary Table 1: Parameters for extracting potential Low Back Pain presentations using the Emergency Department (ED).

Code	Explanatory Description
Presenting Complaint	
17005	Requesting Investigation
17006	Requesting Medication/Certificate/Results
17008	Unplanned ED Representation
21002	Altered Sensation
25005	Back Pain
9047	Mechanical Strain
Primary ICD-10 AM Diagnosis Code	
G95.9	Disease of Spinal Cord, Unspecified
M13.99	Arthritis, Unspecified
M19.99	Arthrosis, Unspecified
M25.59	Pain in a Joint, Site Unspecified
M25.99	Unspecified Joint Disorder
M48.99	Unspecified Spondylopathy
M51.9	Intervertebral Disc Disorder
M53.3	Sacro Coccygeal Disorders
M53.99	Unspecified Dorsopathy
M54.19	Radiculopathy, Site Unspecified
M54.2	Cervicalgia
M54.3	Sciatica
M54.5	Low Back Pain
M54.6	Pain in Thoracic Spine
M54.99	Unspecified Dorsalgia Site Unspecified
M79.19	Myalgia, Site Unspecified
M79.29	Neuralgia and Neuritis, Unspecified
M89.99	Unspecified Disorder of Bone
M95.9	Acquired Musculo Skeletal Deformity
R25.2	Cramp and Spasm
R26.8	Abnormalities of Gait and Mobility
R29.89	Musculo Skeletal System Symptoms / Signs
R52.9	Pain, Unspecified
S22.00	Fracture of Thoracic Vertebra
S23.3	Sprain and Strain of Thoracic Spine
S32.00	Fracture of Lumbar Vertebra
S32.82	Fracture of Lumbosacral Spine
S33.3	Dislocation Lumbar Spine and Pelvis
S33.7	Sprain/Strain Lumbar Spine/Pelvis
S34.1	Other Injury of Lumbar Spinal Cord
S38.1	Crush Injury Abdomen/Lower Back/Pelvis
S39.9	Injury of Abdomen/Lower Back/Pelvis
Z02.7	Issue of Medical Certificate
Z04.9	Examination / Observation Unspecified
Z76.0	Issue of Repeat Prescription

Note: Codes and descriptions relate to the intervention ED location of FASTRACK and control ED location of BLUE or GREEN ZONE.

BMJ Open

Reinforcing Informed Medication prescription for low back pain in the Emergency department (RIME): a controlled interrupted time-series implementation study protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2023-082668.R1
Article Type:	Protocol
Date Submitted by the Author:	31-Jan-2024
Complete List of Authors:	O'Leary, Shaun; University of Queensland; Royal Brisbane and Women's Hospital, Physiotherapy Department Heine, Janelle; Royal Brisbane and Women's Hospital Emergency and Trauma Centre; Royal Brisbane and Women's Hospital, Physiotherapy Department Warren, Jacelle; Metro North Hospital and Health Service, Jamieson Trauma Institute, Royal Brisbane & Women's Hospital (RBWH) Smyth, Tanya; Metro North Hospital and Health Service, Jamieson Trauma Institute, Royal Brisbane & Women's Hospital (RBWH) Ballard, Emma ; QIMR Berghofer Medical Research Institute, Mitchell, Gary; Royal Brisbane and Women's Hospital Emergency and Trauma Centre See, William; The Prince Charles Hospital ED Barlas, Panos; Metro North Hospital and Health Service, Jamieson Trauma Institute, Royal Brisbane & Women's Hospital (RBWH) Machado, Gustavo; University of Sydney - Camperdown and Darlington Campus, School of Public Health Cottrell, Michelle; Royal Brisbane and Women's Hospital, Physiotherapy Department Comans, Tracy; 8. Centre for Health Services Research, The University of Queensland Foster, Nadine; The University of Queensland, Surgical Treatment and Rehabilitation Service
Primary Subject Heading:	Emergency medicine
Secondary Subject Heading:	Emergency medicine, Evidence based practice, Public health
Keywords:	Back pain < ORTHOPAEDIC & TRAUMA SURGERY, ACCIDENT & EMERGENCY MEDICINE, CLINICAL PHARMACOLOGY, Implementation Science, Protocols & guidelines < HEALTH SERVICES ADMINISTRATION & MANAGEMENT



Reinforcing Informed Medication prescription for low back pain in the Emergency department (RIME): a controlled interrupted time-series implementation study protocol

Authors: Shaun O'Leary,^{1,2} Janelle Heine,^{1,3} Jacelle Warren,⁴ Tanya Smyth,⁴ Emma Ballard,⁵ Gary Mitchell,³ William See⁶, Panagiotis Barlas,⁴ Gustavo C Machado,⁷ Michelle Cottrell,¹ Tracy Comans,⁸ Nadine E Foster^{2,9}

Institutional addresses:

1. Physiotherapy Department, Royal Brisbane and Women's Hospital, Metro North Health
2. School of Health and Rehabilitation Sciences, The University of Queensland
3. Emergency and Trauma Centre, Royal Brisbane and Women's Hospital, Metro North Health
4. Jamieson Trauma Institute, Royal Brisbane and Women's Hospital, Metro North Health
5. Statistics Unit, QIMR Berghofer Medical Research Institute, Herston
6. Emergency Department, The Prince Charles Hospital, Metro North Health
7. Institute for Musculoskeletal Health, University of Sydney and Sydney Local Health District
8. Centre for Health Services Research, The University of Queensland
9. Surgical Treatment and Rehabilitation Service (STARS) Education and Research Alliance, The University of Queensland and Metro North Health

Corresponding author: Associate Professor Shaun O'Leary Postal Address: Physiotherapy Department, School of Health and Rehabilitation Sciences, University of Queensland, St Lucia, Queensland, 4072, Australia. Institute Email Address: s.oleary@uq.edu.au
Phone : +61 4 07746299.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Introduction: Management guidelines for low back pain (LBP) recommend exclusion of serious pathology, followed by simple analgesics, superficial heat therapy, early mobilisation, and patient education. An audit in a large metropolitan hospital Emergency Department (ED) revealed high rates of non-recommended medication prescription for LBP (65% of patients prescribed opioids, 17% prescribed benzodiazepines), high inpatient admission rates (20% of ED LBP patients), delayed patient mobilisation (on average 6 hours), and inadequate patient education (48% of patients). This study aims to improve medication prescription for LBP in this ED by implementing an intervention shown previously to improve guideline-based management of LBP in other Australian EDs.

Methods and Analysis: A controlled interrupted time series study will evaluate the intervention in the ED before (24 weeks; March 20 to September 3, 2023) and after (24 weeks; November 27 to May 12, 2024) implementation (12 weeks; September 4 to November 26, 2023), additionally comparing findings with another ED in the same health service. The multicomponent implementation strategy uses a formalised clinical flow chart to support clinical decision-making and aims to change clinician behaviour, through clinician education, provision of alternative treatments, educational resources, audit and feedback, supported by implementation champions. The primary outcome is the percentage of LBP patients prescribed non-recommended medications (opioids, benzodiazepines, and/or gabapentinoids), assessed via routinely collected ED data. Anticipated sample size is 2000 patients (n=1000 intervention, n=1000 control) based on average monthly admissions of LBP presentations in the EDs. Secondary outcomes include inpatient admission rate, time to mobilisation, provision of patient education, imaging requests, re-presentation to the ED within 6 months, and healthcare costs. In nested qualitative research, we will study ED clinicians' perceptions of the implementation and identify how benefits can be sustained over time.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Erasmus Hogeschool

Ethics and dissemination This study received ethical approval from the Metro North Human Research Ethics Committee (HREC/2022/MNHA/87995). Study findings will be published in peer-reviewed journals and presented at international conferences and educational workshops.

Trial registration number: Australia New Zealand Clinical Trials Registry: ACTRN12622001536752.

Keywords: low back pain; opioids; emergency department; guidelines; implementation; interrupted time series

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ARTICLE SUMMARY

Strengths and limitations of this study

- Controlled interrupted time series design will test the effects of the intervention on relevant outcomes while observing and gathering information on implementation in a real-world ED setting.
- Outcomes are not only evaluated before and after implementation at the intervention ED, but also through comparison with a control ED.
- Process evaluation will explore components of the implementation that clinicians found useful, as well as perceived barriers to adoption and sustainability of improved LBP management in the ED.
- Both participating EDs are within the same health service and therefore results may lack generalisability to other settings.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Erasmus Hogeschool

INTRODUCTION

Low back pain (LBP) is the fifth most common presentation to Australian Emergency Departments (EDs).[1] Management guidelines for LBP recommend an initial exclusion of serious pathology (e.g. spinal fracture, infection, cancer), followed by the use of simple analgesics, superficial heat therapy, early mobilisation to improve function, and the provision of patient education to enable self-management.[2] Research to date has evidenced inconsistent use of best practice guidelines for LBP in Australian EDs.[3]

In line with previous research a recent audit of ED routinely collected data at a large metropolitan tertiary hospital in Queensland, Australia, identified a high rate of medication prescription for LBP, particularly opioids (prescribed for 65% of LBP patients) but also muscle relaxants (benzodiazepine, 17%). These are not recommended first line medications for LBP and have known side-effects including dependence, poisoning and death.[4] Other concerning audit findings were that 20% of patients with LBP were admitted as an inpatient (to either the Short Stay Unit or hospital wards), had long waits until they received support to mobilise (6 hours on average until they were supported to walk), and may not have received patient education (48% of patients' records had no documented evidence of education/advice).[5] In response, an intervention was sought to promote guideline based management of LBP in this metropolitan hospital ED to improve patient care, particularly with regard to medication prescription.

A recent randomised trial conducted in other Australian EDs was identified that showed improvement in the use of LBP best practice guidelines by clinicians.[6] The Sydney Health Partners Emergency Department (SHaPED) trial successfully reduced opioid prescription by over 12% (50.5% in intervention EDs v 62.8% in control) (OR=0.57, 95% CI 0.38-0.85), with no adverse effects on patients' pain scores or satisfaction with care.[6] Improvements were also observed regarding clinicians' beliefs and knowledge regarding the management of LBP.[6]

The SHaPED trial specifically intervened to change ED clinician behaviour regarding the use of LBP guidelines.[6] The SHaPED trial used a multicomponent implementation strategy that was informed not only by a review of practice, but also included locally driven strategies that clinicians identified as potential facilitators and barriers to the uptake of LBP guidelines in the participating Sydney EDs.[7]

This protocol paper describes the Reinforcing Informed Medication prescription for low back pain in the Emergency department (RIME) study. The aim of the RIME study is to improve the management of LBP in our Queensland-based ED, by implementing and evaluating the intervention shown to be effective in the SHaPED trial in Sydney EDs. The primary and secondary objectives of the study are as follows.

Primary objective

Objective 1: To evaluate the impact of the intervention on the percentage of patients with LBP who are prescribed non-recommended medications (i.e. opioids, benzodiazepines and/or gabapentinoids) in the ED (primary outcome).

Secondary objectives

Objective 2: Evaluate the impact of the intervention on the percentage of patients with LBP who are admitted to hospital, or who receive support to mobilise (including time to support to walk), appropriate advice/education, lumbar spine imaging, and who re-present to ED within 6 months (secondary outcomes).

Process evaluation

Objective 3: Undertake a cost analysis of implementing the intervention from a hospital resources perspective.

Objective 4: Understand the perspectives (e.g. acceptability, adoption, appropriateness, feasibility, sustainability) of ED clinicians towards the intervention, the implementation strategy, and sustainability of changes in LBP management in the ED.

Objective 5: Determine the fidelity to which each component of the implementation strategy was delivered, including the percentage of appropriate clinician encounters with the RIME training team during implementation.

METHODS AND ANALYSIS

Study design

RIME is a controlled Interrupted Time Series (ITS) study design evaluating outcomes across study phases with nested process evaluation. This design permits evaluation of changes in outcomes both pre- and post-implementation in the intervention ED, as well as intervention and control ED comparisons to exclude confounders due to co-interventions or other events occurring during the study timeline.[8] The study is a prospective effectiveness-implementation hybrid model (Type 1), testing intervention effects on relevant measurable outcomes (Objectives 1-3) while observing and gathering information on implementation (Objectives 4-5) in a real-world setting.[9]

Scheduled activity over the three study phases are shown in Table 1 and include; Phase 1/ Pre-implementation usual care - during which primary and secondary clinical outcome data will be collected over a 6-month period for adult patients (aged 18 years and over) presenting with LBP to either the intervention or control EDs; Phase 2/ Implementation - during which the intervention will be implemented with clinicians over a 12-week period at the intervention ED only; and Phase 3/ Post-implementation care – which repeats the identical data collection procedure of Phase 1 over a 6-month period at both intervention and control EDs.

Table 1 Schedule of procedures during the RIME study with week (W) 1-24 representing the pre-implementation phase (Pre), week 25-36 the implementation phase, and week 37-60 the post-implementation phase (Post). Implementation weeks and implementation strategies are shaded grey. Black arrows denote continuous periods of outcome measurements recorded fortnightly, thus initial recordings are W2 and W38 of the pre-and post-implementation phases, respectively.

	Pre			Implementation								Post					
Procedures/Measurements	W2	W24	W25	W26	W27	W28	W29	W30	W31	W32	W33	W34	W35	W36	W38	W60	
Measurement of Primary and Secondary Outcomes (fortnightly)	✓	➡		✓		✓		✓		✓		✓		✓	✓	➡	✓
Study Information Sheets provided			✓														
Educational seminars delivered			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓			
Educational materials provided			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓			
Provision of alternative LBP treatment options			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓			
Fast-track referral to outpatient services			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓			
Audit and feedback						✓				✓				✓			
Implementation Champion actioned			✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓			

Cost data collection	√	√	√	√	√	√	√	√	√	√	√	√	√	√	√
Audit & Observation Measures Recorded				√		√		√		√		√			
Qualitative Interviews commence														√	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Patient and public involvement

None.

Study setting

The study will be conducted in two large metropolitan tertiary hospital EDs in Queensland, Australia. One ED will serve as the intervention site and the other the control site. Both EDs are governed by the same Hospital and Health Service, and the catchment populations of both are almost identical, projected to be approximately 368,000 persons by 2026.[10] The median age of residents in the intervention ED catchment area is slightly younger (approximately 34 years) than for the control ED (approximately 37 years),[11] however both catchment areas have over 1,000 LBP patients presenting to their ED each year (January to December 2021 data; intervention ED 1,046 patients, control ED 1,485 patients), and each hospital has a similar skill-mix in their ED clinical teams. These clinical teams include senior and junior ED doctors and nurses, as well as Emergency Physiotherapy Practitioners (EPPs). The EPPs work relatively independently within the ED and are usually responsible for the assessment, investigation, and management of LBP patients.

Participants

There are three participant groups in this study.

Patients with LBP: Data from adult patients (18+ years) presenting to either the intervention or control ED with LBP will be included in the study. A waiver of consent has been granted for the use of the anonymous (de-identified) ED patient data. Only routine ED service data will be required for the study, there is no foreseeable risk or additional burden placed on patients, and

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
ErasmusHogeschool

it is impractical to gain individual consent for the use of these routine service data. It was also not deemed necessary to burden patients with patient-reported outcome data collection, given the previous SHaPED trial already showed the intervention to be effective in reducing opioid prescription without compromising patient reported pain or satisfaction.[6]

Potentially eligible LBP patient cases will be identified at the two sites using the Electronic Medical Record (EMR) system - Emergency Department Information System (EDIS) - with the following parameters:

- Seen in a low-acuity area of the ED, AND
- Presenting complaint related to 'Back Pain', 'Altered Sensation', 'Mechanical Strain', 'Requesting Investigation' or 'Requesting Medication/Certificate/Results', OR
- Primary diagnosis code related to LBP (coding system ICD-10-AM)

Exclusions are patient cases with spinal fracture, infections, severe neurological disorders, cauda equina syndrome, major or multi-trauma, malignancy, psychiatric disorders and mental health disorders, and other non-spinal causes of LBP. These exclusions will be identified through screening EDIS information (see Supplementary Table 1 for a full list of EDIS extraction parameters). Potentially eligible LBP patients will then be manually screened by site-based study staff who will review the presenting problem free text in the ED data to further refine the study sample to those with LBP who do not present with one of the exclusions. Reasons for exclusion during manual review will be collected and summarised.

ED Clinicians: All ED clinicians at the intervention ED will be informed about the intervention as a service improvement activity but their level of involvement will be at their discretion. Individual ED clinician consent to participate will not be sought. This approach ensures these ED clinicians do not feel pressured to undertake activities or practices they do not wish to. The

research team will attempt to facilitate uptake by offering all ED clinicians all components of the implementation strategy. Only the intervention ED clinicians will be given a RIME study information sheet to minimise the risk of potential contamination at the control ED.

ED Clinicians who take part in the nested interviews: Following the implementation phase, up to 20 ED clinicians will be invited to participate in a qualitative semi-structured interview exploring their views of the intervention, its implementation and potential sustainability of improvements in practice. This sample size will enable representation from the three ED clinician disciplines (ie. EPPs, nurses, medical) and is similar to our earlier ED clinician qualitative study.[5] Consent from these participants will be implied by their voluntary participation in the interviews.

Sample Size

Based on recommendations[12, 13] and given the number of observations within each time-period and effect on statistical power, the study will use 12 time-periods spaced 2 weeks apart (6 months total) for all outcomes in both the pre- and post-implementation phases (Phases 1 and 3). Results from the SHaPED trial[6] showed a difference in any opioid medication prescription between control (62.8%) and intervention (50.5%) groups of 12.3%. Simulations based on these findings using SAS v9.4 suggests the power to detect at least a 12% reduction in non-recommended medication prescription (the primary outcome) at the intervention ED is between 70% and 80%.[14] Anticipating 500 LBP ED presentations in 6 months at the intervention ED allows for approximately 45 patients in each 2-week time-period before and after implementation. The total anticipated sample size is 2000 (500 in each of the two EDs, in each of the pre- and post-implementation periods); (n=1000 intervention, n=1000 control).

Intervention and Implementation Plan

The intervention addresses ED clinician behaviour to improve their use of best-practice LBP management guidelines, specifically the NSW Agency for Clinical Innovation (ACI) model of care for acute LBP[2] and the more recent Australian LBP Clinical Care Standard.[15]

To best inform the implementation approach for the local ED context, five focus groups with 18 ED clinicians (medical, nursing, EPPs) were previously conducted.[5] These interviews gained the local perspective on current practice, LBP guidelines, and barriers and enablers potentially impacting the uptake of guidelines for LBP management at the intervention ED.[5] Findings indicated that adherence to guidelines is primarily driven by (i) clinician beliefs and behaviours; (ii) patient expectations and behaviours; and (iii) workflow processes. Clinicians' beliefs around role scope and accountability were perceived to serve as barriers to providing recommended treatments such as prescription of simple analgesics and early mobilisation. Patients' expectations of care (e.g., specifically requesting stronger analgesics) were perceived to influence clinical decisions regarding medication prescription. Limited access to out-of-hours physiotherapy, along with National Emergency Access Target pressures, were also felt to influence clinical decisions, particularly inpatient admissions to the Short Stay Unit (SSU).[5]

Through methodological triangulation (audit, survey, focus groups), the focus group data provided a deeper understanding of the previous ED audit results.[5] Strategies suggested to improve best practice within the intervention ED context included materials such as a formalised clinical flow chart (to support clinical decision-making that aligns with best practice), targeted education to medical and nursing staff around safe mobilisation practices and best practice management of LBP patients, and patient education resources to better support patient self-management. Additionally, it was identified that the multifaceted intervention would require organisational support to embed the strategies within clinical

management systems including the EMR, to facilitate clinicians to adopt best practice routinely.[5]

Implementation of the intervention will be underpinned by the Knowledge-to-Action framework (as per the SHaPED trial),[16] incorporating evidence-based implementation strategies specifically targeting the behaviour of ED clinicians at the intervention ED. The Capability-Opportunity-Motivation=Behaviour (COM-B) behaviour change theoretical framework[17] has been used to shape the implementation plans to support behaviour change in ED clinicians to enhance capability, opportunity, and motivation to improve uptake of guidelines when managing LBP patients in the ED.

Figure 1 depicts the multicomponent implementation strategy, including:

1. *Educational Seminars:* These will comprise structured best practice updates from experienced ED clinicians (i.e. Emergency Physiotherapy Practitioners, Emergency Medical Consultants) that focus on knowledge and skills for assessing, managing, educating, and referring patients in line with LBP guidelines.[2] Preparation of these best practice updates will be assisted by the investigator team. These best practice sessions will be offered on numerous occasions throughout the 12-week implementation period in protected ED teaching time. All ED clinicians will be invited to participate and clinician participation in the education sessions will be tracked through a logbook, with reminders sent and personal communication from the study team and ED clinical leads where needed.

2. *Educational materials:* ED clinicians will receive a hard copy of the LBP guidelines and LBP Clinical Standards,[2][15] a link to an established, contemporary evidence-based patient-facing website (<https://mybackpain.org.au/>), and the formalised clinical flow chart to support clinical decision-making such as the appropriate use of analgesic medicines. Posters highlighting key messages about benefits and harms of opioid medicines, lumbar imaging, and

inpatient admission will be displayed throughout the intervention ED. Anonymised patient cases from study phase 1 will be discussed, illustrating examples of non-recommended practice and best practice. Patient educational materials (based on the LBP guidelines) and scripts to guide conversations with patients will be provided so that clinicians can use these with patients.

3. Provision of alternative treatment options for LBP: Non-opioid pain medicines will be made more easily accessible to clinicians as an evidence-based alternative to opioid medicines or muscle relaxants. Heat wraps (used as a non-pharmacological modality for pain relief) will also be made available to clinicians with encouragement to use these as alternatives to non-recommended pharmacological treatments.

4. Fast-track referral to outpatient services: Clinicians will be educated on available referral pathways, for whom these are appropriate, and how to collaboratively facilitate the referral process with the patient. Referral pathway options include; private physiotherapy services within the primary care setting, onsite public physiotherapy outpatient services, or onsite advanced-practice musculoskeletal physiotherapy screening services.

5. Audit and feedback: During the implementation phase, routinely collected ED data will be audited and clinicians provided with structured ED-level- and discipline-level audit and feedback reports, including medication prescription data. These reports are planned to be provided at 4, 8 and 12 weeks after the start of the implementation phase.

6. Support from an ED 'Implementation Champion': A key feature of the implementation is the inclusion of a dedicated 'Implementation Champion'. An experienced ED clinician (investigator Heine) will undertake this role, as champions are considered vital to successful implementation and change in their own sphere of influence, particularly when intrinsically motivated and enthusiastic about the practices they promote.[18] The implementation champion will support reinforcement of implementation aims with ED staff, provide personal feedback sessions, and offer one-to-one or small group discussions with ED staff as needed.

Insert Figure 1 Here

Study Outcomes

Study outcomes include primary and secondary clinical outcomes, as well as a nested process evaluation.

Data related to the primary and secondary clinical outcomes, in addition to key demographic characteristics and ED management data, will be obtained using either routine electronic data from EDIS, or via manual extraction from the medical records. Data validation rules will be implemented to reduce data transcription errors and improve data quality. Following completion of the data extraction, patient identifying information will be removed and replaced with an anonymised ID code.

Data obtained from EDIS include patient presentation date/time, demographic characteristics (age, gender, postcode) and ED management details (triage category, arrival mode [ambulance, walk in etc], primary clinician, consultations performed, ED discharge destination, time to treatment etc). Data obtained via manual extraction from the medical record include previous history of LBP, pain severity, ED treatment details (attempted mobilisation, provision of education/advice, heat pack used, imaging requested etc), and medications prescribed while in the ED and on discharge. Data relating to re-presentations within 6 months will also be obtained from EDIS.

Primary Clinical Outcome (Objective 1)

- Percentage (%) of patients presenting to the ED with LBP prescribed non-recommended medications in the ED. Non-recommended medications are grouped under the Anatomical

Therapeutic Chemical (ATC) classification[19] of N02A (Opioids), N05B (Benzodiazepines) or N02BF (Gabapentinoids).[6, 7]

Secondary Clinical Outcomes (Objective 2)

- Percentage (%) of patients presenting to the ED with LBP prescribed analgesic medications in the following ATC categories: N02BE01 (Paracetamol) or M01A (Anti-inflammatory and Antirheumatic Products, non-steroidal).
- Mean/Median time to mobilisation (time to be supported to walk).
- Percentage (%) of patients receiving education about LBP care.
- Percentage (%) of patients receiving lumbar spine imaging.
- Percentage (%) of patients admitted to hospital or short stay.
- Percentage (%) of patients re-presenting to the same ED facility within 6 months.

Process Evaluation

The process evaluation of the implementation is underpinned by the framework developed by Proctor et al.[20] This framework consists of eight distinct implementation outcomes which serve to understand the implementation process and determine implementation success. Implementation outcomes will be assessed by the approaches shown in Table 2.

Cost Evaluation (Objective 3)

Costs will be estimated based on project staff time to conduct the implementation components/activities, clinical staff time to attend the educational seminars and cost of developing materials. Research specific costs (e.g. semi-structured interviews) will be excluded from this analysis, however, will be reported separately. Direct health system costs

will be estimated using actual cost data obtained from each hospital’s clinical costings department for each eligible ED patient presentation and any related inpatient episodes of care. The cost categories collected include medical, nursing, allied health, pathology, imaging, consumables, hotel and medications. These data will be used to evaluate and compare the cost of treating LBP patients pre- to post-implementation at the intervention ED, and between intervention and control EDs.

Semi-structured interviews (Objective 4)

Semi-structured interviews will be conducted with ED clinicians addressing Proctor et al.’s [20] implementation outcome domains of: acceptability, adoption, appropriateness, feasibility, and sustainability (Table 2). The interview guide will be developed through an iterative review and feedback process involving internal study team consultation and testing. The interview protocol will be tested on available ED clinician/s who have been exposed to the RIME intervention. Interviewing will be conducted following the intervention phase of the study, and a range of ED clinicians (i.e., nurses, medical officers, physiotherapists) will be approached to participate. All interviews will be audio-recorded, transcribed verbatim and coded using a hybrid deductive and inductive approach. Other study team members will independently code a subset of the interviews, and any discrepancies will be discussed and reassessed to reach agreement. The remaining interviews will be coded from the agreed approach. The coded data will be used to develop themes, which will be reported to the study team for discussion. The final agreed thematic analysis will be written into final reporting and manuscripts.

Audit & Observation (Objective 5)

Both penetration and fidelity outcomes, as defined by Proctor et al.[20] (Table 2) of each proposed implementation strategy component will be determined by calculating the percentage (%) of encounters where the strategy was actually used, as well as the extent to which the strategy was delivered as intended (%). As feasible, regular monitoring throughout the 12-week implementation period (Phase 2) will occur through both chart audits (reviewed treatment notes) and peer observation led by the implementation champion. Engagement with the implementation components will be additionally captured by recording attendance to educational sessions and use of resources (recorded within a logbook at a department and discipline level, not individual clinician level) as accurately as possible.

Table 2 Implementation outcomes based on Proctor et al.[20] and adapted for RIME. Time points of measurement include Phase 2 Implementation or Phase 3 Post-implementation

Outcome	Definition from Proctor et al.	Definition as applied to RIME	Timing of Measurement	Measurement Source
Acceptability	Perception among implementation stakeholders that a given treatment, service, practice, or innovation is agreeable, palatable, or satisfactory.	ED clinicians’ perceptions of the RIME intervention components, and implementation strategies, as agreeable, palatable, or satisfactory.	Phase 3 Post-implementation	Clinician interviews
Adoption	The intention, initial decision, or action to try, or employ, an innovation or evidence-based practice. Adoption also may be referred to as ‘uptake’.	The intention and behavioural uptake of the RIME intervention components by ED clinicians to facilitate adherence to LBP Clinical Standards.	Phase 3 Post-implementation	Clinician Interviews
Appropriateness	The perceived fit, relevance, or compatibility of the innovation or evidence-based practice for a given practice setting, provider, or consumer; and/or perceived fit of the innovation to address a particular issue or problem.	The extent to which ED clinicians perceive the fit or relevance of the RIME intervention components within their clinical setting to facilitate adherence to LBP Clinical Standards.	Phase 3 Post-implementation	Clinician Interviews
Feasibility	The extent to which a new treatment, or an innovation, can be successfully used or carried out within a given agency or setting.	The extent to which implementation of the RIME intervention, and implementation strategies, is considered feasible for the clinical setting under study, allowing for individual tailoring given the differing needs of clinicians.	Phase 3 Post-implementation	Clinician Interviews

Fidelity	The degree to which an intervention was implemented as it was prescribed in the original protocol or as it was intended by the program developers.	The degree to which each RIME intervention component was implemented as described within the original protocol.	Phase 2 Implementation	Audit & Observation
Implementation Cost	The cost impact of an implementation effort. The cost of implementing an intervention includes the costs of the intervention, the implementation strategy used, and the service delivery.	The costs involved in the development and implementation of the RIME intervention within the ED setting under study.	Phase 2 Implementation	Health Economic Evaluation
Penetration	The integration of a practice within a service setting and its subsystems. Can also be referred to as 'reach' (using the RE-AIM framework).	The extent to which each RIME intervention component was accessed/used by the ED clinicians.	Phase 2 Implementation	Audit & Observation
Sustainability	The extent to which a newly implemented treatment is maintained or institutionalised within a service setting's ongoing, stable operations.	The extent to which the RIME intervention components could be embedded as standard practice within the ED setting, following the completion of the RIME study.	Phase 3 Post implementation	Clinician Interviews

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Data Analysis

Primary and Secondary Clinical Outcomes: Two analytic approaches will be considered to assess primary and secondary outcomes: 1) a controlled interrupted time series (ITS) analysis, where data are aggregated on a fortnightly basis, and 2) a segmented regression analysis on all data. The change in trend (slope) between pre- and post-implementation periods and the change in level between the end of the pre-implementation period and the beginning of the post-implementation periods and between the beginning of the pre-implementation period and at the end of the post-implementation period will be described for separate models for each hospital and models including hospital ED as a fixed effect. Additionally for models including hospital ED as a fixed effect, the difference between the two EDs for the change in slope and change in level will be described. All estimates will be reported with 95% confidence intervals (CI). Data will be presented visually, overlaid with the regression line for pre- and post-implementation periods for each outcome. Assumptions relating to linearity, autocorrelation, stationarity and heteroscedasticity will be examined, and consideration given to the influence of outliers and the normality of residuals in developing a robust model.

Patient characteristics (e.g age, sex, Socio-Economic Indexes for Area (SEIFA), ED arrival mode, severity (i.e. Australian Triage Scale (ATS) 1 – 5), type of treating clinician and day/time of presentation will be compared between EDs during the pre-implementation period and between pre and post-implementation periods within EDs. Associations between these characteristics and outcomes will also be examined. These comparisons will be done using the Pearson Chi-squared test or Fisher’s exact test for categorical data and Student t-test or Mann-Whitney U test for continuous data. Categorical data will be presented as frequency and percentage and continuous data as mean and standard deviation or median and interquartile range if not normally distributed. The distribution of these characteristics overtime will be

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Erasmus Hogeschool

examined visually and reported. If these characteristics are found to be associated with the intervention or they are associated with the outcome and change over time in both EDs, these variables will be adjusted for in the segmented regression. Stepwise backward elimination will be used to select the most parsimonious model.

Planned exploratory analyses include; a three period (pre, implementation, post) analysis; a two period analysis where data from the implementation period is combined with the post implementation period. Planned sensitivity analyses include; analysis of index presentation only, multilevel segmented regression model to account for clustering of observations within patients and observations within clinician groups, and controlling for differences in the patient base by including the value of the outcome denominator in the ITS model. The planned subgroup analysis is to understand the ED medication usage in patients who are transported to hospital via ambulance (given ambulance staff may provide medications to patients with LBP before their arrival at the ED).

Potential sources of bias exist and will be monitored and addressed as feasible. For example, if change over time is clearly nonlinear, transformation of continuous outcomes would be considered. Seasonality will be examined by plotting outcomes over time, however determining its effect is challenging due to the relatively short pre- and post-implementation periods of 6 months. These short outcome periods also make it difficult to detect observational bias although having a control ED to compare changes over time provides some support.

Other mechanisms have been included to minimise other potential sources of bias such as a data dictionary and formal process for manual extraction of data, data validation rules on variables, and single clinician data entry.

Process Evaluation:

Cost Evaluation: To assess the value for money of the intervention, total healthcare costs will be compared between pre and post-implementation phases, and between intervention and control EDs. The costs will be compared using models appropriate for clustered data and the non-normal nature of the cost and variable (e.g., generalised linear model with robust standard errors). Confounders will be included in the model if assessed as necessary in the statistical analysis. A final parsimonious model will be selected following data examination and measures of model fit. Following the base case analysis, sub-group analysis will be conducted of the main categories of cost (e.g. medication only) to assess where differences have accrued.

Semi-structured interviews: Interviews will be transcribed and analysed using NVivo software. The Framework Method, as described by Gale et al.[21] will be employed to deductively analyse interview data based upon Proctor et al.’s pre-defined implementation outcomes[20] introduced in Table 2. In addition, ‘open-coding’ will be performed for information that does not readily adhere to the pre-defined framework, where themes/concepts will be generated using an inductive thematic analysis approach.

Audit & Observation: Simple descriptive statistics will be employed to measure fidelity and penetration of each intervention component. Fidelity will be measured at regular intervals (approximately every 2 weeks) and descriptive analysis undertaken to explore any changes in fidelity over the implementation period.

ETHICS AND DISSEMINATION

No risks to patients within this study are foreseeable given its intention to improve the management of LBP in the ED, based on best-practice guidelines. The study is aligned with concerted efforts nationally and globally to reduce the use of opioids which continues to be a

major health concern.[4, 22-25] It is hypothesised the intervention will reduce the rate of non-recommended medication prescriptions for patients with LBP in the intervention ED (Objective 1); reduce inpatient admissions, improve early mobilisation, improve the provision of patient education, and reduce re-presentation to the intervention ED (Objective 2); and improve the cost of care from the hospital perspective (Objective 3), all through behavioural change in the clinical practice of ED clinicians (Objective 4). Furthermore, it will provide patients with the resources to better self-manage LBP, reducing recurrence and healthcare dependency.

It is planned to disseminate findings widely including engagement with relevant stakeholders regarding implementation of the intervention at respective sites locally (i.e., the control ED), nationally and internationally. This will be facilitated by planned publications (results paper, process evaluation paper) in open access journals, presentations at high impact ED and LBP conferences, stakeholder engagement via a national ED online symposium and workshops. This includes the provision of training on implementation of the intervention (consistent with the information delivered at the stakeholder engagement), engaging clinicians, and ensuring sustainability in change of practice. It is also planned to produce media resources that will permit the widespread dissemination of the study's findings.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements: The investigators would like to thank the clinical and administrative staff of the Emergency Departments at the participating EDs. We also thank the Physiotherapy Department and Jamieson Trauma Centre at the Royal Brisbane and Women’s Hospital.

Author Contributions: All authors (SO, JH, JW, TS, EB, GM, WS, PB, GM, MC, TC, NEF) contributed to the study design and preparation, data collection planning and process, intervention design and implementation, and data analysis planning. All authors contributed to and approved the final manuscript preparation.

Funding: This study has been funded by a Hospitals Contribution Fund (HCF) Research Foundation Translational Research Grant Program (Grant number: 21TRG3). This funding body had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Competing interests: None declared.

Patient consent for publication: Not required.

Ethics approval: This study received ethical approval from the Metro North Human Research Ethics Committee (HREC/2022/MNHA/87995). Subsequent site specific approval was then gained from each participating hospital site.

Provenance and peer review: Not commissioned; externally peer reviewed.

Open access: This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Erasmus Hogeschool

given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Emergency department care. Australian Institute of Health and Welfare (AIHW), Australian Government: Canberra, 2021.
- 2 Management of people with acute low back pain: model of care. NSW Agency for Clinical Innovation. NSW Health Chatswood, NSW, 2016. p. 39.
- 3 Machado GC, Rogan E, Maher CG. Managing non-serious low back pain in the emergency department: Time for a change? *Emerg Med Australas* 2018;30(2):279-82.
- 4 Makary MA, Overton HN, Wang P. Overprescribing is major contributor to opioid crisis. *BMJ* 2017;359:j4792.
- 5 Heine J, Window P, Hacker S, et al. Adherence to recommended guidelines for low back pain presentations to an Australian emergency department: Barriers and enablers. *Australas Emerg Care* 2023;26(4):326-32.
- 6 Coombs DM, Machado GC, Richards B, et al. Effectiveness of a multifaceted intervention to improve emergency department care of low back pain: a stepped-wedge, cluster-randomised trial. *BMJ Qual Saf* 2021;30(10):825-35.
- 7 Machado GC, Richards B, Needs C, et al. Implementation of an evidence-based model of care for low back pain in emergency departments: protocol for the Sydney Health Partners Emergency Department (SHaPED) trial. *BMJ Open* 2018;8(4):e019052.
- 8 Lopez Bernal J, Cummins S, Gasparrini A. The use of controls in interrupted time series studies of public health interventions. *Int J Epidemiol* 2018;47(6):2082-93.

- 9 Curran GM, Bauer M, Mittman B, et al. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care* 2012;50(3):217-26.
- 10 Health Service Strategy 2021-26. Available from: <https://metronorth.health.qld.gov.au/about-us/publications/health-service-strategy-planning>.
- 11 Brisbane North PHN and Metro North Hospital and Health Service. Population Health Report. Brisbane: Department of Health; 2019.
- 12 Penfold RB, Zhang F. Use of interrupted time series analysis in evaluating health care quality improvements. *Acad Pediatr* 2013;13(6 Suppl):S38-44.
- 13 Wagner AK, Soumerai SB, Zhang F, et al. Segmented regression analysis of interrupted time series studies in medication use research. *J Clin Pharm Ther* 2002;27(4):299-309.
- 14 Interrupted Time Series Power Calculation using DO Loop Simulations (sas.com).
- 15 Australian Commission on Safety and Quality in Health Care (ACSQHC). Low Back Pain Clinical Care Standard. Sydney: ACSQHC; 2022.
- 16 Graham ID, Logan J, Harrison MB, et al. Lost in knowledge translation: time for a map? *J Contin Educ Health Prof* 2006;26(1):13-24.
- 17 Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011;6:42.
- 18 Damschroder LJ, Banaszak-Holl J, Kowalski CP, et al. The role of the champion in infection prevention: results from a multisite qualitative study. *Qual Saf Health Care* 2009;18(6):434-40.
- 19 WHO Collaborating Centre for Drug Statistics Methodology. ATC/DDD Index. Oslo: Norwegian Institute of Public Health; 2023.

- 20 Proctor E, Silmere H, Raghavan R, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Adm Policy Ment Health* 2011;38(2):65-76.
- 21 Gale NK, Heath G, Cameron E, et al. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol* 2013;13:117.
- 22 The Lancet. Managing the opioid crisis in North America and beyond. *Lancet* 2022;399(10324):495.
- 23 Lalic S, Ilomaki J, Bell JS, et al. Prevalence and incidence of prescription opioid analgesic use in Australia. *Br J Clin Pharmacol* 2019;85(1):202-15.
- 24 Donovan PJ, Arroyo D, Pattullo C, et al. Trends in opioid prescribing in Australia: a systematic review. *Aust Health Rev* 2020;44(2):277-87.
- 25 Caughey GE, Gadzhanova S, Shakib S, et al. Concomitant prescribing of opioids and benzodiazepines in Australia, 2012-2017. *Med J Aust* 2019;210(1):39-40.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGENDS

Figure 1 Diagram depicting the 6 components of the implementation plan designed to change clinician behaviour towards the use of best-practice LBP management guidelines, specifically the NSW Agency for Clinical Innovation (ACI) model of care for acute LBP[2] and the more recent Australian LBP Clinical Care Standard.[15]

For peer review only

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Erasmus Hogeschool



Figure 1 Diagram depicting the 6 components of the implementation plan designed to change clinician behaviour towards the use of best-practice LBP management guidelines, specifically the NSW Agency for Clinical Innovation (ACI) model of care for acute LBP[2] and the more recent Australian LBP Clinical Care Standard.[15]

338x190mm (300 x 300 DPI)

Supplementary Table 1: Parameters for extracting potential Low Back Pain presentations using the Emergency Department (ED).

Code	Explanatory Description
Presenting Complaint	
17005	Requesting Investigation
17006	Requesting Medication/Certificate/Results
17008	Unplanned ED Representation
21002	Altered Sensation
25005	Back Pain
9047	Mechanical Strain
Primary ICD-10 AM Diagnosis Code	
G95.9	Disease of Spinal Cord, Unspecified
M13.99	Arthritis, Unspecified
M19.99	Arthrosis, Unspecified
M25.59	Pain in a Joint, Site Unspecified
M25.99	Unspecified Joint Disorder
M48.99	Unspecified Spondylopathy
M51.9	Intervertebral Disc Disorder
M53.3	Sacro Coccygeal Disorders
M53.99	Unspecified Dorsopathy
M54.19	Radiculopathy, Site Unspecified
M54.2	Cervicalgia
M54.3	Sciatica
M54.5	Low Back Pain
M54.6	Pain in Thoracic Spine
M54.99	Unspecified Dorsalgia Site Unspecified
M79.19	Myalgia, Site Unspecified
M79.29	Neuralgia and Neuritis, Unspecified
M89.99	Unspecified Disorder of Bone
M95.9	Acquired Musculo Skeletal Deformity
R25.2	Cramp and Spasm
R26.8	Abnormalities of Gait and Mobility
R29.89	Musculo Skeletal System Symptoms / Signs
R52.9	Pain, Unspecified
S22.00	Fracture of Thoracic Vertebra
S23.3	Sprain and Strain of Thoracic Spine
S32.00	Fracture of Lumbar Vertebra
S32.82	Fracture of Lumbosacral Spine
S33.3	Dislocation Lumbar Spine and Pelvis
S33.7	Sprain/Strain Lumbar Spine/Pelvis
S34.1	Other Injury of Lumbar Spinal Cord
S38.1	Crush Injury Abdomen/Lower Back/Pelvis
S39.9	Injury of Abdomen/Lower Back/Pelvis
Z02.7	Issue of Medical Certificate
Z04.9	Examination / Observation Unspecified
Z76.0	Issue of Repeat Prescription

Note: Codes and descriptions relate to the intervention ED location of FASTRACK and control ED location of BLUE or GREEN ZONE.

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.
Erasmus Hogeschool