BMJ Open

Has there been a change in the rates of sickness certification for back pain over time? An examination of historical data from 2000 to 2010

Journal:	BMJ Open
Manuscript ID	bmjopen-2015-009634
Article Type:	Research
Date Submitted by the Author:	04-Aug-2015
Complete List of Authors:	Wynne-Jones, Gwenllian; Arthritis Research UK Primary Care Centre, Primary Care Sciences Dunn, Kate; Arthritis Research UK Primary Care Centre, Primary Care Sciences
Primary Subject Heading :	Epidemiology
Secondary Subject Heading:	Epidemiology, Occupational and environmental medicine
Keywords:	EPIDEMIOLOGY, OCCUPATIONAL & INDUSTRIAL MEDICINE, Back pain < ORTHOPAEDIC & TRAUMA SURGERY



BMJ Open

Has there been a change in the rates of sickness certification for back pain over time? An examination of historical data from 2000 to 2010

Gwenllian Wynne-Jones*, Kate M Dunn

*Corresponding author Arthritis Research UK Primary Care Centre, Research Institute for Primary Care & Health Sciences, Keele University, Keele, Staffordshire, ST5 5BG Email; g.wynne-jones@keele.ac.uk Phone; 01782 733962

Key words; Back pain, Absence, Rate, Epidemiology Word count; 2759

ABSTRACT

Objectives

This paper aims to investigate historical patterns of sickness certification for back pain from 2000 to 2010.

Design

Electronic medical records from 14 practices that are part of the NIHR Clinical Research Network: West Midlands were reviewed. All records for back pain consultations from 2000 to 2010 were downloaded and matched, by date, to corresponding sickness certification records.

Setting

Primary Care

Results

A total of 93896 back pain consultations were recorded over the eleven year period resulting in 30913 sickness certificates. There was a statistically significant decrease in the rate of certification over the period, falling from 376.8 (95% CI 362.1, 392) per 1000 back pain consultations in 2000 to 246.5 (95% CI 236.5, 332.9) per 1000 back pain consultations in 2010. There was also a statistically significant difference in certification between males and females, with males issued more certificates than females. There was a statistically significant difference in certification by age, with those aged 60 years and over being less likely to be issued a certificate compared to all other age groups.

Conclusions

Rates of sickness certification for back pain demonstrated a downward trend between 2000 and 2010. Whilst the reasons for this are not transparent, it may be related to changing beliefs around working with back pain.

Strengths and limitations of this study

- This medical record review provides the ability to examine the rate of back pain associated sickness certificates over an eleven year period in all adults registered at 14 general practices.
- The database used has previously been demonstrated to be generalisable to the age and gender of the UK population.

BMJ Open

• The reasons for the decrease in sickness certification are not fully clear but may be related to changing beliefs around working with back pain.

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

INTRODUCTION

Low back pain is a common reason for consulting in primary care, with 38% of adults affected in any one year at an estimated cost to the NHS of £1 billion per annum.^[1] Furthermore, musculoskeletal pain, principally back pain, is one of the most common reasons for absence from work.^[2]

The costs of illness to employers, particularly back pain, are large; it has been estimated that 31 million days of absence were taken as a result of musculoskeletal pain in 2013.^[3] The Chartered Institute of Personnel and Development^[4] reports the rate of UK sickness absence in terms of a percentage of lost working hours, estimating sickness absence as 3.0%, 3.4% and 3.2% of working hours in 2012, 2011 and 2010 respectively. The Health and Safety Executive^[5] have estimated the total net cost of sick pay to be £14.2 billion during the year 2012-2013 as a result of illness and workplace injuries. When it is considered that musculoskeletal pain, principally back pain, is one of the most common reasons for absence from work it can be assumed that a large proportion of this sick pay is as a result of back pain.^[4] In the UK, proxy measures are used to estimate rates of sickness certification based on work absence. However, reporting rates of absence as a percentage of lost working hours does not reflect the clinical issues associated with certification in terms of the numbers of individuals absent from the workplace, nor does it represent the number of consultations for certification in general practice.^[6,7]

The health service costs and lost capacity in the workplace has made health and work a key target for public policy in the UK.^[8] The Government is actively aiming to reduce the number of employees signed off sick each year with a multi-agency government programme, launched in 2005, to address the issues of health, work and wellbeing.^[9] Furthermore, in 2010 the Fit Note was introduced to replace sickness certificates. The aim of the Fit Note was to change the focus from what the patient can't do to what they can do in relation to work and to provide the GP with the option to state that patients may be fit for some work with provisions; a phased return to work, altered hours, amended duties and workplace adaptations. However, in order to assess the impact of such initiatives, methods are required to estimate rates of sickness certification over time, including the periods before such initiatives were implemented.

Page 4 of 19

BMJ Open

The National Institute for Health Research Clinical Research Network: West Midlands holds frozen archive data on sickness certification in its Medical Certificates in Primary Care Archive and consultation data in its Consulters in Primary Care Archive. These databases have been validated for assessment of sickness certification,^[10] and the rates of sickness certification for a range of health conditions have been estimated.^[11] However, it is unknown whether the rate of certification for back pain has changed over time or whether there are any trends over time by age and gender. This paper aims to investigate historical patterns of sickness certification for back pain from 2000 to 2010.

METHODS

All consultation records from 2000 to 2010 for individuals with a back pain Read code were downloaded from the Consultation in Primary Care Archive database (CiPCA). Read codes are a hierarchy of morbidity, symptom and process codes, which become more specific further down the hierarchy. A set of Read codes has been established to identify back pain consistently from the medical records, this comprises a total of 589 Read codes plus 20 other terms.^[12] It was these codes that were used to identify patients for this study. Records for working age adults (aged 19-64 years), and those patients who were registered at the practice for the full year included in the analysis, were eligible. Each consultation record includes the unique individual identification number and practice identification number, plus age, gender, and year of consultation, date of consultation, the Read Code which was used to identify the problem with which the patient consulted, and the consultation free text. All sickness certification records between 2000 and 2010 for the same individuals, identified using each patient's unique identification number, were downloaded from the Medical Certificates in Primary Care Archive database (MiPCA). In the UK, a sickness certificate is required from the seventh day of absence. Each sickness certification record includes a unique individual identification number and practice identification number, plus age, gender, year of issue, date of issue, and the Read Code used to identify the record of a sickness certificate. Sickness certification records were then matched to back pain consultation records using the date of issue/consultation, the patient's unique identification number, plus age and gender.

Analysis

Numbers of consultations for back pain each year, and numbers of sickness certificates issued, were calculated using SPSS version 21. The rate of certification was defined as the number of certificates issued for back pain divided by the number of consultations for back pain in each year, presented per 1000 back pain consultations. The crude rate of certification and the rates by age and gender were calculated with 95% Confidence Intervals (CIs). A *t*-test for differences in proportions was calculated for year of certification and gender. An Analysis of Variance (ANOVA) was calculated to examine differences in rates by age.

Ethical approval

North Staffordshire Research Ethics Committee (LREC Project 03/04).

RESULTS

During the 11 year period (2000-2010) there were a total of 93896 consultations for back pain, resulting in 30913 sickness certificates being issued. This gives a sickness certification prevalence of 32.9%; or a third of all consultations for back pain resulting in a sickness certificate. The overall rate of certification was 329.2 certificates per 1000 consultations (95% CI 325.6, 332.9) (table 1 and figure 1).

There appears to be a downward trend in the rate of certification over the study period, with the rate of certification falling from 376.8 (95% CI 362.1, 392) per 1000 back pain consultations in 2000, to 246.5 (95% CI 236.5, 332.9) per 1000 back pain consultations in 2010 (see Figure 1). The rate of certification reaches a peak of 424.8 (95% CI 410.2, 439.7) per 1000 back pain consultations during the year 2003, with another spike in 2006. This decrease in the rate of certification over the eleven year period was statistically significantly (p = <0.001).

BMJ Open

Table 1: Rate of sickness certification per 1000 consultations for back pain, by year and gender

				Sickness certification rate	e per 1000 back pain cons	sultations (95 % CI)
Year	Total number of	Total number of	Prevalence (%)	Total	Males	Females
	certificates issued	consultations				
2000	2455	6515	37.7	376.8 (362.1, 392)	397.1 (375.2,419.9)	358.4 (338.5, 379)
2001	2809	6926	40.5	405.6 (390.7, 420.9)	484.1 (460.9, 508.2)	331.5 (312.8, 350.9)
2002	2863	6801	42.1	420.1 (405.7, 436.7)	473.7 (450.6, 497.7)	370.6 (350.7, 391.4)
2003	3215	7569	42.5	424.8 (410.2, 439.7)	471.4 (449,494.6)	384.1 (365.3, 403.7)
2004	3060	7476	40.9	409.3 (394.9, 424.1)	458.2 (436.5, 480.6)	362.6 (436.5, 480.6)
2005	2720	9338	29.1	291.3 (280.4, 302.4)	347.7 (330.3, 365.8)	242.9 (229.4, 256.9)
2006	3495	9442	37.0	370.1 (357.9382.6)	394.7 (376.3, 413.8)	349 (333, 365.7)
2007	2653	10200	26.4	260.1 (250.3, 270.2)	281.8 (266.6, 297.8)	242.9 (230.2, 256)
2008	2636	10046	26.2	262.4 (252.5, 272.6)	296.1 (280.6, 312.3)	233.7 (221, 246.9)
2009	2731	10349	26.4	263.9 (254.1, 2739)	293.4 (278.1, 309.4)	239.5 (226.9, 252.6)
2010	2276	9234	24.6	246.5 (236.5, 256.8)	275.7 (259.9, 292.3)	223.3 (210.6, 236.6)
All years	30913	93896	32.9	329.2(325.6, 332.9)	370.5 (364.8, 376.3)	293.5 (288.8, 298.3)

Page 7 of 19

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

Examining the data by gender demonstrates that the rate of certification associated with back pain is slightly higher in males compared to females (figure 2). The rate of certification reaches a peak of 484.4 (95% CI 460.9, 508.2) certificates per 1000 back pain consultations in males in 2001 compared to 384.1 (95% CI 365.2, 403.7) certificates per 1000 back pain consultations in females in 2003, falling to a low of 275.7 (95% CI 259.9, 292.3) certificates per 1000 back pain consultations in males during 2010 compared to 223.3 (95% CI 210.6, 236.6) certificates per 1000 consultations in females during 2010 (figure 2). There was a statistically significant difference in the rate of sickness certification between males and females (p = < 0.001).

Comparing the rate of certification per 1000 back pain consultations by age shows that no single age group is reliably recording a higher rate of certification than other groups. However, the 60+ age group consistently record much lower rates of certificates per 1000 back pain consultations compared to the other age groups (figure 3). An ANOVA demonstrates that this difference in rate of certification between the 60+ age group and the other age groups is significant p = <0.001; excluding the 60+ age group from the ANOVA demonstrates that there are no statistical differences in the rate of certification between the age groups from 19 to 59 years.

DISCUSSION

Overall, the rate of sickness certification for back pain has significantly declined between 2000 and 2010. This decline in rates has principally occurred from 2003 onwards. This study demonstrated that approximately one third of consultations for back pain will lead to a sickness certificate; this finding is in line with many studies examining back pain which also find that approximately one third of participants are absent from work.^[13,14]

There was a trend in certification by gender, with men reporting consistently higher rates of certification than women. This is to be expected and has been reported in previous studies of certification using this dataset.^[11,15] The most likely explanation is the increased proportions of males working in the manual sector at 34.6% when compared to females at 10.9%.^[16] On average, manual workers report more absence than non-manual employees, and also report more absence as a result of

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

musculoskeletal conditions than non-manual employees.^[4] The differences in certification for gender may also be associated with differing consultation behaviours, and it has been reported that men are more likely to consult with an episode of back pain than women at a female:male rate ratio of 1:20 (95% Confidence Interval 1.13, 1.28).^[12]

There was no clear trend in certification by age with the exception of the 60+ years age group, who consistently had a lower rate of certification compared to younger age groups. Again, the rate of certification in this group showed a downward trend but it was not as pronounced as in other groups. There are a number of reasons why this difference may be seen in the 60+ age group. First, there are fewer people employed in this age group, in 2011 the Organisation for Economic Co-operation and Development reported that 56% of the population in the 55-64 years age group were employed compared to 80% of the population in the 25-54 years age group¹⁷. Second, there could be a healthy worker effect whereby individuals suffering with back pain are removing themselves from the workplace. Findings are not likely to be related to differential reporting, as reporting of back pain over time is likely to be consistent.^[18]

It is difficult to identify the reason for the decreasing rate of sickness certification in this study. It could be argued that the prevalence of back pain has decreased over time and so there is less need for certification. However, the literature suggests that the prevalence of back pain has remained largely unchanged over the period 1990-2010^[19] and may even have increased.^[20,21] During the period of analysis that this paper spans there have been a number of initiatives, both in the UK and worldwide, surrounding the management of back pain that *may* have contributed to the decline in rates of certification. A review of clinical guidelines for the management of nonspecific back pain assessed guidelines published between 2000 and 2008;^[22] this review is an update of a previous review.^[23] The authors report that the most common advice is to reassure patients and encourage them to remain active, however, in contrast to the earlier review the current guidelines increasingly mention return to work, despite back pain, in their recommendations. In the UK, 2009 saw the publication of two National Institute for Clinical Excellence (NICE) reports, the first considering early management of non-specific back pain^[24] and the second reporting on primary care management of long term sickness absence.^[25] The non-specific back

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

pain guidelines^[24] concur with the review by Koes *et al*^[22] that individuals should be encouraged to remain active and continue normal activities as far as possible. The second report is focussed on promoting the benefits of working with health conditions,^[23] specifically back pain, and again encourages individuals to maintain work despite pain. However, the impact of these recent reports is unlikely to be seen in the current analysis.

Recent research from the Global Burden of Disease study has demonstrated that back pain leads to more years lived with disability than any other condition and this burden of back pain is increasing with an ageing population.^[19] There is also evidence that the number of consultations for back pain has remained static, at least in the population of the United States, for a period of over 10 years from the 1990s to the mid-2000s.^[26] It is interesting then to note that despite the relative stability of consultations for back pain and the relatively adverse outcomes in terms of the burden of back pain that the number of sickness certificates for back pain has decreased in this study. Within the UK there have not been any recent public health initiatives which may account for this decrease in certification; there has however been the sea change in the information that general practitioners are advised to give their patients in regards to working with pain. There is consistent information provided by the Royal College of General Practitioners,^[27] Department for Work and Pensions^[28] and NICE^[24,25] coupled with the availability of information booklets detailing the management of health and work for the GP, patient and employer.^[29,30,31,32] It may be that media campaigns to promote working with musculoskeletal pain could see the rate of sickness certification reduce further. There has been some success with media campaigns in Scotland and Australia demonstrating that positive messages around working with musculoskeletal pain led to improvements in knowledge, attitudes and beliefs at the population level which were maintained, albeit at a reduced level, three years later.^[33,34,35] A Canadian study also utilising a media campaign had a more limited effect on behaviours related to back pain e.g healthcare use, than those run in Scotland and Australia.^[36] However, the Canadian study did find that participants agreement with the statement "if you have back pain you should try to stay active" significantly increased.^[36] It seems then that knowledge does not necessarily translate into behaviour and it has also been reported that provision of information alone is not sufficient to prevent work absence,^[37] but it seems that improving the baseline

Page 10 of 19

BMJ Open

There are a number of limitations to the current study. Firstly, capturing the duration of sickness absence is not possible in the current dataset, meaning it is unclear whether the data are unduly influenced by a large number of short or long term certificates. The most frequently recorded duration for absence is two weeks^[39,40,41,42] and there is no reason to believe that GPs in the current study would differ to any great degree. A second consideration is that this dataset is based in one area of the UK, North Staffordshire, and it could be argued that it is not generalizable to the rest of the population. Previous work with this dataset has demonstrated that crude rates of certification change very little when the data are standardised to the age and gender of the population as a whole, and there is no indication that this should be any different for this study.^[11] Lastly, the data included in the manuscript only goes as far as 2010 as from this date onwards the Read coding for sickness certificates changed as a result of the introduction of the Fit Note. This change in Read coding means that Fit Notes are now coded as "not fit for work" and "may be fit for some work". As this classification is different from the pre-2010 sickness certificates it would make comparison of the data between these two periods difficult in this manuscript because we would not be comparing like with like. As a result we could not be confident that any changes in the rate of certification would be as a result of the change to the Fit Note system or as a result of the change in Read coding i.e. more read codes available to classify not fit for work and may be fit for some work.

The main strength of this study is the large database in which this work was carried out. This dataset allowed the linking of consultation data to certification data to enable trends by age and gender to be seen. This is the first study to map trends in certification for back pain in the UK using an established dataset which has been validated for use in examining both consultation and sickness certification data.^[43,10] There is a need for establishing baseline rates of certification against which any change in policy or strategy at either local or national level can be compared,^[44] and this dataset goes some way to establishing these baseline figures. Linaker *et al*^[44] state that improvement of existing datasets would be more attractive than the development of a new national system to record sickness absence; with the introduction of the e- Fit

Page 11 of 19

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

Note the current data set is being updated to include information on duration of absence and whether or not a patient may be fit for work, further strengthening this dataset for future research. Another strong point of this study is that is appears to be supported by other literature; this lends credibility to our findings. For example, Ruseckaite *et* al^[45] reported a significant decrease in certification for musculoskeletal disorders between 2003 and 2010 in Australia. They also report a decrease in rate of certification for back pain and strains although there is no indication of whether this trend is significant or not. Lambeek *et al*^[46] reported that the costs of back pain in the Netherlands decreased between 2002 and 2007, acknowledging that a large proportion of this cost is made up of absence. In the United Kingdom the Office for National Statistics have reported a decrease in reported absence across all health conditions between 1993 and 2013.^[47] However, work still needs to be carried out to investigate why there has been this change, and whether it is truly related to the available evidence about working with pain.

Conclusions

Rates of certification for back pain demonstrated a significant downward trend over the period 2000 to 2010; the reasons for this are not fully transparent but may be as a result of changing beliefs around working with back pain. These findings may provide a benchmark against which the impact of public health initiatives may be evaluated and monitored. With the new recording of the e-Fit Note this dataset will become more useful in tracking rates of certification over time.

ACKNOWLEDGEMENTS

We would like to thank the NIHR Clinical Research Network: West Midlands, and Ian Thomas, Simon Wathall and Kelvin Jordan for their assistance with data retrieval.

COMPETING INTERESTS

All authors have completed the ICMJE uniform disclosure form at <u>www.icmje.org/coi_disclosure.pdf</u> and declare: all authors had financial support from the British Occupational Health Research Foundation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

FUNDING

This paper presents independent research funded by the National Institute for Health Research (NIHR), under its Programme Grants for Applied Research funding scheme: "Optimal management of spinal pain and sciatica in primary care" (NIHR-RP-PG-0707-10131). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

GW-J is funded by an NIHR Research Post-doctoral Fellowship (PDF-2009-02-54).

ETHICAL APPROVAL

Approval for use of the Medical Certificates and Consultation databases was gained from the North Staffordshire Research Ethics Committee (LREC Project 03/04)

CONTRIBUTORSHIP

GW-J and KD contributed to the design of the study, interpreting the analysis and writing of the manuscript. GW-J conducted the analysis.

DATA SHARING STATEMENT

Researchers wanting to apply for access to individual patient data from archived studies hosted by the Keele Research Institute for Primary Care and Health Sciences

Page 13 of 19

should complete an External Request Form and send an electronic version of the form, together with an outline design of the proposed investigation and a short CV for all study team members to – primarycare.datasharing@keele.ac.uk.

BMJ Open

REFERENCES

- Maniadakis N, and Gray A, The economic burden of back pain. *Pain* 2000;84(1):95-103
- CBI Fit for Purpose: Absence and Workplace Health Survey 2013, http://www.cbi.org.uk/media/2150120/cbi-

pfizer_absence___workplace_health_2013.pdf (accessed 31st March 2015)

- Office For National Statistics, Statistical Bulletin: Labour Market Statistics 2014 http://www.ons.gov.uk/ons/rel/lmac/sickness-absence-in-the-labourmarket/2014/rpt---sickness-absence-in-the-labour-market.html (Accessed 31st March 2015)
- The Chartered Institute for Personnel and Development, Absence Management 2012, http://www.cipd.co.uk/binaries/absence-management_2012.pdf (Accessed 31st March 2015)
- Health and Safety Executive, Health and Safety Statistics, Annual Report for Great Britain (2013/2014), http://www.hse.gov.uk/statistics/overall/hssh1314.pdf (Accessed 31st March
- 6. Hiscock J, Ritchie J. The role of GPs in sickness certification. Leeds: Report number 148, Her Majesties Stationary Office 2001
- Hussey S, Hoddinott P, Wilson P, *et al.* Sickness certification system in the United Kingdom: qualitative study of views of general practitioners in Scotland. *BMJ* 2004;328:88
- Department of Health, Health and Safety Executive, Department for Work and Pensions. Health, work and well-being - Caring for our future. London: Department for Work and Pensions 2005
- Health work and wellbeing. Working for a healthier tomorrow 2008 https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/ 209782/hwwb-working-for-a-healthier-tomorrow.pdf (Accessed 31st March 2015)
- Wynne-Jones G, Mallen C.D, Main C.J. *et al.* Sickness certification in general practice: A comparison of electronic records with self-reported work absence. *Prim Health Res Dev* 2008;9:113-118

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

- Wynne-Jones G, Mallen C.D, Mottram S. *et al.* The identification of U.K. sickness certification rates, standardised for age and sex. *Br J Gen Pract* 2009;59(564):510-515
- Jordan K.P, Kadam U.T, Hayward R, *et al.* Annual consultation prevalence of regional musculoskeletal problems in primary care: an observational study. *BMC Musculoskelet Disord* 2010;11:144
- 13. Urquhart D.M, Kelsall H.L, Hoe V.C. *et al.* Are psychosocial factors associated with low back pain and work absence for low back pain in an occupational cohort?. *Clin J Pain* 2013;29(12):1015-1020
- Wynne-Jones G, Cowen J, Jordan J.L, *et al.* Absence from work and return to work in people with back pain: a systematic review and meta-analysis. *Occup Environ Med* 2014:71(6):448-456
- Wynne-Jones G, Mallen C.D, Dunn K.M. Sickness certification for musculoskeletal conditions. *Clin Rheumatol* 2010;29(5):573-574
- 16. OfficeforNationalStatistics2013http://www.ons.gov.uk/ons/dcp171778_332467.pdf(Accessed 31st March2015)
- 17. Organisation for Economic Co-operation and Development 2013 http://www.oecd-ilibrary.org/sites/factbook-2013en/07/01/02/index.html?contentType=&itemId=/content/chapter/factbook-2013-52-

en&containerItemId=/content/serial/18147364&accessItemIds=&mimeType=t ext/html (Accessed 31st March 2015)

- Lemeunier N, Leboeuf-Yde C, Kjaer P, et al. Stability of low back pain reporting over 8 years in a general population aged 40/41 years at base-line: data from three consecutive cross-sectional surveys. BMC Musculoskelet Disord 2013;14:270
- 19. Hoy D, March L, Brooks P, et al. The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. Ann Rheum Dis 2014;73:968-974
- 20. Großschädl F, Freidl W, Rásky E, *et al.* A 35-Year Trend Analysis for Back Pain in Austria: The Role of Obesity. *PLOS One* 2014;9(9):e107436

BMJ Open

2	of chronic back pain: secular trend analysis of longitudinal medical
	expenditures panel survey data for years 2000 to 2007. J Manipulative Physiol
	<i>Ther</i> 2013:36(1):2-11
2	2. Koes B.W, van Tulder M.W, Lin C-W.C, et al. An updated overview of
	clinical guidelines for the management of non-specific low back pain in
	primary care. Eur Spine J 2010;19:2075-2094
2	3. Koes B.W, van Tulder M.W, Ostelo R, et al. Clinical guidelines for the
	management of low back pain in primary care: an international comparison.
	Spine 2001;26:2504–2513
2	4. National Institute for Health and Care Excellence, Early Management of
	Persistent non-specific low back pain 2009a
2	5. National Institute for Health and Care Excellence, Managing long-term
	sickness and incapacity for work 2009b
2	6. Deyo R.A, Mirza S.K, Martin B.I, Back Pain Prevalence and Visit Rates
	Estimates From U.S. National Surveys. Spine 2002;31(23):2724-2727
2	7. Royal College of General Practitioners, Health and Work in General Practice
	http://www.rcgp.org.uk/courses-and-events/health-and-work-in-general-
	practice-events.aspx (Accessed 31st March 2015)
2	8. Department for Work and Pensions, Royal College of General Practitioners,
	British Medical Association. Statement of fitness for work: a guide for general
	practitioners and other doctors. London: Department for Work and Pensions
	2010
2	9. Waddell G, Burton AK. Work and Health. London: The Stationary Office,
	2006
3	0. Waddell G, Burton AK. Health and Work. London: The Stationary Office,
	2007a
3	1. Waddell G, Burton AK. Advising Patients about Work. London: The
	Stationary Office, 2007b
3	2. Kendall NAS, Burton AK, Main CJ, et al, on behalf of the Flags Think-Tank:
	Tackling Musculoskeletal Problems: a Guide for the Clinic and Workplace -
	Identifying Obstacles Using the Psychosocial Flags Framework 2009, London:
	The Stationary Office
	Page 17 of 19

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

- 33. Waddell G, O'Connor M , Boorman S. *et al.* Working Backs Scotland: a public and professional health education campaign for back pain. *Spine* 2007c;32:2139–43
- Buchbinder R , Jolley D, Wyatt M . Population based intervention to change back pain beliefs and disability: three part evaluation. *BMJ* 2001;322:1516–20
- 35. Buchbinder R , Jolley D. Population based intervention to change back pain beliefs: three year follow up population survey. *BMJ* 2004;328-321
- 36. Gross D.P, Russell A.S, Ferrari R, Battié M.C, et al. Evaluation of a Canadian back pain mass media campaign. Spine 2010;35(8):906-913
- 37. Henrotin YE, Cedraschi C, Duplan B, *et al.* Information and low back pain management: a systematic review. *Spine* 2006;31:E326–E334.
- 38. Lorig K.R , Laurent D.D , Deyo R.A, et al. Can a Back Pain E-mail Discussion Group improve health status and lower health care costs?: A randomized study. Arch Intern Med 2002;162(7):792–6
- 39. Head J, Ferrie J. E, Alexanderson K, *et al.* Diagnosis-specific sickness absence as a predictor of mortality: the Whitehall II prospective cohort study. *BMJ* 2008;337:a1469
- 40. Tellnes G. Days lost by sickness certification. Scand J Prim Health Care 1989;7(4):245–251.
- 41. Reiso H, Gulbrandsen P, Brage S. Doctors' prediction of certified sickness absence. *Fam Pract* 2004;21(2):192–198.
- 42. Bollag U, Rajeswaran A, Ruffieuxf C, *et al.* Sickness certification in primary care—the physician's role. *Swiss Med Wkly* 2007; 137(23–24): 341–346
- 43. Porcheret M, Hughes R, Evans D. *et al.* Data quality of general practice electronic health records: the impact of a program of assessments, feedback, and training. *J Am Med Inform Assoc* 2004;11:78–86
- 44. Linaker C, Harris E.C, Cooper C, *et al.* The burden of sickness absence from musculoskeletal causes in Great Britain. *Occup Med* 2011;61(7):458-464
- 45. Ruseckaite R, Collie A, Bohensky M. *et al*, Trends in Sickness Certification of Injured Workers by General Practitioners in Victoria, Australia. J Occup Rehabil 2014:24:525-532
- 46. Lambeek L.C, van Tulder M.W, Swinkles I.C.S *et al.* The trend in total cost of back pain in the Netherlands in the period 2002 to 2007. *Spine* 2011;36(13):1050-1058

Page 18 of 19

1				
2	47 000	C	NT. (******1	Ct. () (
3	4/. Office	for	National	Statistics
4 5	http://www.ons.gov.	uk/ons/dcp171776	353899.pdf (Accessed	31st March
5	1		1	
7	2015)			
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				
19				
20				
22				
23				
24				
25				
26				
27				
28				
29				
31				
32				
33				
34				
35				
36				
37				
38				
39 40				
40				
42				
43				
44				
45				
46				
4/				
40 70				
51				
52				
53				
54				
55				
56				
5/				
50 50				
59 60		Page 10 of	19	
00		1 ugo 17 01	17	
	For peer review only -	http://bmjopen.bmj	j.com/site/about/guideli	nes.xhtml



Figure 1: Rates of sickness certification associated with back pain per 1000 back pain consultations

210x297mm (300 x 300 DPI)





210x297mm (300 x 300 DPI)



Figure 3: Rate of certification associated with back pain per 1000 back pain consultations by age group

210x297mm (300 x 300 DPI)

BMJ Open

Has there been a change in the rates of UK sickness certification for back pain over time? An examination of historical data from 2000 to 2010

Journal:	BMJ Open
Manuscript ID	bmjopen-2015-009634.R1
Article Type:	Research
Date Submitted by the Author:	16-Dec-2015
Complete List of Authors:	Wynne-Jones, Gwenllian; Arthritis Research UK Primary Care Centre, Primary Care Sciences Dunn, Kate; Arthritis Research UK Primary Care Centre, Primary Care Sciences
Primary Subject Heading :	Epidemiology
Secondary Subject Heading:	Epidemiology, Occupational and environmental medicine, General practice / Family practice
Keywords:	Back pain < ORTHOPAEDIC & TRAUMA SURGERY, Rate, Absence, EPIDEMIOLOGY



BMJ Open

Has there been a change in the rates of UK sickness certification for back pain over time? An examination of historical data from 2000 to 2010

Gwenllian Wynne-Jones*, Kate M Dunn

*Corresponding author Arthritis Research UK Primary Care Centre, Research Institute for Primary Care & Health Sciences, Keele University, Keele, Staffordshire, ST5 5BG Email; g.wynne-jones@keele.ac.uk Phone; 01782 733962

Key words; Back pain, Absence, Rate, Epidemiology Word count; 2759

ABSTRACT

Objectives

This paper aims to investigate historical patterns of sickness certification for back pain from 2000 to 2010.

Design

Electronic medical records from 14 practices that are part of the NIHR Clinical Research Network: West Midlands were reviewed. All records for back pain consultations from 2000 to 2010 were downloaded and matched, by date, to corresponding sickness certification records.

Setting

Primary Care

Results

A total of 93896 back pain consultations were recorded over the eleven year period resulting in 30913 sickness certificates. There was a statistically significant decrease in the rate of certification over the period, falling from 376.8 (95% CI 362.1, 392) per 1000 back pain consultations in 2000 to 246.5 (95% CI 236.5, 332.9) per 1000 back pain consultations in 2010. There was also a statistically significant difference in certification between males and females, with males issued more certificates than females. There was a statistically significant difference in certification by age, with those aged 60 years and over being less likely to be issued a certificate compared to all other age groups.

Conclusions

Rates of sickness certification for back pain demonstrated a downward trend between 2000 and 2010. Whilst the reasons for this are not transparent, it may be related to changing beliefs around working with back pain.

Strengths and limitations of this study

- This medical record review provides the ability to examine the rate of back pain associated sickness certificates over an eleven year period in all adults registered at 14 general practices.
- The database used has previously been demonstrated to be generalisable to the age and gender of the UK population.

BMJ Open

• The reasons for the decrease in sickness certification are not fully clear but may be related to changing beliefs around working with back pain.

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

INTRODUCTION

Low back pain is a common reason for consulting in primary care, with 38% of adults affected in any one year at an estimated cost to the NHS of £1 billion per annum.^[1] Furthermore, musculoskeletal pain, principally back pain, is one of the most common reasons for absence from work.^[2]

The costs of illness to employers, particularly back pain, are large; it has been estimated that 31 million days of absence were taken as a result of musculoskeletal pain in 2013.^[3] The Chartered Institute of Personnel and Development^[4] reports the rate of UK sickness absence in terms of a percentage of lost working hours, estimating sickness absence as 3.0%, 3.4% and 3.2% of working hours in 2012, 2011 and 2010 respectively. The Health and Safety Executive^[5] have estimated the total net cost of sick pay to be £14.2 billion during the year 2012-2013 as a result of work-related illness and injuries. When it is considered that musculoskeletal pain, principally back pain, is one of the most common reasons for absence from work it can be assumed that a large proportion of this sick pay is as a result of back pain.^[4] In the UK, proxy measures are used to estimate rates of sickness certification based on work absence. However, reporting rates of absence as a percentage of lost working hours does not reflect the clinical issues associated with certification in terms of the numbers of individuals absent from the workplace, nor does it represent the number of consultations for certification in general practice.^[6,7]

The health service costs and lost capacity in the workplace has made health and work a key target for public policy in the UK.^[8] The Government is actively aiming to reduce the number of employees signed off sick each year with a multi-agency government programme, launched in 2005, to address the issues of health, work and wellbeing.^[9] Furthermore, in 2010 the Fit Note was introduced to replace sickness certificates. The aim of the Fit Note was to change the focus from what the patient can't do to what they can do in relation to work and to provide the GP with the option to state that patients may be fit for some work with provisions; a phased return to work, altered hours, amended duties and workplace adaptations. However, in order to assess the impact of such initiatives, methods are required to estimate rates of sickness certification over time, including the periods before such initiatives were implemented.

BMJ Open

The National Institute for Health Research Clinical Research Network: West Midlands holds frozen archive data on sickness certification in its Medical Certificates in Primary Care Archive and consultation data in its Consulters in Primary Care Archive. These databases have been validated for assessment of sickness certification,^[10] and the rates of sickness certification for a range of health conditions have been estimated.^[11] However, it is unknown whether the rate of certification for back pain has changed over time or whether there are any trends over time by age and gender. This paper aims to investigate historical patterns of sickness certification for back pain from 2000 to 2010.

METHODS

All consultation records from 2000 to 2010 for individuals with a back pain Read code were downloaded from the Consultation in Primary Care Archive database (CiPCA). Read codes are a hierarchy of morbidity, symptom and process codes, which become more specific further down the hierarchy. A set of Read codes has been established to identify both back pain diagnoses and symptoms consistently from the medical records, this comprises a total of 589 Read codes plus 20 other terms.^[12] It was these codes that were used to identify patients for this study. Records for working age adults (aged 19-64 years), and those patients who were registered at the practice for the full year included in the analysis, were eligible. Each consultation record includes the unique individual identification number and practice identification number, plus age, gender, and year of consultation, date of consultation, the Read Code which was used to identify the problem with which the patient consulted, and the consultation free text. All sickness certification records between 2000 and 2010 for the same individuals, identified using each patient's unique identification number, were downloaded from the Medical Certificates in Primary Care Archive database (MiPCA). In the UK, a sickness certificate is required from the seventh day of absence. Each sickness certification record includes a unique individual identification number and practice identification number, plus age, gender, year of issue, date of issue, and the Read Code used to identify the record of a sickness certificate. Sickness certification records were then matched to back pain consultation records using the date of issue/consultation, the patient's unique identification number, plus age and gender.

Page 5 of 19

Analysis

Numbers of consultations for back pain each year, and numbers of sickness certificates issued, were calculated using SPSS version 21. The rate of certification was defined as the number of certificates issued for back pain divided by the number of consultations for back pain in each year, presented per 1000 back pain consultations. The crude rate of certification and the rates by age and gender were calculated with 95% Confidence Intervals (CIs). A *t*-test for differences in proportions was calculated for year of certification and gender. An Analysis of Variance (ANOVA) was calculated to examine differences in rates by age. For all calculations the significance level was set at p=0.001 and a Bonferroni correction was applied to account for multiple analyses in the ANOVA.

Ethical approval

North Staffordshire Research Ethics Committee (LREC Project 03/04).

RESULTS

During the 11 year period (2000-2010) there were a total of 93896 consultations for back pain, resulting in 30913 sickness certificates being issued. This gives a sickness certification prevalence of 32.9%; or a third of all consultations for back pain resulting in a sickness certificate. The overall rate of certification was 329.2 certificates per 1000 consultations (95% CI 325.6, 332.9) (table 1 and figure 1).

There appears to be a downward trend in the rate of certification over the study period, with the rate of certification falling from 376.8 (95% CI 362.1, 392) per 1000 back pain consultations in 2000, to 246.5 (95% CI 236.5, 332.9) per 1000 back pain consultations in 2010 (see Figure 1). The rate of certification reaches a peak of 424.8 (95% CI 410.2, 439.7) per 1000 back pain consultations during the year 2003, with another spike in 2006. This decrease in the rate of certification over the eleven year period was statistically significantly (p = <0.001).

Table 1: Rate of sickness certification per 1000 consultations for back pain, by year and gender

				Sickness certification rate	e per 1000 back pain cons	ultations (95 % CI)
Year	Total number of	Total number of	Prevalence (%)	Total	Males	Females
	certificates issued	consultations				
2000	2455	6515	37.7	376.8 (362.1, 392)	397.1 (375.2,419.9)	358.4 (338.5, 379)
2001	2809	6926	40.5	405.6 (390.7, 420.9)	484.1 (460.9, 508.2)	331.5 (312.8, 350.9)
2002	2863	6801	42.1	420.1 (405.7, 436.7)	473.7 (450.6, 497.7)	370.6 (350.7, 391.4)
2003	3215	7569	42.5	424.8 (410.2, 439.7)	471.4 (449,494.6)	384.1 (365.3, 403.7)
2004	3060	7476	40.9	409.3 (394.9, 424.1)	458.2 (436.5, 480.6)	362.6 (436.5, 480.6)
2005	2720	9338	29.1	291.3 (280.4, 302.4)	347.7 (330.3, 365.8)	242.9 (229.4, 256.9)
2006	3495	9442	37.0	370.1 (357.9382.6)	394.7 (376.3, 413.8)	349 (333, 365.7)
2007	2653	10200	26.4	260.1 (250.3, 270.2)	281.8 (266.6, 297.8)	242.9 (230.2, 256)
2008	2636	10046	26.2	262.4 (252.5, 272.6)	296.1 (280.6, 312.3)	233.7 (221, 246.9)
2009	2731	10349	26.4	263.9 (254.1, 2739)	293.4 (278.1, 309.4)	239.5 (226.9, 252.6)
2010	2276	9234	24.6	246.5 (236.5, 256.8)	275.7 (259.9, 292.3)	223.3 (210.6, 236.6)
All years	30913	93896	32.9	329.2(325.6, 332.9)	370.5 (364.8, 376.3)	293.5 (288.8, 298.3)

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies

Examining the data by gender demonstrates that the rate of certification associated with back pain is slightly higher in males compared to females (figure 2). The rate of certification reaches a peak of 484.4 (95% CI 460.9, 508.2) certificates per 1000 back pain consultations in males in 2001 compared to 384.1 (95% CI 365.2, 403.7) certificates per 1000 back pain consultations in females in 2003, falling to a low of 275.7 (95% CI 259.9, 292.3) certificates per 1000 back pain consultations in males during 2010 compared to 223.3 (95% CI 210.6, 236.6) certificates per 1000 consultations in females during 2010 (figure 2). There was a statistically significant difference in the rate of sickness certification between males and females (p = < 0.001).

Comparing the rate of certification per 1000 back pain consultations by age shows that no single age group is reliably recording a higher rate of certification than other groups. However, the 60+ age group consistently record much lower rates of certificates per 1000 back pain consultations compared to the other age groups (figure 3). An ANOVA demonstrates that this difference in rate of certification between the 60+ age group and the other age groups is significant p = <0.001; excluding the 60+ age group from the ANOVA demonstrates that there are no statistical differences in the rate of certification between the age groups from 19 to 59 years.

DISCUSSION

Overall, the rate of sickness certification for back pain has significantly declined between 2000 and 2010. This decline in rates has principally occurred from 2003 onwards. This study demonstrated that approximately one third of consultations for back pain will lead to a sickness certificate; this finding is in line with many studies examining back pain which also find that approximately one third of participants are absent from work.^[13,14]

There was a trend in certification by gender, with men reporting consistently higher rates of certification than women. This is to be expected and has been reported in previous studies of certification using this dataset.^[11,15] The most likely explanation is the increased proportions of males working in the manual sector at 34.6% when compared to females at 10.9%.^[16] On average, manual workers report more absence than non-manual employees, and also report more absence as a result of

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

musculoskeletal conditions than non-manual employees.^[4] The differences in certification for gender may also be associated with differing consultation behaviours, and it has been reported that men are more likely to consult with an episode of back pain than women at a female:male rate ratio of 1:20 (95% Confidence Interval 1.13, 1.28).^[12]

There was no clear trend in certification by age with the exception of the 60+ years age group, who consistently had a lower rate of certification compared to younger age groups. Again, the rate of certification in this group showed a downward trend but it was not as pronounced as in other groups. There are a number of reasons why this difference may be seen in the 60+ age group. First, there are fewer people employed in this age group, in 2011 the Organisation for Economic Co-operation and Development reported that 56% of the population in the 55-64 years age group were employed compared to 80% of the population in the 25-54 years age group¹⁷. Secondly, data from 2013 demonstrated that whilst individuals in the 50-64 years age group bracket are less likely to work on a part time basis (just 28% of all workers in this age group) those in the over 64 years age group are more likely to report that they work part time, a total of 66% of all those in this age group^[18]. Thirdly, there could be a healthy worker effect whereby individuals suffering with back pain are removing themselves from the workplace. Findings are not likely to be consistent.^[19]

It is difficult to identify the reason for the decreasing rate of sickness certification in this study. It could be argued that the prevalence of back pain has decreased over time and so there is less need for certification. However, the literature suggests that the prevalence of back pain has remained largely unchanged over the period 1990- $2010^{[20]}$ and may even have increased.^[21, 22] It could also be that the number of consultations for back pain has increased, however the literature again suggests that this is not the case. ^[12, 23, 24] During the period of analysis that this paper spans there have been a number of initiatives, both in the UK and worldwide, surrounding the management of back pain that *may* have contributed to the decline in rates of certification. A review of clinical guidelines for the management of non-specific back pain assessed guidelines published between 2000 and 2008;^[25] this review is an update of a previous review.^[26] The authors report that the most common advice is to

reassure patients and encourage them to remain active, however, in contrast to the earlier review the current guidelines increasingly mention return to work, despite back pain, in their recommendations. In the UK, 2009 saw the publication of two National Institute for Clinical Excellence (NICE) reports, the first considering early management of non-specific back pain^[27] and the second reporting on primary care management of long term sickness absence.^[28] The non-specific back pain guidelines^[27] concur with the review by Koes *et al*^[26] that individuals should be encouraged to remain active and continue normal activities as far as possible. The second report is focussed on promoting the benefits of working with health conditions,^[28] specifically back pain, and again encourages individuals to maintain work despite pain. However, the impact of these recent reports is unlikely to be seen in the current analysis.

Recent research from the Global Burden of Disease study has demonstrated that back pain leads to more years lived with disability than any other condition and this burden of back pain is increasing with an ageing population.^[20] There is also evidence that the number of consultations for back pain has remained static, at least in the population of the United States, for a period of over 10 years from the 1990s to the mid-2000s.^[29] It is interesting then to note that despite the relative stability of consultations for back pain and the relatively adverse outcomes in terms of the burden of back pain that the number of sickness certificates for back pain has decreased in this study. Within the UK there have not been any recent public health initiatives which may account for this decrease in certification; there has however been the sea change in the information that general practitioners are advised to give their patients in regards to working with pain. There is consistent information provided by the Royal College of General Practitioners,^[30] Department for Work and Pensions^[31] and NICE^[27, 28] coupled with the availability of information booklets detailing the management of health and work for the GP, patient and employer.^[32, 33, 34, 35] It may be that media campaigns to promote working with musculoskeletal pain could see the rate of sickness certification reduce further. There has been some success with media campaigns in Scotland and Australia demonstrating that positive messages around working with musculoskeletal pain led to improvements in knowledge, attitudes and beliefs at the population level which were maintained, albeit at a reduced level, three years later.^[36, 37, 38] A Canadian study also utilising a media campaign had a more

Page 10 of 19

Page 11 of 22

BMJ Open

limited effect on behaviours related to back pain e.g healthcare use, than those run in Scotland and Australia.^[39] However, the Canadian study did find that participants agreement with the statement "if you have back pain you should try to stay active" significantly increased.^[39] It seems then that knowledge does not necessarily translate into behaviour and it has also been reported that provision of information alone is not sufficient to prevent work absence,^[40] but it seems that improving the baseline understanding of the general population may enhance any further information provided by healthcare professionals.^[41]

There are a number of limitations to the current study. Firstly, capturing the duration of sickness absence is not possible in the current dataset, meaning it is unclear whether the data are unduly influenced by a large number of short or long term certificates. The most frequently recorded duration for absence is two weeks^{[,42, 43, 44,} ^{45]} and there is no reason to believe that GPs in the current study would differ to any great degree. A second consideration is that this dataset is based in one area of the UK, North Staffordshire, and it could be argued that it is not generalizable to the rest of the population. Previous work with this dataset has demonstrated that crude rates of certification change very little when the data are standardised to the age and gender of the population as a whole, and there is no indication that this should be any different for this study.^[11] Lastly, the data included in the manuscript only goes as far as 2010 as from this date onwards the Read coding for sickness certificates changed as a result of the introduction of the Fit Note. This change in Read coding means that Fit Notes are now coded as "not fit for work" and "may be fit for some work". As this classification is different from the pre-2010 sickness certificates it would make comparison of the data between these two periods difficult in this manuscript because we would not be comparing like with like. As a result we could not be confident that any changes in the rate of certification would be as a result of the change to the Fit Note system or as a result of the change in Read coding i.e. more read codes available to classify not fit for work and may be fit for some work.

The main strength of this study is the large database in which this work was carried out. This dataset allowed the linking of consultation data to certification data to enable trends by age and gender to be seen. This is the first study to map trends in certification for back pain in the UK using an established dataset which has been

Page 11 of 19

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

validated for use in examining both consultation and sickness certification data.^[46,10] There is a need for establishing baseline rates of certification against which any change in policy or strategy at either local or national level can be compared,^[45] and this dataset goes some way to establishing these baseline figures. Linaker *et al*^[47] state that improvement of existing datasets would be more attractive than the development of a new national system to record sickness absence; with the introduction of the e- Fit Note the current data set is being updated to include information on duration of absence and whether or not a patient may be fit for work, further strengthening this dataset for future research. Another strong point of this study is that is appears to be supported by other literature; this lends credibility to our findings. For example, Ruseckaite *et* al^[48] reported a significant decrease in certification for musculoskeletal disorders between 2003 and 2010 in Australia. They also report a decrease in rate of certification for back pain and strains although there is no indication of whether this trend is significant or not. Gabbay et al also reported a decrease in the number of Fit Notes, lasting over 12 weeks, issued over a twelve month period. ^[49] Lambeek *et al*^[50] reported that the costs of back pain in the Netherlands decreased between 2002 and 2007, acknowledging that a large proportion of this cost is made up of absence. In the United Kingdom the Office for National Statistics have reported a decrease in reported absence across all health conditions between 1993 and 2013.^[51] However, work still needs to be carried out to investigate why there has been this change, and whether it is truly related to the available evidence about working with pain.

Conclusions

Rates of certification for back pain demonstrated a significant downward trend over the period 2000 to 2010; the reasons for this are not fully transparent but may be as a result of changing beliefs around working with back pain. These findings may provide a benchmark against which the impact of public health initiatives may be evaluated and monitored. With the new recording of the e-Fit Note this dataset will become more useful in tracking rates of certification over time.

ACKNOWLEDGEMENTS

We would like to thank the NIHR Clinical Research Network: West Midlands, and Ian Thomas, Simon Wathall and Kelvin Jordan for their assistance with data retrieval.

COMPETING INTERESTS

There are no competing interests.

FUNDING

This paper presents independent research funded by the National Institute for Health Research (NIHR), under its Programme Grants for Applied Research funding scheme: "Optimal management of spinal pain and sciatica in primary care" (NIHR-RP-PG-0707-10131). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool .

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

GW-J is funded by an NIHR Research Post-doctoral Fellowship (PDF-2009-02-54).

ETHICAL APPROVAL

Approval for use of the Medical Certificates and Consultation databases was gained from the North Staffordshire Research Ethics Committee (LREC Project 03/04)

CONTRIBUTORSHIP

GW-J and KD contributed to the design of the study, interpreting the analysis and writing of the manuscript. GW-J conducted the analysis.

DATA SHARING STATEMENT

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool .

rasmushogesc

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

No additional data available.



- pfizer absence workplace health 2013.pdf (accessed 31st March 2015)

BMJ Open

- Office For National Statistics, Statistical Bulletin: Labour Market Statistics 2014 http://www.ons.gov.uk/ons/rel/lmac/sickness-absence-in-the-labourmarket/2014/rpt---sickness-absence-in-the-labour-market.html (Accessed 31st March 2015)
- The Chartered Institute for Personnel and Development, Absence Management 2012, http://www.cipd.co.uk/binaries/absence-management_2012.pdf (Accessed 31st March 2015)
- Health and Safety Executive, Health and Safety Statistics, Annual Report for Great Britain (2013/2014), http://www.hse.gov.uk/statistics/overall/hssh1314.pdf (Accessed 31st March
- 6. Hiscock J, Ritchie J. The role of GPs in sickness certification. Leeds: Report number 148, Her Majesties Stationary Office 2001
- Hussey S, Hoddinott P, Wilson P, *et al.* Sickness certification system in the United Kingdom: qualitative study of views of general practitioners in Scotland. *BMJ* 2004;328:88
- Department of Health, Health and Safety Executive, Department for Work and Pensions. Health, work and well-being - Caring for our future. London: Department for Work and Pensions 2005
- Health work and wellbeing. Working for a healthier tomorrow 2008 https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/ 209782/hwwb-working-for-a-healthier-tomorrow.pdf (Accessed 31st March 2015)
- Wynne-Jones G, Mallen C.D, Main C.J. *et al.* Sickness certification in general practice: A comparison of electronic records with self-reported work absence. *Prim Health Res Dev* 2008;9:113-118
- Wynne-Jones G, Mallen C.D, Mottram S. *et al.* The identification of U.K. sickness certification rates, standardised for age and sex. *Br J Gen Pract* 2009;59(564):510-515
- Jordan K.P, Kadam U.T, Hayward R, *et al.* Annual consultation prevalence of regional musculoskeletal problems in primary care: an observational study. *BMC Musculoskelet Disord* 2010;11:144

- 13. Urquhart D.M, Kelsall H.L, Hoe V.C. *et al.* Are psychosocial factors associated with low back pain and work absence for low back pain in an occupational cohort?. *Clin J Pain* 2013;29(12):1015-1020
- 14. Wynne-Jones G, Cowen J, Jordan J.L, *et al.* Absence from work and return to work in people with back pain: a systematic review and meta-analysis. *Occup Environ Med* 2014:71(6):448-456
- Wynne-Jones G, Mallen C.D, Dunn K.M. Sickness certification for musculoskeletal conditions. *Clin Rheumatol* 2010;29(5):573-574
- 16. Office for National Statistics 2013 http://www.ons.gov.uk/ons/dcp171778_332467.pdf (Accessed 31st March 2015)
- 17. Organisation for Economic Co-operation and Development 2013 http://www.oecd-ilibrary.org/sites/factbook-2013en/07/01/02/index.html?contentType=&itemId=/content/chapter/factbook-2013-52-

en&containerItemId=/content/serial/18147364&accessItemIds=&mimeType=t ext/html (Accessed 31st March 2015)

- 18. Department for Work and Pensions, Older Workers Statistical Information Booklet 2013: Official Statistics <u>https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/</u>264899/older-workers-statistical-information-booklet-2013.pdf (Accessed 26th November 2015)
- Lemeunier N, Leboeuf-Yde C, Kjaer P, et al. Stability of low back pain reporting over 8 years in a general population aged 40/41 years at base-line: data from three consecutive cross-sectional surveys. BMC Musculoskelet Disord 2013;14:270
- 20. Hoy D, March L, Brooks P, *et al.* The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis* 2014;73:968-974
- Großschädl F, Freidl W, Rásky E, *et al.* A 35-Year Trend Analysis for Back Pain in Austria: The Role of Obesity. *PLOS One* 2014;9(9):e107436
- 22. Smith M, Davis M.A, Stano M. et al. Aging baby boomers and the rising cost of chronic back pain: secular trend analysis of longitudinal medical

BMJ Open

expenditures panel survey data for years 2000 to 2007. *J Manipulative Physiol Ther* 2013:36(1):2-11

- 23. Jöud A, Petersson I.F, Englund M, Low back pain: Epidemiology of consultations, Arthritis Care & Research 2012, 64(7) 1084–1088
- 24. Brtholomeeusen S, Van Zundert J, Truyers C, Buntinx F, Paulus D. Higher incidence of common diagnoses in patients with low back pain in primary care. *Pain Pract* 2012; 12: 1–6
- 25. Koes B.W, van Tulder M.W, Lin C-W.C, *et al.* An updated overview of clinical guidelines for the management of non-specific low back pain in primary care. *Eur Spine J* 2010;19:2075-2094
- 26. Koes B.W, van Tulder M.W, Ostelo R, *et al.* Clinical guidelines for the management of low back pain in primary care: an international comparison. *Spine* 2001;26:2504–2513
- 27. National Institute for Health and Care Excellence, Early Management of Persistent non-specific low back pain 2009a
- 28. National Institute for Health and Care Excellence, Managing long-term sickness and incapacity for work 2009b
- 29. Deyo R.A, Mirza S.K, Martin B.I, Back Pain Prevalence and Visit Rates Estimates From U.S. National Surveys. *Spine* 2002;31(23):2724-2727
- 30. Royal College of General Practitioners, Health and Work in General Practice http://www.rcgp.org.uk/courses-and-events/health-and-work-in-generalpractice-events.aspx (Accessed 31st March 2015)
- 31. Department for Work and Pensions, Royal College of General Practitioners, British Medical Association. Statement of fitness for work: a guide for general practitioners and other doctors. London: Department for Work and Pensions
- 32. Waddell G, Burton AK. Work and Health. London: The Stationary Office, 2006
- Waddell G, Burton AK. Health and Work. London: The Stationary Office, 2007a
- Waddell G, Burton AK. Advising Patients about Work. London: The Stationary Office, 2007b
- 35. Kendall NAS, Burton AK, Main CJ, *et al*, on behalf of the Flags Think-Tank: Tackling Musculoskeletal Problems: a Guide for the Clinic and Workplace –

Identifying Obstacles Using the Psychosocial Flags Framework 2009, London: The Stationary Office

- 36. Waddell G, O'Connor M , Boorman S. *et al.* Working Backs Scotland: a public and professional health education campaign for back pain. *Spine* 2007c;32:2139–43
- 37. Buchbinder R , Jolley D, Wyatt M . Population based intervention to change back pain beliefs and disability: three part evaluation. *BMJ* 2001;322:1516–20
- Buchbinder R , Jolley D. Population based intervention to change back pain beliefs: three year follow up population survey. *BMJ* 2004;328-321
- Gross D.P, Russell A.S, Ferrari R, Battié M.C, *et al.* Evaluation of a Canadian back pain mass media campaign. *Spine* 2010;35(8):906-913
- 40. Henrotin YE, Cedraschi C, Duplan B, *et al.* Information and low back pain management: a systematic review. *Spine* 2006;31:E326–E334.
- 41. Lorig K.R , Laurent D.D , Deyo R.A, et al. Can a Back Pain E-mail Discussion Group improve health status and lower health care costs?: A randomized study. Arch Intern Med 2002;162(7):792–6
- 42. Head J, Ferrie J. E, Alexanderson K, *et al.* Diagnosis-specific sickness absence as a predictor of mortality: the Whitehall II prospective cohort study. *BMJ* 2008;337:a1469
- 43. Tellnes G. Days lost by sickness certification. Scand J Prim Health Care 1989;7(4):245–251.
- 44. Reiso H, Gulbrandsen P, Brage S. Doctors' prediction of certified sickness absence. *Fam Pract* 2004;21(2):192–198.
- 45. Bollag U, Rajeswaran A, Ruffieuxf C, *et al.* Sickness certification in primary care—the physician's role. *Swiss Med Wkly* 2007; 137(23–24): 341–346
- 46. Porcheret M, Hughes R, Evans D. *et al.* Data quality of general practice electronic health records: the impact of a program of assessments, feedback, and training. *J Am Med Inform Assoc* 2004;11:78–86
- 47. Linaker C, Harris E.C, Cooper C, *et al.* The burden of sickness absence from musculoskeletal causes in Great Britain. *Occup Med* 2011;61(7):458-464
- 48. Ruseckaite R, Collie A, Bohensky M. *et al*, Trends in Sickness Certification of Injured Workers by General Practitioners in Victoria, Australia. J Occup Rehabil 2014:24:525-532

BMJ Open

1 2 3 4 5 6 7 8 9 10	 49. Gabbay M, Shiels certified sickness at 50. Lambeek L.C, van 7 back pain in the 2011;36(13):1050-1 	M, Hillage J, Facto psence in the UK, <i>Oc</i> Fulder M.W, Swinkle Netherlands in 058	ors associated with <i>cup Environ Med</i> , 2 es I.C.S <i>et al</i> . The tr the period 2002	length of fit note- 015:72:467-475 rend in total cost of to 2007. <i>Spine</i>
12	51 Office	for	National	Statistics
13			272000 16 (A	
14	http://www.ons.gov	.uk/ons/dcp1/1//6_	353899.pdf (Acces	ssed 31st March
15	2015)			
17				
18				
20				
21				
22				
23				
25				
26 27				
28				
29				
30 31				
32				
33				
35				
36				
37				
38 39				
40				
41				
43				
44				
45 46				
47				
48				
49 50				
51				
52 53				
54				
55				
56 57				
58				

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.



Figure 1: Rates of sickness certification associated with back pain per 1000 back pain consultations 210x297mm (300 x 300 DPI)



Figure 2: Rate of certification associated with back pain per 1000 back pain consultations by gender





BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.



Figure 3: Rate of certification associated with back pain per 1000 back pain consultations by age group

Figure 3: Rates of certification associated with back pain per 1000 back pain consultations by age group 210x297mm (300 x 300 DPI)

BMJ Open

Has there been a change in the rates of UK sickness certification for back pain over time? An examination of historical data from 2000 to 2010

Journal:	BMJ Open
Manuscript ID	bmjopen-2015-009634.R2
Article Type:	Research
Date Submitted by the Author:	04-Feb-2016
Complete List of Authors:	Wynne-Jones, Gwenllian; Arthritis Research UK Primary Care Centre, Primary Care Sciences Dunn, Kate; Arthritis Research UK Primary Care Centre, Primary Care Sciences
Primary Subject Heading :	Epidemiology
Secondary Subject Heading:	Epidemiology, Occupational and environmental medicine, General practice / Family practice
Keywords:	Rate, Absence, EPIDEMIOLOGY, Back pain < ORTHOPAEDIC & TRAUMA SURGERY



BMJ Open

Has there been a change in the rates of UK sickness certification for back pain over time? An examination of historical data from 2000 to 2010

Gwenllian Wynne-Jones*, Kate M Dunn

*Corresponding author Arthritis Research UK Primary Care Centre, Research Institute for Primary Care & Health Sciences, Keele University, Keele, Staffordshire, ST5 5BG Email; g.wynne-jones@keele.ac.uk Phone; 01782 733962

Key words; Back pain, Absence, Rate, Epidemiology Word count; 2759

ABSTRACT

Objectives

This paper aims to investigate historical patterns of sickness certification for back pain from 2000 to 2010.

Design

Electronic medical records from 14 practices that are part of the NIHR Clinical Research Network: West Midlands were reviewed. All records for back pain consultations from 2000 to 2010 were downloaded and matched, by date, to corresponding sickness certification records.

Setting

Primary Care

Results

A total of 93896 back pain consultations were recorded over the eleven year period resulting in 30913 sickness certificates. There was a statistically significant decrease in the rate of certification over the period, falling from 376.8 (95% CI 362.1, 392) per 1000 back pain consultations in 2000 to 246.5 (95% CI 236.5, 332.9) per 1000 back pain consultations in 2010. There was also a statistically significant difference in certification between males and females, with males issued more certificates than females. There was a statistically significant difference in certification by age, with those aged 60 years and over being less likely to be issued a certificate compared to all other age groups.

Conclusions

Rates of sickness certification for back pain demonstrated a downward trend between 2000 and 2010. Whilst the reasons for this are not transparent, it may be related to changing beliefs around working with back pain.

Strengths and limitations of this study

- This medical record review provides the ability to examine the rate of back pain associated sickness certificates over an eleven year period in all adults registered at 14 general practices.
- The database used has previously been demonstrated to be generalisable to the age and gender of the UK population.

BMJ Open

• The reasons for the decrease in sickness certification are not fully clear but may be related to changing beliefs around working with back pain.

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

INTRODUCTION

Many patients visit a general practitioner (GP) in primary care with symptoms of low back pain, in the United Kingdom (UK) each visit with a GP is termed a consultation. Whilst many patients consult their GP for back pain, 85% will not receive a hard diagnosis,^[1] and this consultation will be recorded on the medical record as a consultation for symptoms of back pain. As so few people with symptoms of back pain receive a hard diagnosis, this paper will use the phrase back pain to include both the majority of patients without a diagnosis, in addition to those who do have a diagnosis associated with their pain.

It has been estimated that 38% of adults are affected by back pain in any one year at an estimated cost to the National Health Service of £1 billion per annum.^[2] Musculoskeletal pain, principally back pain, is one of the most common reasons for absence from work.^[3]The costs of illness to employers, particularly back pain, are large; it has been estimated that 31 million days of absence were taken as a result of musculoskeletal pain in 2013.^[4] The Chartered Institute of Personnel and Development^[5] reports the rate of UK sickness absence in terms of a percentage of lost working hours, estimating sickness absence as 3.0%, 3.4% and 3.2% of working hours in 2012, 2011 and 2010 respectively. The Health and Safety Executive^[6] have estimated the total net cost of sick pay to be $\pounds 14.2$ billion during the year 2012-2013 as a result of work-related illness and injuries. When it is considered that musculoskeletal pain, principally back pain, is one of the most common reasons for absence from work it can be assumed that a large proportion of this sick pay is as a result of back pain.^[5] In the UK, proxy measures are used to estimate rates of sickness certification based on work absence. However, reporting rates of absence as a percentage of lost working hours does not reflect the clinical issues associated with certification in terms of the numbers of individuals absent from the workplace, nor does it represent the number of consultations for certification in general practice.^[7,8]

The health service costs and lost capacity in the workplace has made health and work a key target for public policy in the UK.^[9] The Government is actively aiming to reduce the number of employees signed off sick each year with a multi-agency government programme, launched in 2005, to address the issues of health, work and wellbeing.^[10] Furthermore, in 2010 the Fit Note was introduced to replace sickness

Page 4 of 20

BMJ Open

certificates. The aim of the Fit Note was to change the focus from what the patient can't do to what they can do in relation to work and to provide the GP with the option to state that patients may be fit for some work with provisions; a phased return to work, altered hours, amended duties and workplace adaptations. However, in order to assess the impact of such initiatives, methods are required to estimate rates of sickness certification over time, including the periods before such initiatives were implemented, a measure of the rate of certification per consultation is one method of achieving this.

The National Institute for Health Research Clinical Research Network: West Midlands holds frozen archive data on sickness certification in its Medical Certificates in Primary Care Archive and consultation data in its Consulters in Primary Care Archive. These databases have been validated for assessment of sickness certification,^[11] and the rates of sickness certification for a range of health conditions have been estimated.^[12] However, it is unknown whether the rate of certification for back pain has changed over time or whether there are any trends over time by age and gender. This paper aims to investigate historical patterns of sickness certification for back pain from 2000 to 2010.

METHODS

All consultation records from 2000 to 2010 for individuals with a back pain Read code were downloaded from the Consultation in Primary Care Archive database (CiPCA). Read codes are a hierarchy of morbidity, symptom and process codes, which become more specific further down the hierarchy. A set of Read codes has been established to identify both back pain diagnoses and symptoms consistently from the medical records, this comprises a total of 589 Read codes plus 20 other terms http://www.keele.ac.uk/mrr/morbiditydefinitions/.^[13] It was these codes that were used to identify patients for this study. The inclusion of both symptom and diagnostic codes ensures that both medically unexplained back pain and back pain that has a clear cause are included, as 85% of patients consulting with symptoms of back pain do not receive a diagnosis ^[1] focussing only on those patients with a diagnosis would exclude the vast majority of individuals for whom sickness absence is common.

Records for working age adults (aged 19-64 years), and those patients who were registered at the practice for the full year included in the analysis, were eligible. Each consultation record includes the unique individual identification number and practice identification number, plus age, gender, and year of consultation, date of consultation, the Read Code which was used to identify the problem with which the patient consulted, and the consultation free text. All sickness certification records between 2000 and 2010 for the same individuals, identified using each patient's unique identification number, were downloaded from the Medical Certificates in Primary Care Archive database (MiPCA). In the UK, a sickness certificate is required from the seventh day of absence. Each sickness certification number, plus age, gender, year of issue, date of issue, and the Read Code used to identify the record of a sickness certificate. Sickness certification records were then matched to back pain consultation records using the date of issue/consultation, the patient's unique identification number, plus age and gender.

Analysis

Numbers of consultations for back pain each year, and numbers of sickness certificates issued, were calculated using SPSS version 21. The rate of certification was defined as the number of certificates issued for back pain divided by the number of consultations for back pain in each year, presented per 1000 back pain consultations. The crude rate of certification and the rates by age and gender were calculated with 95% Confidence Intervals (CIs). A *t*-test for differences in proportions was calculated for year of certification and gender. An Analysis of Variance (ANOVA) was calculated to examine differences in rates by age. For all calculations the significance level was set at p=0.001 and a Bonferroni correction was applied to account for multiple analyses in the ANOVA.

Ethical approval

North Staffordshire Research Ethics Committee (LREC Project 03/04).

RESULTS

During the 11 year period (2000-2010) there were a total of 93896 consultations for back pain, resulting in 30913 sickness certificates being issued. This gives a sickness

Page 6 of 20

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

certification prevalence of 32.9%; or a third of all consultations for back pain resulting in a sickness certificate. The overall rate of certification was 329.2 certificates per 1000 consultations (95% CI 325.6, 332.9) (table 1 and figure 1).

There appears to be a downward trend in the rate of certification over the study period, with the rate of certification falling from 376.8 (95% CI 362.1, 392) per 1000 back pain consultations in 2000, to 246.5 (95% CI 236.5, 332.9) per 1000 back pain consultations in 2010 (see Figure 1). The rate of certification reaches a peak of 424.8 (95% CI 410.2, 439.7) per 1000 back pain consultations during the year 2003, with another spike in 2006. This decrease in the rate of certification over the eleven year tically signm... period was statistically significantly (p = <0.001).

Table 1: Rate of sickness certification per 1000 consultations for back pain, by year and gender

				Sickness certification rate	e per 1000 back pain cons	ultations (95 % CI)
Year	Total number of	Total number of	Prevalence (%)	Total	Males	Females
	certificates issued	consultations				
2000	2455	6515	37.7	376.8 (362.1, 392)	397.1 (375.2,419.9)	358.4 (338.5, 379)
2001	2809	6926	40.5	405.6 (390.7, 420.9)	484.1 (460.9, 508.2)	331.5 (312.8, 350.9)
2002	2863	6801	42.1	420.1 (405.7, 436.7)	473.7 (450.6, 497.7)	370.6 (350.7, 391.4)
2003	3215	7569	42.5	424.8 (410.2, 439.7)	471.4 (449,494.6)	384.1 (365.3, 403.7)
2004	3060	7476	40.9	409.3 (394.9, 424.1)	458.2 (436.5, 480.6)	362.6 (436.5, 480.6)
2005	2720	9338	29.1	291.3 (280.4, 302.4)	347.7 (330.3, 365.8)	242.9 (229.4, 256.9)
2006	3495	9442	37.0	370.1 (357.9382.6)	394.7 (376.3, 413.8)	349 (333, 365.7)
2007	2653	10200	26.4	260.1 (250.3, 270.2)	281.8 (266.6, 297.8)	242.9 (230.2, 256)
2008	2636	10046	26.2	262.4 (252.5, 272.6)	296.1 (280.6, 312.3)	233.7 (221, 246.9)
2009	2731	10349	26.4	263.9 (254.1, 2739)	293.4 (278.1, 309.4)	239.5 (226.9, 252.6)
2010	2276	9234	24.6	246.5 (236.5, 256.8)	275.7 (259.9, 292.3)	223.3 (210.6, 236.6)
All years	30913	93896	32.9	329.2(325.6, 332.9)	370.5 (364.8, 376.3)	293.5 (288.8, 298.3)

Page 9 of 23

BMJ Open

Examining the data by gender demonstrates that the rate of certification associated with back pain is slightly higher in males compared to females (figure 2). The rate of certification reaches a peak of 484.4 (95% CI 460.9, 508.2) certificates per 1000 back pain consultations in males in 2001 compared to 384.1 (95% CI 365.2, 403.7) certificates per 1000 back pain consultations in females in 2003, falling to a low of 275.7 (95% CI 259.9, 292.3) certificates per 1000 back pain consultations in males during 2010 compared to 223.3 (95% CI 210.6, 236.6) certificates per 1000 consultations in females during 2010 (figure 2). There was a statistically significant difference in the rate of sickness certification between males and females (p = < 0.001).

Comparing the rate of certification per 1000 back pain consultations by age shows that no single age group is reliably recording a higher rate of certification than other groups. However, the 60+ age group consistently record much lower rates of certificates per 1000 back pain consultations compared to the other age groups (figure 3). An ANOVA demonstrates that this difference in rate of certification between the 60+ age group and the other age groups is significant p = <0.001; excluding the 60+ age group from the ANOVA demonstrates that there are no statistical differences in the rate of certification between the age groups from 19 to 59 years.

DISCUSSION

Overall, the rate of sickness certification for back pain has significantly declined between 2000 and 2010. This decline in rates has principally occurred from 2003 onwards. This study demonstrated that approximately one third of consultations for back pain will lead to a sickness certificate; this finding is in line with many studies examining back pain which also find that approximately one third of participants are absent from work.^[14,15]

There was a trend in certification by gender, with men reporting consistently higher rates of certification than women. This is to be expected and has been reported in previous studies of certification using this dataset.^[12,16] The most likely explanation is the increased proportions of males working in the manual sector at 34.6% when compared to females at 10.9%.^[17] On average, manual workers report more absence than non-manual employees, and also report more absence as a result of

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool . Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

musculoskeletal conditions than non-manual employees.^[5] The differences in certification for gender may also be associated with differing consultation behaviours, and it has been reported that men are more likely to consult with an episode of back pain than women at a female:male rate ratio of 1:20 (95% Confidence Interval 1.13, 1.28).^[13]

There was no clear trend in certification by age with the exception of the 60+ years age group, who consistently had a lower rate of certification compared to younger age groups. Again, the rate of certification in this group showed a downward trend but it was not as pronounced as in other groups. There are a number of reasons why this difference may be seen in the 60+ age group. First, there are fewer people employed in this age group, in 2011 the Organisation for Economic Co-operation and Development reported that 56% of the population in the 55-64 years age group were employed compared to 80% of the population in the 25-54 years age group^[18]. Secondly, data from 2013 demonstrated that whilst individuals in the 50-64 years age bracket are less likely to work on a part time basis (just 28% of all workers in this age group) those in the over 64 years age group are more likely to report that they work part time, a total of 66% of all those in this age group^[19]. Thirdly, there could be a healthy worker effect whereby individuals suffering with back pain are removing themselves from the workplace. Findings are not likely to be related to differential reporting, as reporting of back pain over time is likely to be consistent.^[20]

It is difficult to identify the reason for the decreasing rate of sickness certification in this study. It could be argued that the prevalence of back pain has decreased over time and so there is less need for certification. However, the literature suggests that the prevalence of back pain has remained largely unchanged over the period 1990- $2010^{[21]}$ and may even have increased.^[22,23] It could also be that the number of consultations for back pain has increased, however the literature again suggests that this is not the case. ^[13, 24, 25] During the period of analysis that this paper spans there have been a number of initiatives, both in the UK and worldwide, surrounding the management of back pain that *may* have contributed to the decline in rates of certification. A review of clinical guidelines for the management of non-specific back pain assessed guidelines published between 2000 and 2008;^[26] this review is an update of a previous review.^[27] The authors report that the most common advice is to

Page 10 of 20

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

reassure patients and encourage them to remain active, however, in contrast to the earlier review the current guidelines increasingly mention return to work, despite back pain, in their recommendations. In the UK, 2009 saw the publication of two National Institute for Clinical Excellence (NICE) reports, the first considering early management of non-specific back pain^[28] and the second reporting on primary care management of long term sickness absence.^[29] The non-specific back pain guidelines^[28] concur with the review by Koes *et al*^[27] that individuals should be encouraged to remain active and continue normal activities as far as possible. The second report is focussed on promoting the benefits of working with health conditions,^[29] specifically back pain, and again encourages individuals to maintain work despite pain. However, the impact of these recent reports is unlikely to be seen in the current analysis.

Recent research from the Global Burden of Disease study has demonstrated that back pain leads to more years lived with disability than any other condition and this burden of back pain is increasing with an ageing population.^[21] There is also evidence that the number of consultations for back pain has remained static, at least in the population of the United States, for a period of over 10 years from the 1990s to the mid-2000s.^[30] It is interesting then to note that despite the relative stability of consultations for back pain and the relatively adverse outcomes in terms of the burden of back pain that the number of sickness certificates for back pain has decreased in this study. Within the UK there have not been any recent public health initiatives which may account for this decrease in certification; there has however been the sea change in the information that general practitioners are advised to give their patients in regards to working with pain. There is consistent information provided by the Royal College of General Practitioners,^[31] Department for Work and Pensions^[32] and NICE^[28, 29] coupled with the availability of information booklets detailing the management of health and work for the GP, patient and employer.^[33, 34, 35, 36] It may be that media campaigns to promote working with musculoskeletal pain could see the rate of sickness certification reduce further. There has been some success with media campaigns in Scotland and Australia demonstrating that positive messages around working with musculoskeletal pain led to improvements in knowledge, attitudes and beliefs at the population level which were maintained, albeit at a reduced level, three years later.^[37, 38, 39] A Canadian study also utilising a media campaign had a more

Page 11 of 20

limited effect on behaviours related to back pain e.g healthcare use, than those run in Scotland and Australia.^[40] However, the Canadian study did find that participants agreement with the statement "if you have back pain you should try to stay active" significantly increased.^[40] It seems then that knowledge does not necessarily translate into behaviour and it has also been reported that provision of information alone is not sufficient to prevent work absence,^[41] but it seems that improving the baseline understanding of the general population may enhance any further information provided by healthcare professionals.^[42]

There are a number of limitations to the current study. Firstly, capturing the duration of sickness absence is not possible in the current dataset, meaning it is unclear whether the data are unduly influenced by a large number of short or long term certificates. The data presented are a "pure" rate of certification i.e. a rate of certification per consultation. The data do not take into account the potential for a single patient to be issued multiple certificates for one episode of back pain. The possibility that the rate is artificially influenced as a result of a few patients receiving multiple certificates or deflated as a result of multiple patients receiving one certificate of a long duration cannot be ruled out. However, the most frequently recorded duration for absence is two weeks^[43, 44, 45, 46] and there is no reason to believe that GPs in the current study would differ to any great degree. It is also important to acknowledge that for approximately two thirds of people with an episode of back pain, a return-to-work within one month is expected^[15], indicating that the majority of consultations will be for individual episodes of back pain and therefore be individual certificates.

Secondly this dataset is based in one area of the UK, North Staffordshire, and it could be argued that it is not generalizable to the rest of the population. Previous work with this dataset has demonstrated that crude rates of certification change very little when the data are standardised to the age and gender of the population as a whole, and there is no indication that this should be any different for this study.^[12] Lastly, the data included in the manuscript only goes as far as 2010 as from this date onwards the Read coding for sickness certificates changed as a result of the introduction of the Fit Note. This change in Read coding means that Fit Notes are now coded as "not fit for work" and "may be fit for some work". As this classification is different from the pre-

Page 12 of 20

BMJ Open

2010 sickness certificates it would make comparison of the data between these two periods difficult in this manuscript because we would not be comparing like with like. As a result we could not be confident that any changes in the rate of certification would be as a result of the change to the Fit Note system or as a result of the change in Read coding i.e. more read codes available to classify not fit for work and may be fit for some work.

The main strength of this study is the large database in which this work was carried out. This dataset allowed the linking of consultation data to certification data to enable trends by age and gender to be seen. This is the first study to map trends in certification for back pain in the UK using an established dataset which has been validated for use in examining both consultation and sickness certification data.^{[11,13,} ⁴⁷] There is a need for establishing baseline rates of certification against which any change in policy or strategy at either local or national level can be compared, and this dataset goes some way to establishing these baseline figures. Linaker *et al*^[48] state that improvement of existing datasets would be more attractive than the development of a new national system to record sickness absence; with the introduction of the e- Fit Note the current data set is being updated to include information on duration of absence and whether or not a patient may be fit for work, further strengthening this dataset for future research. Another strong point of this study is that is appears to be supported by other literature; this lends credibility to our findings. For example, Ruseckaite *et* al^[49] reported a significant decrease in certification for musculoskeletal disorders between 2003 and 2010 in Australia. They also report a decrease in rate of certification for back pain and strains although there is no indication of whether this trend is significant or not. Gabbay *et al* also reported a decrease in the number of Fit Notes, lasting over 12 weeks, issued over a twelve month period. ^[50] Lambeek *et al*^[51] reported that the costs of back pain in the Netherlands decreased between 2002 and 2007, acknowledging that a large proportion of this cost is made up of absence. In the United Kingdom the Office for National Statistics have reported a decrease in reported absence across all health conditions between 1993 and 2013.^[52] However, work still needs to be carried out to investigate why there has been this change, and whether it is truly related to the available evidence about working with pain.

Conclusions

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool .

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

Rates of certification for back pain demonstrated a significant downward trend over the period 2000 to 2010; the reasons for this are not fully transparent but may be as a result of changing beliefs around working with back pain. These findings may provide a benchmark against which the impact of public health initiatives may be evaluated and monitored. With the new recording of the e-Fit Note this dataset will become more useful in tracking rates of certification over time.

ACKNOWLEDGEMENTS

racking We would like to thank the NIHR Clinical Research Network: West Midlands, and Ian Thomas, Simon Wathall and Kelvin Jordan for their assistance with data retrieval.

COMPETING INTERESTS

There are no competing interests.

FUNDING

This paper presents independent research funded by the National Institute for Health Research (NIHR), under its Programme Grants for Applied Research funding scheme: "Optimal management of spinal pain and sciatica in primary care" (NIHR-RP-PG-0707-10131). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

GW-J is funded by an NIHR Research Post-doctoral Fellowship (PDF-2009-02-54).

Page 14 of 20

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

ETHICAL APPROVAL

Approval for use of the Medical Certificates and Consultation databases was gained from the North Staffordshire Research Ethics Committee (LREC Project 03/04)

CONTRIBUTORSHIP

GW-J and KD contributed to the design of the study, interpreting the analysis and writing of the manuscript. GW-J conducted the analysis.

DATA SHARING STATEMENT

No additional data available.

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies

REFERENCES

- Deyo RA, Phillips WR. Low back pain. A primary care challenge. Spine. 1996;21(24):2826-32.
- 2. Maniadakis N, and Gray A, The economic burden of back pain. *Pain* 2000;84(1):95-103
- CBI Fit for Purpose: Absence and Workplace Health Survey 2013, http://www.cbi.org.uk/media/2150120/cbi-

pfizer_absence___workplace_health_2013.pdf (accessed 31st March 2015)

- Office For National Statistics, Statistical Bulletin: Labour Market Statistics 2014 http://www.ons.gov.uk/ons/rel/lmac/sickness-absence-in-the-labourmarket/2014/rpt---sickness-absence-in-the-labour-market.html (Accessed 31st March 2015)
- The Chartered Institute for Personnel and Development, Absence Management 2012, http://www.cipd.co.uk/binaries/absence-management_2012.pdf (Accessed 31st March 2015)
- 6. Health and Safety Executive, Health and Safety Statistics, Annual Report for Great Britain (2013/2014), http://www.hse.gov.uk/statistics/overall/hssh1314.pdf (Accessed 31st March
- Hiscock J, Ritchie J. The role of GPs in sickness certification. Leeds: Report number 148, Her Majesties Stationary Office 2001
- Hussey S, Hoddinott P, Wilson P, *et al.* Sickness certification system in the United Kingdom: qualitative study of views of general practitioners in Scotland. *BMJ* 2004;328:88
- Department of Health, Health and Safety Executive, Department for Work and Pensions. Health, work and well-being - Caring for our future. London: Department for Work and Pensions 2005
- Health work and wellbeing. Working for a healthier tomorrow 2008 https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/ 209782/hwwb-working-for-a-healthier-tomorrow.pdf (Accessed 31st March 2015)

Page 16 of 20

BMJ Open

11. Wynne-Jones G, Mallen C.D, Main C.J. et al. Sickness certification in general
practice: A comparison of electronic records with self-reported work absence.
Prim Health Res Dev 2008;9:113-118

- Wynne-Jones G, Mallen C.D, Mottram S. *et al.* The identification of U.K. sickness certification rates, standardised for age and sex. *Br J Gen Pract* 2009;59(564):510-515
- Jordan K.P, Kadam U.T, Hayward R, *et al.* Annual consultation prevalence of regional musculoskeletal problems in primary care: an observational study. *BMC Musculoskelet Disord* 2010;11:144
- 14. Urquhart D.M, Kelsall H.L, Hoe V.C. *et al.* Are psychosocial factors associated with low back pain and work absence for low back pain in an occupational cohort?. *Clin J Pain* 2013;29(12):1015-1020
- 15. Wynne-Jones G, Cowen J, Jordan J.L, *et al.* Absence from work and return to work in people with back pain: a systematic review and meta-analysis. *Occup Environ Med* 2014:71(6):448-456
- Wynne-Jones G, Mallen C.D, Dunn K.M. Sickness certification for musculoskeletal conditions. *Clin Rheumatol* 2010;29(5):573-574
- 17. OfficeforNationalStatistics2013http://www.ons.gov.uk/ons/dcp171778_332467.pdf(Accessed 31st March2015)

18. Organisation for Economic Co-operation and Development 2013 http://www.oecd-ilibrary.org/sites/factbook-2013en/07/01/02/index.html?contentType=&itemId=/content/chapter/factbook-2013-52-

en&containerItemId=/content/serial/18147364&accessItemIds=&mimeType=t ext/html (Accessed 31st March 2015)

- 19. Department for Work and Pensions, Older Workers Statistical Information Booklet 2013: Official Statistics <u>https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/</u> <u>264899/older-workers-statistical-information-booklet-2013.pdf</u> (Accessed 26th November 2015)
- 20. Lemeunier N, Leboeuf-Yde C, Kjaer P, *et al.* Stability of low back pain reporting over 8 years in a general population aged 40/41 years at base-line:

data from three consecutive cross-sectional surveys. *BMC Musculoskelet Disord* 2013;14:270

- 21. Hoy D, March L, Brooks P, et al. The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. Ann Rheum Dis 2014;73:968-974
- 22. Großschädl F, Freidl W, Rásky E, *et al.* A 35-Year Trend Analysis for Back Pain in Austria: The Role of Obesity. *PLOS One* 2014;9(9):e107436
- 23. Smith M, Davis M.A, Stano M. *et al.* Aging baby boomers and the rising cost of chronic back pain: secular trend analysis of longitudinal medical expenditures panel survey data for years 2000 to 2007. *J Manipulative Physiol Ther* 2013:36(1):2-11
- 24. Jöud A, Petersson I.F, Englund M, Low back pain: Epidemiology of consultations, Arthritis Care & Research 2012, 64(7) 1084–1088
- 25. Brtholomeeusen S, Van Zundert J, Truyers C, Buntinx F, Paulus D. Higher incidence of common diagnoses in patients with low back pain in primary care. *Pain Pract* 2012; 12: 1–6
- 26. Koes B.W, van Tulder M.W, Lin C-W.C, *et al.* An updated overview of clinical guidelines for the management of non-specific low back pain in primary care. *Eur Spine J* 2010;19:2075-2094
- 27. Koes B.W, van Tulder M.W, Ostelo R, *et al.* Clinical guidelines for the management of low back pain in primary care: an international comparison. *Spine* 2001;26:2504–2513
- 28. National Institute for Health and Care Excellence, Early Management of Persistent non-specific low back pain 2009a
- 29. National Institute for Health and Care Excellence, Managing long-term sickness and incapacity for work 2009b
- 30. Deyo R.A, Mirza S.K, Martin B.I, Back Pain Prevalence and Visit Rates Estimates From U.S. National Surveys. *Spine* 2002;31(23):2724-2727
- 31. Royal College of General Practitioners, Health and Work in General Practice http://www.rcgp.org.uk/courses-and-events/health-and-work-in-generalpractice-events.aspx (Accessed 31st March 2015)
- 32. Department for Work and Pensions, Royal College of General Practitioners, British Medical Association. Statement of fitness for work: a guide for general

BMJ Open

practitioners and other doctors. London: Department for Work and Pensions 2010

- Waddell G, Burton AK. Work and Health. London: The Stationary Office,
- Waddell G, Burton AK. Health and Work. London: The Stationary Office, 2007a
- 35. Waddell G, Burton AK. Advising Patients about Work. London: The Stationary Office, 2007b
- 36. Kendall NAS, Burton AK, Main CJ, et al, on behalf of the Flags Think-Tank: Tackling Musculoskeletal Problems: a Guide for the Clinic and Workplace – Identifying Obstacles Using the Psychosocial Flags Framework 2009, London: The Stationary Office
- 37. Waddell G, O'Connor M , Boorman S. *et al.* Working Backs Scotland: a public and professional health education campaign for back pain. *Spine* 2007c;32:2139–43
- Buchbinder R , Jolley D, Wyatt M . Population based intervention to change back pain beliefs and disability: three part evaluation. *BMJ* 2001;322:1516–20
- 39. Buchbinder R , Jolley D. Population based intervention to change back pain beliefs: three year follow up population survey. *BMJ* 2004;328-321
- 40. Gross D.P, Russell A.S, Ferrari R, Battié M.C, *et al.* Evaluation of a Canadian back pain mass media campaign. *Spine* 2010;35(8):906-913
- 41. Henrotin YE, Cedraschi C, Duplan B, *et al.* Information and low back pain management: a systematic review. *Spine* 2006;31:E326–E334.
- 42. Lorig K.R , Laurent D.D , Deyo R.A, *et al.* Can a Back Pain E-mail Discussion Group improve health status and lower health care costs?: A randomized study. *Arch Intern Med* 2002;162(7):792–6
- 43. Head J, Ferrie J. E, Alexanderson K, *et al.* Diagnosis-specific sickness absence as a predictor of mortality: the Whitehall II prospective cohort study. *BMJ* 2008;337:a1469
- 44. Tellnes G. Days lost by sickness certification. Scand J Prim Health Care 1989;7(4):245–251.
- 45. Reiso H, Gulbrandsen P, Brage S. Doctors' prediction of certified sickness absence. *Fam Pract* 2004;21(2):192–198.

- 46. Bollag U, Rajeswaran A, Ruffieuxf C, *et al.* Sickness certification in primary care—the physician's role. *Swiss Med Wkly* 2007; 137(23–24): 341–346
- 47. Porcheret M, Hughes R, Evans D. *et al.* Data quality of general practice electronic health records: the impact of a program of assessments, feedback, and training. *J Am Med Inform Assoc* 2004;11:78–86
- 48. Linaker C, Harris E.C, Cooper C, *et al.* The burden of sickness absence from musculoskeletal causes in Great Britain. *Occup Med* 2011;61(7):458-464
- Ruseckaite R, Collie A, Bohensky M. *et al*, Trends in Sickness Certification of Injured Workers by General Practitioners in Victoria, Australia. J Occup Rehabil 2014:24:525-532
- 50. Gabbay M, Shiels M, Hillage J, Factors associated with length of fit notecertified sickness absence in the UK, *Occup Environ Med*, 2015:72:467-475
- 51. Lambeek L.C, van Tulder M.W, Swinkles I.C.S *et al.* The trend in total cost of back pain in the Netherlands in the period 2002 to 2007. *Spine* 2011;36(13):1050-1058

52. Office for National Statistics http://www.ons.gov.uk/ons/dcp171776_353899.pdf (Accessed 31st March 2015)



2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010

Figure 1: Rates of sickness certification associated with back pain per 1000 back pain

Figure 1: Rates of sickness certification associated with back pain per 1000 back pain consultations 205x171mm (300 x 300 DPI)

BMJ Open: first published as 10.1136/bmjopen-2015-009634 on 25 April 2016. Downloaded from http://bmjopen.bmj.com/ on May 15, 2025 at Department GEZ-LTA Erasmushogeschool .

Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies











Figure 3: Rate of certification associated with back pain per 1000 back pain consultations by age group

Figure 3: Rates of certification associated with back pain per 1000 back pain consultations by age group 209x180mm (300 x 300 DPI)