BMJ Open Population characteristics, mechanisms of primary care and premature mortality in England: a cross-sectional study

Richard Baker,¹ Kate Honeyford,¹ Louis S Levene,¹ Arch G Mainous III,^{2,3} David R Jones,¹ M John Bankart,⁴ Tim Stokes⁵

To cite: Baker R,

Honevford K. Levene LS. et al. Population characteristics, mechanisms of primary care and premature mortality in England: a cross-sectional study. BMJ Open 2016;6: e009981. doi:10.1136/ bmjopen-2015-009981

Prepublication history and additional material is available. To view please visit the journal (http://dx.doj.org/ 10.1136/bmiopen-2015-009981).

Received 13 September 2015 Revised 2 December 2015 Accepted 31 December 2015



¹Department of Health Sciences, University of Leicester, Leicester, UK ²Department of Health Services Research, Management and Policy, University of Florida, Gainesville, Florida, USA ³Department of Community Health and Family Medicine, University of Florida, Gainesville, Florida, USA ⁴Department of Primary Care and Health Sciences, University of Keele, Keele, UK ⁵Department of General Practice and Rural Health. Dunedin School of Medicine, University of Otago, Dunedin, New Zealand

Correspondence to

Professor Richard Baker; rb14@le.ac.uk

ABSTRACT

Objectives: Health systems with strong primary care tend to have better population outcomes, but in many countries demand for care is growing. We sought to identify mechanisms of primary care that influence premature mortality.

Design: We developed a conceptual model of the mechanisms by which primary care influences premature mortality, and undertook a cross-sectional study in which population and primary care variables reflecting the model were used to explain variations in mortality of those aged under 75 years. The premature standardised mortality ratios (SMRs) for each practice, available from the Department of Health, had been calculated from numbers of deaths in the 5 years from 2006 to 2010. A regression model was undertaken with explanatory variables for the year 2009/2010, and repeated to check stability using data for 2008/2009 and 2010/2011.

Setting: All general practices in England were eligible for inclusion and, of the total of 8290, complete data were available for 7858.

Results: Population variables, particularly deprivation, were the most powerful predictors of premature mortality, but the mechanisms of primary care depicted in our model also affected mortality. The number of GPs/1000 population and detection of hypertension were negatively associated with mortality. In less deprived practices, continuity of care was also negatively associated with mortality.

Conclusions: Greater supply of primary care is associated with lower premature mortality even in a health system that has strong primary care (England). Health systems need to sustain the capacity of primary care to deliver effective care, and should assist primary care providers in identifying and meeting the needs of socioeconomically deprived groups.

INTRODUCTION

Health systems in many countries face financial constraints, but at the same time, they are also experiencing increased demand because of population ageing and, in some countries, population growth. Health systems

Strengths and limitations of this study

- The study included all general practices in England for which the necessary data were available (7858 practices included out of a total of 8290).
- A conceptual model, informed by previous research evidence, was developed prior to the analysis, to guide the selection of explanatory variables hypothesised to affect premature mortality. The model highlighted the potential roles of (1) the capacity of primary care, (2) access to care (including continuity), (3) clinical care (prevention, detection and management of disease) and (4) the performance of consultations in affecting premature mortality.
- The study was cross-sectional, and this limits the ability to draw conclusions about causality. A longitudinal study was not possible because the necessarv data were not available.
- The study used standardised mortality ratios as the measure of mortality. The use of count data would have been preferable, but these data were not available.
- The regression model explained a high proportion of the variation in premature mortality. We checked the stability of the regression model by using data for explanatory variables for three different years.

with stronger primary care, such as those in England, tend to achieve better health outcomes while restricting costs.^{1 2} However, the mechanisms by which primary care influences outcomes such as mortality are only opartly understood, although health system **g**. configuration, supply of physicians and continuity of care appear important.³ The introduction of a country-wide primary care strategy in Brazil was followed by a decline in cardiovascular mortality⁴; in the USA, supply of primary care physicians has been associated with reduced population mortality;⁵ and in England, relational continuity in primary care has been associated with lower mortality.⁶

BMJ

Starfield postulated six mechanisms by which primary care can affect population health: (1) greater access to services, (2) better quality of care, (3) greater focus on prevention, (4) early clinical management, (5) the effect of primary care characteristics (focus on the person, continuity) and (6) reduction of unnecessary specialist care.³ Primary care interventions may affect various outcomes and, in this paper, we focus on mortality. The characteristics of patients, including age, lifestyles and morbidity, have important effects on premature mortality. In particular, socioeconomic deprivation is a powerful predictor of mortality. However, primary care is expected to incorporate a population health perspective that includes responding to the health needs of deprived groups,⁷ and a key objective for primary care set by the WHO is to contribute to health equity and social justice.⁸

Our aim was to investigate whether a conceptual model representing some of the proposed mechanisms of primary care explains variations in premature mortality in general practice populations, taking particular account of deprivation among population characteristics. We hypothesised that the capacity of primary care (supply of general practitioners and their efficiency), access to care (being able to get an appointment with a general practitioner and relational continuity with a general practitioner) and the delivery of effective clinical care (prevention, detection and management of disease, and performance in consultations), would influence premature mortality.

METHOD

Study design

We undertook a cross-sectional study, in England, of premature mortality rates of patients of general practices (the main providers of English primary care) to identify variations in general practice associated with variations in mortality, while accounting for population characteristics. Since general practices have small populations, wide variations in mortality rates can occur from year to year.⁹ Therefore, published data over a 5-year period, provided by the Department of Health to the Health and Social Care Information Centre,¹⁰ were used to provide more stable mortality rates at practice level. We used data on population and practice characteristics for the year 2009-2010 as explanatory variables, and repeated the analysis with data from 2008/2009 and 2010/2011.¹⁰ All data were publicly available and, therefore, ethics committee approval was not required.

Study sample

All practices for which mortality data were published were eligible for inclusion (n=8290).

Mortality

We used the standardised mortality ratio (SMR) of those under the age of 75 years as the measure of premature

mortality.¹¹ The Office of National Statistics in England has recommended defining premature mortality as death under the age of 75 years.¹² The SMR is the ratio of the observed to expected number of deaths, the expected number of deaths being calculated using agespecific death rates for England and applied to local areas, determined by the Office of National Statistics, known as middle-level super output areas (MSOAs).¹³ MSOAs have an average population of 5000, allowing reliable estimates of mortality rates to be calculated. The number of observed deaths had been calculated at practice level by the government's Department of Health for the period 2006-2010 by taking a weighted average of by copyright, includ the mortality data for each MSOA in which a given practice's patients live. The weights are the percentage of the practice's patients residing in each MSOA.

Predictor selection

To guide the selection of predictor variables and to summarise our hypotheses, we developed a conceptual model explaining how population characteristics and primary care mechanisms influence mortality (figure 1). The model drew on Starfield's mechanisms³ and a model we had devised previously.⁶ In the model, population characteristics are expected to have the major impact. The model postulates a mechanism in which capacity of primary care will influence access, and access will enable patients to obtain preventive care, detection of illnesses and disease management. Our conceptual model focuses on primary care, but it recognises that public health, as well as secondary and social care, also affect mortality, as does the degree of connectedness between sectors.

Population predictors

Demographic characteristics related to premature mortality⁶¹⁴ were considered for inclusion. These included a measure of deprivation-the Index of Multiple Deprivation 2010 (IMD 2010)-which combines indicators in seven domains (income, employment, health, education, housing, crime and environment) into a score for each small area in England.¹⁵ The GP Practice IMD was estimated by taking a weighted average of the IMD scores for each lower-level super output area (LSOA) in which a given practice had registered for patients.¹⁶ A LSOA has a population size of approximately 1500 people, and provides a more discriminating unit for calculation of practice population deprivation scores than use of the larger population unit of MSOAs. We included an estimate of smoking in those with chronic conditions, derived from pay for performance indicators for chronic disease management (the UK quality and outcomes framework for general practice-QOF).¹⁴ To reflect the underlying chronic disease morbidity of the practice population, we selected the prevalence of diabetes, based on QOF registers. General practice diabetes registers are good measures of the prevalence of diabetes,¹⁷ and, in a recent study from

Figure 1 Model of how population and primary care characteristics are postulated to influence population mortality; characteristics of social and secondary care and public health not included. Modified from Levene et al⁶ to include Starfield et al s³ mechanisms: ¹greater access; ²better quality of care; ³greater focus on prevention; ⁴early management of health problems; ⁵primary care characteristics: ⁶reducing unnecessary specialist care.



National features - health system, economy, policies affecting health (e.g. food, road safety)

Scotland, 47% of people with diabetes were found to have at least three other chronic conditions.¹⁸ The percentage of the practice population who were white was taken from the GP patient survey.¹⁹ We undertook sensitivity analyses comparing the use of these data with estimates of the ethnicity of practice populations derived from hospital admissions for 2005–2007.²⁰

Primary care predictors

We used numbers of full-time equivalent general practitioners (GPs)/1000 population, a measure designed by the National Health Service (NHS) Information Centre to describe the capacity of each practice in relation to the population served.²¹ The measure includes all GPs except those in training, and used a measure of the patient population weighted according to patient need. We used the total number of patients registered with the practice list as a measure of efficiency,²² since larger practices potentially have opportunities for efficiencies.²

Our model distinguished between capacity and access,²⁴ for which we used patient reports collected through the annual GP patient survey, specifically the percentage of patients who recalled being able to make an appointment within 48 h.¹⁹ Similarly, we used the percentage of patients who recalled being able to make an appointment with a GP in advance as a measure of continuity. The GP patient survey includes random samples of the adult population of every general practice, and these survey items were selected because they have been shown to be adequately reliable for allocating access incentive payments to practices.²⁵ We also included another survey measure expressing patients'

trust and confidence in their GP as a measure of consultation performance.

Indicators from the QOF were adopted as measures of prevention, detection and management. We used a study of the estimated public health impact of OOF indicators to select the prevention and management indicators with the greatest potential impact on mortality,²⁶ while at the same time limiting the number of indicators, to reduce the risk of identifying spurious associations. For prevention, we used the percentage of patients with diabetes having influenza immunisation (estimated annual mortality reduction 63.7/100000 patients 26), and the proportion of smokers with a chronic condition who were offered help with cessation (estimated annual mortality reduction $10.9/100000^{26}$). For detection, we used the proportion of patients recorded as having atrial fibrillation with the diagnosis confirmed by an ECG or specialist, and the proportion f patients on the practice hypertension register, since freater detection of hypertension is associated with wer coronary heart disease and stroke mortality rates.⁶ For management, underlying achievement of selected **g**. of patients on the practice hypertension register, since greater detection of hypertension is associated with lower coronary heart disease and stroke mortality rates.⁶

QOF indicators was used, that is, the percentage of eli- $\overline{\mathbf{g}}$ gible patients for whom a target has been achieved. We used the percentage of patients with hypertension in whom the last blood pressure reading was 150/ 90 mm Hg or less (estimated annual mortality reduction $48.2/100\ 000^{26}$), the percentage of patients with coronary heart disease currently treated with a β-blocker (estimated annual mortality reduction $45.9/100\ 000^{26}$), the percentage of patients with diabetes in whom the last glycated haemoglobin (HbA1c) level was 7.0%

З

(53.0 mmol/mol) or less (estimated annual mortality reduction $26.5/100\ 000^{26}$), and the percentage of patients on the chronic kidney disease register in whom the last blood pressure reading was $140/85\ \text{mm Hg}$ or less (estimated annual mortality reduction $26.2/100\ 000^{26}$).

Statistical analysis

The analysis sought to determine the potential effect of all candidate explanatory variables on indirectly standardised premature mortality rates, calculated from deaths in 2006–2010. QOF and patient survey data from 2009/ 2010 were used as explanatory variables in the primary analysis. Stata V.11 was used to undertake the analysis (StataCorp. Stata data analysis and statistical software. Release 11. New in Stata 11. http://www.stata.com/ stata11/ (accessed 8 Apr 2015)). No formal adjustment for multiple hypothesis testing (multiplicity) was made.

In stage 1, we undertook univariable analysis of all explanatory variables. In stage 2, we developed a multiple linear regression model of premature SMR including all explanatory variables without transformation, to assess our conceptual model. The errors were assumed to follow a normal distribution, with regression diagnostics supporting this assumption. Counts of deaths were not available, and it was not possible to use a count model.²⁷ Primary care trusts and mortality estimates, based on local areas that overlap for some general practices, cannot be considered independent. We adjusted for the clustering effect by using robust SEs.²⁸

To test the stability of the regression model, we repeated it using QOF and patient survey variables from 2008/2009 and 2010/2011. Data from 2007/2008 and earlier were not used as there were changes to the patient survey in 2008/2009. As the distribution of premature SMR was positively skewed, the log of the SMR was also modelled.

Stage 3 sought to determine the possible interactions between deprivation and primary care characteristics. Interactions between primary care characteristics and deprivation were then examined in the model, one at a time. Aikaike and Bayesian Information Criteria (AIC and BIC)²⁹ were used to determine whether including an interaction improved model fit.

RESULTS

There were 8290 general practices in England in 2009/2010. Of these, 432 were omitted from the analysis for the following reasons: missing data on deprivation of the practice population (n=63); missing data on ethnicity of the practice population (n=153); missing data for QOF indicators (n=29); missing data on exception reporting of QOF indicators (n=9); missing data on numbers of GPs/1000 (n=105); incomplete general practice patient survey data (n=15); practices with no patients to whom QOF indicators applied (n=58). The final number of practices included was 7858.

The median premature SMR was 103 (IQR: 85 to 125), ranging from 34 to 223. Univariable analysis of individual explanatory variables indicated that increasing deprivation, smoking and diabetes prevalence were positively associated, and white ethnicity negatively associated, with premature mortality (table 1). All aspects of patient experience were negatively associated with premature mortality, although the proportion of the variation explained was low. Hypertension detection was negatively associated with premature mortality, but a mixed pattern was found when the other clinical management variables were considered.

The regression model (table 2) explained 84% of the **b** variation in premature mortality. Of the population characteristics, increasing deprivation, increasing proportions of the practice population who were white, increasing proportions with diabetes and increasing proportions who were smokers, were associated with higher mortality.

Of the practice predictors, increasing numbers of GPs/1000 population (capacity), increasing proportions able to get an appointment within 48 h (access), increasing numbers of patients on hypertension registers (detection) and increasing proportions of those with hypertension with blood pressure below 150/90 mm Hg (management), were all associated with lower mortality. Increasing proportions of those with coronary heart disease treated with a β -blocker, increasing proportions ç of patients with diabetes in whom the last HbA1c level e was 7.0% or less and increased proportions of patients and with chronic kidney disease with blood pressure below 140/85 mm Hg were all associated with increasing SMR, although the coefficients for all the management (and some other) variables were very small.

Based on this model, the benefits from primary care may be quite small at the level of a practice, but would ≥ be much greater when aggregated to all of England (table 3). Using MSOA level data on the numbers of deaths in small populations, we have estimated that a practice with a median list size of 6009 patients (see table 1) could be expected to have 20 deaths in 1 year, based on an SMR of between 95 and 105. An increase in <u>0</u> hypertension detection of 1% was associated with an annual reduction of 0.16 premature deaths, but when aggregated to the whole of England, this equates to a potential decrease of 1202 premature deaths. This approach was used to calculate the possible impact on the number of premature deaths in England of a one & unit change in each explanatory variable, shown in table 3. In 2009/2010, the median number of patients per GP was 1800; decreasing this to 1400 (equivalent to an additional 8500 full time GPs in post) would equate to 0.14 fewer premature deaths annually in a practice, and 1078 in England as a whole.

In stage 3 of the analysis, interactions between deprivation and five service characteristics improved the model fit. There were negative interactions between deprivation and GP supply, the proportion of patients able to make

Variable	Median (IQR)	(Minimum, maximum)	Univariable β-coefficient (95% Cl)	p Value	R ²
Population characteristics					
IMD 2010	21.55 (13.57, 31.85)	(2.86, 66.30)	2.00 (1.98 to 2.02)	<0.001	0.81
Proportion of the practice population who are white (%)	91.60 (77.14, 95.24)	(0.56, 100.00)	-0.335 (-0.362 to -0.309)	<0.001	0.074
Proportion of those with chronic conditions who are smokers (%)	16.63 (12.94, 21.24)	(1.13, 93.79)	2.86 (2.80 to 2.93)	<0.001	0.51
Proportion of the practice population who are on the practice's diabetes register (%)	5.37 (4.59, 6.23)	(0.33, 14.77)	6.14 (5.78 to 6.50)	<0.001	0.12
Efficiency					
Registered list size	6009 (3557, 9290)	(406, 39 919)	-0.0012 (-0.0013 to -0.0010)	<0.001	0.034
Supply		· · · · · ·	, ,		
Number of GPs per 1000 patients	0.557 (0.460, 0.668)	(0.000, 5.837)	-7.08 (-9.66 to -4.50)	<0.001	0.004
Access	, , , , , , , , , , , , , , , , , , ,	· · · ·	, , , , , , , , , , , , , , , , , , ,		
Proportion of patients reporting being able to get an appointment within 48 h (%)	83.33 (75.00, 90.07)	(22.01, 100)	-0.59 (-0.64 to -0.54)	<0.001	0.061
Continuity of care and quality of relationship					
Proportion of patients reporting being able to get an appointment in advance (%)	73.89 (62.13, 84.02)	(14.81, 100)	-0.34 (-0.38 to -0.30)	<0.001	0.037
Proportion of patients expressing trust in their GP (%)	95.42 (92.82, 97.05)	(66.67, 100)	-2.02 (-2.17 to -1.88)	<0.001	0.088
Prevention	, , , , , , , , , , , , , , , , , , ,	, , , ,	, , , , , , , , , , , , , , , , , , ,		
DM18 The percentage of patients with diabetes who have had influenza immunisation in the preceding 1 September to 31 March (%)	78.39 (74.28, 82.31)	(25.00, 100)	-0.41 (-0.50 to -0.32)	<0.001	0.010
SM04 The percentage of patients with any or any combination of 10 specified conditions who smoke, and whose notes contain a record that	92.43 (90.31, 95.15)	(33.60, 100)	-0.15 (-0.27 to -0.038)	0.009	0.000
smoking cessation advice or referral to a specialist service, where available,					
nas been offered within the previous 15 months					
		(0.00.400)		0.474	0.000
AF04 The percentage of patients with atrial fibrillation diagnosed after 1 April 2008 with ECG or specialist-confirmed diagnosis	91.30 (85.11, 100.00)	(0.00, 100)	0.018 (-0.031 to 0.066)	0.471	0.000
Proportion of practice population on the QOF hypertension register (%)	13.55 (11.48, 15.68)	(0.16, 36.65)	-1.18 (-1.35 to -1.01)	<0.001	0.024
Clinical management		(
BP05 The percentage of patients with hypertension in whom the last blood	76.16 (71.82, 80.31)	(38.30, 99.45)	-0.22 (-0.31 to -0.13)	<0.001	0.003
pressure (measured in the previous 9 months) is 150/90 or less			/		
CHD10 The percentage of patients with coronary heart disease who are	56.39 (51.62, 61.14)	(16.33, 100)	0.25 (0.18 to 0.33)	<0.001	0.005
currently treated with a β-blocker (unless a contraindication or side effects are recorded)					
DM23 The percentage of patients with diabetes in whom the last HbA1c	46.58 (41.00, 52.44)	(0, 96.40)	-0.22 (-0.29 to -0.16)	<0.001	0.006
level is 7 or less (or equivalent test/reference range depending on local laboratory) in the previous 15 months					
CKD03 The percentage of patients on the CKD register in whom the last blood pressure reading, measured in the previous 15 months, is 140/85 or	69.20 (63.50, 75.00)	(0.00, 100)	0.22 (0.16 to 0.28)	<0.001	0.006

IMD, Index of Multiple Deprivation; QOF, quality and outcomes framework; SMR, standardised mortality ratios.

BMJ Open: first published as 10.1136/bmjopen-2015-00981 on 11 February 2016. Downloaded from http://bmjopen.bmj.com/ on May 11, 2025 at Department GEZ-LTA Erasmushogeschool . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

сл

07

Variable	lices in Lingianu (i	1=7856)	
	β-Coefficient	95% CI	p Value
Population characteristics			
IMD 2010	1.81	(1.69 to 1.94)	<0.001
Proportion of the practice population who are white (%)	0.17	(0.10 to 0.25)	<0.001
Proportion of those with chronic conditions who are smokers (%)	0.49	(0.39 to 0.64)	<0.001
Proportion of the practice population who are on the practice's diabetes	1.30	(0.50 to 2.11)	0.002
register (%)			
Efficiency			
Registered list size	0.00002	(-0.00008 to 0.0001)	0.739
Supply			
Number of GPs per 1000 patients	-4.31	(-6.80 to -1.82)	<0.001
Access			
Proportion of patients reporting being able to get an appointment within	-0.06	(-0.11 to -0.02)	0.006
48 h (%)			
Continuity of care and quality of relationship			
Proportion of patients reporting being able to get an appointment in	-0.01	(-0.04 to 0.01)	0.283
advance (%)		· · · · ·	
Proportion of patients expressing trust in their GP (%)	0.03	(-0.10 to 0.16)	0.635
Prevention		· · · · ·	
DM18 The percentage of patients with diabetes who have had influenza	0.007	(-0.06 to 0.08)	0.849
immunisation in the preceding 1 September to 31 March (%)		X 7	
SM04 The percentage of patients with any or any combination of 10	-0.02	(-0.08 to 0.04)	0.587
specified conditions who smoke, and whose notes contain a record that		(
smoking cessation advice or referral to a specialist service, where			
available, has been offered within the previous 15 months			
Detection			
AF04 The percentage of patients with atrial fibrillation diagnosed after 1	-0.002	(-0.032 to 0.029)	0.918
April 2008 with ECG or specialist confirmed diagnosis	0.002	(0.002 10 0.020)	0.0.0
Proportion of practice population on the QOE hypertension register (%)	-0.77	(-0.98 to -0.56)	<0.001
Clinical management	0.77	(0.00 10 0.00)	
BP05 The percentage of patients with hypertension in whom the last	-0.077	(-0.15 to -0.004)	0.04
blood pressure (measured in the previous nine months) is 150/90 or less	0.077	(0.10 10 0.001)	0.01
CHD10 The percentage of patients with coronary heart disease who are	0.088	(0.018 to 0.16)	0.015
currently treated with a B-blocker (unless a contraindication or side effects	0.000	(0.010 10 0.10)	0.010
are recorded)	,		
DM23 The percentage of patients with diabetes in whom the last HbA1c	0.06	(_0.003 to 0.13)	0.061
lovel is 7 or loss (or equivalent test/reference range depending on local	0.00	(-0.000 10 0.10)	0.001
laboratory) in the provious 15 months			
CKD02 The percentage of patients on the CKD register in whom the last	0.0200	(0,0007 to 0,0977)	0 1 1 6
blood proceure reading measured in the provious 15 menths is 140/85	0.0590	(-0.0037 10 0.0077)	0.110
or loss			
Of itess	a a such faur als sata vieras	No formal adjustes and form	ee a altie lie ite a
Statistical model: multiple linear regression of premature SMR with robust errors to acc	count for clustering.	No formal adjustment for l	multiplicity.
ratios.	ro, giyoalou naomo		montanty
appointment within 48 h or in advance and those Sensitivity	/ analysis		
appointment within 48 h or in advance and those Sensitivity	<mark>/ analysis</mark> ings of analyses (of the 3 years separate	elv (2008)
appointment within 48 h or in advance and those spressing trust in their GP. A positive interaction was The findi	/ analysis ings of analyses (09/2010 - 2010/	of the 3 years separate	ely (2008)
appointment within 48 h or in advance and those spressing trust in their GP. A positive interaction was und between deprivation and the percentage of 2009, 200 vients on the company heart discuss (CUD) parities.	/ analysis ings of analyses (09/2010, 2010/	of the 3 years separate 2011) gave consisten	ely (2008) nt conclu
a appointment within 48 h or in advance and those pressing trust in their GP. A positive interaction was und between deprivation and the percentage of ttients on the coronary heart disease (CHD) register utients of the other than the percentage of the sions with	/ analysis ings of analyses 09/2010, 2010/ h only minor	of the 3 years separate 2011) gave consisten differences. More c	ely (2008/ nt conclu letails are
a appointment within 48 h or in advance and those pressing trust in their GP. A positive interaction was und between deprivation and the percentage of atients on the coronary heart disease (CHD) register eated with β-blocker. The inclusion of the interaction given in	/ analysis ings of analyses 09/2010, 2010/ h only minor the online s	of the 3 years separate 2011) gave consisten differences. More c upplementary apper	ely (2008/ nt conclu letails are ndix. The
a appointment within 48 h or in advance and those pressing trust in their GP. A positive interaction was und between deprivation and the percentage of atients on the coronary heart disease (CHD) register eated with β-blocker. The inclusion of the interaction given in results of the five service characteristics suggests results of the service characteristics sugg	/ analysis ings of analyses 09/2010, 2010/ h only minor the online s the log transfo	of the 3 years separate 2011) gave consisten differences. More c upplementary apper rmed model were al	ely (2008, nt conclu letails are ndix. The so consist

an appointment within 48 h or in advance and those expressing trust in their GP. A positive interaction was found between deprivation and the percentage of patients on the coronary heart disease (CHD) register treated with β -blocker. The inclusion of the interaction term for each of the five service characteristics suggests that the impact of an improvement in supply, access, continuity of care and consultation quality is lower for practices serving a more deprived community. The percentage of patients with CHD treated with a β-blocker was positively associated with mortality, again inclusion of the interaction term reduced the impact of an improvement in management in patients serving more deprived patients. This is illustrated in table 4.

results of the log transformed model were also consistent with the original model, although the fit of the logged model may be inferior. When practice size was modelled as a categorical variable, to indicate the smallest 10% of practices, there was evidence that practice size was associated with premature mortality. Small practices, with a list size of 2367 (identifying the smallest 10% of practices), were associated with higher premature mortality.

	Potential reduction in premature deaths in England in 1 year if there is a 1 unit* change in the explanatory variable (95% Cl)†
Population characteristics	
Change is a decrease in 1 unit that is associated with a reduction in deaths for all variables	
Reduction in IMD of 1 unit	2830 (2642 to 3033)
Reduction of 1% in the proportion of the practice population who are white	269 (156 to 391)
Reduction of 1% in the proportion of those with chronic conditions who are smokers	764 (610 to 1001)
Reduction of 1% in the proportion of the practice population who are on the practice's	2032 (782 to 3299)
diabetes register	
Primary care characteristics	
Change is an increase in 1 unit that is associated with a reduction in deaths for some varial	oles and an increase in deaths
for others	
Efficiency	
Increase in 1000 patients in registered list size*	–25 (–156 to 81)
Supply	
Increase of 1.00 GPs per 1000 patients	6738 (2845 to 10 631)
Increase of 0.16 GPs per 1000 patients*	1078 (455 to 1701)
Access	00(01 + 170)
increase of 1% in the proportion of patients reporting being able to get an appointment	96 (31 to 172)
WILING 48 D	
Continuity of care and quality of relationship	01 (10 + 00)
increase of 1% in the proportion of patients reporting being able to get an appointment in	21 (-16 10 63)
duvalice	$40(.250 \pm 0.156)$
Provention	-49 (-250 10 156)
Increase of 1% in the percentage of patients with diabates who have had influenza	-10 (-125 to 63)
immunisation in the preceding 1 September to 31 March (DM18)	10 (123 10 00)
Increase of 1% in the percentage of natients with any or any combination of 10 specified	25 (-63 to 125)
conditions who smoke, and whose notes contain a record that smoking cessation advice	20 (00 10 120)
or referral to a specialist service, where available, has been offered within the previous	
15 months (SM04)	
Detection	
Increase of 1% in the percentage of patients with atrial fibrillation diagnosed after 1 April	2 (-45 to 50)
2008 with ECG or specialist-confirmed diagnosis (AF04)	, , , , , , , , , , , , , , , , , , ,
Increase of 1% in the proportion of practice population on the QOF hypertension register	1202 (875 to 1532)
Clinical management	
Increase of 1% in the percentage of patients with hypertension in whom the last blood	121 (6 to 234)
pressure (measured in the previous 9 months) is 150/90 or less (BP05)	
Increase of 1% in the percentage of patients with coronary heart disease who are	-137 (-250 to -28)
currently treated with a β -blocker (unless a contraindication or side effects are recorded) (CHD10)	
Increase of 1% in the percentage of patients with diabetes in whom the last HbA1c level is	-100 (-203 to 5)
7 or less (or equivalent test/reference range depending on local laboratory) in the previous	
15 months (DM23)	
Increase of 1% in the percentage of patients on the CKD register in whom the last blood	-61 (-137 to 15)
pressure reading, measured in the previous 15 months, is 140/85 or less (CKD03)	

been chosen. A change in GPs/1000 of 0.16 is equivalent to a reduction in list size where changes that are more applicable to practice of the size of

†Negative figures indicate a negative reduction in deaths, that is, an increase in deaths.

CKD, chronic kidney disease; DM, diabetes mellitus; GPs, general practitioners; HbA1c, glycated haemoglobin; IMD, Index of Multiple Deprivation; SMR, standardised mortality ratios.

DISCUSSION Principal findings

Population variables were the most powerful predictors of mortality. Practices with greater levels of diabetes,

more smokers, more white patients and greater deprivation, had higher levels of premature mortality. Deprivation had the greatest effect. Although non-white ethnicity was associated with higher premature mortality

being able ableBeing able to get an appointment within appointment within appointment within appointment within mpact onbeing able to get an appointment within appointment within appointment within appointment within advance (%)Proportion of patients with CHD who are currently reated with a β -blocker (CHD10) (%)ableGP s per 1000 patients appointment within mpact on SMR for each oneHa (%) (CHD10) (%)CHD10) (%) (CHD10) (%)Impact on SMR for each oneGP s/1000 (GP s/1000Median mpact onMedian mpact on (CHD10) (%)Median (CHD10) (%)Impact on soluteGP s/1000 (GP soluteMedian (GP soluteMedian (GP soluteMedian (GP soluteImpact on soluteGP s/1000 (GP soluteSMR for each (for eachMedian (for each for each for (for each for	ble 4 Inte	erpretation of the	interaction b	etween service c	haracteristics a atients	nd deprivation	patients			The percentage of	patients
Impact on SMR for each one initiabsoluteImpact on 	able	GPs per 1000	patients	being able to g appointment w 48 h (%)	let an ithin	being able to g appointment in advance (%)	get an	Proportion of F expressing tru: GP (%)	atients st in their	with CHD who are treated with a β-blo (CHD10) (%)	cker
ivationunit absolutefor each1% absolutefor each1%for each1%tileincreasequintileincreasequintileincreasequintileincreasequintileincreasequintile -8.28 0.576 -0.104 86.8 -0.0581 79.3 -0.225 96.6 0.175 55.6 -7.09 0.578 -0.0904 85.4 -0.0412 76.5 -0.151 96.2 0.149 56.1 -5.62 0.554 -0.0904 81.2 $+0.00451$ 72.9 -0.0598 95.3 0.108 56.1 -3.86 0.54 -0.0527 81.2 $+0.00451$ 70.7 0.0451 94.6 0.0623 56.9 -1.34 0.537 -0.0233 78.6 $+0.04022$ 69.2 0.207 93.1 -0.00208 56.6		Impact on SMR for each one	Median GPs/1000	Impact on SMR for each	Median	Impact on SMR for each	Median	Impact on SMB for each	Median nercentade	Impact on SMR	Median
-8.28 0.576 -0.104 86.8 -0.0581 79.3 -0.225 96.6 0.175 55.6 -7.09 0.578 -0.0904 85.4 -0.0412 76.5 -0.151 96.2 0.149 56.1 -5.62 0.55 -0.0733 83.1 -0.0205 72.9 -0.0598 95.3 0.108 56.1 -3.86 0.54 -0.0527 81.2 +0.00451 70.7 0.0451 94.6 0.0623 56.9 -1.34 0.537 -0.0233 78.6 +0.0402 69.2 0.0623 56.9	rivation	unit absolute increase	for each quintile	1% absolute increase	for each quintile	1% absolute increase	for each quintile	1% absolute increase	for each quintile	for each 1% absolute increase	for each quintile
-7.09 0.578 -0.0904 85.4 -0.0412 76.5 -0.151 96.2 0.149 56.1 -5.62 0.55 -0.0733 83.1 -0.0205 72.9 -0.0598 95.3 0.108 56.7 -3.86 0.54 -0.0527 81.2 +0.00451 70.7 0.0451 94.6 0.0623 56.9 -1.34 0.537 -0.0233 78.6 +0.0402 69.2 0.207 93.1 -0.00208 56.6		-8.28	0.576	-0.104	86.8	-0.0581	79.3	-0.225	96.6	0.175	55.6
-5.62 0.55 -0.0733 83.1 -0.0205 72.9 -0.0598 95.3 0.108 56.7 -3.86 0.54 -0.0527 81.2 +0.00451 70.7 0.0451 94.6 0.0623 56.9 -1.34 0.537 -0.0233 78.6 +0.0402 69.2 0.207 93.1 -0.00208 56.6		-7.09	0.578	-0.0904	85.4	-0.0412	76.5	-0.151	96.2	0.149	56.1
-3.86 0.54 -0.0527 81.2 +0.00451 70.7 0.0451 94.6 0.0623 56.9 -1.34 0.537 -0.0233 78.6 +0.0402 69.2 0.207 93.1 -0.00208 56.6		-5.62	0.55	-0.0733	83.1	-0.0205	72.9	-0.0598	95.3	0.108	56.7
-1.34 0.537 -0.0233 78.6 +0.0402 69.2 0.207 93.1 -0.00208 56.6		-3.86	0.54	-0.0527	81.2	+0.00451	70.7	0.0451	94.6	0.0623	56.9
		-1.34	0.537	-0.0233	78.6	+0.0402	69.2	0.207	93.1	-0.00208	56.6

BMJ Open: first published as 10.1136/bmjopen-2015-009981 on 11 February 2016. Downloaded from http://bmjopen.bmj.com/ on May 11, 2025 at Department GEZ-LTA Erasmushogeschool ≥

in the univariate analysis, once deprivation had been included in the multivariate analysis, higher proportions of white patients were associated with higher mortality, although the coefficient was small. This effect of adjustment for deprivation has been reported previously,^{30–31} and the finding may also be influenced by characteristics of the IMD and, possibly, inconsistencies in the recording of ethnicity. There is debate about the inclusion of measures of deprivation in assessments of clinical performance,³² but we included deprivation in our model in view of its importance in predicting mortality. Characteristics of primary care indicative of the

Characteristics of primary care indicative of the mechanisms in our model also predicted premature ŝ mortality, although less so than population characteristics. Greater numbers of GPs (capacity of primary care) were associated with lower premature mortality, as was access, although for access, the size of effect of access was negligible. This finding supports proposals to increase GP numbers,² but data at practice level on numbers of primary care nurses and other staff were not available. An increase in the numbers of these staff might reduce the need for an increase in the numbers of GPs. Our measure of efficiency-practice size-did not predict mortality, although sensitivity analysis suggested that smaller practice size, when treated as a categorical variable, was associated with higher mortality. It is possible that greater efficiency does not improve care for those most in need, or that practice size was not a ç good measure of efficiency.^{22 23} An alternative measure e of efficiency, such as the experience of the practice manager or the number of nurse-led clinics provided by the practice, was not available, and therefore this finding should be viewed with caution. Neither variations in the measure of continuity nor those in the measure of consultation performance were associated with mortality, but when the effect was related to variations in deprivation, continuity was associated with lower mortality in less deprived practices. Other studies have identified an association between continuity and mortality.^{6 33} It is possible that the negative finding is due to the measures not adequately reflecting the concepts of continuity and consultation performance, but it may also be explained by the modifying effect of deprivation. Greater deprivation was also associated with the reduced effect of GP supply. Detection of hypertension did predict mortality, as found in other studies restricted to coronary heart disease and stroke.^{6 34} The measures of clinical management had only small and **g**. inconsistent relationships to mortality, and were not 8 stable over the 3 years investigated. A potential explanation is the only limited variability in performance between practices that the QOF has engendered. This finding accords with a recent UK study that demonstrated little impact of QOF on mortality.³⁵ A further potential explanation is that an increase in volume may be associated with an improvement in quality of care, in which case the indicators may, to some extent, reflect prevalence.³⁶

Strengths and limitations

This study included an entire country in a practice level analysis, with repetition of the regression model in 3 years to check its stability, and the use of a conceptual model to explain how the mechanisms of primary care influence outcomes. The mechanisms of the model were derived from review of available evidence,³ and the associations found in the regression analysis tend to support several of them. However, the study was crosssectional only, and it is important to recognise that definitive assumptions about causation cannot be made. Our conceptual model was focused on the role of primary care, but community and specialist secondary care, and public health services, also affect health outcomes. Practice level mortality is, therefore, a result of care by these other components of care plus population characteristics, in addition to the specific contribution of primary care.

The majority of data used for the explanatory variables were obtained from sources that have been used for administrative purposes and in research studies for several years, such as the quality and outcomes framework. The data collection systems are well established, and the advantages and disadvantages of these data have become well understood. Nevertheless, there were some potential limitations in the variables used. For example, the response rates to the general practice patient survey tend to be low, although there is evidence to support reliability,²⁵ and alternative measures of access and continuity were not available. Recorded diabetes prevalence is not an ideal measure of the level of chronic disease morbidity in a population; unfortunately, the use of a composite measure derived from a quality and outcome framework smoking indicator was not possible since the measure included hypertension as one of the chronic conditions taken into account. The study used data from five or more years ago, although these data were the most recent available for practice level mortality. The mortality data related to the 2006-2010 period, although we used data on practice characteristics for 2009/2010. After the introduction of the QOF, performance improved during the first 2 or 3 years, but thereafter remained relatively stable,^{37 38} and therefore changes in practice performance are unlikely to have influenced our findings. The study used practice level data, and further research using patient level data would be desirable. Linear regression of age-adjusted rates when the explanatory variables are not age-adjusted can introduce bias. However, sensitivity analysis including age variables did not impact on the size of effect of deprivation as an explanatory variable or size of the effect. Although negative binomial regression of counts of deaths would be preferred,²⁷ the data available meant that this approach was not possible.

Implications and future research

The population characteristics of deprivation and higher levels of morbidity predicted mortality, but the detection

of hypertension and the capacity of primary care, measured in our study by the supply of GPs, also had effects. Although the effects of primary care were small at practice level, the implications for total annual numbers of deaths are important; the associations between detection of hypertension, and numbers of GPs and mortality, suggest that improvement in these aspects of primary care have potential to make worthwhile reductions in mortality. The finding on numbers of GPs supports the case that general practice in England lacks capacity to meet demand. Levels of detection of hypertension have been shown to be associated with the supply of GPs and patient access.³⁹ An increase in the numbers of GPs is required, therefore, not only to improve the conveni- 2 ence of access but also to improve population health. If 8 steps are taken to increase capacity, a longitudinal study of premature mortality should be undertaken to monitor the impact. Since international comparisons rate access in England highly,⁴⁰ our finding is even more important to countries with health systems where access is more limited. Our findings on the role of deprivation, including the reduced impact of primary care mechanisms, indicate that special attention must be given to the needs of deprived populations.⁴¹ In addition to concentration of resources in practices that care for large numbers of deprived people, targeted public health and social care measures have a role to play. These might be facilitated by improving responsiveness to local population characteristics.42

Contributions RB and KH conceived the study. LSL, RB, AGM, TS and KH developed the conceptual model. All the authors took part in designing the study. Data collection was undertaken by KH and RB. Data analysis was undertaken by KH, with contributions by DRJ and MJB. All the authors contributed to interpretation of the findings. The first draft was prepared by RB and all the authors took part in revision of the manuscript and approved the final version. RB and KH are guarantors of the paper.

Funding The study was funded by a grant from Leicester City Clinical Commissioning Group.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Extra data can be accessed via the Dryad data repository at http://datadryad.org/ with the doi:10.5061/dryad.g2t66.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/

REFERENCES

- Macinko J, Starfield B, Shi L. The contribution of primary care systems to health outcomes within Organization for Economic Cooperation and Development (OECD) countries, 1970–1998. *Health Serv Res* 2003;38:831–65.
- NHS England. *Five year forward view*. London: NHS England, 2014.
 Starfield B, Shji L, Macinko J. Contribution of primary care to health
- systems and health. *Milbank Q* 2005;833:457–502.
- 4. Rasella D, Harhay MO, Pamponet ML, et al. Impact of primary health care on mortality from heart and cerebrovascular diseases in

Open Access

Brazil: a nationwide analysis of longitudinal data. BMJ 2014;349: g4014.

- Shi L, Starfield B. The effect of primary care physician supply and 5. income inequality on mortality among blacks and whites in US metropolitan areas. Am J Public Health 2001;91:1246-50.
- Levene LS. Bankart J. Khunti K. et al. Association of primary care 6 characteristics with variations in mortality rates in England: an observational study. PLoS ONE 2012;7: e47800.
- 7 Thorlby R. Reclaiming a population health perspective. Future challenges for primary care. London: The Nuffield Trust. 2013.
- World Health Organization. World health report 2008. Primary health 8. care. Now more than ever. WHO, 2008.
- 9. Baker R, Jones DR, Baker R, et al. Monitoring mortality rates in general practice after Shipman. BMJ 2003;326:274-6.
- Health and Social Care Information Centre. Indicator Portal. https:// 10 indicators.ic.nhs.uk/webview/ (accessed 8 Apr 2015).
- 11. Health and Social Care Information Centre. Dataset: Estimates of Standardised Mortality Rates (SMR) <65 and 75: 2006–2010. https://indicators.ic.nhs.uk/download/GP%20Practice%20data/ Specification/SMR%20v1.pdf (accessed 8 Apr 2015).
- Wheller L, Baker A, Griffiths C. Trends in premature mortality in England and Wales, 1950–2004. *Health Stat Q* 2006;31:34–41. 12.
- Public Health England. Deaths from all causes under 75 years. 13. http://www.apho.org.uk/resource/item.aspx?RID=97049 (accessed 8 Apr 2015)
- Honeyford K, Baker R, Bankart MJG, et al. Estimating smoking 14. prevalence in general practice using data from the Quality and Outcomes Framework (QOF). BMJ Open 2014;4:e005217.
- Communities and Local Government. The English indices of 15. deprivation 2010. London: Department for Communities and Local Government, 2011. ISBN: 978-1-4098-2924-9
- 16. Health and Social Care Information Centre. Estimates of Index of Multiple Deprivation (IMD) 2010 for GP practices. December 2011. https://indicators.ic.nhs.uk/webview/index/en/MvServer/ NHS-Information-Centre-indicators.c.MyServer/GP-Practice-data.d. 48/Demography.d.77/Deprivation.d.202/Estimates-of-Index-of-Multiple-Deprivation-IMD-2010-for-GP-practices/fStudy/P01112 (accessed 9 Sep 2015).
- Martin D, Wright JA. Disease prevalence in the English population: 17 a comparison of primary care registers and prevalence models. Soc Sci Med 2009;68:266-74.
- Barnett K, Mercer SW, Norbury M, et al. Epidemiology of 18 multimorbidity and implications for health care, research, and medical education: a cross-sectional study. Lancet 2012;380:37-43.
- GP Patient Survey. Surveys and reports. https://gp-patient.co.uk/ 19 surveys-and-reports (accessed 8 Apr 2015).
- Healthcare Commission. Practice ethnicity data for disease 20 prevalence modeling. Eastern Region Public Health Observatory, 2010
- 21. Health and Social Care Information Centre. Number of GPs per registered and weighted practice population. 2010. https://indicators. ic.nhs.uk/download/Infrastructure/Specification/GP%20numbers% 20v2.pdf (accessed 8 Apr 2015).
- Gulliford MC, Jack RH, Adams G, et al. Availability and structure of 22 primary medical care services and population health and health care indicators in England. BMC Health Serv Res 2004;4:12.
- Szczepura A, Davies C, Fletcher J, et al. Efficiency and 23
- effectiveness in general practice. *J Manag Med* 1993;7:36–47. Gulliford M, Figueroa-Munoz J, Morgan M, *et al.* What does 'access 24 to health care' mean?. J Health Serv Res Policy 2002;7:186-8.

- Roland M. Elliott M. Lyratzopoulos G. et al. Reliability of patient 25 responses in pay for performance schemes: analysis of national General Practitioner Patient Survey data in England. BMJ 2009;339: b3851
- Ashworth M, Schofield P, Doran T, et al. The Public Health Impact 26 score: a new measure of public health effectiveness for general practices in England. Br J Gen Pract 2013;63:e291-9.
- 27. Honeyford K, Baker R, Bankart MJG, et al. Modelling factors in primary care quality improvement: a cross-sectional study of premature CHD mortality. BMJ Open 2013;3:e003391.
- Calderón-Larrañaga A, Soljak M, Cecil E et al. Does higher quality 28 of primary healthcare reduce hospital admissions for diabetes complications? A national observational study. Diabet Med 2014;31:657-65.
- Chatterjee G, Hadi AS. Regression analysis by example. 5th edn. 29 Hoboken, NJ: John Wiley and Sons, 2012.
- 30 Sundquist J, Bajeka M, Jarman B, et al. Underprivileged area score, ethnicity, social factors and general mortality in district health authorities in England and Wales. Scand J Prim Health Care 1996.14.79-85
- Association of Public Health Observatories. Indications of Public 31. Health in the English Regions. 4: Ethnicity and Health. http://www. apho.org.uk/resource/view.aspx?QN=IND SERIES (accessed 11 Nov 2015).
- Fiscella K, Burstin HR, Nerenz DR. Quality measures and 32 sociodemographic risk factors: to adjust or not to adjust. JAMA 2014;312:2615-16.
- 33 Shin DW, Cho J, Yang HK, et al. Impact of continuity of care on mortality and health care costs: a nationwide cohort study in Korea. Ann Fam Med 2014;12:534-41.
- 34 Levene SL, Baker R, Bankart MJ, et al. Association of features of primary care with coronary heart disease mortality. JAMA 2010:304:2028-34
- Kontopantelis E, Springate DA, Ashworth M, et al. Investigating the 35 relationship between quality of primary care and premature mortality in England: a spatial whole-population study. BMJ 2015;350:h904.
- 36 Millett C, Car J, Eldred D, et al. Diabetes prevalence, process of care and outcomes in relation to practice size, caseload and deprivation: national cross-sectional study in primary care. J R Soc Med 2007;100:275-83.
- Doran T, Kontopantelis E, Valderas JM, et al. Effect of financial 37. incentives on incentivised and non-incentivised clinical activities: longitudinal analysis of data from the UK Quality and Outcomes Framework. BMJ 2011;342:d3590.
- Doran Tim, Kontopantelis Evangelos, Reeves David, et al. Setting 38 performance targets in pay for performance programmes: what can we learn from QOF? 2014;348:g1595.
- Bankart MJ, Anwar MS, Walker N, et al. Are there enough GPs in 39 England to detect hypertension and maintain access? A cross-sectional study. Br J Gen Pract 2013;63:e339-44.
- 40 Thomson S, Osborn R, Squires D, et al. International profiles of health care systems. New York: The Commonwealth Fund, 2013.
- Marmot Review. Fair Society, Healthy Lives: Strategic Review of 41 Health Inequalities in England Post 2010. London: Marmot Review; 2010
- Tarrant C, Angell E, Baker R, et al. Responsiveness of primary care 42. services: development of a patient-report measure. Health Services and Delivery Research Programme. HMSO 2014. http://www.nets. nihr.ac.uk/__data/assets/pdf_file/0009/117378/FLS-09-1801-1029. pdf (accessed 24 Apr 2015).

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies