

Nutrition Users' Guides: an introduction to structured guides to evaluate the nutrition literature

Bradley C Johnston ^{1,2} Mary Rozga,³ Gordon H Guyatt,^{4,5} Rosa K Hand,^{6,7} Deepa Handu,³ Kevin C Klatt,⁸ Malgorzata M Bala⁹

To cite: Johnston BC, Rozga M, Guyatt GH, *et al.* Nutrition Users' Guides: an introduction to structured guides to evaluate the nutrition literature. *BMJ Nutrition, Prevention & Health* 2025;**0**. doi:10.1136/bmjnp-2023-000832

For numbered affiliations see end of article.

Correspondence to

Dr Bradley C Johnston;
bradley.johnston@tamu.edu

Received 20 November 2023
Accepted 11 March 2025

ABSTRACT

Despite evidence that nutrition can play a substantial role in curbing the burden of chronic disease, findings reported in the nutrition literature have been plagued with debate and uncertainty, including questions about the confidence we can place in evidence from observational studies, the validity of dietary intake data, and the applicability of randomised trials to real-world patients or members of the public. Structured nutrition users' guides (NUGs) to evaluate common research study designs (ie, randomised trials, cohort studies, systematic reviews and clinical practice guidelines) addressing nutrition questions will help clinicians and their patients, as well as health service workers and policy-makers, use the evidence to make more informed decisions on disease management and prevention. In addition, NUGs will provide comprehensive teaching materials for nutrition trainees on how to appraise, interpret and apply the research evidence. We hereby introduce a series of structured NUGs for the literature on nutrients, foods and dietary patterns and programmes. Each article will address three key components when assessing different study designs used to assess nutrition interventions or exposures, including (1) assessing the methodological quality of the study, (2) interpreting study results (magnitude and precision of treatment or exposure effects for outcomes of benefit and harm) and (3) applying the results to unique patient or population scenarios based on their health-related values and preferences related to the potential benefits, harms, convenience and cost of an intervention. This series of articles will serve to empower clinicians, health service workers and health policy-makers to better understand the validity, interpretability and applicability of the nutrition literature, while also helping practitioners and their clients make more evidence-based, value-sensitive and preference-sensitive nutrition decisions.

Worldwide, chronic non-communicable illnesses represent the largest burden of disease. There is good evidence that healthy eating habits represent one of the cornerstones to primary prevention of chronic disease.^{1–3} Nutrition-related deficiencies and disease may have a substantial impact on disability-adjusted life years and death. For instance, one 2017 estimate suggested that dietary habits accounted for 10.9 million deaths globally.⁴

Despite the evidence that nutrition can play an important role in curbing the burden of chronic disease, nutrition plays only a minor role in most clinicians' practice. For those trained in medicine, providing nutritional counselling in clinical practice comes with a unique set of challenges, including a lack of nutrition training in medical school, lack of time and lack of financial compensation for offering such counselling.^{5,6} For those trained in nutrition, there may be considerable variability in evidence-based practice training.⁷

Providing the highest-quality nutrition care depends on a clinicians' knowledge and skills in evidence-based practice, including skills to ask a structured question and acquire, appraise, interpret and apply the best available evidence within the context of clinical expertise and patient values and preferences.^{7–9} However, dietitians, physicians and other healthcare workers may lack these skills,^{7,10,11} with a recent literature review finding no studies among dietitians that have reported on their competencies in interpreting the absolute magnitude (size) of effect, or the certainty of evidence in effect, competencies essential for optimising clinical nutrition decision-making.⁷

For those who lack evidence-based practice skills, or the time to use their skills, pre-appraised and pre-interpreted evidence by methodologists allows clinicians to more accurately inform and guide their patients with clinical decision-making. While many comprehensive up-to-date resources with pre-appraised/interpreted evidence in medicine exist (eg, UpToDate, Dynamed, McMaster Plus; BMJ Best Practice), pre-appraised/interpreted evidence specific to clinical and public health nutrition (ie, Evidence Analysis Library, Practice-based Evidence in Nutrition) are fewer and not always up to date.

In healthcare practice and policy, both clinicians and authorities have accepted



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

the importance of evidence-based practice, including shared decision-making informed by the best available evidence as well as patient or population values and preferences. This is no less true for nutrition care, for which clinicians and policy-makers must consider the best available evidence and its bearing on unique nutrition-related decisions based on the values and preferences of the target patient or population.^{9 12} Value-sensitive and preference-sensitive decisions must ultimately follow from weighing the trade-offs between potential benefits, harms and costs of alternative treatment or prevention strategies, the certainty of evidence informing estimates of these key considerations, and the ability of people to implement and adhere to a nutrition or diet recommendation, particularly if burdensome.

We anticipate that this series of articles will provide a welcome and thought-provoking set of structured users' guides for clinicians and healthcare professionals (eg, health service workers, policy-makers) using nutrition research articles to guide practice and policy, a series of articles that can also be used to support informed, shared decision-making with patients and clients. The series will include guidance on how to appraise, interpret and apply randomised trials, cohort studies, systematic reviews and practice guidelines, while also laying the foundation for future articles considering emerging approaches to nutrition and medical research (eg, Mendelian randomisation; platform and adaptive trials; planetary health). As compared with a variety of other "appraisal" checklists used in nutrition that focus almost exclusively on methodological quality and/or risk of bias (ie, appraisal), particularly unique to our framework is the guidance on interpreting and applying study results including interpreting the absolute magnitude of treatment effect (eg, from trivial to small, moderate or large) for people-important health outcomes^{7 13 14} and applying the evidence together with patient or client values and preferences. A third component of evidence-based practice is health service expertise. For example, expert, experienced clinicians play an important role in interpreting the evidence, including laboratory results, and guiding patients to make informed decisions. Ideally, clinicians are highly skilled in the principles of evidence-based practice⁹ and are particularly skilled in determining the best available evidence (eg, systematic reviews, umbrella reviews, practice guidelines) and engaging with patients based on their unique circumstances, including the patients' health-related values and preferences. A clinical practice based on expertise alone however may, in many instances, be less trustworthy given that clinicians (ie, humans) are prone to many cognitive biases and to clinical time constraints.^{5 6}

With respect to nutrition users' guide articles on summary evidence (systematic reviews, guidelines), this series will adhere to guidance from the Grading of Recommendations Assessment, Development and Evaluation (GRADE) working group that, in a four-category classification of certainty of evidence (high, moderate, low and very low), considers systematic reviews of randomised

controlled trials (RCTs) to start at high certainty evidence, with guidance on five factors (ie, risk of bias, indirectness, inconsistency, imprecision, publication bias) that can justify rating the certainty of evidence down.^{15 16} Observational studies typically start at low certainty evidence,^{9 15} or if they start at high (eg, Risk Of Bias In Nonrandomized Studies of Interventions risk of bias tool), they are typically downgraded for serious issues of bias.¹⁷ The reason for this guidance is that observational studies inevitably suffer from a risk of substantial residual confounding¹⁸ and risk of publication bias,¹⁹ and only in the presence of large associations (eg, relative risk >2 or <0.5) or valid dose-response gradients can they provide higher certainty evidence, provided there are no other serious²⁰ limitations (table 1).

Nutrition evidence is unique in that RCTs may be absent for many clinical or public health nutrition questions, many nutrition RCTs lack genuine placebos and are challenging to blind, RCTs of dietary interventions may be challenging to adhere to and may not be long enough to capture impacts on hard endpoints (eg, mortality), and, unlike pharmaceuticals, many interventions aim to modify existing variables that participants are exposed to.⁹ Thus, some have argued that evidentiary standards should differ for nutrition, and that prospective cohort studies may sometimes be as helpful as RCTs in judging estimates of effect of therapy or exposure.²¹⁻²⁵ Indeed, the GRADE method includes provisions that authors of systematic reviews based on observational studies may rate up the certainty of evidence in certain situations (table 1), resulting in moderate or high certainty evidence, when justified typically based on large associations or a credible dose-response gradient.^{18 20} The GRADE method also includes provisions for downgrading systematic reviews of RCTs²⁶ to account for some of the inherent challenges in nutrition interventions that compromise the certainty of evidence (eg, risk of bias related to poor or infrequent dietary intake assessments, and indirectness when the target intervention is a nutrient evaluated in pragmatic trials that are not isocaloric).

GRADE also provides guidance on moving from the best available evidence (ie, systematic reviews) to decision-making (ie, guideline recommendations) using an evidence-to-decision framework.²⁷ When, for example, it is clear that the desirable outcomes (benefits) outweigh undesirable outcomes (harms) and when the certainty of evidence is high, a strong recommendation is typically most appropriate, particularly if all or almost all patients or members of the public would be willing to take the intervention based on a systematic summary of their values and preferences. If there is a close balance between the benefits and the harms and the certainty of evidence is low to moderate, a conditional (or weak) recommendation is most appropriate. With conditional recommendations, shared decision-making is particularly encouraged, and decisions should be driven by the patients or target populations' values and preferences (box 1).

Table 1 Grading of Recommendations Assessment, Development and Evaluation approach to determine certainty of evidence for each outcome (systematic reviews)³⁸

Study design	(1) Establish initial level of certainty	(2) Consider lowering or raising level of certainty		(3) Final level of certainty rating
	Initial certainty in an estimate of effect	RCTs can be rated down if serious issues with:	Observational studies can be rated up if robust evidence of:	Certainty in an estimate of effect across all considerations graded as:
Randomised controlled trials (RCTs)	High certainty (RCTs start at high certainty & can be rated down)	Risk of bias	Large effect	⊕⊕⊕⊕ High
Observational studies eg cohort, case-control	Low certainty (observational studies start at low certainty & can be rated up)*	Inconsistency	Dose response	⊕⊕⊕○ Moderate
		Indirectness	All plausible confounding & bias would reduce a demonstrated effect, or would suggest a spurious effect if no effect was observed	⊕⊕○○ Low
		Imprecision		⊕○○○ Very low
		Publication bias		

Certainty of estimates of effect for each health outcome or indicator (surrogate):
High: we are very certain that the true effect lies close to that of the estimate of the effect.
Moderate: we are moderately certain in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.
Low: our certainty in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.
Very low: we have very little certainty in the effect estimate: the true effect is likely to be substantially different from the estimate of the effect.

Certainty of evidence is a continuum; any discrete categorisation involves some degree of arbitrariness. Nevertheless, advantages of simplicity, transparency and vividness outweigh these limitations.

*Alternatively, if a systematic review team uses the Risk Of Bias In Nonrandomized Studies of Interventions/Exposures tools to assess a body of evidence of non-randomised studies, the body of evidence starts at high certainty, and is assessed for potential study limitations, typically resulting in rating down due to serious risk of bias issues.

Those who support applying the GRADE approach to nutrition research recognise that biases, such as unaccounted confounders in observational studies, often impact the validity of effect estimates.^{28–30} For instance, a meta-analysis of nine cohort studies including over 290 000 patients claimed a 25% (95% CI 7% to 40%) relative risk reduction in coronary heart disease among men consuming supplemental vitamin C.³¹ In contrast, the Physicians' Health Study II,³² a factorial RCT that enrolled over 14 500 male physicians and followed them for upwards of 10 years, demonstrated no difference in coronary heart disease when comparing vitamin C versus placebo (HR 0.99; 95% CI 0.89 to 1.11). These same studies showed discrepant results between cohort and randomised trials for vitamin E and cardiovascular disease. These examples demonstrate there were confounding variables that accounted for the 25% relative risk reduction in heart disease associated with vitamin C reported in observational studies assessed in the meta-analysis.

Nutrition content expertise is needed to apply the GRADE approach to nutrition research, as interpretation and application require knowledge of topics such as nutritional biochemistry and/or physiology, the strengths and

limitations of nutrition assessment methods (ie, dietary intake data, biomarkers), and knowledge of behavioural science. For instance, in well-designed and conducted nutrition studies, known prognostic factors should include both baseline dietary intakes and indicators of baseline nutrient status if potentially valid biomarkers exist (eg, red blood cell omega-3 fatty acids status, or 25-hydroxyvitamin D status in an omega-3 or vitamin D intervention study).³³ Few observational studies or RCTs in nutrition quantify the exposure of interest present at baseline in the diet and/or endogenously in tissues. Failure to consider baseline intake and/or status as a component of the inclusion criteria or as pre-specified subgroup analysis may put the results of a study at risk of bias, particularly when attempting to assess the effectiveness of a nutrient relative to baseline nutrient status, such as the utility of vitamin D supplementation in those with lower baseline vitamin D status. Indeed, large randomised controlled trials aiming to answer such questions have typically recruited participants irrespective of their baseline vitamin D status, ultimately recruiting individuals with average baseline 25-hydroxyvitamin D levels already in the observed 'protective' range.³⁴

Box 1 GRADE approach to determining strength of recommendation (guidelines)³⁸

Strong recommendation

1. High certainty evidence
2. Desirable (benefits) consequences clearly outweigh undesirable consequences (harms), or vice versa
3. All or almost all fully informed individuals in the target population would choose the recommended course of action.
4. Wording: “We recommend...” “for intervention”, or “against intervention”

Conditional (weak) recommendation

1. Low certainty evidence
2. Close balance between undesirable and desirable consequences.
3. Many fully informed individuals in the target population would choose the recommended course of action but a substantial minority would not.
4. Need for shared decision-making
5. Wording: “We suggest...” “for intervention”, or “against intervention”

Note: When guideline panellists are faced with predominantly ‘moderate’ certainty evidence, conditional recommendations are more likely; however, it is at the discretion of the panel and ideally driven by value and preference data (ie, for a strong recommendation would all or almost all fully informed individuals choose the recommended course of action).

Another nuance of nutrition interventions is that many nutrition interventions lack an inert placebo to compare against. Rather, modifying intakes of specific foods typically occurs via their replacement for other foods (ie, the replacement of saturated fat-rich foods with polyunsaturated fat-rich foods), leading to an inherent substitution effect. Specific substitutions may not be explicit in the intervention’s design, complicating interpretation of observed effects without rigorous dietary assessment methods to identify modifications. Ultimately, multiple studies are typically needed with consistent results to provide high confidence in the relationship between specific foods/nutrients and health outcomes, accounting for their effects relative to various potential comparator foods/nutrients. These unique elements of nutrition interventions are critical to consider in systematic reviews and meta-analyses of the literature, as many similar interventions captured by inclusion criteria may have resulted in variable substitutions and may have occurred in populations with varying baseline intakes/status, resulting in varying results.

As the field of nutrition continues to tackle diet-chronic disease relationships, and embraces ‘precision nutrition’ in an effort to explain inter-individual variation in the response to nutrition interventions, it is critical to ensure evidentiary standards are methodologically robust and transparent (including the need for study protocols for observational studies) to ensure nutrition recommendations provided to the patient or members of the public are optimally trustworthy. Overall, we anticipate that the guidance that this series generates will ultimately further the understanding and discussion of how the nutritional

literature can best guide clinical and population health practice considering issues of validity (risk of bias), the magnitude and precision of exposure or treatment effects on health outcomes, and the applicability of study results to the target patient or population.^{35–37} This series can also be used by those designing courses in critical appraisal and evidence-based nutrition practice, while also serving to empower health service workers, guideline panellists and policy-makers to better understand the strengths and limitations of the nutrition literature.

Author affiliations

¹Department of Nutrition, College of Agriculture and Life Sciences, Texas A&M University, College Station, Texas, USA

²Department of Epidemiology and Biostatistics, School of Public Health, Texas A&M University, College Station, Texas, USA

³Academy of Nutrition and Dietetics, Chicago, Illinois, USA

⁴Department of Medicine, McMaster University, Hamilton, Ontario, Canada

⁵Department of Health Research Methods, Evidence and Impact, McMaster University, Hamilton, Ontario, Canada

⁶Department of Nutrition, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA

⁷Centre for Medical Education, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA

⁸Department of Nutritional Science and Toxicology, University of California Berkeley, Berkeley, California, USA

⁹Chair of Epidemiology and Preventive Medicine, Department of Hygiene and Dietetics, Jagiellonian University Medical College, Krakow, Małopolskie, Poland

X Bradley C Johnston @methodsnerd and Kevin C Klatt @KCKlatt

Contributors BJ conceptualised and drafted the paper. All authors provided critical feedback and suggestions on different iterations of the manuscript. All authors reviewed the semifinal version and approved the final version for publication.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests BCJ has received funds from Texas A&M AgriLife Research to fund investigator-initiated research related to saturated and polyunsaturated fats. The grant was from Texas A&M AgriLife institutional funds from interest and investment earnings, not a sponsoring organisation, industry, or company. BCJ has also received funds from National Institute of Diabetes, Digestive and Kidney Diseases (NIDDK) R25 program to support training in evidence-based nutrition practice. All other authors claim no relevant disclosures.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; internally peer reviewed by Dr Katharine Martyn, University of Brighton, UK.

Data availability statement Data sharing not applicable as no datasets generated and/or analysed for this study.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Bradley C Johnston <http://orcid.org/0000-0001-8872-8626>

REFERENCES

- 1 Karam G, Agarwal A, Sadeghirad B, *et al.* Comparison of seven popular structured dietary programmes and risk of mortality and major cardiovascular events in patients at increased cardiovascular risk: systematic review and network meta-analysis. *BMJ* 2023;380:e072003.

- 2 Afshin A, Sur PJ, Fay KA. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2019;393:1958–72.
- 3 Wang X, Ouyang Y, Liu J, *et al.* Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer: systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ* 2014;349:g4490.
- 4 Global Burden of Disease 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018;392:1923–94.
- 5 Johansson M, Guyatt G, Montori V. Guidelines should consider clinicians' time needed to treat. *BMJ* 2023;380:e072953.
- 6 Privett N, Guerrier S. Estimation of the Time Needed to Deliver the 2020 USPSTF Preventive Care Recommendations in Primary Care. *Am J Public Health* 2021;111:145–9.
- 7 Ghosh NR, Esmaeilnezhad Z, Zajac J, *et al.* Evidence-Based Practice Competencies among Nutrition Professionals and Students: A Systematic Review. *J Nutr* 2024;154:1414–27.
- 8 Hand RK, Davis AM, Thompson KL, *et al.* Updates to the Definition of Evidence-Based (Dietetics) Practice: Providing Clarity for Practice. *J Acad Nutr Diet* 2021;121:1565–73.
- 9 Johnston BC, Seivenpiper JL, Vernooij RWM, *et al.* The Philosophy of Evidence-Based Principles and Practice in Nutrition. *Mayo Clin Proc Innov Qual Outcomes* 2019;3:189–99.
- 10 Hand RK, Abram JK. Sense of Competence Impedes Uptake of New Academy Evidence-Based Practice Guidelines: Results of a Survey. *J Acad Nutr Diet* 2016;116:695–705.
- 11 Johnston BC, Alonso-Coello P, Friedrich JO, *et al.* Do clinicians understand the size of treatment effects? A randomized survey across 8 countries. *CMAJ* 2016;188:25–32.
- 12 Watowicz RP, Hand RK. Call for Standardized Language and Training to Bridge the Appraise/Apply Gap in Evidence-Based Dietetics Practice. *J Acad Nutr Diet* 2023;123:1121–6.
- 13 Tikkinen KAO, Guyatt GH. Understanding of research results, evidence summaries and their applicability-not critical appraisal-are core skills of medical curriculum. *BMJ Evid Based Med* 2021;26:231–3.
- 14 Alonso-Coello P, Carrasco-Labra A, Brignardello-Petersen R, *et al.* Systematic reviews experience major limitations in reporting absolute effects. *J Clin Epidemiol* 2016;72:16–26.
- 15 Guyatt GH, Oxman AD, Vist GE, *et al.* GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924–6.
- 16 Guyatt GH, Oxman AD, Schünemann HJ, *et al.* GRADE guidelines: a new series of articles in the Journal of Clinical Epidemiology. *J Clin Epidemiol* 2011;64:380–2.
- 17 Schünemann HJ, Cuello C, Akl EA, *et al.* GRADE guidelines: 18. How ROBINS-I and other tools to assess risk of bias in nonrandomized studies should be used to rate the certainty of a body of evidence. *J Clin Epidemiol* 2019;111:105–14.
- 18 Verbeek JH, Whaley P, Morgan RL, *et al.* An approach to quantifying the potential importance of residual confounding in systematic reviews of observational studies: A GRADE concept paper. *Environ Int* 2021;157:106868.
- 19 Ioannidis JPA. The importance of potential studies that have not existed and registration of observational data sets. *JAMA* 2012;308:575–6.
- 20 Guyatt GH, Oxman AD, Sultan S, *et al.* GRADE guidelines: 9. Rating up the quality of evidence. *J Clin Epidemiol* 2011;64:1311–6.
- 21 Mann JI. Evidence-based nutrition: Does it differ from evidence-based medicine. *Ann Med* 2010;42:475–86.
- 22 Micha R, Mozaffarian D. Saturated fat and cardiometabolic risk factors, coronary heart disease, stroke, and diabetes: a fresh look at the evidence. *Lipids* 2010;45:893–905.
- 23 Harris WS, Mozaffarian D, Lefevre M, *et al.* Towards establishing dietary reference intakes for eicosapentaenoic and docosahexaenoic acids. *J Nutr* 2009;139:804S–19S.
- 24 Schwingshackl L, Knüppel S, Schwedhelm C, *et al.* Perspective: NutriGrade: A Scoring System to Assess and Judge the Meta-Evidence of Randomized Controlled Trials and Cohort Studies in Nutrition Research. *Adv Nutr* 2016;7:994–1004.
- 25 Meerpohl JJ, Naude CE, Garner P, *et al.* Comment on “Perspective: NutriGrade: A Scoring System to Assess and Judge the Meta-Evidence of Randomized Controlled Trials and Cohort Studies in Nutrition Research”. *Adv Nutr* 2017;8:789–90.
- 26 Balshem H, Helfand M, Schünemann HJ, *et al.* GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011;64:401–6.
- 27 Alonso-Coello P, Schünemann HJ, Moher J, *et al.* GRADE Evidence to Decision (EtD) frameworks: a systematic and transparent approach to making well informed healthcare choices. 1: Introduction. *BMJ* 2016;353:i2016.
- 28 Zeraatkar D, Johnston BC, Guyatt G. Evidence Collection and Evaluation for the Development of Dietary Guidelines and Public Policy on Nutrition. *Annu Rev Nutr* 2019;39:227–47.
- 29 Zähringer J, Schwingshackl L, Movsisyan A, *et al.* Use of the GRADE approach in health policymaking and evaluation: a scoping review of nutrition and physical activity policies. *Implement Sci* 2020;15:37.
- 30 Vernooij R, Guyatt GH, Zeraatkar D, *et al.* Reconciling contrasting guideline recommendations on red and processed meat for health outcomes. *J Clin Epidemiol* 2021;138:215–8.
- 31 Knekt P, Ritz J, Pereira MA, *et al.* Antioxidant vitamins and coronary heart disease risk: a pooled analysis of 9 cohorts. *Am J Clin Nutr* 2004;80:1508–20.
- 32 Sesso HD, Buring JE, Christen WG, *et al.* Vitamins E and C in the prevention of cardiovascular disease in men: the Physicians' Health Study II randomized controlled trial. *JAMA* 2008;300:2123–33.
- 33 Brouwer-Brolsma EM, Brennan L, Dreven CA, *et al.* Combining traditional dietary assessment methods with novel metabolomics techniques: present efforts by the Food Biomarker Alliance. *Proc Nutr Soc* 2017;76:619–27.
- 34 Manson JE, Cook NR, Lee I-M, *et al.* Vitamin D Supplements and Prevention of Cancer and Cardiovascular Disease. *N Engl J Med* 2019;380:33–44.
- 35 Zeraatkar D, de Souza RJ, Guyatt GH, *et al.* Nutrition users' guides: systematic reviews Part 1 -structured guide for methodological assessment, interpretation and application of systematic reviews and meta-analyses of non-randomised nutritional epidemiology studies. *BMJ Nutr Prev Health* 2024;7:e000835.
- 36 Bala MM, Agarwal A, Klatt KC, *et al.* Nutrition Users' Guides: RCTs Part 2 - structured guide for interpreting and applying study results from randomised controlled trials on therapy or prevention questions. *BMJ Nutr Prev Health* 2024;7:e000834.
- 37 Agarwal A, Bala MM, Zeraatkar D, *et al.* Nutrition users' guides: RCTs part 1 - structured guide for assessing risk of bias in randomised controlled trials that address therapy or prevention questions. *BMJ Nutr Prev Health* 2024;7:e000833.
- 38 Schünemann H, Brożek J, Guyatt G, *et al.* GRADE handbook for grading quality of evidence and strength of recommendations (updated October 2013). 2023. Available: <https://gdt.gradeapro.org/app/handbook/handbook.html> [Accessed 21 Dec 2024].