

Prenatal ambient air pollution exposure and the risk of stillbirth: systematic review and meta-analysis of the empirical evidence

Nazeeba Siddika,¹ Hamudat A Balogun,¹ Adeladza K Amegah,^{1,2}
Jouni J K Jaakkola^{1,3}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/oemed-2015-103086>).

¹Faculty of Medicine, Center for Environmental and Respiratory Health Research, University of Oulu, Oulu, Finland

²Department of Biomedical and Forensic Sciences, University of Cape Coast, Cape Coast, Ghana

³Medical Research Center, University of Oulu and Oulu University Hospital, Oulu, Finland

Correspondence to

Prof Jouni J K Jaakkola, Faculty of Medicine, Center for Environmental and Respiratory Health Research, Aapistie 5B, PO Box 5000, University of Oulu, Oulu 90014, Finland; jouni.jaakkola@oulu.fi

Received 21 May 2015

Revised 22 January 2016

Accepted 5 February 2016

Published Online First

24 May 2016

ABSTRACT

Background Individual studies on the relations between ambient air pollution and the risk of stillbirth have provided contradictory results. We conducted a systematic review and meta-analysis to summarise the existing evidence.

Methods We conducted a systematic search of three databases: PubMed, Scopus and Web of Science, from their time of inception to mid-April, 2015. Original studies of any epidemiological design were included. Data from eligible studies were extracted by two investigators. To calculate the summary effect estimates (EE), the random effects model was used with their corresponding 95% CI.

Results 13 studies met the inclusion criteria. Although not reaching statistical significance, all the summary effect estimates for the risk of stillbirth were systematically elevated in relation to mean prenatal exposure to NO₂ per 10 ppb (EE=1.066, 95% CI 0.965 to 1.178, n=3), CO per 0.4 ppm (EE=1.025, 95% CI 0.985 to 1.066, n=3), SO₂ per 3 ppb (EE=1.022, 95% CI 0.984 to 1.062, n=3), PM_{2.5} per 4 µg/m³ (EE=1.021, 95% CI 0.996 to 1.046, n=2) and PM₁₀ per 10 µg/m³ (EE=1.014, 95% CI 0.948 to 1.085, n=2). The effect estimates for SO₂, CO, PM₁₀ and O₃ were highest for the third trimester exposure. Two time series studies used a lag term of not more than 6 days preceding stillbirth, and both found increased effect estimates for some pollutants.

Conclusions The body of evidence suggests that exposure to ambient air pollution increases the risk of stillbirth. Further studies are needed to strengthen the evidence.

INTRODUCTION

Ambient air pollution is a major environmental health problem in developed and in developing countries,¹ and is a major cause of several important diseases including lung cancer,² acute lower respiratory infections,³ cardiovascular diseases,⁴ pregnancy outcomes,⁵ chronic obstructive pulmonary disease COPD⁶ and asthma.⁷ The WHO reports that in 2012⁸ around 7 million people died—one in eight of total global deaths was as a result of air pollution exposure, 3.7 million of these were attributed to ambient air pollution due to exposure to particulate matter (PM₁₀).¹ This estimate is more than twofold of the previous estimates and suggests that air pollution is now the world's largest single environmental health risk.¹

Fetal growth and pregnancy outcome are determined by several factors including maternal

What this paper adds

- Previous reviews on the relations between prenatal exposure to air pollution and the risk of stillbirth were based on only three studies and were inconclusive. Emergence of a substantial number of studies since the last review, called for re-evaluation of the existing evidence.
- This systematic review and meta-analysis suggests an elevated risk of stillbirth in relation to air pollution, although further studies are needed to strengthen the evidence.
- Policies such as control of vehicular emissions, fuel quality improvement and control of industrial waste emission, should be developed and implemented to reduce the risk of air pollutants.

nutrition, environmental exposures and heredity. The prenatal stage of life is a very sensitive period such that exposure to harmful substances can have an adverse effect on the developing fetus. The effects of air pollution on fetal growth and pregnancy outcomes have been studied especially in developed countries and the results have been summarised in several reviews.^{9–12}

The WHO reported that 2.6 million stillbirths occurred worldwide in 2009, according to the first comprehensive set of estimates published in a special series of *Lancet*, 2011. Every day, more than 7200 babies are stillborn.¹³ Previous studies have identified the important causes of stillbirth as umbilical cord accidents, congenital anomalies, placental abruption, maternal disease (diabetes, HIV, syphilis and hypertension), obesity, primiparity and smoking in pregnancy.¹⁴

Two previous reviews have suggested that ambient air pollutant exposure including nitrogen dioxide (NO₂), sulfur dioxide (SO₂), black carbon, carbon monoxide (CO), polycyclic aromatic hydrocarbons (PAH's) and particulate matter (PM), can also be an important cause of stillbirth, but both concluded that the evidence was weak at the time of conduct. There are also recent reviews indicating that air pollution from secondhand smoke¹⁵ and solid fuels¹⁶ increases the risk of stillbirth.

The two previous reviews of the effects of ambient air pollution evaluated three studies each, two of which were common to both reviews, with



► <http://dx.doi.org/10.1136/oemed-2016-103625>



CrossMark

To cite: Siddika N, Balogun HA, Amegah AK, et al. *Occup Environ Med* 2016;**73**:573–581.

Glinianaia *et al*¹⁷ stating that the evidence available is insufficient to assess a possible association between PM and stillbirth. Lacasana *et al*¹⁸ reported a positive but not consistent association between ambient air pollution and stillbirth; this may be due to an insufficient number of studies available. A recent review by Zhu *et al*¹⁹ evaluated the effect of exposure to PM_{2.5} on pregnancy outcomes but only included one study on stillbirth. A substantial number of studies have emerged since the conduct of these reviews, and this certainly calls for an evaluation of the evidence to provide insight into causality and identify gaps in knowledge. The objective of the present study was to assess the effect of prenatal ambient air pollution exposure on the risk of stillbirth through systematic review and meta-analysis.

METHODS

A systematic search of three databases—PubMed, Scopus and Web of Science—was carried out from their time of inception to mid-April, 2015, using Medical Subject Heading (MeSH) terms without any language restriction. The search terms used are listed in table 1.

Two investigators (NS and HAB) independently searched for relevant studies from the databases by first screening the titles of the citations and subsequently reviewing the abstracts of relevant titles. Studies selected after the abstract review were retrieved in full and reviewed with articles selected for inclusion in the study satisfying the following criteria: (1) original articles of any epidemiological design; (2) conducted in a human population; (3) provided effect estimates for the relation between exposure to any outdoor air pollutant and the risk of stillbirth, or reported the occurrence of stillbirth among exposed and unexposed mothers. The reference lists of the included studies were also reviewed to identify additional eligible studies.

Data extraction and study quality appraisal

A data extraction form adapted from our previous review¹⁶ was used independently by two investigators (NS and HAB) to extract the relevant information from the studies meeting the inclusion criteria. The information extracted by the two investigators was compared with any differences resolved by repeatedly checking the original articles and through discussion with the third investigator (AKA) adjudicating in situations where there were disagreements. Methodological quality of the

included studies was assessed by investigating evidence of selection, information and confounding bias, and evaluation of the case ascertainment protocols. The general quality of the studies was assessed using the Newcastle-Ottawa Scale (NOS).²⁰

Statistical analysis

Owing to differences in study design, geographical settings and different study population, we anticipated the inconsistency in the studies and applied the random effect model in summary effect estimates with their corresponding 95% CI. Heterogeneity of the studies was assessed using the I² statistic, with a value >50% being deemed to indicate high heterogeneity, 25–50% indicating moderate and <25% indicating low heterogeneity. Individual studies included in the meta-analysis had their effect estimates with different magnitude of air pollutant exposure (except for NO₂ and PM₁₀), therefore, before estimating the summary effects the individual effect estimates were converted with a common pollutant concentration such as per 3 ppb increase in SO₂, per 0.4 ppm in CO, 4 µg/m³ in PM_{2.5} and 10 ppb increase in O₃ exposure. With regard to the study providing only trimester specific effect estimates, we first combined the three (1st, 2nd and 3rd trimester) estimates using the fixed effects model to get the estimate for the entire pregnancy, and then applied the combined estimate in the overall meta-analysis. Forest plots corresponding to each summary effect estimate were visually assessed. Sensitivity analysis was not conducted due to the small number of studies included in the meta-analysis. Publication bias was assessed by visual inspection of the funnel plots and application of Begg's and Egger's tests. Statistical analyses were performed using Stata V.13.0 (StataCorp LP, College Station, Texas, USA) software.

RESULTS

A total of 13 studies were included in the review. The study selection process is shown in figure 1.

Characteristics of included studies

The characteristics of the included studies are presented in online supplementary table S1.

Of the 13 studies, two studies^{21 22} assessed the effects of short-term air pollution exposure with the remaining studies focusing on long-term air pollution exposure. Six studies^{23–28} assessed maternal exposure to air pollution on a trimester basis.


Of the 11 epidemiological studies with assessment of long-term air pollution exposure, six studies^{23 25–29} employed a cohort design, of which one²⁶ was prospective. Two studies^{30 31} were semicological, that is, the exposure was assessed at group level and the outcome at individual level, and another two studies were cross-sectional^{32 33} with one study²⁴ applying a case control design. Of the two epidemiological studies with assessment of short-term air pollution exposure, one²¹ applied a case-crossover design with the hazard period defined as lag day 2 and reference periods selected by matching on the weekday within the same calendar month, and they also used two through 6 days before delivery as lag days. The other short-term air pollution exposure study²² was a time series analysis using Poisson regression and with lag effect of up to 5 days preceding delivery. Regarding the geographical location of the included studies, five studies were conducted in the USA, one in Latin America, three in East Asia, two in Europe, one in Russia and one in the UK. Seven of the included studies were published after the year 2005.

Eleven^{21–31} of the included studies relied on routine air monitoring data in their respective study areas in estimating maternal

Table 1 Search terms

Exposure MeSH Terms	Free text words	Outcome MeSH Terms	Free text words
Air pollution	Ambient air	Stillbirth	Birth outcome
Environmental pollution	pollution	Perinatal	Intrauterine
Vehicular emission	Outdoor air	mortality	mortality
Particulate matter	pollution	Fetal mortality	
PM	Ambient air	Fetal death	
Ozone	quality	Pregnancy	
O ₃	Traffic exposure	outcome	
Nitrogen dioxide			
NO ₂			
Sulfur dioxide			
SO ₂			
Carbon monoxide			
CO			
Polycyclic aromatic hydrocarbons			
PAH			

Figure 1 Flowchart of article selection process.



air pollution exposure. Landgren³³ categorised each exposure in two ways, (1) above and below the mean exposure value (SO₂: 8.0 µg/g, CH: 6.6 µg/g and NO: 14.7 µg/g) of all the included municipalities and (2) the municipality with the highest exposure level was compared with all other municipalities. Vassilev *et al*³² used statewide combined modelled average concentrations for each census tract and categorised exposures into low (0.040–0.268 mg/m³), medium (0.269–0.610 mg/m³) and high exposure (0.611–2.830 mg/m³) level with low exposure serving as the reference in the analysis. The included studies measured 13 pollutants including SO₂, NO₂, CO, particles (PM_{2.5}, PM₁₀, SPM), O₃, NO, POM, NO_x, hydrocarbon (CH), black smoke, which was taken as equivalent to PM₄ in the study that measured this pollutant,²⁵ and suspended solids.

Of the 11 studies relying on air monitoring station data, seven studies^{21 23–28} and that by Vassilev *et al*³² assigned exposures to mothers based on their residential addresses at the time of delivery. Regarding the studies conducted by Faiz *et al*^{21 23} and DeFranco *et al*,²⁷ mothers had to live within a 10 km radius of the closest monitoring station; and Green *et al*²⁸ used 20 km radius for PM_{2.5}, O₃, SO₂ and 5 km radius for CO and NO₂ to be included in the studies. Whereas Hwang *et al*²⁴ applied a 25 km radius. The exposure assignments in the studies by Kim *et al*,²⁶ Pearce *et al*²⁵ and Vassilev *et al*³² were not based on any fixed radius. The two ecological studies^{30 31} used annual mean concentrations of the pollutants studied to assign exposures.

Dimitriev²⁹ used monthly concentration of air pollutants in the study areas.

Six studies^{21 23 24 27 28 32} used a stillbirth cut-off point of >20 weeks of gestation, with three studies^{22 25 30} using a cut-off of >28 weeks of gestation. Landgren³³ and Sakai³¹ did not provide a case definition in their reports. DeFranco *et al*²⁷ ascertained stillbirth by using both, last menstrual period (LMP) and ultrasound examination methods, whereas two other studies^{25 26} mostly used the LMP method but also used fetal ultrasound examination if there was either uncertainty about the LMP date or discordance between the two estimates. Two studies^{23 28} used the LMP method only, with Hwang *et al*²⁴ applying ultrasound examination. The type of stillbirth studied was not mentioned in any of the included studies except the study by Pereira *et al*,²² which identified the stillbirths as intrauterine mortality, which we assume to be antepartum stillbirth. The source of data on stillbirth was obtained from fetal death certificates in five studies,^{21–23 32 33} birth registry or vital statistics in five studies^{24 27 28 30 31} and hospital records in two studies.^{25 26} Dimitriev²⁹ did not provide any information on how the stillbirths were ascertained.

Methodological quality of the included studies

Selection bias

Selection bias was very minimal in the included studies as most of the studies collected data from fetal death certificates or birth

registries and are likely to have represented their source populations with high response rate also reported. The prospective cohort study,²⁶ however, only included pregnant women who visited the hospital for prenatal care, excluding mothers with missing values for residential address and sociodemographic risk factors. Few studies^{25 28 32} excluded mothers without gestational age information and census tract coding (8%) from the analysis.

Information bias

There was evidence of potential information bias in all the included studies due to the reliance on proximity of maternal homes to the nearest air pollution monitoring station,^{21–31} and use of emission measurement and meteorological data^{32 33} in assessing exposure. Factors such as mother's mobility, change of residence during pregnancy, occupation of mother and air exchange were not considered, and this may lead to a decrease in the accuracy of the exposure assessment and introduce a non-differential misclassification that might lead to an underestimation of the effects of air pollution. Also, the exposure source was not reported by the studies other than those by Hwang *et al*,²⁴ Pearce *et al*,²⁵ Vassilev *et al*³² and Sakai.³¹ For the outcome measurement, the true dates of fetal death were unknown, hence an estimated time using date of delivery was recorded on the fetal death certificate and this may introduce bias. This was observed in the study by Faiz *et al*,^{21 23} Hwang *et al*,²⁴ Kim *et al*,²⁶ Green *et al*,²⁸ Bobak and Leon,³⁰ and Vassilev *et al*.³²

Control of confounding

Any determinant of the risk of stillbirth could be considered as a potential confounder in cohort, case-control and cross-sectional studies. The case-crossover study and time-series analyses eliminate any confounding related to individual characteristics and environmental exposures that are linked to the studied air pollutant. Most studies adjusted for the characteristics of the mother, such as age and family characteristics, at an individual level. Some of the studies applied control of confounding at group levels.³⁰ When estimating the effect of an individual air pollutant, exposure to other air pollutants is a potential confounder. Only five studies fitted more than one air pollutant into the multivariate model, that is, applied multipollutant rather than single pollutant models.^{21 22 24 28 30}

Regarding the ecological studies, Bobak and Leon³⁰ adjusted for several socioeconomic characteristics, which were obtained from the Czech Statistical office, at the district level; whereas Sakai³¹ did not control for confounding. In the time series analysis, Pereira *et al*²² adjusted for season and weather. Faiz *et al*²¹ conducted a case-crossover study where they controlled for mean temperature of the corresponding lag days. Of the nine long-term air pollution exposure studies, Dimitriev²⁹ did not control for any confounding and the confounding control was considered inadequate in one study,³³ as the author only adjusted for year of birth, maternal age and parity. Confounding control was considered adequate in the remaining studies.^{23–28 32} These studies adjusted for a range of confounders including maternal age, race, education, socioeconomic status, season or month of conception, parity, infant sex, prenatal care and lifestyle characteristics. Hwang *et al*²⁴ and Kim *et al*²⁶ further controlled for gestational age and maternal anthropometry, respectively. Vassilev *et al*³² used separate Mantel-Haenszel OR analysis for potential confounding factors and few factors were controlled in the final logistic regression

model, this approach can also be deemed as adequate control of confounding factors.

By applying the NOS scale, three studies^{23 24 28} were rated as very high quality (case-control/cohort—8 or more stars) and one study²⁶ was rated as high quality (cohort study—7 stars). The Newcastle-Ottawa-Scores for all the studies included in the meta-analyses are presented in the online supplementary table S2.

Findings of included studies and summary effect estimates

Summary effect estimates are presented in table 2 and the corresponding forest plot in figure 2.

Three studies^{23 24 28} provided estimates for the relation of SO₂ exposure (per 3 ppb, 1 ppb and 10 ppb increase in mean concentration) to stillbirth for the entire pregnancy period. The summary-effect estimate (EE) per 3 ppb increase in SO₂ exposure in the random effects model was 1.022 (95% CI 0.984 to 1.062), with low heterogeneity between the studies observed ($I^2=19.6\%$). The case-crossover study²¹ reported increased risk of stillbirth with IQR (4.7 ppb) increase in mean SO₂ exposure 2 days before delivery, and also found similar associations on all the lag days; whereas, another time series study²² reported marginal association between daily counts of intrauterine mortality and SO₂ concentration on the same day as delivery. One of the ecological studies³¹ also found significant positive correlation between spontaneous fetal death rate and SO₂ concentration. However, Landgren³³ and other ecological study (per 50 µg/m³ increase in annual mean concentration of SO₂),³⁰ however, did not find any association, even after adjustment with other pollutants SPM and NO_x.

Three studies^{23 24 28} provided estimates for the relation of both NO₂ and CO exposure for the entire pregnancy period, with the summary EE in the random effects model found to be 1.066 (95% CI 0.965 to 1.178) per 10 ppb increase in mean NO₂ concentration and 1.025 (95% CI 0.985 to 1.066) per 0.4 ppm increase in mean CO concentration, respectively. Evidence of high heterogeneity was observed in the NO₂ analysis ($I^2=79.6\%$). The case-crossover²¹ studies reported increased risk of stillbirth with IQR increase of NO₂ (16.4 ppb) and CO (0.54 ppm) in mean concentration, respectively, 2 days before delivery and also found similar associations on all the lag days. The time-series study²² reported strong significant dose-response relationship between daily counts of intrauterine mortality with NO₂ concentration, and marginal association with CO concentration at 5 days and 3 days before delivery, respectively. The findings of CO exposure with stillbirth were, however, less consistent. Sakai³¹ also found a statistically significant positive correlation between NO₂ concentration and spontaneous fetal death rate. Pereira *et al*²² also developed an overall index of air pollution (combination of NO₂, SO₂, CO) and found a very robust significant association with dose-response relationship, whereas Faiz *et al*²¹ used two pollutant models on lag day 2 but found estimates similar to those of the single pollutant models. Green *et al*²⁸ adjusted PM_{2.5} or ozone with NO₂ and the associations were unaffected.

Two studies^{24 26} provided estimates for the relation of stillbirth per 10 µg/m³ increase in average PM₁₀ concentration during the entire pregnancy duration to the summary EE in the random effects model found to be 1.014 (95% CI 0.948 to 1.085). Evidence of high heterogeneity was noted in the analysis ($I^2=85.0\%$). The time series study²² did not find any statistically significant associations.

Two studies^{23 28} provided estimates for the relation of stillbirth per 4 µg/m³ increase in average PM_{2.5} concentration during the entire pregnancy duration to the summary EE in the

Table 2 Summary effect estimates for the relation between ambient air pollution exposure and the risk of stillbirth

Air pollutant	Studies contributing to the summary effect estimate	Fixed effects Summary effect estimates EE (95% CI)	Random effects Summary effect estimates EE (95% CI)	Heterogeneity		
				χ^2	p Value	I^2 (%)
SO ₂ (3 ppb)	Faiz 2012, Green 2015, Hwang 2011	1.019 (0.989 to 1.049)	1.022 (0.984 to 1.062)	2.49	0.288	19.6
1st trimester	Faiz 2012, Green 2015, Hwang 2011,	0.997 (0.975 to 1.020)	1.040 (0.962 to 1.125)	10.34	0.006	80.7
2nd trimester	Faiz 2012, Green 2015, Hwang 2011	1.003 (0.977 to 1.030)	1.003 (0.977 to 1.030)	1.79	0.408	0.0
3rd trimester	Faiz 2012, Green 2015, Hwang 2011	0.996 (0.967 to 1.026)	1.042 (0.951 to 1.142)	11.26	0.004	82.2
NO ₂ (10 ppb)	Faiz 2012, Green 2015, Hwang 2011	1.049 (1.012 to 1.088)	1.066 (0.965 to 1.178)	9.78	0.008	79.6
1st trimester	Faiz 2012, Green 2015, Hwang 2011	1.025 (0.996 to 1.054)	1.035 (0.983 to 1.089)	4.43	0.109	54.8
2nd trimester	Faiz 2012, Green 2015, Hwang 2011	1.005 (0.977 to 1.034)	1.007 (0.948 to 1.071)	5.83	0.054	65.7
3rd trimester	Faiz 2012, Green 2015, Hwang 2011	1.015 (0.980 to 1.051)	1.015 (0.980 to 1.051)	1.88	0.391	0.0
CO (0.4 ppm)	Faiz 2012, Green 2015, Hwang 2011	1.022 (0.995 to 1.050)	1.025 (0.985 to 1.066)	2.52	0.284	20.5
1st trimester	Faiz 2012, Green 2015, Hwang 2011	1.002 (0.983 to 1.022)	1.011 (0.967 to 1.057)	2.92	0.232	31.6
2nd trimester	Faiz 2012, Green 2015, Hwang 2011	1.002 (0.979 to 1.025)	1.015 (0.948 to 1.087)	5.60	0.061	64.3
3rd trimester	Faiz 2012, Green 2015, Hwang 2011	1.014 (0.992 to 1.038)	1.052 (0.973 to 1.138)	10.19	0.006	80.4
PM ₁₀ (10 µg/m ³)	Hwang 2011, Kim 2007	1.012 (0.986 to 1.039)	1.014 (0.948 to 1.085)	6.67	0.010	85.0
1st trimester	Hwang 2011, Kim 2007	1.015 (0.991 to 1.039)	0.998 (0.936 to 1.064)	2.18	0.140	54.1
2nd trimester	Hwang 2011, Kim 2007	0.968 (0.944 to 0.993)	1.005 (0.905 to 1.116)	5.31	0.021	81.2
3rd trimester	Hwang 2011, Kim 2007	0.995 (0.968 to 1.022)	1.021 (0.919 to 1.134)	10.96	0.001	90.9
PM _{2.5} (4 µg/m ³)	Faiz 2012, Green 2015	1.021 (0.996 to 1.046)	1.021 (0.996 to 1.046)	0.18	0.669	0.0
1st trimester	Faiz 2012, Green 2015	1.002 (0.982 to 1.022)	1.042 (0.920 to 1.180)	2.35	0.126	57.4
2nd trimester	Faiz 2012, Green 2015	1.011 (0.996 to 1.026)	1.040 (0.940 to 1.152)	1.92	0.166	47.9
3rd trimester	Faiz 2012, Green 2015	1.00 (0.981 to 1.020)	1.00 (0.981 to 1.020)	0.23	0.631	0.0
O ₃ (10 ppb)	Green 2015, Hwang 2011	1.005 (0.982 to 1.029)	1.002 (0.971 to 1.034)	1.24	0.265	19.6
1st trimester	Green 2015, Hwang 2011	1.001 (0.983 to 1.020)	1.001 (0.983 to 1.020)	0.13	0.714	0.0
2nd trimester	Green 2015, Hwang 2011	1.004 (0.985 to 1.022)	0.991 (0.944 to 1.040)	3.18	0.074	68.6
3rd trimester	Green 2015, Hwang 2011	1.025 (1.006 to 1.043)	1.012 (0.966 to 1.060)	2.72	0.099	63.2

random effects model found to be 1.021 (95% CI 0.996 to 1.046). No heterogeneity was noted in the analysis ($I^2=0.0\%$). DeFranco *et al*²⁷ found non-significant 21% (OR: 1.21, 95% CI 0.96 to 1.53) increased risk in stillbirth with high PM_{2.5} (15.67 µg/m³) exposure during the entire pregnancy.

Two studies^{24, 28} provided estimates for the relation of stillbirth per 10 ppb increase in average O₃ concentration during the entire pregnancy duration to the summary EE in the random effects model found to be 1.002 (95% CI 0.971 to 1.034). Evidence of low heterogeneity was noted in the analysis ($I^2=19.6\%$). The time series study²² evaluated the relation between O₃ exposure and stillbirth, and found no association.

Four studies^{23, 24, 26, 28} provided trimester-specific estimates for the relation of stillbirth risk to six pollutants (SO₂, NO₂, CO, PM₁₀, PM_{2.5}, O₃). With the exception of SO₂ exposure during the second trimester, NO₂ and PM_{2.5} exposure during the third trimester and O₃ exposure during the first trimester, we observed evidence of moderate to substantial heterogeneity in most of the analysis. In the random effect model, almost all these pollutants (per 3 ppb SO₂, 10 ppb NO₂, 0.4 ppm CO, 10 µg/m³ PM₁₀, 4 µg/m³ PM_{2.5} and 10 ppb O₃ increase in mean concentration) showed increased risk in each trimester except SO₂ and NO₂ exposure in the second trimester, PM₁₀ and O₃ exposure in the first and second trimester and PM_{2.5} exposure in the third trimester. For SO₂, CO, PM₁₀ and O₃, third trimester exposure appears to pose the highest risk whereas for NO₂ and PM_{2.5}, first trimester exposure posed the highest risk. The summary EE per 10 µg/m³ increase in PM₁₀ exposure showed an increasing trend whereas for PM_{2.5} per 4 µg/m³ increase in concentration, a decreasing trend was

observed. Regarding SO₂, NO₂, CO and O₃, the trend was inconsistent with the second trimester summary estimates, being the lowest for SO₂, NO₂, O₃ and first trimester for the CO exposure. Faiz *et al*²¹ found a non-significant small increased risk of stillbirth with IQR increase in the mean PM_{2.5} concentration in their time series analysis on all the lag days. Whereas DeFranco *et al*²⁷ found significant 42% (OR 1.42, 95% CI 1.06 to 1.91) increased risk in stillbirth with high PM_{2.5} (16.22 µg/m³) exposure only in the third trimester. Hwang *et al*²⁴ evaluated increased risk of stillbirth in association with SO₂ per 1 ppb and PM₁₀ per 10 µg/m³ increase during the first and second months of pregnancy, and was stable after adjustment for O₃ and either CO or NO₂ in multipollutant models. Green *et al*²⁸ found robust association with third trimester O₃ exposure after addition of PM_{2.5} and NO₂.

Dimitriev²⁹ compared the occurrence of stillbirth in good and worse ecological areas assessed on the basis of monthly concentration of selected pollutants (suspended solids, SO₂, CO, NO₂), and reported risk of stillbirth as 6.63 and 11.03 per 1000 births, respectively. The estimated risk ratio showed a 65% (RR 1.650, 95% CI 1.136 to 2.397) increased risk of stillbirth among mothers resident in polluted ecological areas.

Two studies^{31, 33} that investigated the relation of NO exposure to stillbirth also found no association. The association between stillbirth risk and per 10 µg/m³ relating average weekly black smoke (equivalent to PM₄) exposure was evaluated by Pearce *et al*,²⁵ but did not find any significant association during pregnancy. Bobak and Leon³⁰ evaluated the effects of SPM and NO_x (per 50 µg/m³ increase in concentration), and found no significant association in the single pollutant model nor after




Figure 2 Forest plot showing the effect of ambient air pollutants on stillbirth. ES, effect size; weights are from random effects analysis.

adjustment with all pollutant (SPM, SO₂, NO_x) models. Vassilev *et al*³² investigated POM exposure and, using low exposure as the reference category, found statistically significant increased risk of stillbirth with medium and high exposure (ORs of 1.21 (95% CI 1.04 to 1.40) and 1.19 (95% CI 1.02 to 1.39), respectively). The effect of CH was investigated by Landgren,³³ but no association was found between stillbirth and levels of CH concentration in Swedish municipalities.

Evaluation of publication bias

Figure 3 presents the funnel plots for all the study specific effect estimates used to calculate the summary effect estimates. Online supplementary table S3 presents results from the Begg's and Egger's tests. There was no indication of publication bias present, although these results should be interpreted with caution because they were based on two or three study-specific effect estimates only.

DISCUSSION

Our systematic review and meta-analysis provided evidence that prenatal exposure to air pollution increases the risk of stillbirth. The summary effect estimates from the random effects models were systematically elevated, although they did not reach statistical significance. Per 10 ppb NO₂, 0.4 ppm CO, 3 ppb SO₂, 4 µg/m³ PM_{2.5} and 10 µg/m³ PM₁₀, increase in mean exposure during the entire pregnancy duration, respectively, resulted in a 6.6% (EE=1.066, 95% CI 0.965 to 1.178), 2.5% (EE=1.025, 95% CI 0.985 to 1.066), 2.2% (EE=1.022, 95% CI 0.984 to 1.062), 2.1% (EE=1.021, 95% CI 0.996 to 1.046) and 1.4% (EE=1.014, 95% CI 0.948 to 1.085) increased risk of stillbirth.

The point estimates for the third trimester were slightly elevated for SO₂, CO, PM₁₀ and O₃ consistently, with a hypothesis of a susceptible time window for the adverse effects, although the differences were not statistically significant. Two time series studies used lag day not more than 6 days preceding stillbirth and both found increased risk with certain pollutants.

Validity of results

We included all the studies identified in an extensive systematic search, so missing of important epidemiological studies is less likely to have happened. A significant number of studies have emerged since the last reviews; hence the critical assessment of the evidence is timely. Even though our review contains eight more studies and much more information than the previous reviews, we found a very limited number of estimates for each of the pollutants, and only five studies made attempts to adjust for other air pollutants when presenting effect estimates of each air pollutant. Therefore, we could not include all of the studies in the meta-analyses, and the reliability on the summary effect estimates is further compromised. However, the existing evidence is suggestive of causality for air pollution and stillbirth without precise identification of the timing of exposure. With the limited studies on the relevant topic, our review suggests strong priorities for future research. The visual inspection of the funnel plots and the statistical assessment did not indicate publication bias.

Biological plausibility

Fetuses are more affected by a variety of environmental toxicants because of differential exposure and physiological




Figure 3 Funnel plot with pseudo 95% confidence limits for the relation between ambient air pollutants and stillbirth.

immaturity.³⁴ The biological mechanisms by which exposure to ambient air pollutants leads to fetal death is not very clear. Faiz *et al*²¹ suggested the direct crossing of air pollutants across the placenta, causing irreversible damage to the dividing cells of the growing fetus and triggering hypoxic damage or immune-mediated injury during critical periods of development, as a possible mechanism.

Of all the pollutants, only the mechanism of the toxic effects of CO on the fetus is well established.¹⁷ CO reduces oxygen-carrying capacity of maternal haemoglobin that could seriously affect oxygen delivery to fetal circulation.³⁵ Sangalli *et al*³⁶ revealed that CO crosses the placental barrier and haemoglobin on fetal blood has greater affinity for binding CO than that in an adult; O₂ delivery to fetal tissues is further compromised.³⁷ Moreover, fetal elimination of carbon monoxide is slower than in the mother.³⁸ There is also a significant dose-dependent relationship between CO and COHb, and a developing fetus can be deprived of adequate oxygenation due to high levels of COHb, which may even lead to fetal death.²²

Maternal exposures to particulate (PM) air pollutants during pregnancy can result in increased concentration of DNA adducts or may lower the efficiency of the transplacental function, resulting in decreased fetal health leading to stillbirth.^{39 40}

There is evidence that the presence of air pollution increases blood viscosity and plasma fibrinogen relates to coagulation; these haematological factors might have an influence on blood perfusion of the placenta, which could also lead to impair fetal health.^{41–43}

Synthesis with previous knowledge

In the meta-analysis, we observed a 1.4% (EE 1.014, 95% CI 0.948 to 1.085) increased risk of stillbirth with 10 µg/m³ increase in PM₁₀ exposure during the entire duration of pregnancy, albeit statistically but not significantly. Pereira *et al*²² also did not find any statistically significant association in their time series study. Glinianaia *et al*¹⁷ reported little evidence of association between exposure to PM and stillbirth risk, and thus describes it as insufficient to assess a possible association between PM and stillbirth.

For PM_{2.5}, Zhu *et al* indicated that there was no evidence of statistically significant effect (OR: 1.18, 95% CI 0.69 to 2.04) on stillbirth with an increase of 10 µg/m³; this result is consistent with our review, which reveals (EE 1.021, 95% CI 0.996 to 1.046) with an increase of 4 µg/m³. However, in our meta-analysis, we also found a 4.2% (EE 1.042, 95% CI 0.920 to 1.180) and 4.0% (EE 1.040, 95% CI 0.940 to 1.152) increase per 4 µg/m³, in the first and second trimester, respectively.

We also observed a small and statistically non-significant increased risk of stillbirth with SO₂ 2.2% (EE 1.022, 95% CI 0.984 to 1.062) and CO 2.5% (EE 1.025, 95% CI 0.985 to 1.066) exposure for the entire pregnancy period in the meta-analysis. The short-term air pollution studies on these relationships²¹ and one of the ecological studies³¹ also reported an increased risk of stillbirth related to SO₂ and CO exposure. The findings of CO exposure with stillbirth was, however, less consistent.

Lacasana *et al*¹⁸ stated that SO₂ and CO also showed some, though less consistent, relationship with intrauterine mortality. In our review, we found that there is an increased risk of stillbirth associated with increased ambient concentrations of sulfur dioxide in early pregnancy, with pooled effect estimates of 4.0% (EE 1.040, 95% CI 0.962 to 1.125). Also, we found the highest

pooled estimate of 5.2% (EE 1.052, 95% CI 0.973 to 1.138) for carbon monoxide in the last trimester.

For NO₂, we observed a higher increase risk of stillbirth of 6.6% (EE 1.066, 95% CI 0.965 to 1.178) for the entire pregnancy. This finding is consistent with the study by Lacasana *et al*,¹⁸ which reported an increased risk of stillbirth with exposure to NO₂.

Most of the pollutants in polluted ambient air, such as PM and CO, are also present in indoor air polluted by smoking and solid fuel combustion, although the concentrations are lower. Leonardi-Bee *et al*¹⁵ conducted a meta-analysis of studies on the effects of secondhand smoking on the risk of stillbirth, presenting a 23% risk increase (EE 1.23, 95% CI 1.09 to 1.38). Amegah *et al*¹⁶ synthesised the effect of solid fuels showing an effect estimate of 29% for stillbirth. These findings are consistent with the present pollutant-specific summary effect estimates and thus strengthen the hypothesis that ambient air pollution increases the risk of stillbirth.

Conclusion, recommendations and implications for future research

Our results provide suggestive evidence that ambient air pollution is a risk factor for stillbirth. Pregnant women should be aware of the potential adverse effects of ambient air pollution, although the prevention against exposure to air pollutants generally requires more action by the government than by the individual. The healthcare sector can create awareness and engage other sectors contributing to ambient air pollution (such as the housing sector, transportation sector, industries and the energy sector), to develop and implement policies such as control of vehicular emissions, fuel quality improvement and control of industrial waste emission, to reduce the risk of air pollutants.

Most of the studies reviewed used data from monitoring stations to assess maternal exposure levels. Future studies should integrate the use of personal monitoring methods and also consider the activity of mothers, change in residence, air exchange, mother's occupation and outdoor activities of the mothers. The pregnant women should also be monitored if possible from the first month of pregnancy in order to ascertain the exact period of the effect.

Acknowledgements The authors thank Svetlana Filatova, Medical Research Center Oulu, Oulu University Hospital and University of Oulu, Finland, for helping with the translation of the Russian study included in the review.

Contributors JJKJ conceived the study and designed it with AKA. NS and HAB conducted the search and screened the articles, and contributed to the statistical analyses. AKA contributed to the evaluation of the manuscript, writing of the manuscript, statistical analyses and interpretation of the results. All the authors revised the manuscript for important intellectual content, and the final version has been approved by all the authors. JJKJ takes responsibility for the integrity of the article.

Funding The University of Oulu strategic funds.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data needed for repeating the meta-analyses are presented in tables. Additional information on the methods or preliminary data can be requested from the corresponding author.

REFERENCES

- 1 World Health Organization (WHO). Ambient (outdoor) air quality and health. <http://www.who.int/mediacentre/factsheets/fs313/en/>. Fact sheet N°313. Updated March, 2014.
- 2 Demetriou CA, Raaschou-Nielsen O, Loft S, *et al*. Biomarkers of ambient air pollution and lung cancer: a systematic review. *Occup Environ Med* 2012;69:619–27.

- 3 Darrow LA, Klein M, Flanders WD, *et al.* Air pollution and acute respiratory infections among children 0–4 years of age: an 18-year time-series study. *Am J Epidemiol* 2014;180:968–77.
- 4 Lee BJ, Kim B, Lee K. Air pollution exposure and cardiovascular disease. *Toxicol Res* 2014;30:71–5. Review.
- 5 Bonzini M, Carugno M, Grillo P, *et al.* Impact of ambient air pollution on birth outcomes: systematic review of the current evidences. *Med Lav* 2010;101:341–63.
- 6 Schikowski T, Adam M, Marcon A, *et al.* Association of ambient air pollution with the prevalence and incidence of COPD. *Eur Respir J* 2014;44:614–26.
- 7 Gowers AM, Cullinan P, Ayres JG, *et al.* Does outdoor air pollution induce new cases of asthma? Biological plausibility and evidence; a review evidence; a review. *Respirology* 2012;17:887–98.
- 8 World Health Organization (WHO). *Burden of disease from ambient air pollution for 2012*. Geneva: WHO, 2014. http://www.who.int/phe/health_topics/outdoorair/databases/FINAL_HAP_AAP_BoD_24March2014.pdf. (accessed 14 Nov 2014).
- 9 Sapkota A, Chelikowsky AP, Nachman KE, *et al.* Exposure to particulate matter and adverse birth outcomes: a comprehensive review and meta-analysis. *Air Qual Atmos Health* 2012;5:369–81.
- 10 Maisonet M, Correa A, Misra D, *et al.* A review of the literature on the effects of ambient air pollution on fetal growth. *Environ Res* 2004;95:106–15.
- 11 Vrijheid M, Martinez D, Manzanera S, *et al.* Ambient air pollution and risk of congenital anomalies: a systematic review and meta-analysis. *Environ Health Perspect* 2011;119:598–606.
- 12 Stillerman KP, Mattison DR, Giudice LC, *et al.* Environmental exposures and adverse pregnancy outcomes: a review of the science. *Reprod Sci* 2008;15:631–50.
- 13 Cousens S, Blencowe H, Stanton C, *et al.* National, regional, and worldwide estimates of stillbirth rates in 2009 with trends since 1995: a systematic analysis. *Lancet* 2011;377:1319–30.
- 14 Aminu M, Unkels R, Mdegela M, *et al.* Causes of and factors associated with stillbirth in low- and middle-income countries: a systematic literature review. *BJOG* 2014;121(Suppl 4):141–53.
- 15 Leonardi-Bee J, Britton J, Venn A. Secondhand smoke and adverse fetal outcomes in nonsmoking pregnant women: a meta-analysis. *Pediatrics* 2011;127:734–41.
- 16 Amegah AK, Quansah R, Jaakkola JJ. Household air pollution from solid fuel use and risk of adverse pregnancy outcomes: a systematic review and meta-analysis of the empirical evidence. *PLoS ONE* 2014;9:e113920.
- 17 Glinianaia SV, Rankin J, Bell R, *et al.* Particulate air pollution and fetal health: a systematic review of the epidemiologic evidence. *Epidemiology* 2004;15:36–45.
- 18 Lacasaña M, Esplugues A, Ballester F. Exposure to ambient air pollution and prenatal and early childhood health effects. *Eur J Epidemiol* 2005;20:183–99.
- 19 Zhu X, Liu Y, Chen Y, *et al.* Maternal exposure of fine particulate matter (PM_{2.5}) and pregnancy outcomes: a meta-analysis. *Environ Sci Pollut Res* 2015;22:3383–96.
- 20 Wells G, Shea B, O'Connell D, *et al.* The Newcastle–Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analyses. http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm. (accessed 22 Feb 2013)
- 21 Faiz AS, Rhoads GG, Demissie K, *et al.* Does ambient air pollution trigger stillbirth? *Epidemiology* 2013;24:538–44.
- 22 Pereira LA, Loomis D, Conceição GM, *et al.* Association between air pollution and intrauterine mortality in Sao Paulo, Brazil. *Environ Health Perspect* 1998;106:325–9.
- 23 Faiz AS, Rhoads GG, Demissie K, *et al.* Ambient air pollution and the risk of stillbirth. *Am J Epidemiol* 2012;176:308–16.
- 24 Hwang BF, Lee YL, Jaakkola JJ. Air pollution and stillbirth: a population-based case-control study in Taiwan. *Environ Health Perspect* 2011;119:1345–9.
- 25 Pearce MS, Glinianaia SV, Rankin J, *et al.* No association between ambient particulate matter exposure during pregnancy and stillbirth risk in the north of England, 1962–1992. *Environ Res* 2010;110:118–22.
- 26 Kim OJ, Ha EH, Kim BM, *et al.* PM10 and pregnancy outcomes: a hospital-based cohort study of pregnant women in Seoul. *J Occup Environ Med* 2007;49:1394–402.
- 27 DeFranco E, Hall E, Hossain M, *et al.* Air pollution and stillbirth risk: exposure to airborne particulate matter during pregnancy is associated with fetal death. *PLoS ONE* 2015;10:e0120594.
- 28 Green R, Sarovar V, Malig B, *et al.* Association of stillbirth with ambient air pollution in a California cohort study. *Am J Epidemiol* 2015;181:874–82.
- 29 Dimitriou DA. Effects of air pollution on stillbirth in an industrial town. *Gig Sanit* 2000;5:7–9.
- 30 Bobak M, Leon DA. Pregnancy outcomes and outdoor air pollution: an ecological study in districts of the Czech Republic 1986–8. *Occup Environ Med* 1999;56:539–43.
- 31 Sakai R. Fetal abnormality in a Japanese industrial zone. *Int J Environ Stud* 2007;23:113–20.
- 32 Vassilev ZP, LRobson MG, Klotz JB. Outdoor exposure to airborne polycyclic organic matter and adverse reproductive outcomes: a pilot study. *Am J Ind Med* 2001;40:255–62.
- 33 Landgren O. Environmental pollution and delivery outcome in southern Sweden: a study with central registries. *Acta Paediatr* 1996;85:1361–4.
- 34 Perera FP, Jedrychowski W, Rauh V, *et al.* Molecular epidemiologic research on the effects of environmental pollutants on the fetus. *Environ Health Perspect* 1999;107:451–60.
- 35 Salam MT, Millstein J, Li YF, *et al.* Birth outcomes and prenatal exposure to ozone, carbon monoxide, and particulate matter: results from the Children's health study. *Environ Health Perspect* 2005;113:1638–44.
- 36 Sangalli MR, Mclean AJ, Peek MJ, *et al.* Carbon monoxide disposition and permeability-surface area product in the foetal circulation of the perfused term human placenta. *Placenta* 2003;24:8–11.
- 37 Di Cera E, Doyle ML, Morgan MS, *et al.* Carbon monoxide and oxygen binding to human hemoglobin F0. *Biochemistry* 1989;28:2631–8.
- 38 Hill EP, Hill JR, Power GG, *et al.* Carbon monoxide exchanges between the human fetus and mother: a mathematical model. *Am J Physiol* 1977;232:H311–23.
- 39 Perera FP, Hemminki K, Gryzbowska E, *et al.* Molecular and genetic damage in humans from environmental pollution in Poland. *Nature* 1992;360:256–8.
- 40 Zondervan HA, Oosting J, Hardeman MR, *et al.* The influence of maternal whole blood viscosity on fetal growth. *Eur J Obstet Gynecol Reprod Biol* 1987;25:187–94.
- 41 Ha EH, Kwon HJ. Issues in air pollution epidemiologic studies. *Korean J Pre Med* 2001;34:109–18.
- 42 Peters A, Döring A, Wichmann HE, *et al.* Increased plasma viscosity during an air pollution episode: a link to mortality? *Lancet* 1997;349:1582–7.
- 43 Knottnerus JA, Delgado LR, Knipschild PG, *et al.* Haematological parameters and pregnancy outcome. A prospective cohort study in the third trimester. *J Clin Epidemiol* 1990;43:461–6.