
Journal of Clinical Pathology



Keratocystoma is a cystic squamous parotid gland neoplasm (A) which forms small irregular nests which may mimic invasion (B). Cellular atypia is absent. Mucoepidermoid carcinoma (C.D). Rare forms of mucoepidermoid carcinoma may be ciliated (C), and exhibit acinar differentiation (D).

Contents

Review

289 V

Fusions in salivary gland neoplasms: a review of practical diagnostic applications

J A Bishop

Editorial

Cholangiocarcinoma classification: current approach, relevance and challenges B Goeppert, Y Zen, J Valle, D Klimstra, V Deshpande

Original research

von Meyenburg complexes are more frequently associated with cholangiocarcinoma

D Jain, B Khandakar, P Ni, B Kenney, L Qin, V Deshpande, M I Fiel

307 6

Combined analysis of albumin in situ hybridisation and C reactive protein immunohistochemistry for the diagnosis of intrahepatic cholangiocarcinoma: towards a molecular classification paradigm

T Albrecht, A Rossberg, F Rose, K Breuhahn, E-M Baumann, M Tóth, F Brinkmann, A Charbel, M N Vogel, B Köhler, A Mehrabi, M W Büchler, S Singer, W Solass, B Straub, P Schirmacher, S Roessler, B Goeppert

Viewpoint

320

6

317 Leadership and emotional intelligence: an early-career pathologist's perspective on the laboratory medical director role

Novel scoring system provides high separation of

diagnosis: an adaption of HER2 D-DISH for ploidy

P Corcoran, J Coulter, K O'Donoghue, B Fitzgerald

diploidy and triploidy to aid partial hydatidiform mole

CM Joyce, S Dineen, J Deane, N Conlon, P M O'Shea,

M Asif

Original research



Articles carrying the Unlocked Logo are freely





www.publicationethics.org

327 6

Best practice

Short report

Original research



Morphology combined with HER2 D-DISH ploidy analysis to diagnose partial hydatidiform mole: an evaluation audit using molecular genotyping C M Joyce, G J Maher, S Dineen, N Suraweera, T V McCarthy, J Coulter, K O'Donoghue, M J Seckl, B Fitzgerald

practice

Fifth edition WHO classification: myeloid neoplasms X Chen, N Patkar, P Tembhare, S Papagudi, C Yeung, R Kanagal Shamanna, S Gujral, B Wood, K N Naresh

treport

Prevalence and breakdown of KRAS driver mutations in a large UK non-small cell lung cancer cohort

I C C Niesner, K J Balbi, B Poskitt, C Gemma, J Linares, D Allen, O Shutkever, C R Lindsay, P Bennett, D A Moore

mal research

Single-molecule localisation microscopy (SMLM) is feasible in human and animal formalin fixed paraffin embedded (FFPE) tissues in medical renal disease S F Brockmoeller, H Slaney, A Curd, A Bono, J H Felce, embedded (FFPE) tissues in medical renal disease S F Brockmoeller, H Slaney, A Curd, A Bono, J H Felce, D Arora, A Lewington, A G Miklosi, P Quirke

PostScript


357 **(6)**

Sunitinib induced glomerular thrombotic microangiopathy in a patient with refractory pancreatic neuroendocrine tumour

A Khan, M Consing Gangelhoff, S Moubarak, S Herrmann, K Nooruddin, M Alexander

359 Diagnostic utility of ERG immunostaining in dermatofibroma: be aware of ERG expression in cellular neurothekeoma

F Cordier, L Ferdinande, J Van Dorpe, D Creytens





OPEN ACCESS available online under the BMJ Journals See http://authors.bmi.com/open-access/



This journal is a member of and subscribes to the principles of the Committee on Publication



The online version of this article contains multiple choice questions hosted on BMJ

