

Lung function and onset of cardiometabolic diseases in the longitudinal Burden of Obstructive Lung Disease study

Christer Janson ,¹ James Potts,² Andrei Malinovski ,³ Dhiraj Agarwal,⁴ Rana Ahmed ,⁵ Althea Aquart-Stewart,⁶ Imed Harrabi,⁷ Meriam Denguezli,⁸ Graham Devereux ,⁹ Gregory E Erhabor,¹⁰ Thorarinn Gislason,¹¹ Rain Jogi,¹² Sanjay K Juvekar,¹³ Ben Knox-Brown ,² Parvaiz Koul,¹⁴ Kevin Mortimer,^{15,16} Asaad Ahmed Nafees,¹⁷ Rune Nielsen,¹⁸ Padukudru Anand Mahesh,¹⁹ Stefanni Nonna M Paraguas,²⁰ Anders Ørskov Rotevatn,²¹ Talant Sooronbaev,^{22,23} Peter G J Burney ,² Andre F S Amaral ,² BOLD Collaborative Research Group

To cite: Janson C, Potts J, Malinovski A, *et al.* Lung function and onset of cardiometabolic diseases in the longitudinal Burden of Obstructive Lung Disease study. *BMJ Open Respir Res* 2025;**12**:e002442. doi:10.1136/bmjresp-2024-002442

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjresp-2024-002442>).

Received 21 March 2024
Accepted 5 December 2024



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Christer Janson;
christer.janson@medsci.uu.se

ABSTRACT

Introduction Previous population-based studies, mainly from high-income countries, have shown that a higher forced vital capacity (FVC) is associated with a lower risk of developing cardiometabolic diseases. The aim of this study was to assess the longitudinal association between spirometry measures and the onset of cardiometabolic diseases across sites in low-income, middle-income and high-income countries.

Methods The study population comprised 5916 individuals from 15 countries participating in the Burden of Obstructive Lung Disease baseline and follow-up assessments. Postbronchodilator forced expiratory volume in 1 s (FEV₁), FVC and FEV₁/FVC were measured at baseline. Participants who reported having doctor-diagnosed hypertension, diabetes, heart disease and stroke at follow-up but not at baseline were considered new cases of these diseases. The association between lung function and the onset of participant-reported cardiometabolic diseases was assessed in each site using regression models, and estimates were combined using random effects meta-analysis. Models were adjusted for sex, age, smoking, body mass index and educational level.

Results Participants with greater per cent predicted FVC were less likely to have new-onset diabetes (OR per 10%=0.91, 95% CI 0.84 to 0.99), heart disease (OR per 10%=0.86, 95% CI 0.80 to 0.92) and stroke (OR per 10%=0.81, 95% CI 0.73 to 0.89) during the follow-up period (mean±SD 9.5±3.6 years). A greater percentage of FEV₁ was associated with a lower risk of onset of heart disease and stroke. No significant association was found between FEV₁/FVC and onset of reported cardiometabolic diseases, except for a higher risk of diabetes (OR per 10%=1.21, 95% CI 1.08 to 1.35) in participants with higher FEV₁/FVC.

Conclusions The findings of this study suggest that a low FVC is more important than a low FEV₁/FVC as a risk factor for developing cardiometabolic diseases. The value of including FVC in risk score models to improve their precision in predicting the onset of cardiometabolic diseases should be explored.

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Many patients with COPD have cardiovascular disease and diabetes. Previous studies, however, suggest that cardiometabolic diseases are more closely related to low lung function than airflow obstruction. Most of these studies have been cross-sectional and are from high-income countries.

WHAT THIS STUDY ADDS

⇒ This longitudinal study with almost 6000 randomly selected participants from 15 countries with different income levels showed that persons with a better lung function expressed as forced vital capacity (FVC) were less likely to develop diabetes, heart disease and stroke over 10 years.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ The potential value of including FVC in risk score models to improve their precision in predicting the onset of cardiometabolic diseases should be explored.

INTRODUCTION

Many patients with chronic obstructive pulmonary disease (COPD) have cardiovascular disease and diabetes.^{1 2} COPD is spirometrically characterised by chronic airflow obstruction. In a cross-sectional analysis of the populational-based Burden of Obstructive Lung Disease (BOLD) study, Triest *et al* found no significant association of postbronchodilator airflow obstruction with hypertension, diabetes and cardiovascular disease after adjusting for confounders.³ This suggests that the coexistence of chronic airflow obstruction

and cardiometabolic diseases is due to shared risk factors, such as age and smoking.

Another cross-sectional BOLD analysis found an association between a low forced vital capacity (FVC) and hypertension, diabetes and cardiovascular disease.⁴ In other analyses from European subsamples of BOLD, hypertension was found to be associated with lower FVC and forced expiratory volume in 1 s (FEV₁),⁵ and a greater pulse wave velocity was associated with both low total lung capacity and low FVC.⁶ These findings suggest that a low lung function is more important than airflow obstruction in developing cardiometabolic diseases. This is also supported by other studies showing that having a low FVC and FEV₁ is associated with a higher risk of hypertension,⁷ diabetes,^{8–10} myocardial infarction and cardiovascular death.^{11 12} The biological explanation for the association is unclear, but several studies have shown an association between low FVC and FEV₁ and systemic inflammatory markers such as fibrinogen,¹¹ C reactive protein (CRP)^{13 14} and interleukin-6 (IL-6).^{15 16}

The aim of this investigation was to study associations between lung function and the onset of participant-reported cardiometabolic diseases in a longitudinal setting with participants from low-income, middle-income and high-income countries. We hypothesised that the onset of hypertension, diabetes, heart disease and stroke is more closely associated with low lung function than airflow obstruction.

METHODS

Study population

The baseline BOLD study was conducted between 2003 and 2016 and included 41 sites across Africa, Asia, Europe, North America, the Caribbean and Oceania and collected high-quality prebronchodilator and postbronchodilator spirometry from 28 828 randomly selected participants of the age of 40 years and higher. Between 2019 and 2021, a BOLD follow-up study was conducted in 18 sites across Africa, Asia, Europe and the Caribbean. At baseline, there were 12 502 participants with high-quality spirometry at these sites. A total of 6452 participants were followed up, with 5936 completing the study core questionnaire. The current analyses included 5916 (47.3%) participants with data on cardiometabolic disease at baseline and follow-up (figure 1).¹⁷

Lung function

FVC and FEV₁ were measured at baseline before and after bronchodilation (200 µg salbutamol). The spirometry was conducted with the participant sitting upright, wearing a disposable mouthpiece and a nose clip. FEV₁ and FVC values were obtained by spirometry using the ndd EasyOne (ndd Medizintechnik, Zurich, Switzerland). The postbronchodilator values of FVC, FEV₁ and FEV₁/FVC were expressed as per cent of the predicted using the reference values from the third US National Health and Nutrition Examination Survey (NHANES III)




Figure 1 Study population of the Burden of Obstructive Lung Disease (BOLD) follow-up study.

for adult European American men and women.¹⁸ Airflow obstruction was defined according to the lower limit of normal (LLN, <5th percentile) FEV₁/FVC.³ Restrictive lung function was defined as having a postbronchodilator FVC below LLN.⁴

All spirometry curves were checked centrally at the BOLD Operations Centre, and usable tests had to include at least three acceptable curves, with the two best blows being within 200 mL of each other. Before the start of the surveys, study site staff underwent intensive training covering consenting, questionnaire data collection, spirometry testing and quality control, anthropometry measurements and data transfer.¹⁷

Onset of cardiometabolic diseases

Participants who reported that they had been told by a doctor or healthcare provider they had hypertension, diabetes, heart disease and stroke at follow-up but not at baseline were defined as having an onset of these diseases.

The sites were classified into the level of income using the 2022–2023 World Bank Country Lending Group guidelines <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>.

Potential confounders

Height and weight were measured, and body mass index (BMI) was calculated as weight in kilograms divided by the square of height expressed in metres. Information on smoking history (current, ex-smoker and never-smoker), educational level and cardiometabolic diseases was obtained through a structured interview.

Statistics

Data were analysed using Stata V.17 (Stata). Characteristics and onset of hypertension, diabetes, heart disease and stroke are presented by site and described as n (%) or mean (SD). χ^2 test and unpaired t-test were used to compare responders and non-responders.

The association between postbronchodilator FVC, FEV₁ and FEV₁/FVC expressed as per cent of predicted at baseline with the onset of hypertension, diabetes, heart disease and stroke were assessed by logistic regression in each site and then combined by random effects meta-analyses. The models were adjusted for age, sex, smoking history, BMI and educational level. All regression models were adjusted for sampling weights within each site. Study sites that reported a low number of people with specific comorbidity were excluded from the meta-analysis because these sites could not be fitted into the model. Differences were considered to be significant if the p value was less than 0.05. Heterogeneity across sites was estimated using the I² statistic. I² values of 0%, 25%, 50% and 75% were considered no, low, moderate and high heterogeneity, respectively.

Sex-stratified analyses were performed because of previous findings of sex differences in the association between lung function and cardiometabolic diseases.³ Sensitivity analyses were done by excluding participants with a BMI≥30,¹⁹ those with any cardiometabolic disease and those with an FVC<LLN, respectively. We also used a multilevel (mixed effects) logistic regression model in our analyses. This allowed us to include participants from sites where the number of participants with comorbidities was too low to include in some of the meta-analyses.

Patient and public involvement

Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

RESULTS

The characteristics of the study population are presented in [table 1](#). The mean age of the population at baseline was 54 years, and 55% of the participants were women. There was a large variation in the prevalence of current smokers and level of education across the various study sites. Participants who took part in both surveys were comparatively younger, more likely to be women, had a slightly lower BMI, were more often highly educated and had a lower prevalence of reported cardiometabolic diseases than those who only participated in the first survey. The participants also had slightly higher mean lung function values than those not participating in the follow-up study (online supplemental table S1). The mean follow-up time was 9.5 years ([table 1](#)). During this period, 1144 had an onset of hypertension, 424 had diabetes, 607 had heart disease and 154 had an onset of stroke ([table 2](#)). Of those included, 453 (7.7%) had airflow obstruction, and 2303 (38.8%) had restrictive lung function.

Table 1 Characteristics of the population of the BOLD follow-up study at baseline and length of follow-up (n% and mean±SD)

| Country | Site | n | Age (years) | Women | BMI (kg/m ²) | Current smokers | Follow-up (years) | Country level of income |
|-------------|--------------------|------|-------------|-----------|--------------------------|-----------------|-------------------|-------------------------|
| Malawi | Chikwawa | 371 | 53±10 | 188 (51) | 21.8±3.8 | 63 (18) | 6.2±4.2 | Low |
| Sudan | Khartoum | 56 | 52±9 | 24 (45) | 27.1±5.5 | 12 (21) | 8.1±2.5 | Low |
| Benin | Seme-Kpodji | 117 | 51±9 | 62 (53) | 27.8±5.7 | 1 (1) | 7.4±3.6 | Low middle |
| Nigeria | Ife | 461 | 55±12 | 322 (70) | 25.8±5.7 | 9 (2) | 8.3±3.0 | Low middle |
| Tunisia | Sousse | 269 | 52±8 | 143 (53) | 29.7±5.2 | 62 (23) | 1+0.4±1.2 | Low middle |
| Morocco | Fes | 64 | 53±9 | 28 (44) | 27.7±4.9 | 7 (11) | 10.9±1.9 | Low middle |
| Philippines | Nampicuan-Talugtug | 472 | 52±9 | 259 (55) | 21.8±4.0 | 157 (33) | 10.8±1.4 | Low middle |
| Pakistan | Karachi | 259 | 51±9 | 151 (58) | 26.7±5.5 | 28 (11) | 5.7±3.8 | Low middle |
| India | Mysore | 537 | 46±7 | 313 (58) | 24.7±3.7 | 40 (7) | 7.4±1.5 | Low middle |
| India | Pune | 694 | 51±9 | 303 (44) | 22.1±3.8 | 51 (7) | 10.9±1.5 | Low middle |
| India | Kashmir | 89 | 53±11 | 39 (43) | 22.2±3.7 | 10 (11) | 12.5±5.0 | Low middle |
| Kyrgyzstan | Chui | 469 | 52±9 | 323 (69) | 28.7±5.2 | 88 (19) | 6.2±1.8 | Low middle |
| Kyrgyzstan | Naryn | 622 | 53±10 | 392(63) | 27.2±5.0 | 88 (14) | 6.3±1.9 | Low middle |
| Jamaica | | 95 | 56±11 | 53 (56) | 28.8±6.9 | 8 (8) | 5.9±1.7 | High middle |
| Estonia | Tartu | 391 | 58±11 | 199 (51) | 28.2±4.9 | 74 (19) | 10.8±1.2 | High |
| Iceland | Reykjavik | 378 | 52±8 | 171 (45) | 27.9±4.6 | 65 (17) | 15.2±1.0 | High |
| Sweden | Uppsala | 274 | 55±9 | 125 (46) | 26.7±3.9 | 36 (13) | 12.9±0.7 | High |
| Norway | Bergen | 298 | 54±9 | 138 (46) | 26.3±4.0 | 82 (28) | 15.1±0.8 | High |
| All | | 5916 | 54±11 | 3233 (55) | 26.0±5.5 | 881 (15) | 9.5±3.6 | |

BMI, body mass index; BOLD, Burden of Obstructive Lung Disease.

Table 2 The onset of cardiometabolic disease at different sites among participants without the disease at baseline (n (%))

| Country | Site | Hypertension (n=4820) | Diabetes (n=5624) | Heart disease (n=5480) | Stroke (n=5858) |
|-------------|--------------------|--------------------------|----------------------|---------------------------|--------------------|
| Malawi | Chikwawa | 20 (6) | 5 (1) | 11 (3) | 2 (1) |
| Sudan | Khartoum | 11 (22) | 16 (29) | 4 (7) | 0 |
| Benin | Seme-Kpodji | 16 (19) | 2 (2) | 3 (3) | 2 (2) |
| Nigeria | Ife | 64 (14) | 11 (2) | 5 (1) | 4 (1) |
| Tunisia | Sousse | 41 (19) | 43 (17) | 13 (5) | 9 (3) |
| Morocco | Fes | 6 (15) | 6 (11) | 8 (14) | 3 (5) |
| Philippines | Nampicuan-Talugtug | 124 (33) | 33 (7) | 42 (10) | 21 (5) |
| Pakistan | Karachi | 57 (30) | 43 (19) | 13 (5) | 8 (3) |
| India | Mysore | 28 (6) | 9 (2) | 7 (1) | 0 |
| India | Pune | 77 (12) | 68 (10) | 14 (2) | 13 (2)) |
| India | Kashmir | 25 (40) | 4 (5) | 3 (3) | 0 |
| Kyrgyzstan | Chui | 118 (36) | 19 (4) | 104 (27) | 18 (4) |
| Kyrgyzstan | Naryn | 215 (41) | 43 (7) | 168 (30) | 19 (3) |
| Jamaica | Jamaica | 27 (44) | 27 (32) | 9 (10) | 7 (7) |
| Estonia | Tartu | 72 (30) | 24 (7) | 53 (21) | 20 (5) |
| Iceland | Reykjavik | 99 (34) | 32 (9) | 72 (20) | 8 (2) |
| Sweden | Uppsala | 75 (35) | 25 (9) | 37 (14) | 4 (1) |
| Norway | Bergen | 69 (30) | 14 (5) | 41 (15) | 16 (5) |
| All | | 1144 (24) | 424 (7.5) | 607 (11) | 154 (3) |

Unadjusted analyses

High postbronchodilator FVC and FEV₁ expressed as per cent of predicted at baseline were associated with a lower risk of onset of participant-reported hypertension, diabetes, heart disease and stroke. Higher FEV₁/FVC expressed as per cent of predicted was associated with a higher risk of onset of diabetes (table 3). A moderate site

heterogeneity was found for the associations with higher lung function, diabetes and stroke.

Adjusted analyses

There was a statistically significant association between higher FVC and FEV₁ and a lower risk of onset of heart

Table 3 Association between lung function measures as per cent of predicted and cardiometabolic diseases

| | FVC | I ² | FEV ₁ | I ² | FEV ₁ /FVC | I ² |
|---------------|-------------------------|----------------|-------------------------|----------------|-------------------------|----------------|
| Unadjusted | | | | | | |
| Hypertension | 0.93 (0.88–0.98) | 0% | 0.94 (0.90–0.99) | 16% | 0.95 (0.89–1.02) | 24% |
| Diabetes | 0.85 (0.79–0.92) | 56% | 0.92 (0.87–0.99) | 49% | 1.25 (1.12–1.40) | 39% |
| Heart disease | 0.86 (0.80–0.92) | 0% | 0.88 (0.83–0.93) | 0% | 0.94 (0.88–1.02) | 13% |
| Stroke | 0.80 (0.73–0.88) | 55% | 0.82 (0.76–0.89) | 66% | 0.98 (0.89–1.11) | 54% |
| Adjusted | | | | | | |
| Hypertension | 0.97 (0.92–1.03) | 0% | 0.97 (0.92–1.02) | 15% | 0.94 (0.87–1.01) | 24% |
| Diabetes | 0.91 (0.84–0.99) | 45% | 0.96 (0.90–1.03) | 39% | 1.21 (1.08–1.35) | 54% |
| Heart disease | 0.86 (0.80–0.92) | 4% | 0.88 (0.83–0.94) | 0% | 0.94 (0.86–1.03) | 0% |
| Stroke | 0.81 (0.73–0.89) | 48% | 0.83 (0.76–0.90) | 66% | 1.02 (0.91–1.13) | 58% |

Unadjusted and adjusted OR (95% CI) for 10 units change. The association was assessed in each site and then combined by random effects meta-analyses. Statistically significant estimates are marked with bold numbers. I² values of 0%, 25%, 50% and 75% were considered no, low, moderate and high heterogeneity.

Adjusted for sex, age, smoke history, BMI and educational level baseline.

BMI, body mass index; FEV₁, forced expiratory volume in 1 s; FVC, forced vital capacity.

disease and stroke after adjustment for sex, age, smoking history, BMI and educational level (table 3 and figure 2). High FVC was also associated with a lower risk of diabetes, whereas a high FEV_1/FVC was associated with a higher risk in the adjusted analyses. Moderate heterogeneity across sites was found for the associations of diabetes and stroke with high lung function. The association between FVC and diabetes was larger in the sites from high-income than in low-income and middle-income countries (figure 2). No income-related association was found for other associations (figure 2).

Sensitivity analyses

In the sensitivity analyses, slight reductions were observed in the estimates when participants who were obese or had any cardiometabolic disease were excluded. Excluding participants with a $BMI \geq 30$ kg/m² resulted in the association between FVC and diabetes becoming statistically non-significant (online supplemental table S2). The association between FEV_1/FVC and diabetes remained statistically significant even after excluding participants with $FVC < LLN$ (OR (95% CI) 1.29 (1.10 to 1.51)).

Sex-stratified analyses

The association between a high FVC and high FEV_1/FVC and the onset of diabetes after adjustment for sex, age, smoking history, BMI and educational level was statistically significant in men but not in women. No sex differences were found regarding the association between lung function and heart disease or stroke (online supplemental table S3). Excluding obese participants did not change these associations.

Analyses using multilevel modelling

Analyses using multilevel logistic regression gave estimates that were very close to what was found in the models using meta-analyses (online supplemental table S4).

DISCUSSION

In this longitudinal study, a higher postbronchodilator FVC and FEV_1 were associated with a decreased risk of developing participant-reported heart disease and stroke, and a higher FVC was associated with a lower risk of diabetes. The risk of developing heart disease per 10% predicted FVC decreased by 14%. For the same increase in FVC, the risk of diabetes decreased by 9%, and the risk of stroke decreased by 19%. A high FEV_1/FVC was associated with an increased risk of developing diabetes. The association between lung function and the onset of heart disease was consistent across all geographical locations. However, there was moderate heterogeneity between sites regarding the risk of diabetes and stroke.

In the present study, participants with a high FVC had a decreased risk of having a participant-reported onset of diabetes during a mean follow-up period of

approximately 10 years. This result matches Engström *et al*'s findings in a Swedish population study.⁸ They found that a low FVC was associated with insulin resistance at baseline and a higher incidence of diabetes during the follow-up.²⁰ This is also in line with the results of McKeever *et al*, who reported that plasma glucose level 2 hours after oral glucose administration was inversely related to FVC and FEV_1 .⁹ An inverse association between FVC and the risk of type 2 diabetes was also found in a meta-analysis of 13 cohorts of patients with COPD.²¹ Surprisingly, we found that the association between high FVC and a lower risk of onset of diabetes was significant in men but not in women. In contrast to our results, Lee *et al* found no difference between men and women regarding the association between reduced FVC and 12-year incidence of diabetes,²² and Zaigham *et al* also reported a similar risk of diabetes in men and women with low lung function.²⁰ We have no clear explanation for the difference between our results and those previously reported, except that the present study included participants from countries with different income levels. In contrast, previous studies have been conducted in high-income countries.

In the current study, high FVC and FEV_1 were independently associated with a lower risk of onset of participant-reported heart disease. Previous studies have found that a low FVC is associated with increased coronary heart disease, myocardial infarction and cardiovascular death.^{11 12} The association between lung function and cardiometabolic diseases remained statistically significant in the sensitivity analysis, where we only included participants without any other cardiometabolic disease. We found no difference in the association between low lung function and heart disease between men and women. This is similar to what was reported by Lee *et al*²³ but contrary to the findings of Schroeder *et al*, where the association between FEV_1 and FVC and the 10-year incidence of coronary heart disease was stronger in women than men.¹² The current study found that higher FVC and FEV_1 were associated with a lower risk of stroke onset. This is in line with what has been reported by Li *et al* in a study using data from the UK Biobank.¹⁰ Previous studies have found that lower FEV_1 and FVC were associated with signs of atherosclerosis, such as higher carotid intima-media thickness²⁴ and reduced ankle-brachial pressure index.²⁵

Like Lee *et al* and Cuttica *et al*,^{23 26} we found no association between cardiovascular disease and a low FEV_1/FVC . However, we found that participants with a high FEV_1/FVC were more likely to have an onset of diabetes. This result does not align with that of a meta-analysis of a large number of patients with COPD, where no significant association between FEV_1/FVC and risk of type 2 diabetes was found.²¹ These results are, however, similar to the analyses of the baseline BOLD study, where airflow obstruction was inversely associated with diabetes.³ The association between high FEV_1/FVC and low FVC to a higher risk of diabetes supports a recent meta-analysis that concludes that restrictive but not obstructive lung




Figure 2 Association between forced vital capacity expressed as per cent of predicted and onset of diabetes, heart disease and stroke. Adjusted OR* (95% CI) for 10 units change. The association was assessed in each site and combined by random effects meta-analyses. I^2 values of 0%, 25%, 50% and 75% were considered no, low, moderate and high heterogeneity. *Adjusted for age, sex, smoke history, BMI and educational level baseline. BMI, body mass index.

function impairment is significantly associated with type 2 diabetes.²⁷ As we have not done diagnostic tests such as high-resolution CT scans, we cannot know to what extent this association with diabetes is related to interstitial lung disease or other restrictive lung disorders.

The explanation for the association between low lung function and cardiometabolic disease is not clear. It could be due to residual confounding from risk factors shared by a low lung function and cardiometabolic disease that have not been considered in the current and previous analyses. However, studies using polygenic scores²⁸ and Mendelian Randomisation studies^{29 30} suggest that FVC and FEV₁ are independent risk factors for cardiometabolic disease. Systemic inflammation could be one of the mechanisms connecting low lung function and cardiometabolic disease. In a recent study using data from two large Swedish populations, Rydell *et al* found that a low FVC and, to a lesser extent, FEV₁ were associated with higher levels of many cardiovascular disease-linked plasma proteins. In contrast, no such association was found for FEV₁/FVC.¹⁶ Likewise, Nerpin *et al*, analysing data from the NHANES population, found that a low FVC and FEV₁ but not FEV₁/FVC was associated with higher CRP levels.¹⁴ Another possibility is that the association between low lung and cardiometabolic diseases could result from suboptimal development of the lung and other organs related to environmental influences in utero and early childhood.³¹

The strengths of this study are its longitudinal design and inclusion of several geographical regions. Previous studies have mostly included participants from high-income countries, whereas the present study also includes low-income and middle-income countries. This is also the first study to examine the association between lung function and cardiometabolic diseases using postbronchodilator lung function. Another advantage is the rigorous quality control of the spirometry. The main limitation is that we use self-reported onset of cardiometabolic diseases. This is likely to be particularly problematic in low-income and middle-income countries with poorly developed health services focusing more on communicable diseases than non-communicable diseases. It is, therefore, possible that some of the associations related to diabetes and heart disease could be a consequence of undiagnosed preclinical disease. Diabetes is known to be associated with abnormal lung function, probably because of the effects of glucose on lung vasculature basement membranes.³² Similarly, low FVC and heart disease could reflect the effects of undiagnosed, mild, undiagnosed heart failure that is known to be associated with reduced FVC.³³ The data also do not enable us to distinguish between different heart diseases or stroke. Another limitation is the average response rate below 50%, which we addressed using inverse probability weights. Levels of particulate matter were not included in the regression models. However, in most cases, the associations found in this study were similar across sites with high levels of particulate matter, such as Karachi, and sites with low

levels of particulate matter, such as Reykjavik.³⁴ Another potential limitation is that we have not adjusted for the use of corticosteroids. We know, however, from a previous publication that the proportion of participants using corticosteroids is low in this population.³⁵ Therefore, it is unlikely that the use of corticosteroids has influenced our results substantially.

In the present study, high FVC was independently associated with a lower risk of participant-reported diabetes, heart disease and stroke. To our knowledge, FVC is not included in any risk score for predicting the risk of cardiometabolic events, although data also suggests that FVC predicted mortality more strongly than systolic blood pressure or BMI.³⁶ Our results and several previous studies suggest that including FVC will improve the precision of risk scores used to predict the onset of diabetes and cardiovascular diseases.

Author affiliations

¹Department of Medical Sciences Respiratory Medicine, Uppsala Universitet, Uppsala, Sweden

²National Heart and Lung Institute, Imperial College London, London, UK

³Medical Sciences, Uppsala University, Uppsala, Sweden

⁴KEM Hospital Pune Research Centre, Pune, Maharashtra, India

⁵The Epidemiological Laboratory, Khartoum, Sudan

⁶The University of the West Indies, Kingston, Jamaica

⁷Faculté de Médecine, Sousse, Tunisia

⁸Universite de Sousse Faculté de Medecine de Sousse, Sousse, Tunisia

⁹Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK

¹⁰Department of Medicine, Obafemi Awolowo University, Ife, Nigeria

¹¹Landspítali University Hospital, Reykjavik, Iceland

¹²Lung Clinic, Tartu University Hospital, Tartu, Estonia

¹³Independent Consultant, Pune, Maharashtra, India

¹⁴Pulmonary Medicine, SKIMS, Srinagar, India

¹⁵University of Cambridge, Cambridge, UK

¹⁶Liverpool University Hospitals NHS Foundation Trust, Liverpool, UK

¹⁷Community Health Sciences, Aga Khan University, Karachi, Pakistan

¹⁸University of Bergen, Bergen, Norway

¹⁹Respiratory Medicine, JSS Medical College and Hospital, Mysore, Karnataka, India

²⁰Philippine College of Chest Physicians, Manila, Philippines

²¹Department of Thoracic Medicine, Haukeland University Hospital, Bergen, Norway

²²Department of Respiratory Medicine, National Center for Cardiology and Internal Medicine, Bishkek, Kyrgyzstan

²³Kyrgyz-Swiss High Altitude Clinic and Medical Research Center, Tuja Ashu, Kyrgyzstan

X Rana Ahmed @RanaAttaElmula and Gregory E Erhabor @gregerhabor

Acknowledgements We thank the BOLD Collaborative Research Group: Hasan Hafizi, Anila Aliko, Donika Bardhi, Holta Tafa, Natasha Thanasi, Arian Mezzini, Alma Teferici, Dafina Todri, Jolanda Nikolla, Rezarta Kazasi, Hamid Hacene Cherkaski, Amira Bengrait, Tabarek Haddad, Ibtissem Zgaoula, Maamar Ghit, Abdelhamid Rouhbia, Soumaya Boudra, Feryal Atoui, Randa Yakoubi, Rachid Benali, Abdelghani Bencheikh, Nadia Ait-Khaled, Christine Jenkins, Guy Marks, Tessa Bird, Paola Espinel, Kate Hardaker, Brett Toelle, Michael Studnicka, Torkil Dawes, Bernd Lamprecht, Lea Schirhofer, Akramul Islam, Syed Masud Ahmed, Shayla Islam, Qazi Shafayetul Islam, Mesbah-Ul-Haque, Tridib Roy Chowdhury, Sukantha Kumar Chatterjee, Dulal Mia, Shyamal Chandra Das, Mizanur Rahman, Nazrul Islam, Shahaz Uddin, Nurul Islam, Luiza Khatun, Monira Parvin, Abdul Awal Khan, Maidul Islam, Herve Lawin, Arsene Kpangon, Karl Kpoussou, Gildas Agodokpessi, Paul Ayelo, Benjamin Fayomi, Bertrand Mbathou, Atongno Humphrey Ashu, Wan C. Tan, Wen Wang, NanShan Zhong, Shengming Liu, Jiachun Lu, Pixian Ran, Dali Wang, Jin-ping Zheng, Yumin Zhou, Rain Jogi, Hendrik Laja, Katrin Ulst, Vappu Zobel, Toomas-Julius Lill, Ayola Akim Adegnika, Tobias Welte, Isabelle Bodemann, Henning Geldmacher, Alexandra SchwedaLinow, Thorarinn Gislason, Bryndis Benediktssdottir, Kristin Jorundsdottir, Lovisa Gudmundsdottir, Sigrun

Gudmundsdottir, Gunnar Gudmundsson, Mahesh Rao, Parvaiz A. Koul, Sajjad Malik, Nissar A. Hakim, Umar Hafiz Khan, Rohini Chowgule, Vasant Shetye, Jonelle Raphael, Rosel Almeda, Mahesh Tawde, Rafiq Tadv, Sunil Katkar, Milind Kadam, Rupesh Dhanawade, Umesh Ghurup, Sanjay Juvekar, Siddhi Hirve, Sonmath Sambhudas, Bharat Chaidhary, Meera Tambe, Savita Pingale, Arati Umap, Archana Umap, Nitin Shelar, Sampada Devchakke, Sharda Chaudhary, Suvarna Bondre, Savita Walke, Ashlesha Gawhane, Anil Sapkal, Rupali Argade, Vijay Gaikwad, Sundeeep Salvi, Bill Brashier, Jyoti Londhe, Sapna Madas, Althea Aquart-Stewart, Akosua Francia Aikman, Talant M. Sooronbaev, Bermet M. Estebe-sova, Meerim Akmatallieva, Saadat Usenbaeva, Jypara Kydyrova, Eliza Bostonova, Ulan Sheraliev, Nuridin Marajapov, Nurgul Toktogulova, Berik Emilov, Toktogul Azilova, Gulnara Beishekeeva, Nasyikat Dononbaeva, Ajmal Tabyshova, Kevin Mortimer, Wezzie Nyapigoti, Ernest Mwangoka, Mayamiko Kambwili, Martha Chipeta, Gloria Banda, Suzgo Mkandawire, Justice Banda, Li-Cher Loh, Abdul Rashid, Siti Sholehah, Mohamed C. Benjelloun, Chakib Nejari, Mohamed Elbiaze, Karima El Rhazi, E.F.M. Wouters, G.J. Wesseling, Daniel Obaseki, Gregory Erhabor, Olayemi Awopeju, Olufemi Adewole, Amund Gulsvik, Tina Endresen, Lene Svendsen, Asaad A. Nafees, Muhammad Irfan, Zafar Fatmi, Aysha Zahidie, Natasha Shaukat, Meesha Iqbal, Luisito F. Idol, Teresita S. de Guia, Norberto A. Francisco, Camilo C. Roa, Fernando G. Ayuyao, Cecil Z. Tady, Daniel T. Tan, Sylvia Banal-Yang, Vincent M. Balanag, Jr., Maria Teresita N. Reyes, Renato B. Dantes, Renato B. Dantes, Lourdes Amarillo, Lakan U. Berratio, Lenora C. Fernandez, Norberto A. Francisco, Gerard S. Garcia, Teresita S. de Guia, Luisito F. Idol, Sullian S. Naval, Thessa Reyes, Camilo C. Roa, Jr., Flordeliza Sanchez, Leander P. Simpao, Ewa Nizankowska-Mogilnicka, Jakub Frey, Rafal Harat, Filip Mejza, Pawel Nastalek, Andrzej Pajak, Wojciech Skucha, Andrzej Szczekliak, Magda Twardowska, Cristina Barbara, Fatima Rodrigues, Herminia Dias, Joao Cardoso, João Almeida, Maria Joao Matos, Paula Simão, Moutinho Santos, Reis Ferreira, M. Al Ghobain, H. Alorainy, E. El-Hamad, M. Al Hajjaj, A. Hashi, R. Dela, R. Fanuncio, E. Doloriel, I. Marciano, L. Safia, Eric Bateman, Anamika Jithoo, Desiree Adams, Edward Barnes, Jasper Freeman, Anton Hayes, Siphon Hlengwa, Christine Johannissen, Mariana Koopman, Innocentia Louw, Ina Ludick, Alta Olckers, Johanna Ryck, Janita Storbeck, Kirithi Gunasekera, Rajitha Wickremasinghe, Asma Elsony, Hana A. Elsadig, Nada Bakery Osman, Bandar Salah Noory, Monjida Awad Mohamed, Hasab Alrasoul Akasha Ahmed Osman, Namariq Moham ed Elhassan, Abdel Mu'is El Zain, Marwa Mohamed Mohamaden, Suhaiba Khalifa, Mahmoud Elhadi, Mohand Hassan, Dalia Abdelmonam, Christer Janson, Inga Sif Olafsdottir, Katarina Nisser, Ulrike SpetzNystrom, Gunilla Hagg, GunMarie Lund, Terence Seemungal, Fallon Lutchmansingh, Liane Conyette, Imed Harrabi, Myriam Denguezli, Zouhair Tabka, Hager Daldoul, Zaki Boukheroufa, Firas Chouikha, Wahbi Belhaj Khalifa, Ali Kocabas, Attila Hancioglu, Ismail Hanta, Sedat Kuleci, Ahmet Sinan Turkyilmaz, Sema Umut, Turgay Unalan, Peter G.J. Burney, Anamika Jithoo, Louisa Gnatiuc, Hadia Azar, Jaymini Patel, Caron Amor, James Potts, Michael Tumilty, Fiona McLean, Risha Dudhaiya, A. Sonia Buist, Mary Ann McBurnie, William M. Vollmer, Suzanne Gillespie, Sean Sullivan, Todd A. Lee, Kevin B. Weiss, Robert L. Jensen, Robert Crapo, Paul Enright, David M. Mannino, John Cain, Rebecca Copeland, Dana Hazen, and Jennifer Methvin and all participants and field workers/research assistants for their time and effort in this study.

Contributors CJ and AFSA conceived the study. Under the supervision of AFSA, CJ performed data analysis and prepared the initial draft with input from PGJB. JP assisted with the preparation of the databases and analyses. CJ, JP, AM, DA, RA, AA-S, MD, GD, GEE, TG, IH, RJ, SKJ, BK-B, PK, KM, AAN, RN, PAM, SNMP, AØR, TS, PGJB and AFSA contributed to acquisition of the data and to further drafting and final approval of the paper. CJ is the guarantor for this paper, accepts full responsibility for the work and the conduct of the study, has access to the data and controls the decision to publish.

Funding Funding The Burden of Obstructive Lung Disease (BOLD) study has been supported by grants from the Wellcome Trust (085790/Z/08/Z) and Medical Research Council (MR/R011192/1). The follow-up study in the Nordic centres (Bergen, Reykjavik, Tartu, Uppsala) was supported by an unrestricted grant from AstraZeneca (ESR-17-13417).

Competing interests RN declares support from Astra Zeneca related to and the Norwegian Respiratory Society not related to the manuscript. AØR declares support from Boehringer Ingelheim and the Endowment of Timber Merchant A. Delphin and Wife unrelated to the manuscript. PGJB declares support from the Wellcome Trust related to the manuscript. AFSA declares support from the Colt Foundation not related to the manuscript. None of the other authors declare any conflicts of interest.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval This study involves human participants and was approved by all sites from their local ethics committee, the follow-up study was also approved by Imperial College London Research Ethics Committee (ref. 17IC4272), and participants provided informed consent. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Deidentified participant data and questionnaires of the BOLD study may be shared after publication on a collaborative basis on reasonable request made to AFSA (a.amaral@imperial.ac.uk). Requesting researchers will be required to submit an analysis plan.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Christer Janson <http://orcid.org/0000-0001-5093-6980>
 Andrei Malinovschi <http://orcid.org/0000-0002-4098-7765>
 Rana Ahmed <http://orcid.org/0000-0003-4389-7084>
 Graham Devereux <http://orcid.org/0000-0002-0024-4887>
 Ben Knox-Brown <http://orcid.org/0000-0001-5573-4413>
 Peter G J Burney <http://orcid.org/0000-0001-8635-5678>
 Andre F S Amaral <http://orcid.org/0000-0002-0369-9449>

REFERENCES

- Stållberg B, Janson C, Larsson K, *et al*. Real-world retrospective cohort study ARCTIC shows burden of comorbidities in Swedish COPD versus non-COPD patients. *NPJ Prim Care Respir Med* 2018;28:33.
- Barnett K, Mercer SW, Norbury M, *et al*. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380:37–43.
- Triest FJJ, Studnicka M, Franssen FME, *et al*. Airflow Obstruction and Cardio-metabolic Comorbidities. *COPD* 2019;16:109–17.
- Kulbacka-Ortiz K, Triest FJJ, Franssen FME, *et al*. Restricted spirometry and cardiometabolic comorbidities: results from the international population based BOLD study. *Respir Res* 2022;23:34.
- Margretardottir OB, Thorleifsson SJ, Gudmundsson G, *et al*. Hypertension, systemic inflammation and body weight in relation to lung function impairment—an epidemiological study. *COPD* 2009;6:250–5.
- Amaral AFS, Patel J, Gnatiuc L, *et al*. Association of pulse wave velocity with total lung capacity: A cross-sectional analysis of the BOLD London study. *Respir Med* 2015;109:1569–75.
- Takase M, Yamada M, Nakamura T, *et al*. Association between lung function and hypertension and home hypertension in a Japanese population: the Tohoku Medical Megabank Community-Based Cohort Study. *J Hypertens* 2023;41:443–52.
- Engström G, Hedblad B, Nilsson P, *et al*. Lung function, insulin resistance and incidence of cardiovascular disease: a longitudinal cohort study. *J Intern Med* 2003;253:574–81.
- McKeever TM, Weston PJ, Hubbard R, *et al*. Lung function and glucose metabolism: an analysis of data from the Third National Health and Nutrition Examination Survey. *Am J Epidemiol* 2005;161:546–56.
- Li G, Lu Y, Qiao Y, *et al*. Role of Pulmonary Function in Predicting New-Onset Cardiometabolic Diseases and Cardiometabolic Multimorbidity. *Chest* 2022;162:421–32.
- Engström G, Lind P, Hedblad B, *et al*. Lung function and cardiovascular risk: relationship with inflammation-sensitive plasma proteins. *Circulation* 2002;106:2555–60.
- Schroeder EB, Welch VL, Couper D, *et al*. Lung function and incident coronary heart disease: the Atherosclerosis Risk in Communities Study. *Am J Epidemiol* 2003;158:1171–81.
- Ólafsdóttir IS, Gíslason T, Gudnason V, *et al*. CRP is associated with lung function decline in men but not women: a prospective study. *Respir Med* 2013;107:91–7.

- 14 Nerpin E, Jacinto T, Fonseca JA, *et al.* Systemic inflammatory markers in relation to lung function in NHANES. 2007-2010. *Respir Med* 2018;142:94–100.
- 15 Thorleifsson SJ, Margretardottir OB, Gudmundsson G, *et al.* Chronic airflow obstruction and markers of systemic inflammation: results from the BOLD study in Iceland. *Respir Med* 2009;103:1548–53.
- 16 Rydell A, Nerpin E, Zhou X, *et al.* Cardiovascular disease-linked plasma proteins are mainly associated with lung volume. *ERJ Open Res* 2023;9:00321–2022.
- 17 Amaral AFS, Potts J, Knox-Brown B, *et al.* Cohort Profile: Burden of Obstructive Lung Disease (BOLD) study. *Int J Epidemiol* 2023;52:e364–73.
- 18 Buist AS, McBurnie MA, Vollmer WM, *et al.* International variation in the prevalence of COPD (The BOLD Study): a population-based prevalence study. *The Lancet* 2007;370:741–50.
- 19 Valenzuela PL, Carrera-Bastos P, Castillo-García A, *et al.* Obesity and the risk of cardiometabolic diseases. *Nat Rev Cardiol* 2023;20:475–94.
- 20 Zaigham S, Nilsson PM, Wollmer P, *et al.* The temporal relationship between poor lung function and the risk of diabetes. *BMC Pulm Med* 2016;16:75.
- 21 Peng Y, Zhong G-C, Wang L, *et al.* Chronic obstructive pulmonary disease, lung function and risk of type 2 diabetes: a systematic review and meta-analysis of cohort studies. *BMC Pulm Med* 2020;20:137.
- 22 Lee JH, Lee HS, Lee YJ. Lung function as a predictor of incident type 2 diabetes in community-dwelling adults: A longitudinal finding over 12 years from the Korean Genome and Epidemiology Study. *Diabetes Metab* 2020;46:392–9.
- 23 Lee HM, Liu MA, Barrett-Connor E, *et al.* Association of lung function with coronary heart disease and cardiovascular disease outcomes in elderly: the Rancho Bernardo study. *Respir Med* 2014;108:1779–85.
- 24 Takase M, Yamada M, Nakamura T, *et al.* The Association of Lung Function and Carotid Intima-Media Thickness in a Japanese Population: The Tohoku Medical Megabank Community-Based Cohort Study. *J Atheroscler Thromb* 2023;30:1022–44.
- 25 Sugiura T, Dohi Y, Takagi Y, *et al.* Close Association between Subclinical Atherosclerosis and Pulmonary Function in Middle-Aged Male Smokers. *J Atheroscler Thromb* 2020;27:1230–42.
- 26 Cuttica MJ, Colangelo LA, Dransfield MT, *et al.* Lung Function in Young Adults and Risk of Cardiovascular Events Over 29 Years: The CARDIA Study. *J Am Heart Assoc* 2018;7:e010672.
- 27 Zhou Y, Meng F, Wang M, *et al.* Reduced lung function predicts risk of incident type 2 diabetes: insights from a meta-analysis of prospective studies. *Endocr J* 2022;69:299–305.
- 28 Zaigham S, Gonçalves I, Center RG, *et al.* Polygenic scores for low lung function and the future risk of adverse health outcomes. *Cardiovasc Diabetol* 2022;21:230.
- 29 Au Yeung SL, Borges MC, Lawlor DA, *et al.* Impact of lung function on cardiovascular diseases and cardiovascular risk factors: a two sample bidirectional Mendelian randomisation study. *Thorax* 2022;77:164–71.
- 30 Higbee DH, Granell R, Sanderson E, *et al.* Lung function and cardiovascular disease: a two-sample Mendelian randomisation study. *Eur Respir J* 2021;58:2003196.
- 31 Nobile S, Di Sipio Morgia C, Vento G. Perinatal Origins of Adult Disease and Opportunities for Health Promotion: A Narrative Review. *J Pers Med* 2022;12:157.
- 32 Zheng H, Wu J, Jin Z, *et al.* Potential Biochemical Mechanisms of Lung Injury in Diabetes. *Aging Dis* 2017;8:7–16.
- 33 Baum C, Ojeda FM, Wild PS, *et al.* Subclinical impairment of lung function is related to mild cardiac dysfunction and manifest heart failure in the general population. *Int J Cardiol* 2016;218:298–304.
- 34 Amaral AFS, Burney PGJ, Patel J, *et al.* Chronic airflow obstruction and ambient particulate air pollution. *Thorax* 2021;76:1236–41.
- 35 Gnatiuc L, Buist AS, Kato B, *et al.* Gaps in using bronchodilators, inhaled corticosteroids and influenza vaccine among 23 high- and low-income sites. *Int J Tuberc Lung Dis* 2015;19:21–30.
- 36 Gupta RP, Strachan DP. Ventilatory function as a predictor of mortality in lifelong non-smokers: evidence from large British cohort studies. *BMJ Open* 2017;7:e015381.