



OPEN ACCESS

EULAR points to consider for the diagnosis and management of rheumatic immune-related adverse events due to cancer immunotherapy with checkpoint inhibitors

Marie Kostine ,¹ Axel Finckh ,² Clifton O Bingham 3rd,³ Karen Visser,⁴ Jan Leipe,^{5,6} Hendrik Schulze-Koops,⁶ Ernest H Choy,⁷ Karolina Benesova,⁸ Timothy R D J Radstake,⁹ Andrew P Cope,¹⁰ Olivier Lambotte,¹¹ Jacques-Eric Gottenberg ,¹² Yves Allenbach ,¹³ Marianne Visser,¹⁴ Cindy Rusthoven,¹⁴ Lone Thomasen,¹⁵ Shahin Jamal,¹⁶ Aurélien Marabelle,¹⁷ James Larkin,¹⁸ John B A G Haanen,¹⁹ Leonard H Calabrese ,²⁰ Xavier Mariette,^{21,22} Thierry Schaevebeke¹

Handling editor Josef S Smolen

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2020-217139>).

For numbered affiliations see end of article.

Correspondence to

Dr Marie Kostine, Rheumatology, Centre Hospitalier Universitaire de Bordeaux Groupe hospitalier Pellegrin, Bordeaux 33000, France; marie.kostine@chu-bordeaux.fr

XM and TS are joint senior authors.

Received 10 February 2020
Revised 3 April 2020
Accepted 7 April 2020
Published Online First 23 April 2020



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Kostine M, Finckh A, Bingham 3rd CO, et al. *Ann Rheum Dis* 2021;**80**:36–48.

ABSTRACT

Background Rheumatic and musculoskeletal immune-related adverse events (irAEs) are observed in about 10% of patients with cancer receiving checkpoint inhibitors (CPIs). Given the recent emergence of these events and the lack of guidance for rheumatologists addressing them, a European League Against Rheumatism task force was convened to harmonise expert opinion regarding their identification and management.

Methods First, the group formulated research questions for a systematic literature review. Then, based on literature and using a consensus procedure, 4 overarching principles and 10 points to consider were developed.

Results The overarching principles defined the role of rheumatologists in the management of irAEs, highlighting the shared decision-making process between patients, oncologists and rheumatologists. The points to consider inform rheumatologists on the wide spectrum of musculoskeletal irAEs, not fulfilling usual classification criteria of rheumatic diseases, and their differential diagnoses. Early referral and facilitated access to rheumatologist are recommended, to document the target organ inflammation. Regarding therapeutic, three treatment escalations were defined: (1) local/systemic glucocorticoids if symptoms are not controlled by symptomatic treatment, then tapered to the lowest efficient dose, (2) conventional synthetic disease-modifying antirheumatic drugs, in case of inadequate response to glucocorticoids or for steroid sparing and (3) biological disease-modifying antirheumatic drugs, for severe or refractory irAEs. A warning has been made on severe myositis, a life-threatening situation, requiring high dose of glucocorticoids and close monitoring. For patients with pre-existing rheumatic disease, baseline immunosuppressive regimen should be kept at the lowest efficient dose before starting immunotherapies.

Conclusion These statements provide guidance on diagnosis and management of rheumatic irAEs and aim to support future international collaborations.

INTRODUCTION

Although the concept of immunotherapy in cancer is far from new, monoclonal antibodies targeting immunological checkpoints or ‘checkpoint inhibitors’ (CPIs) represent a growing class of agents across multiple tumour types and at all stages of disease. Agents targeting the T-cell cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) or the programmed cell death-(ligand) 1 (PD-1/PD-L1) coinhibitory receptors marked a turning point in the success of immunotherapeutic approaches.^{1–3} By enhancing antitumour T-cell activity, unprecedented long-lasting tumour responses were observed in patients with unresectable or advanced metastatic disease.^{4–7} The clinical value of these immune CPIs, as single agents or in combination, is being investigated in various solid tumours and haematological malignancies, and their use is expanding rapidly.⁸ So far, the Food and Drug Administration and the European Medicines Agency approved seven immune checkpoint-blocking antibodies in selected cancers: one anti-CTLA-4 (ipilimumab), three anti-PD-1 (nivolumab, pembrolizumab and cemiplimab) and three anti-PD-L1 (atezolizumab, avelumab and durvalumab).

The T-cell activation induced by CPIs commonly promotes inflammatory or autoimmune-like side effects, known as immune-related adverse events (irAEs).⁹ Compared with conventional cancer therapies, this spectrum of toxicities is unique and can affect any organ system, most frequently the skin, gastrointestinal tract, endocrine glands and lung. Among irAEs, specific rheumatic manifestations have been described rather rarely in randomised clinical trials, but are much more common in clinical practice. The clinical features of rheumatic irAEs have been described in a growing number of case series and reports.¹⁰ However, despite the growing interest for irAEs among rheumatologists, evidence is lacking for the optimal diagnostic approach and the management of these patients in ways that also permit effective antitumour therapy to continue. According to a recent survey, a large proportion of

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Ann Rheum Dis: first published as 10.1136/annrheumdis-2020-217139 on 23 April 2020. Downloaded from <http://ard.bmj.com/> on May 20, 2025 at Department GEZ-LTA Erasmus Hogeschool

rheumatologists have limited experience and little confidence in managing rheumatic irAEs, highlighting the need for education and recommendations in this emerging condition.¹¹

In 2017, the European Society for Medical Oncology developed clinical guidelines for the management of immune toxicities and mentioned the paucity of literature on management of rheumatic irAEs.¹² Three other consensus recommendations have been proposed by the Society for Immunotherapy of Cancer, the American Society of Clinical Oncology and The National Comprehensive Cancer Network, which among others included the management of inflammatory arthritis, polymyalgia rheumatica and myositis.^{13–15} This European League Against Rheumatism initiative assembled international experts primarily from the rheumatology and immunology but also the oncology field with the explicit goal of generating the first set of recommendations for the diagnosis and the management of rheumatic irAEs arising as a direct consequence of CPI. Rheumatologists, but also in some countries internists and immunologists, have to play a pivotal role in developing with the oncologists a patient-centred approach to improve the management of rheumatic irAEs. While the initiative primarily set out to guide clinicians, it is noteworthy that there is limited and rapidly changing literature and that future additional studies can drastically change the profile for diagnosis and management. This area will be a continually evolving field; therefore, the accompanying comments may also serve as a framework for future longitudinal cohorts and/or clinical studies.

METHODS

After approval by the European League Against Rheumatism Executive Committee, an international task force was convened to develop points to consider for the diagnosis and the management of rheumatic irAEs due to cancer immunotherapy. Among these members, were 19 clinical experts from Europe and North America (14 rheumatologists including 2 delegates of the European League Against Rheumatism young rheumatologists' network EMEUNET, 2 internists and 3 oncologists), 1 clinical epidemiologist, 1 allied health professional and 2 patient representatives from the PARE network of patient research partners. The process adhered to the updated European League Against Rheumatism standardised operating procedures for the development of recommendations.¹⁶

In July 2018, the first meeting was convened in Zürich, Switzerland, to define the focus of the task force, identify the target population and the research questions for the systematic literature review (SLR). The SLR was performed by the research fellow (MK), with support from the clinical epidemiologist (AF) and a librarian (Catherine Weill), to identify relevant publications through December 2018. Based on the findings of the SLR, a first draft of points to consider including 12 items was prepared by the fellow (MK) and the two convenors (TS and XM).

SLR results were presented at a second meeting that was held in Zürich, Switzerland, in January 2019. Following the evaluation of literature and a group discussion of the first draft of propositions, the task force formulated overarching principles and consensus statements. Each proposal was then submitted to a voting process, requiring at least 75% of votes in the first ballot for each recommendation to be accepted. In case this threshold was not achieved, further discussion and textual changes were proposed for a second round, for which a 67% majority was required. Five members of the task force could not attend this second meeting, but they subsequently commented and voted on each statement by email. The level of evidence (LoE) and

grade of recommendation was based on the Oxford Levels of Evidence.¹⁷ After this second face-to-face meeting, members of the task force were asked to anonymously rate each item in an online survey, on a scale of 0 (absolutely disagree) to 10 (absolutely agree) to assess the level of agreement (LoA). Furthermore, the task force agreed on adding relevant references published between the SLR and the writing of this manuscript. The manuscript was reviewed and approved by all task force members and the European League Against Rheumatism Executive Committee before submission.

RESULTS

Systematic literature review

The literature search strategy and summary of results are detailed in online supplementary data. The first objective was to identify phase III clinical trials to assess the frequency and type of rheumatic and musculoskeletal diseases' (RMDs) complaints associated with CPI compared with the comparator group. The search was performed using Medline, Embase and the Cochrane Library, through December 2018. Among 630 references identified, 22 studies were selected for inclusion. The second objective was to obtain detailed information on rheumatic and musculoskeletal symptoms that have been described under CPI treatment. The third objective was to assess outcomes in patients with pre-existing autoimmune diseases. Therefore, relevant keywords relative to three key domains were used in Medline and Embase databases: immune CPIs, rheumatic and systemic diseases and adverse events. Abstracts from the last two European League Against Rheumatism and American College of Radiology meetings were included, combined with manual searches from references of the selected articles. From among 2156 references identified, 170 were included, including pharmacovigilance registries (n=5), case series (n=51) and case reports (n=114).

After group discussion of the results of the SLR, the consensus process was initiated and the full task force agreed on a final set of 4 overarching principles and 10 points to consider (table 1).

Overarching principles

A. Rheumatic and musculoskeletal immune-related adverse events can occur as manifestations in cancer patients receiving immunotherapy with checkpoint inhibitors (LoE na; LoA 9.6).

Analysis of phase III clinical trials revealed that arthralgia, arthritis, myalgia, myositis, dry mouth, musculoskeletal and back pain were reported in patients receiving CPI. However, their frequency was not significantly different to that of patients receiving chemotherapy or placebo.^{5 18–38} Data from several series, both retrospective and prospective, reporting prevalences of rheumatic irAEs in real life, ranging from 1.5% to 22%, suggest that rheumatic irAEs are under-reported in clinical trials.^{39–54} Of note, an heterogeneous definition of rheumatic irAEs may explain such wide interval. Many clinical trials do not report rheumatic irAEs (by disregarding of musculoskeletal/rheumatic events as a distinct organ system, even in the online supplementary data) or partially only report high-grade and/or frequent adverse events (ie, occurring in $\geq 10\%$ of the patients). Therefore, the task force wanted to emphasise with this first principle that rheumatic and musculoskeletal manifestations are a relevant part of the broad spectrum of irAEs.

B. Management of rheumatic and musculoskeletal immune-related adverse events should be based on a shared decision-making process between patients, oncologists and rheumatologists (LoE na; LoA 9.5).

Recommendation

Table 1 Overarching principles and points to consider for the diagnosis and management of rheumatic irAEs

		LoE	GoR	LoA (0–10) mean (SD)
Overarching principles				
A.	Rheumatic and musculoskeletal immune-related adverse events can occur as manifestations in cancer patients receiving immunotherapy with checkpoint inhibitors.	n.a.	n.a.	9.6 (0.7)
B.	Management of rheumatic and musculoskeletal immune-related adverse events should be based on a shared decision-making process between patients, oncologists and rheumatologists.	n.a.	n.a.	9.5 (1.1)
C.	Rheumatologists should engage with oncologists to contribute to the inter-disciplinary care of patients presenting with musculoskeletal signs and symptoms.	n.a.	n.a.	9.1 (1.2)
D.	The role of rheumatologists is to assist oncologists in differential diagnosis and to relieve rheumatic and musculoskeletal symptoms to an acceptable level enabling patients to maintain effective cancer immunotherapy.	n.a.	n.a.	9.5 (0.9)
Points to consider				
1.	Rheumatologists should be aware of the wide spectrum of clinical presentations of rheumatic and/or systemic immune-related adverse events that often do not fulfil traditional classification criteria of RMDs.	4	C	9.5 (1.2)
2.	Oncologists should be encouraged to consult rheumatologists promptly for assessment when rheumatic musculoskeletal and systemic signs or symptoms are suspected due to immunotherapy, and rheumatologists should provide facilitated access for such patients.	5	D	9.4 (1.3)
3.	Metastases, paraneoplastic syndromes and unrelated rheumatic diseases should be considered as a potential differential diagnosis of rheumatic immune-related events. The comprehensive assessment should be focused on documenting evidence of target organ inflammation, and based on history, clinical features, laboratory tests, imaging and/or biopsy.	4	C	9.5 (0.9)
4.	In case of inefficacy of symptomatic treatment and depending on the disease severity, local and/or systemic glucocorticoids should be considered for immune-related rheumatic and systemic symptoms. Dose regimen and route of administration should be decided according to the clinical entity and activity. When improvement is achieved, systemic glucocorticoids should be tapered to the lowest effective dose to control the symptoms.	4	C	9.4 (1)
5.	csDMARD should be considered in patients with insufficient response to acceptable dose of glucocorticoids or requiring glucocorticoid-sparing.	4	C	9 (1.2)
6.	For patients experiencing severe immune-related rheumatic and systemic immune-related adverse events or with insufficient response to csDMARD, bDMARD may be considered, with TNF or IL-6 inhibitors being the preferred options for inflammatory arthritis.	4	C	8.8 (1.2)
7.	The decision to hold or to continue the cancer immunotherapy should be based on the severity of rheumatic immune-related adverse events, the extent of required immunosuppressive regimen, the tumour response and its duration, as well as the future oncology treatment plan, in a shared decision with the patient.	5	D	9.4 (1)
8.	Myositis may be a severe condition. Immunotherapy withdrawal needs to be discussed. In the presence of life-threatening manifestations (bulbar symptoms (dysphagia, dysarthria, dysphonia), dyspnoea and myocarditis), high dose of glucocorticoids, IVIg and/or plasma exchange should be considered; immunotherapy withdrawal is always necessary.	4	C	8.9 (1.2)
9.	A pre-existing autoimmune rheumatic and/or systemic disease should not preclude the use of cancer immunotherapy. Baseline immunosuppressive regimen should be kept at the lowest dose possible (for glucocorticoids, below 10 mg prednisone per day if possible). However, many patients may have a flare of the underlying condition and/or immune-related adverse events, requiring the use of glucocorticoids and/or DMARDs.	4	C	9 (1.3)
10.	Before initiation of cancer immunotherapy, there is no indication to test every patient for the presence of autoantibodies. In the case of unexplained rheumatic, musculoskeletal or systemic symptoms, a complete rheumatological assessment should be performed.	5	D	9 (1.3)

GoR: A: based on consistent level 1 studies; B: based on consistent level 2 or 3 studies or extrapolations from level 1 studies; C: based on level 4 studies or extrapolations from level 2 or 3 studies; D: based on level 5 studies or on troublingly inconsistent or inconclusive studies of any level.

LoE: 1a: systematic review of RCTs; 1b: individual RCT; 2a: systematic review of cohort studies; 2b: individual cohort study (including low-quality RCT); 3a: systematic review of case-control studies; 3b: individual case-control study; 4: case-series (and poor quality cohort and case-control studies); 5: expert opinion without explicit critical appraisal, or based on physiology, bench research or 'first principles'.

bDMARD, biological disease-modifying antirheumatic drug; csDMARD, conventional synthetic disease-modifying antirheumatic drug; DMARD, disease-modifying antirheumatic drug; GoR, grade of recommendation; IL-6, interleukin 6; irAEs, immune-related adverse events; IVIg, intravenous immunoglobulin; LoA, level of agreement; LoE, level of evidence; RCT, randomised clinical trial; RMD, rheumatic and musculoskeletal disease; TNF, tumour necrosis factor.

Rheumatic and musculoskeletal irAEs occur in a context of cancer; therefore, a dialogue between rheumatologists and oncologists is important to balance the harm and risk of oncology treatment and immunosuppressive drugs. The most important stakeholder is the patient. Shared decision between a patient and his/her rheumatologist is a fundamental principle of RMDs management, as illustrated by its representation as an overarching principle in several European League Against Rheumatism recommendations.^{55–57} Because evidence-based data for irAEs management are limited, and irAEs can have a large impact on the quality of life, patient's preferences and discussions concerning risks and benefits of each treatment option are even more important.

C. Rheumatologists should engage with oncologists to contribute to the inter-disciplinary care of patients presenting with musculoskeletal signs and symptoms (LoE na; LoA 9.1).

irAEs may affect any organ system including the rheumatic and musculoskeletal system. Some patients may even experience multiple organ toxicities in sequence or concurrently. The importance of developing a local multidisciplinary network of oncologists and specialists of all organ system potentially involved in the management of irAEs has been recently highlighted.^{58 59} Rheumatologists should actively engage in these local multidisciplinary networks as valuable members due to their knowledge of clinical immunology, their expertise in multiorgan autoimmune disease and their long-standing

experience with the use of immunosuppressive drugs and biological therapies.⁶⁰ This engagement should also include efforts aimed at improving patient education before or when starting cancer immunotherapy, to prevent delay in diagnosis when rheumatic or musculoskeletal side effects occur. Patients have reported that they were informed about other immune-related side effects more than rheumatic or musculoskeletal symptoms.⁶¹ Furthermore, since the rheumatic and the cancer disease both induce high impact on the patient's life, even when independently considered and when disease activity is controlled (ie, fatigue, pain, functional impairments, emotional problems, secondary effects of the treatments), the value of an interdisciplinary collaboration between the rheumatologist and the oncologist is worthwhile.

D. The role of rheumatologists is to assist oncologists in establishing the diagnosis and to relieve rheumatic and musculoskeletal symptoms to an acceptable level enabling patients to maintain effective cancer immunotherapy (LoE na; LoA 9.5).

This principle aimed to better define the role of rheumatologists as oncologists' partners based on the clinical experience of the task force members. Once a patient with cancer receiving immunotherapy is referred for evaluation of rheumatic or musculoskeletal symptoms, the rheumatologist should consider several potential aetiologies: tumour progression, paraneoplastic syndromes, non-rheumatic events (ie, viral infection, thrombosis, endocrine abnormality), all already considered by the referring oncologist, or rheumatic/systemic irAE or immune non-related adverse events. This aspect of differential diagnosis is also described in more detail in recommendation 3. Once a rheumatic irAE is diagnosed, the supervising rheumatologist should propose an appropriate treatment to relieve patient's symptoms to an acceptable level with the objective of maintaining quality of life and permitting continuation of effective cancer immunotherapy, if this is recommended by the oncologist. This treatment goal is different to classic rheumatic entities, in which usually remission is the targeted treatment outcome.

Points to consider

1. Rheumatologists should be aware of the wide spectrum of clinical presentations of rheumatic and/or systemic immune-related adverse events that often do not fulfil traditional classification criteria of RMDs (LoE 4; LoA 9.5).

While arthralgia and myalgia were the most commonly reported rheumatic irAEs in clinical trials, numerous case series and reports have captured a broader spectrum of de novo rheumatic and systemic manifestations that can occur with cancer immunotherapy.^{62–64} Polymyalgia rheumatica (PMR)-like syndromes and inflammatory arthritis syndromes are two of the major clinical presentations encountered.^{39–41 48 65 66} PMR-like manifestation occurred with a median exposure time to CPI of 60 days, but also much later (IQR 24–210 days). Exposure time to CPI was generally longer for patients experiencing inflammatory arthritis (median 120 days, IQR 48–262 days). In addition, a variety of other rheumatic syndromes have been reported. These include arthralgia; monoarthritis, oligoarthritis or polyarthritis; reactive arthritis; psoriatic arthritis (PsA); remitting seronegative symmetrical synovitis with pitting oedema (RS3PE); tenosynovitis; enthesitis; non-inflammatory musculoskeletal conditions and osteoarthritis.^{41 44 46 51 66–77} Importantly, autoantibodies are often absent. In arthritis, only a few patients are positive for rheumatoid factor (RF; n=20, range 18–246 UI/mL) and/or anti-citrullinated peptide antibodies (ACPAs; n=14, range 18–614 U/mL).⁷⁸ Instead, positivity of antinuclear antibodies

(ANAs) is observed, but often at a low titre (range 1:80 to 1:3200, one patient with ANA 1:12 800 and only 35 patients with ANA >1:160). Similarly, acute phase reactants may be normal in some patients with PMR-like presentations.⁴¹ Overall, around 20% of patients fulfilled classification criteria of rheumatoid arthritis (RA) (55/271) or PMR (11/52). This percentage was higher (55%) for PsA (6/11), as well as in a recent series of PMR-like syndrome (37/49; 75%).⁷⁹ The first observation of recurrent pseudogout flares 7 to 10 days after each nivolumab infusion has been recently reported.⁸⁰

Several cases of myositis have been reported, with frequent limb-girdle myalgia and weakness that may mimic a PMR-like condition.^{81–83} Because it represents a potentially life-threatening complication, the task force decided to formulate a dedicated recommendation on myositis (*recommendation 8*).

Among systemic manifestations, sicca syndrome has been described early on, presenting mainly with dry mouth, and possible associated neurological symptoms in a few patients.^{40 48 65 66 84–87} Two major studies on CPI-induced sicca syndrome were published in 2019 and therefore included in this manuscript. The ImmunoCancer International Registry reported on 26 patients experiencing CPI-associated sicca syndrome. This mainly included men, with frequent organ-specific autoimmune manifestations but lower prevalence of autoantibodies (52% ANA, 20% Ro/SS-A, 9% RF, 8% La/SS-B) in comparison with classical Sjögren's syndrome.⁸⁸ Interestingly, a predominant T-cell infiltrate with acinar destruction has been reported in salivary glands, distinct from the histological profile of idiopathic primary Sjögren's syndrome. Authors hypothesise that CPI therapy may break immune tolerance locally leading to the activation of cytotoxic T cells damaging the salivary epithelium.⁸⁹

Other systemic manifestations have been described, including sarcoidosis or sarcoid-like reactions.^{90–93} The diagnosis is usually suspected through imaging when new hilar lymphadenopathy or pulmonary nodules are detected in imaging, requiring biopsy. Half of patients experienced cutaneous manifestations (nodules, rash), and some patients had cough/dyspnoea (29%) and arthralgia/arthritis (18%). Uveitis, parotitis, hypercalcaemia and neurological symptoms are rarely reported. Some patients experienced systemic sclerosis or scleroderma-like reactions, all presenting with skin thickening, but only one with new-onset Raynaud's phenomenon.^{48 94–96} None tested positive for specific autoantibodies. Since PD-1-deficient mice spontaneously developed lupus-like autoimmune diseases with arthritis and glomerulonephritis, such clinical phenotypes could be expected in patients treated with anti PD-(L)1 agents, but are not observed. A few cases of lupus-like cutaneous reaction and one Jaccoud arthropathy have yet been reported with anti-PD-1 agents, and only one lupus-like nephritis was attributed to anti-CTLA-4 treatment.^{97–102}

All vessel-sized vasculitis (eg, large, medium and small vessels) with various clinical manifestations, including purpura, digital necrosis arthralgia, arthritis, myalgia, fever, fatigue and abdominal pain have also been reported.^{40 48 84 103–115} Of note, ANA, antineutrophil cytoplasmic antibodies (ANCA), cryoglobulin and RF were rarely positive. Analysis of the WHO pharmacovigilance database revealed that temporal arteritis (n=16) was particularly over-reported with ipilimumab monotherapy treatment.¹¹⁴ The first case of granulomatosis with polyangiitis with a high anti-PR3 ANCA titre was reported in 2019.¹¹⁵

Recently, patients experiencing rapid bone loss with CPI leading to multiple fractures were reported, raising the question of a potential influence of immune activation on bone metabolism.¹¹⁶

Importantly, rheumatic and/or systemic irAEs may occur across all classes of CPI, most frequently and severely with combination treatments and may be associated with other organ-specific irAEs.

2. Oncologists should be encouraged to consult rheumatologists promptly for assessment when rheumatic musculoskeletal and systemic signs or symptoms are suspected due to immunotherapy, and rheumatologists should provide facilitated access for such patients (LoE 5; LoA 9.4).

The data that are available regarding the process of referral to a rheumatologist suggests that this is not widely done and might lead to delay in diagnosis. In one series, only 4 out of 12 patients experiencing rheumatic irAEs were reviewed by rheumatologists.⁴³ One cohort reported an average of 9.5 ± 9.3 days between the counselling request and the first rheumatologist visit and 2.5 ± 4.4 months from the start of arthralgias to the confirmation of synovitis.⁶⁷ Two other series reported a median of 34 days (range 16–210 days) and 7 days (range 1–57 days) before a rheumatology appointment.^{66, 117} Rheumatic side effects of CPI appear underappreciated, which probably delays proper assessment and treatment. However, as mentioned in the overarching principles, a prompt rheumatological evaluation should support rapid shared treatment decision to relieve patient symptoms, maintain a good quality of life and allow pursuing an effective cancer immunotherapy.

Currently, algorithms for irAEs management are based on the severity/grade of the irAE according to the Common Terminology Criteria for Adverse Events (CTCAE v5.0). The first international guidelines recommended referral to a rheumatologist in the case of severe symptoms not responding to glucocorticoids (grade 3). Subsequently, prompt referral was proposed as soon as the patient experienced moderate pain associated with signs of inflammation (grade 2).^{12–14} While CTCAE grading is routine for oncologists, and a requirement for clinical trials, rheumatologists are less familiar with this grading which do not accurately reflect the spectrum or severity of rheumatic or systemic manifestations (online supplementary table S1). Accordingly, the task force decided not to use the CTCAE grading system to prioritise referral but instead to recommend prompt assessment, ideally before starting glucocorticoids. For this purpose, rheumatologists should be encouraged to offer facilitated access since they may be able to avoid systemic glucocorticoids or use lower dose than oncologists to manage rheumatic toxicities.

3. Metastases, paraneoplastic syndromes and unrelated rheumatic diseases should be considered as a potential differential diagnosis of rheumatic immune-related events. The comprehensive assessment should be focused on documenting evidence of target organ inflammation, and based on history, clinical features, laboratory tests, imaging and/or biopsy (LoE 4; LoA 9.5).

The first part of this statement has previously highlighted the overarching principle of defining the role of the rheumatologist (*overarching principle D*). While delaying the diagnosis of irAEs and its adequate treatment may result in a worse prognosis regarding both CPI adherence and immune-mediated tissue/organ destruction, focusing only on irAEs without considering other differential diagnoses may also be inappropriate. CPIs are commonly administered to patients with advanced cancer, and so new rheumatic/musculoskeletal symptoms must raise suspicion of cancer progression, as well as the lack of improvement of inflammatory arthritis with glucocorticoids (ie, possibility of metastases or paraneoplastic syndrome).^{118, 119} Advanced imaging, such as CT scan, MRI, bone scintigraphy or positron emission tomography-CT, may be helpful in arriving at such a

diagnosis. The diagnosis of irAEs versus metastasis may become even more challenging as non-malignant resorptive lesions have recently been described, which can mimic metastases.¹¹⁶ Pulmonary sarcoidosis-like lesions may also be first considered as tumour progression.

Immunological toxicities may also manifest as paraneoplastic syndromes. Current literature covers mainly paraneoplastic neurological syndromes with few published data regarding paraneoplastic rheumatic syndromes.^{120, 121} However, based on the clinical experience of task force members, the group agreed to include paraneoplastic syndromes in the differential diagnosis of rheumatic irAEs to inform clinicians that they may encounter newly and not pre-existing paraneoplastic syndromes following CPI therapy, notably hypertrophic osteoarthropathy. RS3PE and dermatomyositis were also reported, either as paraneoplastic syndromes or induced by CPI therapy, but one may not be able to make the distinction when appearing under CPI therapy.

The term ‘unrelated rheumatic diseases’ covers manifestations for which the causal link with cancer immunotherapy is not obvious, such as shoulder tendinitis, lateral epicondylitis, non-inflammatory back pain or complex regional pain syndrome. The task force agrees that it may be difficult to establish when a specific rheumatic feature can be considered related or unrelated to the administration of CPI. Using the adverse drug reaction probability score (Naranjo scale) may help to assess the causal link with CPI therapy.

The task force proposes that the key objective of the diagnostic work-up is to document evidence of target organ inflammation. By adopting the term target organ inflammation, the task force wants to emphasise that priority for the supervising rheumatologist is not only to search for joint inflammation but also to document evidence of any organ inflammation according to the symptoms presented (muscle, fascia, vessels, heart, lung, skin, endocrine glands, salivary glands, etc), either clinically or preferably by using appropriate laboratory tests, imaging and tissue biopsy.

Tissue diagnosis should be decided on a case-by-case basis, based on the type and severity of rheumatic irAE, when other supportive information would not be sufficient to make a clinical decision in terms of therapy. Notably, histopathological data may be frequently indicated in patients presenting with vasculitis, sarcoidosis and myositis, but should not interfere with starting treatment, particularly with myositis or patients presenting with life-threatening irAE. On the other hand, synovial biopsies will not change the acute management of inflammatory arthritis. They may provide insights into targeted therapies with glucocorticoid saving approaches, but are not recommended for daily practice.

4. In case of inefficacy of symptomatic treatment and depending on the disease severity, local and/or systemic glucocorticoids should be considered for immune-related rheumatic and systemic symptoms. Dose regimen and route of administration should be decided according to the clinical entity and activity. When improvement is achieved, systemic glucocorticoids should be tapered to the lowest effective dose to control the symptoms (LoE 4; LoA 9.4).

In the absence of contraindications, symptomatic treatment including non-steroidal anti-inflammatory drugs and/or analgesics should be the initial treatment for mild-to-moderate rheumatic manifestations. There are no data on the efficacy of symptomatic therapies in the context of systemic manifestations. An anti-inflammatory effect of these drugs can be expected within several hours or a few days. Additionally, intra-articular glucocorticoids should be considered in the context of monoarthritis

or oligoarthritis, combined with an analysis of the synovial fluid, whenever possible, to rule out differential diagnoses such as infection, osteoarthritis or crystals.^{40–42 48 66 67 75–77 122–126} If symptomatic treatment is insufficient and tissue inflammation is still evident, systemic glucocorticoids should be considered for both immune-related rheumatic and systemic symptoms. Overall, systemic glucocorticoids were used for 224/296 patients (76%) with arthritis^{39–44 46 48 51 52 65–74 76–78 83 116 122 123 125–143} with a median dosage of 20 mg/day, for 37/65 patients (57%) with sicca syndrome^{39 40 65 66 85–89} with a median dosage of 40 mg/day (16 patients for sicca symptoms, 15 patients for systemic manifestations or associated arthritis, 6 patients for sicca symptoms and associated other irAE), for 22/29 patients (76%) with vasculitis^{48 84 103–106 108–112 144–150} with a median initial dosage of 60 mg/day, for 15/33 patients (45%) with sarcoidosis^{39 44 91–93 151–158} with a median initial dosage of 55 mg/day, for 7/7 patients (100%) with scleroderma^{48 94–96} with an initial dosage of 1 mg/kg/day and for 4/13 patients (31%) with lupus.^{98–101} Subacute cutaneous lupus was mainly treated with topical steroids.⁹⁷ Treatment of patients with myositis is reported in a separate statement (*point to consider* 8).

So far, there are reassuring data regarding the use of glucocorticoids for irAE management.^{159 160} For rheumatic irAEs, patients receiving glucocorticoids equivalent to 10 mg/day of prednisone for 6 weeks concurrent to anti-PD1 therapy had a similar antitumour response.⁷⁷ However, recent preclinical data point out that glucocorticoids markedly impair the activation and the killing ability of tumour-infiltrating lymphocytes.¹⁶¹ Because of concerns of glucocorticoids on antitumour responses, the task force did not recommend using methylprednisolone pulses or high-dose oral glucocorticoids in the absence of life-threatening complications and myositis, even in severe presentations, and favoured the concept of glucocorticoid sparing where rheumatologists have extensive experience with alternative options. Furthermore, the task force members recommended tapering glucocorticoids to the lowest effective dose within weeks or as soon as improvement is achieved was desirable. The objective of reaching a dose less than or equal to 10 mg/day of equivalent prednisone was considered as an acceptable target dose. This target dose as maintenance therapy is based on current preclinical and retrospective clinical data,^{161–163} and higher than the one recommended for the main classical RMDs (online supplementary table S2).

5. csDMARD should be considered in patients with insufficient response to acceptable dose of glucocorticoids or requiring glucocorticoid-sparing (LoE 4; LoA 9).

In case of active rheumatic irAE requiring dose of glucocorticoids higher than 10 mg/day of equivalent prednisone, conventional synthetic disease-modifying antirheumatic drug (csDMARD) should be considered. Several csDMARDs have been used as second-line therapy in the case of an insufficient response to glucocorticoids or for use as steroid sparing agents. So far, no specific biological disease-modifying antirheumatic drug has proven superiority. For the various types of arthritis in cases reported, methotrexate was the most frequently drug prescribed, followed by hydroxychloroquine then sulfasalazine, either as monotherapy or in combination.^{39–41 43 44 48 51 52 65–67 69 70 75–78 122 123 125 128 131 132 134 140–142 164 165} Of note, no safety issues were described regarding long-term use of methotrexate associated with CPI in a few patients, with a median follow-up of over 1 year.⁶⁷ It is noteworthy that a higher proportion of hypersensitivity reactions were reported with sulfasalazine in the context of CPI-induced inflammatory arthritis, suggesting

caution to its use in those situations.¹⁶⁵ One case series reported the initiation of hydroxychloroquine prior to glucocorticoids, limiting glucocorticoid exposure, which would deserve further evaluation.¹⁶⁶ The use of csDMARDs has not been described for patients with CPI-induced sicca syndrome. Two patients received hydroxychloroquine and one the combination of hydroxychloroquine and methotrexate for cutaneous leucocytoclastic vasculitis.¹⁰³ One patient with granulomatosis with polyangiitis was treated with oral cyclophosphamide.¹⁰⁵ For other systemic manifestations, hydroxychloroquine was safely prescribed for patients with CPI-induced lupus and scleroderma and in one patient with sarcoidosis.^{48 95–100 157} Four patients with scleroderma-like syndromes received mycophenolate mofetil.^{48 94 96} Among them, two also received intravenous immunoglobulin. Finally, two patients with neurosarcoidosis were successfully treated with methotrexate after an infusion reaction to infliximab.^{92 93}

6. For patients experiencing severe rheumatic and systemic immune-related adverse events or with insufficient response to csDMARD, bDMARD may be considered, with TNF or IL-6 inhibitors being the preferred options for inflammatory arthritis (LoE 4; LoA 8.8).

Gastroenterologists have safely and successfully administered infliximab for patients with severe CPI-induced colitis who had an insufficient response to glucocorticoids.^{167 168} Based on these data, tumour necrosis factor (TNF) inhibitors (infliximab prevailing on etanercept and adalimumab) have been reported for severe and refractory inflammatory arthritis.^{65 66 116 123 138} However, while patients experiencing colitis required one or two infliximab infusions, patients with arthritis may require long-term administration of TNF inhibitors, which is an important difference of unclear clinical significance at this time. A recent study reported that antitumour responses were not adversely affected in patients treated with TNF inhibitors, with a median follow-up of 9 months, but further data are needed.¹⁶⁹ Preclinical data support the use of TNF inhibitors, since infliximab only had a minor influence on T-cell activation and the killing ability of tumour-infiltrating lymphocytes, whereas even low doses of glucocorticoids markedly impaired this antitumour activity.¹⁶¹ Furthermore, a synergistic effect of TNF inhibitors with CPI has been demonstrated in mouse models.^{170 171} A phase I investigator-initiated trial (TICIMEL, NCT03293784) is currently testing the safety of this combined approach (double immunotherapy plus TNF inhibitor) in patients with melanoma. Results of this study will likely inform the management of rheumatic irAEs. The use of infliximab was also reported in two patients with neurosarcoidosis.^{92 93}

There are also several observations in patients with CPI-induced inflammatory arthritis treated with tocilizumab.^{48 52 68 83} Notably, one patient responded to tocilizumab after infliximab failure.⁵²

Regarding interleukin 17 blockade, the use of secukinumab has been reported in a patient with mismatch-repair-deficient metastatic colon cancer and a previous history of Crohn's disease who experienced colitis, severe psoriatic rash and arthralgia.¹⁷² While providing a dramatic relief of the immune-related skin, rheumatic and gastrointestinal side effects, subsequent tumour progression was observed. A second recent publication described the complete resolution of pembrolizumab-induced psoriasiform eruption with secukinumab in a patient with melanoma, without impact on tumour response.¹⁷³ Due to limited data and concerns about interleukin 17 inhibition on CPI efficacy, the task force agreed not to recommend interleukin 17 blockade for inflammatory arthritis.

For mechanistic reasons, abatacept should also not be considered for the treatment of CPI-induced rheumatic and systemic diseases, owing to the hypothetical risk of antagonising antitumour responses of CPI. However, one may consider its use in cases of life-threatening conditions, as discussed in the statement for myositis (*point to consider* 8).

One patient with neuro-Sjögren's syndrome was successfully treated with rituximab after intravenous pulses of methylprednisolone, immunoglobulins and one dose of cyclophosphamide.⁸⁷ Rituximab was also used in one patient with acral vasculitis without improvement and the need of surgical amputation.¹⁰⁵

7. The decision to hold or to continue the cancer immunotherapy should be based on the severity of rheumatic immune-related adverse events, the extent of required immunosuppressive regimen, the tumour response and its duration, as well as the future oncology treatment plan, in a shared decision with the patient (LoE 5; LoA 9.4).

Currently, decisions regarding CPI and immunosuppressive regimens vary from institution to institution according to local practice, with no randomised trials to provide evidence in choosing between holding CPI and/or introducing an immunosuppressive regimen. Overall, the SLR revealed that CPIs were discontinued in 25% of patients experiencing inflammatory arthritis, 61% of patients with sicca syndrome (a discontinuation of CPI often due to another associated irAE), 80% of patients with vasculitis, 64% of patients with sarcoidosis, 75% of patients with scleroderma and 78% of patients with lupus. It is noteworthy that several studies reported ongoing clinical benefit in patients who discontinue their cancer immunotherapy for irAEs.^{7 174 175} Well-designed prospective trials will be required help to clarify the optimal immunosuppressive regimens.

8. Myositis may be a severe condition. Immunotherapy withdrawal needs to be discussed. In the presence of life-threatening manifestations (bulbar symptoms (dysphagia, dysarthria, dysphonia), dyspnoea and myocarditis), high dose of glucocorticoids, IVIg and/or plasma exchange should be considered; immunotherapy withdrawal is always necessary (LoE 4; LoA 8.9).

Myositis belongs to the spectrum of potentially fatal toxicity associated with CPI, since it is frequently associated with myocarditis and/or myasthenia gravis.^{176–178} Notably, it generally occurs very early after CPI initiation, often within the first month of treatment (median exposure time of 25 days, IQR 25–45 days). Proximal weakness and myalgia are the major symptoms, which can mimic a PMR-like condition.⁸¹ Therefore, a high awareness for myositis is needed among rheumatologists with measurement of creatine kinase (CK) since increased CKs are seen in the majority of patients with myositis (median of 2650 IU/L, ranging from 335 to 20 270 IU/L).^{48 77 81–83 100 117 179–207} Of note, CK levels are usually within the normal range in patients presenting with myalgia.^{42 83} Ptosis and diplopia are also commonly reported and may be related to associated myasthenia gravis.^{81 82 100 117 184 185 189 191 192 195 197 198 204 205 208–210} Of note, some patients present with dropped head syndrome.^{82 198 211} Importantly, one should search for the presence of life-threatening manifestations, including dyspnoea, palpitations, chest pain or syncope, which should alert on a possible concurrent myocarditis.^{39 48 77 81–83 117 177–179 182 184 185 187 190 193 200 207 211–215} Of note, an increased risk of death in patients experiencing CPI-related myositis has been observed compared with patients with idiopathic inflammatory myositis (around 20% vs less than 10%).^{177 216} This increased mortality rate seems to be related to the development of myocarditis. While there is no standardised assessment of myocarditis in large series of idiopathic

inflammatory myopathy, signs of myocardial inflammation cardiac has been reported on magnetic resonance tomography in more than 60% of such patients,²¹⁷ which argue that myocarditis belong to the myositis clinical spectrum and does not represent a different concomitant irAE. Therefore, cardiac evaluation must be systematic for any patient with myositis or suspected myositis. It includes cardiac troponin (troponin T is less specific than troponin I in case of associated skeletal muscle diseases) and electrocardiography. In case of clinical syndrome associated with myocarditis and/or increase cardiac troponin level and/or electrocardiography, a cardiac MRI is necessary.²¹⁸ Of note, normal cardiac enzyme cannot always rule out the possibility of myocarditis. Furthermore, the presence of bulbar symptoms (dysphagia, dysarthria, dysphonia) and/or respiratory failure may be related to myositis or associated myasthenia gravis encountered in 12.5% of patients (57/454 cases reported).^{82 83 100 177 178 186 188 189 192 195 203 204 208 211 212} Of note, the majority of patients will not have a typical skin rash of dermatomyositis, only reported in a few patients.^{199 201 219}

Myositis-associated autoantibodies are mostly negative, though cases with positive ANA, antistriated antibodies, anti-PM/ Scl, anti-SM, anti-TIF1 gamma, anti-PL-7, anti-PL12, anti-Jo1 or anti-SRP have been reported.^{77 83 117 184} Electrodiagnostic studies usually reveal myopathic pattern with musculature enhancement may be observed on MRI. Biopsy is often performed and confirms muscle damage with variable degrees of inflammatory and necrotic changes.⁸¹ Of interest, fasciitis is also increasingly reported clinically and seen on MRI findings.^{40 76 220–224}

Prompt recognition and early management of myositis is imperative. Discontinuation or at least interruption of CPI was reported in more than 85% of patients and is mandatory in the presence of dyspnoea, bulbar symptoms, severe muscle weakness and/or myocarditis. High-dose systemic glucocorticoids are the first-line treatment, usually 1–2 mg/kg/day (median dosage 70 mg/day). Ten per cent of reported patients received intravenous pulses of methylprednisolone. Up to 20% of patients also received intravenous immunoglobulins,^{39 48 77 81–83 181 183–185 188 189 191 198 201 207–209 211 212 220 225 226} and plasma exchanges were performed in around 10% of patients.^{48 81 82 117 150 184 188 189 191 197 204 205 226} As second-line therapy, several csDMARDs have been used: mycophenolate mofetil,^{77 209 225} methotrexate,^{39 44 77 81} azathioprine in one patient but stopped for pancreatitis⁷⁷ and hydroxychloroquine in one patient.⁷⁶ Six patients have been treated with infliximab, but only one successfully.^{82 182 184 225} Importantly, a recent publication reported the resolution of a severe glucocorticoid-refractory myocarditis with abatacept, received after plasma exchanges was unsuccessful.²²⁷ Another T-cell directed therapy, alemtuzumab, has been successfully used in a patient with glucocorticoid-refractory myocarditis.²²⁸ The task force agreed that further evaluation is warranted, most notably on the impact on tumour response; however, due to the lack of effective therapy and the high mortality rate of myositis complicated with myocarditis or severe respiratory failure, one may consider their use as rescue therapy in refractory situations.

9. A pre-existing autoimmune rheumatic and/or systemic disease should not preclude the use of cancer immunotherapy. Baseline immunosuppressive regimen should be kept at the lowest dose possible (for glucocorticoids, below 10 mg prednisone per day if possible). However, many patients may have a flare of the underlying condition and/or immune-related adverse events, requiring the use of glucocorticoids and/or DMARDs (LoE 4; LoA 9).

Patients with pre-existing inflammatory or autoimmune disease have been largely excluded from clinical trials due to the theoretical risk of worsening autoimmune manifestations. However, there are several series reporting on CPI safety in such patients, with either anti-CTLA-4^{229–231} or anti-PD-(L)1.^{77 232–235} Together, a flare of the pre-existing inflammatory or autoimmune disease was observed in half of patients with RA (47/86 patients), PsA (4/8 patients) and myositis (1/2 patients), 64% of patients with PMR (16/25 patients), 31% of patients with SA (4/13 patients) and patients with systemic lupus erythematosus (4/13 patients), 43% of patients with Sjögren's syndrome (3/7 patients), 25% of patients with systemic sclerosis (2/8 patients) and 20% of patients with sarcoidosis (3/15 patients), but less than 10% had to stop CPI therapy during a flare. The patient with pre-existing giant cell arteritis experienced a relapse. There was no flare reported for the few patients with pre-existing seronegative arthritis (n=4), other vasculitis (n=4) and Behçet's disease (n=1). Furthermore, 18 of 104 patients (17%) experienced other irAEs, mainly colitis (n=12), hypophysitis (n=3) and thyroiditis (n=3). One patient with RA developed myositis requiring high dose of glucocorticoids and intravenous immunoglobulins, and another patient with RA developed Sjögren's syndrome with autoantibodies (ANA 1/1280, anti-SSA and SSB). Overall, CPI was discontinued in 8% of patients with pre-existing autoimmune disease due to other irAEs, unrelated to their pre-existing autoimmune disease. In a recent case series of 112 patients with pre-existing autoimmune diseases treated with CPI, a flare of pre-existing autoimmune disease or another irAE occurred in 71% of the patients (47% has a flare of their pre-existing disease and 43% had another irAE).²³⁶ Thus, the occurrence of a flare/irAE was frequent but mostly manageable without CPI discontinuation in 79% of the patients.

In these case series, most flares and irAEs were managed with glucocorticoids, with the need of csDMARDs in some patients, usually hydroxychloroquine, methotrexate, sulfasalazine, either in monotherapy or in combination. The need for TNF inhibitors was only reported in patients with flares of their inflammatory bowel disease flares and in two cases of new-onset colitis. Based on these data, the task force agreed that CPI therapy in patients with pre-existing autoimmune rheumatic and systemic disease was not contraindicated, provided that the patient is well-informed and closely monitored. No preventive treatment is needed. Importantly, this remains a shared decision between the oncologist, rheumatologist and the patient, and whether CPI will be used in a metastatic or adjuvant setting is a major aspect to be considered.

Regarding baseline immunosuppressive regimen, recent preclinical and clinical data highlighted the deleterious impact of baseline glucocorticoids on CPI efficacy, when used at a dosage of greater than 10 mg/day.^{163 237} However, this was in patients treated with steroids for their cancer or cancer-related symptoms and not for autoimmune symptoms. Accordingly, the task force agreed on recommending the lowest immunosuppressive regimen possible at the start of CPI therapy. However, future data on prophylactic TNF inhibition and a possible synergistic effect of TNF inhibitors and CPI, reported in a mouse model and currently evaluated in patients, may challenge this statement over time.¹⁷¹

10. Before initiation of cancer immunotherapy, there is no indication to test every patient for the presence of autoantibodies. In the case of unexplained rheumatic, musculoskeletal or systemic symptoms, a complete rheumatologic assessment should be performed (LoE 5; LoA 9).

Analysis of pretreatment and post-treatment sera of anti-CTLA4-treated patients with melanoma revealed that for

most autoantibodies, including RA-associated antibodies, post-treatment titres increased only marginally and were not associated with the occurrence of irAEs.²³⁸ Similarly, the presence of ANA in serum collected prior to initiating CPI therapy was not found to predict the development of irAEs, except for colitis.^{239 240} One study reported divergent data, with pre-existing antibodies independently associated with the occurrence of irAEs, but also with clinical benefits on advanced non-small cell lung cancer.²⁴¹ Notably, skin reactions were more frequent among patients with pre-existing RF.

Since autoantibodies are not found in the majority of patients experiencing CPI-induced rheumatic and systemic disease, there is no indication to test every patient at baseline. Of note, the presence of ACPAs has been detected in serum samples obtained prior to CPI therapy in few patients who experienced RA and were asymptomatic before the start of CPI.⁷⁸ But this situation might be rare, and the detection of autoantibodies in an asymptomatic patient would not preclude the start of CPI therapy. However, there is the particular situation of patients with thymoma who develop CPI-induced myositis and who all have anti-acetylcholine receptor and antistriated muscle antibodies detected in serum sample obtained prior to CPI therapy.¹⁷⁹ Accordingly, as myositis may evolve into a severe irAE, testing for the presence of these antibodies before starting CPI in a patient with thymoma is recommended to identify a high risk of myositis.

CONCLUSION

These points to consider provide the basis of an European League Against Rheumatism consensus on the diagnosis and the management of rheumatic and systemic irAEs which represent a new and rapidly expanding field. The task force aimed to raise awareness and to assist rheumatologists to improve the diagnosis and the management of patients with irAEs. In contrast to other irAEs, rheumatic irAEs frequently persist over time, specifically inflammatory arthritis was persistent in almost 50% at most recent follow-up with a median of 9 months in a recent study.¹⁶⁹ Thus, irAEs represent a new spectrum of RMDs that rheumatologists should familiarise with. Interestingly, many of these manifestations, either frequent (arthritis, myositis, sicca syndrome) or more exceptionally reported (scleroderma, lupus) are also characteristics of graft versus host disease.²⁴² Early consultation and strong collaboration between the referring oncologist, the treating rheumatologist, potentially other organ specialists and the patient are all required for optimal irAEs management.

These statements, being based almost entirely on low levels of evidence and on experts opinion, will undoubtedly require updating over the next few years, as new data emerge. Indeed, we expect that future oncological data will likely impact our irAEs therapeutic strategy. We also anticipate a better understanding of irAEs mechanisms and pathophysiology. Finally, multicentre collaborative efforts, prospective registries and randomised trials will help to define the optimal treatment strategies to relieve patient symptoms without altering oncological outcomes.

RESEARCH AGENDA

- To better understand pathophysiology of rheumatic and systemic irAEs.
- To develop information on rheumatic and systemic irAEs for patients starting cancer immunotherapy.
- To define optimal glucocorticoid dose and duration according to the type of rheumatic and/or systemic irAE.
- To assess the effect of different immunomodulatory/ immunosuppressive agents already given before the start

of CPI therapy in pre-existing RMDs on the outcome of immunotherapy.

- To assess the effect of different immunomodulatory/immunosuppressive agents administered for de novo rheumatic and systemic irAEs on the outcome of immunotherapy, using prospective registries.
- To develop well-designed trials on irAE management.
- To assess long-term evolution of rheumatic and systemic irAEs.
- To search for predictive factors for rheumatic and systemic irAEs.
- To revise CTCAE grading of rheumatic and systemic irAEs.
- To obtain insights on the initiation and propagation of classical rheumatic diseases.

Author affiliations

- ¹Rheumatology, University Hospital of Bordeaux, Bordeaux, France
²Division of Rheumatology, University Hospital of Geneva, Geneva, Switzerland
³Rheumatology, Johns Hopkins University, Baltimore, Maryland, USA
⁴Rheumatology, Haga Hospital, Den Haag, The Netherlands
⁵Department of Medicine V, Division of Rheumatology, University Hospital Centre, Mannheim, Germany
⁶Department of Internal Medicine IV, Division of Rheumatology and Clinical Immunology, University of Munich, Munich, Germany
⁷Institute of Infection and Immunity, Cardiff University School of Medicine, Cardiff, UK
⁸Rheumatology, University Hospital Heidelberg, Heidelberg, Germany
⁹Rheumatology and Clinical Immunology, Utrecht Medical Center, Utrecht, The Netherlands
¹⁰Academic Department of Rheumatology, King's College London, London, UK
¹¹Internal Medicine and Clinical Immunology, Hôpital Bicêtre, Le Kremlin-Bicêtre, France
¹²Rheumatology, University Hospital of Strasbourg, Strasbourg, France
¹³Internal Medicine and Clinical Immunology, Sorbonne Université, Pitié-Salpêtrière University Hospital, Paris, France
¹⁴EULAR PARE Patient Research Partners, Amsterdam, The Netherlands
¹⁵Aarhus University Hospital, Aarhus, Denmark
¹⁶Rheumatology, The University of British Columbia, Vancouver, British Columbia, Canada
¹⁷Drug Development, Gustave Roussy Cancer Center, Villejuif, France
¹⁸Royal Marsden Hospital NHS Foundation Trust, London, UK
¹⁹The Netherlands Cancer Institute, Amsterdam, Noord-Holland, The Netherlands
²⁰Immunology and Rheumatology, Cleveland Clinic, Cleveland, Ohio, USA
²¹Rheumatology, Assistance Publique-Hôpitaux de Paris (AP-HP), Hôpitaux universitaires Paris-Sud – Hôpital Bicêtre, Le Kremlin Bicêtre, France
²²Université Paris-Sud, Center for Immunology of Viral Infections and Auto-immune Diseases (IMVA), Institut pour la Santé et la Recherche Médicale (INSERM) UMR 1184, Université Paris-Saclay, Le Kremlin Bicêtre, France

Twitter Marie Kostine @MarieKostine

Acknowledgements The task force gratefully thank the librarian Catherine Weill (Bibliothèque Inter-universitaire de Santé, Paris, France) for her contribution to the systematic literature search, and European League Against Rheumatism for financial and logistical support.

Contributors The paper was drafted by MK, XM, AF and TS and all authors contributed with specific comments and revisions to the paper.

Funding This study was funded by European League Against Rheumatism (grant no: CL1106).

Competing interests MK: honoraria from Abbvie, BMS, Lilly, Novartis, Pfizer; TRDJR: grants from AbbVie, Takeda, UCB, Janssen, GSK and honoraria from AbbVie, Pfizer, Takeda, Lilly, Medimmune, Novartis, GSK, BMS, AstraZeneca, Janssen; XM: honoraria from BMS; COB: grants and honoraria from BMS and honoraria from Genetech/Roche, Sanofi/Regeneron; OL: grant from Gilead and honoraria from BMS, MSD, AstraZeneca, Janssen; JLe: grants from Novartis, Pfizer and honoraria from AbbVie, AstraZeneca, BMS, Celgene, Hospira, Janssen-Cilag, LEO Pharma, Lilly, MSD, Novartis, Pfizer, Roche, Sanofi, UCB; AM: grants from BMS, Merus and honoraria from Merck Serono, Lytix, BMS, Symphogen, Amgen, AZ/Medimmune, Servier, Gritstone, Pierre Fabre, Eisai, Sanofi; TS: honoraria from Pfizer, Lilly, Novartis, BMS, AbbVie, Sanofi; EHC: grants from Biogen, grants and honoraria from Amgen, Bio-Cancer, Roche, UCB, Pfizer, honoraria from Chugai Pharma, AbbVie, BMS, Celgene, Eli Lilly, Janssen, ObsEva, Regeneron, Sanofi, SynAct Pharma, Tonix, Gilead; LT: honoraria from Novartis; KB: grants from AbbVie, Novartis, Rheumaliga Baden-Württemberg and honoraria from MSD, AbbVie, BMS, Janssen, Lilly, Mundipharma, Novartis, Pfizer,

Roche, UCB; KV: speaker fees from BMS; JLa: grants from Aveo and Pharmacyclics, grants and honoraria from Achilles Therapeutics, MSD, Nektar, Novartis, Pierre Fabre, Pfizer, Roche/Genetech and Immunocore, honoraria from AstraZeneca, Boston Biomedical, BMS, Eisai, EUSA Pharma, GSK, Ipsen, Imugen, Incyte, iOnctura, Kymab, Merck Serono, Secarna, Vitaccess and Covance; JBAGH: grants from BMS, Novartis and advisory boards and/or lectures for MSD, BMS, Roche, Novartis; J-EG: grants from BMS, UCB, Pfizer, Sanofi and honoraria from BMS, Lilly, Pfizer, Sanofi-Genzyme, UCB.

Patient and public involvement Patients and/or the public were involved in the design, conduct, reporting or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Marie Kostine <http://orcid.org/0000-0002-6729-6200>
 Axel Finckh <http://orcid.org/0000-0002-1210-4347>
 Jacques-Eric Gottenberg <http://orcid.org/0000-0002-9469-946X>
 Yves Allenbach <http://orcid.org/0000-0002-3185-7993>
 Leonard H Calabrese <http://orcid.org/0000-0002-1789-4923>

REFERENCES

- 1 Wolchok J. Putting the immunologic brakes on cancer. *Cell* 2018;175:1452–4.
- 2 Topalian SL, Drake CG, Pardoll DM. Immune checkpoint blockade: a common denominator approach to cancer therapy. *Cancer Cell* 2015;27:450–61.
- 3 Wei SC, Duffy CR, Allison JP. Fundamental mechanisms of immune checkpoint blockade therapy. *Cancer Discov* 2018;8:1069–86.
- 4 Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 2010;363:711–23.
- 5 Reck M, Rodríguez-Abreu D, Robinson AG, et al. Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N Engl J Med* 2016;375:1823–33.
- 6 Motzer RJ, Escudier B, McDermott DF, et al. Nivolumab versus everolimus in advanced renal-cell carcinoma. *N Engl J Med* 2015;373:1803–13.
- 7 Hodi FS, Chiarion-Sileni V, Gonzalez R, et al. Nivolumab plus ipilimumab or nivolumab alone versus ipilimumab alone in advanced melanoma (CheckMate 067): 4-year outcomes of a multicentre, randomised, phase 3 trial. *Lancet Oncol* 2018;19:1480–92.
- 8 Hirsch L, Zitvogel L, Eggermont A, et al. PD-Loma: a cancer entity with a shared sensitivity to the PD-1/PD-L1 pathway blockade. *Br J Cancer* 2019;120:3–5.
- 9 Postow MA, Sidlow R, Hellmann MD. Immune-related adverse events associated with immune checkpoint blockade. *N Engl J Med* 2018;378:158–68.
- 10 Calabrese LH, Calabrese C, Cappelli LC. Rheumatic immune-related adverse events from cancer immunotherapy. *Nat Rev Rheumatol* 2018;14:569–79.
- 11 Kostine M, Cappelli LC, Calabrese C, et al. Addressing immune-related adverse events of cancer immunotherapy: how prepared are rheumatologists? *Ann Rheum Dis* 2019;78:860–2.
- 12 Haanen JBAG, Carbone F, Robert C, et al. Management of toxicities from immunotherapy: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2017;28:iv119–42.
- 13 Puzanov I, Diab A, Abdallah K, et al. Managing toxicities associated with immune checkpoint inhibitors: consensus recommendations from the Society for Immunotherapy of Cancer (SITC) toxicity management Working group. *J Immunother Cancer* 2017;5:95.
- 14 Brahmer JR, Lacchetti C, Schneider BJ, et al. Management of immune-related adverse events in patients treated with immune checkpoint inhibitor therapy: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol* 2018;36:1714–68.
- 15 Thompson JA, Schneider BJ, Brahmer J, et al. Management of Immunotherapy-Related toxicities, version 1.2019. *J Natl Compr Canc Netw* 2019;17:255–89.
- 16 van der Heijde D, Aletaha D, Carmona L, et al. 2014 update of the EULAR standardised operating procedures for EULAR-endorsed recommendations. *Ann Rheum Dis* 2015;74:8–13.
- 17 OCEBM Levels of Evidence Working Group. Oxford centre for evidence-based medicine – levels of evidence (March 2009). Available: <https://www.cebm.net/2009/06/oxford-centre-evidence-based-medicine-levels-evidence-march-2009/> [Accessed 20 Jan 2019].

- 18 Antonia SJ, Villegas A, Daniel D, *et al.* Durvalumab after chemoradiotherapy in stage III non-small-cell lung cancer. *N Engl J Med* 2017;377:1919–29.
- 19 Bellmunt J, de Wit R, Vaughn DJ, *et al.* Pembrolizumab as second-line therapy for advanced urothelial carcinoma. *N Engl J Med* 2017;376:1015–26.
- 20 Borghaei H, Paz-Ares L, Horn L, *et al.* Nivolumab versus docetaxel in advanced Nonsquamous non-small-cell lung cancer. *N Engl J Med* 2015;373:1627–39.
- 21 Brahmer J, Reckamp KL, Baas P, *et al.* Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med* 2015;373:123–35.
- 22 Eggermont AMM, Blank CU, Mandala M, *et al.* Adjuvant pembrolizumab versus placebo in resected stage III melanoma. *N Engl J Med* 2018;378:1789–801.
- 23 Eggermont AMM, Chiarion-Sileni V, Grob J-J, *et al.* Prolonged survival in stage III melanoma with ipilimumab adjuvant therapy. *N Engl J Med* 2016;375:1845–55.
- 24 Gandhi L, Rodríguez-Abreu D, Gadgeel S, *et al.* Pembrolizumab plus chemotherapy in metastatic non-small-cell lung cancer. *N Engl J Med* 2018;378:2078–92.
- 25 Govindan R, Szczesna A, Ahn M-J, *et al.* Phase III trial of ipilimumab combined with paclitaxel and carboplatin in advanced squamous non-small-cell lung cancer. *J Clin Oncol* 2017;35:3449–57.
- 26 Herbst RS, Baas P, Kim D-W, *et al.* Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet* 2016;387:1540–50.
- 27 Kang Y-K, Boku N, Satoh T, *et al.* Nivolumab in patients with advanced gastric or gastro-oesophageal junction cancer refractory to, or intolerant of, at least two previous chemotherapy regimens (ONO-4538-12, ATTRACTION-2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2017;390:2461–71.
- 28 Kwon ED, Drake CG, Scher HI, *et al.* Ipilimumab versus placebo after radiotherapy in patients with metastatic castration-resistant prostate cancer that had progressed after docetaxel chemotherapy (CA184-043): a multicentre, randomised, double-blind, phase 3 trial. *Lancet Oncol* 2014;15:700–12.
- 29 Paz-Ares L, Luft A, Vicente D, *et al.* Pembrolizumab plus chemotherapy for squamous non-small-cell lung cancer. *N Engl J Med* 2018;379:2040–51.
- 30 Powles T, Durán I, van der Heijden MS, *et al.* Atezolizumab versus chemotherapy in patients with platinum-treated locally advanced or metastatic urothelial carcinoma (IMvigor211): a multicentre, open-label, phase 3 randomised controlled trial. *Lancet* 2018;391:748–57.
- 31 Rittmeyer A, Barlesi F, Waterkamp D, *et al.* Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet* 2017;389:255–65.
- 32 Robert C, Long GV, Brady B, *et al.* Nivolumab in previously untreated melanoma without BRAF mutation. *N Engl J Med* 2015;372:320–30.
- 33 Robert C, Schachter J, Long GV, *et al.* Pembrolizumab versus ipilimumab in advanced melanoma. *N Engl J Med* 2015;372:2521–32.
- 34 Robert C, Thomas L, Bondarenko I, *et al.* Ipilimumab plus dacarbazine for previously untreated metastatic melanoma. *N Engl J Med* 2011;364:2517–26.
- 35 Schmid P, Adams S, Rugo HS, *et al.* Atezolizumab and nab-paclitaxel in advanced triple-negative breast cancer. *N Engl J Med* 2018;379:2108–21.
- 36 Socinski MA, Jotte RM, Cappuzzo F, *et al.* Atezolizumab for first-line treatment of metastatic Nonsquamous NSCLC. *N Engl J Med* 2018;378:2288–301.
- 37 Weber J, Mandala M, Del Vecchio M, *et al.* Adjuvant nivolumab versus ipilimumab in resected stage III or IV melanoma. *N Engl J Med* 2017;377:1824–35.
- 38 Weber JS, D'Angelo SP, Minor D, *et al.* Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 2015;16:375–84.
- 39 Le Burel S, Champiat S, Mateus C, *et al.* Prevalence of immune-related systemic adverse events in patients treated with anti-Programmed cell Death 1/anti-Programmed cell Death-Ligand 1 agents: A single-centre pharmacovigilance database analysis. *Eur J Cancer* 2017;82:34–44.
- 40 Narváez J, Juárez-López P, LLuch J, *et al.* Rheumatic immune-related adverse events in patients on anti-PD-1 inhibitors: fasciitis with myositis syndrome as a new complication of immunotherapy. *Autoimmun Rev* 2018;17:1040–5.
- 41 Kostine M, Rouxel L, Barnette T, *et al.* Rheumatic disorders associated with immune checkpoint inhibitors in patients with cancer-clinical aspects and relationship with tumour response: a single-centre prospective cohort study. *Ann Rheum Dis* 2018;77:393–8.
- 42 Zimmer L, Goldinger SM, Hofmann L, *et al.* Neurological, respiratory, musculoskeletal, cardiac and ocular side-effects of anti-PD-1 therapy. *Eur J Cancer* 2016;60:210–25.
- 43 Liew DFL, Leung JLY, Liu B, *et al.* Association of good oncological response to therapy with the development of rheumatic immune-related adverse events following PD-1 inhibitor therapy. *Int J Rheum Dis* 2019;22:297–302.
- 44 Lidar M, Giat E, Garelick D, *et al.* Rheumatic manifestations among cancer patients treated with immune checkpoint inhibitors. *Autoimmun Rev* 2018;17:284–9.
- 45 Lomax AJ, Lim J, Cheng R, *et al.* Immune toxicity with checkpoint inhibition for metastatic melanoma: case series and clinical management. *J Skin Cancer* 2018;2018:9602540.
- 46 Inamo J, Kaneko Y, Takeuchi T. Inflammatory Tenosynovitis and enthesitis induced by immune checkpoint inhibitor treatment. *Clin Rheumatol* 2018;37:1107–10.
- 47 Romero R, Schwartz T, Saxena Beem S, *et al.* Immune Related Adverse Events from Immune Checkpoint Inhibitors: A Retrospective Analysis from 2004-2017 at the University of North Carolina at Chapel Hill [abstract]. *Arthritis Rheumatol* 2018;70.
- 48 Richter MD, Crowson C, Kotschade LA, *et al.* Rheumatic syndromes associated with immune checkpoint inhibitors: a single-center cohort of Sixty-One patients. *Arthritis Rheumatol* 2019;71:468–75.
- 49 Eun Y, Kim I, Kim H, *et al.* Risk factors of immune-related adverse events in patients treated with anti-pd-1 antibody pembrolizumab [abstract]. *Ann Rheum Dis* 2018;77:1205.
- 50 Riudavets M, Barba A, Maroto P, *et al.* Correlation between immune-related adverse events (irAEs) and efficacy in patients with solid tumors treated with immune-checkpoints inhibitors (ICIs). *J Clin Oncol* 2018;36:3064. [abstract].
- 51 Buder-Bakhaya K, Benesova K, Schulz C, *et al.* Characterization of arthralgia induced by PD-1 antibody treatment in patients with metastasized cutaneous malignancies. *Cancer Immunol Immunother* 2018;67:175–82.
- 52 Tucker L, Sacks S, Al-Mossawi H. Inflammatory joint disease triggered by immune checkpoint inhibitors [abstract]. *Ann Rheum Dis* 2017;76:419.
- 53 Mooradian M, Cohen JV, Giobbie-Hurder A, *et al.* Inflammatory arthritis: an under-recognized immune-related adverse effect. *J Clin Oncol* 2017;35:e14565.
- 54 Betof AS, Nipp RD, Giobbie-Hurder A, *et al.* Impact of age on outcomes with immunotherapy for patients with melanoma. *Oncologist* 2017;22:963–71.
- 55 Combe B, Landewe R, Daen CI, *et al.* 2016 update of the EULAR recommendations for the management of early arthritis. *Ann Rheum Dis* 2017;76:948–59.
- 56 Smolen JS, Landewe R, Bijlsma J, *et al.* EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis* 2017;76:960–77.
- 57 Smolen JS, Schöls M, Braun J, *et al.* Treating axial spondyloarthritis and peripheral spondyloarthritis, especially psoriatic arthritis, to target: 2017 update of recommendations by an international Task force. *Ann Rheum Dis* 2018;77:3–17.
- 58 Champiat S, Lambotte O, Barreau E, *et al.* Management of immune checkpoint blockade dysimmune toxicities: a collaborative position paper. *Ann Oncol* 2016;27:559–74.
- 59 Naidoo J, Zhang J, Lipson EJ, *et al.* A multidisciplinary toxicity team for cancer Immunotherapy-Related adverse events. *J Natl Compr Canc Netw* 2019;17:712–20.
- 60 Calabrese L, Mariette X. The evolving role of the rheumatologist in the management of immune-related adverse events (irAEs) caused by cancer immunotherapy. *Ann Rheum Dis* 2018;77:162–4.
- 61 Cappelli L, Grieb S, Orbai A, *et al.* "I Was Prepared for the Other Side Effects; I Wasn't Prepared for This One." A Qualitative Study of the Patients' Experience of Inflammatory Arthritis Due to Immune Checkpoint Inhibitor Therapy for Cancer [abstract]. *Arthritis Rheumatol* 2018;70.
- 62 Abdel-Rahman O, Eltooby M, Oweira H, *et al.* Immune-related musculoskeletal toxicities among cancer patients treated with immune checkpoint inhibitors: a systematic review. *Immunotherapy* 2017;9:1175–83.
- 63 Cappelli LC, Gutierrez AK, Bingham CO, *et al.* Rheumatic and musculoskeletal immune-related adverse events due to immune checkpoint inhibitors: a systematic review of the literature. *Arthritis Care Res* 2017;69:1751–63.
- 64 Benfaremo D, Manfredi L, Luchetti MM, *et al.* Musculoskeletal and rheumatic diseases induced by immune checkpoint inhibitors: a review of the literature. *Curr Drug Saf* 2018;13:150–64.
- 65 Calabrese C, Kirchner E, Kontzias A, *et al.* Rheumatic immune-related adverse events of checkpoint therapy for cancer: case series of a new nosological entity. *RMD Open* 2017;3:e000412.
- 66 Cappelli LC, Gutierrez AK, Baer AN, *et al.* Inflammatory arthritis and sicca syndrome induced by nivolumab and ipilimumab. *Ann Rheum Dis* 2017;76:43–50.
- 67 Leipe J, Christ LA, Arnoldi AP, *et al.* Characteristics and treatment of new-onset arthritis after checkpoint inhibitor therapy. *RMD Open* 2018;4:e000714.
- 68 Kim ST, Tayar J, Trinh VA, *et al.* Successful treatment of arthritis induced by checkpoint inhibitors with tocilizumab: a case series. *Ann Rheum Dis* 2017;76:2061–4.
- 69 Law-Ping-Man S, Martin A, Briens E, *et al.* Psoriasis and psoriatic arthritis induced by nivolumab in a patient with advanced lung cancer. *Rheumatology* 2016;55:2087–9.
- 70 Elosua-González M, Pampín-Franco A, Mazzucchelli-Esteban R, *et al.* A case of de novo palmoplantar psoriasis with psoriatic arthritis and autoimmune hypothyroidism after receiving nivolumab therapy. *Dermatol Online J* 2017;23. [Epub ahead of print: 15 Aug 2017].
- 71 Sapalidis K, Kosmidis C, Michalopoulos N, *et al.* Psoriatic arthritis due to nivolumab administration: a case report and review of the literature. *Respir Med Case Rep* 2018;23:182–7.
- 72 Amini-Adle M, Piperno M, Tordo J, *et al.* Remitting seronegative symmetric synovitis with pitting edema associated with partial melanoma response under anti-CTLA-4 and Anti-Programmed death 1 combination treatment. *Arthritis Rheumatol* 2018;70:1358.
- 73 Gauci M-L, Baroudjian B, Laly P, *et al.* Remitting seronegative symmetrical synovitis with pitting edema (RS3PE) syndrome induced by nivolumab. *Semin Arthritis Rheum* 2017;47:281–7.
- 74 Ngo L, Miller E, Valen P, *et al.* Nivolumab induced remitting seronegative symmetrical synovitis with pitting edema in a patient with melanoma: a case report. *J Med Case Rep* 2018;12:48.

- 75 Chan MMK, Kefford RF, Carlino M, *et al.* Arthritis and tenosynovitis associated with the anti-PD1 antibody pembrolizumab in metastatic melanoma. *J Immunother* 2015;38:37–9.
- 76 Mooradian MJ, Nasrallah M, Gainor JF, *et al.* Musculoskeletal rheumatic complications of immune checkpoint inhibitor therapy: a single center experience. *Semin Arthritis Rheum* 2019;48:1127–32.
- 77 Mitchell EL, Lau PKH, Khoo C, *et al.* Rheumatic immune-related adverse events secondary to anti-programmed death-1 antibodies and preliminary analysis on the impact of corticosteroids on anti-tumour response: a case series. *Eur J Cancer* 2018;105:88–102.
- 78 Belkhir R, Burel SL, Dunogean L, *et al.* Rheumatoid arthritis and polymyalgia rheumatica occurring after immune checkpoint inhibitor treatment. *Ann Rheum Dis* 2017;76:1747–50.
- 79 Calabrese C, Cappelli LC, Kostine M, *et al.* Polymyalgia rheumatica-like syndrome from checkpoint inhibitor therapy: case series and systematic review of the literature. *RMD Open* 2019;5:e000906.
- 80 Kim ST, Bittar M, Kim HJ, *et al.* Recurrent pseudogout after therapy with immune checkpoint inhibitors: a case report with immunoprofiling of synovial fluid at each flare. *J Immunother Cancer* 2019;7:126.
- 81 Touat M, Maisonneuve T, Knauss S, *et al.* Immune checkpoint inhibitor-related myositis and myocarditis in patients with cancer. *Neurology* 2018;91:e985–94.
- 82 Shah M, Tayar JH, Abdel-Wahab N, *et al.* Myositis as an adverse event of immune checkpoint blockade for cancer therapy. *Semin Arthritis Rheum* 2019;48:736–40.
- 83 Moreira A, Loquai C, Pföhler C, *et al.* Myositis and neuromuscular side-effects induced by immune checkpoint inhibitors. *Eur J Cancer* 2019;106:12–23.
- 84 Le Burel S, Champiat S, Routier E, *et al.* Onset of connective tissue disease following anti-PD1/PD-L1 cancer immunotherapy. *Ann Rheum Dis* 2018;77:468–70.
- 85 Takahashi S, Chieko X, Sakai T, *et al.* Nivolumab-induced sialadenitis. *Respirol Case Rep* 2018;6:e00322.
- 86 Teyssonneau D, Cousin S, Italiano A, Gougerot-Sjogren-like syndrome under PD-1 inhibitor treatment. *Ann Oncol* 2017;28:3108.
- 87 Ghosn J, Vicino A, Michielin O, *et al.* A severe case of neuro-Sjögren's syndrome induced by pembrolizumab. *J Immunother Cancer* 2018;6:110.
- 88 Ramos-Casals M, Maria A, Suárez-Almazor ME, *et al.* Sicca/Sjögren's syndrome triggered by PD-1/PD-L1 checkpoint inhibitors. Data from the International ImmunoCancer Registry (ICIR). *Clin Exp Rheumatol* 2019;118:114–22.
- 89 Warner BM, Baer AN, Lipson EJ, *et al.* Sicca syndrome associated with immune checkpoint inhibitor therapy. *Oncologist* 2019;24:1259–69.
- 90 Firwana B, Ravilla R, Raval M, *et al.* Sarcoidosis-like syndrome and lymphadenopathy due to checkpoint inhibitors. *J Oncol Pharm Pract* 2017;23:620–4.
- 91 Lomax AJ, McGuire HM, McNeil C, *et al.* Immunotherapy-induced sarcoidosis in patients with melanoma treated with PD-1 checkpoint inhibitors: case series and immunophenotypic analysis. *Int J Rheum Dis* 2017;20:1277–85.
- 92 Dunn-Pirio AM, Shah S, Eckstein C. Neurosarcoidosis following immune checkpoint inhibition. *Case Rep Oncol* 2018;11:521–6.
- 93 Tan I, Malinzak M, Salama AKS. Delayed onset of neurosarcoidosis after concurrent ipilimumab/nivolumab therapy. *J Immunother Cancer* 2018;6:77.
- 94 Tjarks BJ, Kerkvliet AM, Jassim AD, *et al.* Scleroderma-Like skin changes induced by checkpoint inhibitor therapy. *J Cutan Pathol* 2018;45:615–8.
- 95 Shenoy N, Esplin B, Barbosa N, *et al.* Pembrolizumab induced severe sclerodermoid reaction. *Ann Oncol* 2017;28:432–3.
- 96 Barbosa NS, Wetter DA, Wieland CN, *et al.* Scleroderma induced by pembrolizumab: a case series. *Mayo Clin Proc* 2017;92:1158–63.
- 97 Michot J-M, Fusellier M, Champiat S, *et al.* Drug-induced lupus erythematosus following immunotherapy with anti-programmed death-(ligand) 1. *Ann Rheum Dis* 2019;78:e67.
- 98 Zitouni BN, Arnault J-P, Dadban A, *et al.* Subacute cutaneous lupus erythematosus induced by nivolumab: two case reports and a literature review. *Melanoma Res* 2018;29:212–5.
- 99 Liu RC, Sebaratnam DF, Jakkett L, *et al.* Subacute cutaneous lupus erythematosus induced by nivolumab. *Australas J Dermatol* 2018;59:e152–4.
- 100 de Chabot G, Justeau G, Pinquière F, *et al.* [Unexpected adverse events of immunotherapies in non-small cell lung cancer: About 2 cases]. *Rev Pneumol Clin* 2017;73:326–30.
- 101 Fadel F, El Karoui K, Knebelmann B. Anti-CTLA4 antibody-induced lupus nephritis. *N Engl J Med* 2009;361:211–2.
- 102 de Velasco G, Bermas B, Choueiri TK. Autoimmune arthropathy and uveitis as complications of programmed death 1 inhibitor treatment. *Arthritis Rheumatol* 2016;68:556–7.
- 103 Tomelleri A, Campochiaro C, De Luca G, *et al.* Anti-PD1 therapy-associated cutaneous leucocytoclastic vasculitis: a case series. *Eur J Intern Med* 2018;57:e11–12.
- 104 Comont T, Sibaud V, Mourey L, *et al.* Immune checkpoint inhibitor-related acral vasculitis. *J Immunother Cancer* 2018;6:120.
- 105 Padda A, Schiopu E, Sovich J, *et al.* Ipilimumab induced digital vasculitis. *J Immunother Cancer* 2018;6:12.
- 106 van den Brom RRH, Abdulahad WH, Rutgers A, *et al.* Rapid granulomatosis with polyangiitis induced by immune checkpoint inhibition. *Rheumatology* 2016;55:1143–5.
- 107 Roger A, Groh M, Lorillon G, *et al.* Eosinophilic granulomatosis with polyangiitis (Churg-Strauss) induced by immune checkpoint inhibitors. *Ann Rheum Dis* 2019;78:e82.
- 108 Goldstein BL, Gedmintas L, Todd DJ. Drug-associated polymyalgia rheumatica/giant cell arteritis occurring in two patients after treatment with ipilimumab, an antagonist of ctla-4. *Arthritis Rheumatol* 2014;66:768–9.
- 109 Hid Cadena R, Abdulahad WH, Hospers GAP, *et al.* Checks and balances in autoimmune vasculitis. *Front Immunol* 2018;9:315.
- 110 Micaily I, Chernoff M. An unknown reaction to pembrolizumab: giant cell arteritis. *Ann Oncol* 2017;28:2621–2.
- 111 Kang A, Yuen M, Lee DJ. Nivolumab-induced systemic vasculitis. *JAAD Case Rep* 2018;4:606–8.
- 112 Läubli H, Hench J, Stanczak M, *et al.* Cerebral vasculitis mimicking intracranial metastatic progression of lung cancer during PD-1 blockade. *J Immunother Cancer* 2017;5:46.
- 113 Manusow JS, Khoja L, Pesin N, *et al.* Retinal vasculitis and ocular vitreous metastasis following complete response to PD-1 inhibition in a patient with metastatic cutaneous melanoma. *J Immunother Cancer* 2014;2:41.
- 114 Salem J-E, Manouchehri A, Moey M, *et al.* Cardiovascular toxicities associated with immune checkpoint inhibitors: an observational, retrospective, pharmacovigilance study. *Lancet Oncol* 2018;19:1579–89.
- 115 Sibille A, Alfieri R, Malaise O, *et al.* Granulomatosis with polyangiitis in a patient on programmed death-1 inhibitor for advanced non-small-cell lung cancer. *Front Oncol* 2019;9:478.
- 116 Moseley KF, Naidoo J, Bingham CO, *et al.* Immune-related adverse events with immune checkpoint inhibitors affecting the skeleton: a seminal case series. *J Immunother Cancer* 2018;6:104.
- 117 Liewluck T, Kao JC, Mauermann ML. PD-1 inhibitor-associated myopathies: emerging immune-mediated myopathies. *J Immunother* 2018;41:208–11.
- 118 Albayda J, Bingham CO, Shah AA, *et al.* Metastatic joint involvement or inflammatory arthritis? A conundrum with immune checkpoint inhibitor-related adverse events. *Rheumatology* 2018;57:760–2.
- 119 Schaeferbeke T, Kostine M. Response to: 'Checkpoint inhibitors and arthritis: seeking balance between victories and defeats' by Moura and Moura. *Ann Rheum Dis* 2019;78:e92.
- 120 Manson G, Maria ATJ, Poizeau F, *et al.* Worsening and newly diagnosed paraneoplastic syndromes following anti-PD-1 or anti-PD-L1 immunotherapies, a descriptive study. *J Immunother Cancer* 2019;7:337.
- 121 Shibata C, Kato J, Toda N, *et al.* Paraneoplastic dermatomyositis appearing after nivolumab therapy for gastric cancer: a case report. *J Med Case Rep* 2019;13:168.
- 122 Ruiz-Bañobre J, Pérez-Pampín E, García-González J, *et al.* Development of psoriatic arthritis during nivolumab therapy for metastatic non-small cell lung cancer, clinical outcome analysis and review of the literature. *Lung Cancer* 2017;108:217–21.
- 123 Smith MH, Bass AR. Arthritis after cancer immunotherapy: symptom duration and treatment response. *Arthritis Care Res* 2019;71:362–6.
- 124 Corominas H, Badlissi F, Shmerling RH. Crystal-induced oligoarthritis triggered by pembrolizumab, an immune checkpoint inhibitor. *Joint Bone Spine* 2018;85:647–8.
- 125 Kuswanto WF, MacFarlane LA, Gedmintas L, *et al.* Rheumatologic symptoms in oncologic patients on PD-1 inhibitors. *Semin Arthritis Rheum* 2018;47:907–10.
- 126 Humayun MA, Poole R. A case of multiple immune toxicities from ipilimumab and pembrolizumab treatment. *Hormones* 2016;15:303–6.
- 127 Bernier M, Guillaume C, Leon N, *et al.* Nivolumab causing a polymyalgia rheumatica in a patient with a squamous non-small cell lung cancer. *J Immunother* 2017. doi:10.1097/CJI.000000000000163. [Epub ahead of print: 06 Mar 2017].
- 128 Ennis D, To F, Jamal S. Immune related adverse events (IRAEs) associated with checkpoint inhibitors: 12 cases from a single centre [abstract]. *Ann Rheum Dis* year 2018;77:A506.
- 129 Filetti M, Anselmi E, Macrini S, *et al.* Resolution of remitting seronegative symmetrical synovitis with pitting edema (RS3PE) during nivolumab therapy for non-small cell lung cancer: a case report. *Semin Arthritis Rheum* 2018;48:e17–20.
- 130 Wada N, Uchi H, Furue M. Case of remitting seronegative symmetrical synovitis with pitting edema (RS3PE) syndrome induced by nivolumab in a patient with advanced malignant melanoma. *J Dermatol* 2017;44:e196–7.
- 131 Haikal A, Borba E, Khaja T, *et al.* Nivolumab-induced new-onset seronegative rheumatoid arthritis in a patient with advanced metastatic melanoma: a case report and literature review. *Avicenna J Med* 2018;8:34–6.
- 132 Schmutz J-L. [Psoriasis and psoriatic arthritis induced by nivolumab (Opdivo®)]. *Ann Dermatol Venerol* 2016;143:881–2.
- 133 Kodama S, Kurose K, Mukai T, *et al.* Nivolumab-induced polyarthritis. *BMJ Case Rep* 2017;2017:bcr-2017-223387.
- 134 Godfrey J, Bishop MR, Syed S, *et al.* PD-1 blockade induces remissions in relapsed classical Hodgkin lymphoma following allogeneic hematopoietic stem cell transplantation. *J Immunother Cancer* 2017;5:11.

- 135 Nakamagoe K, Moriyama T, Maruyama H, *et al.* Polymyalgia rheumatica in a melanoma patient due to nivolumab treatment. *J Cancer Res Clin Oncol* 2017;143:1357–8.
- 136 Maniu C, Kobe C, Schlaak M, *et al.* Polymyalgia rheumatica occurring during treatment with ipilimumab. *Eur J Dermatol* 2016;26:513–4.
- 137 Garel B, Kramkimel N, Trouvin A-P, *et al.* Pembrolizumab-induced polymyalgia rheumatica in two patients with metastatic melanoma. *Joint Bone Spine* 2017;84:233–4.
- 138 Mahmoud F, Wilkinson JT, Gizinski A, *et al.* Could knee inflammatory synovitis be induced by pembrolizumab? *J Oncol Pharm Pract* 2018;24:389–92.
- 139 Dasanu CA, Jen T, Skulski R. Late-Onset pericardial tamponade, bilateral pleural effusions and recurrent immune monoarthritis induced by ipilimumab use for metastatic melanoma. *J Oncol Pharm Pract* 2017;23:231–4.
- 140 Spathas N, Economopoulou P, Cheila M, *et al.* Inflammatory arthritis induced by pembrolizumab in a patient with head and neck squamous cell carcinoma. *Front Oncol* 2018;8:409.
- 141 Alperin J, Sarazin J, Fecher L, *et al.* Traditional Disease Modifying Anti-Rheumatic Drugs (tDMARDs), Hydroxychloroquine (HCQ) and/or Sulfasalazine (SSZ), Are Rapidly Effective in Immune Checkpoint Inhibitors-Induced Inflammatory Arthritis [abstract]. *Arthritis Rheumatol* 2017;69.
- 142 Salmon J-H, Lambrecht I, Brochet P, *et al.* A case of arthritis under pembrolizumab. *Joint Bone Spine* 2017;84:243–4.
- 143 Manolios N, Schrieber L. Checkpoint inhibitors and arthritis. *Ann Rheum Dis* 2019;78:e58.
- 144 Tsukamoto J, Monteiro M, Vale S, *et al.* Thromboembolic events related to treatment with checkpoint inhibitors: report of two cases. *Case Rep Oncol* 2018;11:648–53.
- 145 Arellano K, Mosley JC, Moore DC. Case report of ipilimumab-induced diffuse, Nonnecrotizing granulomatous lymphadenitis and granulomatous vasculitis. *J Pharm Pract* 2018;31:227–9.
- 146 Gambichler T, Strutzmann S, Tannappel A, *et al.* Paraneoplastic acral vascular syndrome in a patient with metastatic melanoma under immune checkpoint blockade. *BMC Cancer* 2017;17:327.
- 147 Roy AK, Tathireddy HR, Roy M. Aftermath of induced inflammation: acute periaortitis due to nivolumab therapy. *BMJ Case Rep* 2017;2017:bcr-2017-221852.
- 148 Castillo B, Gibbs J, Brohl AS, *et al.* Checkpoint inhibitor-associated cutaneous small vessel vasculitis. *JAAD Case Rep* 2018;4:675–7.
- 149 Ban B, Crowe J, Graham R. Rheumatology case report immune-related aortitis associated with ipilimumab. *Rheumatol* 2017.
- 150 Kao JC, Liao B, Markovic SN, *et al.* Neurological complications associated with Anti-Programmed death 1 (PD-1) antibodies. *JAMA Neurol* 2017;74:1216–22.
- 151 Berthod G, Lazor R, Letovanec I, *et al.* Pulmonary sarcoid-like granulomatosis induced by ipilimumab. *J Clin Oncol* 2012;30:e156–9.
- 152 Reule RB, North JP. Cutaneous and pulmonary sarcoidosis-like reaction associated with ipilimumab. *J Am Acad Dermatol* 2013;69:e272–3.
- 153 Wilgenhof S, Morlion V, Seghers AC, *et al.* Sarcoidosis in a patient with metastatic melanoma sequentially treated with anti-CTLA-4 monoclonal antibody and selective BRAF inhibitor. *Anticancer Res* 2012;32:1355–9.
- 154 Dimitriou F, Frauchiger AL, Urošević-Maiwald M, *et al.* Sarcoid-like reactions in patients receiving modern melanoma treatment. *Melanoma Res* 2018;28:230–6.
- 155 Montaudou H, Pradelli J, Passeron T, *et al.* Pulmonary sarcoid-like granulomatosis induced by nivolumab. *Br J Dermatol* 2017;176:1060–3.
- 156 Reddy SB, Possick JD, Kluger HM, *et al.* Sarcoidosis following anti-PD-1 and anti-CTLA-4 therapy for metastatic melanoma. *J Immunother* 2017;40:307–11.
- 157 Kim C, Gao J, Shannon VR, *et al.* Systemic sarcoidosis first manifesting in a tattoo in the setting of immune checkpoint inhibition. *BMJ Case Rep* 2016;2016:bcr2016216217.
- 158 Murphy KP, Kennedy MP, Barry JE, *et al.* New-onset mediastinal and central nervous system sarcoidosis in a patient with metastatic melanoma undergoing CTLA4 monoclonal antibody treatment. *Oncol Res Treat* 2014;37:351–3.
- 159 Horvat TZ, Adel NG, Dang T-O, *et al.* Immune-Related adverse events, need for systemic immunosuppression, and effects on survival and time to treatment failure in patients with melanoma treated with ipilimumab at Memorial Sloan Kettering cancer center. *J Clin Oncol* 2015;33:3193–8.
- 160 Garant A, Guilbault C, Ekmekejian T, *et al.* Concomitant use of corticosteroids and immune checkpoint inhibitors in patients with hematologic or solid neoplasms: a systematic review. *Crit Rev Oncol Hematol* 2017;120:86–92.
- 161 Draghi A, Borch TH, Radic HD, *et al.* Differential effects of corticosteroids and anti-TNF on tumor-specific immune responses: implications for the management of irAEs. *Int J Cancer* 2019;145:1408–13.
- 162 Cain DW, Cidlowski JA. Immune regulation by glucocorticoids. *Nat Rev Immunol* 2017;17:233–47.
- 163 Arbour KC, Mezquita L, Long N, *et al.* Impact of baseline steroids on efficacy of programmed cell death-1 and programmed Death-Ligand 1 blockade in patients with non-small-cell lung cancer. *J Clin Oncol* 2018;36:2872–8.
- 164 Ornstein MC, Calabrese C, Wood LS, *et al.* Myalgia and arthralgia immune-related adverse events (irAEs) in patients with genitourinary malignancies treated with immune checkpoint inhibitors. *Clin Genitourin Cancer* 2019;17:177–82.
- 165 Ford M, Sahbudin I, Filer A, *et al.* High proportion of drug hypersensitivity reactions to sulfasalazine following its use in anti-PD-1-associated inflammatory arthritis. *Rheumatology* 2018;57:2244–6.
- 166 Roberts J, Smylie M, Walker J, *et al.* Hydroxychloroquine is a safe and effective steroid-sparing agent for immune checkpoint inhibitor-induced inflammatory arthritis. *Clin Rheumatol* 2019;38:1513–9.
- 167 Friedman CF, Proverbs-Singh TA, Postow MA. Treatment of the immune-related adverse effects of immune checkpoint inhibitors: a review. *JAMA Oncol* 2016;2:1346–53.
- 168 Lesage C, Longvert C, Prey S, *et al.* Incidence and clinical impact of Anti-TNF α treatment of severe immune checkpoint inhibitor-induced colitis in advanced melanoma: the Mecolit survey. *J Immunother* 2019;42:175–9.
- 169 Braaten TJ, Brahmer JR, Forde PM, *et al.* Immune checkpoint inhibitor-induced inflammatory arthritis persists after immunotherapy cessation. *Ann Rheum Dis* 2020;79:332–8.
- 170 Bertrand F, Montfort A, Marcheteau E, *et al.* TNF α blockade overcomes resistance to anti-PD-1 in experimental melanoma. *Nat Commun* 2017;8:2256.
- 171 Perez-Ruiz E, Minute L, Otano I, *et al.* Prophylactic TNF blockade uncouples efficacy and toxicity in dual CTLA-4 and PD-1 immunotherapy. *Nature* 2019;569:428–32.
- 172 Esfahani K, Miller WH. Reversal of autoimmune toxicity and loss of tumor response by interleukin-17 blockade. *N Engl J Med* 2017;376:1989–91.
- 173 Johnson D, Patel AB, Uemura MI, *et al.* IL17A blockade successfully treated psoriasisform dermatologic toxicity from immunotherapy. *Cancer Immunol Res* 2019;7:860–5.
- 174 Schandendorf D, Wolchok JD, Hodi FS, *et al.* Efficacy and safety outcomes in patients with advanced melanoma who discontinued treatment with nivolumab and ipilimumab because of adverse events: a pooled analysis of randomized phase II and III trials. *J Clin Oncol* 2017;35:3807–14.
- 175 Martini DJ, Hamieh L, McKay RR, *et al.* Durable clinical benefit in metastatic renal cell carcinoma patients who discontinue PD-1/PD-L1 therapy for immune-related adverse events (irAEs). *Cancer Immunol Res* 2018;6:402–8.
- 176 Wang DY, Salem J-E, Cohen JV, *et al.* Fatal toxic effects associated with immune checkpoint inhibitors: a systematic review and meta-analysis. *JAMA Oncol* 2018;4:1721–8.
- 177 Anquetil C, Salem J-E, Lebrun-Vignes B, *et al.* Immune checkpoint Inhibitor–Associated myositis. *Circulation* 2018;138:743–5.
- 178 Moslehi JJ, Salem J-E, Sosman JA, *et al.* Increased reporting of fatal immune checkpoint inhibitor-associated myocarditis. *Lancet* 2018;391:933.
- 179 Mammen AL, Rajan A, Pak K, *et al.* Pre-existing antiacetylcholine receptor autoantibodies and B cell lymphopenia are associated with the development of myositis in patients with thymoma treated with avelumab, an immune checkpoint inhibitor targeting programmed death-ligand 1. *Ann Rheum Dis* 2019;78:150–2.
- 180 Kao JC, Brickshawana A, Liawluck T. Neuromuscular complications of programmed cell death-1 (PD-1) inhibitors. *Curr Neurol Neurosci Rep* 2018;18:63.
- 181 Hunter G, Voll C, Robinson CA. Autoimmune inflammatory myopathy after treatment with ipilimumab. *Can J Neurol Sci* 2009;36:518–20.
- 182 Johnson DB, Balko JM, Compton ML, *et al.* Fulminant myocarditis with combination immune checkpoint blockade. *N Engl J Med* 2016;375:1749–55.
- 183 Tauber M, Cohen R, Laly P, *et al.* Severe necrotizing myositis associated with long term anti-neoplastic efficacy following nivolumab plus ipilimumab combination therapy. *Clin Rheumatol* 2019;38:601–2.
- 184 Bilen MA, Subudhi SK, Gao J, *et al.* Acute rhabdomyolysis with severe polymyositis following ipilimumab-nivolumab treatment in a cancer patient with elevated anti-striated muscle antibody. *J Immunother Cancer* 2016;4:36.
- 185 Tay SH, Wong AS, Jeyasekharan AD. A patient with pembrolizumab-induced fatal polymyositis. *Eur J Cancer* 2018;91:180–2.
- 186 Shirai T, Sano T, Kamijo F, *et al.* Acetylcholine receptor binding antibody-associated myasthenia gravis and rhabdomyolysis induced by nivolumab in a patient with melanoma. *Jpn J Clin Oncol* 2016;46:86–8.
- 187 Fellner A, Makranz C, Lotem M, *et al.* Neurologic complications of immune checkpoint inhibitors. *J Neurooncol* 2018;137:601–9.
- 188 Kimura T, Fukushima S, Miyashita A, *et al.* Myasthenic crisis and polymyositis induced by one dose of nivolumab. *Cancer Sci* 2016;107:1055–8.
- 189 Hasegawa Y, Kawai S, Ota T, *et al.* Myasthenia gravis induced by nivolumab in patients with non-small-cell lung cancer: a case report and literature review. *Immunotherapy* 2017;9:701–7.
- 190 Behling J, Kaes J, Münzel T, *et al.* New-onset third-degree atrioventricular block because of autoimmune-induced myositis under treatment with anti-programmed cell death-1 (nivolumab) for metastatic melanoma. *Melanoma Res* 2017;27:155–8.
- 191 Diamantopoulos PT, Tsatsou K, Benopoulou O, *et al.* Inflammatory myopathy and axonal neuropathy in a patient with melanoma following pembrolizumab treatment. *J Immunother* 2017;40:221–3.
- 192 Hibino M, Maeda K, Horiuchi S, *et al.* Pembrolizumab-induced myasthenia gravis with myositis in a patient with lung cancer. *Respirol Case Rep* 2018;6:e00355.
- 193 Monge C, Maeng H, Brofferio A, *et al.* Myocarditis in a patient treated with nivolumab and PROSTVAC: a case report. *J Immunother Cancer* 2018;6:150.
- 194 Yoshioka M, Kambe N, Yamamoto Y, *et al.* Case of respiratory discomfort due to myositis after administration of nivolumab. *J Dermatol* 2015;42:1008–9.

- 195 Chen J-H, Lee K-Y, Hu C-J, *et al.* Coexisting myasthenia gravis, myositis, and polyneuropathy induced by ipilimumab and nivolumab in a patient with non-small-cell lung cancer: a case report and literature review. *Medicine* 2017;96:e9262.
- 196 Badovinac S, Korsic M, Zarkovic K, *et al.* Nivolumab-induced synchronous occurrence of myositis and hypothyroidism in a patient with squamous cell lung cancer. *Immunotherapy* 2018;10:427–31.
- 197 Kang KH, Grubb W, Sawlani K, *et al.* Immune checkpoint-mediated myositis and myasthenia gravis: a case report and review of evaluation and management. *Am J Otolaryngol* 2018;39:642–5.
- 198 Bourgeois-Vionnet J, Joubert B, Bernard E, *et al.* Nivolumab-induced myositis: a case report and a literature review. *J Neurol Sci* 2018;387:51–3.
- 199 Sheik Ali S, Goddard AL, Luke JJ, *et al.* Drug-associated dermatomyositis following ipilimumab therapy: a novel immune-mediated adverse event associated with cytotoxic T-lymphocyte antigen 4 blockade. *JAMA Dermatol* 2015;151:195–9.
- 200 Gibson R, Delaune J, Szady A, *et al.* Suspected autoimmune myocarditis and cardiac conduction abnormalities with nivolumab therapy for non-small cell lung cancer. *BMJ Case Rep* 2016;2016. doi:10.1136/bcr-2016-216228. [Epub ahead of print: 20 Jul 2016].
- 201 Yamaguchi Y, Abe R, Haga N, *et al.* A case of drug-associated dermatomyositis following ipilimumab therapy. *Eur J Dermatol* 2016;26:320–1.
- 202 Fox E, Dabrow M, Ochsner G. A case of Nivolumab-Induced myositis. *Oncologist* 2016;21:e3.
- 203 Gonzalez NL, Puwanant A, Lu A, *et al.* Myasthenia triggered by immune checkpoint inhibitors: new case and literature review. *Neuromuscul Disord* 2017;27:266–8.
- 204 Liao B, Shroff S, Kamiya-Matsuoka C, *et al.* Atypical neurological complications of ipilimumab therapy in patients with metastatic melanoma. *Neuro Oncol* 2014;16:589–93.
- 205 Haddox CL, Shenoy N, Shah KK, *et al.* Pembrolizumab induced bulbar myopathy and respiratory failure with necrotizing myositis of the diaphragm. *Ann Oncol* 2017;28:673–5.
- 206 Vallet H, Gaillet A, Weiss N, *et al.* Pembrolizumab-induced necrotic myositis in a patient with metastatic melanoma. *Ann Oncol* 2016;27:1352–3.
- 207 Norwood TG, Westbrook BC, Johnson DB, *et al.* Smoldering myocarditis following immune checkpoint blockade. *J Immunother Cancer* 2017;5:91.
- 208 Tan RYC, Toh CK, Takano A. Continued Response to One Dose of Nivolumab Complicated by Myasthenic Crisis and Myositis. *J Thorac Oncol* 2017;12:e90–1.
- 209 Pushkarevskaya A, Neuberger U, Dimitrakopoulou-Strauss A, *et al.* Severe ocular myositis after ipilimumab treatment for melanoma: a report of 2 cases. *J Immunother* 2017;40:282–5.
- 210 Lecoufflet M, Verschoore M, Giard C, *et al.* [Orbital myositis associated with ipilimumab]. *Ann Dermatol Venerol* 2013;140:448–51.
- 211 Rota E, Varese P, Agosti S, *et al.* Concomitant myasthenia gravis, myositis, myocarditis and polyneuropathy, induced by immune-checkpoint inhibitors: a life-threatening continuum of neuromuscular and cardiac toxicity. *eNeurologicalSci* 2019;14:4–5.
- 212 Suzuki S, Ishikawa N, Konoeda F, *et al.* Nivolumab-related myasthenia gravis with myositis and myocarditis in Japan. *Neurology* 2017;89:1127–34.
- 213 Heinzerling L, Ott PA, Hodi FS, *et al.* Cardiotoxicity associated with CTLA4 and PD1 blocking immunotherapy. *J Immunother Cancer* 2016;4:50.
- 214 Läubli H, Balmelli C, Bossard M, *et al.* Acute heart failure due to autoimmune myocarditis under pembrolizumab treatment for metastatic melanoma. *J Immunother Cancer* 2015;3:11.
- 215 Baldetti L, Melillo F, Beneduce A, *et al.* Combined checkpoint inhibitor-associated myocarditis and pulmonary vasculitis mimicking acute pulmonary embolism. *Eur Heart J Cardiovasc Imaging* 2019;20:243.
- 216 Dobloug GC, Svensson J, Lundberg IE, *et al.* Mortality in idiopathic inflammatory myopathy: results from a Swedish nationwide population-based cohort study. *Ann Rheum Dis* 2018;77:40–7.
- 217 Rosenbohm A, Buckert D, Gerischer N, *et al.* Early diagnosis of cardiac involvement in idiopathic inflammatory myopathy by cardiac magnetic resonance tomography. *J Neurol* 2015;262:949–56.
- 218 Bonaca MP, Olenchock BA, Salem J-E, *et al.* Myocarditis in the setting of cancer therapeutics: proposed case definitions for emerging clinical syndromes in Cardio-Oncology. *Circulation* 2019;140:80–91.
- 219 Kudo F, Watanabe Y, Iwai Y, *et al.* Advanced lung adenocarcinoma with Nivolumab-associated dermatomyositis. *Intern Med* 2018;57:2217–21.
- 220 Parker MJ, Roberts ME, Lorigan PC, *et al.* Autoimmune fasciitis triggered by the anti-programmed cell death-1 monoclonal antibody nivolumab. *BMJ Case Rep* 2018;2018. doi:10.1136/bcr-2017-223249. [Epub ahead of print: 08 Feb 2018].
- 221 Khoja L, Maurice C, Chappell M, *et al.* Eosinophilic fasciitis and acute encephalopathy toxicity from pembrolizumab treatment of a patient with metastatic melanoma. *Cancer Immunol Res* 2016;4:175–8.
- 222 Daoussis D, Kraniotis P, Liossis S-N, *et al.* Immune checkpoint inhibitor-induced myofasciitis. *Rheumatology* 2017;56:2161.
- 223 Kobak S. Pembrolizumab-Induced seronegative arthritis and fasciitis in a patient with lung adenocarcinoma. *Curr Drug Saf* 2019;14:225–9.
- 224 Toussaint F, Hammon M, Erdmann M, *et al.* Checkpoint inhibitor-induced eosinophilic fasciitis following high eosinophilia associated with complete response. *Rheumatology* 2019;58:1875–7.
- 225 Mahmood SS, Fradley MG, Cohen JV, *et al.* Myocarditis in patients treated with immune checkpoint inhibitors. *J Am Coll Cardiol* 2018;71:1755–64.
- 226 John S, Antonia SJ, Rose TA, *et al.* Progressive hypoventilation due to mixed CD8⁺ and CD4⁺ lymphocytic polymyositis following tremelimumab - durvalumab treatment. *J Immunother Cancer* 2017;5:54.
- 227 Salem J-E, Allenbach Y, Vozy A, *et al.* Abatacept for severe immune checkpoint inhibitor-associated myocarditis. *N Engl J Med* 2019;380:2377–9.
- 228 Esfahani K, Buhlaiga N, Thébault P, *et al.* Alemtuzumab for immune-related myocarditis due to PD-1 therapy. *N Engl J Med* 2019;380:2375–6.
- 229 Johnson DB, Sullivan RJ, Ott PA, *et al.* Ipilimumab therapy in patients with advanced melanoma and preexisting autoimmune disorders. *JAMA Oncol* 2016;2:234–40.
- 230 Lee B, Wong A, Kee D, *et al.* The use of ipilimumab in patients with rheumatoid arthritis and metastatic melanoma. *Ann Oncol* 2016;27:1174–7.
- 231 Kähler KC, Eigentler TK, Gesierich A, *et al.* Ipilimumab in metastatic melanoma patients with pre-existing autoimmune disorders. *Cancer Immunol Immunother* 2018;67:825–34.
- 232 Menzies AM, Johnson DB, Ramanujam S, *et al.* Anti-PD-1 therapy in patients with advanced melanoma and preexisting autoimmune disorders or major toxicity with ipilimumab. *Ann Oncol* 2017;28:368–76.
- 233 Gutzmeh R, Koop A, Meier F, *et al.* Programmed cell death protein-1 (PD-1) inhibitor therapy in patients with advanced melanoma and preexisting autoimmunity or ipilimumab-triggered autoimmunity. *Eur J Cancer* 2017;75:24–32.
- 234 Danlos F-X, Voisin A-L, Dyeve V, *et al.* Safety and efficacy of anti-programmed death 1 antibodies in patients with cancer and pre-existing autoimmune or inflammatory disease. *Eur J Cancer* 2018;91:21–9.
- 235 Leonardi GC, Gainor JF, Altan M, *et al.* Safety of programmed death-1 pathway inhibitors among patients with non-small-cell lung cancer and preexisting autoimmune disorders. *J Clin Oncol* 2018;36:1905–12.
- 236 Tison A, Quéré G, Misery L, *et al.* Safety and efficacy of immune checkpoint inhibitors in patients with cancer and preexisting autoimmune disease: a nationwide, multicenter cohort study. *Arthritis Rheumatol* 2019;71:2100–11.
- 237 Tokunaga A, Sugiyama D, Maeda Y, *et al.* Selective inhibition of low-affinity memory CD8⁺ T cells by corticosteroids. *J Exp Med* 2019;216:2701–13.
- 238 de Moel E, Rozeman E, Kapiteijn E, *et al.* Treatment with Immune Checkpoint Inhibitors and the Development of Autoantibodies [abstract]. *Arthritis Rheumatol* 2018;70.
- 239 Meti N, Esfahani K, Colmegna I, *et al.* Elevated sCD40L As a Predictive Biomarker of Immune-Related Adverse Events in Patients Receiving Immune Checkpoint Inhibitors [abstract]. *Arthritis Rheumatol* 2018;70.
- 240 Sakakida T, Ishikawa T, Chihara Y, *et al.* Safety and efficacy of PD-1/PD-L1 blockade in patients with preexisting antinuclear antibodies. *Clin Transl Oncol* 2019. doi:10.1007/s12094-019-02214-8. [Epub ahead of print: 01 Oct 2019].
- 241 Toi Y, Sugawara S, Sugisaka J, *et al.* Profiling preexisting antibodies in patients treated with anti-PD-1 therapy for advanced non-small cell lung cancer. *JAMA Oncol* 2019;5:376–83.
- 242 Gleichmann E, Pals ST, Rolink AG, *et al.* Graft-Versus-Host reactions: clues to the etiopathology of a spectrum of immunological diseases. *Immunol Today* 1984;5:324–32.

Managing irAEs with checkpoint inhibitors

This is the lay version of the EULAR 'points to consider' for the diagnosis and management of rheumatic immune-related side effects in people taking checkpoint inhibitors for cancer. The original publication can be downloaded from the EULAR website: www.eular.org.

Kostine M, et al. EULAR points to consider for the diagnosis and management of rheumatic immune-related adverse events due to cancer immunotherapy with checkpoint inhibitors. *Ann Rheum Dis* 2021;80:36–48.
[doi:10.1136/annrheumdis-2020-217139](https://doi.org/10.1136/annrheumdis-2020-217139)

Introduction

EULAR recommendations give advice to doctors, nurses and patients about the best way to treat and manage diseases. EULAR has written new 'points to consider' on how to diagnose and manage rheumatic immune-related side effects in people taking a type of cancer drug called a checkpoint inhibitor.

Doctors, nurses, other health professionals and patients worked together to develop this advice. The patients in the team ensured that the patient point of view was included.

What do we already know?

Checkpoint inhibitors act on the immune system to treat several types of cancer. This group of drugs includes ipilimumab, nivolumab, pembrolizumab, cemiplimab, atezolizumab, avelumab and durvalumab.

About 10% of people who take a checkpoint inhibitor drug to treat a cancer develop musculoskeletal side effects called *immune-related adverse events* (shorted to irAEs).

EULAR has written a set of 'points to consider' on how rheumatic irAEs should be diagnosed and managed. These aim to help oncologists and rheumatologists work together to treat people who develop these side effects. At the moment there is a very low level of evidence on this subject, so it is not possible to develop full recommendations.

What do the points say?

In total, there are 4 overarching principles and 10 points to consider. The principles highlight that these rheumatic and musculoskeletal irAEs are common in people taking checkpoint inhibitors, and should be managed on a shared-decision basis between patients, their treating oncologist, and the rheumatology team. Rheumatologists should work to assist the oncologist in differential diagnosis, and help to relieve symptoms to enable patients to maintain their cancer immunotherapy.

Each point is based on the best current knowledge and studies of scientific evidence or expert opinion. The more stars a point has the stronger the evidence is. However, even recommendations with limited scientific evidence may be important, because the experts may still have a strong opinion and the evidence may be lagging behind.

One star (*) means it is a recommendation with limited scientific evidence.

Two stars (**) means it is a recommendation with some scientific evidence.

Three stars (***) means it is a recommendation with quite a lot of scientific evidence.

Four stars (****) means it is a recommendation supported with a lot of scientific evidence.

- **Rheumatologists should be aware of the wide spectrum of rheumatic immune-related adverse events that often do not fulfil traditional criteria of rheumatic diseases.***

There is a lot of variation in the types of irAEs that people get from checkpoint inhibitors. These can include arthritis and joint pain, myositis, dry mouth, skin thickening, fever, fatigue, etc. Rheumatologists should be aware of the possible symptoms that can mimic rheumatic diseases.

- **Oncologists should consult rheumatologists quickly when rheumatic symptoms are suspected due to immunotherapy, and rheumatologists should provide access for these patients.***

If people taking checkpoint inhibitors develop rheumatic irAEs, their treating oncologist should quickly refer them to a rheumatologist, ideally before trying any steroid treatment. Rheumatologists should see these patients quickly as they are best-placed to try low-dose or steroid-sparing treatments.

- **Some side effects of the cancer itself or unrelated rheumatic diseases might look like immune-related events. Clinical evidence, laboratory tests, imaging and biopsies should be collected to search for inflammation and exclude other diseases.***

Checkpoint inhibitors are often used in people with advanced cancer, and so new rheumatic or musculoskeletal symptoms may not always be irAEs. The oncology team should collect clinical evidence to rule out cancer progression. It is important that this is done quickly to ensure the correct treatment can be given. Then, the rheumatologist should collect evidence of inflammation (on joints, on muscles, on vessels, etc...) either by clinical evidence, or blood tests and imaging, and if required with biopsies.

- **If other treatments do not work, steroids can be considered for symptoms that look like immune-related rheumatic or connective tissue diseases.***

If treatment with painkillers or non-steroidal anti-inflammatory drugs does not work to control your symptoms, you may be offered glucocorticoids (steroids). This will depend on your particular set of symptoms, but steroids injections or tablets can be useful for arthritis or joint pain. Once your irAE symptoms have improved, the steroid should be reduced to the lowest dose needed to maintain control. This is because steroids used for a long time or at a high dose can cause side effects or may alter how your tumour responds to treatment.

- **csDMARD should be considered in people with insufficient response to glucocorticoids or those requiring glucocorticoid-sparing.***

If steroids do not work to control your symptoms – or if they only work for you at a high dose (more than 10 mg per day) – then you may be offered a conventional synthetic disease-modifying antirheumatic drug instead (often shortened to csDMARD). These types of drugs include methotrexate, hydroxychloroquine or sulfasalazine.

- **bDMARDs can be considered for people with severe rheumatic and connective tissue disease-like immune-related adverse events or those with insufficient response to csDMARDs.***

People experiencing severe rheumatic and systemic irAEs, or those with an insufficient response to csDMARDs can try a biologic (also called a bDMARD). For people with symptoms of inflammatory arthritis, the preferred options are TNF or IL-6 inhibitors.

- **The decision to stop or to continue cancer immunotherapy should be made with the patient, based on side effect severity, the tumour response and duration, and the treatment plan.***

At the moment there is no agreement about whether people with irAEs should stay on the checkpoint inhibitor. This might vary depending on your country or treating team, but you should be involved in the

discussions and decision-making.

- **Myositis may be a severe condition, and stopping cancer immunotherapy needs to be discussed. If there are life-threatening symptoms, other treatment options can be used instead.***

Myositis is inflammation in a person's muscles, causing weakness and pain. Myositis can also affect the muscles of the heart, which can potentially be fatal. If myositis is suspected, it may be necessary to stop the checkpoint inhibitor. People who have life-threatening symptoms such as dysphagia, dysarthria, dysphonia, dyspnoea or myocarditis can be treated with high-dose glucocorticoids, intravenous immunoglobulin (IVIg) and/or plasma exchange.

- **Cancer immunotherapy can be used in people who have a pre-existing autoimmune rheumatic disease, while keeping their immunosuppressive treatment for the rheumatic disease at the lowest dose possible.***

People with pre-existing inflammatory or autoimmune diseases can take checkpoint inhibitors for cancer, but they and their treating team should be aware that they may experience disease flares. In most cases, these flares can be managed with steroids.

- **There is no need to test for autoantibodies before starting cancer immunotherapy, but these should be checked if there are rheumatic, musculoskeletal or systemic symptoms.***

Autoantibodies are not always present in people with irAEs, so there is no need to test everyone before they start treatment with a checkpoint inhibitor. However, it can be useful as part of a complete rheumatology assessment if you have unexplained rheumatic, musculoskeletal or systemic symptoms.

Summary

Overall, these points to consider provide the basis of a EULAR consensus in a new and rapidly expanding field. Early consultation and strong collaboration between the referring oncologist, the treating rheumatologist, other organ specialists and the patient are all required for optimal management.

Recommendations with just one or two stars are based mainly on expert opinion and may not be backed up sufficiently yet by studies. These may be as important as those with three or four stars.

If you have any questions or concerns about your disease or your medication, you should speak to a health professional involved in your care.