


Short-term dose and duration-dependent glucocorticoid risk for cardiovascular events in glucocorticoid-naïve patients with rheumatoid arthritis

Anthony James Ocon ¹, George Reed,^{2,3} Dimitrios A Pappas ^{3,4,5},
Jeffrey R Curtis ⁶, Joel M Kremer^{3,5,7}

Handling editor Josef S Smolen

► Additional online supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2021-220577>).

For numbered affiliations see end of article.

Correspondence to

Dr Anthony James Ocon, Allergy, Immunology, Rheumatology, University of Rochester Medical Center, Rochester, NY 14642, USA; anthony.ocon@gmail.com

Received 16 April 2021
Accepted 22 June 2021
Published Online First
2 July 2021

ABSTRACT

Objectives Rheumatoid arthritis (RA), along with glucocorticoid use, is associated with cardiovascular disease. Cardiovascular safety of glucocorticoids in RA is controversial and may be related to dose and duration of use. We determined if initiating glucocorticoids in steroid-naïve RA patients would increase cardiovascular event (CVE) risk in a dose and duration-dependent manner over short-term intervals.

Methods Patients enrolled in CorEvitas (formerly Corrona) RA registry. Cox proportional-hazards models estimated adjusted HRs (aHR) for incident CVE in patients who initiated glucocorticoid treatment, adjusting for RA duration, traditional cardiovascular risk factors and time-varying covariates: Clinical Disease activity Index, disease-modifying antirheumatic drugs use and prednisone-equivalent use. Glucocorticoid use assessed current daily dose, cumulative dose and duration of use over rolling intervals of preceding 6 months and 1 year.

Results 19902 patients met criteria. 1106 CVE occurred (1.66/100 person-years). Increased aHR occurred at current doses of ≥ 5 –9 mg 1.56 (1.18–2.06) and ≥ 10 mg 1.91 (1.31–2.79), without increased risk at 0–4 mg 1.04 (0.55–1.59). Cumulative dose over preceding 6 months showed increased aHR at 751–1100 mg 1.43 (1.04–1.98) and >1100 mg 2.05 (1.42–2.94), without increased risk at lower doses; duration of use over preceding 6 months exhibited increased aHR for >81 days of use 1.54 (1.08–2.32), without increased risk at shorter durations. One-year analyses were consistent.

Conclusions Over preceding 6-month and 1-year intervals, initiating glucocorticoids in steroid-naïve RA patients is associated with increased risk of CVE at daily doses ≥ 5 mg and increased cumulative dose and duration of use. No association with risk for CVE was found with daily prednisone of ≤ 4 mg or shorter cumulative doses and durations.

INTRODUCTION

Rheumatoid arthritis (RA) is a systemic autoimmune disease characterised by inflammatory destructive arthritis. Risk for cardiovascular disease (CVD) in RA is increased due to high prevalence of traditional risk factors, accelerated atherosclerosis and chronic inflammation.¹ Disease activity is directly related to cardiovascular risk.² Glucocorticoids (GCs) are commonly prescribed as initial, so-called bridge, treatment for RA but are often employed

Key message

What is already known about this subject?

- Both rheumatoid arthritis (RA) and glucocorticoid use increase risk for cardiovascular events. There is controversy regarding the cardiovascular safety and risk of glucocorticoid use in RA patients. Effects of short-term and low-dose use are not well understood.

What does this study add?

- In a large, real-world clinical registry of patients with long-standing disease, there is a daily dose, cumulative dose and duration of use glucocorticoid threshold for cardiovascular event risk in RA when analysed over short-term intervals of 6 months and 1 year.
- Relative cardiovascular safety was found with <5 mg of prednisone-equivalent daily dose and lower cumulative doses and durations of use.

How might this impact on clinical practice or future developments?

- Physicians should be aware of that low-dose and short-term use of glucocorticoids may increase risk of cardiovascular events when prescribing for treatment of RA in a treat-to-target approach.
- Patient education of this risk threshold is essential.

for intervals that extend beyond the onset of action of other conventional, targeted synthetic or biological disease-modifying antirheumatic drugs (cs/ts/bDMARDs).³ It is often clinically challenging to taper GC. However, GCs are associated with CVD and may potentiate hypertension, hyperlipidaemia, diabetes mellitus, congestive heart failure and obesity.^{4–6} Given that CVD is the major comorbidity of RA,^{1,2} the juxtaposition of these circumstances presents a therapeutic dilemma.

Controversy exists regarding the risks and benefits of GC in RA patients. Previous small studies demonstrated an increased number of adverse events in RA patients over longer intervals with a daily prednisone-equivalent doses of >5 to 10 mg.^{7–10} However, debate remains regarding the detrimental



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Ocon AJ, Reed G, Pappas DA, et al. *Ann Rheum Dis* 2021;**80**:1522–1529.

cardiovascular effects of GC therapy in RA patients.^{11–16} Relative cardiovascular safety is generally assumed with lower dose and shorter durations of use, especially over short-term intervals. However, little data has actually been reported regarding the temporal effects of short-term interval GC use preceding cardiovascular events (CVE).

The 2016 and 2019 European League Against Rheumatism and 2015 and 2021 American College of Rheumatology (ACR) recommend the use of ‘low-dose’ GC for ‘the least amount of time’ in combination with DMARDs for the treatment of RA.^{3, 17–19} Thus, it is important to determine the safety of initiating ‘low-dose’ GC in regard to the development of CVE. Furthermore, CVE in RA may be decreasing due to better control of disease activity following the widespread use of ts/bDMARDs, perhaps making the determination of the contribution of GC to CVE even more challenging in the present era.²⁰

We, therefore, examined the CorEvitas (formerly Corrona) RA registry, a longitudinal database of RA patients, to determine whether there was a relationship between CVE in RA and use of GC in the dose ranges and duration of use that are consistent with published guidelines and routine clinical practice. We sought to determine the relative safety or risk for incident CVE in steroid-naïve patients who initiate ‘low-dose’ GC over short-term intervals of use based on real-world clinical observation, while adjusting for traditional cardiovascular risk factors, RA duration and disease activity and cs/ts/bDMARD use. Given the ubiquitous use of GC in this and other inflammatory conditions in a variety of dose ranges, including what is considered ‘low-dose,’ it was both timely and appropriate to reexamine this association.

METHODS

Study Cohort Entry Criteria

The CorEvitas (formerly Corrona) RA registry was previously described.^{21, 22} Inclusion criteria included age ≥ 18 years old and receiving a diagnosis of RA by a rheumatologist.²³ Data were collected between 1 October 2001 and 31 March 2018. During this period, 48 535 patients enrolled. Exclusion criteria included: any history of current or past GC therapy at or prior to enrolment; absence of a follow-up visit; missing data for either gender, age or duration of RA; or patients that had > 15 months between visits. Patients were treated per their rheumatologist without treatment assignment.

Data collection

Observational data were collected from both treating physicians and patients at registry enrolment and at regular intervals consistent with the frequency of scheduled visits occurring every 2–9 months (median 4.6 months, IQR 3.60–6.24). At enrolment, detailed medical history was obtained from patients and review of medical records to accurately document lifetime comorbidities and prior treatments and medication use, including GC use.

Measure of GC use

GC use after entry into the registry was documented as equivalent milligrams of prednisone. Multiple measures of GC use were assessed. Current daily dose was defined as the most recent recorded dose at the time of a CVE or the most recent recorded dose in the registry for patients without an event. Cumulative total dose was defined as the summation of prednisone-equivalent dosage updated at each visit in a continuous, rolling manner over the preceding 6 months or 1 year (see online appendix efigure 1). Duration of use was defined as the

summation of the absolute number of days a patient was treated with GC in a rolling manner over the preceding 6 months or 1 year (see online appendix efigure 1). Interval ranges for daily and cumulative dose were chosen based on equivalent quartiles of patient-time. Quartiles of duration of use were also chosen to have intervals with equal numbers of patient-time in each.

Event definition and documentation

For this study, CVE were defined as cardiac death, myocardial infarction, stroke, hospitalisation for hypertension, coronary revascularisation procedures, ventricular arrhythmia, unstable angina, congestive heart failure, transient ischaemic attacks, deep vein thrombosis (DVT), peripheral arterial thromboembolic event, urgent peripheral arterial revascularisation, peripheral arterial ischaemia, pulmonary embolism (PE), acute coronary syndrome or ‘other’ event. ‘Other’ events included complex or overlapping events, other arrhythmias or conduction abnormality, cardiomyopathy, unspecified coronary artery disease, or events the reporting physician felt more comfortable categorising as ‘other’ if there was potential overlap with category choices provided.

At follow-up visits, both physician and patient-derived clinical data were updated in detail, including medication and dose changes for GC and cs/ts/bDMARDs.²² Incident comorbidities and targeted medical events, including CVE, were specifically ascertained and collected at each visit by the treating rheumatologist (see online appendix file 1). After the receipt of a report of CVE on the registry form, the site then completed a separate e-form with deidentified primary hospital or cardiologist records confirming and validating the event with description of specific drugs and dose used for treatment (see online appendix efigure 2).²⁴ These forms were reviewed to confirm and validate the event, and ensure that it had not been previously reported, with any duplicate events excluded. In addition, a physician could report an event between formal registry visits. Finally, CVE, in particular CV death, were also reported on registry exit form. Any event that was not confirmed and validated was excluded.

Data analysis

The registry enrolment visit date was the index date. Only the first CVE following enrolment was used. Missing data for covariates were carried forward from the prior visit. If missing GC dose at a visit occurred, the prior dose was carried forward.

Time to first CVE was modelled using Cox proportional-hazards regression models to estimate unadjusted and adjusted HR and 95% CIs. Our model computed cumulative dose or duration of use over the preceding 6 months or 1 year at every daily time point from the index date to the last time point for each patient. This last time point could be a CVE, last registry visit, or dropout from the registry, whichever occurred first. At each time point, the model compared the risk of a CVE in patients at each quartile of prednisone use (current dose, cumulative dose, duration of use) to the risk in patients with no use.

For the adjusted analysis, baseline covariables in the model included age, sex, race, duration of RA, history of CV disease, diabetes mellitus, hyperlipidaemia, hypertension, statin use, NSAID use, tobacco use, year of enrolment, baseline modified Health Assessment Questionnaire score (mHAQ) and the baseline Clinical Disease Activity Index (CDAI) for RA. The CDAI is a validated disease activity metric that includes tender and swollen joints (28 joint count), as well as physician and patient evaluation of global arthritis activity on a 10-point Visual Analogue Scale.²⁵ Additionally, time-varying covariates in the

model included measures of prednisone use as described above, NSAID use, cs, b, tsDMARDS and CDAI, which were updated at each follow-up visit.

A sensitivity analysis excluded all venous thromboembolisms (DVT and/or PE) as CVE to determine if excluding venous events impacted risk. A different sensitivity analysis excluded all patients with prior history of a CVE to assess whether this comorbidity had influenced risk. Another sensitivity analysis excluded 'other' CVE to assess its influence on the outcomes of interest.

Student's t-test or χ^2 test compared data at baseline. All analyses were generated using Stata V.16.1 (StataCorp).

RESULTS

Sample characteristics

A total of 19 902 patients (41%) met entry criteria. Exclusions occurred as follows: 21 162 patients had prior history of prednisone use; 5059 patients had no follow-up; 42 patients were missing information regarding use of prednisone; 1672 patients had time between visits of >15 months; 743 patients had missing information for covariates (age, gender, duration of RA, smoking status, CDAI and/or mHAQ).

For the 19 902 who met criteria, the follow-up included 66 436 patient-years and 127 674 follow-up visits over >16 years. Of these patients, 2500 (12.6%) initiated GC during the follow-up. Median time to first use in the registry was 19 months (IQR: 9.1–38.4) after enrolment.

Assessment of CVE risk with initiating GC use

A total of 1106 CVE occurred, yielding a rate of 1.66 CVE per 100 patient-years (95% CI 1.57 to 1.77). As depicted in [table 1](#) of unadjusted enrolment characteristics prior to any CVE and follow-up interval, patients who developed CVE had a greater prevalence of traditional CV risk factors, more severe RA, and were more likely to use csDMARD. [Table 2](#) displays the frequency of each CVE.

[Table 3](#) displays unadjusted and adjusted HR for daily and cumulative dose and duration of GC use over the preceding 6 month and 1 year intervals. Online appendix table 1 shows the number of patients contributing time to each category. Unadjusted current daily dose of <5 mg was not associated with increased risk, while doses ≥ 5 mg increased risk in a dose-response manner. [Figure 1](#) demonstrates the adjusted risk of CVE based on daily prednisone-equivalent dose with similar findings.

As shown in [table 3](#), cumulative doses of >750 mg over the preceding 6 months were associated with increased unadjusted risk for developing a CVE. [Figure 2](#) shows the risk for developing a CVE remained for cumulative doses of >750 mg after adjustment for covariates. In both unadjusted and adjusted analyses, cumulative doses ≤ 750 mg were not associated with increased risk. Also shown in [table 3](#), cumulative doses of >1110 mg over the preceding 1 year were associated with significant increased unadjusted risk for developing a CVE. [Figure 2](#) shows this risk remained for cumulative doses of >1110 mg after adjustment for covariates. In both unadjusted and adjusted analyses, cumulative doses ≤ 1110 mg over the preceding 1-year were not associated with increased risk.

As shown in [table 3](#), GC use for >80 days over the preceding 6 months interval was associated with increased unadjusted risk for developing a CVE. Shorter use than 80 days was not associated with increased risk. [Figure 3](#) illustrates similar risk when adjusted for covariates. Over the preceding 1-year interval, a

similar, if less smooth, increased unadjusted ([table 3](#)) and adjusted ([figure 2](#)) risk for a CVE after 100 days of use was found.

Sensitivity analyses

Online appendix table 2 shows the results of the sensitivity analysis when DVT and PE were excluded. With exclusion of DVT/PE, 1007 CVE occurred. Results were similar to the primary analysis.

Online appendix table 3 shows the results of the sensitivity analysis when a history of prior CVE was excluded. With this exclusion, the total number of patients was 18 168, with 2300 initiating prednisone. There were 829 CVE in this analysis. Results were similar to the primary analysis.

Online appendix table 4 shows the results of the sensitivity analysis when 'other' CVE were excluded. With this exclusion, 817 events occurred. The results were similar to the primary analysis.

DISCUSSION

We report, for the first time, that the relative cardiovascular safety or risk of initiating GC in a real-world clinical sample of steroid-naïve RA patients with longstanding disease at registry enrolment is associated with a threshold daily dose, cumulative dose, and duration of use when analysed over short-term intervals of the preceding 6 months or 1 year. The risk for CVE increased directly with increasing current daily dose, with the greatest estimated risk at ≥ 5 –9 mg and ≥ 10 mg of prednisone-equivalents. Similarly, the risk for CVE increased in a dose-response manner with increasing cumulative dose over these short-term intervals of analysis. The risk for CVE based on duration of use found increased risk after 80-days of use over the preceding 6 months and 100-days over the preceding 1 year in the dose ranges reported. There is 'noise' in the duration of use data, especially over the preceding 1 year, and it is possible that this is due to a threshold effect related to dose. That is, the duration of use analysis does not necessarily account for dose; thus, similar durations of use may have different total doses, especially with longer use. Of additional clinical importance, we found no increased risk for CVE with current prednisone-equivalent daily doses of <5 mg or cumulative doses of ≤ 750 mg over the preceding 6 months or ≤ 1110 mg over the preceding 1 year. We found no increased risk with duration of use ≤ 80 days over the preceding 6 months or ≤ 100 days over the preceding 1 year. It is critically important to note that these findings remained after adjustment for established cardiovascular risk factors, RA duration, disease activity and cs/ts/bDMARDs.

These novel insights can be immediately employed in clinical practice. The methodology of our analysis allows the application to patients based on their most recent 6 months or 1 year GC use. We believe that our data demonstrate that GC use should be tapered to a dose of <5 mg prednisone-equivalents as expeditiously as possible, while being aware of duration of use and cumulative dose. Thus, clinicians should provide counselling and education of these findings when encountering a reluctance on the part of a pain-free patient to taper GC, or succumb to the temptation to simply increase the dose to make the patient feel better until their next visit.

Both the IMPROVED and CareRA trials established that GCs can be tapered in patients with early RA in a protocolised, supervised, investigational regimen.^{26 27} Verschueren *et al* also reported on a supervised step-down GC taper in 19 patients with early RA.²⁸ The observational data we report on in patients

Table 1 Unadjusted sample characteristics stratified by development of a cardiovascular event (CVE) measured at registry enrolment*

Variable	No CVE (N=18 796)		CVE (N=1106)		P value
	Mean	SD	Mean	SD	
Age (years)	57.82	13.36	65.44	11.07	<0.001
Duration of rheumatoid arthritis (years)	8.84	9.42	11.56	10.76	<0.001
mHAQ	0.32	0.42	0.38	0.47	<0.001
CDAI	12.73	12.50	14.04	12.84	0.001
28 Joint Count: Tender	3.93	5.81	4.44	6.39	0.005
28 Joint Count: Swollen	3.81	5.39	4.18	5.33	0.028
Patient Global Assessment (0–100 scale)	28.95	26.08	31.71	26.44	0.001
Physician Global Assessment (0–100 scale)	21.76	20.38	23.75	20.53	0.002
Body mass index (kg/m ²)	29.44	7.14	29.60	7.00	0.468
	N	%	N	%	
Gender					<0.001
Male	4134	22.0	369	33.4	
Female	14 662	78.0	737	66.6	
Race					0.202
Asian	286	1.5	12	1.1	
Black	1184	6.3	55	5	
Mixed race	223	1.2	14	1.3	
Native American	122	0.6	6	0.5	
Other	105	0.6	3	0.3	
Pacific Islander	21	0.1	2	0.2	
Unknown	132	0.7	3	0.3	
White	16 723	89.0	1011	91.4	
History of cardiovascular disease					<0.001
Yes	1457	7.8	277	25.0	
History of diabetes					<0.001
Yes	1502	8.0	176	15.9	
History of hyperlipidaemia					<0.001
Yes	4154	22.1	333	30.1	
History of hypertension					<0.001
Yes	5583	29.7	511	46.2	
Smoking status					0.001
Never	11 330	60.3	604	54.6	
Previous	4946	26.3	339	30.7	
Current	2520	13.4	163	14.7	
Exercise					<0.001
None	5879	31.3	390	35.3	
1–2 times/week	5313	28.3	277	25.0	
3–4 times/week	4054	21.6	213	19.3	
5–6 times/week	1376	7.3	68	6.1	
Daily	1748	9.3	141	12.7	
Not sure	426	2.3	17	1.5	
Statin use					<0.001
Yes	3750	20.0	303	27.4	
NSAID use					0.024
Yes	10 272	54.6	643	58.1	
Analgesic use					<0.001
Yes	8516	45.3	565	51.1	
Prior or current biologic/targeted DMARD use					0.815
Yes	8633	45.9	504	45.6	
Prior or current conventional DMARDs					<0.001
Yes	16 447	87.5	1019	92.1	
Prior or current methotrexate use					<0.001
Yes	14 027	74.6	893	80.7	

*Patients were enrolled with prevalent disease.

CDAI, Clinical Disease Activity Index; DMARDs, disease-modifying antirheumatic drugs; mHAQ, modified Health Assessment Questionnaire; NSAID, non-steroidal anti-inflammatory drug.

Table 2 Frequency of each cardiovascular event

Events	Frequency	Per cent
Acute coronary syndrome	3	0.27
Cardiac arrest	19	1.72
Congestive heart failure	106	9.58
Cardiovascular death	8	0.72
Deep vein thrombosis	83	7.5
Hospitalisation for hypertension	16	1.45
Myocardial infarction	117	10.58
Other cardiovascular event*	289	26.13
Peripheral arterial event	4	0.36
Peripheral arterial intervention	3	0.27
Pulmonary embolism	26	2.35
Coronary revascularisation	204	18.44
Stroke	136	12.3
Transient ischaemic attack	51	4.61
Unstable angina	16	1.45
Urgent peripheral arterial revascularisation	1	0.09
Ventricular arrhythmia	24	2.17
Total events	1106	100

*Other events included the following: complex or overlapping events (eg, acute coronary syndrome with coronary revascularisation), atrial fibrillation, other supraventricular arrhythmia, unspecified bradycardia, other unspecified conduction abnormality, new but stable angina, unspecified coronary artery disease, cardiac syncope or orthostatic hypotension, cardiomyopathy, other cardiac interventional procedure, abdominal aortic aneurysm, other peripheral arterial disease or cardiac event not otherwise specified.

with longstanding disease are perhaps more representative of a general population of RA patients on GC.

It should also be noted that we stratified our subject sample to see how subjects who developed CVE differed at the baseline time of registry enrolment from those who did not (table 1). Those who developed CVE were older, had more traditional CVD risk factors, greater disease activity, longer disease duration, and more commonly were on non-bDMARDs. Again,




Figure 1 Adjusted risk of cardiovascular (CV) event associated with current daily prednisone-equivalent dose. There is a threshold for increased risk of an event. Prednisone-equivalent doses of 5–9 mg and ≥10 mg were associated with an increased risk. However, doses <5 mg were not associated with increased risk. The risk was adjusted for traditional CV risk factors, rheumatoid arthritis disease activity and duration, and cs/ts/bDMARD use. cs/ts/bDMARD, conventional, targeted synthetic or biological disease-modifying antirheumatic drugs.

Table 3 Unadjusted and adjusted HRs for cardiovascular (CV) event with initiating glucocorticoid use

Daily dose (mg)†	Unadjusted HR (95% CI)	Adjusted* HR (95% CI)
None	1 (ref)	1 (ref)
1-5	1.04 (0.61 to 1.76)	0.94 (0.55 to 1.59)
≥5-9	1.78 (1.35 to 2.35)	1.56 (1.18 to 2.05)
≥10	2.09 (1.44 to 3.05)	1.91 (1.31 to 2.79)
Cumulative dose (mg)†	Unadjusted HR (95% CI)	Adjusted* HR (95% CI)
Over preceding 6 months:		
None	1 (ref)	1 (ref)
1-380	0.93 (0.56 to 1.50)	0.86 (0.53 to 1.40)
381-750	1.31 (0.88 to 1.95)	1.20 (0.81 to 1.79)
751-1100	1.62 (1.18 to 2.24)	1.43 (1.04 to 1.98)
>1110	2.25 (1.57 to 3.22)	2.05 (1.42 to 2.94)
Over preceding 1 year:		
None	1 (ref)	1 (ref)
1-500	0.99 (0.64 to 1.54)	0.93 (0.60 to 1.45)
501-1100	1.28 (0.89 to 1.83)	1.19 (0.83 to 1.70)
1101-2100	1.63 (1.18 to 2.25)	1.47 (1.06 to 2.03)
>2100	1.97 (1.41 to 2.74)	1.74 (1.25 to 2.43)
Duration of use (days)	Unadjusted HR (95% CI)	Adjusted* HR (95% CI)
Over preceding 6 months:		
None	1 (ref)	1 (ref)
1-80	0.77 (0.46 to 1.29)	0.72 (0.60 to 1.45)
81-160	1.66 (1.16 to 2.36)	1.54 (1.08 to 2.20)
161-181	1.71 (0.76 to 3.81)	1.56 (0.70 to 3.48)
>181	1.79 (1.38 to 2.35)	1.57 (1.20 to 2.05)
Over preceding 1 year:		
None	1 (ref)	1 (ref)
1-100	1.08 (0.72 to 1.62)	1.02 (0.68 to 1.53)
101-220	1.50 (1.10 to 2.05)	1.41 (1.03 to 1.93)
221-360	0.99 (0.60 to 1.62)	0.88 (0.54 to 1.44)
>360	2.15 (1.59 to 2.92)	1.88 (1.39 to 2.56)

*Adjusted for age, sex, race, duration of RA, history of CV disease, diabetes mellitus, hyperlipidaemia, hypertension, statin use, NSAID use, tobacco use, year of enrolment, baseline modified health assessment questionnaire score, CDAI and cs, b, tsDMARDs use.

†Prednisone-equivalents.

CDAI, Clinical Disease Activity Index; cs/ts/bDMARDs, conventional, targeted synthetic or biological disease-modifying antirheumatic drugs; NSAID, non-steroidal anti-inflammatory drug; RA, rheumatoid arthritis.

adjustment for these factors found an independent association with GC use.

The methodology of our investigation expands on the prior literature. Both Saag *et al* and Davis *et al* studied adverse events, including CVE, in RA patients receiving GC.^{7 10} However, there are numerous differences in methodology of the present investigation including the historical therapeutic interval and duration of observation,¹⁰ robustness of numbers, as well as specific focus on CVE. Our adjustment for multiple confounding variables further distinguishes our approach from prior studies that did not adjust for all these variables.^{8 9 29-32} Our findings provide context beyond these prior studies by highlighting the relative cardiovascular safety of doses of GCs ≤4 mg daily over the 6-month interval described.^{18 33} Our findings also add evidence to the EULAR and ACR task force recommendations for steroid taper.^{19 34} Huscher *et al* also reported a threshold for GC side effects in RA patients without examining CVE.³⁵

Others have looked at the effects of short-term GC use. George *et al* reported on the effect of dosages on serious infectious events (SIEs) using a Medicare claims database.³⁶ They found a




Figure 2 Top: adjusted risk of cardiovascular (CV) events associated with total glucocorticoid use over preceding 6 month interval. Bottom: adjusted risk of CV event associated with total glucocorticoid use over preceding 1-year interval. There was a dose-response increase in risk for CV event. Over the preceding 6 months of use, cumulative prednisone-equivalent doses of 751–1100 mg and >1100 mg were associated with increased risk for a CV event. Over the preceding 1 year of use, cumulative doses of 1101–2011 mg and >2100 mg were associated with increased risk for a CV event. This risk was adjusted for traditional CV risk factors, rheumatoid arthritis disease activity and duration, and cs/ ts/bDMARD use. cs/ts/bDMARD, conventional, targeted synthetic or biological disease-modifying antirheumatic drugs.

robust relationship of SIEs with increasing doses, although they were not able to determine the effect of actual disease activity. Similarly, Yao *et al* used a national insurance database to assess the effects of short ‘burst’ courses (≤ 14 -days) of prednisone on incident adverse events of gastrointestinal (GI) bleeding, sepsis, and heart failure at 5–30 days and 31–90 days from use, finding a higher incidence rate of these events at both time points in the general population who used prednisone.³⁷ Together, the findings of these studies of predominantly non-cardiac adverse events support our conclusion that detrimental effects of GC are strongly associated with short-term intervals preceding the event.

Our study has several additional strengths. Our investigation spanned a 16-year period, while prior studies were based much shorter duration of observation.^{8 9 29–32 38–40} The data were derived from over 700 participating sites of real-world clinical observation in the USA. The protocol independently confirmed and validated events with hospital records. We then analysed




Figure 3 Top: adjusted risk of cardiovascular (CV) event associated with total duration of use of glucocorticoids over preceding 6-month interval. Bottom: adjusted risk of CV event associated with total duration of use of glucocorticoids over preceding 1-year interval. There was a duration of use threshold for increased risk. Over the preceding 6 months, prednisone-equivalent use for ≥ 81 days was associated with an increased risk for a CV event. Over the preceding 1 year, use for 101–220, and >360 days was associated with an increased risk for a CV event, with trend towards increased risk between 221 and 260 days. The risk was adjusted for traditional cardiovascular risk factors, rheumatoid arthritis disease activity and duration, and cs/ ts/bDMARD use. cs/ts/bDMARD, conventional, targeted synthetic or biological disease-modifying antirheumatic drugs.

reported events, not limiting to MACE, increasing the real-world clinical relevance for practising physicians. We excluded prevalent and past GC users, in an effort to minimise the effect of past use. Sensitivity analyses demonstrated that having a history of prior CVE, venous-related events, or the ‘other’ category of CVE did not influence the results, a particularly important design feature of our analysis.^{41 42}

Our study is not without limitations. Patients in observational registries are treated without assignment of interventions. Although we adjusted for multiple confounding factors, there is still a risk of channelling bias or residual confounding. While GC dose and usage were updated in the registry at each visit, there is potential for patient reporting to be limited by recall bias. However, the percentage of patients we studied without prior GC use at the time of registry enrolment (41%) is similar to that reported in the ARAMIS registry, supporting our robust data collection methods.⁴³ It is not clear whether our findings are

applicable to early RA patients treated with GC since our cohort contained predominantly RA patients with longer disease duration. A similar observational analysis of the success of GC taper in early RA patients not participating in a supervised protocol would be of interest.

A possible limitation of this report is that we were not able to adjust our findings for time-varying changes in C reactive protein, which is a known risk factor for CVE, or for either rheumatoid factor or anticitrullinated protein antibody status as laboratory values are not mandated in this observational registry. However, we adjusted for other traditional cardiovascular risk factors, as well as time-varying changes in actual RA disease activity measured at the time of each registry visit. As is the case with long-term observational data, the cohort of patients at different time points may differ as patients enter and exit the registry. The risk was established with a very large number of patients over very long observational intervals that greatly increased our statistical power. Nevertheless, assignment of risk to a specific individual may not be appropriate. In addition, with any statistical association, we cannot determine causality with absolute certainty. While a prospective randomised controlled trial of GC dosing would be ideal, it is highly unlikely that this kind of trial will ever be conducted given the pragmatic challenges with the number of patients required, study duration, funding, ethics, and other challenges associated with the complete absence of steroid use in a control group.

In conclusion, we reported that daily doses of ≥ 5 mg of prednisone-equivalents, elevated cumulative dose and extended duration of use of GC over the preceding 6-month and 1-year intervals are associated with an increased risk for incident CVE in steroid-naïve patients with RA. We also emphasise the relative absence of CVE with dosing of ≤ 4 mg per day, lower cumulative dose and a duration of use of only 6 months prior to an event. Physicians treating patients with RA should consider these threshold ranges of GC use when prescribing prednisone as a part of a treat-to-target regimen.

Author affiliations

¹Medicine and Allergy, Immunology, Rheumatology, University of Rochester Medical Center, Rochester, New York, USA

²Medicine, Preventative and Behavioral Medicine, University of Massachusetts Medical School, Worcester, Massachusetts, USA

³Corrona Research Foundation, LLC, Waltham, MA, USA

⁴Medicine and Rheumatology, Columbia University, New York, New York, USA

⁵CorEvitas (formerly CORRONA), LLC, Waltham, Massachusetts, USA

⁶Department of Medicine, University of Alabama at Birmingham, Birmingham, Alabama, USA

⁷Medicine and Rheumatology, Albany Medical College, The Center for Rheumatology, LLC, Albany, New York, USA

Contributors AJO and JMK designed the study. Statistical analysis by GR. All authors (AJO, GR, DAP, JRC and JMK) participated in data interpretation, writing, editing and final approval.

Funding The investigation was funded by the Corrona Research Foundation, a not for profit, 501(C)(3) independent charitable foundation, with no industry financial ties

Competing interests AJO has no financial conflicts of interest. GR and JMK are consultants for Corrona, LLC. GR is a consultant for the Corrona Research Foundation (CRF), while JMK is an officer of the CRF who serves without any form of remuneration. The CRF is a not for profit, 501(C)(3) independent charitable foundation, with no industry financial ties. DAP is an employee and shareholder of Corrona, and a consultant for Regeneron, Novartis and Roche (unrelated work), and is a member of the board of directors for CFR. JRC has research grants and/or consulting from Abbvie, Amgen, Corrona, Janssen, Lilly, Pfizer, Sanofi (for unrelated work).

Patient and public involvement statement Patients were involved upon signing informed consent to participate in the registry, understanding that information would be recorded longitudinally and used for a variety of research outcomes that could be relevant to care. Patients were informed about the time

required to complete forms at each visit. Outcomes of CorEvitas studies are shared with patients when viewed as potentially impacting their welfare. Patients were not involved in the design of or recruitment for this study.

Patient consent for publication Not required.

Ethics approval New England Institutional Review Board, ID number 120160610. All patients gave written informed consent.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. Deidentified participant data may be obtained from CorEvitas (formerly Corrona) and are not publicly available. Reuse only with permission from CorEvitas. Contact: info@corevitas.com

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Anthony James Ocon <http://orcid.org/0000-0003-2903-5429>

Dimitrios A Pappas <http://orcid.org/0000-0001-8338-027X>

Jeffrey R Curtis <http://orcid.org/0000-0002-8907-8976>

REFERENCES

- Jagpal A, Navarro-Millán I. Cardiovascular co-morbidity in patients with rheumatoid arthritis: a narrative review of risk factors, cardiovascular risk assessment and treatment. *BMC Rheumatol* 2018;2:10.
- Solomon DH, Reed GW, Kremer JM, et al. Disease activity in rheumatoid arthritis and the risk of cardiovascular events. *Arthritis Rheumatol* 2015;67:1449–55.
- Singh JA, Saag KG, Bridges SL, et al. 2015 American College of rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Care Res* 2016;68:1–25.
- Fardet L, Fève B. Systemic glucocorticoid therapy: a review of its metabolic and cardiovascular adverse events. *Drugs* 2014;74:1731–45.
- Roubille C, Martel-Pelletier J, Davy J-M, et al. Cardiovascular adverse effects of anti-inflammatory drugs. *Antiinflamm Antiallergy Agents Med Chem* 2013;12:55–67.
- Wei L, MacDonald TM, Walker BR. Taking glucocorticoids by prescription is associated with subsequent cardiovascular disease. *Ann Intern Med* 2004;141:764–70.
- Saag KG, Koehnke R, Caldwell JR, et al. Low dose long-term corticosteroid therapy in rheumatoid arthritis: an analysis of serious adverse events. *Am J Med* 1994;96:115–23.
- Sihvonen S, Korpela M, Mustonen J, et al. Mortality in patients with rheumatoid arthritis treated with low-dose oral glucocorticoids. A population-based cohort study. *J Rheumatol* 2006;33:1740–6.
- del Rincón I, Battafarano DF, Restrepo JF, et al. Glucocorticoid dose thresholds associated with all-cause and cardiovascular mortality in rheumatoid arthritis. *Arthritis Rheumatol* 2014;66:264–72.
- Davis JM, Maradit Kremers H, Crowson CS, et al. Glucocorticoids and cardiovascular events in rheumatoid arthritis: a population-based cohort study. *Arthritis Rheum* 2007;56:820–30.
- Hafström I, Rohani M, Deneberg S, et al. Effects of low-dose prednisolone on endothelial function, atherosclerosis, and traditional risk factors for atherosclerosis in patients with rheumatoid arthritis—a randomized study. *J Rheumatol* 2007;34:1810–6.
- Mazzantini M, Torre C, Miccoli M, et al. Adverse events during long-term low-dose glucocorticoid treatment of polymyalgia rheumatica: a retrospective study. *J Rheumatol* 2012;39:552–7.
- van Everdingen AA, Jacobs JW, Siewertsz Van Reesema DR, et al. Low-Dose prednisone therapy for patients with early active rheumatoid arthritis: clinical efficacy, disease-modifying properties, and side effects: a randomized, double-blind, placebo-controlled clinical trial. *Ann Intern Med* 2002;136:1–12.
- Ruyssen-Witrand A, Fautrel B, Saraux A, et al. Cardiovascular risk induced by low-dose corticosteroids in rheumatoid arthritis: a systematic literature review. *Joint Bone Spine* 2011;78:23–30.
- Solomon DH, Greenberg J, Curtis JR, et al. Derivation and internal validation of an expanded cardiovascular risk prediction score for rheumatoid arthritis: a consortium of rheumatology researchers of North America registry study. *Arthritis Rheumatol* 2015;67:1995–2003.
- Da Silva JAP, Jacobs JW, Kirwan JR, et al. Safety of low dose glucocorticoid treatment in rheumatoid arthritis: published evidence and prospective trial data. *Ann Rheum Dis* 2006;65:285–93.

- 17 Smolen JS, Landewé R, Bijlsma J, *et al.* EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis* 2017;76:960–77.
- 18 Smolen JS, Landewé RBM, Bijlsma JWJ, *et al.* EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis* 2020;79:685–99.
- 19 Fraenkel L, Bathon JM, England BR. 2021 American College of rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Rheumatol* 2021.
- 20 Arts EE, Fransen J, Den Broeder AA, *et al.* Low disease activity (DAS28 \leq 3.2) reduces the risk of first cardiovascular event in rheumatoid arthritis: a time-dependent COX regression analysis in a large cohort study. *Ann Rheum Dis* 2017;76:1693–9.
- 21 Kremer JM. The CORRONA database. *Autoimmun Rev* 2006;5:46–54.
- 22 Kremer JM. The Corona us registry of rheumatic and autoimmune diseases. *Clin Exp Rheumatol* 2016;34:S96–9.
- 23 Arnett FC, Edworthy SM, Bloch DA, *et al.* The American rheumatism association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988;31:315–24.
- 24 Pappas DA, Nyberg F, Kremer JM, *et al.* Prevalence of cardiovascular disease and major risk factors in patients with rheumatoid arthritis: a multinational cross-sectional study. *Clin Rheumatol* 2018;37:2331–40.
- 25 Aletaha D, Nell VPK, Stamm T, *et al.* Acute phase reactants add little to composite disease activity indices for rheumatoid arthritis: validation of a clinical activity score. *Arthritis Res Ther* 2005;7:R796–806.
- 26 Akdemir G, Heimans L, Bergstra SA, *et al.* Clinical and radiological outcomes of 5-year drug-free remission-steered treatment in patients with early arthritis: improved study. *Ann Rheum Dis* 2018;77:111–8.
- 27 Stouten V, Westhovens R, Pazmino S, *et al.* Five-Year treat-to-target outcomes after methotrexate induction therapy with or without other csDMARDs and temporary glucocorticoids for rheumatoid arthritis in the CareRA trial. *Ann Rheum Dis* 2021;80:965–73.
- 28 Verschueren P, Esselens G, Westhovens R. Daily practice effectiveness of a step-down treatment in comparison with a tight step-up for early rheumatoid arthritis. *Rheumatology* 2008;47:59–64.
- 29 Movahedi M, Costello R, Lunt M, *et al.* Oral glucocorticoid therapy and all-cause and cause-specific mortality in patients with rheumatoid arthritis: a retrospective cohort study. *Eur J Epidemiol* 2016;31:1045–55.
- 30 Nadareishvili Z, Michaud K, Hallenbeck JM, *et al.* Cardiovascular, rheumatologic, and pharmacologic predictors of stroke in patients with rheumatoid arthritis: a nested, case-control study. *Arthritis Rheum* 2008;59:1090–6.
- 31 Solomon DH, Avorn J, Katz JN, *et al.* Immunosuppressive medications and hospitalization for cardiovascular events in patients with rheumatoid arthritis. *Arthritis Rheum* 2006;54:3790–8.
- 32 Wolfe F, Michaud K. The risk of myocardial infarction and pharmacologic and nonpharmacologic myocardial infarction predictors in rheumatoid arthritis: a cohort and nested case-control analysis. *Arthritis Rheum* 2008;58:2612–21.
- 33 Luis M, Freitas J, Costa F, *et al.* An updated review of glucocorticoid-related adverse events in patients with rheumatoid arthritis. *Expert Opin Drug Saf* 2019;18:581–90.
- 34 Strehl C, Bijlsma JWJ, de Wit M, *et al.* Defining conditions where long-term glucocorticoid treatment has an acceptably low level of harm to facilitate implementation of existing recommendations: viewpoints from an EULAR Task force. *Ann Rheum Dis* 2016;75:952–7.
- 35 Huscher D, Thiele K, Gromnica-Ihle E, *et al.* Dose-Related patterns of glucocorticoid-induced side effects. *Ann Rheum Dis* 2009;68:1119–24.
- 36 George MD, Baker JF, Winthrop K, *et al.* Risk for serious infection with low-dose glucocorticoids in patients with rheumatoid arthritis : A cohort study. *Ann Intern Med* 2020;173:870–878.
- 37 Yao T-C, Huang Y-W, Chang S-M, *et al.* Association between oral corticosteroid bursts and severe adverse events : A nationwide population-based cohort study. *Ann Intern Med* 2020;173:325–30.
- 38 Ajeganova S, Svensson B, Hafström I, *et al.* Low-Dose prednisolone treatment of early rheumatoid arthritis and late cardiovascular outcome and survival: 10-year follow-up of a 2-year randomised trial. *BMJ Open* 2014;4:e004259.
- 39 Aviña-Zubieta JA, Abrahamowicz M, Choi HK, *et al.* Risk of cerebrovascular disease associated with the use of glucocorticoids in patients with incident rheumatoid arthritis: a population-based study. *Ann Rheum Dis* 2011;70:990–5.
- 40 Gonzalez-Gay MA, Gonzalez-Juanatey C, Lopez-Diaz MJ, *et al.* HLA-DRB1 and persistent chronic inflammation contribute to cardiovascular events and cardiovascular mortality in patients with rheumatoid arthritis. *Arthritis Rheum* 2007;57:125–32.
- 41 Costello RE, Yimer BB, Roads P, *et al.* Glucocorticoid use is associated with an increased risk of hypertension. *Rheumatology* 2021;60:132–9.
- 42 Curtis JR, Westfall AO, Allison J, *et al.* Population-Based assessment of adverse events associated with long-term glucocorticoid use. *Arthritis Rheum* 2006;55:420–6.
- 43 Chester Wasko M, Dasgupta A, Ilse Sears G, *et al.* Prednisone use and risk of mortality in patients with rheumatoid arthritis: moderation by use of disease-modifying antirheumatic drugs. *Arthritis Care Res* 2016;68:706–10.

Supplemental Appendix

June 21, 2021

Section 1. Title Page

Dose and duration-dependent corticosteroid risk for cardiovascular events in corticosteroid-naïve patients with rheumatoid arthritis

Anthony J. Ocon, MD, PhD¹; George Reed, PhD^{2,3}; Dimitrios Pappas, MD^{3,4}; Jeff Curtis, MD, MS, MPH⁵; Joel M. Kremer, MD^{3,6}

1. Department of Medicine, Division of Allergy, Immunology, and Rheumatology, University of Rochester Medical Center, Rochester, NY, USA

2. Department of Medicine, Division of Preventative and Behavioral Medicine, University of Massachusetts Medical School, Worcester, MA

3. The Corrona Research Foundation, LLC, Waltham, MA, USA

4. Department of Medicine, Division of Rheumatology, Columbia University, New York, NY, USA

5. Department of Medicine, Division of Clinical Immunology and Rheumatology, University of Alabama at Birmingham, Birmingham, AL, USA

6. Department of Medicine, Division of Rheumatology, Albany Medical College; The Center for Rheumatology, Albany, NY, USA

Section 2. Table of Contents

1. Title page	Page: 1
2. Table of contents	Page: 2
3. Methods of calculating cumulative prednisone dose/duration; eFigure 1	Page: 3
4. Event collection protocol for CorEvitas (formerly Corrona); eFigure 2	Page: 4
5. Absolute number of patients for each risk assessment, eTable 1	Page: 5
6. Sensitivity analysis: Exclusion of VTE as CVE; eTable 2	Page: 6
7. Sensitivity analysis: Exclusion of patients with prior CVE, eTable 3	Page: 7
8. Sensitivity analysis: Exclusion of “Other” CVE, eTable 4	Page: 8


Section 3. Methods of calculating cumulative prednisone dose/duration of use

Time was divided into days to cover all events and changes of dose or duration of use across all patients. Prednisone dose and usage were updated at each visit for a patient. The assumption was made that a patient was on the prior dose with continuous use until the next visit unless the patient or physician reported a specific date of dose change or cessation of use.

As a time-varying analysis, for each time point after the index date (enrollment), we computed cumulative dose or duration of use over the preceding 6-months or 1-year. This is illustrated in eFigure 1A with an example patient and shows how the preceding 6-month cumulative dose is updated in a time-varying manner. Cumulative dose and duration of use were divided into equivalent quartiles for analysis.

For any given series of days, eFigure 1B illustrated the computation of cumulative total dose or duration of use. The cumulative total dose (or duration of use) was calculated as the area under the dose curve for the preceding 6-months or 1-year interval.

Appendix eFigure 1. Computational model for total prednisone dose and duration.




Section 4. Event collection protocol for CorEvitas (formerly Corrona) registry

The supplemental eFigure 2 below illustrates the collection of targeted adverse events (TAEs) in the CorEvitas (formerly Corrona) registry. All outcomes of interest are TAEs within CorEvitas.

Primary analysis of events was operationalized as follows. The registry used all events reported on physician forms except for events that were 'not confirmed' by the TAE form. Approximately 90% of reported events are validated with a TAE. Of the remaining 10%, approximately 4% were later ruled out and converted to "not an event"; 5-6% were incomplete in the system pending additional site follow-up/TAE completion.

Appendix eFigure 2. Targeted Adverse Events



Section 5. Absolute Number of Patients For Each Risk Assessment

Note that patients may contribute to more than one category over time due to the time varying analysis. Thus, the total number of patients will be greater than the number included in the study.

Appendix eTable 1: Patient Counts

Daily Dose (mg) †	Patients contributing
none	19902
1-<5	722
≥5-9	1598
≥10	1077
Cumulative Dose (mg) †	
Over Preceding 6-Months	
none	19902
1-380	2500
381-750	2396
751-1100	2003
>1110	1001
Over Preceding 1-Year	
none	19902
1-500	2500
501-1100	2314
1101-2100	1710
>2100	919
Duration of Use (days)	
Over Preceding 6-Months	
none	19902
1-80	2500
81-160	2440
161-181	1864
>181	1705
Over Preceding 1-Year	
none	19902
1-100	2500
101-220	2274
221-360	1388
>360	858

Section 6. Sensitivity analysis: Exclusion of VTE as CVE

Appendix eTable 2. Unadjusted and adjusted Hazard Ratios after Exclusion of Deep Vein Thrombosis and Pulmonary Embolism as Cardiovascular Events

Risk for Cardiovascular Events Excluding DVT/PE				
Daily Dose (mg)†	Unadjusted Hazard Ratio	95% Confidence Interval	Adjusted* Hazard Ratio	95% Confidence Interval
none	1	ref	1	ref
1-<5	0.96	0.54-1.70	0.87	0.49-1.54
≥5-9	1.71	1.28-2.29	1.50	1.12-2.01
≥10	2.03	1.37-3.03	1.83	1.23-2.74
Cumulative Dose (mg)†	Unadjusted Hazard Ratio	95% Confidence Interval	Adjusted* Hazard Ratio	95% Confidence Interval
Over Preceding 6-Months				
none	1	ref	1	ref
1-380	0.83	0.49-1.41	0.78	0.46-1.32
381-750	1.14	0.73-1.78	1.05	0.67-1.64
751-1100	1.62	1.16-2.27	1.43	1.02-2.00
>1110	2.44	1.70-3.50	2.19	1.52-3.14
Over Preceding 1-Year				
none	1	ref	1	ref
1-500	0.97	0.61-1.55	0.92	0.58-1.47
501-1100	1.26	0.86-1.83	1.17	0.80-1.71
1101-2100	1.63	1.17-2.28	1.47	1.05-2.07
>2100	1.95	1.38-2.75	1.71	1.20-2.42
Duration of Use (days)	Unadjusted Hazard Ratio	95% Confidence Interval	Adjusted* Hazard Ratio	95% Confidence Interval
Over Preceding 6-Months				
none	1	ref	1	ref
1-80	0.56	0.30-1.05	0.53	0.28-0.99
81-160	1.69	1.18-2.44	1.58	1.10-2.28
160-181	2.18	1.04-4.59	1.98	0.94-4.18
>181	1.77	1.34-2.34	1.54	1.16-2.04
Over Preceding 1-Year				
none	1	ref	1	ref
1-100	1.13	0.75-1.72	1.08	0.71-1.63
101-220	1.48	1.07-2.05	1.39	1.00-1.93
221-360	0.94	0.55-1.59	0.83	0.49-1.42
>360	2.10	1.53-2.89	1.83	1.33-2.51

DVT, deep vein thrombosis; PE, pulmonary embolism.

†, prednisone-equivalents

*adjusted for age, sex, race, duration of RA, history of CV disease, diabetes mellitus, hyperlipidemia, hypertension, statin use, NSAID use, tobacco use, year of enrollment, baseline modified health assessment questionnaire score, CDAI, and cs, b, tsDMARDS use.

Section 7. Sensitivity analysis: Exclusion of patients with prior CV events

With exclusion of patients who had a history of prior CVE, 829 events occurred. The event rate was 1.34 events per 100 patient-years. The total number of patients was 18,168, with 2300 initiating prednisone.

Appendix eTable 3. Unadjusted and adjusted Hazard Ratios after exclusion of patients with a history of prior CVE

Risk for CVE Excluding Patients with Prior History of CVE				
Daily Dose (mg)†	Unadjusted Hazard Ratio	95% Confidence Interval	Adjusted* Hazard Ratio	95% Confidence Interval
none	1	ref	1	ref
1-<5	0.93	0.50-1.74	0.81	0.43-1.51
≥5-9	1.94	1.44-2.62	1.68	1.24-2.27
≥10	2.19	1.43-3.36	2.04	1.33-3.14
Cumulative Dose (mg)†	Unadjusted Hazard Ratio	95% Confidence Interval	Adjusted* Hazard Ratio	95% Confidence Interval
Over Preceding 6-Months				
none	1	ref	1	ref
1-380	0.86	0.49-1.53	0.81	0.46-1.43
381-750	1.34	0.86-2.09	1.20	0.77-1.87
751-1100	1.79	1.27-2.54	1.57	1.11-2.22
>1110	2.38	1.60-3.55	2.19	1.46-3.28
Over Preceding 1-Year				
none	1	ref	1	ref
1-500	0.98	0.59-1.64	0.92	0.51-1.53
501-1100	1.44	0.98-2.12	1.35	0.91-1.98
1101-2100	1.60	1.11-2.31	1.44	0.99-2.07
>2100	2.23	1.57-3.18	1.98	1.38-2.82
Duration of Use (days)	Unadjusted Hazard Ratio	95% Confidence Interval	Adjusted* Hazard Ratio	95% Confidence Interval
Over Preceding 6-Months				
none	1	ref	1	ref
1-80	0.75	0.42-1.36	0.71	0.39-1.29
81-160	1.76	1.19-2.60	1.64	1.11-2.43
160-181	1.47	0.55-3.92	1.34	0.50-3.61
>181	1.92	1.44-2.57	1.65	1.23-2.21
Over Preceding 1-Year				
none	1	ref	1	ref
1-100	1.14	0.73-1.81	1.09	0.69-1.72
101-220	1.63	1.16-2.30	1.54	1.09-2.17
221-360	1.03	0.60-1.79	0.91	0.54-1.58
>360	2.26	1.62-3.15	1.94	1.39-2.71

†, prednisone-equivalents

*adjusted for age, sex, race, duration of RA, history of CV disease, diabetes mellitus, hyperlipidemia, hypertension, statin use, NSAID use, tobacco use, year of enrollment, baseline modified health assessment questionnaire score, CDAI, and cs, b, tsDMARDS use.

Section 8: Sensitivity analysis: Exclusion of “Other” CVE

This sensitivity analysis excluded “Other” CVE to assess for impact on results. With exclusion of “Other” events, a total of 817 events occurred.

Appendix eTable 4. Unadjusted and adjusted Hazard Ratios after Exclusion of Other CVE

Daily Dose (mg) †	Unadjusted Hazard Ratio [95% CI]	Adjusted* Hazard Ratio [95% CI]
none	1 [ref]	1 [ref]
1-<5	1.14 [0.63-2.06]	1.01 [0.56-1.84]
≥5-9	1.86 [1.36-2.55]	1.65 [1.20-2.27]
≥10	2.03 [1.40-3.17]	1.91 [1.22-2.99]
Cumulative Dose (mg) †	Unadjusted Hazard Ratio [95% CI]	Adjusted* Hazard Ratio [95% CI]
Over Preceding 6-Months		
none	1 [ref]	1 [ref]
1-380	0.90 [0.51-1.59]	0.84 [0.47-1.49]
381-750	1.23 [0.76-2.00]	1.13 [0.70-1.83]
751-1100	1.90 [1.34-2.71]	1.69 [1.19-2.41]
>1110	2.30 [1.51-3.48]	2.16 [1.42-3.29]
Over Preceding 1-Year		
none	1 [ref]	1 [ref]
1-500	1.02 [0.61-1.70]	0.96 [0.58-1.61]
501-1100	1.42 [0.95-2.12]	1.32 [0.88-1.97]
1101-2100	1.73 [1.20-2.50]	1.58 [1.09-2.29]
>2100	2.09 [1.43-3.07]	1.90 [1.29-2.79]
Duration of Use (days)	Unadjusted Hazard Ratio [95% CI]	Adjusted* Hazard Ratio [95% CI]
Over Preceding 6-Months		
none	1 [ref]	1 [ref]
1-80	0.77 [0.43-1.40]	0.73 [0.40-1.33]
81-160	1.63 [1.07-2.47]	1.52 [1.00-2.31]
161-181	1.17 [0.38-3.63]	1.10 [0.35-3.42]
>181	2.04 [1.52-2.74]	1.80 [1.33-2.43]
Over Preceding 1-Year		
none	1 [ref]	1 [ref]
1-100	1.17 [0.74-1.85]	1.12 [0.70-1.75]
101-220	1.42 [0.98-2.07]	1.36 [0.93-1.98]
221-360	1.29 [0.77-2.15]	1.16 [0.69-1.94]
>360	2.38 [1.69-3.37]	2.08 [1.47-2.95]

†, prednisone-equivalents

*adjusted for age, sex, race, duration of RA, history of CV disease, diabetes mellitus, hyperlipidemia, hypertension, statin use, NSAID use, tobacco use, year of enrollment, baseline modified health assessment questionnaire score, CDAI, and cs, b, tsDMARDS use.

Steroid dose and duration linked to cardiovascular events in RA



Dose and duration are associated with an increased risk for incident cardiovascular events in steroid-naïve patients

INTRODUCTION

Rheumatoid arthritis is a chronic inflammatory disease that affects a person's joints, and may cause pain and disability. Rheumatoid arthritis can affect people of all ages, but it most often starts between the ages of 30 and 50 – although this can depend on where you live. Rheumatoid arthritis is more common in women than men.

People with rheumatoid arthritis have an increased risk of suffering from cardiovascular diseases such as heart attacks, heart failure or stroke. This is because the inflammation involved in rheumatoid arthritis can have an effect on other systems in the body, as well as the joints. Some medicines used to treat rheumatoid arthritis can also increase the risk of cardiovascular disease.

Glucocorticoids are a type of steroid medicine used to treat rheumatoid arthritis and other inflammatory conditions. Although most doctors are aware of the risk of using high doses of these medicines, many think that low doses are safe. However, one long-term risk of using glucocorticoids is an increase in cardiovascular events. There is not much information about the risks of low-dose or relatively short-term use of these medicines.

WHAT DID THE AUTHORS HOPE TO FIND?

The authors wanted to find out the safety and risk of using relatively low doses of glucocorticoids for short periods. Since people with rheumatoid arthritis already have an increased risk of cardiovascular disease, they wondered what effect this would have in terms of new cardiovascular events.

WHO WAS STUDIED?

The study looked at over 19,000 people with rheumatoid arthritis who had not taken glucocorticoids before. Everyone was over the age of 18 and living in the United States.

HOW WAS THE STUDY CONDUCTED?

This was a retrospective observational study, which means that the authors used an existing database of patient records to look back and find people for the study. There was no interventional treatment given. The database used was from the CorEvitas registry, which regularly collects information from people with rheumatoid arthritis and their doctors. People were included in this study if they started taking a glucocorticoid for the first time after they enrolled in the CorEvitas registry. The authors used this information to compare the risk of new cardiovascular events in people who started glucocorticoids to those who did not. The results were analysed based on the dose and duration of use.

WHAT WERE THE MAIN FINDINGS OF THE STUDY?

The main finding was that daily glucocorticoid doses of less than 5 mg of prednisone-equivalent were not associated with an increased risk, but all higher daily doses were. The risk increased directly with dose, with the greatest estimated risks for people taking 5–9 mg or more than 10 mg per day.

As well as the daily dose, the authors looked at cumulative use over the past 6 months for each person. They found there was a dose threshold that was associated with an increased risk of cardiovascular events over time. However, there was no increased risk if people took less than 750 mg over 6 months, or less than 1100 mg over 1 year.

The authors also found a threshold for how long these medicines should be used for. The results showed no increased risk in people who used glucocorticoids for up to 80 days over the preceding 6 months, or up to 100 days over 1 year. There was no increased risk for cardiovascular events in people taking glucocorticoid medicines for fewer days.

ARE THESE FINDINGS NEW?

Yes. This is the first time these thresholds have been described for people with rheumatoid arthritis taking glucocorticoid medicines.

WHAT ARE THE LIMITATIONS OF THE STUDY?

One limitation is that the study used retrospective information from an existing database study. People taking part in these kinds of observational registries are treated without being assigned to a specific intervention. The authors took into account other factors that might have affected the results, but there is still a risk of possible bias, or that other external factors might have had an impact.

WHAT DO THE AUTHORS PLAN ON DOING WITH THIS INFORMATION?

The authors hope this information will enhance education for doctors and their patients on the risks of low-dose, short-term glucocorticoids. Additional studies are needed to see if the risk is the same in other diseases that are treated with these medicines.

WHAT DOES THIS MEAN FOR ME?

If you have rheumatoid arthritis, you should be aware of the risk of cardiovascular events even with relatively short-term use of steroid medicines. Your doctor should try to taper (reduce) the amount of medicine you take to the lowest possible dose that still controls your rheumatoid arthritis.

It is important that you do not change the dose of your medicine yourself. If you have any concerns about your disease or its treatment, you should talk to your doctor.

Disclaimer: This is a summary of a scientific article written by a medical professional (“the Original Article”). The Summary is written to assist non medically trained readers to understand general points of the Original Article. It is supplied “as is” without any warranty. You should note that the Original Article (and Summary) may not be fully relevant nor accurate as medical science is constantly changing and errors can occur. It is therefore very important that readers not rely on the content in the Summary and consult their medical professionals for all aspects of their health care and only rely on the Summary if directed to do so by their medical professional. Please view our full Website Terms and Conditions. <http://www.bmj.com/company/legal-information/>

Date prepared: November 2021

Summary based on research article published on: 2 July 2021

From: Ocon AJ, *et al.* Short-term dose and duration-dependent glucocorticoid risk for cardiovascular events in glucocorticoid-naïve patients with rheumatoid arthritis. *Ann Rheum Dis* 2021;80:1522–29. doi:10.1136/annrheumdis-2021-220577

Copyright © 2021 BMJ Publishing Group Ltd & European League Against Rheumatism. Medical professionals may print copies for their and their patients and students non commercial use. Other individuals may print a single copy for their personal, non commercial use. For other uses please contact our [Rights and Licensing Team](#).