Supplemental material

Author Year	Effect Size S	tandard err	or	HR	95%-CI	Weight
IFG-ADA Hubbard, 2019 Parrinello, 2016 Deedwania, 2013 Parrinello, 2016 Jansky, 2008 Overall effect Heterogeneity: $I^2 = 38$	-0.63 -0.24 -0.02 -0.01 0.12 5%, <i>p</i> = 0.19	0.4019 0.1179 0.0749 0.0719 0.1624		0.79 0.98 0.99 1.13	[0.24; 1.17] [0.63; 1.00] [0.85; 1.13] [0.86; 1.14] [0.82; 1.55] [0.84; 1.07]	
IGT Fang, 2019 Overall effect Heterogeneity: not ap	1.62 oplicable	0.8252			[1.00; 25.40] <b>[1.00; 25.40]</b>	
HbA1c-ADA Hubbard, 2019 Parrinello, 2016 Parrinello, 2016 Overall effect Heterogeneity: $I^2 = 6$	-0.63 0.12 0.25 1%, p = 0.08	0.4019 0.1163 0.0757	+	1.13 1.28	[0.24; 1.17] [0.90; 1.42] [1.10; 1.48] [0.87; 1.45]	3.1% 14.3% 17.7% <b>35.2%</b>
<b>Overall effect</b> Heterogeneity: $I^2 = 6$ Residual heterogenei		0.08	0.5 1 2 4 Hazard Ratio		[0.87; 1.17]	100.0%